SOLVABLE LIE GROUPS OF NEGATIVE RICCI CURVATURE

Y. NIKOLAYEVSKY AND YU.G. NIKONOROV

ABSTRACT. We consider the question of whether a given solvable Lie group admits a left-invariant metric of strictly negative Ricci curvature. We give necessary and sufficient conditions of the existence of such a metric for the Lie groups the nilradical of whose Lie algebra is either abelian or Heisenberg or standard filiform, and discuss some open questions.

1. INTRODUCTION

The fundamental question of Riemannian geometry is whether (and when) a given manifold admits a Riemannian metric with a particular sign of the curvature. Naturally, in the context of homogeneous geometry, the same question is being asked for leftinvariant metrics. In that case the curvature is entirely expressed in terms of the algebraic structure of the given homogeneous space and one expects the answer to be stated in both topological and algebraic terms.

The conditions on the sign of the sectional curvature K_{σ} are quite restrictive and the homogeneous spaces whose sectional curvature has a particular sign are mainly understood. For $K_{\sigma} > 0$ the question was settled in [Wal, BB1] who showed that a homogeneous space admits a left-invariant metric with $K_{\sigma} > 0$ if and only if it is diffeomorphic to either CROSS or to a space from a short finite list (so-called Wallach and Allof-Wallach spaces). By [Ale, Hei], a homogeneous space of negative sectional curvature is isometric to a solvmanifold the nilradical \mathfrak{n} of whose Lie algebra \mathfrak{g} has codimension one and there exists $Y \in \mathfrak{g} \setminus \mathfrak{n}$ such that all the eigenvalues of the restriction of ad_Y to \mathfrak{n} have positive real part. Flat homogeneous spaces were completely described in [Ale, BB2]: every such space is isometric to a solvmanifold the nilradical \mathfrak{n} of whose Lie algebra \mathfrak{g} is abelian and the restrictions of all ad_Y , $Y \in \mathfrak{g} \setminus \mathfrak{n}$, to \mathfrak{n} are skew-symmetric.

For the Ricci curvature, the positive case was settled down by Milnor [Mil] (for Lie groups), and by Berestovskii [Ber] (in the general case), who proved that a homogeneous space admits a left-invariant metric with Ric > 0 if and only if it is compact and has

²⁰¹⁰ Mathematics Subject Classification. Primary 53C30, 22E25.

Key words and phrases. Solvable Lie algebra, nilradical, negative Ricci curvature.

The first author is partially supported by ARC Discovery Grant DP130103485. The second author is supported in part by the State Maintenance Program for the Leading Scientific Schools of the Russian Federation (grant NSh-921.2012.1) and by Federal Target Grant "Scientific and educational personnel of innovative Russia" for 2009-2013 (agreement no. 8206, application no. 2012-1.1-12-000-1003-014).

a finite fundamental group. By a beautiful result of [AK], any Ricci-flat homogeneous space is flat.

Much less is known, however, about Riemannian homogeneous spaces of negative Ricci curvature. By the following theorem, no unimodular solvable Lie group (in particular, no nilpotent group) admits a left-invariant metric with Ric < 0:

Theorem 1 ([DM]). Any left-invariant metric with $\text{Ric} \leq 0$ on a solvable unimodular Lie group is Ricci-flat.

By [AK], any such metric is flat. Further in [DLM] it was proved that a unimodular Lie group which admits a left-invariant metric with Ric < 0 is noncompact and semisimple. Examples of such metrics were constructed on SL (n, \mathbb{R}) , $n \ge 3$ [LDM] and on some complex simple Lie groups [DLM]. To the best of our knowledge, however, the general (nonunimodular) case has not been studied in the literature, although an important subclass of left-invariant metrics with negative Ricci curvature, the Einstein metrics of negative scalar curvature, has been extensively studied in the past decades by many authors, including the present authors.

In this paper we ask the following question: Which solvable Lie groups admit a leftinvariant metric of negative Ricci curvaturee?

The Ricci curvature of a left-invariant metric on a Lie group G can be entirely computed from the algebraic data: the structure of the Lie algebra \mathfrak{g} of G and the inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{g} (see Section 2.1 for details). With a slight abuse of terminology, we will speak of the Ricci curvature of the metric Lie algebra $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$.

After some preliminaries (Section 2), we prove in Section 3 the following theorem, which gives some necessary conditions and some sufficient conditions for the existence of an inner product of negative Ricci curvature on a solvable Lie algebra.

Theorem 2. Suppose \mathfrak{g} is a solvable Lie algebra. Let \mathfrak{n} be the nilradical of \mathfrak{g} and \mathfrak{z} be the centre of \mathfrak{n} . Then

- (1) If \mathfrak{g} admits an inner product of negative Ricci curvature, then there exists $Y \in \mathfrak{g}$ such that $\operatorname{Tr} \operatorname{ad}_Y > 0$ and all the eigenvalues of the restriction of the operator ad_Y to \mathfrak{z} have a positive real part;
- (2) If there exists $Y \in \mathfrak{g}$ such that all the eigenvalues of the restriction of ad_Y to \mathfrak{n} have positive real part, then \mathfrak{g} admits an inner product of negative Ricci curvature.

The following is an immediate consequence of Theorem 2.

Corollary. A solvable Lie algebra \mathfrak{g} with an abelian nilradical \mathfrak{n} admits an inner product of negative Ricci curvature if and only if there exists $Y \in \mathfrak{g}$ such that all the eigenvalues of the restriction of ad_Y to \mathfrak{n} have positive real part.

Note that the latter property of \mathfrak{g} in the Corollary is equivalent to the following one: There exists $Y' \in \mathfrak{g}$ such that the restriction of $\operatorname{ad}_{Y'}$ to \mathfrak{n} is a stable linear operator, i. e. all its eigenvalues have negative real parts (see e. g. [Gant] or [LiW] for a discussion on stable linear operators and stable matrices).

 $\mathbf{2}$

We further develop the approach taken in the Corollary (obtaining the condition for Ric < 0 for classes of solvable algebras with a given nilradical) for two important classes of nonabelian nilpotent algebras serving as nilradicals: the Heisenberg algebra and the standard filiform algebra.

Recall that the Heisenberg Lie algebra \mathfrak{h}_{2p+1} of dimension l = 2p+1, $p \geq 1$, is defined by the relations $[X_i, X_{p+i}] = Z$ for $i = 1, \ldots, p$, relative to a basis $\{X_1, \ldots, X_{2p}, Z\}$, where $[X_i, X_j] = 0$ if $|i - j| \neq p$, and Z spans the centre \mathfrak{z} of \mathfrak{h}_{2p+1} . Let \mathfrak{g} be a solvable Lie algebra with the nilradical $\mathfrak{n} = \mathfrak{h}_{2p+1}$. For any $X \in \mathfrak{g}$, the vector Z is an eigenvector of the restriction of ad_X to \mathfrak{n} , so that $[X, Z] = \lambda(X)Z$ for a one-form λ on \mathfrak{g} , and moreover, $(\mathrm{ad}_X)_{|\mathfrak{n}}$ descends to a well-defined linear map $\Phi(X) \in \mathrm{End}(\mathfrak{n}/\mathfrak{z})$. Let $d_i(X) \in \mathbb{C}, i = 1, \ldots, 2p$, be the eigenvalues of $\Phi(X)$, each listed with its algebraic multiplicity. In Section 4 we prove the following theorem.

Theorem 3. A solvable Lie algebra \mathfrak{g} with the Heisenberg nilradical admits an inner product of negative Ricci curvature if and only if there exists $Y \in \mathfrak{g}$ such that in the above notation,

$$\lambda(Y) + \sum_{i:\operatorname{Re} d_i(Y) < 0} \operatorname{Re} d_i(Y) > 0.$$

Further on, in Section 5 we consider solvable Lie algebras whose nilradical is *filiform* (has the maximal possible degree of nilpotency) [Ver]. More specifically, we require the nilradical to be *standard filiform Lie algebra*, which is defined as the *l*-dimensional Lie algebra L_l having a basis X_1, \ldots, X_l such that $[X_1, X_i] = X_{i+1}, i = 2, \ldots, l-1, [X_1, X_l] = 0$, and $[X_i, X_j] = 0$ when $i, j \geq 2$. Note that any filiform Lie algebra admits a basis for which the former relations are satisfied (but in general, not the latter ones).

Let \mathfrak{g} be a solvable Lie algebra with the nilradical L_l . We can assume that $l \geq 4$ (as L_2 is abelian and L_3 is the Heisenberg algebra). The algebra L_l has a (unique) codimension one abelian ideal $\mathfrak{i} = \operatorname{Span}(X_2, \ldots, X_l)$ and the one-dimensional centre $\mathbb{R}X_l$. Both of them are characteristic ideals of L_l (they are invariant under the action of any derivation on L_l ; see Section 5). Define the one-forms λ and ι on \mathfrak{g} as follows: for $Y \in \mathfrak{g}$, $[Y, X_l] = \lambda(Y)X_l$ and $\iota(Y) = \operatorname{Tr}((\operatorname{ad}_Y)_{|\mathfrak{i}})$. We have the following theorem.

Theorem 4. Let \mathfrak{g} be a solvable Lie algebra with the nilradical $\mathfrak{n} = L_l$, $l \geq 4$. The algebra \mathfrak{g} admits an inner product of negative Ricci curvature if and only if there exists $Y \in \mathfrak{g}$ such that $\lambda(Y) > 0$ and $\iota(Y) > 0$.

In the last section we collect some open questions and conjectures. The main of them, motivated by the above theorems is the following (in a slightly vague formulation). Is it true that a solvable Lie algebra \mathfrak{g} admits an inner product with Ric < 0 if and only if there exists a vector $Y \in \mathfrak{g}$ such that the real parts of the eigenvalues of the restriction of ad_Y to the nilradical \mathfrak{n} of \mathfrak{g} satisfy certain linear inequalities which are determined by the structure of \mathfrak{n} ? Speculating a little further we might suggest that such inequalities represent the fact that $\mathrm{Re}(\mathrm{ad}_Y)_{|\mathfrak{n}}$ belongs to a certain open convex hull (see the details in Section 6).

2. Preliminaries

2.1. The Ricci operator. Let G be a Lie group with a left-invariant metric Q obtained by the left translations from an inner product $\langle \cdot, \cdot \rangle$ on the Lie algebra \mathfrak{g} of G. Let B be the Killing form of \mathfrak{g} , and let $H \in \mathfrak{g}$ be the mean curvature vector defined by $\langle H, X \rangle = \operatorname{Tr} \operatorname{ad}_X$.

The Ricci curvature ric of the metric Lie group (G, Q) at the identity is given by

$$\operatorname{ric}(X) = -\langle [H, X], X \rangle - \frac{1}{2}B(X, X) - \frac{1}{2}\sum_{i} \| [X, E_i] \|^2 + \frac{1}{4}\sum_{i,j} \langle [E_i, E_j], X \rangle^2,$$

for $X \in \mathfrak{g}$, where $\{E_i\}$ is an orthonormal basis for $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ (e.g. [Ale]). Equivalently, one can define the Ricci operator Ric of the metric Lie algebra $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$, the symmetric operator associated to ric, by

(1)
$$\operatorname{Ric} = -\frac{1}{2} \sum_{i} \operatorname{ad}_{E_{i}}^{t} \operatorname{ad}_{E_{i}} + \frac{1}{4} \sum_{i} \operatorname{ad}_{E_{i}} \operatorname{ad}_{E_{i}}^{t} - \frac{1}{2} B - (\operatorname{ad}_{H})^{s},$$

where A^t is the operator adjoint to A and $(ad_H)^s = \frac{1}{2}(ad_H + ad_H^t)$ is the symmetric part of ad_H .

If $(\mathfrak{n}, \langle \cdot, \cdot \rangle)$ is a nilpotent metric Lie algebra, then H = 0 and B = 0, and we get

(2)
$$\langle \operatorname{Ric}^{\mathfrak{n}} X, Y \rangle = \frac{1}{4} \sum_{i,j} \langle X, [E_i, E_j] \rangle \langle Y, [E_i, E_j] \rangle - \frac{1}{2} \sum_{i,j} \langle [X, E_i], E_j \rangle \langle [Y, E_i], E_j] \rangle.$$
$$\operatorname{Ric}^{\mathfrak{n}} = -\frac{1}{2} \sum_i \operatorname{ad}_{E_i}^t \operatorname{ad}_{E_i} + \frac{1}{4} \sum_i \operatorname{ad}_{E_i} \operatorname{ad}_{E_i}^t.$$

We will use a more explicit form of (1) in the case when \mathfrak{g} is solvable. Denote \mathfrak{n} the nilradical of \mathfrak{g} , the maximal nilpotent ideal of \mathfrak{g} . Clearly, $[\mathfrak{g}, \mathfrak{g}] \subset \mathfrak{n}$, but in general, $[\mathfrak{g}, \mathfrak{g}] \neq \mathfrak{n}$. It is known that dim $(\mathfrak{n}) \geq \frac{1}{2} (\dim(\mathfrak{g}) + \dim(\mathfrak{g}(\mathfrak{g})))$, where $\mathfrak{g}(\mathfrak{g})$ is the center of \mathfrak{g} [VGO, Theorem 5.2]. Denote $\mathfrak{a} = \mathfrak{n}^{\perp}$ and put $l = \dim \mathfrak{n}$ and $m = \dim \mathfrak{a}$ (we call m the rank of \mathfrak{g}), with $l + m = n = \dim \mathfrak{g}$. Choose orthonormal bases $\{e_i\}$ for \mathfrak{n} , and $\{f_k\}$ for \mathfrak{a} in such a way that

$$t := \operatorname{Tr}(\operatorname{ad}(f_1)) \ge 0, \quad \operatorname{Tr}(\operatorname{ad}(f_j)) = 0, \quad 2 \le j \le m$$

It is easy to see that for a non-unimodular Lie algebra \mathfrak{g} we have $f_1 = ||H||^{-1}H$, $t = \operatorname{Tr}(\operatorname{ad}(f_1)) = ||H|| > 0$. If \mathfrak{g} is unimodular, we can choose $f_1 \in \mathfrak{a}$ arbitrarily (and t = 0).

Relative to the basis $\{e_1, ..., e_l, f_1, ..., f_m\}$, the matrices of the operators ad_{f_j} and ad_{e_i} have the form

(3)
$$\operatorname{ad}_{f_j} = \begin{pmatrix} A_j & B_j \\ 0 & 0 \end{pmatrix}, \quad \operatorname{ad}_{e_i} = \begin{pmatrix} D_i & C_i \\ 0 & 0 \end{pmatrix},$$

for some $(l \times l)$ -matrices A_j , D_i and $(l \times m)$ -matrices B_j , C_i , and the matrix of the Ricci operator of the solvable metric Lie algebra $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ has the form (see the proof of [NN,

Theorem 3])

(4)
$$\operatorname{Ric} = \left(\begin{array}{cc} R_1 & R_2 \\ R_2^t & R_3 \end{array}\right),$$

where

(5)
$$R_1 = \operatorname{Ric}^{\mathfrak{n}} + \frac{1}{2} \sum_{j=1}^{m} [A_j, A_j^t] + \frac{1}{4} \sum_{j=1}^{m} B_j B_j^t - t A_1^s,$$

(6)
$$R_2 = -\frac{1}{2} \Big(\sum_{i=1}^l D_i^t C_i + \sum_{j=1}^m A_j^t B_j + t B_1 \Big),$$

(7)
$$R_3 = -\frac{1}{2} \sum_{j=1}^m B_j^t B_j - L,$$

where L is an $(m \times m)$ -matrix with the entries $L_{pq} = \text{Tr}(A_p^s A_q^s)$, $A_j^s = \frac{1}{2}(A_j^t + A_j)$ is the symmetric part of A_j , $t = \text{Tr}(A_1) = \text{Tr}(A_1^s) \ge 0$ and Ricⁿ is the matrix of the Ricci operator of the metric nilpotent Lie algebra $(\mathfrak{n}, \langle \cdot, \cdot \rangle_{\mathfrak{n}})$ relative to the basis $\{e_1, \ldots, e_l\}$ which from (2) is easily seen to be given by

(8)
$$\operatorname{Ric}^{\mathfrak{n}} = -\frac{1}{2} \sum_{i=1}^{l} D_{i}^{t} D_{i} + \frac{1}{4} \sum_{i=1}^{l} D_{i} D_{i}^{t}.$$

2.2. Orbit closure. Let \mathfrak{g} be a Lie algebra with the underlying linear space \mathbb{R}^n . We denote μ the Lie bracket of \mathfrak{g} , so that $\mu : \Lambda^2 \mathbb{R}^n \to \mathbb{R}^n$ is the defined by $\mu(X,Y) = [X,Y]$, for $X, Y \in \mathbb{R}^n$. The map μ is an element of the space $\mathcal{L} \subset \Lambda^2(\mathbb{R}^{n*}) \otimes \mathbb{R}^n$ of Lie brackets on \mathbb{R}^n (skew-symmetric bilinear maps satisfying the Jacobi identity). The space \mathcal{L} is acted upon by the group $\operatorname{GL}(n)$ as follows ("change of basis"): for $T \in \operatorname{GL}(n)$ and $\mu \in \mathcal{L}$ we define $T.\mu \in \mathcal{L}$ by $(T.\mu)(X,Y) = T^{-1}\mu(TX,TY)$. It is clear that the Lie algebra defined by the bracket $T.\mu$ on \mathbb{R}^n is isomorphic to the one defined by the bracket μ . As \mathcal{L} is defined by polynomial equations, any element of the closure of the orbit $\operatorname{GL}(n).\mu$ of μ (in the usual topology of $\Lambda^2(\mathbb{R}^{n*}) \otimes \mathbb{R}^n$) is again a Lie bracket, but the corresponding Lie algebra $\overline{\mathfrak{g}}(\mathbb{R}^n, \nu)$ may not be isomorphic to (\mathbb{R}^n, μ) . We say that $\overline{\mathfrak{g}}$ is a *degeneration* of \mathfrak{g} ($\overline{\mathfrak{g}}$ is usually "more abelian" than \mathfrak{g} , see e.g. [Bur, NP]).

The following proposition is elementary, but useful.

Proposition 1. Suppose μ and ν are Lie brackets on \mathbb{R}^n such that ν belongs to the closure of the GL(n) orbit of μ . If the Lie algebra (\mathbb{R}^n, ν) admits an inner product of negative Ricci curvature, then so does the Lie algebra (\mathbb{R}^n, μ) .

Proof. Let $\langle \cdot, \cdot \rangle$ be such an inner product on (\mathbb{R}^n, ν) . As the Ricci tensor depends continuously on the structural constants of the Lie algebra relative to a fixed basis, there is a $T \in \operatorname{GL}(n)$ such that the metric Lie algebra $(\mathbb{R}^n, T.\mu, \langle \cdot, \cdot \rangle)$ has negative Ricci curvature. But the latter Lie algebra is isomorphic to the Lie algebra (\mathbb{R}^n, μ) .

2.3. Technical lemmas. We will need the following two facts.

The first one is a "real" modification of the Lie Theorem (this should be well-known; we supply a short proof for the sake of completeness).

Lemma 1. Let \mathfrak{s} be a real solvable Lie algebra and V be a real \mathfrak{s} -module. Then there exists a basis \mathcal{B} for V relative to which all the elements $g \in \mathfrak{s}$ are block-lower-triangular, with the diagonal blocks of sizes either 1×1 with the corresponding diagonal entry $\lambda_i(g)$,

or 2×2, with the corresponding diagonal block $G_i(g) = \begin{pmatrix} \alpha_i(g) & \beta_i(g) \\ -\beta_i(g) & \alpha_i(g) \end{pmatrix}$, where $\lambda_j, \alpha_i, \beta_i$ are linear forms on \mathfrak{s} such that $\beta_i \neq 0$.

Proof. The proof essentially mimics that for the classical Lie Theorem. All the elements $g \in \mathfrak{s}$ have a common eigenvector $X \in V^{\mathbb{C}}$. If X is real, then $gX = \lambda(g)X$ for some linear form λ on V, and $V/(\mathbb{R}X)$ is again an \mathfrak{s} -module. Otherwise, $X = X_1 + X_2 \mathfrak{i}$, where $X_1, X_2 \in V$ are non-collinear, in which case $L = \operatorname{Span}(X_1, X_2)$ is an \mathfrak{s} -module and the representation G of \mathfrak{s} on L relative to the basis $\{X_1, X_2\}$ is given by $G(g) = \begin{pmatrix} \alpha(g) & \beta(g) \\ -\beta(g) & \alpha(g) \end{pmatrix}$, for some linear forms α, β on $\mathfrak{s}, \beta \neq 0$. Then V/L is again a \mathfrak{s} -module and the proof follows by induction.

We will also need the following "general position" lemma.

Lemma 2. Let V be a real linear space and $\mathfrak{s} \subset \mathfrak{gl}(V)$ a solvable subalgebra. Then for an open, dense set of inner products on V, no nonzero element of \mathfrak{s} is skew-symmetric.

Proof. Let \mathcal{B}_0 be an arbitrary basis for V and let $\langle \cdot, \cdot \rangle_0$ be the inner product for which \mathcal{B}_0 is orthonormal. Then, relative to \mathcal{B}_0 , the inner products $\langle \cdot, \cdot \rangle$ on V are in one-to-one correspondence with symmetric positive definite matrices Q, so that $\langle X, Y \rangle = \langle QX, Y \rangle_0$, for all $X, Y \in V$. Choose a basis for \mathfrak{s} represented by the matrices M_1, \ldots, M_m relative to \mathcal{B}_0 . The fact that for the inner product corresponding to a matrix Q the subalgebra \mathfrak{s} contains a nonzero skew-symmetric operator is equivalent to the fact that the matrices $QM_i + M_i^t Q$ are linearly dependent, which is a polynomial condition for the entries of Q. It follows that it suffices to find at least one Q for which it is violated, which is equivalent to finding a basis for V relative to which no nonzero element of \mathfrak{s} is represented by a skew-symmetric matrix. Take a basis \mathcal{B} constructed in Lemma 1 and modify it as follows: for every pair of basis vectors corresponding to a 2×2 diagonal block G_i , multiply one of them by 2. Then no nonzero matrix from \mathfrak{s} is skew-symmetric relative to the resulting basis.

3. Abelian Nilradical and the proof of Theorem 2

Theorem 2 implies the Corollary which answers our question for solvable Lie algebras with the abelian nilradical. For the convenience of the reader, we reproduce it here (with a slight change of notation). **Theorem 2.** Suppose \mathfrak{g} is a solvable Lie algebra. Let \mathfrak{n} be the nilradical of \mathfrak{g} and \mathfrak{z} be the centre of \mathfrak{n} . Then

- (1) If \mathfrak{g} admits an inner product of negative Ricci curvature, then there exists $X \in \mathfrak{g}$ such that $\operatorname{Tr} \operatorname{ad}_X > 0$ and all the eigenvalues of the restriction of the operator ad_X to \mathfrak{z} have a positive real part;
- (2) If there exists $X \in \mathfrak{g}$ such that all the eigenvalues of the restriction of ad_X to \mathfrak{n} have positive real part, then \mathfrak{g} admits an inner product of negative Ricci curvature.

Proof. (1) First of all, note that \mathfrak{z} is an ideal of \mathfrak{g} , so for any $Y \in \mathfrak{g}$, the restriction of ad_Y to \mathfrak{z} is well-defined. Moreover, as \mathfrak{z} is abelian, all such restrictions commute. Suppose that for an inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{g} , the Ricci curvature is negative.

Take $X = f_1$. Let $\xi \in \mathbb{C}$ be an eigenvalue of the restriction of ad_{f_1} to \mathfrak{z} and let $V_{\xi} \subset \mathfrak{z}^{\mathbb{C}}$ be the corresponding eigenspace. As the operators $(\mathrm{ad}_{f_j})_{|\mathfrak{z}|}$ commute, the subspace V_{ξ} is ad_{f_j} -invariant, for all $j = 1, \ldots, m$, and moreover, the restrictions of ad_{f_j} to V_{ξ} again commute. By the Lie Theorem (or by Frobenius Theorem, [New]), the latter have a common eigenvector, so that there exists $Z \in V_{\xi} \setminus 0$ such that $\mathrm{ad}_{f_j} Z = \xi_j Z$, and $\xi_1 = \xi$. Let $Z = Z_1 + Z_{2i}$, $Z_1, Z_2 \in \mathfrak{z}$. Multiplying Z by a nonzero complex number we can assume that $||Z_1|| = 1$ and that $Z_1 \perp Z_2$ (note that Z_2 can be zero). We have $A_j Z_1 = \alpha_j Z_1 - \beta_j Z_2$, $A_j Z_2 = \alpha_j Z_2 + \beta_j Z_1$, where $\xi_j = \alpha_j + \beta_j i$, $j = 1, \ldots, m$. Computing $\langle R_1 Z_1, Z_1 \rangle + \langle R_1 Z_2, Z_2 \rangle$ by (5) and (8) we obtain $\langle (-\frac{1}{2} \sum_{i=1}^{l} D_i^t D_i + \frac{1}{4} \sum_{i=1}^{l} D_i D_i^t) Z_k, Z_k \rangle = \frac{1}{4} \sum_{i=1}^{l} ||D_i^t Z_k||^2 \ge 0$, for k = 1, 2, as $D_i Z_k = 0$, so $\langle \operatorname{Ric}^n Z_k, Z_k \rangle \ge 0$. Moreover, $\langle B_j B_j^t Z_k, Z_k \rangle \ge 0$. Furthermore, for $j = 1, \ldots, m$ we have $\langle ([A_j, A_j^t] Z_1, Z_1) + \langle ([A_j, A_j^t] Z_2, Z_2) \rangle = ||A_j^t Z_1||^2 - ||A_j Z_1||^2 + ||A_j^t Z_2||^2 - ||A_j Z_2||^2 = \sum_{i=1}^{l} \langle A_j e_i, Z_1 \rangle^2 + \langle A_j e_i, Z_2 \rangle^2 - ||A_j Z_1||^2 - ||A_j Z_2||^2$. Specifying the orthonormal basis $\{e_i\}$ in such a way that $e_1 = Z_1$ and $e_2 = ||Z_2||^{-1} Z_2$ if $Z_2 \neq 0$ (or arbitrary otherwise) we find that the latter expression equals $\sum_{i=3}^{l} \langle A_j e_i, Z_1 \rangle^2 + \langle A_j e_i, Z_2 \rangle^2 \rangle = \langle A_1 Z_1, Z_1 \rangle + \langle (I_A, A_j^t |Z_1, Z_1 \rangle + \langle (I_A, A_j^t |Z_2, Z_2 \rangle \geq 0$. Finally we have $\langle A_1^s Z_1, Z_1 \rangle + \langle A_1^s Z_2, Z_2 \rangle \geq \langle A_1 Z_1, Z_1 \rangle + \langle A_1 Z_2, Z_2 \rangle \geq \alpha_1 (1 + ||Z_2||^2)$. It follows that $\langle R_1 Z_1, Z_1 \rangle + \langle R_1 Z_2, Z_2 \rangle \geq \alpha_1 (1 + ||Z_2||^2)$ Tr A_1 , which implies $\operatorname{Re}(\xi) \operatorname{Tr} A_1 < 0$.

(2) By Lemma 1, we can choose a basis $\mathcal{B}' = \{X_i\}_{i=1}^l$ for \mathfrak{n} relative to which all the operators $(\mathrm{ad}_Y)_{|\mathfrak{n}}, Y \in \mathfrak{g}$, are block-lower-triangular, with p (1 × 1)-blocks and q (2 × 2)-blocks, where $2q + p = l, q, p \ge 0$.

For $a = 1, \ldots, p + q$, let $l_a \in \{1, 2\}$ be the dimension of the *a*-th diagonal block (counting from the top-left to the bottom-right corner), and let N be a diagonal matrix (relative to \mathcal{B}') defined by $N = \text{diag}(d_1I_{l_1}, d_2I_{l_2}, \ldots, d_{p+q}I_{l_{p+q}})$, where d_a , $a = 1, \ldots, p+q$, is a sequence of positive numbers such that $d_a + d_b < d_{\max(a,b)+1}$ (say $d_a = 3^a$).

Extend the basis \mathcal{B}' to a basis \mathcal{B} for \mathfrak{g} by elements $\{Y_k\}_{k=1}^m$, and for $s \in \mathbb{R}$, define $T_s \in \operatorname{End}(\mathfrak{g})$ by $T_s X = e^{sN} X$ for $X \in \mathfrak{n}$ and $T_s Y_k = Y_k$. Let μ be the Lie bracket of \mathfrak{g} and let $\nu = \lim_{s \to \infty} T_s . \mu$. By the choice of the d_a , it follows that $\nu(X_i, X_j) = 0$; moreover, as $d_a > 0$, we have $\nu(Y_k, Y_r) = 0$. Again, by the choice of d_a , it follows from that for the

bracket ν the matrices of $(\operatorname{ad}_{Y_k})_{|\mathfrak{n}}$ have the block-diagonal form, with the same diagonal blocks as that for the bracket μ .

It follows that the Lie algebra $\bar{\mathfrak{g}}$ defined on the linear space of \mathfrak{g} by the bracket ν is solvable, with the abelian nilradical \mathfrak{n} and with an abelian linear complement $\operatorname{Span}(Y_1, \ldots, Y_m)$ to \mathfrak{n} (note that the nilradical does not increase, as the complex eigenvalues of $(\operatorname{ad}_{Y_k})_{|\mathfrak{n}}$ do not change when we pass from μ to ν).

To finish the proof, by Proposition 1, it suffices to construct an inner product on $\bar{\mathfrak{g}}$ whose Ricci curvature is negative, provided that for some $Y \in \text{Span}(Y_1, \ldots, Y_m)$ all the eigenvalues of $(\text{ad}_Y)_{|\mathfrak{n}}$ (relative to ν) have positive real part. Without loss of generality, suppose that the latter condition is satisfied for $Y = Y_1$ and first choose an inner product $\langle \cdot, \cdot \rangle_0$ on $\bar{\mathfrak{g}}$ such that the basis \mathcal{B} is orthonormal. From the above, for the matrices defined by (3) we have $D_i = 0$, $B_j = 0$, and moreover, the matrices A_j are normal. Then by (6), $R_2 = 0$ and by (5) and (8), $R_1 = -tA_1^s$, which is negative definite by assumption. Now by Lemma 2 applied to the abelian subalgebra $\text{Span}(Y_1, \ldots, Y_m) \subset \mathfrak{gl}(\mathfrak{n})$, we can slightly perturb the inner product $\langle \cdot, \cdot \rangle_0$ on \mathfrak{n} only in such a way that no nontrivial linear combination of the matrices A_j is skew-symmetric. Then by (7), the matrix R_3 for the resulting inner product is negative definite. Moreover, R_2 is still zero, as we only change the inner product on \mathfrak{n} , and R_1 remains negative definite, if the perturbation is small enough.

4. Heisenberg Nilradical: proof of Theorem 3

Let \mathfrak{g} be a solvable Lie algebra whose nilradical \mathfrak{n} is the Heisenberg Lie algebra of dimension l = 2p + 1, $p \geq 1$ (see eg [RW] for more details of such algebras). Let $\mathcal{B} = \{X_1, \ldots, X_{2p}, Z\}$ be a basis for \mathfrak{n} such that $[X_i, X_{p+i}] = Z$ for $i = 1, \ldots, p$ (and $[X_i, X_j] = 0$ if $|i - j| \neq p$) and let Z span the centre \mathfrak{z} of \mathfrak{n} . Then for every $Y \in \mathfrak{g} \setminus \mathfrak{n}$ there is a well-defined number $\lambda(Y)$ such that $[Y, Z] = \lambda(Y)Z$ and a well-defined linear map $\Phi(Y) := \pi(\mathrm{ad}_Y)_{|\mathfrak{n}}\pi^{-1} \in \mathrm{End}(\mathfrak{n}/\mathfrak{z})$, where $\pi : \mathfrak{n} \to \mathfrak{n}/\mathfrak{z}$ is the natural projection (formally one may think that $\pi^{-1}(X)$ is any $\widetilde{X} \in \mathfrak{n}$ such that $\pi(\widetilde{X}) = X$ in the definition of Φ). Let $d_i(Y) \in \mathbb{C}$, $i = 1, \ldots, 2p$, be the eigenvalues of $\Phi(Y)$, each listed with its algebraic multiplicity. The following theorem is stated in the Introduction.

Theorem 3. A solvable Lie algebra \mathfrak{g} with the Heisenberg nilradical admits an inner product of negative Ricci curvature if and only if there exists $Y_+ \in \mathfrak{g}$ such that in the above notation,

(9)
$$\lambda(Y_{+}) + \sum_{i: \operatorname{Re} d_{i}(Y_{+}) < 0} \operatorname{Re} d_{i}(Y_{+}) > 0.$$

Remark 1. For every element $Y \in \mathfrak{g} \setminus \mathfrak{n}$, the matrix of the restriction A_Y of ad_Y to \mathfrak{n} relative to \mathcal{B} has the form $A_Y = \begin{pmatrix} N_Y & 0 \\ v_Y^t & \lambda(Y) \end{pmatrix}$, where λ is a one-form on \mathfrak{g} which vanishes on \mathfrak{n} , and $JN_Y + N_Y^t J = \lambda(Y)J$, where the $2p \times 2p$ matrix J is defined by $J = \begin{pmatrix} 0 & I_p \\ -I_p & 0 \end{pmatrix}$. Then $J^2 = -I_{2p}$ and it easily follows that $\operatorname{Tr} N_Y = p\lambda(Y)$. Moreover,

the matrices N_Y , $Y \in \mathfrak{g} \setminus \mathfrak{n}$, commute by the Jacobi identity, as for any $Y, Y' \in \mathfrak{g} \setminus \mathfrak{n}$ we have $[Y, Y'] \in \mathfrak{n}$, so $[A_Y, A_{Y'}] = \begin{pmatrix} 0_{2p \times 2p} & 0 \\ * & 0 \end{pmatrix}$. Note that N_Y is the matrix of $\Phi(Y)$ relative to the basis $\{X_1, \dots, X_{2p}\}$.

Proof. Sufficiency. Choose a basis Y_j , j = 1, ..., m, which complements \mathcal{B} to a basis for \mathfrak{g} in such a way that $Y_1 = Y_+$ (this is possible as $\lambda(Y_+) > 0$ by (9), so $Y_+ \notin \mathfrak{n}$) and $\lambda(Y_2) = \ldots = \lambda(Y_m) = 0$. Adding to Y_j 's appropriate linear combinations of the X_i 's we can always assume that $v_{Y_i} = 0$ (note that this does not violate condition (9) for Y_1). Then we have $[A_{Y_j}, A_{Y_k}] = 0$ (by Remark 1), so $[Y_j, Y_k] = \alpha_{jk}Z$, for all $j, k = 1, 2, \ldots, m$. But then the Jacobi identity on the triples (Y_1, Y_j, Y_k) , 1 < j < k, implies $\alpha_{jk} = 0$, so we get $[Y_1, Y_j] = a_j Z$ and $[Y_j, Y_k] = 0$ for $j, k \ge 2$.

To construct an inner product of negative Ricci curvature on \mathfrak{g} we use Proposition 1. On the first step, we eliminate the a_j 's as follows. For $s \in \mathbb{R}$, define $T_s \in GL(\mathfrak{g})$ by $T_sY_j = Y_j, \ T_sX_i = e^sX_i, \ T_sZ = e^{2s}Z$ for $j = 1, \ldots, m, \ i = 1, \ldots, 2p$. If μ is the Lie bracket of \mathfrak{g} and $\nu = \lim_{s \to \infty} T_s \cdot \mu$, then for the Lie algebra $\overline{\mathfrak{g}}$ of ν relative to the basis \mathcal{B} we have

(10)
$$[Y_j, Y_k] = 0, \quad (\mathrm{ad}_{Y_j})_{|\mathfrak{n}|} = \begin{pmatrix} N_j & 0\\ 0 & \lambda_j \end{pmatrix}, \quad [X_i, X_r] = j_{ir}Z,$$

for $i, r = 1, \ldots, 2p, j, k = 1, \ldots, m$, where the matrices N_j commute and satisfy the equation $JN_j + N_j^t J = \lambda_j J$ (and in particular, $\operatorname{Tr} N_j = p \lambda_j$), $J = (j_{ir})_1^{2p}$. From the latter equation it follows that the matrices $M_j = N_j - \frac{1}{2}\lambda_j I$ span an *m*-dimensional abelian subalgebra in $\mathfrak{sp}(2p,\mathbb{R})$. Denote S_j the semisimple part of M_j . Note that the matrices S_j commute, are linearly independent (otherwise the nilradical of \mathfrak{g} would be bigger) and the maps $\begin{pmatrix} S_j + \frac{1}{2}\lambda_j I & 0\\ 0 & \lambda_j \end{pmatrix}$ (relative to \mathcal{B}) are still derivations of \mathfrak{n} (see e. g. [Che, §14]). From [Ric, Proposition 11.14] it follows that the closure of the orbit of $\operatorname{Span}(M_i)$ under the simultaneous adjoint action of the group $\operatorname{Sp}(2p,\mathbb{R})$ contains the abelian subalgebra $\text{Span}(S_i)$. Let $h_N \in \text{Sp}(2p, \mathbb{R}), N \in \mathbb{N}$, be a sequence of matrices such that $\lim_{N\to\infty} h_N^{-1} M_j h_N = S_j$, for all $j = 1, \ldots, m$, and define the operators H_N by $H_N Y_i = Y_i, \ H_N Z = Z, \ H_N X_i = h_N X_i.$ As the group $\operatorname{Sp}(2p, \mathbb{R})$ acts by automorphisms on the Heisenberg Lie algebra \mathfrak{n} we obtain that the Lie bracket $\rho = \lim_{N \to \infty} H_N \nu$ is given by (10), with the N_j 's being replaced by their semisimple parts $S_j + \frac{1}{2}\lambda_j I$. By Proposition 1, it is sufficient to find an inner product for which the Ricci curvature of the Lie algebra with the bracket ρ is negative. To simplify the notation, in the remaining part of the proof we will keep the notation μ (instead of ρ) for this new Lie bracket and \mathfrak{g} for the corresponding Lie algebra.

Recall that the vectors Y_i were chosen in such a way that $\lambda_1 > 0$, $\lambda_2 = \cdots = \lambda_m = 0$. The following lemma shows that it suffices to construct an inner product of negative Ricci curvature for the one-dimensional extension of \mathfrak{n} by Y_1 .

Lemma 3. Let $\langle \cdot, \cdot \rangle$ be an inner product on the subalgebra $\mathfrak{g}_1 = \mathbb{R}Y_1 \oplus \mathfrak{n}$ such that $Y_1 \perp \mathfrak{n}$ and $Z \perp \mathfrak{m} := \operatorname{Span}(X_1, \ldots, X_{2p})$. In the notation of Section 2.1 choose an orthonormal basis $\{e_1, \ldots, e_{2p}, e_l, f_1\}$ for $(\mathfrak{g}_1, \langle \cdot, \cdot \rangle)$ in such a way that $f_1 \parallel Y_1$ and $e_l \parallel Z$. Suppose that the matrix R_1 defined by (5) for $(\mathfrak{g}_1, \langle \cdot, \cdot \rangle)$ relative to the basis $\{e_1, \ldots, e_{2p}, e_l, f_1\}$ is negative definite. Then \mathfrak{g} admits an inner product of negative Ricci curvature.

Proof. Extend the inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{g}_1 to the inner product $\langle \cdot, \cdot \rangle_{\varepsilon}$ on \mathfrak{g} defined by the ortonormal basis $\{e_1, \ldots, e_{2p}, e_l, f_1, f_2 = \varepsilon Y_2, \ldots, f_m = \varepsilon Y_m\}$, where $\varepsilon > 0$ (so that, in particular, the restriction of $\langle \cdot, \cdot \rangle_{\varepsilon}$ to \mathfrak{g}_1 coincides with $\langle \cdot, \cdot \rangle$). As $[Y_i, Y_j] = 0$, we have $B_i = 0$ in the notation of (3). Moreover, $D_l = 0$ and for i < l, the matrix D_i defined by (3) may only have nonzero entries in its bottom row, while the bottom row of C_i is zero, as $[Y_i, \mathfrak{m}] \subset \mathfrak{m}$. It follows that $R_2 = 0$. Furthermore, as the inner product $\langle \cdot, \cdot \rangle_{\varepsilon}$ coincides with $\langle \cdot, \cdot \rangle$ on \mathfrak{g}_1 , it follows from (5) and the assumption that the matrix R_1 for $(\mathfrak{g}, \langle \cdot, \cdot \rangle_{\varepsilon})$ can be made negative definite by choosing ε small enough. Finally, by (7), we have $(R_3)_{jk} = -\operatorname{Tr}(A_j^s A_k^s)$. This matrix is negative semidefinite; it has a zero eigenvalue if and only if the matrices A_i^s are linearly dependent, that is, if and only if a nontrivial linear combination of the A_i 's is skew-symmetric. By Lemma 2 applied to the restrictions of the A_j to \mathfrak{m} (which span an abelian subalgebra of $\mathfrak{gl}(\mathfrak{m})$), we can slightly perturb the inner product $\langle \cdot, \cdot \rangle_{\varepsilon}$ on \mathfrak{m} only, so that R_3 will become negative definite. Note that R_1 will remain negative definite, if the perturbation is small enough and R_2 will remain zero, since we are only changing the inner product on \mathfrak{m} .

To finish the proof it remains to construct an inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{g}_1 such that $Y_1 \perp \mathfrak{n}, Z \perp \mathfrak{m}$ and R_1 is negative definite. Note that for $Y_1 \in \mathfrak{g}_1$, the eigenvalues of $(\operatorname{ad}_{Y_1})_{|\mathfrak{n}}$ are the same as those for $Y_1 \in \mathfrak{g}$. From now on we omit the subscript 1 in Y_1, N_1, S_1 and λ_1 . According to (10) we have $(\operatorname{ad}_Y)_{|\mathfrak{m}} = N$, $[Y, Z] = \lambda Z$, with $\operatorname{Tr} N = p\lambda$ and the matrix $S = N - \frac{1}{2}\lambda I$ being Hamiltonian and semisimple. By [LM, Section 3], there exists a basis \mathcal{B}' for \mathfrak{m} relative to which the matrices S and J simultaneously have a (canonical) block-diagonal form $S = \operatorname{diag}(G_1, \ldots, G_r), J = \operatorname{diag}(J_{2q_1}, \ldots, J_{2q_r})$, where $q_i \in \{1, 2\}, J_q = \begin{pmatrix} 0 & I_q \\ -I_q & 0 \end{pmatrix}$ and the diagonal blocks G_i have one of the following forms $G_i^{(1)}, G_i^{(2)}, G_i^{(3)}$:

(11)
$$\begin{array}{c} G_i^{(1)} = \begin{pmatrix} 0 & \nu_i \\ -\nu_i & 0 \end{pmatrix} \text{ or } \\ G_i^{(2)} = \begin{pmatrix} \mu_i & 0 \\ 0 & -\mu_i \end{pmatrix} \text{ if } q_i = 1, \qquad G_i^{(3)} = \begin{pmatrix} \mu_i & \nu_i & 0 & 0 \\ -\nu_i & \mu_i & 0 & 0 \\ 0 & 0 & -\mu_i & -\nu_i \\ 0 & 0 & \nu_i & -\mu_i \end{pmatrix} \text{ if } q_i = 2,$$

where $\mu_i \geq 0$, $\nu_i \neq 0$. Then the matrix N has the same block-diagonal decomposition, with the diagonal blocks $G_i + \frac{1}{2}\lambda I_{2q_i}$. The eigenvalues of N are respectively $\frac{1}{2}\lambda \pm \nu_i$ or $\pm \mu_i + \frac{1}{2}\lambda$, for every 2×2 block $G_i + \frac{1}{2}\lambda I_2$ and $\frac{1}{2}\lambda \pm \mu_i \pm \nu_i$, for every 4×4 block $G_i + \frac{1}{2}\lambda I_2$.

Introduce an inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{g}_1 as follows. In the notation of Section 2.1, set $f_1 = Y$ and $e_m = \xi Z$, $\xi \neq 0$. Furthermore, define the inner product on \mathfrak{m} so

that its matrix Q relative to the basis \mathcal{B}' has the block-diagonal decomposition $Q = \text{diag}(Q_1, \ldots, Q_r)$ which agrees with that for S and J, where $Q_i = a_i I_{2q_i}$, $a_i > 0$ (so Q is diagonal and the basis \mathcal{B}' is orthogonal). We now define e_i , $i \leq 2p$, to be proportional to the elements of \mathcal{B}' with the corresponding coefficients (so that e_i are unit) and compute R_1 according to (5). Note that relative to the chosen orthonormal basis the matrix A is normal, its symmetric part is diagonal and $t = \text{Tr } A = (p+1)\lambda$. A direct computation shows that the matrix R_1 is diagonal, with

(12)
$$(R_1)_{mm} = \frac{1}{2}\xi^{-2}\sum_{i=1}^r q_i a_i^{-2} - (p+1)\lambda^2,$$

and with the $(2p) \times (2p)$ -submatrix \bar{R} in the top left-hand corner having the blockdiagonal decomposition $\bar{R} = \text{diag}(\bar{R}_1, \ldots, \bar{R}_r)$ which agrees with that for S and J, where \bar{R}_i is the $(2q_i) \times (2q_i)$ matrix of the forms

$$\begin{split} \bar{R}_i^{(1)} &= \left(-\frac{1}{2}a_i^{-2}\xi^{-2} - \frac{1}{2}(p+1)\lambda^2\right)I_2,\\ \bar{R}_i^{(2)} &= \operatorname{diag}\left(-\frac{1}{2}a_i^{-2}\xi^{-2} - (p+1)\lambda(\frac{1}{2}\lambda + \mu_i), -\frac{1}{2}a_i^{-2}\xi^{-2} - (p+1)\lambda(\frac{1}{2}\lambda - \mu_i)\right),\\ \bar{R}_i^{(3)} &= \operatorname{diag}\left(\left(-\frac{1}{2}a_i^{-2}\xi^{-2} - (p+1)\lambda(\frac{1}{2}\lambda + \mu_i)\right)I_2, \left(-\frac{1}{2}a_i^{-2}\xi^{-2} - (p+1)\lambda(\frac{1}{2}\lambda - \mu_i)\right)I_2\right), \end{split}$$

for the corresponding blocks in (11). It follows that $\bar{R}_i^{(1)}$ is always negative definite. Moreover, if $\mu_i \leq \frac{1}{2}\lambda$, then $\bar{R}_i^{(2)}$ and $\bar{R}_i^{(3)}$ are also negative definite (recall that $\mu_i \geq 0$). If $\mu_i > \frac{1}{2}\lambda$, then for $\bar{R}_i^{(2)}$ (respectively, $\bar{R}_i^{(3)}$) to be negative definite, we have to choose a_i so that $a_i^{-2}\xi^{-2} > -2(p+1)\lambda(\frac{1}{2}\lambda-\mu_i)$. Finally, according to (12), for $(R_1)_{mm}$ to be negative, we can choose $a_i^{-2} > 0$ to be arbitrarily small for those *i*'s which correspond to the blocks $\bar{R}_i^{(1)}$ and to the blocks $\bar{R}_i^{(2)}$, $\bar{R}_i^{(3)}$ with $\mu_i \leq \frac{1}{2}\lambda$. Using the above inequalities for the remaining *i*'s we obtain that for $(R_1)_{mm} < 0$ it is sufficient that $\lambda + \sum_{i:\mu_i > \frac{1}{2}\lambda} q_i(\frac{1}{2}\lambda-\mu_i) > 0$. But every $\frac{1}{2}\lambda - \mu_i$ is the real part of an eigenvalue of the restriction of ad_Y to \mathfrak{m} , with the corresponding multiplicity q_i , so that the latter inequality is equivalent to (9).

Necessity. Let \mathfrak{g} be a solvable Lie algebra whose nilradical \mathfrak{n} is the Heisenberg Lie algebra of dimension l = 2p + 1. Suppose \mathfrak{g} admits an inner product $\langle \cdot, \cdot \rangle$ of negative Ricci curvature. Choose an orthonormal basis $\{e_i, f_j\}$ for \mathfrak{g} as in Section 2.1. We will prove that the vector f_1 satisfies inequality (9). The proof only uses the fact that the matrix R_1 defined by (5) is negative definite.

Specify the basis $\{e_i\}$ further, so that e_{2p+1} spans the centre of \mathfrak{n} and let $[f_1, e_{2p+1}] = \lambda e_{2p+1}$ (without loss of generality we may assume that $\lambda > 0$). Note that $t = \operatorname{Tr} \operatorname{ad}_{f_1} = (p+1)\lambda$ (> 0). Denote $\mathfrak{m} = \mathfrak{n} \cap e_{2p+1}^{\perp}$, and introduce a skew-symmetric operator $K \in \operatorname{End}(\mathfrak{m})$ by $[X_1, X_2] = \langle KX_1, X_2 \rangle e_{2p+1}$ for $X_1, X_2 \in \mathfrak{m}$.

Relative to the basis e_i for \mathfrak{n} the matrices A_j has the form $A_j = \begin{pmatrix} N_j & 0 \\ v_j^t & \lambda_j \end{pmatrix}$, where $v_j \in \mathbb{R}^{2p}$, $\lambda_1 = \lambda$ and $\lambda_j = 0$ for j > 1 (Remark 1). For every eigenvalue $d \in \mathbb{C}$ of the operator N_1 acting on $\mathfrak{m}^{\mathbb{C}}$ let $V_d \subset \mathfrak{m}^{\mathbb{C}}$ be its root subspace, so that $V_d = \bigcup_{k=1}^{\infty} \operatorname{Ker}(N_1 - d\operatorname{id}_{\mathfrak{m}})^k$.

Clearly $V_{\bar{d}} = \overline{V_d}$ and $\mathfrak{m}^{\mathbb{C}} = \bigoplus_d V_d$. Let $V_- = \bigoplus_{d:\operatorname{Re}(d)<0} V_d$ and denote $\mathfrak{m}_- = V_- \cap \mathfrak{m}$ and $m_- = \dim \mathfrak{m}_-$. Note that m_- is the number of the eigenvalues of N with negative real part, counted with their algebraic multiplicity. Denote π_- the orthogonal projection to \mathfrak{m}_- (both from \mathfrak{m} and from \mathfrak{n} – the meaning will be clear from the context).

We want to compute $Tr(R_1\pi_-)$. To estimate it we use the following Lemma.

Lemma 4.

- (a) $[\mathfrak{m}_{-},\mathfrak{m}_{-}]=0$, so that $\langle KX_{1},X_{2}\rangle=0$ for $X_{1},X_{2}\in\mathfrak{m}_{-}$.
- (b) $\operatorname{Tr}(K^t K \pi_-) \leq \frac{1}{2} \operatorname{Tr}(K^t K).$
- (c) The subspace \mathfrak{m}_{-} is an invariant subspace of all the N_{j} .
- (d) $\operatorname{Tr}([A_i, A_i^t]\pi_-) \ge -\|\pi_- v_i\|^2$, for all $j = 1, \dots, m$.
- (e) $\operatorname{Tr}(A_1^s \pi_{-}) = \sum_{d:\operatorname{Re} d < 0} d = \sum_{d:\operatorname{Re} d < 0} \operatorname{Re} d$, where the sum is taken by all the eigenvalues d of N_1 , counting their algebraic multiplicity.

Proof. (a) The operator $A_1 = \begin{pmatrix} N_1 & 0 \\ v_1^t & \lambda \end{pmatrix}$ is a derivation of \mathfrak{n} . Adding an appropriate

 $\operatorname{ad}_X, X \in \mathfrak{n}$, we can eliminate v_1 , so that the operator $C = \begin{pmatrix} N_1 & 0 \\ 0 & \lambda \end{pmatrix}$ is again a derivation

of **n**. Then C is a derivation of the complexified algebra $\mathbf{n}^{\mathbb{C}}$. Our arguments are similar to [Hel, Lemma III.3.2]. Let $X_1 \in V_{d_1}$, $X_2 \in V_{d_2}$, $\operatorname{Re} d_i < 0$. Then $(N_1 - d_1 \operatorname{id})^{k_1} X_1 = (N_1 - d_2 \operatorname{id})^{k_2} X_2 = 0$, for some $k_1, k_2 \in \mathbb{N}$, so $(C - d_i \operatorname{id})^{k_i} X_i = 0$, i = 1, 2. As C is a derivation we obtain by induction that $(C - (d_1 + d_2)\operatorname{id})^k [X_1, X_2] = \sum_{i=0}^k {k \choose i} [(C - d_1 \operatorname{id})^i X_1, (C - d_2 \operatorname{id})^{k-i} X_2]$. The right-hand side vanishes for k large enough, while the left-hand side equals $(\lambda - (d_1 + d_2))^k [X_1, X_2]$, as $[X_1, X_2]$ is a multiple of e_{2p+1} . But $\lambda > 0$ and $\operatorname{Re} d_1, \operatorname{Re} d_2 < 0$, so $[X_1, X_2] = 0$. It follows that $[V_-, V_-] = 0$, hence $[\mathfrak{m}_-, \mathfrak{m}_-] = 0$.

Then for $X_1, X_2 \in \mathfrak{m}_-$ we have $\langle KX_1, X_2 \rangle = \langle [X_1, X_2], e_{2p+1} \rangle = 0.$

(b) Let $\{e_i\}_{i=1}^{2p}$ be an orthonormal basis for \mathfrak{m} such that $\{e_i\}_{i=1}^{m_-}$ is an orthonormal basis for \mathfrak{m}_- . We have $\langle Ke_i, e_j \rangle = 0$ for $i, j \leq m_-$ by (a). Then

$$\operatorname{Tr}(K^{t}K) = \sum_{i,j=1}^{2p} \langle Ke_{i}, e_{j} \rangle^{2} = \sum_{i,j=m_{-}+1}^{2p} \langle Ke_{i}, e_{j} \rangle^{2} + 2 \sum_{i \leq m_{-} < j} \langle Ke_{i}, e_{j} \rangle^{2},$$
$$\operatorname{Tr}(K^{t}K\pi_{-}) = \sum_{i=1}^{m_{-}} \langle K^{t}Ke_{i}, e_{i} \rangle = \sum_{i=1}^{m_{-}} \sum_{j=1}^{2p} \langle Ke_{i}, e_{j} \rangle^{2} = \sum_{i \leq m_{-} < j} \langle Ke_{i}, e_{j} \rangle^{2},$$

and the claim follows.

(c) As N_j commutes with N_1 , for any $j = 1, \ldots, m$ (Remark 1), every root subspace $V_d \subset \mathfrak{m}^{\mathbb{C}}$ of N_1 is N_j -invariant. Then $V_- \subset \mathfrak{m}^{\mathbb{C}}$ is also N_j -invariant, as is \mathfrak{m}_- .

(d) Choose an orthonormal basis $\{e_i\}_{i=1}^{2p}$ for \mathfrak{m} as in the proof of (b) above and extend it to the basis for \mathfrak{n} by the vector e_{2p+1} . We have

$$\operatorname{Tr}([A_j, A_j^t]\pi_{-}) = \sum_{i=1}^{m_{-}} \langle [A_j, A_j^t] e_i, e_i \rangle = \sum_{i=1}^{m_{-}} (\|A_j^t e_i\|^2 - \|A_j e_i\|^2)$$
$$= \sum_{i=1}^{m_{-}} \sum_{s=1}^{2p+1} \langle A_j e_s, e_i \rangle^2 - \sum_{i=1}^{m_{-}} \sum_{s=1}^{2p+1} \langle A_j e_i, e_s \rangle^2.$$

But the first sum equals $\sum_{i=1}^{m_-} \sum_{s=1}^{2p} \langle A_j e_s, e_i \rangle^2$ (as e_{2p+1} is an eigenvector of A_j) and the second one, $\sum_{i,s=1}^{m_-} \langle A_j e_i, e_s \rangle^2 + \sum_{i=1}^{m_-} \langle A_j e_i, e_{2p+1} \rangle^2 = \sum_{i,s=1}^{m_-} \langle A_j e_s, e_i \rangle^2 + \sum_{i=1}^{m_-} \langle v_j, e_i \rangle^2$, by (c), so the claim follows.

(e) With the same choice of basis as in the proof of (d) above we have $\operatorname{Tr}(A_1^s \pi_-) = \sum_{i=1}^{m_-} \langle A_1 e_i, e_i \rangle = \sum_{i=1}^{m_-} \langle N_1 e_i, e_i \rangle = \operatorname{Tr}(N_{1|\mathfrak{m}_-})$. Extending N_1 to $\mathfrak{m}^{\mathbb{C}}$ we get $\operatorname{Tr}(N_{1|\mathfrak{m}_-}) = \operatorname{Tr}(N_{1|V_-}) = \sum_{d:\operatorname{Re} d < 0} \operatorname{Tr}(N_{1|V_d}) = \sum_{d:\operatorname{Re} d < 0} d \dim V_d$, where the sum is taken by all the eigenvalues d of N_1 without counting the multiplicity.

Choosing an orthonormal basis $\{e_i\}$ for \mathfrak{n} as in the proof of Lemma 4 (so that $e_1, \ldots e_{m_-}$ is a basis for \mathfrak{m}_-) we obtain by (8) $\operatorname{Tr}(\operatorname{Ric}^{\mathfrak{n}} \pi_-) = -\frac{1}{2} \sum_{s=1}^{2p+1} \sum_{i=1}^{m_-} ||[e_s, e_i]||^2 = -\frac{1}{2} \sum_{i=1}^{m_-} ||Ke_i||^2 = -\frac{1}{2} \operatorname{Tr}(K^t K \pi_-).$

Then using Lemma 4 we get from (5)

(13)
$$\operatorname{Tr}(R_1\pi_-) \ge -\frac{1}{4}\operatorname{Tr}(K^tK) - \frac{1}{2}\sum_{j=1}^m \|\pi_-v_j\|^2 - t\sum_{d:\operatorname{Re}d<0}\operatorname{Re}d$$

where the sum is taken by all the eigenvalues d of N_1 , counting their algebraic multiplicity.

On the other hand, from (5) and (8) (and using the fact that e_{2p+1} is a common eigenvector of all the A_i) we have

$$\langle R_1 e_{2p+1}, e_{2p+1} \rangle \geq \frac{1}{4} \sum_{i,s=1}^{2p} \langle [e_i, e_s], e_{2p+1} \rangle^2 + \frac{1}{2} \sum_{j=1}^m (\|A_j^t e_{2p+1}\|^2 - \|A_j e_{2p+1}\|^2) - t\lambda$$

$$= \frac{1}{4} \sum_{i,s=1}^{2p} \langle K e_i, e_s \rangle^2 + \frac{1}{2} \sum_{j=1}^m \sum_{i=1}^{2p} \langle A_j^t e_{2p+1}, e_i \rangle^2 - t\lambda$$

$$= \frac{1}{4} \operatorname{Tr}(K^t K) + \frac{1}{2} \sum_{j=1}^m \|v_j\|^2 - t\lambda.$$

Adding this to (13) and using the fact that π_{-} is positive semidefinite and R_{1} is negative definite we get $\lambda + \sum_{d:\text{Re } d < 0} \text{Re } d > 0$, which is equivalent to (9).

5. FILIFORM NILRADICAL: PROOF OF THEOREM 4

A nilpotent Lie algebra of dimension l is called *filiform* if it has the maximal possible degree of nilpotency (\Leftrightarrow the longest possible lower central series). Filiform Lie algebras have been introduced in [Ver] and have been given a great deal of attention thereafter.

In this section, we consider solvable Lie algebras \mathfrak{g} whose nilradical is the so called standard filiform Lie algebra. The latter is defined as the *l*-dimensional Lie algebra L_l having a basis X_1, \ldots, X_l such that $[X_1, X_i] = X_{i+1}$, $i = 2, \ldots, l-1$, $[X_1, X_l] = 0$, and $[X_i, X_j] = 0$ when $i, j \geq 2$. On the question of how restrictive the assumption of standardness is, note that any filiform Lie algebra admits a basis for which the former relations are satisfied (but in general, not the latter ones), and that any filiform algebra of dimension l degenerates to L_l . Note also that a "typical" filiform Lie algebra of dimension $l \geq 8$ is characteristically nilpotent, hence cannot be the nilradical of anything except for itself (any solvable extension of it is nilpotent).

Let \mathfrak{g} be a solvable Lie algebra with the nilradical L_l . We can assume that $l \geq 4$ (as L_2 is abelian and L_3 is the Heisenberg algebra). The algebra L_l has a (unique) codimension

one abelian ideal $\mathfrak{i} = \operatorname{Span}(X_2, \ldots, X_l)$ and the one-dimensional centre $\mathbb{R}X_l$. Both of them are characteristic ideals of L_l (they are invariant under the action of any derivation on L_l ; see Remark 2 below). Let λ and ι be one-forms on \mathfrak{g} defined as follows: for $Y \in \mathfrak{g}$, $[Y, X_l] = \lambda(Y)X_l$ and $\iota(Y) = \operatorname{Tr}((\operatorname{ad}_Y)_{|\mathfrak{i}})$. The following theorem is stated in the Introduction.

Theorem 4. Let \mathfrak{g} be a solvable Lie algebra with the nilradical $\mathfrak{n} = L_l$, $l \geq 4$. The algebra \mathfrak{g} admits an inner product of negative Ricci curvature if and only if there exists $Y \in \mathfrak{g}$ such that $\lambda(Y) > 0$ and $\iota(Y) > 0$.

Remark 2. It is easy to see (and is well-known) that relative to the basis $\mathcal{B} = \{X_i\}$ for L_l , any derivation of L_l has the lower-triangular form

(14)
$$\begin{pmatrix} a & & 0 \\ & d & & 0 \\ & & a+d & & \\ & & & 2a+d & \\ & & & \ddots & \\ & & & & (l-2)a+d \end{pmatrix}$$

where $a, d \in \mathbb{R}$ (with some additional relations on the entries below the diagonal). It follows that any solvable extension of L_l of rank $m \geq 3$ contains a nilpotent derivation $\operatorname{ad}_Y, Y \notin L_l$, so that its nilradical is bigger than L_l . Therefore m = 1 or m = 2. But if m = 2, there exists $Y \in \mathfrak{g}$ such that $(\operatorname{ad}_Y)_{|\mathfrak{n}}$ has the form (14), with a, d > 0. Then the inequalities $\lambda(Y) > 0$, $\iota(Y) > 0$ are trivially satisfied for such Y and the existence of an inner product of negative Ricci curvature follows from Theorem 2(2).

We can therefore assume that m = 1. Then for an arbitrary $Y \notin \mathfrak{n}$, the matrix $(\operatorname{ad}_Y)_{|\mathfrak{n}}$ has the form (14), with a, d not simultaneously zero, and the inequalities $\lambda(Y) > 0$ and $\iota(Y) > 0$ are equivalent to

(15)
$$(l-2)a + d > 0, \quad (l-2)a + 2d > 0,$$

respectively.

Proof. By Remark 2 we can assume that \mathfrak{g} is a one-dimensional extension of L_l . Choose and fix a vector $Y \in \mathfrak{g} \setminus \mathfrak{n}$. Then $(\mathrm{ad}_Y)_{|\mathfrak{n}}$ has the form (14), and $\mathrm{Tr} \, \mathrm{ad}_Y = (l-1)d + (\frac{1}{2}(l-1)(l-2)+1)a = \frac{2}{l-2}((l-2)a+d) + \frac{l(l-3)}{2(l-2)}((l-2)a+2d)$. If $\mathrm{Tr} \, \mathrm{ad}_Y = 0$, then (15) is violated and no inner product with negative Ricci curvature exists, as \mathfrak{g} is unimodular (Theorem 1). We can therefore assume that $t = \mathrm{Tr} \, \mathrm{ad}_Y > 0$.

Sufficiency. Suppose the inequalities (15) are satisfied. Let N be a positive derivation of \mathfrak{n} which is diagonal relative to \mathcal{B} , say $N = \operatorname{diag}(1, 2, \ldots, l)$. For s > 0 define $T_s \in$ End(\mathfrak{g}) by $T_sY = Y$ and $T_sX = e^{sN}X$, for $X \in \mathfrak{n}$. When $s \to \infty$, the Lie algebra \mathfrak{g} degenerates to the Lie algebra \overline{g} with the same nilradical $\mathfrak{n} = L_l$ and with $(\operatorname{ad}_Y)_{|\mathfrak{n}} =$ $\operatorname{diag}(a, d, a + d, 2a + d, \ldots, (l-2)a + d)$ (relative to \mathcal{B}). By Proposition 1 it suffices to construct an inner product of negative Ricci curvature on $\overline{\mathfrak{g}}$. In the notation of Section 2.1 take $f_1 = Y$ and $e_i = a_i X_i$, $a_i > 0$, for i = 1, ..., l. An easy calculation shows that $R_2 = 0$ and $R_3 < 0$. Furthermore, from (5) we obtain that the matrix R_1 is diagonal, with the entries

$$(R_1)_{11} = -\sum_{i+2}^{t-1} \xi_i - ta, \quad (R_1)_{22} = -\xi_2 - td, \quad (R_1)_{ll} = \xi_{l-1} - t((l-2)a + d)),$$

$$(R_1)_{ii} = \xi_{i-1} - \xi_i - t((i-1)a + d), \text{ for } i = 3, \dots, l-1,$$

where $\xi_i = \frac{1}{2}a_1^2a_i^2a_{i+1}^{-2}$, $i = 2, \ldots, l-1$, and $t = \text{Tr } A = (l-1)d + (\frac{1}{2}(l-1)(l-2)+1)a > 0$. In a Euclidean space \mathbb{R}^l , with an orthonormal basis E_i , introduce the vectors $F_1 = -E_1 - E_2 + E_3$, $F_2 = -E_1 - E_3 + E_4, \ldots, F_{l-2} = -E_1 - E_{l-1} + E_l$ and the vectors $V_1 = E_1 + \sum_{i=3}^l (i-2)E_i$, $V_2 = \sum_{i=2}^l E_i$. Then the fact that R_1 is negative definite is equivalent to the fact that all the components of the vector $\sum_{i=1}^{l-2} \xi_i F_i - taV_1 - tdV_2$ are negative, that is, to the fact that the vector $aV_1 + dV_2 - t^{-1}\sum_{i=1}^{l-2}\xi_iF_i$ belongs to the first octant of \mathbb{R}^n . Note that given any $\xi_i > 0$ we can easily find the corresponding a_i 's by taking $a_1, a_2 > 0$ arbitrarily and defining $a_{i+1} = a_1a_i(2\xi_i)^{1/2}$ for $i = 2, \ldots, l-1$, so for the existence of an inner product such that R_1 is negative definite it is sufficient that $aV_1 + dV_2$ belongs to the open convex hull of the vectors $F_1, F_2, \ldots, F_{l-2}, E_1, E_2, \ldots, E_l$. But $E_l = F_{l-2} + E_{l-1} + E_{l-2}$, and then $E_{l-1} = F_{l-3} + E_{l-2} + E_{l-3}$, and so on, up to $E_3 = F_1 + E_2 + E_1$, so that the convex hull of the vectors $F_1, F_2, \ldots, F_{l-2}, E_1, E_2, \ldots, E_l$ is the same as the convex hull of the vectors $F_1, F_2, \ldots, F_{l-2}, E_1, E_2, \ldots, E_l$

$$V_{1} = \sum_{i=1}^{l-2} \left(\sum_{j=i}^{l-2} j \right) F_{i} + \left(\frac{1}{6} (l-2)(l-1)(2l-3) + 1 \right) E_{1} + \frac{1}{2} (l-2)(l-1) E_{2},$$

$$V_{2} = \sum_{i=1}^{l-2} (l-1-i) F_{i} + \frac{1}{2} (l-2)(l-1) E_{1} + (l-1) E_{2},$$

so the vector $aV_1 + dV_2$ lies in the open convex hull of the vectors $F_1, F_2, \ldots, F_{l-2}, E_1, E_2$ if and only if the following inequalities are satisfied:

$$\left(\sum_{j=i}^{l-2} j\right)a + (l-1-i)d > 0, \quad i = 1, \dots, l-2,$$
$$\left(\frac{1}{6}(l-2)(l-1)(2l-3) + 1\right)a + \frac{1}{2}(l-2)(l-1)d > 0, \quad \frac{1}{2}(l-2)(l-1)a + (l-1)d > 0.$$

These inequalities are equivalent to the inequalities $\kappa_i a + d > 0$, where $\kappa_i > 0$ are defined by $\kappa_i = (\sum_{j=i}^{l-2} j)/(l-1-i)$ for $i = 1, \ldots, l-2$, and $\kappa_{l-1} = (\frac{1}{6}(l-2)(l-1)(2l-3)+1)(\frac{1}{2}(l-2)(l-1))^{-1}$, $\kappa_l = \frac{1}{2}(l-2)$, and hence are equivalent to the two inequalities $\kappa_{\max} a + d > 0$, $\kappa_{\min} a + d > 0$, where $\kappa_{\max} = \max \kappa_i$, $\kappa_{\min} = \min \kappa_i$. As κ_i increase when $i = 1, \ldots, l-2$ and $\kappa_l < \kappa_1$, and moreover, $\kappa_l < \kappa_{l-1} < \kappa_{l-2}$, we get $\kappa_{\max} = \kappa_{l-2} = l-2$, $\kappa_{\min} = \kappa_l = \frac{1}{2}(l-2)$. This gives (l-2)a + d > 0, (l-2)a + 2d > 0, which is satisfied by the assumption (see (15), Remark 2).

Necessity. Suppose $\langle \cdot, \cdot \rangle$ is an inner product of negative Ricci curvature on \mathfrak{g} . Take f_1 to be the unit vector orthogonal to \mathfrak{n} with $\operatorname{Tr} \operatorname{ad}_{f_1} > 0$. The inequality $\lambda(f_1) > 0$ follows from Theorem 2(1). To prove the inequality $\iota(f_1) > 0$ introduce an orthonormal

basis e_i for \mathfrak{n} in such a way that $e_1 \perp \mathfrak{i}$. Note that $[e_i, e_j] = 0$ for i, j > 1, as \mathfrak{i} is abelian, and that $[e_1, \mathfrak{i}] \subset \mathfrak{i}$, as \mathfrak{i} is an ideal. Then by (8) we obtain

$$\sum_{j=2}^{l} \langle \operatorname{Ric}^{\mathfrak{n}} e_{j}, e_{j} \rangle = -\frac{1}{2} \sum_{j=2}^{l} \sum_{i=1}^{l} ||[e_{i}, e_{j}]||^{2} + \frac{1}{4} \sum_{j=2}^{l} \sum_{i,k=1}^{l} \langle [e_{i}, e_{k}], e_{j} \rangle^{2}$$
$$= -\frac{1}{2} \sum_{j=2}^{l} ||[e_{1}, e_{j}]||^{2} + \frac{1}{2} \sum_{j,k=2}^{l} \langle [e_{1}, e_{k}], e_{j} \rangle^{2}$$
$$= 0.$$

It now follows from (5) that

$$\sum_{j=2}^{l} \langle R_1 e_j, e_j \rangle = \sum_{j=2}^{l} \langle (\frac{1}{2} [A_1, A_1^t] - tA_1^s) e_j, e_j \rangle$$

= $\operatorname{Tr}(\frac{1}{2} [A_1, A_1^t] - tA_1^s) - \langle (\frac{1}{2} [A_1, A_1^t] - tA_1^s) e_1, e_1 \rangle$
= $-t(\operatorname{Tr} A_1 - \langle A_1 e_1, e_1 \rangle) - \frac{1}{2} (\|A_1^t e_1\|^2 - \|A_1 e_1\|^2).$

The matrix A_1 relative to the basis e_i is similar to the matrix (14), with some $a, d \in \mathbb{R}$ such that $t = \text{Tr}(A_1) > 0$, and as $\mathfrak{i} = \text{Span}(X_2, \ldots, X_l) = \text{Span}(e_2, \ldots, e_l)$ we have

$$A_1 = \left(\begin{array}{cc} a & 0 \dots 0 \\ v & M \end{array}\right),$$

where $v \in \mathbb{R}^{l-1}$ and M is an $(l-1) \times (l-1)$ -matrix with the eigenvalues $d, a+d, 2a+d, \ldots, (l-2)a+d$. In particular, $\iota(f_1) = \operatorname{Tr} M = \operatorname{Tr} A_1 - \langle A_1e_1, e_1 \rangle$ and $||A_1^te_1||^2 - ||A_1e_1||^2 = -||v||^2$. Then from the above

$$\sum_{j=2}^{l} \langle R_1 e_j, e_j \rangle = -t\iota(f_1) + \frac{1}{2} ||v||^2,$$

so $\iota(f_1) > 0$, as required.

6. Open questions

In this section, motivated by the above theorems (and by the theory of Einstein solvmanifolds [Heb, LW]), we collect some open questions and conjectures.

- (1) Is it true that we can always reduce the rank? In other words, is the following true (or at least, one way true): "a solvable algebra \mathfrak{g} with the nilradical \mathfrak{n} admits an inner product with negative Ricci curvature if and only if there exists $Y \notin \mathfrak{n}$ such that the subalgebra $\mathbb{R}Y \oplus \mathfrak{n} \in \mathfrak{g}$ admits such an inner product"?
- (2) Are there some "best" solvable metric Lie algebras of negative Ricci curvature? More specifically: can we modify a given solvable Lie algebra with Ric < 0 by degeneration and then choose an inner product in such a way that the resulting algebra is nicer – e.g., the restrictions $(ad_Y)_{|\mathfrak{n}}$, $Y \in \mathfrak{a}$, are semisimple, the subspace \mathfrak{a} is abelian and \mathfrak{n} and α are invariant subspaces of Ric (so that in the notation of Section 2.1, $R_2 = 0$ and hence the negativity of Ric only depends on R_1)?

(3) A stronger form of the first question: is it true that a solvable Lie algebra \mathfrak{q} admits an inner product with Ric < 0 if and only if there exists a vector $Y \in \mathfrak{g}$ such that the real parts of the eigenvalues of the restriction of ad_{Y} to the nilradical \mathfrak{n} satisfy certain linear inequalities which are determined by the structure of \mathfrak{n} ? Such inequalities may represent the fact that $\operatorname{Re}(\operatorname{ad}_Y)_{|\mathfrak{n}}$ belongs to a certain open convex hull. An alternative description may be the following. For a linear operator A denote $A^{\mathbb{R}}$ its real semisimple part. If A is a derivation of a Lie algebra \mathfrak{n} , then $A^{\mathbb{R}}$ also is (see e.g. [Heb, Section 3.3]). Relative to some basis for \mathfrak{n} , $A^{\mathbb{R}}$ is diagonal. Consider the space \mathfrak{t} of all the diagonal matrices relative to that basis (more invariantly, the Cartan subalgebra of the centraliser of $A^{\mathbb{R}}$ in $\mathfrak{gl}(\mathfrak{n})$) and call A semipositive if $A^{\mathbb{R}}$ belongs to the orthogonal projection of the first octant of \mathfrak{t} (the cone of the matrices with all the diagonal entries positive) to $\mathfrak{t} \cap \operatorname{Der}(\mathfrak{n})$ via the restriction of the Killing form of $\mathfrak{gl}(\mathfrak{n})$ to \mathfrak{t} . Is it true then that a solvable Lie algebra \mathfrak{g} admits an inner product of negative Ricci curvature if and only if there exists a vector $Y \in \mathfrak{g}$ such that the restriction of ad_Y to the nilradical \mathfrak{n} is semipositive?

References

- [Ale] Alekseevskii D.V. Homogeneous Riemannian spaces of negative curvature, Mat. Sb. (N. S.), 96 (1975), 93 - 117 (in Russian). English translation in: Math. USSR-Sb., 25 (1976), 87 - 109.
- [AK] Alekseevskii D.V., Kimel'fel'd B. N. Structure of homogeneous Riemannian spaces with zero Ricci curvature, Functional Anal. Appl. 9 (1975), 297 – 339.
- [BB1] Bérard Bergery L. Les variétés riemanniennes homogènes simplement connexes de dimension impaire à courbure strictement positive, J. Math. Pures Appl. 55 (1976), 47 – 67.
- [BB2] Bérard Bergery L. Sur la courbure des métriques riemanniennes invariantes des groupes de Lie et des espaces homogènes, Ann. Sci. École Norm. Sup. (4) 11 (1978), 543 – 576.
- [Ber] Berestovskii V. N., Homogeneous Riemannian manifolds of positive Ricci curvature (Russian), Mat. Zametki 58 (1995), 334 – 340; English translation in: Math. Notes 58 (1995), 905 – 909.
- [Bur] Burde D. Degenerations of nilpotent Lie algebras, J. Lie Theory 9(1999), 193 202.
- [Che] Chevalley C. Théorie des groupes de Lie. Tome II. Groupes algébriques, Actualités Sci. Ind. 1152, (1951), Hermann & Cie., Paris.
- [DLM] Dotti Miatello I., Leite M. L., Miatello R. J. Negative Ricci curvature on complex simple Lie groups, Geom. Dedicata 17 (1984), 207 – 218.
- [DM] Dotti Miatello I. Ricci curvature of left invariant metrics on solvable unimodular Lie groups, Math. Z. 180 (1982), 257 – 263.
- [Gant] Gantmacher F. R. Applications of the theory of matrices, Interscience Publishers, Inc., New York; Interscience Publishers Ltd., London, 1959.
- [Heb] Heber J., Noncompact homogeneous Einstein spaces, Invent. Math., 133 (1998), 279 352.
- [Hei] Heintze E. On Homogeneous manifolds of negative curvature, Mat. Ann. **211** (1974), 23 34.
- [Hel] Helgason S. Differential geometry, Lie groups, and symmetric spaces, Pure and Applied Mathematics, 80. Academic Press, Inc. New York–London, 1978.
- [LDM] Leite M. L., Dotti Miatello I. Metrics of negative Ricci curvature on $SL(n, \mathbb{R}), n \ge 3$, J. Differential Geom. 17 (1982), 635 641.

- [LM] Laub A.J., Meyer K. Canonical forms for symplectic and Hamiltonian matrices, Celestial Mech., 9 (1974), 213 – 238.
- [LW] Lauret J., Will C., Einstein solvmanifolds: existence and non-existence questions, Math. Ann. 350 (2011), 199 – 225.
- [LiW] Li Michael Y., Wang Liancheng, A criterion for stability of matrices, J. Math. Anal. Appl. 225 (1998), 249 – 264.
- [Mil] Milnor J. Curvatures of left invariant metrics on Lie groups, Adv. Math. 21 (1976), 293 329.
- [NP] Nesterenko M., Popovych R. Contractions of low-dimensional Lie algebras, J. Math. Phys. 47 (2006), 123515, 45 pp.
- [New] Newman M. Two classical theorems on commuting matrices, J. Res. Nat. Bur. Standards Sect. B 71B (1967), 69 – 71.
- [NN] Nikitenko E.V., Nikonorov Yu.G. Six-dimensional Einstein solvmanifolds, Mat. tr., 8 (2005), 71 – 121 (in Russian). English translation in: Siberian Adv. Math., 16:1 (2006), 66 – 112.
- [Ric] Richardson R. W. Conjugacy classes of n-tuples in Lie algebras and algebraic groups, Duke Math. J. 57 (1988), 1 – 35.
- [RW] Rubin J. L., Winternitz P. Solvable Lie algebras with Heisenberg ideals, J. Phys. A 26 (1993), 1123 – 1138.
- [Ver] Vergne M. Cohomologie des algèbres de Lie nilpotentes. Application à l'étude de la variété des algèbres de Lie nilpotentes, Bull. Soc. Math. France 98 (1970), 81 – 116.
- [VGO] Vinberg E.B., Gorbatsevich V.V., Onishchik A.L. Lie Groups and Lie Algebras, III. Structure of Lie Groups and Lie Algebras, Encyclopedia of Math. Sciences V. 41, Springer-Verlag, Berlin, 1994; Translated from Russian: Structure of Lie Groups and Algebras, Itogi Nauki i Tekhniki (Sovrem. Probl. Mat., Fundam. Napravleniya), 41, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform. (VINITI), Moscow, 1990.
- [Wal] Wallach N. R. Compact homogeneous Riemannian manifolds with strictly positive curvature, Ann. of Math. (2) 96 (1972), 277 – 295.

Y. Nikolayevsky, Department of Mathematics and Statistics, La Trobe University, Melbourne, Australia $3086\,$

E-mail address: y.nikolayevsky@latrobe.edu.au

YU.G. NIKONOROV, SOUTH MATHEMATICAL INSTITUTE OF VSC RAS, 22 Markus St, Vladikavkaz, Russia362027

E-mail address: nikonorov2006@mail.ru