
E.A. Kolganova

Ultracold Scattering and Universal Correlations

Abstract Particle-dimer scattering below and above the three-body threshold is studied using Faddeev differential equations. Correlations between the observables are shown and some analogies between three-nucleon and three-atom systems are discussed.

Keywords Three-body systems · Faddeev equations · scattering and resonances

Ultracold scattering of different atoms and universal correlations have recently received considerable attention, in both experiment and theory. A remarkable linear correlation between the triton binding energy and the doublet nucleon - deuteron scattering length was first pointed out by Phillips [1], and is known as Phillips line. A similar approximate linear correlation between the binding energy of the helium trimer and the ^4He -dimer scattering length can be observed in the three-atomic system [2]. The Phillips line has not been explained for many years and only two decades later Efimov and Tkachenko [3] showed that the approximate linearity of the Phillips line is due to the fact that the interval covered by the calculations with various NN-potentials is not wide enough. Roudnev and Cavagnero in [2] suggested a modified Phillips line with a dimensionless variables

$$\alpha = \ell_{\text{sc}} \sqrt{-2\mu\epsilon_d/\hbar}, \quad \omega = 1/\sqrt{\frac{E_3}{\epsilon_d} - 1}, \quad (1)$$

where α is a dimensionless scattering length and ω is a dimensionless parameter characterizing the three-body binding energy; ϵ_d and E_3 define the energies of the two- and three-body bound states, respectively; μ is the reduced mass of the particle-dimer system, and ℓ_{sc} is the particle-dimer scattering length. Their idea to use such parametrization was based on the paper by Efimov and Tkachenko [3] who noted that for a weakly bound three-body state the approximate correlation $\epsilon_d - E_3 \approx \hbar^2/(2\mu\ell_{\text{sc}})$ should hold. The universal properties of systems with a large scattering length can well be described by an effective field theory. As shown in [4], a zero-range model formulated in field theoretical terms is able to simulate the scattering situation. Applications of the effective field theory to systems we deal with below could be found in recent review articles [5; 6].

Here we study this problem using the Faddeev formalism [7] in the hard core model [8]. In our approach, in order to change the value of the scattering length, we modify the interatomic potential by multiplying the original interatomic HFD-B potential from [9] by the strength factor λ : $V(x) = \lambda V_{\text{HFD-B}}(x)$. This strategy is extensively used in literature [10; 11; 12] because it allows one to deal with a wide range of scattering lengths.

Based on a talk presented at the 22nd European Conference on Few-Body Problems in Physics (September 9–13, 2013, Cracow, Poland). The paper is to be published in *Few-Body Systems*. This work was supported in part by the Heisenberg-Landau Program

E.A. Kolganova
 Bogoliubov Laboratory of Theoretical Physics
 Joint Institute for Nuclear Research
 Joliot-Curie 6,141980 Dubna, Moscow region, Russia
 E-mail: kea@theor.jinr.ru




Fig. 1 The modified Phillips line plotted with the rescaled dimensionless parameters (1) for the HFD-B potential [9]. Solid line and dashed line correspond to the binding energy of the helium trimer and the virtual state energy, respectively.

Following Refs.[11; 13], we use the Faddeev formalism [7]. After the partial-wave and angular analysis the Faddeev differential equations read

$$\left[-\frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2} + l(l+1) \left(\frac{1}{x^2} + \frac{1}{y^2} \right) - E \right] \Phi_l(x, y) = \begin{cases} -V(x)\Psi_l(x, y), & x > c \\ 0, & x < c. \end{cases} \quad (2)$$

Here x and y stand for the standard Jacobi variables, c for the core range, and l for partial angular momentum. $V(x)$ is the He-He central potential acting outside the core domain. The partial wave function $\Psi_l(x, y)$ is related to the Faddeev components $\Phi_l(x, y)$ (see details in [8]).

The functions $\Phi_l(x, y)$ satisfy the boundary conditions

$$\Phi_l(x, y) |_{x=0} = \Phi_l(x, y) |_{y=0} = 0, \quad \Psi_l(c, y) = 0. \quad (3)$$

In addition, for the bound state problem, the partial wave Faddeev components $\Phi_l(x, y)$ should satisfy the asymptotic boundary condition given by

$$\Phi_l(x, y) = \delta_{l0} \psi_d(x) \exp(i\sqrt{E_3 - \varepsilon_d}y) \left[a_0 + o(y^{-1/2}) \right] + \frac{\exp(i\sqrt{E_3}\rho)}{\sqrt{\rho}} \left[A_l(\theta) + o(\rho^{-1/2}) \right], \quad (4)$$

while for the problem of two-fragment scattering states, the asymptotic boundary condition for the function $\Phi_l(x, y)$ reads

$$\Phi_l(x, y; p) = \delta_{l0} \psi_d(x) \left\{ \sin(py) + \exp(ip y) \left[a_0(p) + o(y^{-1/2}) \right] \right\} + \frac{\exp(i\sqrt{E}\rho)}{\sqrt{\rho}} \left[A_l(\theta, p) + o(\rho^{-1/2}) \right]. \quad (5)$$

Here $\psi_d(x)$ denotes the dimer wave function, E stands for the scattering energy given by $E = \varepsilon_d + p^2$ with ε_d being the dimer energy, and p corresponds to the relative momentum conjugate to the variable y . The variables $\rho = \sqrt{x^2 + y^2}$ and $\theta = \arctan \frac{y}{x}$ are the hyperradius and hyperangle, respectively. One of the important parameters, which determines the behavior of asymptotic boundary conditions (5), is the coefficient $a_0(p)$ that is the elastic scattering amplitude. The functions $A_l(p, \theta)$ provide us, at $E > 0$, with the corresponding partial-wave Faddeev breakup amplitudes. The scattering length ℓ_{sc} is given by

$$\ell_{sc} = -\frac{\sqrt{3}}{2} \lim_{p \rightarrow 0} \frac{a_0(p)}{p}. \quad (6)$$

Now let us discuss the results obtained by the solution of the Faddeev integro-differential equations (2)-(5). In Fig. 1 we plot the dependence of the dimensionless correlation α on ω computed for the HFD-B potential where the scattering length varies from -7000 \AA to 8500 \AA . Almost linear relation between the




Fig. 2 Root locus curves of the real and imaginary parts of the scattering matrix $S_0(z)$ in the case of helium trimer (a) and np system (b). The solid lines correspond to $\text{Re}S_0(z) = 0$ while the tiny dashed lines to $\text{Im}S_0(z) = 0$. Curves 1 and 2 denote the boundaries of the domains where the scattering matrix $S_0(z)$ and Faddeev components $\Phi_l(x, y; z)$ can be analytically continued in z , respectively.

rescaled parameters α and ω is held for large values of ω , for which a weakly bound three-body state exists (solid curve in Fig.1). The nonlinear regime is observed when the scaled scattering length α is small. This nonlinearity originates from the presence of the near-threshold virtual states shown by the dashed curve line in Fig.1. This virtual state violates the linear dependence $\alpha(\omega)$ at the small scattering length. When the scattering length is large and negative, the virtual state turns into a bound state. We studied this mechanism in detail in [11]. In this limit of the large but negative scattering length the linear dependence is again restored. We note that similar results for the weakly bound three-body state were obtained in Ref.[2] where in order to demonstrate the universality of the linear dependence $\alpha(\omega)$ the curve was plotted for different two-body potentials.

Recently, Kievsky and Gattobigio [12] investigated the universal behavior in the elastic scattering below the dimer breakup threshold calculating the ^4He atom-dimer effective-range function. They showed, that the same parametrization of the effective-range function also described the nucleon-deuteron scattering below the deuteron breakup threshold.

The similarity of properties of the atomic helium trimer and nuclear np systems can also be seen in the S-matrix formalism. Let us compare the results obtained by the S-matrix formalism for atomic helium trimer and nuclear np system. To compute the s-state np and $^4\text{He}_3$ resonances, we use the algorithm described in Refs. [11; 14]. We note that the resonances lying on the two-body unphysical sheet are the poles of the S-matrix. The unphysical sheet is connected with the physical one by crossing the spectral interval $(\epsilon_d, 0)$ between the two-body (deuteron or dimer) energy $z = \epsilon_d$ and breakup threshold $z = 0$. Therefore, as it was analytically proved in Ref. [15], these resonances are the roots of the S-matrix on the physical sheet. Using this fact, we solve two-dimensional Faddeev integro-differential equations (2) with the boundary conditions (3) and (5) at complex energies z and extract the truncated s-state scattering matrix

$$S_0(z) = 1 + 2ia_0(z). \quad (7)$$

We start our analysis by considering the $^4\text{He}_3$ system in the only state with the angular momentum $l = 0$. As the interatomic He-He-interaction we employ the semiempirical potential HFD-B [9]. We assumed $\hbar^2/m = 12.12 \text{ K}\text{\AA}^2$ where m stands for the mass of the ^4He atom. Notice that for the HFD-B potential the ^4He -dimer energy is -1.685 mK [13].

In this case, we study graph surfaces of the real and imaginary parts of the scattering matrix $S_0(z)$ in the domain of its holomorphy (curve 1 on Fig.2, see for details [11]). The root lines of the functions $\text{Re}S_0(z)$ and $\text{Im}S_0(z)$ presented in Fig. 2a in the dimensionless case were obtained for the grid parameters $N_\theta = N_\rho = 600$ and $\rho_{\max} = 600 \text{ \AA}$. The resonances are associated with the intersection points of the curves $\text{Re}S_0(z) = 0$ and $\text{Im}S_0(z) = 0$.

Next we study the np system using the Malfliet-Tjon potential MT I-III [16] as NN-interaction. Contrary to the previous case, this system is a fermionic one, nevertheless, it is described by the set of equations similar to (2) - (5). The only difference is that for the doublet nd scattering the function $\Phi_l(x, y)$ is the two-component vector. Therefore, we have to solve the system of two coupled integro-differential equations. The details of calculation and the procedure how to compute the np system can be found in Refs. [7; 14].

In this case, we have succeeded to find one root $z_{\text{res}} = E_{\text{T}}^v$ of the function $S_0(z)$ corresponding to the known virtual state of the ntp system at the total spin $S=1/2$. For the grid described by $N_\theta = N_\rho = 1400$, $\rho_{\text{max}} = 100$ fm we have found $z_{\text{res}} = -2.699$ MeV. We see that it is situated 0.475 MeV to the left from the nd threshold $\varepsilon_d = -2.224$ MeV (in the MT I–III model [16]). Our result agrees well with the shift $\varepsilon_d - z_{\text{res}} = 0.48$ MeV found from experimental data on nd scattering [17]. Moreover, this value is in good agreement with the previous theoretical results known from the literature [18; 19; 20]: the value 0.48 MeV was obtained in [18] as a pole of the T-matrix with a separable version of the MT potential; in [19] the authors studied an effective-range function for the doublet nd scattering while in [20] the complex scaling method was used.

Concerning the structure of the S-matrix in the ntp system, we also studied the root locus lines of the real and imaginary parts of the scattering matrix $S_0(z)$ obtained for the grid parameters $N_\theta = N_\rho = 240$ and $\rho_{\text{max}} = 100$ fm (see Fig.2b). Qualitatively, the structure of the S-matrix for two very different systems presented in Fig.2 a) and b) is very similar. This finding strongly supports the results by Kievsky and Gattobigio [12], and also connects the universal behavior of atomic systems with the large two-body scattering length to the known physics of the nuclear systems with the near-threshold virtual state.

We have investigated theoretically the particle-dimer scattering below and above the three-body threshold. We considered two systems – atomic helium trimer and nuclear ntp systems – and showed that despite the different nature of these systems they show universal correlation. This universality is captured by using Faddeev differential equations and studying the S-matrix structure.

Acknowledgements The author is grateful to A.K.Motovilov and W.Sandhas for their interest in this work and constant support.

References

1. A.C. Phillips (1968) Consistency of the low-energy three-nucleon observables and the separable interaction model, Nucl. Phys. A **107**, 209–216.
2. V. Roudnev and M. Cavagnero (2012) Approaching Universality in Weakly Bound Three-Body Systems, Phys. Rev. Lett. **108**, 110402.
3. V. Efimov and E.G. Tkachenko (1985) Explanation of the Phillips line in the three-nucleon problem, Phys. Lett. B **157**, 108 – 114; V. Efimov and E.G. Tkachenko (1988) On the correlation between the triton binding energy and the neutron-deuteron doublet scattering length. Few-Body Systems **4**, 71 – 88.
4. E. Braaten and H.W. Hammer (2006) Universality in few-body systems with large scattering length, Phys. Rept. **428**, 259–390.
5. L. Platter (2009) Low-Energy Universality in Atomic and Nuclear Physics, Few-Body Systems **46**, 139–171.
6. H.W.Hammer and L. Platter (2010) Efimov States in Nuclear and Particle Physics, Ann. Rev. Nucl. Part. Sci. **60**, 207236.
7. L.D. Faddeev and S.P. Merkuriev (1993) *Quantum scattering theory for several particle systems*. Kluwer Academic Publishers, 404 pp.
8. E.A. Kolganova, A.K. Motovilov, and S.A. Sofianos (1998) Three-body configuration space calculations with hard-core potentials. J. Phys. B **31**, 1279–1302.
9. R.A. Aziz, F.R.W. McCourt, and C.C.K. Wong (1987) A new determination of the ground state interatomic potential for He₂. Mol. Phys. **61**, 1487–1511.
10. B.D. Esry, C.D. Lin, and C.H. Greene (1996) Adiabatic hyperspherical study of the helium trimer, Phys. Rev. A **54**, 394–401.
11. A.K. Motovilov and E.A. Kolganova (1999) Structure of T- and S- matrices in unphysical sheets and resonances in three-body systems, Few-Body Syst. Suppl. **10**, 75–84; E.A. Kolganova and A.K. Motovilov (1999) Mechanism of the emergence of Efimov states in the He₄ trimer, Phys. Atom. Nucl. **62**, 1179–1192 [Yad. Fiz. **62**, 1253–(1997)].
12. A. Kievsky and M. Gattobigio (2013) Universal nature and finite-range correlations in elastic atom-dimer scattering below the dimer breakup threshold. Phys. Rev. A **87**, 052719.
13. E.A. Kolganova, A.K. Motovilov, and W. Sandhas (2009) Ultra cold Collisions in the system of three helium atoms, Phys. Part. Nucl. **40**, 206–235.
14. E.A. Kolganova and A.K. Motovilov (1997) Using Faddeev differential equations to calculate three-body resonances, Phys. Atom. Nucl. **60**, 177–185 [Yad. Fiz. **60**, 235 (1997)].
15. A.K. Motovilov (1997) Representation for the three-body T-matrix, scattering matrices and resolvent in unphysical energy sheets, Math. Nachr. **187**, 147–210.
16. R.A. Malfliet and J.A. Tjon (1969) Solution of the Faddeev equations for the triton problem using local two-particle interactions, Nucl. Phys. A **127**, 161–168.
17. J.E. Purcell, J.H. Kelley, E. Kwan, C.G. Sheu, H.R. Weller (2010) Energy levels of light nuclei image, Nucl. Phys. A **848**, 1–74.
18. Yu.V. Orlov, V.V. Turovtsev (1984) Integral equations for resonance and virtual states, Sov. Phys. JETP **59**, 934–943.
19. Yu.V. Orlov, L.I. Nikitina (2006) Effective-range function for doublet nd scattering from an analysis of modern data, Phys. At. Nucl. **69**, 607–622 [Yad. Fiz. **69**, 631 (2006)].
20. E.A. Kolganova, A.K. Motovilov and Y.K.Ho (2001) Complex Scaling of the Faddeev Operator, Nucl. Phys. A **684**, 623–625.