

Congruences for $q^{[p/8]} \pmod{p}$ II

ZHI-HONG SUN

School of Mathematical Sciences, Huaiyin Normal University,
Huaian, Jiangsu 223001, PR China
E-mail: zhihongsun@yahoo.com
Homepage: <http://www.hytc.edu.cn/xsjl/szh>

ABSTRACT. Let \mathbb{Z} be the set of integers, and let p be a prime of the form $4k + 1$. Suppose $q \in \mathbb{Z}$, $2 \nmid q$, $p \nmid q$, $p = c^2 + d^2$, $c, d \in \mathbb{Z}$ and $c \equiv 1 \pmod{4}$. In this paper we continue to discuss congruences for $q^{[p/8]} \pmod{p}$ and present new reciprocity laws, but we assume $4p = x^2 + qy^2$ or $p = x^2 + 2qy^2$, where $[\cdot]$ is the greatest integer function and $x, y \in \mathbb{Z}$.

MSC: Primary 11A15, Secondary 11A07, 11E25

Keywords: Reciprocity law; octic residue; congruence; quartic Jacobi symbol

1. Introduction.

Let \mathbb{Z} be the set of integers, $i = \sqrt{-1}$ and $\mathbb{Z}[i] = \{a + bi \mid a, b \in \mathbb{Z}\}$. For any positive odd number m and $a \in \mathbb{Z}$ let $\left(\frac{a}{m}\right)$ be the (quadratic) Jacobi symbol. (We also assume $\left(\frac{a}{1}\right) = 1$.) For our convenience we also define $\left(\frac{a}{-m}\right) = \left(\frac{a}{m}\right)$. Then for any two odd numbers m and n with $m > 0$ or $n > 0$ we have the following general quadratic reciprocity law: $\left(\frac{m}{n}\right) = (-1)^{\frac{m-1}{2} \cdot \frac{n-1}{2}} \left(\frac{n}{m}\right)$.

For $a, b, c, d \in \mathbb{Z}$ with $2 \nmid c$ and $2 \mid d$, one can define the quartic Jacobi symbol $\left(\frac{a+bi}{c+di}\right)_4$ as in [S1,S2,S4]. From [IR] we know that $\overline{\left(\frac{a+bi}{c+di}\right)_4} = \left(\frac{a-bi}{c-di}\right)_4 = \left(\frac{a+bi}{c+di}\right)_4^{-1}$, where \bar{x} means the complex conjugate of x . In Section 2 we list main properties of the quartic Jacobi symbol. See also [IR], [BEW] and [S4]. For the history of quartic reciprocity laws, see [Lem].

Let p be a prime of the form $4k + 1$, $q \in \mathbb{Z}$, $2 \nmid q$ and $p \nmid q$. Suppose that $p = c^2 + d^2 = x^2 + qy^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod{4}$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod{4}$. Assume that $(c, x + d) = 1$ or $(d_0, x + c) = 1$, where (m, n) is the greatest common divisor of m and n . In [S5], using the quartic reciprocity law the author deduced some congruences for $q^{[p/8]} \pmod{p}$ in terms of c, d, x and y , where $[\cdot]$ is the greatest integer function. For example, if p and q are primes of the form $8k + 1$, $p = c^2 + d^2 = x^2 + qy^2$, $q = a^2 + b^2$, $a, b, c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod{4}$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod{4}$, then

$$q^{\frac{p-1}{8}} \equiv (-1)^{\frac{d}{4} + \frac{xy}{4}} \left(\frac{d}{c}\right)^m \pmod{p} \iff \left(\frac{ac + bd}{ac - bd}\right)^{\frac{q-1}{8}} \equiv \left(\frac{b}{a}\right)^m \pmod{q}.$$

If $p \equiv 1, 9, 25 \pmod{56}$ is a prime and so $p = c^2 + d^2 = x^2 + 7y^2$ for some $c, d, x, y \in \mathbb{Z}$, by [S5, Corollary 4.2] we have

$$7^{\frac{p-1}{8}} \equiv \begin{cases} -(-1)^{\frac{y}{4}} \pmod{p} & \text{if } 7 \mid c, \\ (-1)^{\frac{y}{4}} \pmod{p} & \text{if } 7 \mid d, \\ \pm(-1)^{\frac{y}{4}} \frac{c}{d} \pmod{p} & \text{if } c \equiv \pm d \pmod{7}. \end{cases}$$

Let p be a prime of the form $4k + 1$, $q \in \mathbb{Z}$, $2 \nmid q$ and $p \nmid q$. Suppose that $p = c^2 + d^2$ with $c, d, x, y \in \mathbb{Z}$ and $c \equiv 1 \pmod{4}$. In this paper we continue to discuss congruences for $q^{\lfloor p/8 \rfloor} \pmod{p}$, but we assume $4p = x^2 + qy^2$ or $p = x^2 + 2qy^2$. Here are some typical results:

★ Let p be a prime of the form $8k + 1$ and $\left(\frac{-11}{p}\right) = 1$, $p = c^2 + d^2$, $4p = x^2 + 11y^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod{4}$ and $d = 2^r d_0 (2 \nmid d_0)$. Assume that $(c, x + 2d) = 1$ or $(d_0, x + 2c) = 1$. Then

$$(-11)^{\frac{p-1}{8}} \equiv \begin{cases} \pm(-1)^{\lfloor \frac{x}{4} \rfloor} \pmod{p} & \text{if } 2 \nmid x \text{ and } x \equiv \pm 4c, \pm 9c \pmod{11}, \\ \pm(-1)^{\lfloor \frac{x}{4} \rfloor} \frac{d}{c} \pmod{p} & \text{if } 2 \nmid x \text{ and } x \equiv \pm 4d, \pm 9d \pmod{11}, \\ \mp(-1)^{\lfloor \frac{x}{8} \rfloor + \frac{y}{8}} \pmod{p} & \text{if } 2 \mid x \text{ and } x \equiv \pm 4c, \pm 9c \pmod{11}, \\ \mp(-1)^{\lfloor \frac{x}{8} \rfloor + \frac{y}{8}} \frac{d}{c} \pmod{p} & \text{if } 2 \mid x \text{ and } x \equiv \pm 4d, \pm 9d \pmod{11}. \end{cases}$$

★ Let p and q be primes such that $p \equiv 1 \pmod{8}$, $q \equiv 7 \pmod{8}$, $p = c^2 + d^2 = x^2 + 2qy^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod{4}$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod{4}$. Assume $(c, x + d) = 1$ or $(d_0, x + c) = 1$. Then

$$(-q)^{\frac{p-1}{8}} \equiv \left(\frac{d}{c}\right)^m \pmod{p} \iff \left(\frac{c - di}{c + di}\right)^{\frac{q+1}{8}} \equiv i^m \pmod{q}.$$

★ Let p be a prime of the form $8k + 1$ and $a \in \mathbb{Z}$ with $2 \nmid a$. Suppose that $p = c^2 + d^2 = x^2 + (a^2 + 1)y^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod{4}$, $d = 2^r d_0$, $d_0 \equiv 1 \pmod{4}$ and $4 \mid y$. Assume $(c, x + d) = 1$ or $(d_0, x + c) = 1$. Then

$$(a + \sqrt{a^2 + 1})^{\frac{p-1}{4}} \equiv (-1)^{\frac{d}{4} + \frac{y}{4}} \pmod{p}.$$

★ Let $p \equiv 1, 9 \pmod{40}$ be a prime and so $p = c^2 + d^2 = x^2 + 10y^2$ with $c, d, x, y \in \mathbb{Z}$. Suppose $c \equiv 1 \pmod{4}$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod{4}$. Assume $(c, x + d) = 1$ or $(d_0, x + c) = 1$. Then

$$(-5)^{\frac{p-1}{8}} \equiv \begin{cases} \pm(-1)^{\frac{d}{4} + \frac{x-1}{2} + \frac{y}{2}} \pmod{p} & \text{if } \frac{2c-d}{x} \equiv \pm 2 \pmod{5}, \\ \pm(-1)^{\frac{d}{4} + \frac{x-1}{2} + \frac{y}{2}} \frac{c}{d} \pmod{p} & \text{if } \frac{2c-d}{x} \equiv \pm 1 \pmod{5}. \end{cases}$$

Throughout this paper, if $n \in \mathbb{Z}$, $2^\alpha \mid n$ and $2^{\alpha+1} \nmid n$, we then write that $2^\alpha \parallel n$.

2. Basic lemmas.

Lemma 2.1 ([S4, Proposition 2.1]). *Let $a, b \in \mathbb{Z}$ with $2 \nmid a$ and $2 \mid b$. Then*

$$\left(\frac{i}{a+bi}\right)_4 = i^{\frac{a^2+b^2-1}{4}} = (-1)^{\frac{a^2-1}{8}} i^{(1-(-1)^{\frac{b}{2}})/2}$$

and

$$\left(\frac{1+i}{a+bi}\right)_4 = \begin{cases} i^{((-1)^{\frac{a-1}{2}}(a-b)-1)/4} & \text{if } 4 \mid b, \\ i^{\frac{(-1)^{\frac{a-1}{2}}(b-a)-1}{4}-1} & \text{if } 2 \parallel b. \end{cases}$$

Lemma 2.2 ([S4, Proposition 2.2]). *Let $a, b \in \mathbb{Z}$ with $2 \nmid a$ and $2 \mid b$. Then*

$$\left(\frac{-1}{a+bi}\right)_4 = (-1)^{\frac{b}{2}} \quad \text{and} \quad \left(\frac{2}{a+bi}\right)_4 = i^{(-1)^{\frac{a-1}{2}} \frac{b}{2}}.$$

Lemma 2.3 ([S4, Proposition 2.3]). *Let $a, b, c, d \in \mathbb{Z}$ with $2 \nmid ac$, $2 \mid b$ and $2 \mid d$. If $a+bi$ and $c+di$ are relatively prime elements of $\mathbb{Z}[i]$, we have the following general law of quartic reciprocity:*

$$\left(\frac{a+bi}{c+di}\right)_4 = (-1)^{\frac{b}{2} \cdot \frac{c-1}{2} + \frac{d}{2} \cdot \frac{a+b-1}{2}} \left(\frac{c+di}{a+bi}\right)_4.$$

In particular, if $4 \mid b$, then

$$\left(\frac{a+bi}{c+di}\right)_4 = (-1)^{\frac{a-1}{2} \cdot \frac{d}{2}} \left(\frac{c+di}{a+bi}\right)_4.$$

Lemma 2.4 ([E], [S1, Lemma 2.1]). *Let $a, b, m \in \mathbb{Z}$ with $2 \nmid m$ and $(m, a^2+b^2) = 1$. Then*

$$\left(\frac{a+bi}{m}\right)_4^2 = \left(\frac{a^2+b^2}{m}\right).$$

Lemma 2.5 ([S3, Lemma 4.3]). *Let $a, b \in \mathbb{Z}$ with $2 \nmid a$ and $2 \mid b$. For any integer x with $(x, a^2+b^2) = 1$ we have*

$$\left(\frac{x^2}{a+bi}\right)_4 = \left(\frac{x}{a^2+b^2}\right).$$

For a given odd prime p let \mathbb{Z}_p denote the set of those rational numbers whose denominator is not divisible by p . Following [S1,S2] we define

$$Q_r(p) = \left\{ k \mid k \in \mathbb{Z}_p, \left(\frac{k+i}{p}\right)_4 = i^r \right\} \quad \text{for } r = 0, 1, 2, 3.$$

Lemma 2.6 ([S1, Theorem 2.3]). *Let p be an odd prime, $r \in \{0, 1, 2, 3\}$, $k \in \mathbb{Z}_p$ and $k^2 + 1 \not\equiv 0 \pmod{p}$.*

(i) *If $p \equiv 1 \pmod{4}$ and $t^2 \equiv -1 \pmod{p}$ with $t \in \mathbb{Z}_p$, then $k \in Q_r(p)$ if and only if $(\frac{k+t}{k-t})^{(p-1)/4} \equiv t^r \pmod{p}$.*

(ii) *If $p \equiv 3 \pmod{4}$, then $k \in Q_r(p)$ if and only if $(\frac{k-i}{k+i})^{(p+1)/4} \equiv i^r \pmod{p}$.*

Lemma 2.7 ([S5, Lemma 2.9]). *Suppose $c, d, m, x \in \mathbb{Z}$, $2 \nmid m$, $x^2 \equiv c^2 + d^2 \pmod{m}$ and $(m, x(x+d)) = 1$. Then*

$$\left(\frac{c+di}{m}\right)_4 = \left(\frac{x(x+d)}{m}\right).$$

Lemma 2.8. *Let p be a prime of the form $8k+1$, $q \in \mathbb{Z}$, $2 \nmid q$ and $p \nmid q$. Suppose that $p = c^2 + d^2 = x^2 + 2qy^2$ with $c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod{4}$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod{4}$. If $(\frac{x/y}{c+di})_4 = (-1)^{s+\frac{p-1}{8}} i^n$, then*

$$(-q)^{\frac{p-1}{8}} \equiv \begin{cases} (-1)^s (\frac{d}{c})^{n-1} \pmod{p} & \text{if } 8 \mid d-4, \\ (-1)^{s+\frac{d}{8}} (\frac{d}{c})^n \pmod{p} & \text{if } 8 \mid d. \end{cases}$$

Proof. As $c \equiv 1 \pmod{4}$ and $4 \mid d$ we see that $c+di$ is primary in $\mathbb{Z}[i]$. Since $i \equiv d/c \pmod{c+di}$ we have

$$\left(\frac{x}{y}\right)^{\frac{p-1}{4}} \equiv \left(\frac{x/y}{c+di}\right)_4 = (-1)^{s+\frac{p-1}{8}} i^n \equiv (-1)^{s+\frac{p-1}{8}} \left(\frac{d}{c}\right)^n \pmod{c+di}.$$

As $(\frac{x}{y})^2 \equiv -q \pmod{p}$ and the norm of $c+di$ is p , from the above we deduce that

$$(-2q)^{\frac{p-1}{8}} \equiv \left(\frac{x}{y}\right)^{\frac{p-1}{4}} \equiv (-1)^{s+\frac{p-1}{8}} \left(\frac{d}{c}\right)^n \pmod{p}.$$

By [L] or [HW, (1.4) and (1.5)],

$$(2.1) \quad (-2)^{\frac{p-1}{8}} \equiv \left(\frac{c}{d}\right)^{-\frac{d}{4}} \equiv \begin{cases} (\frac{c}{d})^{-d_0} \equiv (\frac{c}{d})^{-1} = \frac{d}{c} \pmod{p} & \text{if } 8 \mid d-4, \\ (-1)^{\frac{d}{8}} \pmod{p} & \text{if } 8 \mid d. \end{cases}$$

Thus the result follows.

3. Congruences for $q^{[p/8]} \pmod{p}$ with $4p = x^2 + qy^2$.

Theorem 3.1. *Let p be a prime of the form $4k+1$, $q \in \mathbb{Z}$, $q \equiv 3 \pmod{8}$ and $p \nmid q$. Suppose that $p = c^2 + d^2$, $4p = x^2 + qy^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv y \equiv 1 \pmod{4}$, $(c, x+2d) = 1$ and $(\frac{2c/(x+2d)+i}{q})_4 = i^k$. Then*

$$(-q)^{[p/8]} \equiv \begin{cases} (-1)^{[\frac{x}{4}] + \frac{q-3}{8}} (\frac{d}{c})^{k-1} \pmod{p} & \text{if } 8 \mid p-1, \\ -(-1)^{\frac{x^2-1}{8} + \frac{q-3}{8}} (\frac{d}{c})^{k-1} \frac{y}{x} \pmod{p} & \text{if } 8 \mid p-5. \end{cases}$$

Proof. Since $4p = x^2 + qy^2$ and $y \equiv 1 \pmod{4}$ we see that $2 \nmid x$, $(x, y) = 1$ and $p \nmid x$. Thus $(x, (2c)^2 + (x + 2d)^2) = (x, 4p) = 1$. As $qy^2 = (2c)^2 + (x + 2d)(2d - x)$ we see that $(qy, x + 2d) \mid 4c^2$ and so $(qy, x + 2d) = 1$. Also,

$$\begin{aligned} & (qy^2, (2c)^2 + (x + 2d)^2) \\ &= ((2c)^2 + (x + 2d)^2 - 2x(x + 2d), (2c)^2 + (x + 2d)^2) = (2x(x + 2d), (2c)^2 + (x + 2d)^2) \\ &= (x + 2d, (2c)^2 + (x + 2d)^2) = (x + 2d, 4c^2) = 1. \end{aligned}$$

Now using Lemmas 2.1-2.5 and the fact that $(\frac{a}{n})_4 = 1$ for $a, n \in \mathbb{Z}$ with $2 \nmid n$ and $(a, n) = 1$ we see that

$$\begin{aligned} i^k &= \left(\frac{2c + (x + 2d)i}{q} \right)_4 = \left(\frac{i}{q} \right)_4 \left(\frac{x + 2d - 2ci}{q} \right)_4 \\ &= (-1)^{\frac{q^2-1}{8}} \cdot (-1)^{\frac{q-1}{2}} \left(\frac{q}{x + 2d - 2ci} \right)_4 = \left(\frac{qy^2}{x + 2d - 2ci} \right)_4 \left(\frac{y^2}{x + 2d - 2ci} \right)_4 \\ &= \left(\frac{(x + 2d)^2 + (2c)^2 - 2x(x + 2d)}{x + 2d - 2ci} \right)_4 \left(\frac{y}{(x + 2d)^2 + 4c^2} \right)_4 \\ &= \left(\frac{-2x(x + 2d)}{x + 2d - 2ci} \right)_4 \left(\frac{y}{(x + 2d)^2 + 4c^2} \right)_4 \\ &= - \left(\frac{2}{x + 2d - 2ci} \right)_4 \left(\frac{x(x + 2d)}{x + 2d - 2ci} \right)_4 \left(\frac{y}{(x + 2d)^2 + 4c^2} \right)_4 \\ &= -i^{(-1)^{(x+1)/2}} (-1)^{\frac{x(x+2d)-1}{2}} \left(\frac{x + 2d - 2ci}{x(x + 2d)} \right)_4 \left(\frac{(x + 2d)^2 + 4c^2}{y} \right)_4 \\ &= (-1)^{\frac{x-1}{2}} i \left(\frac{2(d - ci)}{x} \right)_4 \left(\frac{-2ci}{x + 2d} \right)_4 \left(\frac{2x(x + 2d) + qy^2}{y} \right)_4 \\ &= (-1)^{\frac{x-1}{2}} i \left(\frac{d - ci}{x} \right)_4 \left(\frac{i}{x + 2d} \right)_4 \left(\frac{2x(x + 2d)}{y} \right)_4. \end{aligned}$$

Thus, applying Lemma 2.7 we see that

$$\begin{aligned} i^k &= (-1)^{\frac{x-1}{2}} i \left(\frac{-i}{x} \right)_4 \left(\frac{c + di}{x} \right)_4 (-1)^{\frac{(x+2d)^2-1}{8}} \cdot (-1)^{\frac{y^2-1}{8}} \left(\frac{x(x + 2d)}{y} \right)_4 \\ &= (-1)^{\frac{x-1}{2}} i \cdot (-1)^{\frac{x^2-1}{8}} \left(\frac{c + di}{x} \right)_4 (-1)^{\frac{x^2-1}{8} + \frac{d}{2}} \cdot (-1)^{\frac{4p-x^2-q}{8}} \left(\frac{\frac{x}{2}(\frac{x}{2} + d)}{y} \right)_4 \\ &= (-1)^{\frac{x-1}{2} + \frac{d}{2}} i \cdot (-1)^{\frac{x-1}{2} \cdot \frac{d}{2}} \left(\frac{x}{c + di} \right)_4 (-1)^{\frac{x^2-1}{8} + \frac{q-3}{8}} \left(\frac{c + di}{y} \right)_4 \\ &= -(-1)^{\frac{x+1}{2}(\frac{d}{2}+1)} i \left(\frac{x}{c + di} \right)_4 (-1)^{\frac{x^2-1}{8} + \frac{q-3}{8}} \left(\frac{y}{c + di} \right)_4 \\ &= -(-1)^{\frac{x+1}{2}(\frac{d}{2}+1)} \cdot (-1)^{\frac{x^2-1}{8} + \frac{q-3}{8}} i \left(\frac{x}{c + di} \right)_4 \left(\frac{y^{-1}}{c + di} \right)_4 \left(\frac{y^2}{c + di} \right)_4 \\ &= -(-1)^{\frac{x+1}{2}(\frac{d}{2}+1)} \cdot (-1)^{\frac{x^2-1}{8} + \frac{q-3}{8}} i \left(\frac{x/y}{c + di} \right)_4 \left(\frac{y}{c^2 + d^2} \right)_4. \end{aligned}$$

As $\left(\frac{y}{c^2+d^2}\right) = \left(\frac{c^2+d^2}{y}\right) = \left(\frac{4(c^2+d^2)}{y}\right) = \left(\frac{x^2+qy^2}{y}\right) = \left(\frac{x^2}{y}\right) = 1$, from the above we deduce that

$$\left(\frac{x/y}{c+di}\right)_4 = -(-1)^{\frac{x+1}{2}(\frac{d}{2}+1)} \cdot (-1)^{\frac{x^2-1}{8} + \frac{q-3}{8}} i^{k-1}.$$

Clearly $(-1)^{\frac{d}{2}} = (-1)^{\frac{p-1}{4}}$ and $i \equiv d/c \pmod{c+di}$. Since $c+di$ or $-c-di$ is primary in $\mathbb{Z}[i]$, we have

$$\left(\frac{x}{y}\right)^{(p-1)/4} \equiv \left(\frac{x/y}{c+di}\right)_4 \equiv -(-1)^{\frac{x+1}{2} \cdot \frac{p-5}{4} + \frac{x^2-1}{8} + \frac{q-3}{8}} \left(\frac{d}{c}\right)^{k-1} \pmod{c+di}$$

and so

$$\left(\frac{x}{y}\right)^{(p-1)/4} \equiv -(-1)^{\frac{x+1}{2} \cdot \frac{p-5}{4} + \frac{x^2-1}{8} + \frac{q-3}{8}} \left(\frac{d}{c}\right)^{k-1} \pmod{p}.$$

Note that $(x/y)^2 \equiv -q \pmod{p}$. We then have

$$(-q)^{[p/8]} \equiv \begin{cases} \left(\frac{x}{y}\right)^{\frac{p-1}{4}} \equiv (-1)^{\frac{x-1}{2} + \frac{x^2-1}{8} + \frac{q-3}{8}} \left(\frac{d}{c}\right)^{k-1} \pmod{p} & \text{if } 8 \mid p-1, \\ \left(\frac{x}{y}\right)^{\frac{p-1}{4}} \frac{y}{x} \equiv -(-1)^{\frac{x^2-1}{8} + \frac{q-3}{8}} \left(\frac{d}{c}\right)^{k-1} \frac{y}{x} \pmod{p} & \text{if } 8 \mid p-5. \end{cases}$$

This yields the result.

Corollary 3.1. *Let p be a prime of the form $12k+1$, $p = c^2 + d^2$, $4p = x^2 + 27y^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv y \equiv 1 \pmod{4}$ and $(c, x+2d) = 1$.*

(i) *If $p \equiv 1 \pmod{24}$, then*

$$(-3)^{\frac{p-1}{8}} \equiv \begin{cases} (-1)^{[\frac{p}{4}]} \frac{d}{c} \pmod{p} & \text{if } c \equiv 0 \pmod{3}, \\ \pm(-1)^{[\frac{p}{4}]} \pmod{p} & \text{if } x-d \equiv \pm c \pmod{3}. \end{cases}$$

(ii) *If $p \equiv 13 \pmod{24}$, then*

$$(-3)^{\frac{p-5}{8}} \equiv \begin{cases} -(-1)^{\frac{x^2-1}{8}} \frac{3dy}{cx} \pmod{p} & \text{if } c \equiv 0 \pmod{3}, \\ \pm(-1)^{\frac{x^2-1}{8}} \frac{3y}{x} \pmod{p} & \text{if } x-d \equiv \pm c \pmod{3}. \end{cases}$$

Proof. Clearly

$$\left(\frac{i}{3}\right)_4 = -1, \quad \left(\frac{1+i}{3}\right)_4 = -i, \quad \left(\frac{-1+i}{3}\right)_4 = i.$$

and

$$(-3)^{[p/8]} = 3^{4[p/8]} (-27)^{-[p/8]} \equiv 3^{(-1)^{(p-1)/4} - 1} (-27)^{-[p/8]} \pmod{p}.$$

Now taking $q = 27$ in Theorem 3.1 and then applying the above we deduce the result.

Theorem 3.2. *Let p be a prime of the form $4k + 1$, $q \in \mathbb{Z}$, $q \equiv 3 \pmod{8}$ and $p \nmid q$. Suppose that $p = c^2 + d^2$, $4p = x^2 + qy^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv y \equiv 1 \pmod{4}$, $(d, x + 2c) = 1$ and $(\frac{-2d/(x+2c)+i}{q})_4 = i^k$. Then*

$$(-q)^{[p/8]} \equiv \begin{cases} (-1)^{[\frac{x}{4}]} (\frac{d}{c})^k \pmod{p} & \text{if } 8 \mid p - 1, \\ -(-1)^{\frac{x^2-1}{8}} (\frac{d}{c})^k \frac{y}{x} \pmod{p} & \text{if } 8 \mid p - 5. \end{cases}$$

Proof. Since $4p = x^2 + qy^2$ and $y \equiv 1 \pmod{4}$ we see that $2 \nmid x$, $(x, y) = 1$ and $p \nmid x$. Thus $(x, (2d)^2 + (x + 2c)^2) = (x, 4p) = 1$. As $qy^2 = (2d)^2 + (x + 2c)(2c - x)$ we see that $(qy, x + 2c) \mid (2d)^2$ and so $(qy, x + 2c) = 1$. Also,

$$\begin{aligned} & (qy^2, (2d)^2 + (x + 2c)^2) \\ &= ((2d)^2 + (x + 2c)^2 - 2x(x + 2c), (2d)^2 + (x + 2c)^2) = (2x(x + 2c), (2d)^2 + (x + 2c)^2) \\ &= (x + 2c, (2d)^2 + (x + 2c)^2) = (x + 2c, 4d^2) = 1. \end{aligned}$$

Now using Lemmas 2.1-2.5 and the fact that $(\frac{a}{n})_4 = 1$ for $a, n \in \mathbb{Z}$ with $2 \nmid n$ and $(a, n) = 1$ we see that

$$\begin{aligned} i^k &= \left(\frac{-2d + (x + 2c)i}{q} \right)_4 = \left(\frac{i}{q} \right)_4 \left(\frac{x + 2c + 2di}{q} \right)_4 \\ &= (-1)^{\frac{q^2-1}{8}} \left(\frac{q}{x + 2c + 2di} \right)_4 = - \left(\frac{qy^2}{x + 2c + 2di} \right)_4 \left(\frac{y^2}{x + 2c + 2di} \right)_4 \\ &= - \left(\frac{(x + 2c)^2 + (2d)^2 - 2x(x + 2c)}{x + 2c + 2di} \right)_4 \left(\frac{y}{(x + 2c)^2 + 4d^2} \right)_4 \\ &= - \left(\frac{2}{x + 2c + 2di} \right)_4 \left(\frac{x(x + 2c)}{x + 2c + 2di} \right)_4 \left(\frac{y}{(x + 2c)^2 + 4d^2} \right)_4 \\ &= -(-1)^{\frac{d}{2}} \left(\frac{x + 2c + 2di}{x(x + 2c)} \right)_4 \left(\frac{(x + 2c)^2 + 4d^2}{y} \right)_4 \\ &= -(-1)^{\frac{p-1}{4}} \left(\frac{2(c + di)}{x} \right)_4 \left(\frac{2di}{x + 2c} \right)_4 \left(\frac{2x(x + 2c) + qy^2}{y} \right)_4 \\ &= -(-1)^{\frac{p-1}{4}} \left(\frac{c + di}{x} \right)_4 \left(\frac{i}{x + 2c} \right)_4 \left(\frac{2x(x + 2c)}{y} \right)_4. \end{aligned}$$

Thus, applying Lemma 2.7 we see that

$$\begin{aligned}
i^k &= -(-1)^{\frac{p-1}{4}} \left(\frac{c+di}{x}\right)_4 (-1)^{\frac{(x+2c)^2-1}{8}} \left(\frac{2}{y}\right) \left(\frac{x(x+2c)}{y}\right) \\
&= -(-1)^{\frac{p-1}{4}} \left(\frac{c+di}{x}\right)_4 (-1)^{\frac{x^2-1}{8} + \frac{x+c}{2}} \left(\frac{i}{y}\right)_4 \left(\frac{\frac{x}{2}(\frac{x}{2}+c)}{y}\right) \\
&= -(-1)^{\frac{p-1}{4}} \cdot (-1)^{\frac{x-1}{2} \cdot \frac{d}{2}} \left(\frac{x}{c+di}\right)_4 (-1)^{\frac{x^2-1}{8} + \frac{x+1}{2}} \left(\frac{i}{y}\right)_4 \left(\frac{d+ci}{y}\right)_4 \\
&= -(-1)^{(1+\frac{x-1}{2})(1+\frac{p-1}{4})} \cdot (-1)^{\frac{x^2-1}{8}} \left(\frac{x}{c+di}\right)_4 \left(\frac{-c+di}{y}\right)_4 \\
&= -(-1)^{\frac{x+1}{2} \cdot \frac{p+3}{4} + \frac{x^2-1}{8}} \left(\frac{x}{c+di}\right)_4 \left(\frac{c+di}{y}\right)_4^{-1} \\
&= -(-1)^{\frac{x+1}{2} \cdot \frac{p+3}{4} + \frac{x^2-1}{8}} \left(\frac{x}{c+di}\right)_4 \left(\frac{y}{c+di}\right)_4^{-1} \\
&= -(-1)^{\frac{x+1}{2} \cdot \frac{p+3}{4} + \frac{x^2-1}{8}} \left(\frac{x/y}{c+di}\right)_4.
\end{aligned}$$

Clearly $(-1)^{\frac{d}{2}} = (-1)^{\frac{p-1}{4}}$ and $i \equiv d/c \pmod{c+di}$. Since $c+di$ or $-c-di$ is primary in $\mathbb{Z}[i]$, we have

$$\left(\frac{x}{y}\right)^{(p-1)/4} \equiv \left(\frac{x/y}{c+di}\right)_4 \equiv -(-1)^{\frac{x+1}{2} \cdot \frac{p+3}{4} + \frac{x^2-1}{8}} \left(\frac{d}{c}\right)^k \pmod{c+di}$$

and so

$$\left(\frac{x}{y}\right)^{(p-1)/4} \equiv -(-1)^{\frac{x+1}{2} \cdot \frac{p+3}{4} + \frac{x^2-1}{8}} \left(\frac{d}{c}\right)^k \pmod{p}.$$

Note that $(x/y)^2 \equiv -q \pmod{p}$. We then have

$$(-q)^{\lfloor p/8 \rfloor} \equiv \begin{cases} \left(\frac{x}{y}\right)^{\frac{p-1}{4}} \equiv -(-1)^{\frac{x+1}{2} + \frac{x^2-1}{8}} \left(\frac{d}{c}\right)^k = (-1)^{\lfloor \frac{x}{4} \rfloor} \left(\frac{d}{c}\right)^k \pmod{p} & \text{if } 8 \mid p-1, \\ \left(\frac{x}{y}\right)^{\frac{p-1}{4}} \frac{y}{x} \equiv -(-1)^{\frac{x^2-1}{8}} \left(\frac{d}{c}\right)^k \frac{y}{x} \pmod{p} & \text{if } 8 \nmid p-1. \end{cases}$$

This is the result.

Remark 3.1 We note that the k in Theorem 3.1 depends only on $\frac{c}{x+2d} \pmod{q}$, and the k in Theorem 3.2 depends only on $\frac{d}{x+2c} \pmod{q}$.

Theorem 3.3. *Let p and q be primes such that $p \equiv 1 \pmod{4}$ and $q \equiv 3 \pmod{8}$. Suppose $p = c^2 + d^2$, $4p = x^2 + qy^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv y \equiv 1 \pmod{4}$, and $\left(\frac{2(c-di)}{x}\right)^{\frac{q+1}{4}} \equiv i^m \pmod{q}$. Assume $(c, x+2d) = 1$ or $(d, x+2c) = 1$. Then*

$$(-q)^{\lfloor p/8 \rfloor} \equiv \begin{cases} (-1)^{\lfloor \frac{x}{4} \rfloor + 1} \left(\frac{d}{c}\right)^m \pmod{p} & \text{if } p \equiv 1 \pmod{8}, \\ (-1)^{\frac{x^2-1}{8}} \left(\frac{d}{c}\right)^m \frac{y}{x} \pmod{p} & \text{if } p \equiv 5 \pmod{8}. \end{cases}$$

Proof. Clearly $q \nmid x$ and x is odd. We first assume $(c, x + 2d) = 1$. By the proof of Theorem 3.1, $(q, (x + 2d)(c^2 + (x + 2d)^2)) = 1$. It is easily seen that $\frac{2c/(x+2d)-i}{2c/(x+2d)+i} = \frac{2c-(x+2d)i}{2c+(x+2d)i} \equiv \frac{2(c-di)}{ix} \pmod{q}$. Thus, for $k = 0, 1, 2, 3$, using Lemma 2.6 we get

$$\begin{aligned} \left(\frac{2c/(x+2d)+i}{q}\right)_4 &= i^k \\ \iff \frac{2c}{x+2d} \in Q_k(q) &\iff \left(\frac{\frac{2c}{x+2d}-i}{\frac{2c}{x+2d}+i}\right)^{\frac{q+1}{4}} \equiv i^k \pmod{q} \\ \iff \left(\frac{2(c-di)}{ix}\right)^{\frac{q+1}{4}} &\equiv i^k \pmod{q} \iff \left(\frac{2(c-di)}{x}\right)^{\frac{q+1}{4}} \equiv i^{\frac{q+1}{4}+k} \pmod{q}. \end{aligned}$$

Since $\left(\frac{2(c-di)}{x}\right)^{\frac{q+1}{4}} \equiv i^m \pmod{q}$, from the above we deduce that

$$\left(\frac{2c/(x+2d)+i}{q}\right)_4 = i^{m-\frac{q+1}{4}} = (-1)^{\frac{q-3}{8}} i^{m-1}.$$

Now, applying Theorem 3.1 we derive the result.

Now we assume $(d, x + 2c) = 1$. By the proof of Theorem 3.2, $(q, x + 2c) = (q, d^2 + (x + 2c)^2) = 1$. It is easily seen that $\frac{2d+(x+2c)i}{2d-(x+2c)i} \equiv \frac{2(c-di)}{-x} \pmod{q}$. Thus, for $k = 0, 1, 2, 3$, using Lemma 2.6 we get

$$\begin{aligned} \left(\frac{-2d/(x+2c)+i}{q}\right)_4 &= i^k \\ \iff -\frac{2d}{x+2c} \in Q_k(q) &\iff \left(\frac{-\frac{2d}{x+2c}-i}{-\frac{2d}{x+2c}+i}\right)^{\frac{q+1}{4}} \equiv i^k \pmod{q} \\ \iff \left(\frac{2d+(x+2c)i}{2d-(x+2c)i}\right)^{\frac{q+1}{4}} &\equiv i^k \pmod{q} \iff \left(\frac{2(c-di)}{-x}\right)^{\frac{q+1}{4}} \equiv i^k \pmod{q} \\ \iff \left(\frac{2(c-di)}{x}\right)^{\frac{q+1}{4}} &\equiv i^{\frac{q+1}{2}+k} \pmod{q}. \end{aligned}$$

Since $\left(\frac{2(c-di)}{x}\right)^{\frac{q+1}{4}} \equiv i^m \pmod{q}$, from the above we deduce that $\left(\frac{-2d/(x+2c)+i}{q}\right)_4 = i^{m-\frac{q+1}{2}} = (-1)^{\frac{q+1}{4}} i^m = -i^m$. Now applying Theorem 3.2 we deduce the result. The proof is now complete.

Theorem 3.4. *Let p and q be primes such that $p \equiv 1 \pmod{4}$ and $q \equiv 3 \pmod{8}$. Suppose $p = c^2 + d^2$, $4p = x^2 + qy^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv y \equiv 1 \pmod{4}$, and $q \mid cd$. Assume that $(c, x + 2d) = 1$ or $(d, x + 2c) = 1$. Then*

$$(-q)^{\left[\frac{p}{8}\right]} \equiv \begin{cases} \pm(-1)^{\left[\frac{x}{4}\right]+1} \pmod{p} & \text{if } p \equiv 1 \pmod{8} \text{ and } x \equiv \pm 2c \pmod{q}, \\ \pm(-1)^{\frac{q-3}{8}+\left[\frac{x}{4}\right]} \frac{d}{c} \pmod{p} & \text{if } p \equiv 1 \pmod{8} \text{ and } x \equiv \pm 2d \pmod{q}, \\ \pm(-1)^{\frac{x^2-1}{8}} \frac{y}{x} \pmod{p} & \text{if } p \equiv 5 \pmod{8} \text{ and } x \equiv \pm 2c \pmod{q}, \\ \mp(-1)^{\frac{q-3}{8}+\frac{x^2-1}{8}} \frac{dy}{cx} \pmod{p} & \text{if } p \equiv 5 \pmod{8} \text{ and } x \equiv \pm 2d \pmod{q}, \end{cases}$$

Proof. As $(\frac{x}{2})^2 \equiv c^2 + d^2 \pmod{q}$ we see that $x \equiv \pm 2c \pmod{q} \Leftrightarrow q \mid d$ and that $x \equiv \pm 2d \pmod{q} \Leftrightarrow q \mid c$. Thus,

$$\left(\frac{c-di}{x/2}\right)_4^{\frac{q+1}{4}} \equiv \begin{cases} (\pm 1)^{\frac{q+1}{4}} = \pm 1 \pmod{q} & \text{if } x \equiv \pm 2c \pmod{q}, \\ (\mp i)^{\frac{q-1}{4}} = \mp(-1)^{\frac{q-3}{8}}i \pmod{q} & \text{if } x \equiv \pm 2d \pmod{q}. \end{cases}$$

Now applying Theorem 3.3 we derive the result.

Theorem 3.5. *Let p be a prime of the form $4k+1$ and $(\frac{-11}{p}) = 1$, $p = c^2 + d^2$, $4p = x^2 + 11y^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod{4}$, $d = 2^r d_0 (2 \nmid d_0)$, $y = 2^t y_0$ and $y_0 \equiv 1 \pmod{4}$. Assume that $(c, x+2d) = 1$ or $(d_0, x+2c) = 1$.*

(i) *If $p \equiv 1 \pmod{8}$, then*

$$(-11)_{\left[\frac{p}{8}\right]} \equiv \begin{cases} \pm(-1)^{\left[\frac{x}{4}\right]} \pmod{p} & \text{if } 2 \nmid x \text{ and } x \equiv \pm 4c, \pm 9c \pmod{11}, \\ \pm(-1)^{\left[\frac{x}{4}\right]} \frac{d}{c} \pmod{p} & \text{if } 2 \nmid x \text{ and } x \equiv \pm 4d, \pm 9d \pmod{11}, \\ \mp(-1)^{\left[\frac{x}{8}\right] + \frac{y}{8}} \pmod{p} & \text{if } 2 \mid x \text{ and } x \equiv \pm 4c, \pm 9c \pmod{11}, \\ \mp(-1)^{\left[\frac{x}{8}\right] + \frac{y}{8}} \frac{d}{c} \pmod{p} & \text{if } 2 \mid x \text{ and } x \equiv \pm 4d, \pm 9d \pmod{11}. \end{cases}$$

(ii) *If $p \equiv 5 \pmod{8}$, then*

$$(-11)_{\left[\frac{p}{8}\right]} \equiv \begin{cases} \mp(-1)^{\frac{x^2-1}{8}} \frac{y}{x} \pmod{p} & \text{if } 2 \nmid x \text{ and } x \equiv \pm 4c, \pm 9c \pmod{11}, \\ \mp(-1)^{\frac{x^2-1}{8}} \frac{dy}{cx} \pmod{p} & \text{if } 2 \nmid x \text{ and } x \equiv \pm 4d, \pm 9d \pmod{11}, \\ \mp(-1)^{\frac{p-5}{8}} \frac{y}{x} \pmod{p} & \text{if } 2 \mid x \text{ and } x \equiv \pm 4c, \pm 9c \pmod{11}, \\ \mp(-1)^{\frac{p-5}{8}} \frac{dy}{cx} \pmod{p} & \text{if } 2 \mid x \text{ and } x \equiv \pm 4d, \pm 9d \pmod{11}. \end{cases}$$

Proof. As

$$\left(\frac{x}{2}\right)^2 \equiv c^2 + d^2 \pmod{11} \quad \text{and} \quad (c-di)^3 = c(c^2 - 3d^2) + d(d^2 - 3c^2)i,$$

we see that

$$\left(\frac{c-di}{x/2}\right)_4^3 \equiv \begin{cases} \mp 1 \pmod{p} & \text{if } x \equiv \pm 4c, \pm 9c \pmod{11}, \\ \mp i \pmod{p} & \text{if } x \equiv \pm 4d, \pm 9d \pmod{11}. \end{cases}$$

When $2 \nmid x$, from the above and Theorem 3.3 (with $q = 11$) we deduce the result. When $2 \mid x$ and $p \equiv 1 \pmod{8}$, we have $8 \mid y$ and so $(-1)^{\frac{p-1}{8} + \frac{x/2-1}{2}} = (-1)^{\frac{(x/2)^2-1}{8} + \frac{x/2-1}{2}} = (-1)^{\left[\frac{x}{8}\right]}$. Thus, applying the above and [S5, Theorem 4.1 (with $q = 11$)] we obtain the result.

Remark 3.2 Let p be a prime of the form $4k+1$, $q \in \mathbb{Z}$, $q \equiv 3 \pmod{8}$, $p \nmid q$, $p = c^2 + d^2$, $4p = x^2 + qy^2$, $c, d, x, y \in \mathbb{Z}$, $2 \nmid xy$, $c \equiv 1 \pmod{4}$, we conjecture that one can always choose the sign of x such that $(c, x+2d) = 1$ or $(d, x+2c) = 1$. Thus the condition $(c, x+2d) = 1$ or $(d, x+2c) = 1$ in Theorems 3.1-3.5 can be canceled. See also [S5, Remark 4.1].

4. Congruences for $(-q)^{(p-1)/8} \pmod{p}$ with $p = x^2 + 2qy^2$.

Theorem 4.1. *Let p be a prime of the form $8k + 1$, $q \in \mathbb{Z}$, $2 \nmid q$ and $p \nmid q$. Suppose that $p = c^2 + d^2 = x^2 + 2qy^2$ with $c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod{4}$, $d = 2^r d_0$, $d_0 \equiv 1 \pmod{4}$ and $(c, x + d) = 1$. Assume that $(\frac{c/(x+d)+i}{q})_4 = i^k$. Then*

$$(-q)^{\frac{p-1}{8}} \equiv \begin{cases} (-1)^{\frac{q-1}{8} + \frac{d}{4} + \frac{y}{2}} \left(\frac{d}{c}\right)^k \pmod{p} & \text{if } q \equiv 1 \pmod{8}, \\ (-1)^{\frac{q-3}{8} + \frac{x-1}{2}} \left(\frac{d}{c}\right)^{k+1} \pmod{p} & \text{if } q \equiv 3 \pmod{8}, \\ (-1)^{\frac{q-5}{8} + \frac{d}{4} + \frac{x-1}{2} + \frac{y}{2}} \left(\frac{d}{c}\right)^{k-1} \pmod{p} & \text{if } q \equiv 5 \pmod{8}, \\ (-1)^{\frac{q+1}{8}} \left(\frac{d}{c}\right)^k \pmod{p} & \text{if } q \equiv 7 \pmod{8}. \end{cases}$$

Proof. Suppose $y = 2^t y_0$ ($2 \nmid y_0$). We may assume $y_0 \equiv 1 \pmod{4}$. Since $p = c^2 + d^2 = x^2 + 2qy^2 \equiv 1 \pmod{8}$ we see that $2 \nmid x$, $2 \mid y$, $4 \mid d$, $(x, y) = 1$ and $p \nmid x$. Thus $(x, c^2 + (x+d)^2) = (x, p) = 1$. As $2qy^2 = c^2 + (d+x)(d-x)$ we see that $(qy, x+d) \mid c^2$ and so $(qy, x+d) = 1$. Also,

$$\begin{aligned} & (qy^2, (c^2 + (x+d)^2)/2) \\ &= ((c^2 + (x+d)^2)/2 - x(x+d), (c^2 + (x+d)^2)/2) = (x(x+d), (c^2 + (x+d)^2)/2) \\ &= (x(x+d), c^2 + (x+d)^2) = (x+d, c^2) = 1. \end{aligned}$$

It is easily seen that

$$c + (x+d)i = i^{\frac{1+i}{2}} (1+i) \left(\frac{x+d \pm c}{2} + \frac{\pm(x+d) - c}{2} i \right)$$

and so

$$\left(\frac{x+d \pm c}{2} \right)^2 + \left(\frac{\pm(x+d) - c}{2} \right)^2 = \frac{c^2 + (x+d)^2}{2}.$$

Set $\varepsilon = (-1)^{\frac{x-1}{2}}$. As $4 \mid d$ and $4 \mid c-1$ we have $x+d \equiv \varepsilon \pmod{4}$ and $4 \mid (\varepsilon(x+d) - c)$. Using Lemmas 2.1-2.5 and [S5, Lemma 2.10(ii)] and the above we see that

$$\begin{aligned} i^k &= \left(\frac{c + (x+d)i}{q} \right)_4 = \left(\frac{i}{q} \right)_4^{\frac{1-\varepsilon}{2}} \left(\frac{1+i}{q} \right)_4 \left(\frac{\frac{x+d+\varepsilon c}{2} + \frac{\varepsilon(x+d)-c}{2} i}{q} \right)_4 \\ &= (-1)^{\frac{q - (\frac{-1}{q})}{4} \cdot \frac{x-1}{2}} i^{\frac{(-1)}{q} q-1} (-1)^{\frac{q-1}{2} \cdot \frac{\varepsilon(x+d)-c}{4}} \left(\frac{q}{\frac{x+d+\varepsilon c}{2} + \frac{\varepsilon(x+d)-c}{2} i} \right)_4 \end{aligned}$$

and

$$\begin{aligned}
& \left(\frac{q}{\frac{x+d+\varepsilon c}{2} + \frac{\varepsilon(x+d)-c}{2}i} \right)_4 \\
&= \left(\frac{qy^2}{\frac{x+d+\varepsilon c}{2} + \frac{\varepsilon(x+d)-c}{2}i} \right)_4 \left(\frac{y^2}{\frac{x+d+\varepsilon c}{2} + \frac{\varepsilon(x+d)-c}{2}i} \right)_4 \\
&= \left(\frac{(c^2 + (x+d)^2)/2 - x(x+d)}{\frac{x+d+\varepsilon c}{2} + \frac{\varepsilon(x+d)-c}{2}i} \right)_4 \left(\frac{y}{(\frac{x+d+\varepsilon c}{2})^2 + (\frac{\varepsilon(x+d)-c}{2})^2} \right) \\
&= \left(\frac{-x(x+d)}{\frac{x+d+\varepsilon c}{2} + \frac{\varepsilon(x+d)-c}{2}i} \right)_4 \left(\frac{y}{(c^2 + (x+d)^2)/2} \right) \\
&= (-1)^{\frac{\varepsilon(x+d)-c}{4}} \left(\frac{\frac{x+d+\varepsilon c}{2} + \frac{\varepsilon(x+d)-c}{2}i}{x(x+d)} \right)_4 (-1)^{\frac{c^2-(x+d)^2}{8}t + \frac{d}{4}t} \left(\frac{y^{-1}}{c+di} \right)_4.
\end{aligned}$$

Clearly,

$$(-1)^{\frac{c^2-(x+d)^2}{8}} = (-1)^{\frac{c^2-x^2-2dx}{8}} = (-1)^{\frac{c-\varepsilon x}{4} \cdot \frac{c+\varepsilon x}{2} - \frac{d}{4}\varepsilon} = (-1)^{\frac{c-\varepsilon x}{4} - \frac{d}{4}\varepsilon} = (-1)^{\frac{\varepsilon(x+d)-c}{4}}.$$

It is easily seen that

$$\begin{aligned}
& \left(\frac{\frac{x+d+\varepsilon c}{2} + \frac{\varepsilon(x+d)-c}{2}i}{x(x+d)} \right)_4 \\
&= \left(\frac{x+d+\varepsilon c + (\varepsilon(x+d)-c)i}{x} \right)_4 \left(\frac{x+d+\varepsilon c + (\varepsilon(x+d)-c)i}{x+d} \right)_4 \\
&= \left(\frac{d+\varepsilon c + (\varepsilon d - c)i}{x} \right)_4 \left(\frac{\varepsilon c - ci}{x+d} \right)_4 = \left(\frac{(\varepsilon - i)(c+di)}{x} \right)_4 \left(\frac{\varepsilon - i}{x+d} \right)_4 \\
&= \left(\frac{\varepsilon - i}{x(x+d)} \right)_4 \left(\frac{c+di}{x} \right)_4 = \left(\frac{i^{\frac{5+\varepsilon}{2}}(1+i)}{x(x+d)} \right)_4 \left(\frac{c+di}{x} \right)_4 \\
&= \left(\frac{i}{x(x+d)} \right)_4^{\frac{5+\varepsilon}{2}} \left(\frac{1+i}{x(x+d)} \right)_4 \left(\frac{x}{c+di} \right)_4 \\
&= (-1)^{\frac{x(x+d)-1}{4} \cdot \frac{5+\varepsilon}{2}} i^{\frac{x(x+d)-1}{4}} \left(\frac{x}{c+di} \right)_4 \\
&= (-1)^{\frac{d}{4} \cdot \frac{x+1}{2} + \frac{x^2-1}{8}} i^{\frac{dx}{4}} \left(\frac{x}{c+di} \right)_4.
\end{aligned}$$

Hence

$$\begin{aligned}
\left(\frac{q}{\frac{x+d+\varepsilon c}{2} + \frac{\varepsilon(x+d)-c}{2}i} \right)_4 &= (-1)^{\frac{\varepsilon(x+d)-c}{4}(1+t) + \frac{d}{4}t} \cdot (-1)^{\frac{d}{4} \cdot \frac{x+1}{2} + \frac{x^2-1}{8}} i^{\frac{dx}{4}} \left(\frac{x/y}{c+di} \right)_4 \\
&= (-1)^{\frac{\varepsilon x - c}{4}(1+t) + \frac{d}{4} \cdot \frac{x-1}{2} + \frac{x^2-1}{8}} i^{\frac{dx}{4}} \left(\frac{x/y}{c+di} \right)_4.
\end{aligned}$$

Therefore

$$i^k = (-1)^{\frac{q - \left(\frac{-1}{q}\right) \cdot \frac{x-1}{2} + \frac{q-1}{2} \left(\frac{\varepsilon x - c}{4} + \frac{d}{4}\right)} i^{\frac{\left(\frac{-1}{q}\right)_{q-1}}{4}} \\ \times (-1)^{\frac{\varepsilon x - c}{4} (1+t) + \frac{d}{4} \cdot \frac{x-1}{2} + \frac{x^2-1}{8}} i^{\frac{dx}{4}} \left(\frac{x/y}{c + di} \right)_4.$$

It is clear that

$$i^{\frac{dx}{4}} = i^{\frac{d}{4}(x-1) + \frac{d}{4}} = (-1)^{\frac{d}{4} \cdot \frac{x-1}{2}} i^{\frac{d}{4}}, \quad (-1)^{\frac{x^2-1}{8}} = (-1)^{\frac{p-1-2qy^2}{8}} = (-1)^{\frac{p-1}{8} + \frac{y}{2}}, \\ (-1)^{\frac{\varepsilon x - c}{4}} = (-1)^{\frac{\varepsilon x - c}{4} \cdot \frac{\varepsilon x + c}{2}} = (-1)^{\frac{x^2 - c^2}{8}} = (-1)^{\frac{d^2 - 8q\left(\frac{y}{2}\right)^2}{8}} = (-1)^{\frac{y}{2}}$$

and so $(-1)^{\frac{\varepsilon x - c}{4}(1+t)} = (-1)^{\frac{y}{2}(1+t)} = 1$. Thus,

$$\left(\frac{x/y}{c + di} \right)_4 = (-1)^{\frac{q - \left(\frac{-1}{q}\right) \cdot \frac{x-1}{2} + \frac{q-1}{2} \left(\frac{y}{2} + \frac{d}{4}\right) + \frac{p-1}{8} + \frac{y}{2}} i^{k - \frac{d}{4} - \frac{q\left(\frac{-1}{q}\right)_{-1}}{4}}.$$

Now applying Lemma 2.8 we deduce the result.

Theorem 4.2. *Let p be a prime of the form $4k + 1$, $q \in \mathbb{Z}$, $2 \nmid q$ and $p \nmid q$. Suppose that $p = c^2 + d^2 = x^2 + 2qy^2$ with $c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod{4}$, $d = 2^r d_0$, $d_0 \equiv 1 \pmod{4}$, $(d_0, x + c) = 1$ and $\left(\frac{-d/(x+c)+i}{q}\right)_4 = i^k$. Then*

$$(-q)^{\frac{p-1}{8}} \equiv \begin{cases} (-1)^{\frac{x+1}{2} \cdot \frac{q-1}{4} + \frac{d}{4} + \frac{y}{2}} \left(\frac{d}{c}\right)^k \pmod{p} & \text{if } q \equiv 1 \pmod{4}, \\ (-1)^{\frac{x+1}{2} \cdot \frac{q+1}{4}} \left(\frac{d}{c}\right)^k \pmod{p} & \text{if } q \equiv 3 \pmod{4}. \end{cases}$$

Proof. Suppose $2^m \parallel (x+c)$ and $y = 2^t y_0$ ($2 \nmid y_0$). We may assume $y_0 \equiv 1 \pmod{4}$. As $(x+c, d_0) = 1$, by Lemma 2.11 we have $(qy_0, x+c) = 1$ and $(qy_0^2, (x+c)^2 + d^2) = 1$. Note that $(x, y)^2 \mid p$. We also have $(x, y) = 1$. We prove the theorem by considering the following three cases:

Case 1. $m < r$. Using Lemmas 2.1-2.5 and the fact $\left(\frac{a}{q}\right)_4 = 1$ for $a \in \mathbb{Z}$ with $(a, q) = 1$ we see that

$$\left(\frac{d - (x+c)i}{q} \right)_4 = \left(\frac{-2^m}{q} \right)_4 \left(\frac{i}{q} \right)_4 \left(\frac{\frac{x+c}{2^m} + \frac{d}{2^m}i}{q} \right)_4 = (-1)^{\frac{q^2-1}{8} + \frac{q-1}{2} \cdot \frac{d}{2^{m+1}}} \left(\frac{q}{\frac{x+c}{2^m} + \frac{d}{2^m}i} \right)_4 \\ = (-1)^{\frac{q^2-1}{8} + \frac{q-1}{2} \cdot \frac{d}{2^{m+1}}} \left(\frac{qy^2}{\frac{x+c}{2^m} + \frac{d}{2^m}i} \right)_4 \left(\frac{y^2}{\frac{x+c}{2^m} + \frac{d}{2^m}i} \right)_4 \\ = (-1)^{\frac{q^2-1}{8} + \frac{q-1}{2} \cdot \frac{d}{2^{m+1}}} \left(\frac{((x+c)^2 + d^2)/2 - x(x+c)}{\frac{x+c}{2^m} + \frac{d}{2^m}i} \right)_4 \left(\frac{y}{\frac{(x+c)^2 + d^2}{2^{2m}}} \right) \\ = (-1)^{\frac{q^2-1}{8} + \frac{q-1}{2} \cdot \frac{d}{2^{m+1}}} \left(\frac{-2^m x(x+c)/2^m}{\frac{x+c}{2^m} + \frac{d}{2^m}i} \right)_4 \left(\frac{y}{\frac{(x+c)^2 + d^2}{2^{2m}}} \right).$$

By [S5, p.15],

$$\left(\frac{y}{((x+c)^2 + d^2)/2^{2m}} \right) = (-1)^{\frac{d}{2^{m+1}}t + \frac{d}{4}t} \left(\frac{y^{-1}}{c+di} \right)_4.$$

We also have

$$\left(\frac{2^m}{\frac{x+c}{2^m} + \frac{d}{2^m}i} \right)_4 = \left(\frac{2}{\frac{x+c}{2^m} + \frac{d}{2^m}i} \right)_4^m = i^{\frac{x+c}{2^m} \cdot \frac{dm}{2^{m+1}}}$$

and

$$\begin{aligned} \left(\frac{-x(x+c)/2^m}{\frac{x+c}{2^m} + \frac{d}{2^m}i} \right)_4 &= (-1)^{\frac{-x(x+c)/2^{m-1}}{2} \cdot \frac{d}{2^{m+1}}} \left(\frac{\frac{x+c}{2^m} + \frac{d}{2^m}i}{-x(x+c)/2^m} \right)_4 \\ &= (-1)^{\frac{x(x+c)/2^{m+1}}{2} \cdot \frac{d}{2^{m+1}}} \left(\frac{\frac{x+c}{2^m} + \frac{d}{2^m}i}{-x} \right)_4 \left(\frac{\frac{x+c}{2^m} + \frac{d}{2^m}i}{(x+c)/2^m} \right)_4 \\ &= (-1)^{\frac{x(x+c)/2^{m+1}}{2} \cdot \frac{d}{2^{m+1}}} \left(\frac{x+c+di}{x} \right)_4 \left(\frac{\frac{d}{2^m}i}{(x+c)/2^m} \right)_4 \\ &= (-1)^{\frac{x(x+c)/2^{m+1}}{2} \cdot \frac{d}{2^{m+1}}} \left(\frac{c+di}{x} \right)_4 \left(\frac{i}{(x+c)/2^m} \right)_4 \\ &= (-1)^{\frac{x(x+c)/2^{m+1}}{2} \cdot \frac{d}{2^{m+1}}} \left(\frac{x}{c+di} \right)_4 (-1)^{\frac{(\frac{x+c}{2^m})^2 - 1}{8}}. \end{aligned}$$

Thus,

$$(4.1) \quad \begin{aligned} i^k &= \left(\frac{d - (x+c)i}{q} \right)_4 = (-1)^{\frac{q^2-1}{8} + \frac{q-1}{2} \cdot \frac{d}{2^{m+1}}} i^{\frac{x+c}{2^m} \cdot \frac{dm}{2^{m+1}}} \\ &\times (-1)^{\frac{x(x+c)/2^{m+1}}{2} \cdot \frac{d}{2^{m+1}} + \frac{(\frac{x+c}{2^m})^2 - 1}{8}} (-1)^{\frac{d}{2^{m+1}}t + \frac{d}{4}t} \left(\frac{x/y}{c+di} \right)_4. \end{aligned}$$

As $2qy^2 = d^2 - (x+c)^2 + 2c(x+c)$ we have

$$(4.2) \quad q \frac{y^2}{2^m} = 2^{2r-m-1} d_0^2 - 2^{m-1} \left(\frac{x+c}{2^m} \right)^2 + c \cdot \frac{x+c}{2^m}.$$

Suppose $x \equiv 1 \pmod{4}$. Then $m = 1 < r$. From (4.2) we see that $2^{2t-1}q \equiv 2^{2r-2} - 1 + c \cdot \frac{x+c}{2} \pmod{8}$ and so $\frac{x+c}{2} \equiv c(2^{2t-1}q - 2^{2r-2} + 1) \pmod{8}$. If $8 \nmid d$, then $r = 2$ and $\frac{x+c}{2} \equiv c(2^{2t-1}q - 3) \pmod{8}$. Thus,

$$\begin{aligned} (-1)^{\frac{(\frac{x+c}{2})^2 - 1}{8}} &= (-1)^{\frac{c^2-1}{8} + \frac{(2^{2t-1}q-3)^2-1}{8}} = (-1)^{\frac{c^2-1}{8} + \frac{(2^{2t-2}q-2)(2^{2t-2}q-1)}{2}} \\ &= (-1)^{\frac{c^2-1}{8} + 1 + \frac{q+1}{2} \cdot \frac{y}{2}} = (-1)^{\frac{p-1}{8} + 1 + \frac{q+1}{2} \cdot \frac{y}{2}} \end{aligned}$$

and

$$(-1)^{\frac{(x+c)/2+1}{2} \cdot \frac{x+c}{2}} = (-1)^{2^{2t-2}q-1} i^{2^{2t-1}q-3} = -(-1)^{2^{2t-2}} \cdot (-1)^{2^{2t-2}} i = -i.$$

Hence, from (4.1) we deduce that

$$\begin{aligned} i^k &= (-1)^{\frac{q^2-1}{8} + \frac{q-1}{2}} i^{\frac{x+c}{2}} (-1)^{\frac{(x+c)/2+1}{2}} \cdot (-1)^{\frac{(\frac{x+c}{2})^2-1}{8}} \left(\frac{x/y}{c+di} \right)_4 \\ &= (-1)^{\frac{q^2-1}{8} + \frac{q-1}{2}} (-i) (-1)^{\frac{p-1}{8} + 1 + \frac{q+1}{2} \cdot \frac{y}{2}} \left(\frac{x/y}{c+di} \right)_4. \end{aligned}$$

That is,

$$\left(\frac{x/y}{c+di} \right)_4 = (-1)^{\frac{q^2-1}{8} + \frac{q-1}{2} + \frac{p-1}{8} + \frac{q+1}{2} \cdot \frac{y}{2}} i^{k-1}.$$

Now applying Lemma 2.8 we obtain the result.

If $8 \mid d$, from (4.2) we see that $2^{2t-1}q \equiv 2q(\frac{y}{2})^2 \equiv -1 + c \cdot \frac{x+c}{2} \pmod{8}$ and so $\frac{x+c}{2} \equiv c(2^{2t-1}q + 1) \pmod{8}$. Thus,

$$\begin{aligned} (-1)^{\frac{(\frac{x+c}{2})^2-1}{8}} &= (-1)^{\frac{c^2-1}{8} + \frac{(2^{2t-1}q+1)^2-1}{8}} = (-1)^{\frac{c^2+d^2-1}{8} + \frac{2^{2t-2}q(2^{2t-2}q+1)}{2}} \\ &= (-1)^{\frac{p-1}{8} + \frac{q+1}{2} \cdot \frac{y}{2}}. \end{aligned}$$

Now, from (4.1) and the above we derive that

$$\begin{aligned} i^k &= (-1)^{\frac{q^2-1}{8}} i^{\frac{x+c}{2} \cdot \frac{d}{4}} (-1)^{\frac{(\frac{x+c}{2})^2-1}{8}} \left(\frac{x/y}{c+di} \right)_4 \\ &= (-1)^{\frac{q^2-1}{8} + \frac{d}{8} + \frac{p-1}{8} + \frac{q+1}{2} \cdot \frac{y}{2}} \left(\frac{x/y}{c+di} \right)_4. \end{aligned}$$

Now applying Lemma 2.8 we obtain

$$(-q)^{\frac{p-1}{8}} \equiv (-1)^{\frac{q^2-1}{8} + \frac{q+1}{2} \cdot \frac{y}{2}} \left(\frac{d}{c} \right)^k = \begin{cases} (-1)^{\frac{q-1}{4} + \frac{d}{4} + \frac{y}{2}} \left(\frac{d}{c} \right)^k \pmod{p} & \text{if } 4 \mid q-1, \\ \left(\frac{d}{c} \right)^k \pmod{p} & \text{if } 4 \mid q-3. \end{cases}$$

This is the result.

Now assume $r > m \geq 2$. Then $x \equiv 3 \pmod{4}$, $2r - m - 1 \geq 2(m+1) - m - 1 = m+1 \geq 3$ and so $q \frac{y^2}{2^m} \equiv -2^{m-1} + c \cdot \frac{x+c}{2^m} \pmod{8}$. Hence $2^m \parallel y^2$, $m = 2t$ and so $q \equiv -2^{m-1} + c \cdot \frac{x+c}{2^m} \pmod{8}$. That is, $\frac{x+c}{2^m} \equiv c(2^{m-1} + q) \pmod{8}$. Thus,

$$\begin{aligned} &(-1)^{\frac{q^2-1}{8} + \frac{q-1}{2} \cdot \frac{d}{2^{m+1}}} (-1)^{\frac{x(x+c)/2^{m+1}}{2} \cdot \frac{d}{2^{m+1}} + \frac{(\frac{x+c}{2^m})^2-1}{8}} \cdot (-1)^{\frac{d}{2^{m+1}}t + \frac{d}{4}t} i^{\frac{x+c}{2^m} \cdot \frac{dm}{2^{m+1}}} \\ &= (-1)^{\frac{q^2-1}{8} + \frac{q-1}{2} \cdot \frac{d}{2^{m+1}}} (-1)^{\frac{-(2^{m-1}+q)+1}{2} \cdot 2^{r-m-1}d_0 + \frac{c^2(2^{m-1}+q)^2-1}{8}} (-1)^{\frac{d}{2^{m+1}}t + \frac{d}{4}t} (-1)^{\frac{dt}{2^{m+1}}} \\ &= (-1)^{\frac{q^2-1}{8} + \frac{q-1}{2} \cdot 2^{r-m-1}d_0} \cdot (-1)^{(2^{m-2} + \frac{q-1}{2})2^{r-m-1} + \frac{c^2-1}{8} + \frac{(2^{m-1}+q)^2-1}{8}} \cdot (-1)^{\frac{d}{4}t} \\ &= (-1)^{\frac{q^2-1}{8} + \frac{q-1}{2} \cdot 2^{r-m-1}} \cdot (-1)^{(2^{m-2} + \frac{q-1}{2})2^{r-m-1} + \frac{p-1}{8} + \frac{q^2-1}{8} + 2^{m-3}(2^{m-2}+q)} \\ &= (-1)^{2^{r-3} + \frac{p-1}{8} + 2^{m-3}(2^{m-2}+q)} = \begin{cases} (-1)^{\frac{d}{8} + \frac{p-1}{8} + \frac{q+1}{2}} & \text{if } m = 2, \\ (-1)^{\frac{p-1}{8}} & \text{if } m > 2. \end{cases} \end{aligned}$$

Hence, from (4.1) and the above we get

$$\left(\frac{x/y}{c+di}\right)_4 = \begin{cases} (-1)^{\frac{d}{8} + \frac{p-1}{8} + \frac{q+1}{2}} i^k & \text{if } r > m = 2, \\ (-1)^{\frac{p-1}{8}} i^k & \text{if } r > m > 2. \end{cases}$$

Now applying Lemma 2.8 and the fact $m = 2t$ we obtain

$$\begin{aligned} (-q)^{\frac{p-1}{8}} &\equiv \begin{cases} (-1)^{\frac{q+1}{2}} \left(\frac{d}{c}\right)^k \pmod{p} & \text{if } r > m = 2, \\ (-1)^{\frac{d}{8}} \left(\frac{d}{c}\right)^k = \left(\frac{d}{c}\right)^k \pmod{p} & \text{if } r > m > 2 \end{cases} \\ &= (-1)^{\frac{q+1}{2} \cdot \frac{y}{2}} \left(\frac{d}{c}\right)^k = (-1)^{\frac{q+1}{2} \left(\frac{d}{4} + \frac{y}{2}\right)} \left(\frac{d}{c}\right)^k \pmod{p} \quad \text{for } r > m \geq 2. \end{aligned}$$

This yields the result.

Case 2. $m = r$. As $p \equiv 1 \pmod{8}$ we have $4 \mid d$, $m = r \geq 2$, $4 \mid x + c$ and so $x \equiv 3 \pmod{4}$. It is clear that

$$\begin{aligned} (4.3) \quad i^k &= \left(\frac{-d/(x+c) + i}{q}\right)_4 = \left(\frac{-\frac{d}{2^r} + \frac{x+c}{2^r}i}{q}\right)_4 \\ &= \left(\frac{1+i}{q}\right)_4 \left(\frac{\frac{x+c-d}{2^{r+1}} + \frac{x+c+d}{2^{r+1}}i}{q}\right)_4 = \left(\frac{1+i}{q}\right)_4 \left(\frac{\frac{x+c-d}{2^{r+1}} - \frac{x+c+d}{2^{r+1}}i}{q}\right)_4^{-1} \\ &= \left(\frac{i}{q}\right)_4 \left(\frac{\frac{x+c}{2^r} + \frac{d}{2^r}i}{q}\right)_4 = \left(\frac{2}{q}\right) \left(\frac{1+i}{q}\right)_4 \left(\frac{\frac{x+c+d}{2^{r+1}} - \frac{x+c-d}{2^{r+1}}i}{q}\right)_4. \end{aligned}$$

As $2^r \parallel x+c$ and $2^r \parallel d$ we have $2^{r+1} \mid x+c \pm d$ and $\frac{d+x+c}{2^r} \cdot \frac{d-x-c}{2^r} = \left(\frac{d}{2^r}\right)^2 - \left(\frac{x+c}{2^r}\right)^2 \equiv 0 \pmod{8}$. Since $2qy^2 = (d+x+c)(d-x-c) + 2c(x+c)$ and $r \geq 2$ we see that

$$(4.4) \quad c \frac{x+c}{2^r} = \frac{qy^2}{2^r} - \frac{(d+x+c)(d-x-c)}{2^{r+1}} \equiv q \frac{y^2}{2^r} \pmod{2^{r+2}}.$$

Thus, $r = 2t$, $(-1)^{\frac{d}{4} + \frac{y}{2}} = 1$ and $\frac{x+c}{2^r} \equiv cq \pmod{8}$. Hence

$$\frac{x+c}{2^r} \equiv q \equiv (-1)^{\frac{q-1}{2}} \equiv (-1)^{\frac{q-1}{2}} d_0 \equiv (-1)^{\frac{q-1}{2}} \frac{d}{2^r} \pmod{4}.$$

Set

$$A = \frac{x+c + (-1)^{\frac{q-1}{2}} d}{2^{r+1}} \quad \text{and} \quad B = \frac{x+c - (-1)^{\frac{q-1}{2}} d}{2^{r+1}}.$$

Then

$$2 \nmid A, \quad 2 \mid B \quad \text{and} \quad A^2 + B^2 = \frac{(x+c)^2 + d^2}{2^{2r+1}}.$$

Thus,

$$\begin{aligned}
\left(\frac{A-Bi}{q}\right)_4 &= (-1)^{\frac{q-1}{2} \cdot \frac{B}{2}} \left(\frac{q}{A-Bi}\right)_4 = (-1)^{\frac{q-1}{2} \cdot \frac{B}{2}} \left(\frac{qy^2}{A-Bi}\right)_4 \left(\frac{y^2}{A-Bi}\right)_4 \\
&= (-1)^{\frac{q-1}{2} \cdot \frac{B}{2}} \left(\frac{((x+c)^2 + d^2)/2 - x(x+c)}{A-Bi}\right)_4 \left(\frac{y}{A^2+B^2}\right)_4 \\
&= (-1)^{\frac{q-1}{2} \cdot \frac{B}{2}} \left(\frac{-2^r x(x+c)/2^r}{A-Bi}\right)_4 \left(\frac{y}{((x+c)^2 + d^2)/2^{2r+1}}\right)_4 \\
&= (-1)^{\frac{q-1}{2} \cdot \frac{B}{2}} \left(\frac{2}{A-Bi}\right)_4^r \left(\frac{-x(x+c)/2^r}{A-Bi}\right)_4 \left(\frac{2^t y_0}{((x+c)^2 + d^2)/2^{2r+1}}\right)_4 \\
&= (-1)^{\frac{q-1}{2} \cdot \frac{B}{2}} i^{-\frac{AB}{2}r} (-1)^{\frac{-x(x+c)/2^r-1}{2} \cdot \frac{B}{2}} \left(\frac{A-Bi}{-x(x+c)/2^r}\right)_4 \\
&\quad \times (-1)^{\frac{\frac{1}{2}((\frac{x+c}{2^r})^2 + (\frac{d}{2^r})^2) - 1}{4}} t \left(\frac{((x+c)^2 + d^2)/2^{2r+1}}{y_0}\right)_4.
\end{aligned}$$

Since $x \equiv 3 \pmod{4}$ and $\frac{x+c}{2^r} \equiv q \pmod{4}$ we have $(-1)^{\frac{-x(x+c)/2^r-1}{2}} = (-1)^{\frac{q-1}{2}}$. As $\frac{x+c}{2^r} \equiv cq \pmod{8}$, we see that $\frac{B}{2} = \frac{1}{4}(x+c - (-1)^{\frac{q-1}{2}} \frac{d}{2^r}) \equiv \frac{cq - (-1)^{\frac{q-1}{2}} d_0}{4} \pmod{2}$ and

$$\begin{aligned}
(-1)^{\frac{\frac{1}{2}((\frac{x+c}{2^r})^2 + (\frac{d}{2^r})^2) - 1}{4}} &= (-1)^{\frac{(\frac{x+c}{2^r})^2 - 1}{8} + \frac{(\frac{d}{2^r})^2 - 1}{8}} = (-1)^{\frac{(cq)^2 - 1}{8} + \frac{d_0^2 - 1}{8}} \\
&= (-1)^{\frac{c^2 - 1}{8} + \frac{q^2 - 1}{8} + \frac{d_0^2 - 1}{8}} = (-1)^{\frac{p-1}{8} + \frac{q^2 - 1}{8} + \frac{d_0^2 - 1}{8}}.
\end{aligned}$$

Recall that $r = 2t$. From the above we deduce that

$$\begin{aligned}
&\left(\frac{A-Bi}{q}\right)_4 \\
&= (-1)^{\frac{AB}{2}t} \cdot (-1)^{(\frac{p-1}{8} + \frac{q^2-1}{8} + \frac{d_0^2-1}{8})t} \left(\frac{A-Bi}{x(x+c)/2^r}\right)_4 \left(\frac{((x+c)^2 + d^2)/2}{y_0}\right)_4 \\
&= (-1)^{(\frac{cq - (-1)^{(q-1)/2} d_0}{4} + \frac{p-1}{8} + \frac{q^2-1}{8} + \frac{d_0-1}{4})t} \left(\frac{A-Bi}{x(x+c)/2^r}\right)_4 \left(\frac{((x+c)^2 + d^2)/2}{y_0}\right)_4 \\
&= (-1)^{(\frac{cq - (-1)^{(q-1)/2}}{4} + \frac{p-1}{8} + \frac{q^2-1}{8})t} \\
&\quad \times \left(\frac{x+c + (-1)^{\frac{q-1}{2}} d - (x+c - (-1)^{\frac{q-1}{2}} d)i}{x(x+c)/2^r}\right)_4 \left(\frac{qy^2 + x(x+c)}{y_0}\right)_4 \\
&= (-1)^{(\frac{c-1}{4} + \frac{q - (-1)^{(q-1)/2}}{4} + \frac{p-1}{8} + \frac{q^2-1}{8})t} \\
&\quad \times \left(\frac{c + (-1)^{\frac{q-1}{2}} d - (c - (-1)^{\frac{q-1}{2}} d)i}{x}\right)_4 \left(\frac{(-1)^{\frac{q-1}{2}} d + (-1)^{\frac{q-1}{2}} di}{(x+c)/2^r}\right)_4 \left(\frac{x(x+c)}{y_0}\right)_4 \\
&= \left(\frac{-i(1+i)(c + (-1)^{\frac{q-1}{2}} di)}{x}\right)_4 \left(\frac{1+i}{(x+c)/2^r}\right)_4 \left(\frac{x(x+c)}{y_0}\right)_4.
\end{aligned}$$

Recall that $\frac{x+c}{2^r} \equiv cq \pmod{8}$. From the above and Lemma 2.7 we deduce that

$$\begin{aligned} \left(\frac{A-Bi}{q}\right)_4 &= \left(\frac{i}{x}\right)_4 \left(\frac{1+i}{x}\right)_4 \left(\frac{c+(-1)^{\frac{q-1}{2}}di}{x}\right)_4 i^{\frac{(-1)^{\frac{(x+c)/2^r-1}{2}} \frac{x+c}{2^r}-1}{4}} \left(\frac{x(x+c)}{y_0}\right) \\ &= (-1)^{\frac{x+1}{4}} \cdot i^{\frac{-x-1}{4}} \left(\frac{x}{c+(-1)^{\frac{q-1}{2}}di}\right)_4 i^{\frac{(-1)^{(q-1)/2} \frac{x+c}{2^r}-1}{4}} \left(\frac{d+ci}{y_0}\right)_4 \end{aligned}$$

Using (4.4) we see that

$$\begin{aligned} i^{\frac{(-1)^{\frac{q-1}{2}} \frac{x+c}{2^r}-1}{4}} &= i^{\frac{(-1)^{\frac{q-1}{2}} c \frac{x+c}{2^r}-c}{4}} = i^{\frac{(-1)^{\frac{q-1}{2}} qy_0^2-c}{4}} = i^{\frac{(-1)q(y_0^2-1)+(-\frac{1}{q})-c}{4}} \\ &= (-1)^{\frac{y_0^2-1}{8}} i^{\frac{(-\frac{1}{q})q-c}{4}} = \left(\frac{i}{y_0}\right)_4 i^{\frac{(-\frac{1}{q})q-c}{4}}. \end{aligned}$$

Also,

$$i^{-\frac{x+1}{4}} = i^{-\frac{x+c}{4}} \cdot i^{\frac{c-1}{4}} = \begin{cases} i^{\frac{c-1}{4}} & \text{if } r > 2, \\ i^{-q+\frac{c-1}{4}} = (-1)^{\frac{q+1}{2}} i^{1+\frac{c-1}{4}} & \text{if } r = 2 \end{cases}$$

and

$$\begin{aligned} \left(\frac{i}{y_0}\right)_4 \left(\frac{d+ci}{y_0}\right)_4 &= \left(\frac{c-di}{y_0}\right)_4 = \left(\frac{y_0}{c-di}\right)_4 = \left(\frac{2}{c-di}\right)_4^{-t} \left(\frac{y}{c+di}\right)_4 \\ &= i^{\frac{d}{2}t} \left(\frac{y}{c-di}\right)_4 = (-1)^{\frac{d}{4} \cdot \frac{t}{2}} \left(\frac{y}{c-di}\right)_4 \\ &= (-1)^{\frac{d}{4}} \left(\frac{y}{c+di}\right)_4^{-1} = (-1)^{\frac{d}{4}} \left(\frac{y^{-1}}{c+di}\right)_4. \end{aligned}$$

Thus,

$$\begin{aligned} \left(\frac{A-Bi}{q}\right)_4 &= (-1)^{\frac{x+1}{4}} i^{-\frac{x+1}{4} + \frac{(-\frac{1}{q})q-c}{4}} \left(\frac{x}{c+(-1)^{\frac{q-1}{2}}di}\right)_4 (-1)^{\frac{d}{4}} \left(\frac{y^{-1}}{c+di}\right)_4 \\ &= (-1)^{\frac{x+1}{4} + \frac{d}{4}} i^{-\frac{x+c}{4} + \frac{(-\frac{1}{q})q-1}{4}} \left(\frac{x}{c+(-1)^{\frac{q-1}{2}}di}\right)_4 \left(\frac{y^{-1}}{c+di}\right)_4. \end{aligned}$$

For $r = 2$ we have $\frac{x+c}{4} \equiv q \pmod{4}$ and so $i^{-\frac{x+c}{4}} = i^{-q} = (-1)^{\frac{q+1}{2}} i$. For $r > 2$ we have $r = 2t \geq 4$, $2^4 \mid x+c$ and so $i^{-\frac{x+c}{4}} = 1$. If $q \equiv 1 \pmod{4}$, from (4.3) and the above we get

$$\begin{aligned} i^k &= \left(\frac{2}{q}\right) \left(\frac{1+i}{q}\right)_4 \left(\frac{A-Bi}{q}\right)_4 \\ &= (-1)^{\frac{q-1}{4}} i^{\frac{q-1}{4}} (-1)^{\frac{x+1}{4} + \frac{d}{4}} i^{-\frac{x+c}{4} + \frac{q-1}{4}} \left(\frac{x}{c+di}\right)_4 \left(\frac{y^{-1}}{c+di}\right)_4 \\ &= (-1)^{\frac{x+1}{4} + \frac{d}{4}} i^{-\frac{x+c}{4}} \left(\frac{x/y}{c+di}\right)_4 = (-1)^{\frac{x+c}{4} + \frac{d}{4} + \frac{1-c}{4}} i^{-\frac{x+c}{4}} \left(\frac{x/y}{c+di}\right)_4 \\ &= (-1)^{\frac{1-c}{4}} i^{-\frac{x+c}{4}} \left(\frac{x/y}{c+di}\right)_4 = (-1)^{\frac{p-1}{8}} i^{-\frac{x+c}{4}} \left(\frac{x/y}{c+di}\right)_4. \end{aligned}$$

and so

$$\left(\frac{x/y}{c+di}\right)_4 = \begin{cases} (-1)^{\frac{p-1}{8}} i^{k+1} & \text{if } r = 2, \\ (-1)^{\frac{p-1}{8}} i^k & \text{if } r > 2. \end{cases}$$

Now applying Lemma 2.8 we deduce the result. If $q \equiv 3 \pmod{4}$, then $\left(\frac{y}{c^2+d^2}\right) = \left(\frac{y_0}{p}\right) = \left(\frac{p}{y_0}\right) = \left(\frac{x^2+2qy^2}{y_0}\right) = 1$. From (4.3) and the above we deduce that

$$\begin{aligned} \left(\frac{A-Bi}{q}\right)_4 &= (-1)^{\frac{x+1}{4}+\frac{d}{4}} i^{-\frac{x+c}{4}+\frac{-q-1}{4}} \left(\frac{x}{c-di}\right)_4 \left(\frac{y^{-1}}{c+di}\right)_4 \\ &= (-1)^{\frac{x+c}{4}+\frac{d}{4}+\frac{1-c}{4}} i^{-\frac{x+c}{4}-\frac{q+1}{4}} \left(\frac{x}{c+di}\right)_4^{-1} \left(\frac{y}{c+di}\right)_4 \left(\frac{y^{-2}}{c+di}\right)_4 \\ &= (-1)^{\frac{1-c}{4}} i^{-\frac{x+c}{4}-\frac{q+1}{4}} \left(\frac{x^{-1}y}{c+di}\right)_4 \left(\frac{y^2}{c+di}\right)_4^{-1} \\ &= (-1)^{\frac{c^2-1}{8}} i^{-\frac{x+c}{4}-\frac{q+1}{4}} \left(\frac{x/y}{c+di}\right)_4^{-1} \left(\frac{y}{c^2+d^2}\right)^{-1} \\ &= (-1)^{\frac{p-1}{8}} i^{-\frac{x+c}{4}-\frac{q+1}{4}} \left(\frac{x/y}{c+di}\right)_4^{-1}. \end{aligned}$$

Thus, using (4.3) we see that

$$i^k = \left(\frac{1+i}{q}\right)_4 \left(\frac{A-Bi}{q}\right)_4^{-1} = i^{\frac{-q-1}{4}} \cdot (-1)^{\frac{p-1}{8}} i^{\frac{x+c}{4}+\frac{q+1}{4}} \left(\frac{x/y}{c+di}\right)_4$$

and so

$$\left(\frac{x/y}{c+di}\right)_4 = (-1)^{\frac{p-1}{8}} i^{k-\frac{x+c}{4}} = \begin{cases} (-1)^{\frac{p-1}{8}} i^{k-q} = (-1)^{\frac{p-1}{8}} i^{k+1} & \text{if } r = 2, \\ (-1)^{\frac{p-1}{8}} i^k & \text{if } r > 2. \end{cases}$$

Applying Lemma 2.8 we deduce the result.

Case 3. $m > r$. Using Lemmas 2.1-2.5 we see that

$$\begin{aligned} i^k &= \left(\frac{-d/(x+c)+i}{q}\right)_4 = \left(\frac{\frac{d}{2^r} - \frac{x+c}{2^r}i}{q}\right)_4 = (-1)^{\frac{q-1}{2} \cdot \frac{x+c}{2^{r+1}}} \left(\frac{q}{\frac{d}{2^r} - \frac{x+c}{2^r}i}\right)_4 \\ &= (-1)^{\frac{q-1}{2} \cdot \frac{x+c}{2^{r+1}}} \left(\frac{qy^2}{\frac{d}{2^r} - \frac{x+c}{2^r}i}\right)_4 \left(\frac{y^2}{\frac{d}{2^r} - \frac{x+c}{2^r}i}\right)_4 \\ &= (-1)^{\frac{q-1}{2} \cdot \frac{x+c}{2^{r+1}}} \left(\frac{((x+c)^2+d^2)/2 - x(x+c)}{\frac{d}{2^r} - \frac{x+c}{2^r}i}\right)_4 \left(\frac{y}{\left(\frac{d}{2^r}\right)^2 + \left(\frac{x+c}{2^r}\right)^2}\right) \\ &= (-1)^{\frac{q-1}{2} \cdot \frac{x+c}{2^{r+1}}} \left(\frac{-2^m x(x+c)/2^m}{\frac{d}{2^r} - \frac{x+c}{2^r}i}\right)_4 \left(\frac{y}{((x+c)^2+d^2)/2^{2r}}\right) \\ &= (-1)^{\frac{q-1}{2} \cdot \frac{x+c}{2^{r+1}}} \left(\frac{2}{\frac{d}{2^r} - \frac{x+c}{2^r}i}\right)_4^m \left(\frac{-x(x+c)/2^m}{\frac{d}{2^r} - \frac{x+c}{2^r}i}\right)_4 \left(\frac{2^t y_0}{((x+c)^2+d^2)/2^{2r}}\right) \\ &= (-1)^{\frac{q-1}{2} \cdot \frac{x+c}{2^{r+1}}} i^{-\frac{d}{2^r} \cdot \frac{x+c}{2^{r+1}} \cdot m} (-1)^{\frac{-x(x+c)/2^m - 1}{2} \cdot \frac{x+c}{2^{r+1}}} \\ &\quad \times \left(\frac{\frac{d}{2^r} - \frac{x+c}{2^r}i}{-x(x+c)/2^m}\right)_4 (-1)^{\frac{x+c}{2^{r+1}}t} \left(\frac{(x+c)^2+d^2}{y_0}\right). \end{aligned}$$

Applying Lemma 2.7 we derive that

$$\begin{aligned}
& \left(\frac{\frac{d}{2r} - \frac{x+c}{2^r}i}{-x(x+c)/2^m} \right)_4 \left(\frac{(x+c)^2 + d^2}{y_0} \right) \\
&= \left(\frac{d - (x+c)i}{-x} \right)_4 \left(\frac{d - (x+c)i}{(x+c)/2^m} \right)_4 \left(\frac{2qy^2 + 2x(c+x)}{y_0} \right) \\
&= \left(\frac{d-ci}{-x} \right)_4 \left(\frac{d}{(x+c)/2^m} \right)_4 \left(\frac{2x(c+x)}{y_0} \right) \\
&= \left(\frac{-i}{x} \right)_4 \left(\frac{c+di}{x} \right)_4 \left(\frac{2}{y_0} \right) \left(\frac{d+ci}{y_0} \right)_4 = (-1)^{\frac{x+1}{4}} \left(\frac{x}{c+di} \right)_4 \left(\frac{i(d+ci)}{y_0} \right)_4 \\
&= (-1)^{\frac{x+1}{4}} \left(\frac{x}{c+di} \right)_4 \left(\frac{c-di}{y_0} \right)_4 = (-1)^{\frac{x+1}{4}} \left(\frac{x}{c+di} \right)_4 \left(\frac{y_0}{c-di} \right)_4 \\
&= (-1)^{\frac{x+1}{4}} \left(\frac{x}{c+di} \right)_4 \left(\frac{y_0}{c+di} \right)_4^{-1} = (-1)^{\frac{x+c}{4} + \frac{1-c}{4}} \left(\frac{x}{c+di} \right)_4 \left(\frac{2^t/y}{c+di} \right)_4 \\
&= (-1)^{\frac{c-1}{4} + \frac{d}{4}t} \left(\frac{x/y}{c+di} \right)_4 = (-1)^{\frac{p-1}{8} + \frac{d}{4}t} \left(\frac{x/y}{c+di} \right)_4.
\end{aligned}$$

Therefore,

$$\begin{aligned}
i^k &= (-1)^{\frac{q-1}{2} \cdot \frac{x+c}{2^{r+1}}} i^{-\frac{x+c}{2^{r+1}} \cdot m} (-1)^{\frac{x(x+c)/2^{m+1} \cdot \frac{x+c}{2^{r+1}}}{2}} \cdot (-1)^{\frac{x+c}{2^{r+1}}t + \frac{p-1}{8} + \frac{d}{4}t} \left(\frac{x/y}{c+di} \right)_4 \\
&= \begin{cases} (-1)^{\frac{q-1}{2} + \frac{x(x+c)/2^{r+1} + 1}{2} + \frac{p-1}{8} + (1+\frac{d}{4})t} i^{-\frac{x+c}{2^{r+1}}(r+1)} \left(\frac{x/y}{c+di} \right)_4 & \text{if } m = r+1, \\ (-1)^{m \frac{x+c}{2^{r+2}} + \frac{p-1}{8} + \frac{d}{4}t} \left(\frac{x/y}{c+di} \right)_4 & \text{if } m > r+1. \end{cases}
\end{aligned}$$

Since $2^t \parallel y$, $2^r \parallel d$, $2^m \parallel x+c$, $m > r \geq 2$ and $2qy^2 - d^2 = (x+c)^2 - 2x(x+c)$, we see that $2^{m+1} \parallel 2qy^2 - d^2$. That is, $2^{m+1} \parallel 2^{2t+1}qy_0^2 - 2^{2r}d_0^2$. Hence $m+1 = \min\{2t+1, 2r\} \leq 2r$ and so $t > 1$. If $m+1 = 2r$, then $t \geq r$ and

$$\begin{aligned}
x \frac{x+c}{2^m} &= 2^{m-1} \left(\frac{x+c}{2^m} \right)^2 - 2^{2t+1-2r} qy_0^2 + d_0^2 \\
&\equiv 1 - 2^{2t+1-2r} q \equiv \begin{cases} 1 \pmod{4} & \text{if } t > r, \\ 3 \pmod{4} & \text{if } t = r. \end{cases}
\end{aligned}$$

If $m+1 < 2r$, then $m = 2t$, $t < r$ and

$$\begin{aligned}
x \frac{x+c}{2^m} &= 2^{m-1} \left(\frac{x+c}{2^m} \right)^2 - qy_0^2 + 2^{2r-1-2t} d_0^2 \\
&\equiv -q + 2^{2r-2t-1} \equiv \begin{cases} -q \pmod{4} & \text{if } t < r-1, \\ 2-q \pmod{4} & \text{if } t = r-1. \end{cases}
\end{aligned}$$

For $r = 2$ we must have $m = 3$ and so

$$\begin{aligned}
i^k &= (-1)^{\frac{q-1}{2} + \frac{x(x+c)/8+1}{2} + \frac{p-1}{8} + (1+\frac{d}{4})t} i^{-\frac{x+c}{8} \cdot 3} \left(\frac{x/y}{c+di} \right)_4 \\
&= (-1)^{\frac{q-1}{2} + \frac{x(x+c)/8+1}{2} + \frac{p-1}{8}} i^{\frac{x+c}{8}} \left(\frac{x/y}{c+di} \right)_4 = (-1)^{\frac{q-1}{2} + \frac{p-1}{8}} i \left(\frac{x/y}{c+di} \right)_4.
\end{aligned}$$

Now assume $r > 2$. If $m + 1 = 2r$, then $m > r + 1$ and so

$$i^k = (-1)^{\frac{x+c}{2r+2} + \frac{p-1}{8}} \left(\frac{x/y}{c+di} \right)_4 = (-1)^{2r-3 + \frac{p-1}{8}} \left(\frac{x/y}{c+di} \right)_4 = (-1)^{\frac{d}{8} + \frac{p-1}{8}} \left(\frac{x/y}{c+di} \right)_4.$$

If $m + 1 < 2r$ and $m = r + 1$, then $t = \frac{m}{2} = \frac{r+1}{2}$, $x \frac{x+c}{2r+1} \equiv -q + 2^{2r-2t-1} = -q + 2^{r-2} \equiv -q + \frac{d}{4} \pmod{4}$ and so

$$\begin{aligned} i^k &= (-1)^{\frac{q-1}{2} + \frac{x(x+c)/2r+1+1}{2} + \frac{p-1}{8} + t} i^{-\frac{x+c}{2r+1}(r+1)} \left(\frac{x/y}{c+di} \right)_4 \\ &= (-1)^{\frac{q-1}{2} + \frac{1-q+d}{2} + \frac{p-1}{8} + \frac{r+1}{2}} (-1)^{\frac{r+1}{2}} \left(\frac{x/y}{c+di} \right)_4 = (-1)^{\frac{p-1}{8} + \frac{d}{8}} \left(\frac{x/y}{c+di} \right)_4. \end{aligned}$$

If $m + 1 < 2r$ and $m > r + 1$, then $r > 3$, $m = 2t$ and so $i^k = (-1)^{\frac{p-1}{8}} \left(\frac{x/y}{c+di} \right)_4 = (-1)^{\frac{p-1}{8} + \frac{d}{8}} \left(\frac{x/y}{c+di} \right)_4$. Hence

$$\left(\frac{x/y}{c+di} \right)_4 = \begin{cases} (-1)^{\frac{q-1}{2} + \frac{p-1}{8}} i^{k-1} & \text{if } r = 2, \\ (-1)^{\frac{p-1}{8} + \frac{d}{8}} i^k & \text{if } r > 2. \end{cases}$$

Now applying Lemma 2.8 we obtain the result.

By the above, the theorem is proved.

Theorem 4.3. *Let p and q be primes such that $p \equiv 1 \pmod{8}$ and $q \equiv 3 \pmod{4}$. Suppose $p = c^2 + d^2 = x^2 + 2qy^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod{4}$, $d = 2^r d_0$, $d_0 \equiv 1 \pmod{4}$ and $\left(\frac{c-di}{x}\right)^{\frac{q+1}{4}} \equiv i^m \pmod{q}$. Assume $(c, x+d) = 1$ or $(d_0, x+c) = 1$. Then*

$$(-q)^{\frac{p-1}{8}} \equiv (-1)^{\frac{x-1}{2} \cdot \frac{q+1}{4}} \left(\frac{d}{c}\right)^m \pmod{p}.$$

Proof. Clearly $q \nmid x$ and x is odd. We first assume $(c, x+d) = 1$. By the proof of Theorem 4.1, $(q, (x+d)(c^2 + (x+d)^2)) = 1$. It is easily seen that $\frac{c/(x+d)-i}{c/(x+d)+i} = \frac{c-(x+d)i}{c+(x+d)i} \equiv \frac{c-di}{ix} \pmod{q}$. Thus, for $k = 0, 1, 2, 3$, using Lemma 2.6 we get

$$\begin{aligned} \left(\frac{c/(x+d)+i}{q} \right)_4 &= i^k \\ \iff \frac{c}{x+d} \in Q_k(q) &\iff \left(\frac{\frac{c}{x+d}-i}{\frac{c}{x+d}+i} \right)^{\frac{q+1}{4}} \equiv i^k \pmod{q} \\ \iff \left(\frac{c-di}{ix} \right)^{\frac{q+1}{4}} &\equiv i^k \pmod{q} \iff \left(\frac{c-di}{x} \right)^{\frac{q+1}{4}} \equiv i^{\frac{q+1}{4}+k} \pmod{q}. \end{aligned}$$

Since $\left(\frac{c-di}{x}\right)^{\frac{q+1}{4}} \equiv i^m \pmod{q}$, from the above we deduce that

$$\left(\frac{c/(x+d)+i}{q} \right)_4 = i^{m - \frac{q+1}{4}} = \begin{cases} (-1)^{\frac{q+5}{8}} i^{m+1} & \text{if } q \equiv 3 \pmod{8}, \\ (-1)^{\frac{q+1}{8}} i^m & \text{if } q \equiv 7 \pmod{8}. \end{cases}$$

Now, applying Theorem 4.1 we derive the result.

Now we assume $(d_0, x + c) = 1$. By the proof of Theorem 4.2, $(q, x + c) = (q, d^2 + (x + c)^2) = 1$. It is easily seen that $\frac{d+(x+c)i}{d-(x+c)i} \equiv \frac{c-di}{-x} \pmod{q}$. Thus, for $k = 0, 1, 2, 3$, using Lemma 2.6 we get

$$\begin{aligned} \left(\frac{-d/(x+c)+i}{q}\right)_4 &= i^k \\ \iff -\frac{d}{x+c} \in Q_k(q) &\iff \left(\frac{-\frac{d}{x+c}-i}{-\frac{d}{x+c}+i}\right)^{\frac{q+1}{4}} \equiv i^k \pmod{q} \\ \iff \left(\frac{d+(x+c)i}{d-(x+c)i}\right)^{\frac{q+1}{4}} &\equiv i^k \pmod{q} \iff \left(\frac{c-di}{-x}\right)^{\frac{q+1}{4}} \equiv i^k \pmod{q} \\ \iff \left(\frac{c-di}{x}\right)^{\frac{q+1}{4}} &\equiv i^{\frac{q+1}{2}+k} \pmod{q}. \end{aligned}$$

Since $\left(\frac{c-di}{x}\right)^{\frac{q+1}{4}} \equiv i^m \pmod{q}$, from the above we deduce that $\left(\frac{-d/(x+c)+i}{q}\right)_4 = i^{m-\frac{q+1}{2}} = (-1)^{\frac{q+1}{4}} i^m$. Now applying Theorem 4.2 we deduce the result. The proof is now complete.

Corollary 4.1. *Let p and q be primes such that $p \equiv 1 \pmod{8}$ and $q \equiv 3 \pmod{8}$. Suppose $p = c^2 + d^2 = x^2 + 2qy^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod{4}$, $d = 2^r d_0$, $q \mid cd$ and $d_0 \equiv 1 \pmod{4}$. Assume $(c, x + d) = 1$ or $(d_0, x + c) = 1$. Then*

$$(-q)^{\frac{p-1}{8}} \equiv \begin{cases} \pm(-1)^{\frac{x-1}{2}} \pmod{p} & \text{if } x \equiv \pm c \pmod{q}, \\ \mp(-1)^{\frac{q-3}{8}+\frac{x-1}{2}} \frac{d}{c} \pmod{p} & \text{if } x \equiv \pm d \pmod{q}. \end{cases}$$

Proof. If $x \equiv \pm c \pmod{q}$, then $q \mid d$ and so $\left(\frac{c-di}{x}\right)^{\frac{q+1}{4}} \equiv (\pm 1)^{\frac{q+1}{4}} = \pm 1 \pmod{q}$. If $x \equiv \pm d \pmod{q}$, then $q \mid c$ and so $\left(\frac{c-di}{x}\right)^{\frac{q+1}{4}} \equiv (\mp i)^{\frac{q+1}{4}} = \mp(-1)^{\frac{q-3}{8}} i \pmod{q}$. Now applying Theorem 4.3 we deduce the result.

As an example, taking $q = 3$ in Corollary 4.1 we see that if p is a prime of the form $24k + 1$ and so $p = c^2 + d^2 = x^2 + 6y^2$, then

$$(4.5) \quad (-3)^{\frac{p-1}{8}} \equiv \begin{cases} \pm(-1)^{\frac{x-1}{2}} \pmod{p} & \text{if } x \equiv \pm c \pmod{3}, \\ \mp(-1)^{\frac{x-1}{2}} \frac{d}{c} \pmod{p} & \text{if } x \equiv \pm d \pmod{3}. \end{cases}$$

Theorem 4.4. *Let p and q be primes such that $p \equiv 1 \pmod{8}$, $q \equiv 7 \pmod{8}$, $p = c^2 + d^2 = x^2 + 2qy^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod{4}$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod{4}$. Assume $(c, x + d) = 1$ or $(d_0, x + c) = 1$. Then*

$$(-q)^{\frac{p-1}{8}} \equiv \left(\frac{d}{c}\right)^m \pmod{p} \iff \left(\frac{c-di}{c+di}\right)^{\frac{q+1}{8}} \equiv i^m \pmod{q}.$$

Proof. Observe that

$$\left(\frac{c-di}{c+di}\right)^{\frac{q+1}{8}} = \frac{(c-di)^{\frac{q+1}{4}}}{(c^2+d^2)^{\frac{q+1}{8}}} = \frac{(c-di)^{\frac{q+1}{4}}}{(x^2+2qy^2)^{\frac{q+1}{8}}} \equiv \left(\frac{c-di}{x}\right)^{\frac{q+1}{4}} \pmod{q}.$$

The result follows from Theorem 4.3.

We note that if $q \nmid d$, then the m in Theorem 4.4 depends only on $\frac{c}{d} \pmod{q}$.

Corollary 4.2. *Let $p \equiv 1 \pmod{8}$ and $q \equiv 7 \pmod{8}$ be primes such that $p = c^2 + d^2 = x^2 + 2qy^2$ with $c, d, x, y \in \mathbb{Z}$ and $q \mid cd(c^2 - d^2)$. Suppose $c \equiv 1 \pmod{4}$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod{4}$. Assume $(c, x+d) = 1$ or $(d_0, x+c) = 1$. Then*

$$q^{\frac{p-1}{8}} \equiv \begin{cases} (-1)^{\frac{q+1}{8}} \pmod{p} & \text{if } q \mid c, \\ 1 \pmod{p} & \text{if } q \mid d, \\ \pm(-1)^{\frac{q+9}{16}} \frac{d}{c} \pmod{p} & \text{if } 16 \mid (q-7) \text{ and } c \equiv \pm d \pmod{q}, \\ (-1)^{\frac{q+1}{16}} \pmod{p} & \text{if } 16 \mid (q-15) \text{ and } c \equiv \pm d \pmod{q}. \end{cases}$$

Proof. Clearly

$$\frac{c-di}{c+di} \equiv \begin{cases} -1 \pmod{q} & \text{if } q \mid c, \\ 1 \pmod{q} & \text{if } q \mid d, \\ -i \pmod{q} & \text{if } c \equiv d \pmod{q}, \\ i \pmod{q} & \text{if } c \equiv -d \pmod{q}. \end{cases}$$

Thus the result follows from Theorem 4.4.

Theorem 4.5. *Let p and q be distinct primes, $p \equiv 1 \pmod{8}$, $q \equiv 1 \pmod{4}$, $p = c^2 + d^2 = x^2 + 2qy^2$, $q = a^2 + b^2$, $a, b, c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod{4}$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod{4}$. Assume $(c, x+d) = 1$ or $(d_0, x+c) = 1$. Suppose $\left(\frac{ac+bd}{ax}\right)^{\frac{q-1}{4}} \equiv \left(\frac{b}{a}\right)^m \pmod{q}$. Then*

$$(-q)^{\frac{p-1}{8}} \equiv (-1)^{\frac{x-1}{2} \cdot \frac{q-1}{4} + \frac{d}{4} + \frac{y}{2}} \left(\frac{d}{c}\right)^m \pmod{p}.$$

Proof. Clearly $q \nmid x$. We first assume $(c, x+d) = 1$. By the proof of Theorem 4.1, $(q, (x+d)(c^2 + (x+d)^2)) = 1$. It is easily seen that $\frac{ac+b(x+d)}{ac-b(x+d)} \equiv \frac{ac+bd}{ax} \cdot \frac{b}{a} \pmod{q}$. Thus, for $k = 0, 1, 2, 3$, using Lemma 2.6 we get

$$\begin{aligned} \left(\frac{c/(x+d)+i}{q}\right)_4 = i^k &\iff \frac{c}{x+d} \in Q_k(q) \iff \left(\frac{\frac{c}{x+d} + \frac{b}{a}}{\frac{c}{x+d} - \frac{b}{a}}\right)^{\frac{q-1}{4}} \equiv \left(\frac{b}{a}\right)^k \pmod{q} \\ &\iff \left(\frac{ac+b(x+d)}{ac-b(x+d)}\right)^{\frac{q-1}{4}} \equiv \left(\frac{b}{a}\right)^k \pmod{q} \\ &\iff \left(\frac{ac+bd}{ax} \cdot \frac{b}{a}\right)^{\frac{q-1}{4}} \equiv \left(\frac{b}{a}\right)^k \pmod{q} \\ &\iff \left(\frac{ac+bd}{ax}\right)^{\frac{q-1}{4}} \equiv \left(\frac{b}{a}\right)^{k-\frac{q-1}{4}} \pmod{q}. \end{aligned}$$

Since $(\frac{ac+bd}{ax})^{\frac{q-1}{4}} \equiv (\frac{b}{a})^m \pmod{q}$, from the above we get $(\frac{c/(x+d)+i}{q})_4 = i^{m+\frac{q-1}{4}}$. Now the result follows from Theorem 4.1 immediately.

Suppose $(d_0, x+c) = 1$. By the proof of Theorem 4.2, $(q, (x+c)(d^2+(x+c)^2)) = 1$. It is easily seen that $\frac{ad-b(x+c)}{ad+b(x+c)} \equiv \frac{ac+bd}{-ax} \pmod{q}$. Thus, for $k = 0, 1, 2, 3$, using Lemma 2.6 we get

$$\begin{aligned} & \left(\frac{-d/(x+c)+i}{q}\right)_4 = i^k \\ \iff & -\frac{d}{x+c} \in Q_k(q) \iff \left(\frac{-\frac{d}{x+c} + \frac{b}{a}}{-\frac{d}{x+c} - \frac{b}{a}}\right)^{\frac{q-1}{4}} \equiv \left(\frac{b}{a}\right)^k \pmod{q} \\ \iff & \left(\frac{ad-b(x+c)}{ad+b(x+c)}\right)^{\frac{q-1}{4}} \equiv \left(\frac{b}{a}\right)^k \pmod{q} \iff \left(\frac{ac+bd}{-ax}\right)^{\frac{q-1}{4}} \equiv \left(\frac{b}{a}\right)^k \pmod{q} \\ \iff & \left(\frac{ac+bd}{ax}\right)^{\frac{q-1}{4}} \equiv \left(\frac{b}{a}\right)^{\frac{q-1}{2}+k} \pmod{q}. \end{aligned}$$

Since $(\frac{ac+bd}{ax})^{\frac{q-1}{4}} \equiv (\frac{b}{a})^m \pmod{q}$, by the above we get $(\frac{-d/(x+c)+i}{q})_4 = i^{m-\frac{q-1}{2}}$. Now applying Theorem 4.2 we derive the result. The proof is now complete.

Corollary 4.3. *Let $p \equiv 1 \pmod{8}$ and $q \equiv 5 \pmod{8}$ be primes such that $p = c^2 + d^2 = x^2 + 2qy^2$ with $c, d, x, y \in \mathbb{Z}$ and $q \mid cd$. Suppose $c \equiv 1 \pmod{4}$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod{4}$. Assume $(c, x+d) = 1$ or $(d_0, x+c) = 1$. Then*

$$(-q)^{\frac{p-1}{8}} \equiv \begin{cases} \pm(-1)^{\frac{d}{4} + \frac{x-1}{2} + \frac{y}{2}} \pmod{p} & \text{if } x \equiv \pm c \pmod{q}, \\ \pm(-1)^{\frac{q-5}{8} + \frac{d}{4} + \frac{x-1}{2} + \frac{y}{2}} \frac{d}{c} \pmod{p} & \text{if } x \equiv \pm d \pmod{q}. \end{cases}$$

Proof. Suppose $q = a^2 + b^2$ with $a, b \in \mathbb{Z}$. If $x \equiv \pm c \pmod{q}$, then $q \mid d$ and so $(\frac{ac+bd}{ax})^{\frac{q-1}{4}} \equiv (\frac{c}{x})^{\frac{q-1}{4}} \equiv (\pm 1)^{\frac{q-1}{4}} = \pm 1 \pmod{q}$. If $x \equiv \pm d \pmod{q}$, then $q \mid c$ and so $(\frac{ac+bd}{ax})^{\frac{q-1}{4}} \equiv (\frac{bd}{ax})^{\frac{q-1}{4}} \equiv (\pm \frac{b}{a})^{\frac{q-1}{4}} \equiv \pm(-1)^{\frac{q-5}{8}} \frac{b}{a} \pmod{q}$. Now putting the above with Theorem 4.5 we deduce the result.

Theorem 4.6. *Let p and q be distinct primes such that $p \equiv 1 \pmod{8}$, $q \equiv 1 \pmod{8}$, $p = c^2 + d^2 = x^2 + 2qy^2$, $q = a^2 + b^2$, $a, b, c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod{4}$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod{4}$. Assume $(c, x+d) = 1$ or $(d_0, x+c) = 1$. Then*

$$(-q)^{\frac{p-1}{8}} \equiv (-1)^{\frac{d}{4} + \frac{y}{2}} \left(\frac{d}{c}\right)^m \pmod{p} \iff \left(\frac{ac+bd}{ac-bd}\right)^{\frac{q-1}{8}} \equiv \left(\frac{b}{a}\right)^m \pmod{q}.$$

Proof. Observe that $b^2 \equiv -a^2 \pmod{q}$, $p \equiv x^2 \pmod{q}$ and so

$$\left(\frac{ac+bd}{ac-bd}\right)^{\frac{q-1}{8}} = \frac{(ac+bd)^{\frac{q-1}{4}}}{(a^2c^2 - b^2d^2)^{\frac{q-1}{8}}} \equiv \frac{(ac+bd)^{\frac{q-1}{4}}}{(a^2p)^{\frac{q-1}{8}}} \equiv \left(\frac{ac+bd}{ax}\right)^{\frac{q-1}{4}} \pmod{q}.$$

The result follows from Theorem 4.5.

We note that if $q \nmid d$, then the m in Theorem 4.6 depends only on $\frac{c}{d} \pmod{q}$.

Corollary 4.4. *Let p and q be distinct primes of the form $8k + 1$ such that $p = c^2 + d^2 = x^2 + 2qy^2$ with $c, d, x, y \in \mathbb{Z}$ and $q \mid cd(c^2 - d^2)$. Suppose $c \equiv 1 \pmod{4}$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod{4}$. Assume $(c, x + d) = 1$ or $(d_0, x + c) = 1$. Then*

$$(-q)^{\frac{p-1}{8}} \equiv \begin{cases} (-1)^{\frac{q-1}{8} + \frac{d}{4} + \frac{y}{2}} \pmod{p} & \text{if } q \mid c, \\ (-1)^{\frac{d}{4} + \frac{y}{2}} \pmod{p} & \text{if } q \mid d, \\ (-1)^{\frac{q-1}{16} + \frac{d}{4} + \frac{y}{2}} \pmod{p} & \text{if } 16 \mid (q-1) \text{ and } c \equiv \pm d \pmod{q}, \\ \pm(-1)^{\frac{q-9}{16} + \frac{d}{4} + \frac{y}{2}} \frac{d}{c} \pmod{p} & \text{if } 16 \mid (q-9) \text{ and } c \equiv \pm d \pmod{q}. \end{cases}$$

Proof. Suppose that $q = a^2 + b^2$ with $a, b \in \mathbb{Z}$. Then clearly

$$\left(\frac{ac + bd}{ac - bd}\right)^{\frac{q-1}{8}} \equiv \begin{cases} (-1)^{\frac{q-1}{8}} \pmod{q} & \text{if } q \mid c, \\ 1 \pmod{q} & \text{if } q \mid d, \\ (-1)^{\frac{q-1}{16}} \pmod{q} & \text{if } 16 \mid (q-1) \text{ and } c \equiv \pm d \pmod{q}, \\ \pm(-1)^{\frac{q-9}{16}} \frac{b}{a} \pmod{q} & \text{if } 16 \mid (q-9) \text{ and } c \equiv \pm d \pmod{q}. \end{cases}$$

Thus the result follows from Theorem 4.6.

Theorem 4.7. *Let $p \equiv 1 \pmod{8}$ be a prime, $p = c^2 + d^2 = x^2 + 2(a^2 + b^2)y^2$, $a, b, c, d, x, y \in \mathbb{Z}$, $a \neq 0$, $2 \mid a$, $(a, b) = 1$, $c \equiv 1 \pmod{4}$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod{4}$. Assume $(c, x + d) = 1$ or $(d_0, x + c) = 1$. Then*

$$\begin{aligned} (-a^2 - b^2)^{\frac{p-1}{8}} &\equiv \begin{cases} (-1)^{\frac{d}{4} + \frac{y}{2}} \left(\frac{c}{d}\right)^m \pmod{p} & \text{if } 4 \mid a, \\ (-1)^{\frac{b-1}{2} + \frac{d}{4} + \frac{y}{2} + \frac{x-1}{2}} \left(\frac{c}{d}\right)^{m-1} \pmod{p} & \text{if } 4 \mid a - 2. \end{cases} \\ \iff \left(\frac{(ac + bd)/x}{b + ai}\right)_4 &= i^m. \end{aligned}$$

Proof. Suppose $q = a^2 + b^2$ and $\left(\frac{ac+bd}{b+ai}\right)_4 = i^m$. Then clearly $q \equiv 1 \pmod{4}$ and $p \nmid q$. We first assume $(c, x + d) = 1$. By the proof of Theorem 4.1, $(q, x + d) = (q, c^2 + (x + d)^2) = 1$. Since $\frac{c-(x+d)i}{c+(x+d)i} \equiv \frac{c-di}{ix} \pmod{q}$, from [S5, p.24] we know that

$$\left(\frac{c/(x+d) + i}{q}\right)_4 = (-1)^{\frac{b+1}{2} \cdot \frac{a}{2} + \lceil \frac{q}{8} \rceil} i^{-m}.$$

This together with Theorem 4.1 yields the result in this case.

Now we assume $(d_0, x + c) = 1$. By the proof of Theorem 4.2, $(q, x + c) = (q, (x + c)^2 + d^2) = 1$. Since $\frac{d+(x+c)i}{d-(x+c)i} \equiv \frac{c-di}{-x} \pmod{q}$, from [S5, p.24] we know that

$$\left(\frac{-d/(x+c) + i}{q}\right)_4 = \begin{cases} (-1)^{\frac{b+1}{2}} i^{1-m} & \text{if } 4 \mid a - 2, \\ i^{-m} & \text{if } 4 \mid a. \end{cases}$$

Now applying Theorem 4.2 we deduce the result in this case. So the theorem is proved.

Corollary 4.5. *Let $p \equiv 1, 9 \pmod{40}$ be a prime and so $p = c^2 + d^2 = x^2 + 10y^2$ with $c, d, x, y \in \mathbb{Z}$. Suppose $c \equiv 1 \pmod{4}$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod{4}$. Assume $(c, x + d) = 1$ or $(d_0, x + c) = 1$. Then*

$$(-5)^{\frac{p-1}{8}} \equiv \begin{cases} \pm(-1)^{\frac{d}{4} + \frac{x-1}{2} + \frac{y}{2}} \pmod{p} & \text{if } \frac{2c-d}{x} \equiv \pm 2 \pmod{5}, \\ \pm(-1)^{\frac{d}{4} + \frac{x-1}{2} + \frac{y}{2}} \frac{c}{d} \pmod{p} & \text{if } \frac{2c-d}{x} \equiv \pm 1 \pmod{5}. \end{cases}$$

Proof. Since $(\frac{\pm 1}{1+2i})_4 = \pm 1$ and $(\frac{\pm 2}{1+2i})_4 = \pm i$, taking $a = 2$ and $b = 1$ in Theorem 4.7 we derive the result.

We remark that Corollary 4.5 partially solves [S4, Conjecture 9.8].

5. Congruences for $U_{\frac{p-1}{4}}(2a, -1)$ and $V_{\frac{p-1}{4}}(2a, -1) \pmod{p}$.

For two numbers P and Q the Lucas sequences $\{U_n(P, Q)\}$ and $\{V_n(P, Q)\}$ are defined by

$$\begin{aligned} U_0(P, Q) &= 0, \quad U_1(P, Q) = 1, \quad U_{n+1}(P, Q) = PU_n(P, Q) - QU_{n-1}(P, Q) \quad (n \geq 1), \\ V_0(P, Q) &= 2, \quad V_1(P, Q) = P, \quad V_{n+1}(P, Q) = PV_n(P, Q) - QV_{n-1}(P, Q) \quad (n \geq 1). \end{aligned}$$

Set $D = P^2 - 4Q$. It is well known that

$$(5.1) \quad U_n(P, Q) = \frac{1}{\sqrt{D}} \left\{ \left(\frac{P + \sqrt{D}}{2} \right)^n - \left(\frac{P - \sqrt{D}}{2} \right)^n \right\} \quad (D \neq 0)$$

and

$$(5.2) \quad V_n(P, Q) = \left(\frac{P + \sqrt{D}}{2} \right)^n + \left(\frac{P - \sqrt{D}}{2} \right)^n.$$

Theorem 5.1. *Let p be a prime of the form $8k + 1$ and $a \in \mathbb{Z}$ with $2 \nmid a$. Suppose that $p = c^2 + d^2 = x^2 + (a^2 + 1)y^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod{4}$, $d = 2^r d_0$, $y = 2^\beta y_0$ and $d_0 \equiv y_0 \equiv 1 \pmod{4}$. Assume $(c, x + d) = 1$ or $(d_0, x + c) = 1$. Then*

$$U_{\frac{p-1}{4}}(2a, -1) \equiv \begin{cases} (-1)^{\frac{a-1}{2} + \frac{d}{4} + \frac{x-1}{2}} \frac{y}{x} \pmod{p} & \text{if } 4 \mid y - 2, \\ 0 \pmod{p} & \text{if } 4 \mid y \end{cases}$$

and

$$V_{\frac{p-1}{4}}(2a, -1) \equiv \begin{cases} 0 \pmod{p} & \text{if } 4 \mid y - 2, \\ 2(-1)^{\frac{d}{4} + \frac{y}{4}} \pmod{p} & \text{if } 4 \mid y. \end{cases}$$

Proof. Set $a_1 = (1 - (-1)^{\frac{a-1}{2}} a)/2$ and $b_1 = (1 + (-1)^{\frac{a-1}{2}} a)/2$. Then $2 \mid a_1$, $2 \nmid b_1$ and $a^2 + 1 = 2(a_1^2 + b_1^2)$. It is clear that

$$\left(\frac{(a_1 c + b_1 d) / ((-1)^{\frac{x-1}{2}} x)}{b_1 + a_1 i} \right)_4 = (-1)^{\frac{x-1}{2} \cdot \frac{a_1}{2}} \left(\frac{(a_1 c + b_1 d) / x}{b_1 + a_1 i} \right)_4.$$

We first assume $a \equiv 1 \pmod{4}$. Replacing d, x with $-d, (-1)^{\frac{x-1}{2}}x$ in [S4, Theorem 8.3(i)] we obtain

$$U_{\frac{p-1}{4}}(2a, -1) \equiv \begin{cases} \mp(-1)^{\frac{x-1}{2} \cdot \frac{a-1}{4}} (-a_1^2 - b_1^2)^{\frac{p-1}{8}} \left(-\frac{c}{d}\right)^{(1-(-1)^{\frac{a-1}{4}})/2} (-1)^{\frac{x-1}{2}} \frac{y}{x} \pmod{p} \\ \quad \text{if } 4 \mid y - 2 \text{ and } \left(\frac{(a_1c+b_1d)/x}{b_1+a_1i}\right)_4 = \pm 1, \\ \mp(-1)^{\frac{x-1}{2} \cdot \frac{a-1}{4}} (-a_1^2 - b_1^2)^{\frac{p-1}{8}} \left(-\frac{c}{d}\right)^{1+(1-(-1)^{\frac{a-1}{4}})/2} (-1)^{\frac{x-1}{2}} \frac{y}{x} \pmod{p} \\ \quad \text{if } 4 \mid y - 2 \text{ and } \left(\frac{(a_1c+b_1d)/x}{b_1+a_1i}\right)_4 = \pm i, \\ 0 \pmod{p} \text{ if } 4 \mid y \end{cases}$$

and

$$V_{\frac{p-1}{4}}(2a, -1) \equiv \begin{cases} \pm 2(-1)^{\frac{x-1}{2} \cdot \frac{a-1}{4} + \frac{y}{4}} (-a_1^2 - b_1^2)^{\frac{p-1}{8}} \left(-\frac{c}{d}\right)^{(1-(-1)^{\frac{a-1}{4}})/2} \pmod{p} \\ \quad \text{if } 4 \mid y \text{ and } \left(\frac{(a_1c+b_1d)/x}{b_1+a_1i}\right)_4 = \pm 1, \\ \pm 2(-1)^{\frac{x-1}{2} \cdot \frac{a-1}{4} + \frac{y}{4}} (-a_1^2 - b_1^2)^{\frac{p-1}{8}} \left(-\frac{c}{d}\right)^{1+(1-(-1)^{\frac{a-1}{4}})/2} \pmod{p} \\ \quad \text{if } 4 \mid y \text{ and } \left(\frac{(a_1c+b_1d)/x}{b_1+a_1i}\right)_4 = \pm i, \\ 0 \pmod{p} \text{ if } 4 \mid y - 2. \end{cases}$$

From Theorem 4.7 we know that

$$\begin{aligned} & (-a_1^2 - b_1^2)^{\frac{p-1}{8}} \\ & \equiv \begin{cases} \pm(-1)^{\frac{d}{4} + \frac{y}{2}} \pmod{p} & \text{if } 4 \mid a_1 \text{ and } \left(\frac{(a_1c+b_1d)/x}{b_1+a_1i}\right)_4 = \pm 1, \\ \pm(-1)^{\frac{b_1-1}{2} + \frac{d}{4} + \frac{y}{2} + \frac{x-1}{2}} \frac{d}{c} \pmod{p} & \text{if } 4 \mid a_1 - 2 \text{ and } \left(\frac{(a_1c+b_1d)/x}{b_1+a_1i}\right)_4 = \pm 1, \\ \pm(-1)^{\frac{d}{4} + \frac{y}{2}} \frac{c}{d} \pmod{p} & \text{if } 4 \mid a_1 \text{ and } \left(\frac{(a_1c+b_1d)/x}{b_1+a_1i}\right)_4 = \pm i, \\ \pm(-1)^{\frac{b_1-1}{2} + \frac{d}{4} + \frac{y}{2} + \frac{x-1}{2}} \pmod{p} & \text{if } 4 \mid a_1 - 2 \text{ and } \left(\frac{(a_1c+b_1d)/x}{b_1+a_1i}\right)_4 = \pm i. \end{cases} \end{aligned}$$

Now putting the above together we deduce the result in the case $a \equiv 1 \pmod{4}$. The case $a \equiv 3 \pmod{4}$ can be proved similarly by using [S4, Theorem 8.3(ii)] and Theorem 4.7.

Corollary 5.1. *Let p be a prime of the form $8k + 1$ and $a \in \mathbb{Z}$ with $2 \nmid a$. Suppose that $p = c^2 + d^2 = x^2 + (a^2 + 1)y^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod{4}$, $d = 2^r d_0$, $d_0 \equiv 1 \pmod{4}$ and $4 \mid y$. Assume $(c, x + d) = 1$ or $(d_0, x + c) = 1$. Then*

$$(a + \sqrt{a^2 + 1})^{\frac{p-1}{4}} \equiv (-1)^{\frac{d}{4} + \frac{y}{4}} \pmod{p}.$$

Proof. From (5.1), (5.2) and Theorem 5.1 we see that

$$\begin{aligned} (a + \sqrt{a^2 + 1})^{\frac{p-1}{4}} &= \frac{1}{2} V_{\frac{p-1}{4}}(2a, -1) + \sqrt{a^2 + 1} U_{\frac{p-1}{4}}(2a, -1) \\ &\equiv \frac{1}{2} V_{\frac{p-1}{4}}(2a, -1) \equiv (-1)^{\frac{d}{4} + \frac{y}{4}} \pmod{p}. \end{aligned}$$

This proves the corollary.

Corollary 5.2. *Let p be a prime of the form $8k + 1$ and $a \in \mathbb{Z}$ with $2 \nmid a$. Suppose that $p = c^2 + d^2 = x^2 + (a^2 + 1)y^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod{4}$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod{4}$. Assume $(c, x + d) = 1$ or $(d_0, x + c) = 1$. Then*

$$p \mid U_{\frac{p-1}{8}}(2a, -1) \iff 4 \mid y \quad \text{and} \quad \frac{p-1}{8} \equiv \frac{d}{4} + \frac{y}{4} \pmod{2}.$$

Proof. By [S4, (1.5)],

$$p \mid U_{\frac{p-1}{8}}(2a, -1) \iff V_{\frac{p-1}{4}}(2a, -1) \equiv 2(-1)^{\frac{p-1}{8}} \pmod{p}.$$

Now applying Theorem 5.1 we obtain the result.

Remark 5.1 Theorem 5.1 and Corollary 5.2 were conjectured by the author in [S4, Conjectures 9.17 and 9.19].

REFERENCES

- [BEW] B.C. Berndt, R.J. Evans and K.S. Williams, *Gauss and Jacobi Sums*, Wiley, New York, 1998.
- [E] R.J. Evans, *Residuacity of primes*, Rocky Mountain J. Math. **19** (1989), 1069-1081.
- [HW] R.H. Hudson and K.S. Williams, *Extensions of theorems of Cunningham-Aigner and Hasse-Evans*, Pacific J. Math. **104** (1983), 111-132.
- [IR] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory, 2nd ed.*, Springer, New York, 1990.
- [L] E. Lehmer, *On Euler's criterion*, J. Austral. Math. Soc. **1** (1959), 64-70.
- [Lem] F. Lemmermeyer, *Reciprocity Laws: From Euler to Eisenstein*, Springer, Berlin, 2000.
- [S1] Z.H. Sun, *Supplements to the theory of quartic residues*, Acta Arith. **97** (2001), 361-377.
- [S2] Z.H. Sun, *Quartic residues and binary quadratic forms*, J. Number Theory **113** (2005), 10-52.
- [S3] Z.H. Sun, *On the quadratic character of quadratic units*, J. Number Theory **128** (2008), 1295-1335.
- [S4] Z.H. Sun, *Quartic, octic residues and Lucas sequences*, J. Number Theory **129** (2009), 499-550.
- [S5] Z.H. Sun, *Congruences for $q^{[p/8]} \pmod{p}$* , Acta Arith. **159** (2013), 1-25.