

COMPACT ARITHMETIC QUOTIENTS OF THE COMPLEX 2-BALL AND A CONJECTURE OF LANG

MLADEN DIMITROV AND DINAKAR RAMAKRISHNAN

INTRODUCTION

Let $\mathcal{H}_{\mathbb{C}}^2$ be the 2-dimensional complex hyperbolic space, represented by the unit ball in \mathbb{C}^2 equipped with the Bergmann metric of constant holomorphic sectional curvature -1 , on which the Lie group $U(2, 1)$ acts in a natural way. Let F be a totally real number field of degree d and ring of integers \mathcal{O}_F , and let M be a CM quadratic extension of F . Let G be a unitary group over F defined by a 3-dimensional M -vector space V endowed with an anisotropic hermitian form $\langle \ , \ \rangle$ of signature $(2, 1)$ at one infinite place and $(3, 0)$ at the others. Let Γ be a discrete subgroup of $G(F)$ which is arithmetic, i.e., commensurable with $G(\mathcal{O}_F)$. Then the quotient $X_{\Gamma} := \Gamma \backslash \mathcal{H}_{\mathbb{C}}^2$ is a compact complex analytic space, which can be given the structure of a projective normal surface, in fact over a number field containing M . Denote by $q = q(X_{\Gamma})$ the *irregularity* of X_{Γ} , given by the dimension of $H^0(X_{\Gamma}, \Omega_{X_{\Gamma}}^1)$.

Any projective surface X defined over a number field k is said to be *Mordellic* if, and only if, for any finite extension k' of k , the set $X(k')$ of k' -rational points of X is finite. Lang conjectured in [L, Conjecture VIII.1.2], (see also [T, p.xviii]) that any *hyperbolic*, smooth X is Mordellic; this is consistent with the general philosophy of Vojta.

The following result makes use of certain key theorems of Faltings [F1, F2] (see also [V]) and Rogawski [R1, R2], as well as the hyperbolicity of X_{Γ} .

Theorem 0.1. *Let G be as above and $\Gamma \subset G(F)$ be an arithmetic subgroup. Then we have the following:*

- (i) *There exists a subgroup Γ' of finite index in Γ such that $X_{\Gamma'}$ is Mordellic;*
- (ii) *Suppose X_{Γ} is smooth and that it does not admit a dominant map to its Albanese variety. Then X_{Γ} is itself Mordellic.*

The paper is organized as follows. In §1 we give a precise definition of X_{Γ} and give a criterion for its smoothness that can always be achieved by taking a

finite cover. In this (smooth) case it seems to be a folklore result that X_Γ is of general type, which we verify for completeness in §2 using the Enriques-Kodaira classification and hyperbolicity. The *Bombieri-Lang conjecture* predicts that $X_\Gamma(k)$ is not Zariski dense, implying that the k -rational points are contained in a finite union of curves. This we establish in §3 when $q > 2$ using a deep Theorem due to Faltings. In §4 we recall Rogawski's classification of cohomological automorphic forms on G contributing to $H^1(X_\Gamma)$ for congruence subgroups Γ , from which we deduce that q becomes positive, and in fact arbitrarily large, over a finite cover of X_Γ , for any arithmetic Γ . The importance of this is evident given Mazur's path breaking work on the modular curves in the seventies. Finally in §5 we prove our main Theorem 0.1 using Propositions 3.1 and 4.1, the hyperbolicity of X_Γ , as well as Faltings' proof of the Mordell's conjecture for curves.

Note that even though our theorem only concerns arithmetic subgroups, because F and M can vary, it can be applied to infinitely many pairwise non-commensurable cocompact lattices in $U(2, 1)$.

Our results do *not* apply, however, to the analogous case of (a cocompact discrete subgroup of) a unitary group G' defined by a division algebra of dimension 9 over an imaginary quadratic field M with an involution of the second kind. In that case it is known that q is zero for any congruence subgroup $\Gamma \subset G'(\mathbb{Z})$; this was proved by Rapoport and Zink [RZ] under a ramification hypothesis, and later by Rogawski [R1] using a different method, without the hypothesis.

In a related paper [DR], we analyze the more complicated case when G is defined by an isotropic hermitian form in 3 variables on M^3 . Then, for the arithmetic quotient $Y_\Gamma := \Gamma \backslash \mathcal{H}_\mathbb{C}$ to be a surface, it is necessary that $F = \mathbb{Q}$, and Y_Γ is always *non-compact*. Its toroidal compactification X_Γ is not hyperbolic, being a union of Y_Γ with a finite number of elliptic curves E_κ indexed by the cusps κ . If we proceed as above, one cannot rule out having curves of genus ≤ 1 on X_Γ (in fact there are rational curves at low levels), except to know that by the hyperbolicity of the open surface Y_Γ , they must meet some E_κ on the boundary; the nature of the meeting points is also crucial. Nevertheless we exhibit in [DR] an infinite class \mathcal{C} of X_Γ of congruence type which are not covers of each other, such that $X_\Gamma(M)$ is finite; of course X_Γ cannot be Mordellic in this situation as over some finite extension the E_κ will have positive Mordell-Weil rank. For the finiteness statement over M for the members of \mathcal{C} , we do not appeal to Faltings, and instead employ the full force

of the theory of automorphic forms on $U(3)$, exhibiting suitable abelian variety quotients A of $\text{Alb}(X_\Gamma)$ with finite Mordell-Weil group (as in Mazur's work on modular curves), and establish a formal immersion over \mathbb{C} at the torsion points of E_κ where any curve C in the inverse image of $A(k)$ may meet E_κ . We do the last part by analyzing the Fourier-Jacobi coefficients of certain associated *residual* Picard modular forms.

Acknowledgments. We would like to thank Don Blasius, Laurent Clozel, Najmuddin Fakhruddin, Dick Gross, Haruzo Hida and David Rohrlich for helpful conversations. In fact it was Fakhruddin who suggested our use of Lang's conjecture for abelian varieties. Needless to say, this Note owes much to the deep results of Faltings. Thanks are also due to Serge Lang (posthumously), and to John Tate, for getting one of us interested in the conjectural Mordellic property of hyperbolic varieties. Finally, we are also happy to acknowledge partial support the following sources: the Agence Nationale de la Recherche grants ANR-10-BLAN-0114 and ANR-11-LABX-0007-01 for the first author (M.D.), and from the NSF grant DMS-1001916 for the second author (D.R.).

1. NEAT ARITHMETIC SUBGROUPS

We keep the notation from the introduction. The reductive group G over F is precisely defined by

$$G = \{g \in \text{GL}(V) \mid \langle g(v), g(v') \rangle = \langle v, v' \rangle, \text{ for all } v, v' \in V\}.$$

For example, if M is any quadratic CM extension of a real quadratic field F of discriminant D , then one might consider a hermitian form given by $\begin{pmatrix} 1 & \\ & 1 \\ & & \sqrt{D} \end{pmatrix}$.

Let $F_\infty = F \otimes_{\mathbb{Q}} \mathbb{R} \simeq \mathbb{R}^d$ and let $\iota : F \hookrightarrow \mathbb{R}$ be the embedding corresponding to the first factor in the identification $G(F_\infty) = U(2, 1) \times U(3)^{d-1}$. For simplicity we keep the same notation for a fixed extension $\iota : M \hookrightarrow \mathbb{C}$. We will complete this extension ι to a CM type Φ of M .

Let $\mathcal{H}_{\mathbb{C}}^2$ be the hermitian symmetric space of negative lines in $V \otimes_{F, \iota} \mathbb{R} \simeq \mathbb{C}^3$. By choosing a basis of $V \otimes_{F, \iota} \mathbb{R}$ where the hermitian form is represented by the matrix $\begin{pmatrix} 1 & & \\ & 1 & \\ & & -1 \end{pmatrix}$ one can see that $\mathcal{H}_{\mathbb{C}}^2$ is isometric to the open unit complex 2-ball.

Since G is defined by a hermitian form on M^3 , we have an embedding $G(F) \hookrightarrow \text{GL}(3, M)$, through which we may view elements of $G(F)$ as 3×3 matrices. When we compose this embedding with (the extension to M of) ι , the image of $\iota(G(F))$ lands in $U(2, 1) \subset \text{GL}(3, \mathbb{C})$.

The standard left action of $\mathrm{GL}_3(\mathbb{C})$ on $\mathbb{P}^2(\mathbb{C})$ yields a left transitive action of the first component $G_\iota = \mathrm{U}(2, 1)$ of $G(F_\infty)$ on $\mathcal{H}_{\mathbb{C}}^2$. The standard maximal compact subgroup $K_\infty \subset G(F_\infty)$ is defined as the stabilizer of a distinguished point in $\mathcal{H}_{\mathbb{C}}^2$ and we have $\mathcal{H}_{\mathbb{C}}^2 \simeq G(F_\infty)/K_\infty$. Since K_∞ also stabilizes the orthogonal of the corresponding negative line, one has $K_\infty \simeq \mathrm{U}(2) \times \mathrm{U}(1) \times \mathrm{U}(3)^{d-1}$.

Any arithmetic subgroup Γ of $G(F)$ acts properly discontinuously on $\mathcal{H}_{\mathbb{C}}^2$ (via ι), and the quotient $X_\Gamma = \Gamma \backslash \mathcal{H}_{\mathbb{C}}^2$ is a complex analytic space, which admits, by the Baily-Borel theorem, the structure of a normal, projective variety over \mathbb{C} . By a theorem of Kazhdan (see for example [NR]), for any automorphism σ of \mathbb{C} , the σ -conjugate of X_Γ is again a locally symmetric variety. In fact, since it will, like X_Γ , satisfy $c_1^2 = 3c_2$ ([H]), where c_1, c_2 are the Chern classes of the complex tangent bundle of the surface, we may realize it in the form $\Gamma' \backslash \mathcal{H}_{\mathbb{C}}^2$, for a cocompact lattice Γ' by Miyaoka [M] and Yau [Y]. It follows that an open subgroup of $\mathrm{Aut}(\mathbb{C})$ preserves X_Γ , allowing one to conclude that X_Γ is defined over a number field.

A subgroup Γ of $G(F)$ is called a *congruence* subgroup if there exists an integer N such that Γ contains the principal congruence subgroup of level N , defined as:

$$\Gamma(N) = \ker (G(\mathcal{O}_F) \rightarrow G(\mathcal{O}_F/N\mathcal{O}_F)).$$

In that case X_Γ is (a connected component of) a Shimura surface (see §4) and there is an explicite rule specifying its field of definition.

Definition 1.1. An arithmetic group $\Gamma \subset G(F)$ is *neat* if $\bar{\Gamma} = \Gamma/\Gamma \cap \mathrm{U}(1)$ is torsion free, where $\mathrm{U}(1)$ is centrally embedded in $G(F)$.

Lemma 1.2. (i) *The variety X_Γ is an orbifold.*

(ii) *The variety X_Γ is a manifold if and only if Γ is neat.*

(iii) *Any finite index subgroup Γ' of a neat group Γ is neat. Moreover, if it is also normal, then the natural morphism $X_{\Gamma'} \rightarrow X_\Gamma$ is an étale covering of smooth varieties of group $\Gamma/\Gamma'(\Gamma \cap \mathrm{U}(1))$.*

Proof. (i) The stabiliser of any point of $\mathcal{H}_{\mathbb{C}}^2$ in $G(F_\infty)$ is of the form $g_\infty K_\infty g_\infty^{-1}$ for some element $g_\infty \in G(F_\infty)$, hence is a compact group. Since Γ is a discrete subgroup of $G(F_\infty)$ it follows that $g_\infty K_\infty g_\infty^{-1} \cap \Gamma$ is finite.

(ii) If Γ is neat, then all finite order elements of Γ are in $\Gamma \cap \mathrm{U}(1)$, which, being central, acts trivially on $\mathcal{H}_{\mathbb{C}}^2$, hence X_Γ is a manifold. For the other direction, a direct computation shows that a finite order element γ of Γ is semi-simple and

necessarily fixes a point in $\mathcal{H}_{\mathbb{C}}^2$. Since X_{Γ} is smooth at that point, γ should belong to $\Gamma \cap \mathrm{U}(1)$ as desired.

(iii) The claim follows directly from the exact sequence of groups:

$$1 \rightarrow \bar{\Gamma}' \rightarrow \bar{\Gamma} \rightarrow \Gamma/\Gamma'(\Gamma \cap \mathrm{U}(1)) \rightarrow 1.$$

□

Since X_{Γ} is an orbifold, it is a standard result that it has a finite cover which is a manifold. The following lemma, constructs such a cover explicitly.

Lemma 1.3. *The principal congruence subgroup $\Gamma(3)$ is neat.*

Proof. Since $\Gamma(3) \cap \mathrm{U}(1)$ is finite, it suffices to check that $\Gamma(3)$ does not contain any finite order non-trivial elements. Suppose that $\gamma \in \Gamma(3)$ has prime order ℓ . Since at least one of the eigenvalues ζ of γ is a root of unity of order ℓ , the congruence condition on γ implies that $\ell = 3$. Denote by a the highest power of 3 dividing all the coefficients of $\gamma - 1$. Then the highest power of 3 dividing all the coefficients of $\gamma^3 - 1$ is $a + 1$, leading to a contradiction. □

2. GENERAL TYPE

Proposition 2.1. *Suppose that Γ is neat. Then X_{Γ} is of general type.*

We may, and we will, work over \mathbb{C} . Note also that Γ need not be arithmetic in this section, only a torsion-free (cocompact) lattice in $\mathrm{U}(2, 1)$.

Proof. Since $X = X_{\Gamma}$ is hyperbolic, it cannot contain any rational curves, implying in particular that it is a minimal surface.

We will use the Enriques-Kodaira classification of smooth projective surfaces. The *Kodaira dimension* κ of X has four possible values, namely in $\{-\infty, 0, 1, 2\}$, and we have to rule out the occurrence of the first three situations, as X is of general type if, and only if, $\kappa = 2$. Denote by $\mathcal{K} = \mathcal{K}_X$ the canonical divisor of X .

First suppose $\kappa = -\infty$. Then all the plurigena $P_m := \dim H^0(X, m\mathcal{K})$ vanish. If $q = 0$, then by the Castelnuovo criterion, having $P_2 = q = 0$ implies that X is a rational surface. If $q > 0$, then the fibers of the Albanese map $X \rightarrow \mathrm{Alb}(X)$ are rational curves, i.e., X is ruled. In either case, we would get a non-constant holomorphic map $\mathbb{C} \rightarrow X$, which is impossible. Kodaira's class VII surfaces do not occur here as X is algebraic.

Next look at the case when $\kappa = 0$. In this case $\mathcal{K}^2 = c_1^2 = 0$. The only algebraic surfaces in characteristic zero are among the following types: (i) $K3$, (ii) Enriques, (iii) abelian, and (iv) hyperelliptic. Suppose we are in case (i). Then one knows (see [MM]) that there is always a rational curve in any complex $K3$ surface, which is impossible in our situation. In case (ii), X is necessarily a quotient of a $K3$ surface X' by an involution, which would again give a non-constant holomorphic map from \mathbb{C} to X , furnishing a contradiction. In case (iii), X is a quotient of \mathbb{C}^2 by a lattice, which is again impossible. Finally, in case (iv), such a surface is, over \mathbb{C} , a quotient of a product of two elliptic curves by a finite group of automorphisms. One again this would result in a non-constant holomorphic map $\mathbb{C} \rightarrow X$, and we are done eliminating $\kappa \leq 0$.

Finally, let $\kappa = 1$. Here X is an elliptic surface, admitting a fibration over a base curve with all but finitely many fibers being elliptic curves. Again this is impossible by the hyperbolicity of X . \square

3. BOMBIERI-LANG FOR X

Proposition 3.1. *Suppose that X_Γ is smooth and that it does not admit a dominant map to its Albanese variety. Then the points of X_Γ over any number field k , containing its field of definition, are not Zariski dense in X_Γ .*

This is as predicted by the Bombieri-Lang conjecture as X_Γ is of general type.

The proof below will show that it is a consequence of the following powerful *result of Faltings* on the rational points of subvarieties of abelian varieties.

Theorem F (Faltings [F2], [V]). Suppose A is an abelian variety over a number field k , $Z \subset A$ a closed subvariety. Then there are finitely many translates Z_i of k -rational abelian subvarieties of A , such that $Z_i \subset Z$, and such that each k -rational point of Z lies on one of the Z_i .

Proof of Proposition 3.1. Write $X = X_\Gamma$. If $X(k)$ is empty, there is nothing to prove. Otherwise, use a k -rational point of X to define the Albanese map over k :

$$j : X \rightarrow \text{Alb}(X).$$

Then $Z = j(X)$ is a closed, irreducible subvariety of $\text{Alb}(X)$. Applying Faltings' Theorem F with $A = \text{Alb}(X)$, we get a finite number, say $m \geq 1$, of k -rational

translates Z_i of abelian subvarieties of $\text{Alb}(X)$ such that

$$Z(k) \subset \bigcup_{i=1}^m Z_i(k) \quad \text{and} \quad Z_i \subset Z.$$

Since the Albanese map is defined over k , all the k -rational points of X are contained in those $j^{-1}(Z_i)$. It remains to show that each $j^{-1}(Z_i)$ is either a point or a curve (possibly singular and reducible). Otherwise, since X is irreducible, one would have had $Z = Z_i = \text{Alb}(X)$, contradicting the assumption that X does not admit a dominant map to its Albanese variety. \square

Remark 3.2. Suppose X is a smooth projective surface over \mathbb{C} of general type such that $c_1^2 = 3c_2$. Then by Miyaoka [M, Proposition 2.1.1] and Yau [Y, Theorem 3], X is of the form $\Gamma \backslash \mathcal{H}_{\mathbb{C}}^2$, for a cocompact lattice Γ in $\text{U}(2, 1)$, which may not be arithmetic. Let k be a finitely generated field (over \mathbb{Q}) over which X is defined. Suppose $q(X) > 2$. Then, noting that Faltings' theorem on abelian varieties [F2] holds over finitely generated fields, we can conclude, as in the proof of Proposition 3.1, that $X(k)$ is contained in a finite union of curves on X .

4. POSITIVITY OF q FOR A CONGRUENCE COVER OF X_{Γ}

By the Hodge decomposition of $H^1(X_{\Gamma})$, the first Betti number of X_{Γ} is $2q(X_{\Gamma})$. The positivity of $q(X_{\Gamma})$ is an essential ingredient in the proof of our Diophantine results and is the object of following proposition which will be deduced from the results of Rogawski [R1, R2]. Many examples were known earlier, via the theta correspondence, by the works of Kazhdan [K] and Shimura [Sh, Theorem 8.1].

Proposition 4.1. *For any congruence subgroup Γ of $G(F)$ and for any $r > 0$, there is a subgroup Γ' of finite index in Γ such that $q(X_{\Gamma'}) > r$.*

Proof. There is a well known decomposition

$$(1) \quad H^1(X_{\Gamma}, \mathbb{C}) \simeq \bigoplus_{\pi} H^1(\text{Lie}(G(F_{\infty})), K_{\infty}; \pi)^{\oplus m(\pi, \Gamma)},$$

where π runs over irreducible unitary representations of $G(F_{\infty})$ occurring with multiplicity $m(\pi, \Gamma)$ in $L^2(\Gamma \backslash G(F_{\infty}))$, which is discrete by virtue of Γ being cocompact.

At the distinguished archimedean place ι where $G_{\iota} = \text{U}(2, 1)$ there are exactly two irreducible non-tempered unitary representations of G_{ι} , denoted π^+ and π^- ,

with non-zero relative Lie algebra cohomology in degree 1; π^+ occurs in $H^{1,0}$ and π^- in $H^{0,1}$. So we need to show that $m(\pi^+, \Gamma')$ is non-zero for sufficiently many $\Gamma' \subset \Gamma$. In fact it will suffice to show that there are infinitely many $\Gamma' \subset \Gamma$ of relatively prime auxiliary levels (that is to say levels outside the level of Γ) such that $m(\pi^+, \Gamma') \neq 0$.

The restriction to \mathbb{C}^* of the Langlands parameter of π^+ is given by (see [La, p.64]):

$$(2) \quad z \mapsto \begin{pmatrix} z^{-1} & 0 & 0 \\ 0 & z/\bar{z} & 0 \\ 0 & 0 & \bar{z} \end{pmatrix} \in {}^L G^0 = \mathrm{GL}_3(\mathbb{C}).$$

We will work adelicly and construct infinitely many automorphic representations $\pi = \pi_\infty \otimes \pi_f$ of $G(\mathbb{A}_F)$ of relatively prime auxiliary levels such that $\pi_\ell = \pi^+$. As usual, $\mathbb{A}_F = F_\infty \times \mathbb{A}_{F,f}$ denotes the adèle ring of F , and K is a open compact subgroup of $G(\mathbb{A}_{F,f})$. The adelic quotient

$$(3) \quad X_K := G(F) \backslash G(\mathbb{A}_F) / K_\infty K$$

is a finite disjoint union of surfaces, each of the form X_Γ for some congruence subgroup $\Gamma \subset G(F)$. The decomposition (1) can be rewritten as:

$$H^1(X_K, \mathbb{C}) \simeq \bigoplus_\pi H^1(\mathrm{Lie}(G(F_\infty)), K_\infty; \pi_\infty)^{\oplus m(\pi_f, K)},$$

and by the Multiplicity One Theorem of Rogawski one deduces that:

$$(4) \quad q(X_K) = \sum_{\substack{\pi \text{ automorphic} \\ \pi_\ell \simeq \pi^+}} \dim(\pi_f^K) = \sum_{\substack{\pi \text{ automorphic} \\ \pi_\ell \simeq \pi^-}} \dim(\pi_f^K).$$

Again by Rogawski [R2], we have an explicit description of the automorphic representations π of $G(\mathbb{A})$ such that $\pi_\ell \simeq \pi^+$, that we will now present: Let W_F (resp. W_M) be the global Weil group of F (resp. M). (Due to the simple nature of our situation, we may write the parameter in terms of W_F instead of the conjectural global Langlands group \mathcal{L}_F .) Let ρ denote the non-trivial automorphism of M/F , written as $x \rightarrow \bar{x}$. The corresponding global Arthur parameter

$$\psi : W_F \times \mathrm{SL}(2, \mathbb{C}) \rightarrow {}^L G = \mathrm{GL}(3, \mathbb{C}) \rtimes \{1, \rho\}$$

is such that its restriction ψ_M to W_M is given by the 3-dimensional representation $(\lambda \otimes \mathrm{St}) \oplus (\chi_M \otimes 1)$, where St (resp. 1) is the standard 2-dimensional (resp. trivial) representation of $\mathrm{SL}(2, \mathbb{C})$, λ is a character of W_M whose transfer to W_F

is the quadratic character $\delta_{M/F}$ associated to M/F , and $\chi_M = \chi/\chi \circ \rho$ is the base change of an idele class character χ of $U(1)/M$. We have $\text{Hom}_{\mathbb{Q}}(M, \mathbb{C}) = \Phi \cup \Phi^\rho$, and at the distinguished place ι in Φ , $\psi_{M,\iota}$ agrees with (2) on \mathbb{C}^* . For reasons which will become apparent when we do the transfer from the quasi-split group, we take ψ_M to have the same type at all $v \in \Phi$. Then λ is an algebraic (unitary) Hecke character of M of weight 1, whereas χ is an algebraic character of $U(1)$ of weight -1 .

Let \tilde{G} denote the quasi-split unitary group associated to M/F , so that G is an inner form of \tilde{G} . (In [R1], [R2], the author uses (G, G') for our (\tilde{G}, G) .) Let $\tilde{\Pi}$ be the global Arthur packet associated to ψ , with $\tilde{\Pi}_v$ denoting the local Arthur packet at each place v . It arises by endoscopy from $U(2) \times U(1)$, and the embedding depends on the choice of a Hecke character μ of M whose restriction to F is the quadratic character $\delta_{M/F}$. When v is finite and remains prime in M , $\tilde{\Pi}_v = \{\pi_{s,v}, \pi_{n,v}\}$, where $\pi_{n,v}$ is a certain non-tempered representation and $\pi_{s,v}$ is in the discrete series. At a split place, $\tilde{\Pi}_v = \{\pi_{n,v}\}$, and for almost all v , $\pi_{n,v}$ is unramified. An element π of $\tilde{\Pi}$ is a tensor product $\otimes_v \pi_v$ with each π_v in $\tilde{\Pi}_v$ such that $\pi_v \simeq \pi_{n,v}$ for almost all v .

Since our G is an inner form defined by a hermitian form on M^3 , we know from Rogawski (see [R1, §14.4] and [R2, p.397]) that there will be a corresponding Arthur packet Π of representations of $G(\mathbb{A}_F)$ *if, and only if*, certain condition is satisfied at each of the $d - 1$ archimedean places v where G_v is the compact real group $U(3)$. When such a correspondence is possible, we will say that Π exists, in which case $\Pi_v = \tilde{\Pi}_v$ at all the places v outside the $d - 1$ archimedean places, and in each of the latter, Π_v is a singleton consisting of a finite-dimensional representation σ_v of $U(3)$. As in [R2, p.397], we will write (at each $v \in \Phi$): $\mu_v(z) = (z/|z|)^{2t+1}$, and the one-dimensional representation of $U(2) \times U(1)$ associated to ψ at v by $(h, u) \rightarrow \det(h)^{r-t}u^s$. Then in our case, λ , resp. χ , being of weight 1, resp. -1 , implies that $r = -1$ and $s = 1$. Moreover, an explicit calculation shows in our case that μ has weight -1 , implying that $t = 0$. Since $s > r$, by the recipe on the same page of [R2], the associated triple (a, b, c) is given by $(1, 0, -1)$. Since it is regular, i.e., with three distinct coordinates, the local condition is satisfied at every $v \neq \iota$ in Φ , and there is a corresponding σ_v of $U(3)$. The regularity of (a, b, c) implies that there is a square integrable representation $\pi_{s,v}$ of $U(2, 1)$ in $\tilde{\Pi}_v$, which corresponds to σ_v . In fact, since the highest weight of σ_v is given by $\text{diag}(x, y, z) \rightarrow x^{a-1}y^bz^{c+1}$,

it is the trivial representation. In any case, the global Arthur packet Π exists for our ψ . Now let $\pi \in \Pi$ be such that $\pi_\iota = \pi^+ \in \Pi_\iota$. By [R2, Theorem 1.2], for such a $\pi \in \Pi$, we have

$$(5) \quad \pi \text{ automorphic} \iff W(\lambda\chi_M^{-1}) = (-1)^{d-1+s(\pi)},$$

where $W(\lambda\chi_M^{-1}) \in \{\pm 1\}$ is the root number of the weight 3 algebraic Hecke character $\lambda\chi_M^{-1}$ of M , and $s(\pi)$ is the number of finite inert places v where $\pi_v \simeq \pi_{s,v}$.

Claim. For any CM extension M/F and any CM type Φ , there exists an algebraic Hecke character λ of weight 1 and CM type Φ , and an algebraic character χ of $U(1)$ of weight -1 .

Proof. Consider the character on M_∞^* given by $\lambda_\infty(z) = \prod_{v \in \Phi} \frac{\bar{z}_v}{|z_v|}$. Since M/F is a CM extension, the index m of $(\mathcal{O}_F^*)^2$ in \mathcal{O}_M^* is finite, and λ_∞ is trivial on $(\mathcal{O}_M^*)^m$. By [C, Théorème 1] there exists an open compact subgroup U of $\mathbb{A}_{M,f}$ such that $U \cap \mathcal{O}_M^* \subset (\mathcal{O}_M^*)^m$, hence λ_∞ can be extended (trivially) to $M^*UM_\infty^*$. Finally, since $\mathbb{A}_M^*/M^*UM_\infty^*$ is a finite abelian (class) group, there exists a character λ of \mathbb{A}_M^*/M^* extending λ_∞ .

Since $(\lambda \circ \rho)/\lambda$ is an algebraic Hecke character of weight -2 with trivial restriction to F , it is the base change of a weight -1 algebraic character χ of $U(1)$. \square

Let Π be the global Arthur packet associated to a pair of characters (λ, χ) as in the above Claim. Fix a place v_0 of F , inert in M which is relatively prime to the conductors of λ and χ , and consider the open compact subgroup $K_s = \prod_v K_{s,v}$ of $G(\mathbb{A}_{F,f})$ such that $\pi_{s,v_0}^{K_{s,v_0}} \neq 0$ and $\pi_{n,v}^{K_{s,v}} \neq 0$ for all finite places $v \neq v_0$, with $K_{s,v}$ being the standard hyperspecial maximal compact, for almost all v .

Denote by \mathcal{P} the set of rational primes p which are totally split in F and such that every prime \mathfrak{p} of F dividing p remains inert in M . Note that \mathcal{P} is infinite, since M is a CM extension of F . Let $p \in \mathcal{P}$ be relatively prime to $2v_0$ and to the conductors of λ and χ , and denote by ε_p the unique quadratic Dirichlet character ramified only at p .

The quadratic Hecke character $\varepsilon_p \circ N_{M/\mathbb{Q}}$ is trivial on F hence equal to the base change $\nu_{p,M}$ of a quadratic character ν_p of $U(1)$, ramified only at places above p .

Consider the global Arthur packet Π^p attached to $(\lambda, \chi\nu_p)$.

Claim. For every prime \mathfrak{p} of F dividing p , the representation $\pi_{n,\mathfrak{p}} \in \Pi_{\mathfrak{p}}^p$ has non-zero invariants under the open compact subgroup:

$$K_0^1(\mathfrak{p}) = \left\{ k \in G(\mathcal{O}_{F,\mathfrak{p}}), k \equiv \begin{pmatrix} * & * & * \\ 0 & 1 & * \\ 0 & 0 & * \end{pmatrix} \pmod{\mathfrak{p}} \right\}.$$

Proof. By [R1, §12.2] the representation of $G(F_{\mathfrak{p}})$ obtained by parabolic induction of the character

$$(\alpha, \beta) \mapsto \lambda_{\mathfrak{p}}(\alpha) |\alpha|_{M_{\mathfrak{p}}}^{3/2} \chi_{\mathfrak{p}}(\beta) \nu_{p,\mathfrak{p}}(\beta)$$

of the standard torus $M_{\mathfrak{p}}^* \times \mathrm{U}(1)(F_{\mathfrak{p}})$ has $\pi_{n,\mathfrak{p}}$ as a unique irreducible quotient (the irreducible sub-representation being $\pi_{s,\mathfrak{p}}$). Since $\lambda_{\mathfrak{p}}$ and $\chi_{\mathfrak{p}}$ are unramified and $\nu_{p,\mathfrak{p}}$ is tamely ramified, the claim follows by explicit computation [DR]. \square

Consider an element $\pi^p = \otimes_v \pi_v^p \in \Pi^p$ such that $\pi_{\iota}^p = \pi^+$, $\pi_v^p = 1$ for every infinite $v \neq \iota$, $\pi_v^p = \pi_{n,v}$ for every finite $v \neq v_0$, and finally:

$$(6) \quad \pi_{v_0}^p = \begin{cases} \pi_{n,v_0} & , \text{ if } W(\lambda \chi_M^{-1} \nu_{p,M}^{-1}) = (-1)^{d-1}, \text{ and} \\ \pi_{s,v_0} & , \text{ if } W(\lambda \chi_M^{-1} \nu_{p,M}^{-1}) = (-1)^d. \end{cases}$$

By (5) π^p is automorphic, and by the claim $(\pi_f^p)^{K_s^p} \neq 0$, where K_s^p is the subgroup of K_s with the maximal compact $K_{s,\mathfrak{p}}$ being replaced by $K_0^1(\mathfrak{p})$ at all places \mathfrak{p} of F dividing p .

Consider a strictly increasing sequence $(p_i)_{i \geq 1}$ of elements $p \in \mathcal{P}$ as above and for each $r \geq 1$ denote by X_r the adelic quotient (3) corresponding to the open compact subgroup $\bigcap_{i=1}^r K_s^{p_i}$.

By (4) the automorphic representations $(\pi^{p_i})_{1 \leq i \leq r}$ all contribute to $q(X_r)$ and since they are pairwise distinct we have that $q(X_r) \geq r$, for all r .

Although X_r may have several connected components, it is important to observe that the connected components remain constant as r grows, since by the above claim $\det(K_s^p) = \det(K_s)$ for all $p \in \mathcal{P}$. The proposition then follows immediately. \square

Remark 4.2. (a) The computation of an exact level K at which the automorphic representation π attached to a pair of characters (λ, χ) contributes to the H^1 is analyzed in detail in [DR] when M is an imaginary quadratic

- field. In particular, if λ has ramification only at the primes p dividing the discriminant of M , we check that the level subgroup at any such p is precisely the one conjectured by B. Gross, namely the index 2 subgroup of the maximal parahoric subgroup with reductive quotient $\mathrm{PGL}(2)$.
- (b) One consequence of Rogawski's theory is that the Albanese variety is of CM type for any congruence subgroup (see [MR]). If M is imaginary quadratic of discriminant a prime p , we will show in [DR] that the factor of the Albanese corresponding to the minimally ramified λ turns out (at an appropriate prime to p level) to be isogenous to the CM abelian variety $B(p)$ defined by B. Gross in [G].
 - (c) When Γ is not a congruence subgroup, there are examples of C. Schoen where the Albanese is not of CM type (see [S]).

5. MORDELLICITY OF X

We will first prove that Theorem 0.1(i) follows from Theorem 0.1(ii) applied to a neat congruence subgroup.

Note that in order to check that a surface X over a number field k is Mordellic, it suffices to make a finite base change k'/k and show that X is Mordellic over k' . This will be useful as we will consider diagrams of finite covers, where the common field of definition will be a finite extension of the field we start with. Note further that if X_Γ is Mordellic, then $X_{\Gamma'}$ is Mordellic for any finite index subgroup Γ' of Γ .

By Proposition 4.1 there exists a finite index subgroup Γ' of $\Gamma(3)$ such that $q(X_{\Gamma'}) > 2$ and by lemmas 1.2 and 1.3 the variety $X_{\Gamma'}$ is smooth. It follows then from Theorem 0.1(ii) that $X_{\Gamma'}$ is Mordellic. Since Γ is by hypothesis arithmetic, the intersection $\Gamma'' = \Gamma \cap \Gamma'$ has finite index in both Γ and Γ' . Hence $X_{\Gamma''}$, being a finite cover of $X_{\Gamma'}$, is also Mordellic.

It remains to prove part (ii) of our Theorem 0.1. Let k be a number field over which X_Γ is defined. By Proposition 3.1, $X(k)$ is contained (up to a finite number of points) in a finite union of curves on X_Γ . After replacing k by a finite extension, we may assume that any irreducible curve C occurring in this union is defined over k . If its normalization \tilde{C} is of genus $g \leq 1$, we will get a non-constant holomorphic map from \mathbb{C} to X which is impossible by hyperbolicity. So $g \geq 2$ for each C . Now by Faltings's celebrated proof of Mordell's conjecture [F1], $C(k)$ is finite. Since the set of C is finite, we get, as asserted, that $X(k)$ is finite. \square

Remark 5.1. When Lang originally made his conjecture on Mordellicity, his definition of a variety X over $k \subset \mathbb{C}$ being hyperbolic required the Kobayashi semi-distance on $X(\mathbb{C})$ to be in fact a metric. Later it was established by R. Brody [B] that this was equivalent to requiring that there is no non-constant holomorphic map from \mathbb{C} to $X(\mathbb{C})$. Thus any quotient $\Gamma \backslash \mathcal{H}_{\mathbb{C}}^2$ is hyperbolic. It is expected that every surface X of general type over $k \subset \mathbb{C}$ containing no curve of genus ≤ 1 is hyperbolic.

REFERENCES

- [B] R. BRODY, *Compact manifolds and hyperbolicity*, Trans. Amer. Math. Soc., 235 (1978), pp. 213–219.
- [C] C. CHEVALLEY, *Deux théorèmes d’arithmétique*, J. Math. Soc. Japan, 3 (1951), pp. 36–44.
- [DR] M. DIMITROV AND D. RAMAKRISHNAN, *A finiteness theorem for rational points on certain Picard modular surfaces*, in preparation (2013).
- [F1] G. FALTINGS, *Endlichkeitssätze für abelsche Varietäten über Zahlkörpern*, Invent. Math., 73 (1983), pp. 349–366.
- [F2] ———, *The general case of S. Lang’s conjecture*, in Barsotti Symposium in Algebraic Geometry (Abano Terme, 1991), vol. 15 of Perspect. Math., Academic Press, San Diego, CA, 1994, pp. 175–182.
- [G] B. H. GROSS, *Arithmetic on elliptic curves with complex multiplication*, vol. 776 of Lecture Notes in Mathematics, Springer, Berlin, 1980. With an appendix by B. Mazur.
- [H] F. HIRZEBRUCH, *Automorphe Formen und der Satz von Riemann-Roch*, In Symposium internacional de topología algebraica International symposium on algebraic topology, Universidad Nacional Autónoma de México and UNESCO, Mexico City, 1958, pp. 129–144.
- [K] D. KAZHDAN, *Some applications of the Weil representation*, J. Analyse Mat., 32 (1977), pp. 235–248.
- [L] S. LANG, *Number theory. III*, vol. 60 of Encyclopaedia of Mathematical Sciences, Springer-Verlag, Berlin, 1991. Diophantine geometry.
- [La] R. P. LANGLANDS, *Les débuts d’une formule des traces stable*, vol. 13 of Publications Mathématiques de l’Université Paris VII [Mathematical Publications of the University of Paris VII], Université de Paris VII U.E.R. de Mathématiques, Paris, 1983.
- [M] Y. MIYAOKA, *The maximal number of quotient singularities on surfaces with given numerical invariants*, Math. Ann., 268 (1984), pp. 159–171.
- [MM] S. MORI AND S. MUKAI, *The uniruledness of the moduli space of curves of genus 11*, in Algebraic geometry (Tokyo/Kyoto, 1982), vol. 1016 of Lecture Notes in Math., Springer, Berlin, 1983, pp. 334–353.
- [MR] V. K. MURTY AND D. RAMAKRISHNAN, *The Albanese of unitary shimura varieties*, in The zeta functions of Picard modular surfaces, R. Langlands and D. Ramakrishnan, eds., Les Publications CRM, Montreal, 1992, pp. 445–464.

- [NR] M. V. NORI AND M. S. RAGHUNATHAN, *On conjugation of locally symmetric arithmetic varieties*, in Proceedings of the Indo-French Conference on Geometry (Bombay, 1989), Delhi, 1993, Hindustan Book Agency, pp. 111–122.
- [RZ] M. RAPOPORT AND T. ZINK, *Über die lokale Zetafunktion von Shimuravarietäten. Monodromiefiltration und verschwindende Zyklen in ungleicher Charakteristik*, Invent. Math., 68 (1982), pp. 21–101.
- [R1] J. D. ROGAWSKI, *Automorphic representations of unitary groups in three variables*, vol. 123 of Annals of Mathematics Studies, Princeton University Press, Princeton, NJ, 1990.
- [R2] ———, *The multiplicity formula for A -packets*, in The zeta functions of Picard modular surfaces, Univ. Montréal, Montreal, QC, 1992, pp. 395–419.
- [S] C. SCHOEN, *An arithmetic ball quotient surface whose Albanese variety is not of CM type*, preprint (2013).
- [Sh] G. SHIMURA, *Automorphic forms and the periods of abelian varieties*, J. Math. Soc. Japan, 31 (1979), pp. 561–592.
- [T] J. TATE, *Introduction [Brief biography of Serge Lang]*, in Number theory, analysis and geometry, Springer, New York, 2012, pp. xv–xx.
- [V] P. VOJTA, *Arithmetic of subvarieties of abelian and semiabelian varieties*, in Advances in number theory (Kingston, ON, 1991), Oxford Sci. Publ., Oxford Univ. Press, New York, 1993, pp. 233–238.
- [Y] S. T. YAU, *Calabi’s conjecture and some new results in algebraic geometry*, Proc. Nat. Acad. Sci. U.S.A., 74 (1977), pp. 1798–1799.

UNIVERSITÉ LILLE 1, UMR CNRS 8524, UFR MATHÉMATIQUES, 59655 VILLENEUVE
D’ASCQ CEDEX, FRANCE

E-mail address: mladen.dimitrov@gmail.com

MATHEMATICS 253-37, CALTECH, PASADENA, CA 91125, USA

E-mail address: dinakar@caltech.edu