

The normal matrix model with a monomial potential, a vector equilibrium problem, and multiple orthogonal polynomials on a star

Arno B.J. Kuijlaars* and Abey López-García†

September 10, 2018

Abstract

We investigate the asymptotic behavior of a family of multiple orthogonal polynomials that is naturally linked with the normal matrix model with a monomial potential of arbitrary degree $d + 1$. The polynomials that we investigate are multiple orthogonal with respect to a system of d analytic weights defined on a symmetric $(d + 1)$ -star centered at the origin. In the first part we analyze in detail a vector equilibrium problem involving a system of d interacting measures (μ_1, \dots, μ_d) supported on star-like sets in the plane. We show that in the subcritical regime, the first component μ_1^* of the solution to this problem is the asymptotic zero distribution of the multiple orthogonal polynomials. It also characterizes the domain where the eigenvalues in the normal matrix model accumulate, in the sense that the Schwarz function associated with the boundary of this domain can be expressed explicitly in terms of μ_1^* . The second part of the paper is devoted to the asymptotic analysis of the multiple orthogonal polynomials. The asymptotic results are obtained again in the subcritical regime, and they follow from the Deift/Zhou steepest descent analysis of a Riemann-Hilbert problem of size $(d + 1) \times (d + 1)$. The vector equilibrium problem and the Riemann-Hilbert problem that we investigate are generalizations of those studied recently by Bleher-Kuijlaars in the case $d = 2$.

Keywords: Multiple orthogonal polynomial, vector equilibrium problem, normal matrix model, Riemann-Hilbert problem.

Contents

1	Introduction	3
1.1	Normal matrix model and Laplacian growth	3
1.2	Approach based on sesquilinear forms	4
1.3	Aim of the paper	6
1.4	From 2D orthogonality to orthogonality on contours	6
1.5	Orthogonality on $(d + 1)$ -star and multiple orthogonality	8

*Department of Mathematics, University of Leuven (KU Leuven), Celestijnenlaan 200B, B-3001 Leuven, Belgium. email: arno.kuijlaars@wis.kuleuven.be

†Department of Mathematics and Statistics, University of South Alabama, 411 University Blvd North, ILB 325, Mobile, AL 36688, USA, email: lopezgarcia@southalabama.edu

2	Statement of results	10
2.1	The vector equilibrium problem	10
2.2	Subcritical case	11
2.3	Asymptotics of $P_{n,n}$ in subcritical case	12
2.4	Laplacian growth and the spectral curve	13
2.5	Outline of the paper	14
3	Proofs of Theorems 2.2, 2.6 and 2.7	14
3.1	The Riemann surface in the subcritical case	14
3.2	Asymptotics of functions ξ_k	17
3.3	Construction of the vector equilibrium measure in terms of the functions ξ_k	18
3.4	Proof of Proposition 3.3	20
3.4.1	Total masses	20
3.4.2	Positivity of μ_1^*	21
3.4.3	Identities for Cauchy transforms	23
3.4.4	The variational equalities	24
3.4.5	Positivity of measures μ_k^* with $k \geq 2$	25
3.5	Conclusion of the proof of Proposition 3.3	26
3.6	Proof of Theorem 2.2	26
3.7	Proof of Theorem 2.6	27
3.8	Proof of Theorem 2.7	28
4	The multiple orthogonal polynomials $P_{n,n}$ and the associated Riemann-Hilbert problem	30
4.1	The Riemann-Hilbert problem	31
5	First transformation $\mathbf{Y} \mapsto \mathbf{X}$	32
5.1	Definition and properties of \mathbf{P}	33
5.2	Definition and properties of \mathbf{F}	36
5.3	Definition and properties of \mathbf{Q}	38
5.4	Transformation $\mathbf{Y} \mapsto \mathbf{X}$	39
6	Second transformation $\mathbf{X} \mapsto \mathbf{U}$	40
7	The third transformation $\mathbf{U} \mapsto \mathbf{T}$	42
7.1	The g -functions	42
7.2	The φ -functions	44
7.3	Transformation $\mathbf{U} \mapsto \mathbf{T}$	47
8	Fourth transformation $\mathbf{T} \mapsto \mathbf{S}$	49
9	Global parametrix	56
10	The final transformation $\mathbf{S} \mapsto \mathbf{R}$ and conclusion of the steepest descent analysis	60
10.1	Proof of Theorem 2.3	62
11	Acknowledgements	63

1 Introduction

1.1 Normal matrix model and Laplacian growth

The starting point of this work is the analysis of the *normal matrix model*. This is a probability measure on the space of $n \times n$ normal matrices M of the form

$$\frac{1}{Z_n} e^{-n \operatorname{Tr} \mathcal{V}(M)} \mathrm{d}M. \quad (1.1)$$

A standard expression for the potential \mathcal{V} in (1.1) is

$$\mathcal{V}(M) = \frac{1}{t_0} (MM^* - V(M) - \overline{V}(M^*)), \quad t_0 > 0, \quad (1.2)$$

where V is a polynomial and \overline{V} is the polynomial obtained from V by conjugating the coefficients. This model has attracted considerable interest in recent years and important connections have been established with other problems in mathematics and physics, most notably the study of Laplacian growth [27, 24, 34, 36], two-dimensional Coulomb gases [15, 19] and Toda lattice systems [21], see [38] for an overview. Recently, eigenvalue statistics in the normal matrix model are considered in [3, 4, 30] and connections with orthogonal polynomials on the complex plane are also in [5, 14, 20, 33].

According to the theory that has been mainly developed by Wiegmann and Zabrodin, the eigenvalues of a matrix M with distribution (1.1)–(1.2) fill out, as $n \rightarrow \infty$, a two-dimensional domain Ω (the *droplet*) with uniform density. Moreover, the boundary of Ω evolves, as the *time* parameter $t_0 > 0$ increases, according to the model of Laplacian growth, see [21]. At a critical time for t_0 , the boundary of Ω is expected to develop singularities and a breakdown takes place in the Laplacian growth evolution.

The model (1.1)–(1.2) requires a certain regularization in order to be well-defined since the integral

$$Z_n = \int e^{-n \operatorname{Tr} \mathcal{V}(M)} \mathrm{d}M \quad (1.3)$$

diverges for any polynomial V of degree ≥ 3 . A natural approach is to use a cut-off, as proposed by Elbau and Felder [14, 15]. This approach consists of restricting the model to those normal matrices with spectrum confined in a fixed two-dimensional bounded domain containing the droplet Ω . Another possible approach is to modify V outside of the droplet in such a way that (1.3) converges, while still keeping the essential features of the model (which take place on the droplet). This is the approach followed by Ameur-Hedenmalm-Makarov [3, 4, 19].

In these models the eigenvalues of M are distributed according to a determinantal point process with a correlation kernel that is constructed in terms of polynomials that have a two-dimensional orthogonality. In the cut-off model of [14, 15] with cut-off domain D the orthogonality is associated with the scalar product (depending on n)

$$\langle f, g \rangle_D = \iint_D f(z) \overline{g(z)} e^{-n \mathcal{V}(z)} \mathrm{d}A(z), \quad (1.4)$$

where

$$\mathcal{V}(z) = \frac{1}{t_0} (|z|^2 - V(z) - \overline{V(z)}), \quad t_0 > 0, \quad (1.5)$$

and $\mathrm{d}A$ denotes area measure on D . For each n , if $(Q_{k,n})_{k=0}^\infty$ is the sequence of monic polynomials (i.e., $Q_{k,n}(z) = z^k + \dots$) satisfying

$$\langle Q_{k,n}, Q_{j,n} \rangle_D = h_{k,n} \delta_{j,k}, \quad (1.6)$$

then the correlation kernel for the determinantal point process is given by

$$K_n(w, z) = e^{-\frac{n}{2}(\mathcal{V}(z) + \mathcal{V}(w))} \sum_{k=0}^{n-1} \frac{Q_{k,n}(z) \overline{Q_{k,n}(w)}}{h_{k,n}}.$$

Elbau and Felder also showed in [14, 15] that for any polynomial V of the form

$$V(z) = \sum_{k=1}^{d+1} \frac{t_k}{k} z^k, \quad t_k \in \mathbb{C}, \quad (1.7)$$

with $t_1 = 0$ and $|t_2| < 1$, there exists a compact domain D with 0 in its interior such that for all $t_0 > 0$ small enough, the eigenvalues of M in the model with cut-off D indeed accumulate in a domain $\Omega \subset D$ as $n \rightarrow \infty$. The boundary $\partial\Omega$ of $\Omega = \Omega(t_0; t_1, \dots, t_{d+1})$ is moreover characterized as the only positively oriented polynomial curve of degree at most d satisfying

$$\frac{1}{2\pi i} \oint_{\partial\Omega} \frac{\bar{z}}{z^k} dz = \begin{cases} t_k, & k \in \{1, \dots, d+1\}, \\ 0, & k \in \mathbb{N} \setminus \{1, \dots, d+1\}, \end{cases} \quad (1.8)$$

and enclosing a domain with area πt_0 . The equations (1.8) are characteristic for the model of Laplacian growth [36].

1.2 Approach based on sesquilinear forms

Motivated by certain boundary integral estimates in the cut-off model and an analysis of the algebraic properties of the scalar product (1.4), Bleher and Kuijlaars introduced in [7] a new construction which replaces the cut-off domain D by a system of unbounded contours in the complex plane. This construction leads to a sequence of sesquilinear forms $\langle \cdot, \cdot \rangle_n : \mathcal{P} \times \mathcal{P} \rightarrow \mathbb{C}$ defined on the vector space \mathcal{P} of all polynomials in one variable. Such sesquilinear forms are defined as follows. Given an integer $d \geq 2$, let us consider the directions at infinity

$$\infty_\ell = e^{\frac{(2\ell+1)\pi i}{d+1}} \infty, \quad \ell = 0, \dots, d, \quad (1.9)$$

and let Γ_ℓ denote any unbounded oriented contour from $\infty_{\ell-1}$ to ∞_ℓ , see Figure 1 for the case $d = 3$. By $\bar{\Gamma}_\ell$ we indicate the image of Γ_ℓ under the map $z \mapsto \bar{z}$, with the induced orientation. Then, given $t_0 > 0$ and a polynomial V as in (1.7) with $t_{d+1} > 0$, the sesquilinear forms are defined by

$$\langle f, g \rangle_n = \sum_{j=0}^d \sum_{k=0}^d C_{j,k} \int_{\Gamma_j} dz \int_{\bar{\Gamma}_k} dw f(z) \bar{g}(w) e^{-\frac{n}{t_0}(wz - V(z) - \bar{V}(w))}, \quad f, g \in \mathcal{P}, \quad (1.10)$$

where \bar{g} is the polynomial obtained from g by conjugating the coefficients and $C = (C_{j,k})_{j,k=0}^d$ is a complex matrix of coefficients.

The matrix C is naturally assumed in [7] to be Hermitian and circulant, in order to guarantee the hermiticity and rotational invariance (with respect to the angle $2\pi/(d+1)$) of the sesquilinear forms. In the cubic monomial case

$$V(z) = \frac{t_3}{3} z^3, \quad t_3 > 0, \quad (1.11)$$

an asymptotic analysis of the orthogonal polynomials associated with the corresponding sesquilinear forms was performed in [7], for a suitable choice of the matrix C . The choice of C is governed




Figure 1: The contours Γ_ℓ and the points at infinity ∞_ℓ for $\ell = 0, 1, \dots, d$ in the case $d = 3$.

by the desire to recover the main features of the normal matrix model. In the cubic case (1.11), it was shown in [7] that it is indeed possible to find a matrix $C = (C_{j,k})_{j,k=0}^2$ such that, in a subcritical regime, the orthogonal polynomials associated with the corresponding sesquilinear forms have the same asymptotic behavior as the orthogonal polynomials associated with the corresponding scalar product (1.4)–(1.5).

The study in [7] involved a characterization of the orthogonal polynomials as multiple orthogonal polynomials with respect to two orthogonality weights defined on the sets Γ_0 , Γ_1 and Γ_2 , and built out of Airy functions. The multiple orthogonality implies a characterization of the orthogonal polynomials in terms of a 3×3 matrix valued Riemann-Hilbert problem, which is analyzed in the large n limit. A major role in the asymptotic analysis of [7] is played by a vector equilibrium problem for two measures and an associated three sheeted Riemann surface. The first component μ_1^* of the minimizer of the vector equilibrium problem gives the limiting distribution of the zeros of the orthogonal polynomials. The support of this measure is a three-star

$$\text{supp}(\mu_1^*) = [0, x^*] \cup [0, e^{\frac{2\pi i}{3}} x^*] \cup [0, e^{\frac{4\pi i}{3}} x^*]$$

for a certain x^* that depends on t_0 and t_3 . It is also shown in [7] how the domain Ω that evolves according to the Laplacian growth can be recovered from μ_1^* . This only applies to the subcritical regime, that is for t_0 less than a critical value $t_{0,\text{crit}}$, depending on $t_3 > 0$.

This approach based on sesquilinear forms and multiple orthogonality is extended to the supercritical regime in the recent work [23], where it is found that an analogue of the domain Ω can still be defined but this domain is no longer growing as t_0 increases, but instead it shrinks to a point at a second critical value. Lee et al. [25] argue that the Painlevé I equation provides a model for the shock phenomenon at the critical value $t_{0,\text{crit}}$. This connection with Painlevé I is also likely to appear, if one would try to extend the Riemann-Hilbert analysis from [7] to the critical case. Indeed, it would then be necessary to construct a local parametrix out of the Lax pair solutions associated with Painlevé I, as it is done for example in [11]. However, this has not been worked out yet.

1.3 Aim of the paper

In this paper we study monomial V of higher degree, that is,

$$V(z) = \frac{t_{d+1}}{d+1} z^{d+1}, \quad t_{d+1} > 0, \quad (1.12)$$

where $d \geq 3$. Our initial expectation was that the approach of [7] for the case $d = 2$ would carry through, with proper but non-essential modifications. That is, we expected a formulation of the orthogonal polynomials as multiple orthogonal polynomials with d weights built out of solutions of the differential equation

$$p^{(d)}(z) = (-1)^d z p(z) \quad (1.13)$$

of order d , which for $d = 2$ is the Airy differential equation. This gives rise to a RH problem of size $(d+1) \times (d+1)$. An ingredient for the asymptotic analysis of this RH problem would be an appropriate vector equilibrium problem with d measures which is related to a $(d+1)$ -sheeted Riemann surface. The first component μ_1^* of the minimizer of the vector equilibrium problem would be the limiting zero distribution of the orthogonal polynomials. The support of μ_1^* is a $(d+1)$ -star

$$\text{supp}(\mu_1^*) = \bigcup_{j=0}^d [0, \omega^j x^*], \quad \omega = \exp\left(\frac{2\pi i}{d+1}\right)$$

for some $x^* > 0$, and from μ_1^* one would recover the domain $\Omega = \Omega(t_0; 0, \dots, 0, t_{d+1}, 0, \dots)$ satisfying (1.8) and evolving according to the Laplacian growth model.

It came somewhat as a surprise to us that for the case $d = 3$ we could no longer select a circulant Hermitian matrix $C = (C_{j,k})_{j,k=0}^3$ that gives a sesquilinear form (1.10) for which we can do an asymptotic analysis of the corresponding orthogonal polynomials in the way described above. In addition to being circulant and Hermitian, the matrix C should give rise to multiple orthogonality involving on the interval $[0, x^*]$ the recessive solution of (1.13) as $z \rightarrow \infty$, $\arg z = 0$. This yields a number of conditions on the coefficients $C_{j,k}$ which turned out to be incompatible with the conditions to be circulant and Hermitian. We do not have a conceptual proof why this happens.

1.4 From 2D orthogonality to orthogonality on contours

For this reason, we had to devise another approach, which is inspired by the work of Balogh et al. [5], and consists of first transforming the two-dimensional orthogonality on D given by (1.4)–(1.5) into orthogonality over contours by means of Green's theorem. The contours consist of ∂D together with a certain number of contours within D that for the monomial potential (1.12) we take as a $(d+1)$ -star

$$\Sigma := \{z \in D \mid z^{d+1} \in \mathbb{R}^+\}, \quad (1.14)$$

see Figure 2. We let

$$\omega := \omega_{d+1} = \exp\left(\frac{2\pi i}{d+1}\right) \quad (1.15)$$

be the primitive $(d+1)$ -st root of unity, and this notation will be used throughout the paper. We also continue to use the notation ∞_ℓ as in (1.9) and the contours Γ_ℓ that appear in (1.10), where ℓ is considered modulo $d+1$, so that

$$\infty_{-\ell} = \infty_{d+1-\ell}, \quad \Gamma_{-\ell} = \Gamma_{d+1-\ell}.$$

Proposition 1.1. *Let $V(z)$ be the monomial (1.12) with $d \geq 2$. Let D be a simply connected two-dimensional Jordan domain with 0 in its interior, that is invariant under rotation $z \mapsto \omega z$, as well as under reflection $z \mapsto \bar{z}$ in the real axis and let Σ be as in (1.14). Then, for a polynomial Q and an integer $j \geq 0$, we have*

$$2i \iint_D Q(z) \bar{z}^j e^{-\frac{n}{t_0}(|z|^2 - V(z) - \overline{V(z)})} dA(z) = \int_{\Sigma} Q(z) w_{j,n}(z) dz + \oint_{\partial D} Q(z) \tilde{w}_{j,n}(z) dz, \quad (1.16)$$

where the functions $w_{j,n}(z)$ and $\tilde{w}_{j,n}(z)$ have the following expressions:

$$w_{j,n}(z) = \int_{\Gamma_{-\ell}} s^j e^{-\frac{n}{t_0}(sz - V(s) - V(z))} ds, \quad \arg z = \frac{2\pi}{d+1}\ell, \quad (1.17)$$

$$\tilde{w}_{j,n}(z) = \int_{\infty-\ell-1}^{\bar{z}} s^j e^{-\frac{n}{t_0}(sz - V(s) - V(z))} ds, \quad \frac{2\pi}{d+1}\ell < \arg z < \frac{2\pi}{d+1}(\ell+1), \quad (1.18)$$

where $\ell = 0, \dots, d+1$. In (1.16), each segment of Σ is given the outward orientation (i.e., away from 0) and ∂D is given the positive counterclockwise orientation.

Proof. Let us define the functions

$$F_{j,\ell}(z) = \int_{\infty-\ell-1}^{\bar{z}} s^j e^{-\frac{n}{t_0}(sz - V(s))} ds, \quad j \geq 0, \quad 0 \leq \ell \leq d. \quad (1.19)$$

Let D_ℓ be the part of D given by

$$D_\ell = \{z \in D \mid \frac{2\pi}{d+1}\ell < \arg z < \frac{2\pi}{d+1}(\ell+1)\}. \quad (1.20)$$

Noting that

$$\frac{\partial F_{j,\ell}}{\partial \bar{z}} = \bar{z}^j e^{-\frac{n}{t_0}(|z|^2 - \overline{V(z)})}$$

we find by Green's theorem in the complex plane

$$\begin{aligned} \iint_{D_\ell} Q(z) \bar{z}^j e^{-\frac{n}{t_0}(|z|^2 - V(z) - \overline{V(z)})} dA(z) &= \iint_{D_\ell} Q(z) e^{\frac{n}{t_0}V(z)} \frac{\partial F_{j,\ell}(z)}{\partial \bar{z}} dA(z) \\ &= \frac{1}{2i} \oint_{\partial D_\ell} Q(z) e^{\frac{n}{t_0}V(z)} F_{j,\ell}(z) dz. \end{aligned} \quad (1.21)$$

The boundary ∂D_ℓ consists of the part of ∂D with $\frac{2\pi}{d+1}\ell < \arg z < \frac{2\pi}{d+1}(\ell+1)$ and the parts in Σ with $\arg z = \frac{2\pi}{d+1}\ell$ and $\arg z = \frac{2\pi}{d+1}(\ell+1)$, the latter one traversed towards the origin.

We take the sum of (1.21) for $\ell = 0, \dots, d$ and we find that the contribution of ∂D is

$$\frac{1}{2i} \oint_{\partial D} Q(z) \tilde{w}_{j,n}(z) dz \quad (1.22)$$

since $e^{\frac{n}{t_0}V(z)} F_{j,\ell}(z) = \tilde{w}_{j,n}(z)$ for $z \in \partial D_\ell \setminus \Sigma$, see (1.18) and (1.19).

Each segment in Σ is part of the boundary of two subdomains, say D_ℓ and $D_{\ell-1}$ for some ℓ . We observe by (1.17) and (1.19) that

$$e^{\frac{n}{t_0}V(z)} (F_{j,\ell}(z) - F_{j,\ell-1}(z)) = w_{j,n}(z) \quad \text{for } \arg z = \frac{2\pi}{d+1}\ell,$$




Figure 2: A possible domain D and the star Σ in case $d = 3$. The star has endpoints $\omega^\ell \hat{x}$ for $\ell = 0, \dots, d$.

since by definition $\Gamma_{-\ell}$ is a contour from $\infty_{-\ell-1}$ to $\infty_{-\ell}$. Thus the total contribution of Σ in the sum of (1.21)

$$\frac{1}{2i} \int_{\Sigma} Q(z) w_{j,n}(z) dz. \quad (1.23)$$

In total we get the sum of (1.22) and (1.23), which proves (1.16). \square

It is a consequence of Proposition 1.1 that the orthogonal polynomials $Q_{k,n}$ also satisfy

$$\int_{\Sigma} Q_{k,n}(z) w_{j,n}(z) dz + \oint_{\partial D} Q_{k,n}(z) \tilde{w}_{j,n}(z) dz = 0, \quad j = 0, \dots, k-1. \quad (1.24)$$

1.5 Orthogonality on $(d+1)$ -star and multiple orthogonality

Guided by the idea that in the subcritical case, the domain D can be chosen in such a way that boundary contributions are negligible in the large n limit, we now take the approach to drop the integral over ∂D , and consider instead of the polynomials $Q_{k,n}$ a new set of polynomials $P_{k,n}$ that are orthogonal on the $(d+1)$ -star Σ with respect to the weights (1.17). Since these polynomials will be the focus of the present paper we introduce them with the following formal definition.

Definition 1.2. Let $d \geq 2$, and $\hat{x}, t_0, t_{d+1} > 0$. Then for every $k, n \in \mathbb{N}$, we let $P_{k,n}(z) = z^k + \dots$ be the monic polynomial that satisfies

$$\int_{\Sigma} P_{k,n}(z) w_{j,n}(z) dz = 0, \quad j = 0, \dots, k-1, \quad (1.25)$$

where

$$\Sigma = \bigcup_{j=0}^d [0, \omega^j \hat{x}]$$

and with $w_{j,n}(z)$ given in (1.17).

The polynomials depend on the endpoint \widehat{x} of the star. In what follows we are going to take it in a suitable way, depending on t_0 and t_{d+1} .

Note that $w_{j,n}$ is analytic on each segment of Σ . If we put

$$p_\ell(z) = \frac{1}{2\pi i} \int_{\Gamma_\ell} e^{-sz + \frac{1}{d+1}s^{d+1}} ds \quad (1.26)$$

then it is easy to see from (1.17) that

$$w_{0,n}(z) = 2\pi i d_n e^{\frac{n}{t_0}V(z)} p_{-\ell}(c_n z), \quad \arg z = \frac{2\pi}{d+1}\ell. \quad (1.27)$$

with constants

$$c_n = \left(\frac{n^d}{t_0^d t_{d+1}} \right)^{\frac{1}{d+1}}, \quad d_n = \left(\frac{t_0}{n t_{d+1}} \right)^{\frac{1}{d+1}}, \quad (1.28)$$

and for every $j \geq 1$,

$$w_{j,n}(z) = 2\pi i (-1)^j d_n^{j+1} e^{\frac{n}{t_0}V(z)} p_{-\ell}^{(j)}(c_n z), \quad \arg z = \frac{2\pi}{d+1}\ell. \quad (1.29)$$

The functions (1.26) satisfy (1.13) and $p_{-\ell}$ is the recessive solution of (1.13) as $z \rightarrow \infty$ in the sector $\frac{2\pi}{d+1}(\ell - \frac{1}{2}) < \arg z < \frac{2\pi}{d+1}(\ell + \frac{1}{2})$.

Inserting the definition (1.17) of $w_{j,n}$ we can rewrite the orthogonality (1.25) in terms of a double integral

$$\sum_{\ell=0}^d \int_0^{\omega^\ell \widehat{x}} dz \int_{\Gamma_{-\ell}} ds P_{k,n}(z) s^j e^{-\frac{n}{t_0}(sz - V(z) - V(s))} = 0, \quad j = 0, 1, \dots, k-1.$$

This can be viewed as orthogonality with respect to a sesquilinear form

$$\langle f, g \rangle = \sum_{\ell=0}^d \int_0^{\omega^\ell \widehat{x}} dz \int_{\Gamma_{-\ell}} ds f(z) \bar{g}(s) e^{-\frac{n}{t_0}(sz - V(z) - V(s))}, \quad (1.30)$$

comparable to (1.10). The new sesquilinear form (1.30) is not Hermitian, but it does satisfy the structure relation

$$t_0 \langle f, g' \rangle - n \langle z f, g \rangle + n \langle f, V' g \rangle = 0, \quad (1.31)$$

as can be seen from an integration by parts on the s -integral in (1.30), cf. also [7, section 2.1].

As in [7] it follows from (1.31) that the polynomials $P_{n,n}$ are multiple orthogonal with respect to the system of d weights $\{w_{j,n}\}_{j=0}^{d-1}$.

Lemma 1.3. *The polynomials $P_{n,n}$ are characterized by the multiple orthogonality conditions*

$$\int_{\Sigma} P_{n,n}(z) z^k w_{j,n}(z) dz = 0, \quad k = 0, \dots, \left\lceil \frac{n-j}{d} \right\rceil - 1, \quad j = 0, \dots, d-1. \quad (1.32)$$

Proof. This follows from the structure relation (1.31) in the same manner as in [7, Lemma 5.1]. \square

Multiple orthogonal polynomials on star-like sets have been studied in recent years under different frameworks. Some references on this subject are [1, 2, 10, 26, 31]. See also [13, 18] for the related study of Faber polynomials associated with hypocycloidal domains and stars.

Lemma 1.3 allows us to obtain the asymptotic properties of the polynomials $P_{n,n}$ through the analysis of a matrix-valued Riemann-Hilbert problem of size $(d+1) \times (d+1)$ that encodes the multiple orthogonality conditions (1.32). The first characterization of multiple orthogonality in terms of a RH problem appeared in [35]. The RH problem we will analyze is described in Section 4.

The analysis is basically along the same lines as in [7]. As in [7] a major role is played by a vector equilibrium problem that we will describe first.

2 Statement of results

In this section we state our main results.

2.1 The vector equilibrium problem

In this paper we will obtain the asymptotic behavior of the polynomials $P_{n,n}$ as $n \rightarrow \infty$ in a *subcritical* regime. It is characterized through the solution of a vector equilibrium problem. In order to describe this vector equilibrium problem we need to introduce the star-like sets, where we recall that $\omega = \omega_{d+1} = \exp(2\pi i/(d+1))$,

$$\Sigma_1 := \bigcup_{\ell=0}^d [0, \omega^\ell \hat{x}], \quad \hat{x} > 0, \quad (2.1)$$

and for $2 \leq k \leq d$,

$$\Sigma_k := \begin{cases} \{z \in \mathbb{C} \mid z^{d+1} \in \mathbb{R}^-\} & \text{if } k \text{ is even,} \\ \{z \in \mathbb{C} \mid z^{d+1} \in \mathbb{R}^+\} & \text{if } k \text{ is odd.} \end{cases} \quad (2.2)$$

Thus Σ_k is an unbounded set consisting of $d+1$ halfrays for each $k \geq 2$, and they alternate between Σ_2 and Σ_3 . Note also that $\Sigma_1 \subset \Sigma_3$.

Given two positive measures μ, ν , we use the standard notations from logarithmic potential theory

$$I(\mu) = \iint \log \frac{1}{|x-y|} d\mu(x) d\mu(y), \quad I(\mu, \nu) = \iint \log \frac{1}{|x-y|} d\mu(x) d\nu(y),$$

for the logarithmic energy of μ and the mutual logarithmic energy of μ and ν , respectively. Main references in logarithmic potential theory are [8, 28, 29, 32].

Definition 2.1. Fix $\hat{x}, t_0, t_{d+1} > 0$. The vector equilibrium problem asks for the minimization of the energy functional

$$\sum_{k=1}^d I(\mu_k) - \sum_{k=1}^{d-1} I(\mu_k, \mu_{k+1}) + \frac{1}{t_0} \int \left(\frac{d}{d+1} \frac{1}{t_{d+1}^{1/d}} |z|^{\frac{d+1}{d}} - \frac{t_{d+1}}{d+1} z^{d+1} \right) d\mu_1(z), \quad (2.3)$$

among all positive Borel measures μ_1, \dots, μ_d satisfying the following conditions:

- (1) For each $k = 1, \dots, d$, the measure μ_k has total mass

$$\|\mu_k\| = 1 - \frac{k-1}{d}. \quad (2.4)$$

(2) For each $k = 1, \dots, d$,

$$\text{supp}(\mu_k) \subset \Sigma_k, \quad (2.5)$$

where the sets Σ_k are given in (2.1)–(2.2).

Observe that this vector equilibrium problem depends exclusively on $\hat{x}, t_0, t_{d+1} > 0$. For any choice of these parameters, this vector equilibrium problem is weakly admissible in the sense of [17], and therefore it admits a unique minimizer $(\mu_1^*, \dots, \mu_d^*)$. See also [6] for a similar class of vector equilibrium problems. As we indicated before, in this paper we are interested in the solution of the stated problem under special assumptions on the parameters involved, that is, in a subcritical situation. The vector equilibrium problem was analyzed in [7] in the case $d = 2$.

2.2 Subcritical case

Theorem 2.2. *Let $d \geq 2$ be an arbitrary integer. Fix $t_{d+1} > 0$ and set*

$$t_{0,\text{crit}} = (d^{-\frac{2}{d-1}} - d^{-\frac{d+1}{d-1}}) t_{d+1}^{-\frac{2}{d-1}} > 0. \quad (2.6)$$

Let $0 < t_0 < t_{0,\text{crit}}$ and define

$$x^* = (d+1) d^{-\frac{d}{d+1}} t_{d+1}^{\frac{1}{d+1}} r^{\frac{2d}{d+1}}, \quad (2.7)$$

where r denotes the smallest positive solution of the equation

$$t_0 = r^2 - d t_{d+1}^2 r^{2d} \quad (2.8)$$

(which exists because of $t_0 < t_{0,\text{crit}}$). Then there is an $x^{**} > x^*$ such that for every $\hat{x} \in [x^*, x^{**}]$, the unique minimizer $(\mu_1^*, \dots, \mu_d^*)$ of the vector equilibrium problem of Definition 2.1 satisfies the following.

- (a) All measures are invariant with respect to rotation $z \mapsto \omega z$.
- (b) μ_1^* is supported on

$$\text{supp}(\mu_1^*) = \Sigma_1^* = \bigcup_{j=0}^d [0, \omega^j x^*] \quad (2.9)$$

and the density of μ_1^* is positive on $[0, x^*)$ and vanishes like a square root at x^* .

- (c) For $k = 2, \dots, d$, the measure μ_k^* is supported on

$$\text{supp}(\mu_k^*) = \Sigma_k^* = \Sigma_k, \quad (2.10)$$

see (2.2), with a positive density.

The condition $t_0 < t_{0,\text{crit}}$ represents the subcritical case. In the critical case $t_0 = t_{0,\text{crit}}$, we can take $\hat{x} = x^*$ and then all of the above still holds, except that the density of μ_1^* vanishes to higher order at the endpoints.

2.3 Asymptotics of $P_{n,n}$ in subcritical case

We assume $t_{d+1} > 0$ is fixed and we let $0 < t_0 < t_{0,\text{crit}}$, where $t_{0,\text{crit}}$ is given by (2.6). Let $x^* > 0$ and $r > 0$ be as in (2.7)–(2.8). Then, for a choice of $\hat{x} > x^*$ sufficiently close to x^* we can perform an asymptotic analysis of the RH problem for the polynomials $P_{n,n}$. This analysis occupies a large part of the paper.

As an outcome of the analysis we have strong and uniform asymptotic expansions for the polynomials $P_{n,n}$ as $n \rightarrow \infty$ in all domains in the complex plane. These include the exponential asymptotics in the domain $\mathbb{C} \setminus \Sigma_1^*$, the oscillatory asymptotics on Σ_1^* and turning point asymptotics involving Airy functions at the endpoints $\omega^j x^*$. We do not spell out all results in detail but we only give the exponential asymptotics (which is the simplest), which takes the following form.

Theorem 2.3. *Let $d \geq 2$ be an integer. Assume that $t_{d+1} > 0$, $0 < t_0 < t_{0,\text{crit}}$, and let $x^* > 0$ be given by (2.7)–(2.8). Then, for all $\hat{x} > x^*$ sufficiently close to x^* , the polynomials $P_{n,n}$ exist and are unique for all n large enough that are a multiple of d . The polynomials $P_{n,n}$ satisfy*

$$P_{n,n}(z) = \mathbf{M}_{1,1}(z) e^{ng_1(z)} (1 + O(1/n)), \quad \text{as } n \rightarrow \infty, \quad (2.11)$$

uniformly on compact subsets of $\mathbb{C} \setminus \Sigma_1^*$. Here g_1 is given by

$$g_1(z) = \int \log(z-t) d\mu_1^*(t), \quad z \in \mathbb{C} \setminus \Sigma_1^*,$$

and $\mathbf{M}_{1,1}$ is an analytic function with no zeros in $\mathbb{C} \setminus \Sigma_1^*$.

The Szegő type prefactor $\mathbf{M}_{1,1}$ is denoted this way since it is the $(1,1)$ entry of a certain matrix-valued function \mathbf{M} . This function \mathbf{M} is the solution to a certain Riemann-Hilbert problem described in Section 9. An explicit formula for $\mathbf{M}_{1,1}$ is given in (9.25) in terms of a certain meromorphic differential $\boldsymbol{\eta}$ defined on a $(d+1)$ -sheeted Riemann surface.

Theorem 2.3 is similar to [7, Lemma 6.1] which is for polynomials that are orthogonal with respect to a sesquilinear form (1.10) with $d = 2$. The proof is also along the same lines as the proof in [7]. However, since we are dealing with arbitrary d , some issues arise at a number of places. These issues could be dealt with in an ad hoc way for $d = 2$, but now we have to resolve them more systematically.

The restriction to n being a multiple of d is made for purposes of presentation only. There will be no essential difficulty to extend the proof to general values of n . However, doing so would lengthen the exposition even more (it is already quite long) and also complicate the notation, since one would have to distinguish between $\lfloor \frac{n}{d} \rfloor$ and $\lceil \frac{n}{d} \rceil$ at many places in the paper. Therefore we decided to restrict ourselves to values of n that are multiple of d .

As a corollary of Theorem 2.3 we have the following asymptotic zero distribution for the sequence of polynomials $P_{n,n}$. If P is a polynomial of degree n with zeros $\{z_j\}_{j=1}^n \subset \mathbb{C}$, we let

$$\nu(P) = \frac{1}{n} \sum_{j=1}^n \delta_{z_j}$$

denote the associated normalized zero counting measure.

Corollary 2.4. *Under the same conditions as in Theorem 2.3, we have that all zeros of $P_{n,n}$ accumulate on the star Σ_1^* given by (2.1) as $n \rightarrow \infty$, and the sequence $(\nu(P_{n,n}))$ converges weakly to μ_1^* , the first component of the solution to the vector equilibrium problem given in Definition 2.1.*




Figure 3: The boundary $\partial\Omega$ of the domain Ω and the star Σ_1^* that supports the measure μ_1^* for the case $d = 3$ and $t_4 = 2$ at a subcritical time $t_0 = \frac{1}{20}$ (on the left) and at the critical time $t_0 = t_{0,\text{crit}} = \frac{1}{9}$ (on the right).

2.4 Laplacian growth and the spectral curve

Given the data t_0 , t_{d+1} and x^* as in Theorem 2.3, we let μ_1^* be the first component of the minimizer of the corresponding vector equilibrium problem.

Definition 2.5. We define $\xi_1 : \mathbb{C} \setminus \Sigma_1^* \rightarrow \mathbb{C}$ by

$$\xi_1(z) = t_{d+1} z^d + t_0 \int \frac{d\mu_1^*(\zeta)}{z - \zeta}. \quad (2.12)$$

Theorem 2.6. Let $t_{d+1} > 0$ and $0 < t_0 < t_{0,\text{crit}}$. Then there is a simply connected bounded domain Ω , whose boundary satisfies the equation

$$\xi_1(z) = \bar{z}, \quad z \in \partial\Omega. \quad (2.13)$$

The star Σ_1^* is contained in Ω . For an integer $k \geq 0$ we have

$$\frac{1}{2\pi i} \oint_{\partial\Omega} \frac{\bar{z}}{z^k} dz = \begin{cases} t_0, & \text{if } k = 0, \\ t_{d+1}, & \text{if } k = d + 1, \\ 0, & \text{otherwise,} \end{cases} \quad (2.14)$$

where $\partial\Omega$ is given the positive orientation. Moreover, μ_1^* and the normalized area measure on Ω are related by

$$\int \frac{d\mu_1^*(\zeta)}{z - \zeta} = \frac{1}{\pi t_0} \iint \frac{dA(\zeta)}{z - \zeta}, \quad z \in \mathbb{C} \setminus \Omega. \quad (2.15)$$

Theorem 2.6 generalizes the result obtained in [7, Theorem 2.6] in the case $d = 2$. Observe that (2.13) indicates that ξ_1 is the Schwarz function associated with the curve $\partial\Omega$. The property (2.14) with $k = 0$ means that

$$\text{area}(\Omega) = \pi t_0,$$

and so in terms of the parameter t_0 , the domain Ω evolves according to the equations of Laplacian growth, see Figure 3 for an illustration in the case $d = 3$.

Our final result is that ξ_1 satisfies an algebraic equation, the so-called spectral curve.

Theorem 2.7. *The Schwarz function (2.12) satisfies an algebraic equation of degree $d + 1$ with real coefficients of the form*

$$P(z, \xi) = \xi^{d+1} + z^{d+1} - \sum_{k=1}^d c_k z^k \xi^k + \beta = 0, \quad (2.16)$$

where $c_k > 0$ for all $k = 1, \dots, d$, and $\beta > 0$.

The coefficients c_d and c_{d-1} in (2.16) have the form

$$c_d = t_{d+1}, \quad (2.17)$$

$$c_{d-1} = \begin{cases} t_0 t_{d+1}, & \text{for } d \geq 3, \\ t_0 t_3 + \frac{1}{t_3}, & \text{for } d = 2. \end{cases} \quad (2.18)$$

The function $\xi_1(z)$ is the only solution of (2.16) that can be defined as an analytic function in the exterior of Σ_1^* . Because of (2.16), it has an analytic continuation to a $(d + 1)$ -sheeted Riemann surface. The analytic continuations are also important for the asymptotic analysis that follows.

2.5 Outline of the paper

The rest of the paper is organized as follows. In Section 3 we analyze the properties of the solution to the vector equilibrium problem and prove Theorems 2.2, 2.6 and 2.7. In Section 4 we describe the RH problem associated with the multiple orthogonal polynomials $P_{n,n}$. As a result of the steepest descent analysis of this RH problem, we will obtain the strong asymptotic behavior of the polynomials $P_{n,n}$ outside the star Σ_1^* , see Theorem 2.3. The steepest descent analysis is developed in Sections 5–10.

3 Proofs of Theorems 2.2, 2.6 and 2.7

In this section we analyze the vector equilibrium problem introduced in Definition 2.1, in the subcritical regime and thereby prove the Theorems 2.2, 2.6 and 2.7.

3.1 The Riemann surface in the subcritical case

Let $t_{d+1} > 0$ be fixed. Then it is easy to see that the function

$$r \mapsto r^2 - dt_{d+1}^2 r^{2d}$$

attains its maximum value on the positive real axis at the point $r_{\text{crit}} = (dt_{d+1})^{-\frac{1}{d-1}}$, and the maximum value is exactly given by $t_{0,\text{crit}}$ in (2.6). The function is increasing on $[0, r_{\text{crit}}]$. See also Figure 4. Thus for $t_0 \in (0, t_{0,\text{crit}}]$, there is a unique $r \in [0, r_{\text{crit}}]$ such that (2.8) holds. Then x^* is defined by (2.7) as in Theorem 2.2. From now on, we assume that $t_0 < t_{0,\text{crit}}$, r and x^* are fixed.




Figure 4: The graph of the function $r \mapsto r^2 - dt_{d+1}^2 r^{2d}$ for $r \geq 0$.

Our starting point is the compact Riemann surface \mathcal{R} of genus zero given by the collection of all pairs (z, ξ) of the form

$$\begin{aligned} z &= \psi(w) = rw + \frac{t_{d+1}r^d}{w^d}, & w &\in \overline{\mathbb{C}}. \\ \xi &= \psi\left(\frac{1}{w}\right) = \frac{r}{w} + t_{d+1}r^d w^d, \end{aligned} \quad (3.1)$$

Let us first examine the sheet structure of this Riemann surface.

The finite branch points of \mathcal{R} are precisely the endpoints $\omega^\ell x^*$, $\ell = 0, \dots, d$ of the star Σ_1^* . Indeed, the derivative $\psi'(w) = r - \frac{dt_{d+1}}{w^{d+1}}r^d$ has zeros $\omega^\ell w^*$, $\ell = 0, \dots, d$, where $w^* = (dt_{d+1}r^{d-1})^{\frac{1}{d+1}}$ and $\psi(\omega^\ell w^*) = \omega^\ell x^*$.

For any z , the equation

$$z = rw + \frac{t_{d+1}r^d}{w^d} \quad (3.2)$$

has $d+1$ solutions $w_k(z)$, $k = 1, \dots, d+1$, which we label such that

$$|w_1(z)| \geq |w_2(z)| \geq \dots \geq |w_{d+1}(z)| > 0. \quad (3.3)$$

The values $w_k(z)$ are unambiguously defined at points z where all inequalities in (3.3) are strict. If an equality occurs, we assign an arbitrary labeling such that (3.3) holds. It follows from [10, Thm. 2.2] (see also [2, Prop. 1]) that the star-like sets (2.9)–(2.10) are such that

$$\Sigma_k^* = \{z \in \mathbb{C} : |w_k(z)| = |w_{k+1}(z)|\}, \quad k = 1, \dots, d. \quad (3.4)$$

This implies that for each k , the solution $w_k(z)$ of (3.2) defines an analytic function on $\mathbb{C} \setminus (\Sigma_{k-1}^* \cup \Sigma_k^*)$ (with the understanding that $\Sigma_0^* = \Sigma_{d+1}^* = \emptyset$) and it is analytically continued by $w_{k+1}(z)$ through Σ_k^* .

We use the branches w_k to define the sheet structure on \mathcal{R} . Thus \mathcal{R} has sheets $\mathcal{R}_1, \dots, \mathcal{R}_{d+1}$ given by

$$\begin{aligned} \mathcal{R}_1 &= \overline{\mathbb{C}} \setminus \Sigma_1^*, \\ \mathcal{R}_k &= \overline{\mathbb{C}} \setminus (\Sigma_{k-1}^* \cup \Sigma_k^*), \quad 2 \leq k \leq d, \\ \mathcal{R}_{d+1} &= \overline{\mathbb{C}} \setminus \Sigma_d^*. \end{aligned} \quad (3.5)$$

For each $k = 1, \dots, d$, the sheets \mathcal{R}_k and \mathcal{R}_{k+1} are glued together through the cut Σ_k in the usual crosswise manner. See Figure 5 for a visualization of the Riemann surface in the case $d = 3$.




Figure 5: The Riemann surface \mathcal{R} in the case $d = 3$. The Riemann surface has $d + 1$ sheets and $z \mapsto \xi_k(z)$ is considered as an analytic function on the k th sheet.

Definition 3.1. We define, in accordance with (3.1), the functions

$$\xi_k(z) = \psi\left(\frac{1}{w_k(z)}\right) = \frac{r}{w_k(z)} + t_{d+1} r^d w_k(z)^d, \quad (3.6)$$

for $k = 1, \dots, d + 1$.

We consider $\xi_k(z)$ to be defined on the k th sheet of the Riemann surface, and then these functions are branches of a meromorphic function on \mathcal{R} .

It easily follows from (3.2) that for each $k = 1, \dots, d + 1$,

$$w_k(\bar{z}) = \overline{w_k(z)}, \quad (3.7)$$

$$w_k(\omega z) = \omega w_k(z). \quad (3.8)$$

We also deduce easily that as $z \rightarrow \infty$,

$$w_1(z) = \frac{z}{r} - \frac{t_{d+1} r^{2d-1}}{z^d} + O\left(\frac{1}{z^{2d+1}}\right), \quad (3.9)$$

$$w_k(z) = O\left(\frac{1}{z^{1/d}}\right), \quad \text{for } k = 2, \dots, d + 1. \quad (3.10)$$

The properties (3.7)–(3.10) and (3.6) imply that for $k = 1, \dots, d + 1$,

$$\xi_k(\bar{z}) = \overline{\xi_k(z)}, \quad (3.11)$$

$$\xi_k(\omega z) = \omega^{-1} \xi_k(z), \quad (3.12)$$

and as $z \rightarrow \infty$,

$$\xi_1(z) = t_{d+1} z^d + \frac{t_0}{z} + O\left(\frac{1}{z^{d+2}}\right), \quad (3.13)$$

$$\xi_k(z) = O(z^{1/d}), \quad \text{for } k = 2, \dots, d + 1, \quad (3.14)$$

where in (3.13) we used the relation $t_0 = r^2 - dt_{d+1}^2 r^{2d}$ in identifying the coefficient of $1/z$. We will now give a more detailed description of the asymptotic behavior (3.14).




Figure 6: The eight sectors S_ℓ^\pm in the case $d = 3$.

3.2 Asymptotics of functions ξ_k

In the complex plane we define the infinite sectors

$$S_\ell = \left\{ z \in \mathbb{C} : \frac{(2\ell - 1)\pi}{d+1} < \arg z < \frac{(2\ell + 1)\pi}{d+1} \right\}, \quad \ell = 0, \dots, d, \quad (3.15)$$

and we subdivide each sector S_ℓ into two parts

$$\begin{aligned} S_\ell^+ &= \left\{ z \in \mathbb{C} : \frac{2\ell\pi}{d+1} < \arg z < \frac{(2\ell + 1)\pi}{d+1} \right\}, \\ S_\ell^- &= \left\{ z \in \mathbb{C} : \frac{(2\ell - 1)\pi}{d+1} < \arg z < \frac{2\ell\pi}{d+1} \right\}. \end{aligned} \quad (3.16)$$

We are going to use the index ℓ with period $d+1$, so that $S_\ell = S_{\ell+d+1}$. In total we have $2(d+1)$ sectors S_ℓ^\pm . Frequently, and especially in Lemma 3.2 below, we will not number the sectors from S_0^- to S_d^+ but from $S_{-\frac{d}{2}}^-$ to $S_{\frac{d}{2}}^+$ in case d is even and from $S_{-\frac{d+1}{2}}^+$ to $S_{\frac{d+1}{2}}^-$ in case d is odd. See Figures 6 and 7 for the cases $d = 3$ and $d = 4$.

Thus, we have

$$\mathbb{C} \setminus (\Sigma_2 \cup \Sigma_3) = \bigcup_{\ell=0}^d (S_\ell^+ \cup S_\ell^-),$$

cf. (2.2) and (3.16).

In Lemma 3.2 and throughout the paper we use the principal branch of fractional exponents, that is, with a branch cut along the negative real axis.

Lemma 3.2. *For $k = 2, \dots, d+1$, we have*

$$\xi_k(z) = \kappa_{k,\ell}^\pm z^{1/d} - \frac{t_0}{d} z^{-1} + O\left(z^{-2-1/d}\right), \quad \text{as } z \rightarrow \infty \text{ in } S_\ell^\pm \quad (3.17)$$

where

$$\begin{aligned} \kappa_{k,\ell}^+ &= \omega_d^{-\ell + (-1)^k \lfloor \frac{k-1}{2} \rfloor} t_{d+1}^{-1/d}, & -\lceil d/2 \rceil \leq \ell \leq \lfloor d/2 \rfloor, \\ \kappa_{k,\ell}^- &= \omega_d^{-\ell - (-1)^k \lfloor \frac{k-1}{2} \rfloor} t_{d+1}^{-1/d}, & -\lceil d/2 \rceil \leq \ell \leq \lfloor d/2 \rfloor, \end{aligned} \quad (3.18)$$

and $\omega_d = \exp\left(\frac{2\pi i}{d}\right)$.




Figure 7: The ten sectors S_ℓ^\pm in the case $d = 4$.

Proof. Let $2 \leq k \leq d + 1$. Since the function $w_k(z)$ vanishes at infinity and satisfies

$$w_k(z)^{d+1} - \frac{z}{r} w_k(z)^d + t_{d+1} r^{d-1} = 0,$$

we deduce that as $z \rightarrow \infty$ in one of the sectors S_ℓ^\pm ,

$$w_k(z) = c t_{d+1}^{1/d} r z^{-1/d} + c^2 t_{d+1}^{2/d} r^3 d^{-1} z^{-1-2/d} + O(z^{-2-3/d}), \quad (3.19)$$

where c is a d -th root of unity (i.e., $c^d = 1$) that depends on k and on the sector. In order to respect the ordering (3.3) we have

$$\begin{aligned} w_k(z) &= \omega_d^{\ell - (-1)^k \lfloor \frac{k-1}{2} \rfloor} t_{d+1}^{1/d} r z^{-1/d} + O(z^{-1-2/d}) & \text{as } z \rightarrow \infty \text{ in } S_\ell^+, \\ w_k(z) &= \omega_d^{\ell + (-1)^k \lfloor \frac{k-1}{2} \rfloor} t_{d+1}^{1/d} r z^{-1/d} + O(z^{-1-2/d}) & \text{as } z \rightarrow \infty \text{ in } S_\ell^-. \end{aligned} \quad (3.20)$$

Then applying (3.6) and (3.19)-(3.20) we arrive at (3.17). \square

Note that the coefficient of z^{-1} in (3.17) does not vary with k and the sector.

3.3 Construction of the vector equilibrium measure in terms of the functions ξ_k

We use the functions ξ_k defined in (3.6) to explicitly solve the vector equilibrium problem. As already mentioned, the vector equilibrium problem belongs to the class of weakly admissible problems considered in [17]. Consequently, it has a unique minimizer $(\mu_1^*, \dots, \mu_d^*)$. The uniqueness readily implies that all the measures μ_k^* are invariant under the rotation $z \mapsto \omega z$.

Given a positive measure μ , let us denote by U^μ its logarithmic potential, that is,

$$U^\mu(z) = \int \log \frac{1}{|z-t|} d\mu(t), \quad z \in \mathbb{C}. \quad (3.21)$$

In what follows we describe some general properties of the vector equilibrium measure $(\mu_1^*, \dots, \mu_d^*)$ that are valid regardless of the values of t_0 , t_{d+1} and x^* . Later we will focus on the subcritical regime for these parameters.

Observe from (2.3) that the first measure μ_1^* minimizes the energy functional

$$\mu_1 \mapsto I(\mu_1) - I(\mu_1, \mu_2^*) + \frac{1}{t_0} \int_{\Sigma_1} \left(\frac{d}{(d+1)t_{d+1}^{1/d}} |z|^{\frac{d+1}{d}} - \frac{t_{d+1}}{d+1} z^{d+1} \right) d\mu_1(z)$$

among all probability measures on Σ_1 . This means that μ_1^* is the equilibrium measure in the presence of an external field, cf. [32], therefore μ_1^* is characterized by the variational conditions

$$-2U^{\mu_1^*}(z) + U^{\mu_2^*}(z) - \frac{1}{t_0} \left(\frac{d}{(d+1)t_{d+1}^{1/d}} |z|^{\frac{d+1}{d}} - \frac{t_{d+1}}{d+1} z^{d+1} \right) \begin{cases} = \ell_1, & z \in \text{supp}(\mu_1^*), \\ \leq \ell_1, & z \in \Sigma_1 \setminus \text{supp}(\mu_1^*), \end{cases} \quad (3.22)$$

for a certain constant ℓ_1 .

Similarly, for a fixed k with $2 \leq k \leq d-1$, the measure μ_k^* minimizes the energy functional

$$\mu_k \mapsto I(\mu_k) - I(\mu_{k-1}^*, \mu_k) - I(\mu_k, \mu_{k+1}^*),$$

among all positive measures μ_k satisfying (2.4)–(2.5). Consequently, μ_k^* is the balayage of $(\mu_{k-1}^* + \mu_{k+1}^*)/2$ onto Σ_k . (See e.g. [32] for the notion of balayage of a measure in logarithmic potential theory.) This in turn implies that $\text{supp}(\mu_k^*) = \Sigma_k$ and μ_k^* is characterized by the condition

$$2U^{\mu_k^*}(z) = U^{\mu_{k-1}^*}(z) + U^{\mu_{k+1}^*}(z), \quad z \in \Sigma_k. \quad (3.23)$$

Analogously, we obtain that μ_d^* is the balayage of $\mu_{d-1}^*/2$ onto Σ_d , and so $\text{supp}(\mu_d^*) = \Sigma_d$ and μ_d^* is characterized by the condition

$$2U^{\mu_d^*}(z) = U^{\mu_{d-1}^*}(z), \quad z \in \Sigma_d. \quad (3.24)$$

In what follows we choose an orientation on the stars Σ_1^* and Σ_k , $k = 2, \dots, d$ that on each segment is pointing away from the origin. It induces a $+$ and $-$ side on these contours, where the $+$ side lies on the left and the $-$ side on the right as one moves away from the origin. For an oriented contour Σ and a function f defined on a neighborhood of Σ in the complex plane, we use f_{\pm} to denote the boundary values on Σ as one approaches Σ from the \pm side.

Proposition 3.3. *Let $t_{d+1} > 0$, $t_0 \in (0, t_{0,\text{crit}})$ and x^* be as in Theorem 2.2 and let $(\mu_1^*, \dots, \mu_d^*)$ be the minimizer of the vector equilibrium problem of Definition 2.1 with $\hat{x} = x^*$. Then we have the following:*

(a) *The measure μ_1^* is given by*

$$d\mu_1^*(z) = \frac{1}{2\pi i t_0} (\xi_{1,-}(z) - \xi_{1,+}(z)) dz, \quad z \in \Sigma_1^*, \quad (3.25)$$

where dz is the complex line element on Σ_1^ ,*

(b) *For $k \geq 2$, we have that μ_k^* is given by*

$$d\mu_k^*(z) = \frac{1}{2\pi i t_0} (\eta_{k,-}(z) - \eta_{k,+}(z)) dz, \quad z \in \Sigma_k, \quad (3.26)$$

where

$$\eta_k(z) = \xi_k(z) - \kappa_{k,\ell}^{\pm} z^{1/d}, \quad z \in S_{\ell}^{\pm}, \quad (3.27)$$

and the numbers $\kappa_{k,\ell}^{\pm}$ are as in Lemma 3.2.

Note that it follows from (3.25) and (3.13) that

$$\xi_1(z) = t_{d+1}z^d + t_0 \int \frac{d\mu_1^*(s)}{z-s}$$

by Plemelj's formula for the boundary values of a Cauchy transform. So the definition of ξ_1 in (3.6) is consistent with the earlier Definition 2.5.

To prove Proposition 3.3 we assume that μ_k^* for $k = 1, \dots, d$ are given by the right-hand sides of (3.25)–(3.26) and we verify that they are real and positive measures with total masses

$$\int_{\Sigma_k} d\mu_k^*(z) = 1 - \frac{k-1}{d}, \quad k = 1, \dots, d, \quad (3.28)$$

satisfying the variational conditions (3.22), (3.23) and (3.24). Note that due to (3.17) and (3.27) we have

$$\eta_{k,-}(z) - \eta_{k,+}(z) = O\left(z^{-2-1/d}\right) \text{ as } z \rightarrow \infty$$

so that the right-hand side of (3.26) is a finite, but a priori complex, measure on Σ_k .

The proof of Proposition 3.3 is somewhat lengthy and it is subdivided into a number of steps.

3.4 Proof of Proposition 3.3

3.4.1 Total masses

We start by establishing (3.28).

Lemma 3.4. *The (a priori complex) measures μ_k^* defined in (3.25)–(3.26) are real-valued, rotationally invariant, and have the total masses (3.28).*

Proof. The property (3.11) implies that $\xi_{1,-}(x) = \overline{\xi_{1,+}(x)}$ for every $x \in (0, x^*)$, hence μ_1^* is real-valued on $[0, x^*]$. But (3.12) implies immediately that μ_1^* is rotationally invariant, so this measure is real-valued everywhere.

Applying (3.11)–(3.12) for $z \in S_0^-$ and using (3.17), we deduce that $\kappa_{k,0}^+ = \overline{\kappa_{k,0}^-}$ and $\kappa_{k,1}^- = \omega_d^{-1} \kappa_{k,0}^-$, hence $\kappa_{k,1}^- = \omega_d^{-1} \overline{\kappa_{k,0}^+}$. We also deduce from (3.11)–(3.12) that for $z \in \Sigma_2$ with $\arg z = \frac{\pi}{d+1}$, we have $\xi_{k,+}(z) = \omega_{d+1}^{-1} \overline{\xi_{k,-}(z)}$. With this information we conclude that for k odd, $k \geq 3$, the measure μ_k^* is real-valued on $[0, \infty)$, and for k even, $k \geq 2$, the measure μ_k^* is real-valued on $e^{\frac{\pi i}{d+1}} [0, \infty)$. The rotational invariance is then a consequence of (3.12).

We now prove (3.28). Firstly, since ξ_1 is analytic in $\mathbb{C} \setminus \Sigma_1^*$, using (3.25), (3.13) and Cauchy's theorem, we obtain

$$\int_{\Sigma_1^*} d\mu_1^*(z) = \frac{1}{2\pi i t_0} \int_{\Sigma_1^*} (\xi_{1,-}(z) - \xi_{1,+}(z)) dz = \frac{1}{2\pi i t_0} \oint_{\gamma_1} \xi_1(z) dz = 1, \quad (3.29)$$

where γ_1 is any positively oriented closed curve surrounding Σ_1^* , as in Figure 8. This proves (3.28) for $k = 1$.

In the case $k = 2$, we let γ_2 be a system of unbounded contours around $\Sigma_1^* \cup \Sigma_2$ as shown in the right part of Figure 8, with orientation as also indicated in Figure 8.

Noting that $\eta_2(z) = -\frac{t_0}{d}z^{-1} + O(z^{-2-1/d})$ as $z \rightarrow \infty$, see (3.17) and (3.27), we find by deforming the contour to infinity in $\mathbb{C} \setminus (\Sigma_1^* \cup \Sigma_2)$ that

$$\int_{\gamma_2} \eta_2(z) dz = -2\pi i \frac{t_0}{d}. \quad (3.30)$$




Figure 8: The contour γ_1 around Σ_1^* (on the left) and the system of unbounded contours γ_2 around $\Sigma_1^* \cup \Sigma_2$ (on the right).

On the other hand, we have by deforming γ_2 towards $\Sigma_1^* \cup \Sigma_2$ that

$$\int_{\gamma_2} \eta_2(z) dz = \int_{\Sigma_2} (\eta_{2,-}(z) - \eta_{2,+}(z)) dz + \int_{\Sigma_1^*} (\eta_{2,-}(z) - \eta_{2,+}(z)) dz. \quad (3.31)$$

On Σ_1^* we have

$$\eta_{2,-}(z) - \eta_{2,+}(z) = \xi_{2,-}(z) - \xi_{2,+}(z), \quad z \in \Sigma_1^*$$

which follows from (3.17) and (3.27) since $\kappa_{2,\ell}^+ = \kappa_{2,\ell}^-$, see (3.18). On Σ_1^* we also have $\xi_{2,\mp} = \xi_{1,\pm}$, so that by (3.26), (3.30) and (3.31),

$$\begin{aligned} 2\pi i t_0 \int d\mu_2^*(z) &= \int_{\Sigma_2} (\eta_{2,-}(z) - \eta_{2,+}(z)) dz \\ &= -2\pi i \frac{t_0}{d} - \int_{\Sigma_1^*} (\xi_{1,+}(z) - \xi_{1,-}(z)) dz \\ &= -2\pi i \frac{t_0}{d} + 2\pi i t_0 \end{aligned} \quad (3.32)$$

where in the last step we used (3.29). This proves (3.28) for $k = 2$.

The case $k \geq 3$ follows from a similar argument based on induction and the use of a contour γ_k around $\Sigma_{k-1} \cup \Sigma_k$. Here it is important that

$$\eta_{k,\pm}(z) = \eta_{k-1,\mp}(z), \quad z \in \Sigma_{k-1},$$

which holds since $\xi_{k-1,\mp}$ is the analytic continuation of $\xi_{k,\pm}$ across Σ_{k-1} , and therefore the leading coefficients in their expansions (3.17) also agree if we move via one of the half-rays in Σ_{k-1} from one sector to the next. Thus by (3.27) also η_{k-1} is the analytic continuation of η_k across Σ_{k-1} .

The fact that the leading coefficients agree comes down to the relations $\kappa_{k,\ell}^\pm = \kappa_{k-1,\ell}^\mp$ if k is even, and $\kappa_{k,\ell}^\pm = \kappa_{k-1,\ell\pm 1}^\mp$ if k is odd, which can also be verified directly from (3.18). \square

3.4.2 Positivity of μ_1^*

The next step is to show that μ_1^* is a positive measure. The proof that the other measures are positive as well will come later. We need a lemma about the function $\psi(w) = rw + t_{d+1} r^d w^{-d}$ introduced in (3.1) that is important later as well.

Lemma 3.5. (a) *The function ψ from (3.1) is one-to one in the domain $\{w \in \mathbb{C} \mid |w| \geq \rho\}$ where*

$$\rho := (dt_{d+1}r^{d-1})^{1/(d+1)}. \quad (3.33)$$

(b) *We have*

$$\rho \leq 1 \quad (3.34)$$

with equality if and only if $t_0 = t_{0,\text{crit}}$.

(c) *ψ satisfies*

$$\begin{aligned} \operatorname{Im} w > 0 &\implies \operatorname{Im} \psi(w) > 0, \\ \operatorname{Im} w < 0 &\implies \operatorname{Im} \psi(w) < 0, \end{aligned} \quad \text{for } |w| \geq \rho. \quad (3.35)$$

(d) *For $z = \psi(w)$ with $|w| \geq \rho$, we have*

$$\xi_1(z) = \psi\left(\frac{1}{\psi^{-1}(z)}\right). \quad (3.36)$$

(e) *We have*

$$\Sigma_1^* \subset \mathbb{C} \setminus \psi(\{w \in \mathbb{C} \mid |w| > \rho\}).$$

Proof. (a) Let us put $a = t_{d+1}r^d$. Assume to get a contradiction that $\psi(w_1) = \psi(w_2)$ with $w_1 \neq w_2$, $|w_1| \geq |w_2| \geq \rho$ where ρ is given by (3.33). Thus

$$r(w_1 - w_2) = a \left(\frac{1}{w_2^d} - \frac{1}{w_1^d} \right) = a \frac{w_1^d - w_2^d}{w_1^d w_2^d},$$

which since $w_1 \neq w_2$ leads to

$$r w_1^d w_2^d = a(w_1^{d-1} + w_1^{d-2}w_2 + \cdots + w_2^{d-1}). \quad (3.37)$$

Using $a/r = \frac{1}{d}\rho^{d+1}$, and setting $\lambda_j = w_j/\rho$, for $j = 1, 2$, we can rewrite this further as

$$\lambda_1^d \lambda_2^d = \frac{1}{d}(\lambda_1^{d-1} + \lambda_1^{d-2}\lambda_2 + \cdots + \lambda_2^{d-1}) \quad (3.38)$$

with $|\lambda_1| \geq |\lambda_2| \geq 1$ and $\lambda_1 \neq \lambda_2$. Taking absolute values on both sides of (3.38), we obtain

$$\begin{aligned} |\lambda_1|^d &\leq |\lambda_1^d \lambda_2^d| = \frac{1}{d} |\lambda_1^{d-1} + \lambda_1^{d-2}\lambda_2 + \cdots + \lambda_2^{d-1}| \\ &\leq \frac{1}{d} (|\lambda_1|^{d-1} + |\lambda_1|^{d-2}|\lambda_2| + \cdots + |\lambda_2|^{d-1}) \leq |\lambda_1|^{d-1}, \end{aligned} \quad (3.39)$$

and from this it follows that $|\lambda_1| = |\lambda_2| = 1$.

It also follows that equality should hold in every inequality in (3.39) and in particular

$$|\lambda_1^{d-1} + \lambda_1^{d-2}\lambda_2 + \cdots + \lambda_2^{d-1}| = |\lambda_1|^{d-1} + |\lambda_1|^{d-2}|\lambda_2| + \cdots + |\lambda_2|^{d-1}.$$

This implies that the complex numbers $\lambda_1^{d-1-j}\lambda_2^j$, $j = 0, \dots, d-1$ have the same argument, and it is easily seen that this is impossible if $|\lambda_1| = |\lambda_2|$ with $\lambda_1 \neq \lambda_2$. This contradiction proves that ψ is one-to-one on the closed set $|w| \geq \rho$.

(b) For part (b) we note that $r \leq r_{\text{crit}} = (dt_{d+1})^{-1/(d-1)}$, as is easily seen from Figure 4. In view of (3.33) we get the inequality (3.34). Equality holds if and only if $r = r_{\text{crit}}$ and this happens if and only if $t_0 = t_{0,\text{crit}}$.

(c) Let $|w| \geq \rho$ and $w \in \mathbb{C} \setminus \mathbb{R}$. Since $\psi(\overline{w}) = \overline{\psi(w)}$ and ψ is one-to-one, it follows that $\psi(w) \in \mathbb{C} \setminus \mathbb{R}$, and then part (c) follows.

(d) Let $z = \psi(w_0)$ with $|w_0| \geq \rho$. Since ψ is one-to-one for $|w| \geq \rho$, it is then clear that w_0 is the largest in absolute value solution of (3.2). Thus $w_0 = w_1(z)$ as in (3.3) and part (d) follows because of the definition (3.6).

(e) If $z = \psi(w_0)$ with $|w_0| > \rho$, then $w_0 = w_1(z)$ as in the proof of part (d), and since ψ is one-to-one for $|w| > \rho$, we have $|w_2(z)| \leq \rho$. Then $z \notin \Sigma_1^*$ because of the characterization (3.4) of Σ_1^* . \square

It follows in particular that for $t_0 < t_{0,\text{crit}}$ the curve $\psi(|w| = 1)$ is a closed analytic curve that surrounds a domain containing Σ_1^* .

Lemma 3.6. *The measure μ_1^* is positive with $\text{supp}(\mu_1^*) = \Sigma_1^*$.*

Proof. By the rotational invariance of μ_1^* , see Lemma 3.4, it is enough to show that for $0 < x < x^*$,

$$\frac{d\mu_1^*(x)}{dx} = \frac{\xi_{1,-}(x) - \xi_{1,+}(x)}{2\pi i t_0} = \frac{\text{Im } \xi_{1,-}(x)}{\pi t_0} > 0, \quad (3.40)$$

see (3.25). By (3.6) we have

$$\xi_{1,-}(x) = \psi\left(\frac{1}{w_{1,-}(x)}\right), \quad (3.41)$$

where $w_{1,-}(x)$ is a solution of $\psi(w) = x$ of highest absolute value. We have that $\text{Im } w_{1,-}(x) \neq 0$.

Since $x \in \Sigma_1^*$ we find by Lemma 3.5 (e) that $|w_{1,-}(x)| \leq \rho$. Thus $\frac{1}{|w_{1,-}(x)|} \geq \frac{1}{\rho} \geq \rho$ (since $\rho \leq 1$, see Lemma 3.5 (b)). Applying Lemma 3.5 (c) and (3.41), we find $\text{Im } \xi_{1,-}(x) \neq 0$. Thus by (3.40) the density of μ_1^* does not vanish on $(0, x^*)$. By Lemma 3.4 and the rotational symmetry we already know that

$$\int_0^{x^*} d\mu_1^*(x) = \frac{1}{d+1} > 0.$$

The density is real and continuous, and since it does not vanish we conclude it is positive. This proves (3.40).

It is clear that $\text{supp}(\mu_1^*) = \Sigma_1^*$. \square

3.4.3 Identities for Cauchy transforms

The next steps depend on the properties of the Cauchy transforms of the measures μ_k^* that we define as follows.

Definition 3.7. For $k = 1, \dots, d$, we define F_k as

$$F_k(z) := \int \frac{d\mu_k^*(t)}{z-t}, \quad z \in \mathbb{C} \setminus \Sigma_k, \quad k = 1, \dots, d. \quad (3.42)$$

These functions are closely related to the functions ξ_k as shown in the following lemma.

Lemma 3.8. *The following relations hold between the functions ξ_k defined in (3.6) and the functions F_k defined in (3.42):*

$$\xi_1(z) = t_{d+1} z^d + t_0 F_1(z), \quad z \in \mathbb{C} \setminus \Sigma_1^*, \quad (3.43)$$

and for $k = 2, \dots, d+1$,

$$\xi_k(z) = t_0(F_k(z) - F_{k-1}(z)) + \kappa_{k,\ell}^\pm z^{1/d}, \quad z \in S_\ell^\pm, \quad (3.44)$$

where $F_{d+1} \equiv 0$, and the coefficients $\kappa_{k,\ell}^\pm$ are given in (3.18).

Proof. With the contour γ_1 introduced in the proof of Lemma 3.4, see also Figure 8, we have by (3.42) and (3.25) for any $z \in \mathbb{C} \setminus \Sigma_1^*$ in the exterior of γ_1 ,

$$t_0 F_1(z) = \frac{1}{2\pi i} \int_{\Sigma_1^*} \frac{\xi_{1,-}(s) - \xi_{1,+}(s)}{z-s} ds = \frac{1}{2\pi i} \int_{\gamma_1} \frac{\xi_1(s)}{z-s} ds.$$

Moving the contour to infinity we pick up a residue contribution at $s = z$, which is $\xi_1(z)$, and a contribution at infinity, which because of (3.13) is equal to $-t_{d+1}z^d$. This proves (3.43).

To prove (3.44) for $k \geq 2$ we use a contour γ_2 as in Figure 8. Then for $z \in \mathbb{C} \setminus (\Sigma_1^* \cup \Sigma_2)$, we find (where we use η_2 as in (3.27))

$$2\pi i t_0 (F_2(z) - F_1(z)) = \int_{\Sigma_2} \frac{\eta_{2,-}(s) - \eta_{2,+}(s)}{z-s} ds - \int_{\Sigma_1^*} \frac{\xi_{1,-}(s) - \xi_{1,+}(s)}{z-s} ds.$$

We have $-\xi_{1,-} + \xi_{1,+} = -\xi_{2,+} + \xi_{2,-} = -\eta_{2,+} + \eta_{2,-}$ on Σ_1^* , which implies that

$$2\pi i t_0 (F_2(z) - F_1(z)) = \int_{\gamma_2} \frac{\eta_2(s)}{z-s} ds,$$

provided that z is in the exterior of γ_2 . Moving γ_2 to infinity we pick up a residue contribution $\eta_2(z)$ at $s = z$, and now there is no contribution at infinity, since $\eta_2(s) = O(s^{-1})$ as $s \rightarrow \infty$. This proves (3.44).

The proof for general k is similar. \square

3.4.4 The variational equalities

Lemma 3.9. *The identities (3.22), (3.23) and (3.24) hold where the measures μ_k^* are introduced in (3.25) and (3.26).*

Proof. Using the identities (3.43)–(3.44) and the fact that $\xi_{k,\pm} = \xi_{k+1,\mp}(z)$ on Σ_k , and taking note of the constants (3.18), we easily obtain the following for the Cauchy transforms F_k ,

$$F_{1,+}(z) + F_{1,-}(z) = F_2(z) + \frac{\omega^{\ell d}}{t_0} \left(\frac{1}{t_{d+1}^{1/d}} |z|^{1/d} - t_{d+1} |z|^d \right), \quad z \in (0, \omega^\ell x^*] \subset \Sigma_1, \quad (3.45)$$

$$F_{k,+}(z) + F_{k,-}(z) = F_{k-1}(z) + F_{k+1}(z), \quad z \in \Sigma_k, \quad k = 2, \dots, d-1, \quad (3.46)$$

$$F_{d,+}(z) + F_{d,-}(z) = F_{d-1}(z), \quad z \in \Sigma_d. \quad (3.47)$$

For instance, using (3.43), (3.44) for $k = 2$, and $\kappa_{2,0}^\pm = t_{d+1}^{-1/d}$, the relation $\xi_{1,+} = \xi_{2,-}$ on $(0, x^*)$ implies

$$t_{d+1} x^d + t_0 F_{1,+}(x) = t_0 (F_2(x) - F_{1,-}(x)) + \frac{x^{1/d}}{t_{d+1}^{1/d}}, \quad x \in (0, x^*),$$

which gives (3.45) for $\ell = 0$. Then the symmetry property $F_k(\omega z) = \omega^d F_k(z)$, $z \in \mathbb{C} \setminus \Sigma_k$, implies (3.45) for every ℓ .

From (3.21) and (3.42) we get $F_k = -\left(\frac{\partial}{\partial x} - i\frac{\partial}{\partial y}\right) U^{\mu_k^*}$. Then in view of the symmetry $F_k(\bar{z}) = \overline{F_k(z)}$, we obtain for any $z \in \mathbb{C} \setminus \Sigma_k$,

$$-2\frac{\partial}{\partial x} U^{\mu_k^*}(z) = F_k(z) + F_k(\bar{z}), \quad -2i\frac{\partial}{\partial y} U^{\mu_k^*}(z) = F_k(\bar{z}) - F_k(z). \quad (3.48)$$

This shows that for any $x \in (0, x^*)$,

$$-2 \frac{d}{dx} U^{\mu_1^*}(x) = F_{1,+}(x) + F_{1,-}(x), \quad -\frac{d}{dx} U^{\mu_2^*}(x) = F_2(x),$$

hence from (3.45) we obtain

$$-2 \frac{d}{dx} U^{\mu_1^*}(x) + \frac{d}{dx} U^{\mu_2^*}(x) = \frac{1}{t_0} \left(\frac{1}{t_{d+1}^{1/d}} x^{1/d} - t_{d+1} x^d \right), \quad x \in [0, x^*].$$

Integrating this relation we get

$$-2U^{\mu_1^*}(x) + U^{\mu_2^*}(x) - \frac{1}{t_0} \left(\frac{d}{(d+1)t_{d+1}^{1/d}} x^{\frac{d+1}{d}} - \frac{t_{d+1}}{d+1} x^{d+1} \right) = \ell_1, \quad x \in [0, x^*],$$

for some constant ℓ_1 . Using the rotational invariance of the measures μ_k^* we deduce that (3.22) holds with equality everywhere on Σ_1^* .

Similarly, using (3.46) and the first relation in (3.48), we argue that for k odd, (3.23) holds for $z \in \mathbb{R}_+$ and hence by rotational symmetry it also holds for every $z \in \Sigma_k$. Here the constant of integration vanishes, as can be seen by inspecting the behavior as $z \rightarrow \infty$. The same proof is valid for (3.24) if d is odd. The other cases follow in a similar way. \square

3.4.5 Positivity of measures μ_k^* with $k \geq 2$

The only piece of information that is still missing is that the measures μ_k^* are positive for $k \geq 2$. Recall that the positivity of μ_1^* was established in Lemma 3.6.

Lemma 3.10. *The measures μ_k^* are positive for $k \geq 2$ with $\text{supp}(\mu_k^*) = \Sigma_k^*$.*

Proof. Given the positive measure μ_1^* (3.25) on Σ_1 , we consider the auxiliary vector equilibrium problem consisting of minimizing the energy functional

$$\sum_{k=2}^d I(\mu_k) - I(\mu_1^*, \mu_2) - \sum_{k=2}^{d-1} I(\mu_k, \mu_{k+1})$$

among all positive Borel measures μ_2, \dots, μ_d satisfying $\text{supp}(\mu_k) \subset \Sigma_k$ and $\|\mu_k\| = 1 - \frac{k-1}{d}$ for every $k = 2, \dots, d$.

Let us denote by $(\hat{\mu}_2, \dots, \hat{\mu}_d)$ the minimizer to this new problem. Since μ_1^* is positive, we deduce that $\hat{\mu}_2$ is the balayage of $(\mu_1^* + \hat{\mu}_3)/2$ onto Σ_2 , therefore we have $\text{supp}(\hat{\mu}_2) = \Sigma_2$ and

$$2U^{\hat{\mu}_2}(z) = U^{\mu_1^*}(z) + U^{\hat{\mu}_3}(z), \quad z \in \Sigma_2. \quad (3.49)$$

Similarly we deduce that for each $k = 3, \dots, d$, the measure $\hat{\mu}_k$ is the balayage of $(\hat{\mu}_{k-1} + \hat{\mu}_{k+1})/2$ onto Σ_k , and therefore $\text{supp}(\hat{\mu}_k) = \Sigma_k$ and

$$2U^{\hat{\mu}_k}(z) = U^{\hat{\mu}_{k-1}}(z) + U^{\hat{\mu}_{k+1}}(z), \quad z \in \Sigma_k, \quad (3.50)$$

where $U^{\hat{\mu}_{d+1}} \equiv 0$.

Using (3.23) for $k = 2$ and (3.49), we obtain

$$2(U^{\hat{\mu}_2}(z) - U^{\mu_2^*}(z)) = U^{\hat{\mu}_3}(z) - U^{\mu_3^*}(z), \quad z \in \Sigma_2. \quad (3.51)$$

Analogously, from (3.23)–(3.24) and (3.50) we obtain for each $k = 3, \dots, d-1$,

$$2(U^{\widehat{\mu}_k}(z) - U^{\mu_k^*}(z)) = U^{\widehat{\mu}_{k-1}}(z) - U^{\mu_{k-1}^*}(z) + U^{\widehat{\mu}_{k+1}}(z) - U^{\mu_{k+1}^*}(z), \quad z \in \Sigma_k, \quad (3.52)$$

and

$$2(U^{\widehat{\mu}_d}(z) - U^{\mu_d^*}(z)) = U^{\widehat{\mu}_{d-1}}(z) - U^{\mu_{d-1}^*}(z), \quad z \in \Sigma_d. \quad (3.53)$$

Let us now define the constants

$$m_k := \inf_{z \in \mathbb{C}} \left(U^{\widehat{\mu}_k}(z) - U^{\mu_k^*}(z) \right), \quad M_k := \sup_{z \in \mathbb{C}} \left(U^{\widehat{\mu}_k}(z) - U^{\mu_k^*}(z) \right), \quad k = 2, \dots, d. \quad (3.54)$$

Since $U^{\widehat{\mu}_k} - U^{\mu_k^*}$ is continuous and vanishes at infinity, we have $-\infty < m_k \leq 0 \leq M_k < \infty$ for every $k = 2, \dots, d$.

Note that $U^{\widehat{\mu}_k} - U^{\mu_k^*}$ is harmonic in $\mathbb{C} \setminus \Sigma_k$, which means by the maximum and minimum principles for harmonic functions, that the maximum and minimum are taken on Σ_k . Then (3.51)–(3.53) lead to the inequalities between the numbers M_k , namely

$$2M_2 \leq M_3, \quad (3.55)$$

$$2M_k \leq M_{k-1} + M_{k+1}, \quad \text{for } k = 3, \dots, d-1, \quad (3.56)$$

$$2M_d \leq M_{d-1}. \quad (3.57)$$

Inductively, we obtain from (3.55) and (3.56) that

$$kM_k \leq (k-1)M_{k+1} \quad \text{for } k = 2, \dots, d-1. \quad (3.58)$$

Taking $k = d-1$ in (3.58) and combining this with (3.57) we conclude that $M_d = M_{d-1} = 0$. Then $M_k = 0$ for every k by (3.58).

In a similar way we prove that $m_k = 0$ for every k , and therefore $U^{\mu_k^*} \equiv U^{\widehat{\mu}_k}$. This implies that $\mu_k^* = \widehat{\mu}_k$ for every $k = 2, \dots, d$ by the uniqueness theorem for logarithmic potentials, see e.g. [32, Theorem II.2.1]. In particular μ_k^* is a positive measure. \square

3.5 Conclusion of the proof of Proposition 3.3

We proved in Lemmas 3.6 and 3.10 that the measures are real and positive. The total masses (3.28) are established in Lemma 3.4, and the variational conditions (3.22)–(3.24) are satisfied by Lemma 3.9.

3.6 Proof of Theorem 2.2

Recall from (3.1) and the paragraph that follows it, that $\psi'(w^*) = 0$ and $x^* = \psi(w^*)$. Then as $w \rightarrow w^*$,

$$z = \psi(w) = x^* + c_1(w - w^*)^2 + O(w - w^*)^3 \quad (3.59)$$

where $c_1 = \frac{1}{2}\psi''(w^*) > 0$. Also from (3.1) we have

$$\xi = \psi\left(\frac{1}{w}\right) = \xi_1(x^*) - c_2(w - w^*) + O(w - w^*)^2 \quad (3.60)$$

with $c_2 = \frac{1}{(w^*)^2}\psi'\left(\frac{1}{w^*}\right) > 0$, since $t_0 < t_{0,\text{crit}}$ (which implies that $r < r_{\text{crit}} = (dt_{d+1})^{-1/(d-1)}$, see Figure 4). Inverting (3.59) for z on the first sheet, we get

$$w - w^* = \frac{1}{\sqrt{c_1}}(z - x^*)^{1/2} + O(z - x^*) \quad \text{as } z \rightarrow x^*, \quad (3.61)$$

with the principal branch of the square root since $w > w^*$ for real $z > x^*$. Thus by (3.61) and (3.60)

$$\xi_1(z) = \xi_1(x^*) - c_3(z - x^*)^{1/2} + O(z - x^*) \quad (3.62)$$

as $z \rightarrow x^*$ with $c_3 = \frac{c_2}{\sqrt{c_1}} > 0$.

Then by (3.25) we have for $z \in (0, x^*)$,

$$\frac{d\mu_1^*(z)}{dz} = \frac{1}{\pi t_0} \text{Im} \xi_{1,-}(z) = c_4 \sqrt{x^* - z} \quad \text{as } z \nearrow x^*.$$

where $c_4 = \frac{c_3}{\pi t_0} > 0$. Thus the density of μ_1^* vanishes as a square root at the endpoint x^* . By rotational symmetry the density vanishes as a square root at each of the endpoints.

Similarly to (3.62) we also have

$$\xi_2(z) = \xi_1(x^*) + c_3(z - x^*)^{1/2} + O(z - x^*), \quad (3.63)$$

as $z \rightarrow x^*$. Then

$$\xi_1(z) - \xi_2(z) = -2c_3(z - x^*)^{1/2} + O(z - x^*). \quad (3.64)$$

Now it is easy to calculate that for $x > x^*$,

$$\begin{aligned} \frac{d}{dx} \left(-2U^{\mu_1^*}(x) + U^{\mu_2^*}(x) - \frac{1}{t_0} \left(\frac{d}{(d+1)t_{d+1}^{1/d}} x^{\frac{d+1}{d}} - \frac{t_{d+1}}{d+1} x^{d+1} \right) \right) \\ = 2F_1(x) - F_2(x) - \frac{1}{t_0} \left(\frac{1}{t_{d+1}^{1/d}} x^{1/d} - t_{d+1} x^d \right) \\ = \frac{1}{t_0} (\xi_1(x) - \xi_2(x)), \end{aligned} \quad (3.65)$$

where in the last step we used (3.43), (3.44) and (3.18). By (3.64) there is $x^{**} > x^*$ such that $\xi_1(z) - \xi_2(z) < 0$ for $x \in (x^*, x^{**}]$. Then the variational inequality, see (3.22),

$$-2U^{\mu_1^*}(x) + U^{\mu_2^*}(x) - \frac{1}{t_0} \left(\frac{d}{(d+1)t_{d+1}^{1/d}} x^{\frac{d+1}{d}} - \frac{t_{d+1}}{d+1} x^{d+1} \right) < \ell_1, \quad x \in (x^*, x^{**}],$$

follows from (3.65) and the fact that we have equality at $x = x^*$.

It now follows that for any choice of $\hat{x} \in [x^*, x^{**}]$ the measures $(\mu_1^*, \mu_2^*, \dots, \mu_d^*)$ are the minimizers for the vector equilibrium problem. Theorem 2.2 is now fully proved.

3.7 Proof of Theorem 2.6

By Lemma 3.5 we have that $\psi(|w| = 1)$ is a simple closed curve containing Σ_1^* in its interior if $t_0 < t_{0,\text{crit}}$.

We define Ω as the domain enclosed by the curve $\psi(|w| = 1)$. By parts (b) and (e) of Lemma 3.5 we then have that Σ_1^* is contained in Ω if $t_0 < t_{0,\text{crit}}$. If $z \in \partial\Omega$ then $z = \psi(w)$ with $|w| = 1$. By part (d) of Lemma 3.5, we have $\xi_1(z) = \psi(\frac{1}{w})$. Using $|w| = 1$ and the fact that the coefficients of ψ are real, we then find $\xi_1(z) = \bar{z}$ for $z \in \partial\Omega$.

Using (2.13), we find for an integer $k \geq 0$,

$$\frac{1}{2\pi i} \oint_{\partial\Omega} \frac{\bar{z}}{z^k} dz = \frac{1}{2\pi i} \oint_{\partial\Omega} \frac{\xi_1(z)}{z^k} dz.$$

Since $\xi_1(z)$ is analytic in the exterior of $\partial\Omega$, by deforming this contour and applying (3.13) we obtain immediately (2.14). Note that (2.14) with $k = 0$ implies by Green's theorem that the domain Ω has area πt_0 .

For $z \in \mathbb{C} \setminus \Omega$, we have by Green's theorem and (2.13)

$$\frac{1}{\pi} \iint_{\Omega} \frac{dA(\zeta)}{z - \zeta} = \frac{1}{2\pi i} \oint_{\partial\Omega} \frac{\bar{\zeta}}{z - \zeta} d\zeta = \frac{1}{2\pi i} \oint_{\partial\Omega} \frac{\xi_1(\zeta)}{z - \zeta} d\zeta.$$

Using (3.13) again and Cauchy's integral formula, the last integral is easily seen to be equal to $\xi_1(z) - t_{d+1}z^d$, and this is $t_0 F_1(z)$ by (3.43). This proves (2.15).

3.8 Proof of Theorem 2.7

We write

$$P(z, \xi) = \prod_{k=1}^{d+1} (\xi - \xi_k(z)), \quad (3.66)$$

where as before we consider $\xi_k(z)$ as being defined on the k th sheet of the Riemann surface. The equation $P(z, \xi) = 0$ is an equation of the Riemann surface with rational parametrization (3.1).

By changing $w \mapsto 1/w$ in the parametrization (3.1) we see that $P(z, \xi) = 0 \implies P(\xi, z) = 0$. This means that P is symmetric in z and ξ , and since P is of degree $d+1$ in ξ , we find

$$P(z, \xi) = \sum_{j,k=0}^{d+1} a_{j,k} z^j \xi^k \quad (3.67)$$

for certain real constants $a_{j,k}$ that satisfy $a_{k,j} = a_{j,k}$, and so in particular

$$a_{0,d+1} = a_{d+1,0} = 1.$$

It is also clear by changing $w \mapsto \omega w$ in (3.1) where $\omega = \omega_{d+1}$, that $P(z, \xi) = 0 \implies P(\omega z, \omega^{-1} \xi) = 0$. This implies

$$a_{j,k} \omega^{j-k} = 0, \quad \text{for all } j, k = 0, \dots, d \text{ with } \omega^{j-k} \neq 1.$$

Hence the only $a_{j,k}$ that are possibly non-zero are those with $j = k$ or $j = 0, k = d+1$, or $j = d+1, k = 0$. It thus follows that P has the form (2.16).

Inserting (3.1) into (2.16), we obtain a Laurent polynomial in w . Setting the coefficients in this expansion equal to 0, we obtain recursive relations that determine the coefficients c_k and β . In particular we find (2.17). It remains to show that the coefficients are positive.

In what follows we use the elementary symmetric function

$$e_k(x_1, \dots, x_n) = \sum_{j_1 < \dots < j_k} x_{j_1} x_{j_2} \cdots x_{j_k}$$

and we note that

$$c_{d+1-k} z^{d+1-k} = (-1)^{k-1} e_k(\xi_1(z), \dots, \xi_{d+1}(z)), \quad k = 1, \dots, d, \quad (3.68)$$

and

$$z^{d+1} + \beta = (-1)^{d+1} e_{d+1}(\xi_1(z), \dots, \xi_{d+1}(z)) = (-1)^{d+1} \prod_{j=1}^{d+1} \xi_j(z). \quad (3.69)$$

From (3.69) we get

$$\beta = (-1)^{d+1} \lim_{z \rightarrow 0} \prod_{j=1}^{d+1} \xi_j(z).$$

By (3.2) we have that $z = 0$ corresponds to w values that are solutions of $rw^{d+1} + a = 0$ where $a = t_{d+1}r^d$. This implies

$$\prod_{j=1}^{d+1} w_j(0) = (-1)^{d+1} \frac{a}{r}$$

Also by (3.1) we have

$$\xi_j(0) = \frac{aw_j(0)^{d+1} + r}{w_j(0)} = \frac{r^2 - a^2}{rw_j(0)}$$

and so

$$\beta = (-1)^{d+1} \prod_{j=1}^{d+1} \frac{r^2 - a^2}{rw_j(0)} = \left(\frac{r^2 - a^2}{r} \right)^{d+1} \frac{r}{a}. \quad (3.70)$$

Since $r < r_{\text{crit}} = (dt_{d+1})^{-1/(d-1)}$, see Figure 4, we easily get $dt_{d+1}r^{d-1} < 1$, and so

$$a = t_{d+1}r^d < \frac{r}{d} < r.$$

Then from (3.70) we see that $\beta > 0$.

In order to show that $c_k > 0$ for $k = 1, \dots, d-1$, we are going to prove the following claim.

Claim: For $k = 1, \dots, d-1$, we have that $(-1)^k e_k(\xi_2(z), \dots, \xi_{d+1}(z))$ has a Laurent expansion at infinity of the form

$$(-1)^k e_k(\xi_2(z), \dots, \xi_{d+1}(z)) = \frac{1}{z^k} \sum_{j=0}^{\infty} b_{j,k} z^{-j(d+1)} \quad (3.71)$$

with $b_{j,k} > 0$ for all j .

Proof of the claim: We use induction.

For $k = 1$, we note that by (3.68)

$$-e_1(\xi_2(z), \dots, \xi_{d+1}(z)) = \xi_1(z) - c_d z^d = t_0 F_1(z) \quad (3.72)$$

where we used (3.43) and the fact that $c_d = t_{d+1}$. Since F_1 is the Cauchy transform of μ_1^* we have

$$F_1(z) = \frac{1}{z} \left(1 + \sum_{j=1}^{\infty} s_j z^{-j(d+1)} \right) \quad \text{with} \quad s_j = \int_{\Sigma_1^*} z^{j(d+1)} d\mu_1^*(z). \quad (3.73)$$

By the rotational symmetry and the positivity of μ_1^* we have

$$s_j = (d+1) \int_0^{x^*} x^{j(d+1)} d\mu_1^*(x) > 0. \quad (3.74)$$

Combining (3.72)–(3.74) we have proved the claim for $k = 1$.

Now let $k \geq 2$ with $k \leq d-1$ and assume that the claim is true for $k-1$. The following basic relation for the elementary symmetric polynomials

$$e_k(\xi_2(z), \dots, \xi_{d+1}(z)) = e_k(\xi_1(z), \xi_2(z), \dots, \xi_{d+1}(z)) - \xi_1(z)e_{k-1}(\xi_2(z), \dots, \xi_{d+1}(z))$$

leads by (3.68), (3.71) and (3.72) to

$$(-1)^k e_k(\xi_2(z), \dots, \xi_{d+1}(z)) = -c_{d+1-k} z^{d+1-k} + (c_d z^d + t_0 F_1(z)) \frac{1}{z^{k-1}} \sum_{j=0}^{\infty} b_{j,k-1} z^{-j(d+1)}. \quad (3.75)$$

The term with z^{d+1-k} in the right-hand side cancels out, since it is clear from the asymptotic behavior (3.17) of the ξ_j functions that the elementary symmetric polynomial cannot grow like z^{d+1-k} as $z \rightarrow \infty$. This fact leads to

$$c_{d+1-k} = b_{0,k-1} c_d. \quad (3.76)$$

We then obtain from (3.73), (3.74) and the induction hypothesis that the right-hand side of (3.75) has a Laurent expansion of the form (3.71) with positive coefficients $b_{j,k}$, which proves the claim.

Having proved the claim, we know in particular that $b_{0,k} > 0$ for $k = 2, \dots, d-1$, and the relation (3.76) also holds for $k = 2, \dots, d-1$. These two facts imply that $c_k > 0$ for $k = 2, \dots, d-1$.

The relation (3.76) does not hold for $k = d$, instead we have

$$c_1 = \frac{1}{c_d} + b_{0,d-1} c_d = \frac{1}{t_{d+1}} + b_{0,d-1} t_{d+1}, \quad (3.77)$$

which implies $c_1 > 0$. So we finish this proof by justifying (3.77).

The relation (3.75) also holds for $k = d$, and it implies that

$$(-1)^d e_d(\xi_2(z), \dots, \xi_{d+1}(z)) = (-1)^d \prod_{j=2}^{d+1} \xi_j(z) = (-c_1 + c_d b_{0,d-1}) z + O(z^{-d})$$

as $z \rightarrow \infty$. On the other hand, from (3.69) we deduce

$$(-1)^d e_d(\xi_2(z), \dots, \xi_{d+1}(z)) = -\frac{z^{d+1} + \beta}{\xi_1(z)} = -(z^{d+1} + \beta) \left(\frac{1}{t_{d+1}} z^{-d} + O(z^{-2d-1}) \right),$$

as $z \rightarrow \infty$. Identifying the leading coefficients in both expansions we obtain (3.77).

4 The multiple orthogonal polynomials $P_{n,n}$ and the associated Riemann-Hilbert problem

In this section we start the proof of Theorem 2.3. We fix $t_{d+1} > 0$, $0 < t_0 < t_{0,\text{crit}}$ and let x^* be as (2.7). We also take $\hat{x} > x^*$ with $\hat{x} < x^{**}$ as in Theorem 2.2 and let $(\mu_1^*, \dots, \mu_d^*)$ be the minimizer of the vector equilibrium problem as in Theorem 2.2. We also use the notions that were developed in Sections 3.1 and 3.3, namely the functions ξ_k given in (3.6) and the Cauchy transforms F_k given in (3.42) that satisfy the conditions of Lemma 3.9. These functions will come to play a role in the third transformation of the steepest descent analysis.

Throughout we assume that n is a multiple of d .

4.1 The Riemann-Hilbert problem

We start with the formulation of the Riemann-Hilbert problem for the polynomials $P_{n,n}$. This is based on the multiple orthogonality stated in Lemma 1.3. As a result [35] there is a Riemann-Hilbert problem of size $(d+1) \times (d+1)$.

Recall that the weight functions $w_{j,n}$ are related to the functions p_ℓ as in (1.26)–(1.29). The functions p_ℓ are solutions of (1.13). There are $d+1$ different functions p_ℓ and any d of them form a basis for the solution space of (1.13). It is immediate from (1.26) that

$$\sum_{\ell=0}^d p_\ell \equiv 0, \quad (4.1)$$

and

$$p_\ell(z) = \omega^\ell p_0(\omega^\ell z), \quad \ell = 0, \dots, d, \quad (4.2)$$

recall $\omega = \omega_{d+1} = e^{\frac{2\pi i}{d+1}}$.

It will be convenient to forget about the constant prefactors in (1.29), and therefore we introduce the following definition.

Definition 4.1. For $n \in \mathbb{N}$, and $j = 0, \dots, d-1$, we put

$$v_{j,n}(z) = e^{\frac{nV(z)}{t_0}} p_{-\ell}^{(j)}(c_n z), \quad \arg z = \frac{2\pi}{d+1}\ell. \quad (4.3)$$

By (1.29) and (4.3), the weight $v_{j,n}$ is a constant multiple of $w_{j,n}$, and consequently the polynomial $P_{n,n}$ may be defined alternatively through the multiple orthogonality conditions

$$\int_{\Sigma} P_{n,n}(z) z^k v_{j,n}(z) dz = 0, \quad j = 0, \dots, \left\lceil \frac{n-j}{d} \right\rceil - 1, \quad j = 0, \dots, d-1. \quad (4.4)$$

We now introduce the RH problem that is associated with the polynomials $P_{n,n}$ and which will be the subject of analysis during the rest of the paper. Recall that the star Σ is given the outward radial orientation. This induces a $+$ and $-$ side on each segment $[0, \omega^j \hat{x}]$ of Σ , where the $+$ side ($-$ side) is on the left (right) as we traverse the ray according to its orientation. Recall that the star Σ is also denoted by Σ_1 .

RH problem 4.2. Find a function $\mathbf{Y} : \mathbb{C} \setminus \Sigma \rightarrow \mathbb{C}^{(d+1) \times (d+1)}$ with the following properties:

- \mathbf{Y} is analytic in $\mathbb{C} \setminus \Sigma$.
- For every $z \in \Sigma$, the function \mathbf{Y} has boundary values $\mathbf{Y}_+(z)$ and $\mathbf{Y}_-(z)$ and they are related by $\mathbf{Y}_+(z) = \mathbf{Y}_-(z)\mathbf{J}_Y(z)$, where

$$\mathbf{J}_Y(z) = \begin{pmatrix} 1 & v_{0,n}(z) & v_{1,n}(z) & \cdots & v_{d-1,n}(z) \\ & 1 & & & \\ & & 1 & & \\ & & & \ddots & \\ & & & & 1 \end{pmatrix}, \quad z \in \Sigma. \quad (4.5)$$

- As $z \rightarrow \infty$,

$$\mathbf{Y}(z) = \left(I + O\left(\frac{1}{z}\right) \right) \text{diag} \left(z^n, z^{-\frac{n}{d}}, z^{-\frac{n}{d}}, \dots, z^{-\frac{n}{d}} \right), \quad (4.6)$$

where I denotes the identity matrix of size $(d+1) \times (d+1)$.

- As z approaches one of the endpoints $\omega^\ell \hat{x}$ of Σ ,

$$\mathbf{Y}(z) = \begin{pmatrix} O(1) & O(\log(z - \omega_{d+1}^\ell \hat{x})) & \cdots & O(\log(z - \omega_{d+1}^\ell \hat{x})) \\ O(1) & O(\log(z - \omega_{d+1}^\ell \hat{x})) & \cdots & O(\log(z - \omega_{d+1}^\ell \hat{x})) \\ \vdots & \vdots & & \vdots \\ O(1) & O(\log(z - \omega_{d+1}^\ell \hat{x})) & \cdots & O(\log(z - \omega_{d+1}^\ell \hat{x})) \end{pmatrix}, \quad (4.7)$$

and $\mathbf{Y}(z) = O(1)$ as $z \rightarrow 0$.

The relevance of this RH problem relies on the fact that it characterizes the multiple orthogonal polynomials $P_{n,n}$, as it was already indicated. This characterization result is now standard in the theory of RH problems associated with multiple orthogonal polynomials, and was first obtained in [35] in the context of multiple orthogonality on the real line. It generalizes the RH problem for orthogonal polynomials [8, 16].

Lemma 4.3. *The polynomial $P_{n,n}$ exists and is unique if and only if the RH problem 4.2 is solvable. In this case we have $\mathbf{Y}_{1,1} = P_{n,n}$.*

We obtain the large n asymptotics of the RH problem 4.2 by a steepest descent analysis. The analysis is rather involved, since we are dealing with a RH problem of size $(d+1) \times (d+1)$, where $d \geq 2$ can be any natural number. See [9] for an earlier example of a steepest descent analysis for RH problems of arbitrary size.

One of the outcomes of the steepest descent analysis is that the RH problem 4.2 has a solution for n large enough, and so the polynomials $P_{n,n}$ exist for n large enough. The main outcome is the asymptotic formula (2.11) for the polynomials $P_{n,n}$, as given in Theorem 2.3.

The steepest descent analysis proceeds via a number of transformations.

The sectors S_ℓ and their subsectors S_ℓ^\pm were introduced in (3.15) and (3.16), see also Figures 6–7. There are $d+1$ sectors S_ℓ and ℓ is considered modulo $d+1$. In many cases it does not matter which value modulo $d+1$ we take, but in some cases it does matter. The canonical choice is indicated in Figures 6–7. Roughly speaking the index ℓ runs from $-\frac{d}{2}$ to $\frac{d}{2}$. To be precise, we use the following convention

$$\begin{aligned} \text{for } S_\ell^+ : \quad \ell &= -\lceil \frac{d}{2} \rceil, \dots, \lfloor \frac{d}{2} \rfloor \\ \text{for } S_\ell^- : \quad \ell &= -\lfloor \frac{d}{2} \rfloor, \dots, \lceil \frac{d}{2} \rceil. \end{aligned} \quad (4.8)$$

We use $\omega = \omega_{d+1} = e^{\frac{2\pi i}{d+1}}$ as before.

We also let $E_{j,k}$ be the elementary matrix with 1 in position (j, k) and 0 elsewhere. The size of $E_{j,k}$ will be clear from the context. It is either size $d \times d$ or $(d+1) \times (d+1)$.

The case $d = 2$ is essentially done in [7]. In this proof we assume $d \geq 3$. Then

$$\mathbb{C} \setminus \bigcup_{\ell=0}^d (S_\ell^+ \cup S_\ell^-) = \Sigma_2 \cup \Sigma_3.$$

For a function that is analytic in $\bigcup_{\ell=0}^d (S_\ell^+ \cup S_\ell^-)$ we describe its boundary values separately on Σ_2 and Σ_3 .

5 First transformation $\mathbf{Y} \mapsto \mathbf{X}$

The first transformation is based on the construction of a $d \times d$ -matrix valued function $\mathbf{F} : \mathbb{C} \setminus (\Sigma_2 \cup \Sigma_3) \rightarrow \mathbb{C}^{d \times d}$ out of solutions of the ODE (1.13). In each sector we pick a basis

f_1, \dots, f_d of solutions of (1.13) and we define \mathbf{F} as the Wronskian matrix

$$\mathbf{F} := \begin{pmatrix} f_1 & f_2 & \cdots & f_d \\ f'_1 & f'_2 & \cdots & f'_d \\ \vdots & \vdots & \ddots & \vdots \\ f_1^{(d-1)} & f_2^{(d-1)} & \cdots & f_d^{(d-1)} \end{pmatrix}. \quad (5.1)$$

5.1 Definition and properties of \mathbf{P}

We start from the functions p_ℓ from (1.26) that are special solutions of (1.13). We use cyclic notation here so that

$$p_\ell = p_{d+1+\ell}.$$

The solution p_0 is recessive in the sector S_0 . Indeed a classical steepest descent analysis shows that

$$p_0(z) = \frac{1}{\sqrt{2\pi d}} z^{-\frac{d-1}{2d}} e^{-\frac{d}{d+1}z^{\frac{d+1}{d}}} (1 + O(z^{-\frac{d+1}{d}})) \quad (5.2)$$

as $z \rightarrow \infty$ with $-\pi < \arg z < \pi$. Since

$$p_\ell(z) = \omega^\ell p_0(\omega^\ell z), \quad \omega = \omega_{d+1},$$

we also find

$$p_\ell(z) = \frac{1}{\sqrt{2\pi d}} e^{\frac{\ell}{d}\pi i} z^{-\frac{d-1}{2d}} e^{-\frac{d}{d+1}\omega^\ell z^{\frac{d+1}{d}}} (1 + O(z^{-\frac{d+1}{d}})), \quad \omega_d = e^{\frac{2\pi i}{d}}, \quad (5.3)$$

as $z \rightarrow \infty$ with $-\frac{2\ell}{d+1}\pi - \pi < \arg z < -\frac{2\ell}{d+1}\pi + \pi$. Thus $p_{-\ell}$ is recessive as $z \rightarrow \infty$ in sector S_ℓ .

The asymptotic behaviors (5.2)-(5.3) show that in each sector S_ℓ^\pm all possible asymptotic behaviors occur among the functions p_0, \dots, p_d . We order them according to their absolute value as $z \rightarrow \infty$ in that sector. Ordering them from smallest to largest, we obtain

$$\begin{aligned} \text{in } S_\ell^+ : & \quad p_{-\ell}, p_{-\ell-1}, p_{-\ell+1}, p_{-\ell-2}, \dots, p_{-\ell-(-1)^d \lfloor \frac{d}{2} \rfloor}, p_{-\ell+(-1)^d \lceil \frac{d}{2} \rceil} \\ \text{in } S_\ell^- : & \quad p_{-\ell}, p_{-\ell+1}, p_{-\ell-1}, p_{-\ell+2}, \dots, p_{-\ell+(-1)^d \lfloor \frac{d}{2} \rfloor}, p_{-\ell-(-1)^d \lceil \frac{d}{2} \rceil}. \end{aligned} \quad (5.4)$$

There are $d+1$ entries in each list, but for the construction that follows we do not use the last one.

Definition 5.1. We define $\mathbf{P} : \mathbb{C} \setminus (\Sigma_2 \cup \Sigma_3) \rightarrow \mathbb{C}^{d \times d}$ as the piecewise analytic matrix valued function defined by

$$\mathbf{P} = \begin{pmatrix} p_{-\ell} & p_{-\ell \mp 1} & p_{-\ell \pm 1} & p_{-\ell \mp 2} & \cdots \\ p'_{-\ell} & p'_{-\ell \mp 1} & p'_{-\ell \pm 1} & p'_{-\ell \mp 2} & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ p_{-\ell}^{(d-1)} & p_{-\ell \mp 1}^{(d-1)} & p_{-\ell \pm 1}^{(d-1)} & p_{-\ell \mp 2}^{(d-1)} & \cdots \end{pmatrix}_{d \times d} \quad \text{in } S_\ell^\pm. \quad (5.5)$$

The ordering of the columns in (5.5) is according to (5.4). The last entries in (5.4) do not appear as a column in the matrix \mathbf{P} .

Note that \mathbf{P} has size $d \times d$ and so the last column of \mathbf{P} is

$$\begin{pmatrix} p_{-\ell \mp \frac{d}{2}} & \cdots & p_{-\ell \mp \frac{d}{2}}^{(d-1)} \end{pmatrix}^t \quad \text{if } d \text{ is even} \quad (5.6)$$

and

$$\begin{pmatrix} p_{-\ell \pm \frac{d-1}{2}} & \cdots & p_{-\ell \pm \frac{d-1}{2}}^{(d-1)} \end{pmatrix}^t \quad \text{if } d \text{ is odd.} \quad (5.7)$$

Then \mathbf{P} satisfies a RH problem that we state next.

RH problem 5.2. $\mathbf{P} : \mathbb{C} \setminus (\Sigma_2 \cup \Sigma_3) \rightarrow \mathbb{C}^{d \times d}$ satisfies

- \mathbf{P} is analytic in $\mathbb{C} \setminus (\Sigma_2 \cup \Sigma_3)$,
- $\mathbf{P}_+ = \mathbf{P}_- \mathbf{J}_P$ with jump matrices

$$\mathbf{J}_P = \begin{cases} \text{diag}(1, \sigma_1, \dots, \sigma_1, 0) - \sum_{j=1}^d E_{j,d} & \text{if } d \text{ is even,} \\ \text{diag}(1, \sigma_1, \dots, \sigma_1) & \text{if } d \text{ is odd,} \end{cases} \quad \text{on } \Sigma_3, \quad (5.8)$$

$$\mathbf{J}_P = \begin{cases} \text{diag}(\sigma_1, \dots, \sigma_1) & \text{if } d \text{ is even,} \\ \text{diag}(\sigma_1, \dots, \sigma_1, 0) - \sum_{j=1}^d E_{j,d} & \text{if } d \text{ is odd,} \end{cases} \quad \text{on } \Sigma_2, \quad (5.9)$$

$$\text{where } \sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

- As $z \rightarrow \infty$ in S_ℓ^\pm we have

$$\mathbf{P}(z) = \frac{1}{\sqrt{2\pi d}} \left(I_d + O(z^{-\frac{2}{d}}) \right) \mathbf{D}(z)^{-1} B_\ell^\pm \exp(-\Theta^\pm(\omega^{-\ell} z)) \quad (5.10)$$

where

$$\mathbf{D}(z) = \text{diag} \left(z^{\frac{d-1}{2d}}, z^{\frac{d-3}{2d}}, \dots, z^{-\frac{d-1}{2d}} \right) \quad (5.11)$$

$$\Theta^\pm(z) = \text{diag} \left(1, \omega_d^{\mp 1}, \omega_d^{\pm 1}, \omega_d^{\mp 2}, \dots \right) \frac{d}{d+1} z^{\frac{d+1}{d}}, \quad (5.12)$$

and constant matrices B_ℓ^\pm given by

$$B_0^\pm = \text{diag}(1, -1, 1, -1, \dots) VDM(1, \omega_d^{\mp 1}, \omega_d^{\pm 1}, \omega_d^{\mp 2}, \dots) \text{diag}(1, \omega_{2d}^{\mp 1}, \omega_{2d}^{\pm 1}, \omega_{2d}^{\mp 2}, \dots) \quad (5.13)$$

where VDM denotes a Vandermonde matrix

$$VDM(x_1, \dots, x_d) = (x_k^{j-1})_{j,k=1,\dots,d},$$

$$\omega_d = e^{\frac{2\pi i}{d}}, \omega_{2d} = e^{\frac{\pi i}{d}}, \text{ and}$$

$$B_\ell^\pm = \text{diag}(\omega_{2d}^{-\ell}, \omega_{2d}^{-3\ell}, \dots, \omega_{2d}^{-(2d-1)\ell}) B_0^\pm. \quad (5.14)$$

Proof. The jump matrix of \mathbf{P} on Σ_3 is $\mathbf{J}_P = \mathbf{P}_-^{-1} \mathbf{P}_+$ where \mathbf{P}_- is the limit of \mathbf{P} taken from S_ℓ^- and \mathbf{P}_+ is the limit from S_ℓ^+ onto Σ_3 , for some ℓ . Then the jump matrix (5.8) is immediate from the definition (5.5) in case d is odd. If d is even then the last column (5.6) in S_ℓ^+ is associated with $p_{-\ell-\frac{d}{2}}$ and this column does not appear in \mathbf{P} in S_ℓ^- . Because of the relation (4.1) the last

column in S_ℓ^+ is equal to minus the sum of all columns in S_ℓ^- , and this accounts for the term $-\sum_{j=1}^d E_{j,d}$ in the jump matrix (5.8) in case d is even.

The proof of (5.9) is similar.

To prove (5.10) we use the asymptotic behavior (5.3) of the functions p_ℓ and their derivatives, which can be obtained by taking the derivatives of the leading term in (5.3). Using this in (5.5) in the sector S_0^\pm it then follows that

$$\mathbf{P}(z) = \frac{1}{\sqrt{2\pi d}} \mathbf{D}(z)^{-1} B_0^\pm \left(I_d + O(z^{-\frac{d+1}{d}}) \right) \exp(-\Theta^\pm(z)) \quad (5.15)$$

as $z \rightarrow \infty$ in the sectors S_0^\pm , with $\mathbf{D}(z)$, $\Theta^\pm(z)$ and B_0^\pm as in (5.11), (5.12), (5.13). Moving the O -term to the left we obtain (5.10) for $\ell = 0$ since

$$\mathbf{D}(z)^{-1} O(z^{-\frac{d+1}{d}}) = O(z^{-\frac{2}{d}}) \mathbf{D}(z)^{-1}$$

as $\mathbf{D}(z) = O(z^{\frac{d-1}{2d}})$ and $\mathbf{D}(z)^{-1} = O(z^{\frac{d-1}{2d}})$ as $z \rightarrow \infty$.

We finally use the identity

$$\mathbf{P}(z) = \text{diag}(\omega, \omega^2, \dots, \omega^d) \mathbf{P}(\omega z), \quad \omega = \omega_{d+1}$$

to obtain the asymptotic behavior (5.10) with matrices B_ℓ^\pm as in (5.14) in the other sectors. \square

The jump matrices in (5.8) and (5.9) have a block diagonal form, except partially for the last column. The term $-\sum_{j=1}^d E_{j,d}$ means that the last column is filled with -1 's.

Lemma 5.3. *Let*

$$\mathbf{B}(z) = \mathbf{D}(z)^{-1} B_\ell^\pm, \quad \text{for } z \in S_\ell^\pm. \quad (5.16)$$

where $\mathbf{D}(z)$ is as in (5.11) and the constant matrices B_ℓ^\pm are given in (5.13)–(5.14). Then \mathbf{B} is analytic in $\mathbb{C} \setminus (\Sigma_2 \cup \Sigma_3)$ and $\mathbf{B}_+ = \mathbf{B}_- \mathbf{J}_B$ with

$$\mathbf{J}_B = \begin{cases} \text{diag}(1, \sigma_1, \dots, \sigma_1, -1) & \text{if } d \text{ is even} \\ \text{diag}(1, \sigma_1, \dots, \sigma_1) & \text{if } d \text{ is odd} \end{cases} \quad \text{on } \Sigma_3, \quad (5.17)$$

$$\mathbf{J}_B = \begin{cases} \text{diag}(\sigma_1, \sigma_1, \dots, \sigma_1) & \text{if } d \text{ is even} \\ \text{diag}(\sigma_1, \sigma_1, \dots, \sigma_1, -1) & \text{if } d \text{ is odd} \end{cases} \quad \text{on } \Sigma_2. \quad (5.18)$$

Proof. It is a straightforward calculation from (5.13) to show that

$$(\mathbf{B}_0^-)^{-1} \mathbf{B}_0^+ = \begin{cases} \text{diag}(1, \sigma_1, \dots, \sigma_1, -1) & \text{if } d \text{ is even} \\ \text{diag}(1, \sigma_1, \dots, \sigma_1) & \text{if } d \text{ is odd} \end{cases} \quad (5.19)$$

and using also (5.14) that

$$(\mathbf{B}_0^+)^{-1} \mathbf{B}_1^- = \begin{cases} \text{diag}(\sigma_1, \dots, \sigma_1) & \text{if } d \text{ is even} \\ \text{diag}(\sigma_1, \dots, \sigma_1, -1) & \text{if } d \text{ is odd.} \end{cases} \quad (5.20)$$

It is then easy to see from the definitions (5.14) and (5.16) that we immediately obtain the jump matrices (5.17) on $\Sigma_3 \setminus \mathbb{R}_-$ and (5.18) on $\Sigma_2 \setminus \mathbb{R}_-$.

To compute the jump on the negative real axis (which is oriented from right to left) we note that

$$\begin{aligned} \mathbf{J}_B &= \mathbf{B}_-^{-1}(z)\mathbf{B}_+(z) \\ &= \begin{cases} \left(B_{\frac{d}{2}}^+\right)^{-1} \mathbf{D}_-(z)\mathbf{D}_+(z)^{-1}B_{-\frac{d}{2}}^- & \text{if } d \text{ is even} \\ \left(B_{\frac{d+1}{2}}^-\right)^{-1} \mathbf{D}_-(z)\mathbf{D}_+(z)^{-1}B_{-\frac{d+1}{2}}^+ & \text{if } d \text{ is odd} \end{cases} \quad z \in \mathbb{R}_- \end{aligned} \quad (5.21)$$

where due to (5.11) and the choice of principle branches of the fractional powers

$$\mathbf{D}_-(z)\mathbf{D}_+(z)^{-1} = \text{diag}(\omega_{2d}^{d-1}, \omega_{2d}^{d-3}, \dots, \omega_{2d}^{-(d-1)}), \quad z \in \mathbb{R}_-. \quad (5.22)$$

Using (5.14) and (5.22) in (5.21) we find

$$\begin{aligned} \mathbf{J}_B &= (B_0^+)^{-1} \text{diag}(\omega_{2d}^{\frac{d}{2}}, \omega_{2d}^{\frac{3d}{2}}, \dots) \text{diag}(\omega_{2d}^{d-1}, \omega_{2d}^{d-3}, \dots, \omega_{2d}^{-(d-1)}) \text{diag}(\omega_{2d}^{\frac{d}{2}}, \omega_{2d}^{\frac{3d}{2}}, \dots) B_0^- \\ &= (B_0^+)^{-1} \text{diag}(\omega_{2d}^{-1}, \omega_{2d}^{-3}, \dots) B_0^- \\ &= (B_0^+)^{-1} B_1^- \end{aligned}$$

if d is even, while

$$\begin{aligned} \mathbf{J}_B &= (B_0^-)^{-1} \text{diag}(\omega_{2d}^{\frac{d+1}{2}}, \omega_{2d}^{\frac{3d+1}{2}}, \dots) \text{diag}(\omega_{2d}^{d-1}, \omega_{2d}^{d-3}, \dots, \omega_{2d}^{-(d-1)}) \text{diag}(\omega_{2d}^{\frac{d+1}{2}}, \omega_{2d}^{\frac{3d+1}{2}}, \dots) B_0^+ \\ &= (B_0^-)^{-1} B_0^+ \end{aligned}$$

if d is odd. In view of (5.19) and (5.20) we then have

$$\mathbf{J}_B = \begin{cases} \text{diag}(\sigma_1, \sigma_1, \dots, \sigma_1) & \text{if } d \text{ is even} \\ \text{diag}(1, \sigma_1, \sigma_1, \dots, \sigma_1) & \text{if } d \text{ is odd} \end{cases} \quad \text{on } \mathbb{R}_-.$$

Since $\mathbb{R}_- \subset \Sigma_2$ if d is even, and $\mathbb{R}_- \subset \Sigma_3$ if d is odd, we verified the jump matrix (5.17) and (5.18) also for the part \mathbb{R}_- . \square

5.2 Definition and properties of \mathbf{F}

The jumps (5.8)–(5.9) are not convenient for us. We modify \mathbf{P} by multiplying it on the right by certain constant matrices R^\pm that we define first. We use the elementary symmetric functions on n variables:

$$e_{0,n}(x_1, \dots, x_n) = 1 \quad (5.23)$$

$$e_{k,n}(x_1, \dots, x_n) = \sum_{1 \leq j_1 < j_2 < \dots < j_k \leq n} x_{j_1} \cdots x_{j_k}, \quad 1 \leq k \leq n, \quad (5.24)$$

and define numbers

$$e_{k,n}^+ := e_{k,n}(\omega, \omega^{-1}, \omega^2, \omega^{-2}, \omega^3, \dots), \quad e_{k,n}^- := e_{k,n}(\omega^{-1}, \omega, \omega^{-2}, \omega^2, \omega^{-3}, \dots), \quad \omega = \omega_{d+1}, \quad (5.25)$$

with the understanding that $e_{0,0}^+ = e_{0,0}^- = 1$ and $e_{k,n}^+ = e_{k,n}^- = 0$ for $k < 0$ or $k > n$. Then we put for $j, k = 1, \dots, d$,

$$(R^+)_{j,k} := (-1)^{1+\lfloor \frac{j}{2} \rfloor + \lceil \frac{k}{2} \rceil} \times \begin{cases} e_{\lfloor \frac{k}{2} \rfloor + \frac{j-1}{2}, k-1}^+ & \text{if } j \text{ is odd} \\ e_{\lfloor \frac{k}{2} \rfloor - \frac{j}{2}, k-1}^+ & \text{if } j \text{ is even} \end{cases} \quad (5.26)$$

$$(R^-)_{j,k} := (-1)^{\lfloor \frac{j}{2} \rfloor + \lceil \frac{k}{2} \rceil} \times \begin{cases} e_{\lfloor \frac{k}{2} \rfloor + \frac{j-1}{2}, k-1}^- & \text{if } j \text{ is odd} \\ e_{\lfloor \frac{k}{2} \rfloor - \frac{j}{2}, k-1}^- & \text{if } j \text{ is even.} \end{cases} \quad (5.27)$$

For example, for $d = 4$ we have $\omega = \omega_5 = \exp(2\pi i/5)$ and

$$R^+ = \begin{pmatrix} 1 & e_{1,1}(\omega) & -e_{1,2}(\omega, \omega^{-1}) & -e_{2,3}(\omega, \omega^{-1}, \omega^2) \\ 0 & -1 & 1 & e_{1,3}(\omega, \omega^{-1}, \omega^2) \\ 0 & 0 & e_{2,2}(\omega, \omega^{-1}) & e_{3,3}(\omega, \omega^{-1}, \omega^2) \\ 0 & 0 & 0 & -1 \end{pmatrix}$$

$$R^- = \begin{pmatrix} 1 & -e_{1,1}(\omega^{-1}) & -e_{1,2}(\omega^{-1}, \omega) & e_{2,3}(\omega^{-1}, \omega, \omega^{-2}) \\ 0 & 1 & 1 & -e_{1,3}(\omega^{-1}, \omega, \omega^{-2}) \\ 0 & 0 & e_{2,2}(\omega^{-1}, \omega) & -e_{3,3}(\omega^{-1}, \omega, \omega^{-2}) \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

It can be readily verified that R^\pm are upper triangular matrices with diagonal entries

$$(R^+)_{k,k} = (-1)^{k-1}, \quad (R^-)_{k,k} = 1. \quad (5.28)$$

Definition 5.4. We define $\mathbf{F} : \mathbb{C} \setminus (\Sigma_2 \cup \Sigma_3) \rightarrow \mathbb{C}^{d \times d}$ by

$$\mathbf{F} = \mathbf{P}R^\pm \quad \text{in } S_\ell^\pm \quad (5.29)$$

where \mathbf{P} is defined in (5.5) and R^\pm are the right upper triangular matrices defined above.

Then \mathbf{F} has the form (5.1) with functions f_1, \dots, f_d where f_k in S_ℓ^\pm is a linear combination of the first k functions in $p_{-\ell}, p_{-\ell \mp 1}, p_{-\ell \pm 1}, \dots$. It turns out that \mathbf{F} is the solution of a RH problem with lower triangular jump matrices with a simple 2×2 block structure.

RH problem 5.5. \mathbf{F} is the solution of the following RH problem:

- \mathbf{F} is analytic in $\mathbb{C} \setminus (\Sigma_2 \cup \Sigma_3)$.
- $\mathbf{F}_+ = \mathbf{F}_- \mathbf{J}_F$ on $\Sigma_2 \cup \Sigma_3$ where

$$\mathbf{J}_F = I - \sum_{j=1}^{\lfloor \frac{d-1}{2} \rfloor} E_{2j+1, 2j} \quad \text{on } \Sigma_3, \quad (5.30)$$

$$\mathbf{J}_F = \text{diag}(\omega, \omega^{-1}, \omega^2, \omega^{-2}, \dots) - \sum_{j=1}^{\lfloor \frac{d}{2} \rfloor} E_{2j, 2j-1} \quad \text{on } \Sigma_2. \quad (5.31)$$

- \mathbf{F} has the following asymptotic behavior as $z \rightarrow \infty$,

$$\mathbf{F}(z) = \frac{1}{\sqrt{2\pi d}} \left(I_d + O(z^{-\frac{2}{d}}) \right) \mathbf{D}(z)^{-1}$$

$$\times \begin{cases} B_\ell^+ \text{diag}(1, -1, 1, -1, \dots) \exp(-\Theta^+(\omega^{-\ell} z)), & \text{in } S_\ell^+ \text{ with } -\lceil \frac{d}{2} \rceil \leq \ell \leq \lfloor \frac{d}{2} \rfloor, \\ B_\ell^- \exp(-\Theta^-(\omega^{-\ell} z)), & \text{in } S_\ell^- \text{ with } -\lfloor \frac{d}{2} \rfloor \leq \ell \leq \lceil \frac{d}{2} \rceil, \end{cases} \quad (5.32)$$

with $\mathbf{D}(z)$, $\Theta^\pm(z)$ and B_ℓ^\pm as defined in (5.11), (5.12) and (5.13)–(5.14).

Proof. The jump matrix is given by $\mathbf{J}_F = (R^-)^{-1}\mathbf{J}_P R^+$ on Σ_3 and $\mathbf{J}_F = (R^+)^{-1}\mathbf{J}_P R^-$ on Σ_2 . Here we insert \mathbf{J}_P as given by (5.8)–(5.9) and the definitions (5.26) and (5.27) of R^+ and R^- . Then after straightforward calculations we find that this indeed leads to (5.30)–(5.31).

The asymptotic condition comes from (5.10) and the fact that f_k is a linear combination of the first k of the p_j functions. In each sector these are ordered according to their asymptotic behavior at infinity in that sector, see (5.4). Thus the asymptotic behavior of f_k is given by that of the k th member times a constant coming from the diagonal entry of R^+ or R^- . The diagonal entries are ± 1 according to (5.28) and this leads to (5.32) because of (5.10) and (5.28). \square

5.3 Definition and properties of \mathbf{Q}

Let us introduce the matrix \mathbf{Q} that is employed in the first transformation of the RH problem 4.2.

Definition 5.6. We define \mathbf{Q} as

$$\mathbf{Q}(z) := \mathbf{F}(z)^{-t}, \quad (5.33)$$

that is, the inverse transpose of (5.29).

From the RH problem 5.5 we obtain the following RH problem for \mathbf{Q} .

RH problem 5.7. \mathbf{Q} is the solution of the following RH problem:

- \mathbf{Q} is analytic in $\mathbb{C} \setminus (\Sigma_2 \cup \Sigma_3)$.
- $\mathbf{Q}_+ = \mathbf{Q}_- \mathbf{J}_Q$ on $\Sigma_2 \cup \Sigma_3$ where

$$\mathbf{J}_Q = I_d + \sum_{j=1}^{\lfloor \frac{d-1}{2} \rfloor} E_{2j, 2j+1} \quad \text{on } \Sigma_3, \quad (5.34)$$

$$\mathbf{J}_Q = \text{diag}(\omega^{-1}, \omega, \omega^{-2}, \omega^2, \dots) + \sum_{j=1}^{\lfloor \frac{d}{2} \rfloor} E_{2j-1, 2j} \quad \text{on } \Sigma_2. \quad (5.35)$$

- \mathbf{Q} has the asymptotic behavior

$$\mathbf{Q}(z) = \sqrt{2\pi d} \left(I_d + O(z^{-\frac{2}{d}}) \right) \mathbf{D}(z) A_\ell^\pm \exp(\Theta^\pm(\omega^{-\ell} z)) \quad (5.36)$$

as $z \rightarrow \infty$ in S_ℓ^\pm , where

$$\begin{aligned} A_\ell^+ &= (B_\ell^+)^{-t} \text{diag}(1, -1, 1, -1, \dots), \\ A_\ell^- &= (B_\ell^-)^{-t}. \end{aligned} \quad (5.37)$$

Note that the jump matrices for \mathbf{Q} are upper triangular with a 2×2 block diagonal structure. For later use we define

$$\mathbf{A}(z) = \mathbf{D}(z) A_\ell^\pm \quad \text{for } z \in S_\ell^\pm. \quad (5.38)$$

The following is immediate from Lemma 5.3 and (5.37)–(5.38).

Lemma 5.8. \mathbf{A} is analytic in $\mathbb{C} \setminus (\Sigma_2 \cup \Sigma_3)$ and $\mathbf{A}_+ = \mathbf{A}_- \mathbf{J}_A$ with

$$\mathbf{J}_A = \begin{cases} \text{diag}(1, i\sigma_2, \dots, i\sigma_2, 1) & \text{if } d \text{ is even} \\ \text{diag}(1, i\sigma_2, \dots, i\sigma_2) & \text{if } d \text{ is odd} \end{cases} \quad \text{on } \Sigma_3, \quad (5.39)$$

$$\mathbf{J}_A = \begin{cases} \text{diag}(i\sigma_2, i\sigma_2, \dots, i\sigma_2) & \text{if } d \text{ is even} \\ \text{diag}(i\sigma_2, i\sigma_2, \dots, i\sigma_2, -1) & \text{if } d \text{ is odd} \end{cases} \quad \text{on } \Sigma_2. \quad (5.40)$$

where $i\sigma_2 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$.

Proof. We have $\mathbf{J}_A = (\mathbf{J}_B)^{-t} \text{diag}(1, -1, 1, -1, \dots)$ on Σ_3 and $\mathbf{J}_A = \text{diag}(1, -1, 1, -1, \dots)(\mathbf{J}_B)^{-t}$ on Σ_2 . Then use (5.17) and (5.18). \square

5.4 Transformation $\mathbf{Y} \mapsto \mathbf{X}$

Definition 5.9. Let $\Sigma_X = \Sigma_2 \cup \Sigma_3$ and define $\mathbf{X} : \mathbb{C} \setminus \Sigma_X \rightarrow \mathbb{C}^{(d+1) \times (d+1)}$ by

$$\mathbf{X}(z) := \begin{pmatrix} 1 & 0 \\ 0 & \frac{1}{\sqrt{2\pi d}} \mathbf{D}(c_n)^{-1} \end{pmatrix} \mathbf{Y}(z) \begin{pmatrix} 1 & 0 \\ 0 & \mathbf{Q}(c_n z) \end{pmatrix} \quad (5.41)$$

where c_n is given in (1.28), \mathbf{D} is the diagonal matrix (5.11), and \mathbf{Q} is the matrix (5.33).

Note that \mathbf{D} and \mathbf{Q} are matrices of size $d \times d$, and (5.41) is written in a block form with diagonal blocks of size 1×1 and $d \times d$.

RH problem 5.10. The matrix $\mathbf{X}(z)$ is the solution to the following RH problem:

- \mathbf{X} is analytic in $\mathbb{C} \setminus \Sigma_X$.
- $\mathbf{X}_+ = \mathbf{X}_- \mathbf{J}_X$ on $\Sigma_2 \cup \Sigma_3$ with jump matrix

$$\mathbf{J}_X = I_{d+1} + e^{\frac{n}{i_0} V} E_{1,2} + \sum_{j=2}^{\lfloor \frac{d}{2} \rfloor} E_{2j-1, 2j} \quad \text{on } \Sigma_1 \quad (5.42)$$

$$\mathbf{J}_X = I_{d+1} + \sum_{j=2}^{\lfloor \frac{d}{2} \rfloor} E_{2j-1, 2j} \quad \text{on } \Sigma_3 \setminus \Sigma_1 \quad (5.43)$$

$$\mathbf{J}_X = \text{diag}(1, \omega^{-1}, \omega, \omega^{-2}, \omega^2, \dots) + \sum_{j=1}^{\lfloor \frac{d}{2} \rfloor} E_{2j, 2j+1} \quad \text{on } \Sigma_2. \quad (5.44)$$

- As z tends to infinity in the sector S_ℓ^\pm ,

$$\mathbf{X}(z) = \left(I_{d+1} + O\left(z^{-\frac{2}{d}}\right) \right) \begin{pmatrix} 1 & 0 \\ 0 & \mathbf{A}(z) \end{pmatrix} \begin{pmatrix} z^n & 0 \\ 0 & z^{-\frac{n}{d}} \exp(\Theta^\pm(\omega^{-\ell} c_n z)) \end{pmatrix}, \quad (5.45)$$

where $\mathbf{A}(z)$ is given by (5.38).

- \mathbf{X} satisfies the same endpoint condition (4.7) as \mathbf{Y} .

Proof. For $z \in \Sigma_1$, using (4.5) and (5.41) we find after simple computations that

$$\mathbf{X}_-^{-1}(z) \mathbf{X}_+(z) = \begin{pmatrix} 1 & (v_{0,n}(z) & \cdots & v_{d-1,n}(z)) \mathbf{Q}_+(c_n z) \\ 0 & \mathbf{Q}_-^{-1}(c_n z) \mathbf{Q}_+(c_n z) \end{pmatrix}. \quad (5.46)$$

By the jump (5.34) of \mathbf{Q} we have that the right lower block is

$$\mathbf{Q}_-^{-1}(c_n z) \mathbf{Q}_+(c_n z) = \mathbf{J}_Q = I_d + \sum_{j=1}^{\lfloor \frac{d-1}{2} \rfloor} E_{2j,2j+1}.$$

From the definitions (5.29) and (5.5), and the fact that R^\pm is upper triangular with (1, 1) entry equal to 1, see (5.28), we obtain that

$$\mathbf{F}(z) \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} = \mathbf{P}(z) \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} = \begin{pmatrix} p_{-\ell}(z) \\ p'_{-\ell}(z) \\ \vdots \\ p_{-\ell}^{(d-1)}(z) \end{pmatrix}, \quad z \in S_\ell.$$

Then by (4.3)

$$\mathbf{F}_+(c_n z) \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} = e^{-\frac{n}{t_0} V(z)} \begin{pmatrix} v_{0,n}(z) \\ v_{1,n}(z) \\ \vdots \\ v_{d-1,n}(z) \end{pmatrix}, \quad z \in \Sigma_1,$$

and therefore, since $\mathbf{Q} = \mathbf{F}^{-t}$, see (5.33),

$$(v_{0,n}(z) \quad \cdots \quad v_{d-1,n}(z)) \mathbf{Q}_+(c_n z) = e^{\frac{n}{t_0} V(z)} (1 \quad 0 \quad \cdots \quad 0),$$

which gives the vector in the first row of (5.46). This proves (5.42).

The computation of \mathbf{J}_X on $\Sigma_3 \setminus \Sigma_1$ and Σ_2 is straightforward, since \mathbf{Y} is analytic on these sets, and we just get the jump $\mathbf{J}_X = \text{diag}(1, \mathbf{J}_Q)$ with \mathbf{J}_Q given by (5.34)–(5.35).

The asymptotic condition (5.45) comes from combining the asymptotic conditions (4.6) and (5.36) in the RH problems for \mathbf{Y} and \mathbf{Q} .

Finally, it is clear that \mathbf{X} satisfies the same endpoint conditions as \mathbf{Y} . \square

6 Second transformation $\mathbf{X} \mapsto \mathbf{U}$

The goal of the next transformation is to simplify the asymptotic condition (5.45).

Definition 6.1. We define $\mathbf{U}(z)$ by

$$\mathbf{U}(z) := \mathbf{X}(z) \begin{pmatrix} 1 & 0 \\ 0 & \exp(-\Theta^\pm(\omega^{-\ell} c_n z)) \end{pmatrix}, \quad z \in S_\ell^\pm. \quad (6.1)$$

In order to formulate the jump conditions for \mathbf{U} it is convenient to introduce here the following functions:

$$\lambda_{k,n}(z) = \exp\left(\frac{2 \operatorname{ind}}{(d+1)t_0 t_{d+1}^{1/d}} \sin\left(\frac{\pi k}{d}\right) |z|^{\frac{d+1}{d}}\right), \quad k \in \mathbb{N}. \quad (6.2)$$

We will use $\lambda_{2k,n}$ on Σ_3 and $\lambda_{2k-1,n}$ on Σ_2 . Note that $|\lambda_{k,n}(z)| = 1$ for all $z \in \mathbb{C}$, but when considered on Σ_2 or Σ_3 they allow for analytic continuation into a neighborhood.

RH problem 6.2. The matrix valued function $\mathbf{U}(z)$ is the solution to the following RH problem:

- \mathbf{U} is analytic in $\mathbb{C} \setminus \Sigma_U$ where $\Sigma_U = \Sigma_X = \Sigma_2 \cup \Sigma_3$.
- $\mathbf{U}_+ = \mathbf{U}_- \mathbf{J}_U$ on Σ_U with jump matrix

$$\mathbf{J}_U = \text{diag}(1, 1, \lambda_{2,n}, \lambda_{2,n}^{-1}, \lambda_{4,n}, \dots) + e^{-\frac{n}{t_0} V_1} E_{1,2} + \sum_{j=2}^{\lceil \frac{d}{2} \rceil} E_{2j-1, 2j} \quad \text{on } \Sigma_1 \quad (6.3)$$

where

$$V_1(z) = \frac{d}{(d+1)t_{d+1}^{1/d}} |z|^{\frac{d+1}{d}} - \frac{t_{d+1}}{d+1} z^{d+1}, \quad (6.4)$$

and

$$\mathbf{J}_U = \text{diag}(1, 1, \lambda_{2,n}, \lambda_{2,n}^{-1}, \lambda_{4,n}, \dots) + \sum_{j=2}^{\lceil \frac{d}{2} \rceil} E_{2j-1, 2j} \quad \text{on } \Sigma_3 \setminus \Sigma_1 \quad (6.5)$$

$$\mathbf{J}_U = \text{diag}(1, \omega^{-1} \lambda_{1,n}, \omega \lambda_{1,n}^{-1}, \omega^{-2} \lambda_{3,n}, \dots) + \sum_{j=1}^{\lfloor \frac{d}{2} \rfloor} E_{2j, 2j+1} \quad \text{on } \Sigma_2. \quad (6.6)$$

- As $z \rightarrow \infty$,

$$\mathbf{U}(z) = \left(I_{d+1} + O\left(z^{-\frac{2}{d}}\right) \right) \begin{pmatrix} 1 & 0 \\ 0 & \mathbf{A}(z) \end{pmatrix} \begin{pmatrix} z^n & 0 \\ 0 & z^{-\frac{n}{d}} I_d \end{pmatrix}. \quad (6.7)$$

- \mathbf{U} satisfies the same endpoint condition (4.7) as \mathbf{Y} .

Remark 6.3. The jump matrices (6.3), (6.5) and (6.6) have size $(d+1) \times (d+1)$. It means that the pattern on the diagonal repeats until we reach size $(d+1) \times (d+1)$. If d is even then the diagonals in (6.3) and (6.5) end with $\lambda_{d,n} \equiv 1$, see (6.2). If d is odd, then the diagonal in (6.6) ends with $\omega^{-(d+1)/2} \lambda_{d,n} \equiv -1$.

Proof. Since $\exp(-\Theta^\pm(\omega^{-\ell} c_n z))$ is a diagonal matrix it follows from (6.1) that the non-zero entries in \mathbf{J}_U are in the same positions as those of \mathbf{J}_X .

Let $z \in \Sigma_3$ with $z \in S_\ell$. Then $(\omega^{-\ell} z)^{\frac{d+1}{d}} = |z|^{\frac{d+1}{d}}$, which means that by (5.12) and (1.28)

$$\exp(-\Theta^\pm(\omega^{-\ell} c_n z)) = \exp\left(-\frac{nd}{(d+1)t_0 t_{d+1}^{1/d}} \text{diag}(1, \omega_d^{\mp 1}, \omega_d^{\pm 1}, \omega_d^{\mp 2}, \dots) |z|^{\frac{d+1}{d}}\right). \quad (6.8)$$

It then follows from (6.1) that the $(1, 2)$ entry in the jump matrix (5.42) on Σ_1 is multiplied by the first entry of (6.8) which by (6.4) results in

$$(\mathbf{J}_U(z))_{1,2} = e^{\frac{n}{t_0} V(z)} e^{-\frac{nd}{(d+1)t_0 t_{d+1}^{1/d}} |z|^{\frac{d+1}{d}}} = e^{-\frac{n}{t_0} V_1(z)}, \quad z \in \Sigma_1.$$

For $j \geq 2$, and $z \in \Sigma_3 \cap S_\ell$, we have that the $(2j-1, 2j)$ entry of \mathbf{J}_X is multiplied by the $2j-1$ entry of $\exp(-\Theta^+(\omega^{-\ell} c_n z))$ and by the inverse of the $2j-2$ entry of $\exp(-\Theta^-(\omega^{-\ell} c_n z))$. These factors cancel out because of the way the exponents $\omega_d^{\mp j}, \omega_d^{\pm j}$ of ω_d appear in (5.12). Thus

$$(\mathbf{J}_U(z))_{2j-1, 2j} = (\mathbf{J}_X(z))_{2j-1, 2j} = 1, \quad z \in \Sigma_3,$$

see (5.42)–(5.43).

The diagonal entry $(2j+1, 2j+1)$ of \mathbf{J}_X is multiplied by the $2j$ entry of $\exp(-\Theta^+(\omega^{-\ell}c_n z))$ and by the inverse of the $2j$ entry of $\exp(-\Theta^-(\omega^\ell c_n z))$. This leads to a combination

$$-\omega_d^{-j} + \omega_d^j = 2i \sin \frac{2\pi j}{d}$$

in the exponentials. Then by (5.42)–(5.43) and the definition (6.2) we get

$$(\mathbf{J}_U(z))_{2j+1, 2j+1} = \lambda_{2j, n}(z), \quad z \in \Sigma_3,$$

and similarly

$$(\mathbf{J}_U(z))_{2j+2, 2j+2} = \lambda_{2j, n}^{-1}(z), \quad z \in \Sigma_3.$$

This establishes (6.3) and (6.5).

The jump condition (6.6) on Σ_2 follows from similar considerations.

The other conditions in the RH problem 6.2 are immediate from (6.7) and the RH problem 5.10 for \mathbf{X} . \square

7 The third transformation $\mathbf{U} \mapsto \mathbf{T}$

7.1 The g -functions

In the third transformation we make use of the equilibrium measures described in Section 3. Let $(\mu_1^*, \dots, \mu_d^*)$ be the vector of measures solving the vector equilibrium problem (2.3)–(2.5). We associate to these measures the so-called g -functions.

Definition 7.1. We define for $k = 1, \dots, d$,

$$g_k(z) := \int \log(z-t) d\mu_k^*(t) = -U^{\mu_k^*}(z) + i \int \arg(z-t) d\mu_k^*(t), \quad z \in \mathbb{C} \setminus (\Sigma_k^* \cup \mathbb{R}^-), \quad (7.1)$$

where for each $t \in \Sigma_k^*$, $z \mapsto \arg(z-t)$ is defined with a branch cut along $\mathbb{R}^- \cup [0, t]$.

It follows from this definition that g_k is analytic in $\mathbb{C} \setminus (\Sigma_k^* \cup \mathbb{R}^-)$, with symmetry relations

$$g_k(\omega^\ell z) = g_k(z) + \frac{2\pi\ell i}{d+1} \|\mu_k^*\|, \quad z \in S_0, \quad \ell = \pm 1, \pm 2, \dots, \pm \lfloor \frac{d}{2} \rfloor, \quad (7.2)$$

where we recall that $\|\mu_k^*\| = 1 - \frac{k-1}{d}$.

Note also that $g'_k = F_k$ where F_k is the Cauchy transform of the measure μ_k^* as defined in (3.42). The g -functions therefore satisfy certain jump relations, which when differentiated give rise to (3.45)–(3.47). In the next lemma we state the relations for the g -function, where the main issue is the determination of the constants of integration.

Lemma 7.2. *The following relations hold:*

(a) For $z \in [0, \omega^\ell x^*] \subset \Sigma_1$, $\ell = 0, \pm 1, \dots, \pm \lfloor \frac{d}{2} \rfloor$,

$$g_{1,+}(z) + g_{1,-}(z) - g_2(z) = \frac{1}{t_0} \left(\frac{d}{(d+1)t_{d+1}^{1/d}} |z|^{\frac{d+1}{d}} - \frac{t_{d+1}}{d+1} z^{d+1} \right) + \frac{2\pi\ell i}{d} + \ell_1, \quad (7.3)$$

where ℓ_1 is the variational constant from (3.22).

In addition, in case d is odd, and $z \in [-x^*, 0] \subset \Sigma_1$,

$$g_{1,+}(z) + g_{1,-}(z) - g_{2,\pm}(z) = \frac{1}{t_0} \left(\frac{d}{(d+1)t_{d+1}^{1/d}} |z|^{\frac{d+1}{d}} - \frac{t_{d+1}}{d+1} z^{d+1} \right) \pm \frac{\pi i(d-1)}{d} + \ell_1. \quad (7.4)$$

(b) For $k \geq 2$,

$$g_{k,+}(z) + g_{k,-}(z) = g_{k+1}(z) + g_{k-1}(z), \quad z \in \Sigma_k \setminus \mathbb{R}^-, \quad (7.5)$$

where Σ_k is defined in (2.2), and in case that $k = d$, we understand $g_{d+1} \equiv 0$.

(c) For $k \geq 2$, and $k \equiv d \pmod{2}$.

$$g_{k,+}(z) + g_{k,-}(z) = g_{k+1,\pm}(z) + g_{k-1,\pm}(z) \pm 2\pi i \|\mu_k^*\|, \quad z \in \mathbb{R}^-. \quad (7.6)$$

In case $k = d$ we again understand $g_{d+1} \equiv 0$.

Proof. (a) From the definition of g_1 we deduce that for $z \in [0, x^*]$,

$$g_{1,+}(z) + g_{1,-}(z) = -2U^{\mu_1^*}(z).$$

Since g_2 is real-valued on \mathbb{R}^+ , we have $g_2(z) = -U^{\mu_2^*}(z)$ for $z \in (0, x^*]$. Applying now (3.22) and the fact $\text{supp}(\mu_1^*) = \Sigma_1^*$, we obtain (7.3) for $z \in [0, x^*]$. The remaining cases in (7.3) and (7.4) are then obtained by using the symmetry relations (7.2) of g_1 and g_2 .

(b) and (c) In order to show (7.5) and (7.6) it is convenient to consider the four cases k and d even/odd separately. We consider the case where k is even and d is odd. The other cases are analyzed in a similar manner.

So we assume that k is even and d is odd. We first check that (7.5) is valid for $x \in \Sigma_2$ with $\arg x = \frac{\pi}{d+1}$. By definition of g_k , we have

$$g_{k,+}(x) + g_{k,-}(x) = -2U^{\mu_k^*}(x) + \lim_{z \rightarrow x^-} i \int_{\Sigma_2} \arg(z-t) d\mu_k^*(t) + \lim_{z \rightarrow x^+} i \int_{\Sigma_2} \arg(z-t) d\mu_k^*(t). \quad (7.7)$$

Observe that since d is odd, $e^{\frac{\pi i}{d+1}} \mathbb{R} \subset \Sigma_2$ and we can write

$$\begin{aligned} \lim_{z \rightarrow x^-} \int_{\Sigma_2} \arg(z-t) d\mu_k^*(t) &= \int_{\Sigma_2 \setminus e^{\frac{\pi i}{d+1}} \mathbb{R}} \arg(x-t) d\mu_k^*(t) \\ &\quad + \frac{\pi}{d+1} \mu_k^* \left(e^{\frac{\pi i}{d+1}} \mathbb{R} \setminus (x, e^{\frac{\pi i}{d+1}} \infty) \right) + \left(-\pi + \frac{\pi}{d+1} \right) \mu_k^* \left((x, e^{\frac{\pi i}{d+1}} \infty) \right), \end{aligned}$$

where $-\pi < \arg(x-t) < \pi$. By the rotational invariance of μ_k^* ,

$$\int_{\Sigma_2 \setminus e^{\frac{\pi i}{d+1}} \mathbb{R}} \arg(x-t) d\mu_k^*(t) = \frac{(d-1)\pi}{(d+1)^2} \|\mu_k^*\|.$$

Since $\mu_k^*(e^{\frac{\pi i}{d+1}} \mathbb{R}) = 2\|\mu_k^*\|/(d+1)$, from the previous two identities we get

$$\lim_{z \rightarrow x^-} \int_{\Sigma_2} \arg(z-t) d\mu_k^*(t) = \frac{\pi \|\mu_k^*\|}{d+1} - \pi \mu_k^* \left((x, e^{\frac{\pi i}{d+1}} \infty) \right). \quad (7.8)$$

On the other hand, it is easy to see that

$$\lim_{z \rightarrow x^+} \int_{\Sigma_2} \arg(z-t) d\mu_k^*(t) = \lim_{z \rightarrow x^-} \int_{\Sigma_2} \arg(z-t) d\mu_k^*(t) + 2\pi \mu_k^* \left((x, e^{\frac{\pi i}{d+1}} \infty) \right). \quad (7.9)$$

Therefore from (7.7), (7.8) and (7.9) we obtain

$$g_{k,+}(x) + g_{k,-}(x) = -2U^{\mu_k^*}(x) + \frac{2\pi i \|\mu_k^*\|}{d+1}. \quad (7.10)$$

Now, the functions g_{k-1} and g_{k+1} are analytic on $e^{\frac{\pi i}{d+1}}\mathbb{R}^+$, and reasoning as before we deduce that

$$g_j(x) = -U^{\mu_j^*}(x) + \frac{\pi i \|\mu_j^*\|}{d+1}, \quad j = k-1, k+1. \quad (7.11)$$

It then follows from (3.23) and (7.10)–(7.11) that (7.5) holds for $x \in \Sigma_2$ with $\arg x = \pi/(d+1)$. By applying the symmetry relations (7.2) we deduce from this that (7.5) holds everywhere in $\Sigma_2 \setminus \mathbb{R}^- = \Sigma_2$. \square

7.2 The φ -functions

The jump matrices in the RH problem that follows can be expressed in a very convenient way in terms of certain functions that we now introduce.

Definition 7.3. Let φ_1 be the analytic function defined on $\mathbb{C} \setminus (\Sigma_1^* \cup \Sigma_2)$ by

$$\varphi_1(z) = \frac{1}{2t_0} \int_{\omega^\ell x^*}^z (\xi_1(s) - \xi_2(s)) ds, \quad z \in S_\ell \setminus [0, \omega^\ell x^*], \quad (7.12)$$

where for $z \in S_\ell \setminus [0, \omega^\ell x^*]$, integration is carried out in (7.12) along a path in $S_\ell \setminus [0, \omega^\ell x^*]$.

For $k \geq 2$, let φ_k be the analytic function on $\mathbb{C} \setminus (\Sigma_2 \cup \Sigma_3)$ given by

$$\varphi_k(z) = \frac{1}{2t_0} \int_0^z (\xi_k(s) - \xi_{k+1}(s)) ds \mp (-1)^k \frac{\pi i}{(d+1)} \|\mu_k^*\|, \quad z \in S_\ell^\pm, \quad (7.13)$$

where for $z \in S_\ell^\pm$, integration in (7.13) takes place along a path contained in S_ℓ^\pm .

Lemma 7.4. *The following relations hold:*

(a) For every $k = 1, \dots, d$,

$$\varphi_k(\omega z) = \varphi_k(z), \quad z \in \mathbb{C} \setminus (\Sigma_2 \cup \Sigma_3). \quad (7.14)$$

(b) For $z \in \Sigma_1^* \setminus [-x^*, 0]$,

$$g_{1,+}(z) - g_{1,-}(z) = \pm 2\varphi_{1,\pm}(z), \quad (7.15)$$

and, in case d is odd, for $z \in [-x^*, 0] \subset \Sigma_1^*$,

$$g_{1,+}(z) - g_{1,-}(z) = \pm 2\varphi_{1,\pm}(z) - 2\pi i. \quad (7.16)$$

(c) Let $2 \leq k \leq d$. Then for $z \in \Sigma_k \setminus \mathbb{R}^-$,

$$g_{k,+}(z) - g_{k,-}(z) = \pm 2\varphi_{k,\pm}(z) + \frac{2id \sin((k-1)\pi/d)}{(d+1)t_0 t_{d+1}^{1/d}} |z|^{\frac{d+1}{d}}, \quad (7.17)$$

and for $z \in \mathbb{R}^- \subset \Sigma_k$,

$$g_{k,+}(z) - g_{k,-}(z) = \pm 2\varphi_{k,\pm}(z) + \frac{2id \sin((k-1)\pi/d)}{(d+1)t_0 t_{d+1}^{1/d}} |z|^{\frac{d+1}{d}} - \frac{2\pi i(d-k+1)}{d}. \quad (7.18)$$

(d) For $z \in S_\ell \setminus [0, \omega^\ell x^*]$, $\ell = 0, \pm 1, \dots, \pm \lfloor d/2 \rfloor$,

$$2g_1(z) - g_2(z) - \ell_1 = 2\varphi_1(z) + \frac{1}{t_0} \left(\frac{d\omega_d^{-\ell}}{(d+1)t_{d+1}^{1/d}} z^{\frac{d+1}{d}} - \frac{t_{d+1}}{d+1} z^{d+1} \right) + \frac{2\pi\ell i}{d}, \quad (7.19)$$

and in case d is odd and for $z \in S_{\frac{d+1}{2}}^- \cup S_{-\frac{d+1}{2}}^+$,

$$2g_1(z) - g_2(z) - \ell_1 = 2\varphi_1(z) + \frac{1}{t_0} \left(\frac{d\omega_d^{\pm \frac{d+1}{2}}}{(d+1)t_{d+1}^{1/d}} z^{\frac{d+1}{d}} - \frac{t_{d+1}}{d+1} z^{d+1} \right) \mp \frac{\pi i(d+1)}{d}. \quad (7.20)$$

(e) For $z \in S_0^\pm$ and $2 \leq k \leq d$,

$$2g_k(z) - g_{k-1}(z) - g_{k+1}(z) = 2\varphi_k(z) + (-1)^k \frac{d(\omega_d^{\mp \lceil \frac{k-1}{2} \rceil} - \omega_d^{\pm \lfloor \frac{k-1}{2} \rfloor})}{(d+1)t_0 t_{d+1}^{1/d}} z^{\frac{d+1}{d}}, \quad (7.21)$$

with the convention $g_{d+1} \equiv 0$.

Proof. (a) The symmetry property (7.14) is immediate from (3.12) and (7.12)–(7.13).

(b) We first check that (7.15) holds for $x \in [0, x^*]$. Observe that for $s \in S_0 \setminus [0, x^*]$,

$$\xi_1(s) - \xi_2(s) = 2t_0 F_1(s) - t_0 F_2(s) - \frac{s^{1/d}}{t_{d+1}^{1/d}} + t_{d+1} s^d, \quad (7.22)$$

cf. (3.43) and (3.44). Therefore applying (3.45) we obtain from (7.12) and (7.1)

$$\begin{aligned} 2\varphi_{1,\pm}(x) &= -\frac{1}{t_0} \int_x^{x^*} \left(2t_0 F_{1,\pm}(s) - t_0 F_2(s) - \frac{s^{1/d}}{t_{d+1}^{1/d}} + t_{d+1} s^d \right) ds \\ &= -\int_x^{x^*} (F_{1,\pm}(s) - F_{1,\mp}(s)) ds \\ &= \pm 2\pi i \mu_1^*([x, x^*]) = \pm (g_{1,+}(x) - g_{1,-}(x)), \end{aligned}$$

and (7.15) follows for $z = x \in [0, x^*]$. Because of (7.14), we have for every $z \in \Sigma_1^*$,

$$\varphi_{1,+}(z) = \varphi_{1,+}(|z|) = -\varphi_{1,-}(|z|) = -\varphi_{1,-}(z).$$

Using now the symmetry property (7.2) we obtain (7.15)–(7.16).

(c) The proof of (7.17)–(7.18) is analogous. For instance, if we let

$$\widehat{\varphi}_{2j}(z) = \frac{1}{2t_0} \int_0^z (\xi_{2j}(s) - \xi_{2j+1}(s)) ds, \quad z \in \mathbb{C} \setminus (\Sigma_2 \cup \Sigma_3),$$

it follows from the definitions of ξ_{2j} and ξ_{2j+1} on the sectors S_0^+ and S_1^- , and relations (3.46) and (3.47), that for $x \in \Sigma_2$ with $\arg x = \pi/(d+1)$,

$$\begin{aligned} 2\widehat{\varphi}_{2j,\pm}(x) &= \frac{1}{t_0} \int_0^{|x|} \left(t_0 (F_{2j,\pm}(e^{\frac{\pi i}{d+1}} t) - F_{2j,\mp}(e^{\frac{\pi i}{d+1}} t)) \pm \frac{\omega_d^{-j} - \omega_d^{j-1}}{t_{d+1}^{1/d}} e^{\frac{\pi i}{d+1}} t^{1/d} \right) e^{\frac{\pi i}{d+1}} dt \\ &= \mp 2\pi i \mu_{2j}^*([0, x]) \mp \frac{2id \sin((2j-1)\pi/d)}{(d+1)t_0 t_{d+1}^{1/d}} |x|^{\frac{d+1}{d}}. \end{aligned} \quad (7.23)$$

Since

$$g_{2j,+}(x) - g_{2j,-}(x) = 2\pi i \mu_{2j}^*([x, e^{\frac{\pi i}{d+1}} \infty)) = \frac{2\pi i (d - 2j + 1)}{d(d+1)} - 2\pi i \mu_{2j}^*([0, x]), \quad (7.24)$$

we deduce from (7.23) and (7.24) that (7.17) holds for $k = 2j$ even. We use again the symmetry properties of the g and φ functions to check that the relations (7.15)–(7.16) are valid on the remaining rays of Σ_2 in case k is even. The case k odd can be proved in a similar way.

(d) Using $g'_k = F_k$ and $2\varphi'_1 = (\xi_1 - \xi_2)/t_0$, we can rewrite (7.22) as

$$2g'_1(z) - g'_2(z) = 2\varphi'_1(z) - \frac{t_{d+1}}{t_0} z^d + \frac{z^{1/d}}{t_0 t_{d+1}^{1/d}}, \quad z \in S_0 \setminus [0, x^*],$$

and integration yields

$$2g_1(z) - g_2(z) - \ell_1 = 2\varphi_1(z) + \frac{1}{t_0} \left(\frac{d}{(d+1)t_{d+1}^{1/d}} z^{\frac{d+1}{d}} - \frac{t_{d+1}}{d+1} z^{d+1} \right) + C,$$

for some constant C . It follows from (7.3) that $C = 0$, and this proves (7.19) for $\ell = 0$. The remaining cases in (7.19) and (7.20) follow again from symmetry considerations.

(e) The proof of (7.21) is analogous to the proof of (7.19)–(7.20) and uses the relations (7.5) and (7.17). \square

Lemma 7.5. *The following relations hold:*

(a) For $z \in \Sigma_1^*$,

$$e^{n(g_{1,-}(z) - g_{1,+}(z))} = e^{-2n\varphi_{1,+}(z)} = e^{2n\varphi_{1,-}(z)}. \quad (7.25)$$

(b) For $z \in \Sigma_2$,

$$e^{n(g_{2k,-}(z) - g_{2k,+}(z))} \lambda_{2k-1,n}(z) = e^{-2n\varphi_{2k,+}(z)} = e^{2n\varphi_{2k,-}(z)}, \quad 1 \leq k \leq \lfloor \frac{d}{2} \rfloor. \quad (7.26)$$

(c) For $z \in \Sigma_3$,

$$e^{n(g_{2k+1,-}(z) - g_{2k+1,+}(z))} \lambda_{2k,n}(z) = e^{-2n\varphi_{2k+1,+}(z)} = e^{2n\varphi_{2k+1,-}(z)}, \quad 1 \leq k \leq \lceil \frac{d}{2} \rceil - 1. \quad (7.27)$$

The functions $\lambda_{k,n}(z)$ are defined in (6.2).

Proof. The relations (7.25)–(7.27) are immediate from (7.15)–(7.18), since n is a multiple of d . \square

Lemma 7.6. *The following relations hold:*

(a) For $z \in \Sigma_1^*$ and k even, $2 \leq k \leq d$,

$$\varphi_{k,+}(z) - \varphi_{k,-}(z) = \varphi_{k+1,-}(z) + \varphi_{k-1,-}(z), \quad (7.28)$$

where $\varphi_{d+1} \equiv 0$.

(b) For $z \in \Sigma_2$ and k odd, we have

$$\varphi_{k,+}(z) - \varphi_{k,-}(z) = \begin{cases} \varphi_{2,-}(z) - \frac{\pi i}{d}, & \text{if } k = 1, \\ \varphi_{k+1,-}(z) + \varphi_{k-1,-}(z), & \text{if } k \geq 3, \end{cases} \quad (7.29)$$

where $\varphi_{d+1} \equiv 0$.

Proof. (a) By (7.14) it is enough to prove (7.28) on $(0, x^*)$. Assume that k is even with $k \geq 4$; the same proof with a slight variation can be used for $k = 2$. Applying (7.21) for $z \in S_0^+$,

$$g_{k-1}(z) + g_{k+1}(z) - g_{k-2}(z) - g_{k+2}(z) = 2 \sum_{i=k-1}^{k+1} \varphi_i(z) + \frac{d(\omega_d^{\frac{k}{2}} - \omega_d^{-(\frac{k}{2}-1)})z^{\frac{d+1}{d}}}{(d+1)t_0 t_{d+1}^{1/d}}, \quad (7.30)$$

and similarly for $z \in S_0^-$,

$$g_{k-1}(z) + g_{k+1}(z) - g_{k-2}(z) - g_{k+2}(z) = 2 \sum_{i=k-1}^{k+1} \varphi_i(z) + \frac{d(\omega_d^{-\frac{k}{2}} - \omega_d^{(\frac{k}{2}-1)})z^{\frac{d+1}{d}}}{(d+1)t_0 t_{d+1}^{1/d}}. \quad (7.31)$$

We let z tend to the interval $(0, x^*)$ in both (7.30)–(7.31), which leads to a $+$ -boundary value in (7.30) and a $-$ -boundary value in (7.31). We subtract the results and we get for $z \in (0, x^*)$,

$$\begin{aligned} & g_{k-1,+}(z) - g_{k-1,-}(z) + g_{k+1,+}(z) - g_{k+1,-}(z) \\ &= 2 \sum_{i=k-1}^{k+1} (\varphi_{i,+}(z) - \varphi_{i,-}(z)) + \frac{2id(\sin(\pi k/d) + \sin(\pi(k-2)/d))}{(d+1)t_0 t_{d+1}^{1/d}}, \end{aligned}$$

where we used the fact that for $i = k-2, k+2$, the boundary values $g_{i,\pm}(z)$ are the same. Now (7.28) follows immediately from (7.17). In the case $k = 2$ one also uses (7.19).

(b) The proof of (7.29) is completely analogous, and again by (7.14) it is sufficient to prove these relations for $z \in \Sigma_2$ with $\arg z = \pi/(d+1)$. From (7.21) and the symmetry properties of the g and φ functions one deduces that for $z \in S_1^-$ and $2 \leq k \leq d$,

$$2g_k(z) - g_{k-1}(z) - g_{k+1}(z) = 2\varphi_k(z) + (-1)^k \frac{d(\omega_d^{\lceil \frac{k-3}{2} \rceil} - \omega_d^{-\lfloor \frac{k+1}{2} \rfloor})z^{\frac{d+1}{d}}}{(d+1)t_0 t_{d+1}^{1/d}}.$$

The details are left to the reader. □

7.3 Transformation $\mathbf{U} \mapsto \mathbf{T}$

We now define the third transformation of the steepest descent analysis.

Definition 7.7. The matrix valued function $\mathbf{T}(z)$ is defined by

$$\mathbf{T}(z) = L^n \mathbf{U}(z) \mathbf{G}(z)^n L^{-n}, \quad z \in \mathbb{C} \setminus (\Sigma_2 \cup \Sigma_3), \quad (7.32)$$

where $\mathbf{G}(z)$ and L are the diagonal matrices of size $(d+1) \times (d+1)$ given by

$$\begin{aligned} \mathbf{G}(z) &= \text{diag}(e^{-g_1(z)}, e^{g_1(z)-g_2(z)}, e^{g_2(z)-g_3(z)}, \dots, e^{g_{d-1}(z)-g_d(z)}, e^{g_d(z)}), \\ L &= \text{diag}(e^{-\ell_1}, 1, 1, \dots, 1), \end{aligned}$$

where ℓ_1 is the constant in (3.22).

RH problem 7.8. The matrix \mathbf{T} is the solution to the following RH problem:

- \mathbf{T} is analytic in $\mathbb{C} \setminus \Sigma_T$, where $\Sigma_T = \Sigma_U = \Sigma_2 \cup \Sigma_3$.

- $\mathbf{T}_+ = \mathbf{T}_- \mathbf{J}_T$ on $\Sigma_2 \cup \Sigma_3$ with jump matrix (where we put $\varphi_{d+1} \equiv 0$)

$$\mathbf{J}_T = \text{diag}(e^{-2n\varphi_{1,+}}, e^{-2n\varphi_{1,-}}, e^{-2n\varphi_{3,+}}, e^{-2n\varphi_{3,-}}, \dots) + \sum_{j=1}^{\lfloor \frac{d}{2} \rfloor} E_{2j-1,2j} \quad \text{on } \Sigma_1^*, \quad (7.33)$$

$$\mathbf{J}_T = \text{diag}(1, 1, e^{-2n\varphi_{3,+}}, e^{-2n\varphi_{3,-}}, \dots) + e^{2n\varphi_1} E_{1,2} + \sum_{j=2}^{\lfloor \frac{d}{2} \rfloor} E_{2j-1,2j} \quad \text{on } \Sigma_1 \setminus \Sigma_1^*, \quad (7.34)$$

$$\mathbf{J}_T = \text{diag}(1, 1, e^{-2n\varphi_{3,+}}, e^{-2n\varphi_{3,-}}, \dots) + \sum_{j=2}^{\lfloor \frac{d}{2} \rfloor} E_{2j-1,2j} \quad \text{on } \Sigma_3 \setminus \Sigma_1, \quad (7.35)$$

$$\mathbf{J}_T = \text{diag}(1, \omega^{-1} e^{-2n\varphi_{2,+}}, \omega e^{-2n\varphi_{2,-}}, \omega^{-2} e^{-2n\varphi_{4,+}}, \dots) + \sum_{j=1}^{\lfloor \frac{d}{2} \rfloor} E_{2j,2j+1} \quad \text{on } \Sigma_2. \quad (7.36)$$

- As $z \rightarrow \infty$

$$\mathbf{T}(z) = \left(I_{d+1} + O\left(z^{-\frac{2}{d}}\right) \right) \begin{pmatrix} 1 & 0 \\ 0 & \mathbf{A}(z) \end{pmatrix}. \quad (7.37)$$

- \mathbf{T} satisfies the same endpoint condition (4.7) as \mathbf{Y} .

Remark 7.9. The jump matrices are of size $(d+1) \times (d+1)$ and so the pattern on the diagonal in (7.33)–(7.36) continues until we reach size $(d+1) \times (d+1)$. If d is even, then the last diagonal entry in (7.33)–(7.35) is $e^{-2n\varphi_{d+1,+}} \equiv 1$, because of our convention that $\varphi_{d+1} \equiv 0$. If d is odd, then the last diagonal entry in (7.36) is $\omega^{d+1} e^{-2n\varphi_{d+1,+}} \equiv -1$.

Proof. By definition (7.32), the matrix $\mathbf{J}_T(z)$ is given by

$$\mathbf{J}_T(z) = \mathbf{T}_-^{-1}(z) \mathbf{T}_+(z) = L^n \mathbf{G}_-(z)^{-n} \mathbf{J}_U(z) \mathbf{G}_+(z)^n L^{-n},$$

and recall that $\mathbf{G}(z)$ and L are diagonal matrices.

The k th diagonal entry of \mathbf{J}_U is multiplied by

$$e^{n(g_{k-1,+} - g_{k,+})} e^{-n(g_{k-1,-} - g_{k,-})}$$

where we put $g_0 \equiv 0$ and $g_{d+1} \equiv 0$. On Σ_k we have that $e^{ng_{k-1,+}} = e^{ng_{k-1,-}}$ and then it reduces to multiplication by $e^{n(g_{k,-} - g_{k,+})}$. Lemma 7.5 expresses this in terms of $e^{-2n\varphi_{k,+}}$ and $e^{-2n\varphi_{k,-}}$. It all combines nicely with the entries $\lambda_{k,n}$ and $\lambda_{k,n}^{-1}$ and this gives the diagonal entries in the jump matrices (7.33)–(7.36).

The (1, 2) entry in the jump matrix on Σ_1 is multiplied by $e^{n(g_{1,+}(z) + g_{1,-}(z) - g_{2,+}(z) - \ell_1)}$, and so we obtain

$$e^{n(-\frac{1}{i_0} V_1(z) + g_{1,+}(z) + g_{1,-}(z) - g_{2,+}(z) - \ell_1)} = \begin{cases} 1 & \text{on } \Sigma_1^*, \\ e^{2n\varphi_1(z)} & \text{on } \Sigma_1 \setminus \Sigma_1^*, \end{cases}$$

by (7.3) and (7.19). We also use here that n is a multiple of d . This gives the (1, 2) entries in the jumps in (7.33) and (7.34).

The entry $(k, k+1)$ with $k \geq 2$ is multiplied by $e^{n(g_{k,+}(z) + g_{k,-}(z) - g_{k+1,+}(z) - g_{k-1,-}(z))}$. The relations (7.5)–(7.6) show that this is 1 for $z \in \Sigma_k$. The expressions with even k are relevant on

Σ_2 and those with odd k on Σ_3 . All together this implies that $(\mathbf{J}_U)_{k,k+1} = 1$ on Σ_2 if k is even and $(\mathbf{J}_U)_{k,k+1} = 1$ on Σ_3 is $k \geq 3$ is odd. This proves the jumps in the RH problem.

The asymptotic condition in the RH problem for \mathbf{T} follows from (6.7) and the fact that

$$\mathbf{G}(z)^n = \text{diag}(z^{-n}, z^{\frac{n}{d}} I_d) \times (I_{d+1} + O(z^{-\frac{d+1}{d}})) \quad \text{as } z \rightarrow \infty$$

because of the normalizations of the measures $\|\mu_k^*\| = 1 - \frac{k-1}{d}$, and

$$\begin{aligned} g_1(z) &= \log z + O(z^{-d-1}) \\ g_k(z) &= \|\mu_k^*\| \log z + O(z^{-\frac{d+1}{d}}) \quad \text{for } k \geq 2 \end{aligned}$$

as $z \rightarrow \infty$. □

8 Fourth transformation $\mathbf{T} \mapsto \mathbf{S}$

In the fourth transformation of the RH steepest descent analysis we construct lenses around the stars Σ_1^* , Σ_2 and Σ_3 . Before describing this construction we analyze certain properties of the φ -functions.

Recall that φ_1 is an analytic function on $\mathbb{C} \setminus (\Sigma_1^* \cup \Sigma_2)$. As a consequence of (7.15)–(7.16) we know that $\text{Re } \varphi_{1,\pm} = 0$ on Σ_1^* . Moreover, since $\varphi_{2,\pm}$ is purely imaginary on Σ_2 (cf. (7.17)–(7.18)), it follows from (7.29) that $\text{Re } \varphi_{1,+} = \text{Re } \varphi_{1,-}$ on Σ_2 . Therefore, the harmonic function $\text{Re } \varphi_1$ on $\mathbb{C} \setminus (\Sigma_1^* \cup \Sigma_2)$ can be extended continuously to the whole complex plane.

Consider now the set

$$D := \{z \in \mathbb{C} : \text{Re } \varphi_1(z) < 0\}.$$

By the maximum principle for harmonic functions, the boundary ∂D of D consists of a finite union of analytic arcs that start and end on $\Sigma_1^* \cup \Sigma_2$ or at infinity. Since $\varphi_{1,\pm}(z) = \pm \pi i \mu_1^*([z, \omega^\ell x^*])$ for $z \in [0, \omega^\ell x^*] \subset \Sigma_1^*$, $\ell = 0, \dots, d$, the Cauchy-Riemann equations allow us to conclude that for every $z \in [0, \omega^\ell x^*]$, there exists a neighborhood U of z such that $\text{Re } \varphi_1 > 0$ on $U \setminus [0, \omega^\ell x^*]$. Here we use that the density of μ_1^* is strictly positive at every point $z \in [0, \omega^\ell x^*]$. Hence the closed set $\mathbb{C} \setminus D$ contains in its interior the segments $[0, \omega^\ell x^*]$.

Recall that $-i(\xi_2 - \xi_1)_+ > 0$ on $[0, x^*]$, since this is the density of μ_1^* up to a positive constant, see (3.25) with $\xi_{1,-} = \xi_{2,+}$ on Σ_1^* . It also vanishes as a square root at x^* . Then it follows that $\xi_2(x) - \xi_1(x) > 0$ for $x \in (x^*, \hat{x}]$ provided we take \hat{x} sufficiently close to x^* . We deduce from (7.12) that $(x^*, \hat{x}] \subset D$ and $x^* \in \partial D$. Note also that

$$\varphi_1(z) = -c(z - x^*)^{3/2} + O((z - x^*)^{5/3}), \quad \text{as } z \rightarrow x^*,$$

where $c > 0$, which implies that ∂D makes angles $\pm \pi/3$ with $[0, x^*]$ at x^* . By symmetry, the same properties hold for the other segments $(\omega^\ell x^*, \omega^\ell \hat{x}]$ and $[0, \omega^\ell x^*]$.

On the other hand, observe that

$$2\varphi_1(z) = \frac{t_{d+1}}{(d+1)t_0} z^{d+1} + O(z^{\frac{d+1}{d}}), \quad z \rightarrow \infty, \quad (8.1)$$

which follows from (7.19)–(7.20). From (8.1) it is easy to deduce that the analytic arcs of ∂D that start at the points $\omega^\ell x^*$ cannot end at infinity and therefore necessarily end at points on Σ_2 .

As a consequence of (7.17)–(7.18) we know that for $z \in \Sigma_k$,

$$i\varphi_{k,+}(z) = -i\varphi_{k,-}(z) = \pi \mu_k^*([0, z]) + \frac{d \sin(\frac{(k-1)\pi}{d})}{(d+1)t_0 t_{d+1}^{1/d}} |z|^{\frac{d+1}{d}} - \frac{\pi}{(d+1)} \|\mu_k^*\|,$$

hence $\pm i\varphi_{k,\pm}$ is strictly increasing along each ray of Σ_k . Again from the Cauchy-Riemann equations it follows that $\operatorname{Re} \varphi_k > 0$ on both sides of Σ_k .

We now describe the construction of the lenses around the stars Σ_1^* , Σ_2 and Σ_3 . They are shown in Figure 9 in the case $d = 3$. We use

$$L_1 = L_1^+ \cup L_1^- \cup \Sigma_1^*$$

to denote the lens around Σ_1^* , where L_1^+ (L_1^-) is the part of L_1 that lies on the $+$ side ($-$ side) of Σ_1^* . The boundary ∂L_1 of L_1 intersects Σ_3 at the points $\omega^\ell x^*$, $\ell = 0, \dots, d$ and intersects Σ_2 at points at a positive distance $\delta_1 > 0$ from the origin. By the properties of the φ -functions discussed above, it is possible to take L_1 so that $\partial L_1 \setminus \{\omega^\ell x^*\}_{\ell=0}^d$ is contained in the regions where $\operatorname{Re} \varphi_k > 0$, for every odd $k \geq 1$.

Analogously we construct the lens

$$L_2 = L_2^+ \cup L_2^- \cup \Sigma_2$$

around Σ_2 . It is chosen so that ∂L_2 lies in the regions where $\operatorname{Re} \varphi_k > 0$, for every even $k \geq 2$. The boundary ∂L_2 consists of $2(d+1)$ rays that meet Σ_1 at points at a positive distance $\delta_2 > 0$ from the origin, and have asymptotic angles $\frac{(2\ell+1)\pi}{d+1} \pm \varepsilon$, for some small $\varepsilon > 0$.

Finally, we construct a similar lens

$$L_3 = L_3^+ \cup L_3^- \cup \Sigma_3$$

around Σ_3 . We choose it such that $L_1 \subset L_3$, and the boundary ∂L_3 consists of $\partial L_1 \cap L_2$ and infinite rays that emanate from the intersection points of ∂L_1 and ∂L_2 , see Figure 9. These rays are chosen so that they are contained in the regions where $\operatorname{Re} \varphi_k > 0$, for every odd $k \geq 3$, and with asymptotic angles $\frac{2\pi\ell}{d+1} \pm \varepsilon$.

Definition 8.1. Let $\mathbf{S}(z)$ be the matrix-valued function defined as follows. We set $\mathbf{S}(z) = \mathbf{T}(z)$ for z outside the lenses L_2 and L_3 , and

$$\mathbf{S} = \mathbf{T} \left(I_{d+1} \mp \sum_{j=1}^{\lceil \frac{d}{2} \rceil} e^{-2n\varphi_{2j-1}} E_{2j,2j-1} \right) \quad \text{in } L_1^\pm \setminus L_2, \quad (8.2)$$

$$\mathbf{S} = \mathbf{T} \left(I_{d+1} \mp \sum_{j=2}^{\lceil \frac{d}{2} \rceil} e^{-2n\varphi_{2j-1}} E_{2j,2j-1} \right) \quad \text{in } L_3^\pm \setminus L_1, \quad (8.3)$$

$$\mathbf{S} = \mathbf{T} \left(I_{d+1} \mp \sum_{j=1}^{\lfloor \frac{d}{2} \rfloor} \omega^{\mp j} e^{-2n\varphi_{2j}} E_{2j+1,2j} \right) \quad \text{in } L_2^\pm \setminus L_1. \quad (8.4)$$

$$\mathbf{S} = \mathbf{T} \left(I_{d+1} + \sum_{j=2}^{d+1} \sum_{k=1}^{j-1} c_{j,k}^\pm e^{-2n(\varphi_k + \dots + \varphi_{j-1})} E_{j,k} \right) \quad \text{in } L_1^\pm \cap L_2^\mp. \quad (8.5)$$

The definition of \mathbf{S} in the intersection of the lenses, see (8.5), involves certain constants $c_{j,k}^\pm$ with $1 \leq k < j \leq d+1$, that are not yet defined. We collect these numbers in two lower triangular matrices

$$C^\pm = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ c_{2,1}^\pm & 1 & 0 & \cdots & 0 \\ c_{3,1}^\pm & c_{3,2}^\pm & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ c_{d+1,1}^\pm & c_{d+1,2}^\pm & c_{d+1,3}^\pm & \cdots & 1 \end{pmatrix}. \quad (8.6)$$




Figure 9: The lenses L_1 , L_2 and L_3 around the stars Σ_1^* , Σ_2 and Σ_3 , in the case $d = 3$.

Recall that σ_2 is the Pauli matrix $\sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$ so that

$$i\sigma_2 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

This matrix will appear on the diagonal in the block diagonal matrices (8.7)-(8.8).

Lemma 8.2. *There exist two lower triangular matrices C^+ and C^- of the form (8.6) such that*

$$\left(I_{d+1} + \sum_{j=1}^{\lceil \frac{d}{2} \rceil} E_{2j-1, 2j} \right) C^+ = C^- \times \begin{cases} \text{diag}(i\sigma_2, \dots, i\sigma_2) & \text{if } d \text{ is odd,} \\ \text{diag}(i\sigma_2, \dots, i\sigma_2, 1) & \text{if } d \text{ is even,} \end{cases} \quad (8.7)$$

and

$$\left(\text{diag}(1, \omega^{-1}, \omega, \omega^{-2}, \dots) + \sum_{j=1}^{\lfloor \frac{d}{2} \rfloor} E_{2j, 2j+1} \right) C^- = C^+ \times \begin{cases} \text{diag}(1, i\sigma_2, \dots, i\sigma_2, -1) & \text{if } d \text{ is odd,} \\ \text{diag}(1, i\sigma_2, \dots, i\sigma_2) & \text{if } d \text{ is even.} \end{cases} \quad (8.8)$$

Proof. In order to define the matrices C^\pm we use the elementary symmetric polynomials $e_{k,n}(x_1, \dots, x_n)$ as in (5.23)–(5.24) and we put

$$e_{k,n} = e_{k,n}(1, \omega^{-1}, \omega, \omega^{-2}, \omega^2, \omega^{-3}, \dots), \quad (8.9)$$

$$e'_{k,n} = e_{k,n}(1, \omega, \omega^{-1}, \omega^2, \omega^{-2}, \omega^3, \dots). \quad (8.10)$$

Recall that n denotes the number of variables appearing in the symmetric function.

Then we construct lower triangular matrices C^\pm as in (8.6) with entries

$$\begin{aligned}
c_{j+1,j-2k}^+ &= (-1)^{j-k} e'_{j-k,j} && \text{for } j = 1, \dots, d \text{ and } k = 0, 1, \dots, \lfloor \frac{j-1}{2} \rfloor, \\
c_{j+1,j+1-2k}^+ &= (-1)^k e'_{k,j} && \text{for } j = 1, \dots, d \text{ and } k = 0, 1, \dots, \lfloor \frac{j}{2} \rfloor, \\
c_{2j+1,2j+1-2k}^- &= (-1)^k e_{k,2j} && \text{for } j = 1, \dots, \lfloor \frac{d}{2} \rfloor \text{ and } k = 1, \dots, j, \\
c_{2j+1,2j-2k}^- &= (-1)^{k+1} e_{2j-k,2j} && \text{for } j = 1, \dots, \lfloor \frac{d}{2} \rfloor \text{ and } k = 0, 1, \dots, j-1, \\
c_{2j,2j-1-2k}^- &= (-1)^k e_{k,2j-1} && \text{for } j = 1, \dots, \lfloor \frac{d}{2} \rfloor \text{ and } k = 0, 1, \dots, j-1, \\
c_{2j,2j-2k}^- &= (-1)^k e_{2j-1-k,2j-1} && \text{for } j = 1, \dots, \lfloor \frac{d}{2} \rfloor \text{ and } k = 1, 2, \dots, j-1.
\end{aligned}$$

For example, for $d = 5$ we have

$$C^+ = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ -e'_{1,1} & 1 & 0 & 0 & 0 & 0 \\ -e'_{1,2} & e'_{2,2} & 1 & 0 & 0 & 0 \\ e'_{2,3} & -e'_{1,3} & -e'_{3,3} & 1 & 0 & 0 \\ e'_{2,4} & -e'_{3,4} & -e'_{1,4} & e'_{4,4} & 1 & 0 \\ -e'_{3,5} & e'_{2,5} & e'_{4,5} & -e'_{1,5} & -e'_{5,5} & 1 \end{pmatrix}$$

$$C^- = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ e_{0,1} & 1 & 0 & 0 & 0 & 0 \\ -e_{1,2} & -e_{2,2} & 1 & 0 & 0 & 0 \\ -e_{1,3} & -e_{2,3} & e_{0,3} & 1 & 0 & 0 \\ e_{2,4} & e_{3,4} & -e_{1,4} & -e_{4,4} & 1 & 0 \\ e_{2,5} & e_{3,5} & -e_{1,5} & -e_{4,5} & e_{0,5} & 1 \end{pmatrix}.$$

Then it is a tedious task to verify that the identities (8.7) and (8.8) are indeed satisfied. The verification depends on the following properties of the quantities $e_{k,n}$ and $e'_{k,n}$ defined in (8.9)–(8.10):

- (i) If n is odd then $e_{k,n} = e'_{k,n}$ and $e_{n-k,n} = e_{k,n}$.
- (ii) If n is even then $e_{k,n} = \omega^{-n/2} e'_{n-k,n}$ and $e'_{k,n} = \omega^{n/2} e_{n-k,n}$.
- (iii) For every n ,

$$e_{k,n+1} = e_{k,n} + \begin{cases} \omega^{n/2} e_{k-1,n} & \text{if } n \text{ is even,} \\ \omega^{-(n+1)/2} e_{k-1,n} & \text{if } n \text{ is odd.} \end{cases} \quad (8.11)$$

These identities are easy consequences of the definitions (8.9) and (8.10). For example, the identity in (8.11) relies on the fact that the elementary symmetric polynomial $e_{k,n+1}(x_1, \dots, x_{n+1})$ on $n+1$ variables is the sum of $e_{k,n}(x_1, \dots, x_n)$ which has the terms without x_{n+1} , and $x_{n+1} e_{k-1,n}(x_1, \dots, x_n)$ which has the terms containing x_{n+1} .

Now we are going to verify the identity (8.7). The identity (8.7) for the entries in rows $2j-1, 2j$ and columns $2k-1, 2k$ (where $1 \leq k \leq j \leq \lfloor \frac{d+1}{2} \rfloor$) is exactly

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} c_{2j-1,2k-1}^+ & c_{2j-1,2k}^+ \\ c_{2j,2k-1}^+ & c_{2j,2k}^+ \end{pmatrix} = \begin{pmatrix} c_{2j-1,2k-1}^- & c_{2j-1,2k}^- \\ c_{2j,2k-1}^- & c_{2j,2k}^- \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix},$$

which when written in terms of the elementary symmetric polynomials is equivalent to

$$(-1)^{j+k} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} e'_{j-k,2j-2} & -e'_{j+k-1,2j-2} \\ -e'_{j+k-1,2j-1} & e'_{j-k,2j-1} \end{pmatrix} = (-1)^{j+k} \begin{pmatrix} e_{j-k,2j-2} & e_{j+k-1,2j-2} \\ e_{j-k,2j-1} & e_{j+k-1,2j-1} \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \quad (8.12)$$

The equality for the entries in the second row of (8.12) follows from item (i) above, while the equality for the entries in the first row relies on all three items. For example, the equality for the first entry follows from (where $n = 2j - 2$ is even)

$$\begin{aligned} e'_{j-k, 2j-2} - e'_{j+k-1, 2j-1} &= e'_{j-k, n} - e'_{j+k-1, n+1} \\ &= \omega^{n/2} e_{j+k-2, n} - e_{j+k-1, n+1} && \text{by (ii) and (i)} \\ &= -e_{j+k-1, n} && \text{by (iii)}. \end{aligned}$$

This argument establishes (8.7) completely if d is odd. If d is even, then the last row needs separate verification. The equality follows from

$$-e'_{d+1-k, d} = e_{k, d}, \quad \text{for } k = 1, \dots, d \text{ and } d \text{ is even.} \quad (8.13)$$

To see that (8.13) indeed holds we calculate

$$\begin{aligned} e'_{d+1-k, d} + e_{k, d} &= \omega^{d/2} e_{k-1, d} + e_{k, d} && \text{by item (ii)} \\ &= e_{k, d+1} && \text{by (iii)}. \end{aligned}$$

Now we recall that ω is a $(d+1)$ -st root of unity, and by the definition (8.9) of $e_{k, d+1}$,

$$e_{k, d+1} = e_{k, d+1}(1, \omega, \omega^2, \dots, \omega^d) = 0, \quad k = 1, \dots, d.$$

Thus (8.13) holds, and this gives the identity of the last row of (8.7) in case d is even.

The proof of (8.8) follows along similar lines. \square

We return to the matrix-valued function \mathbf{S} defined in (8.2)–(8.5). The coefficients $c_{j, k}^{\pm}$ are taken from the matrices (8.6) whose existence is proved in Lemma 8.2.

RH problem 8.3. The matrix $\mathbf{S}(z)$ is the solution to the following RH problem:

- \mathbf{S} is analytic in $\mathbb{C} \setminus \Sigma_S$, where $\Sigma_S = \Sigma_2 \cup \Sigma_3 \cup \bigcup_{j=1}^3 \partial L_j$.
- $\mathbf{S}_+ = \mathbf{S}_- \mathbf{J}_S$ on Σ_S with jump matrix

$$\mathbf{J}_S = \begin{cases} \text{diag}(i\sigma_2, i\sigma_2, \dots, i\sigma_2, 1) & \text{if } d \text{ is even} \\ \text{diag}(i\sigma_2, i\sigma_2, \dots, i\sigma_2) & \text{if } d \text{ is odd} \end{cases} \quad \text{on } \Sigma_1^*, \quad (8.14)$$

$$\mathbf{J}_S = e^{2n\varphi_1} E_{1,2} + \begin{cases} \text{diag}(1, 1, i\sigma_2, \dots, i\sigma_2, 1) & \text{if } d \text{ is even} \\ \text{diag}(1, 1, i\sigma_2, \dots, i\sigma_2) & \text{if } d \text{ is odd} \end{cases} \quad \text{on } \Sigma_1 \setminus \Sigma_1^*, \quad (8.15)$$

$$\mathbf{J}_S = \begin{cases} \text{diag}(1, 1, i\sigma_2, \dots, i\sigma_2, 1) & \text{if } d \text{ is even} \\ \text{diag}(1, 1, i\sigma_2, \dots, i\sigma_2) & \text{if } d \text{ is odd} \end{cases} \quad \text{on } \Sigma_3 \setminus \Sigma_1, \quad (8.16)$$

$$\mathbf{J}_S = \begin{cases} \text{diag}(1, i\sigma_2, \dots, i\sigma_2) & \text{if } d \text{ is even} \\ \text{diag}(1, i\sigma_2, \dots, i\sigma_2, -1) & \text{if } d \text{ is odd} \end{cases} \quad \text{on } \Sigma_2, \quad (8.17)$$

and on the lips of the lenses,

$$\mathbf{J}_S = I_{d+1} + e^{-2n\varphi_1} E_{2,1} \quad \text{on } \partial L_1 \setminus L_2 \quad (8.18)$$

$$\mathbf{J}_S = I_{d+1} + \sum_{j=2}^{\lfloor \frac{d}{2} \rfloor} e^{-2n\varphi_{2j-1}} E_{2j,2j-1} \quad \text{on } \partial L_3 \setminus L_2 \quad (8.19)$$

$$\mathbf{J}_S = I_{d+1} + \sum_{j=1}^{\lfloor \frac{d}{2} \rfloor} \omega^{\mp j} e^{-2n\varphi_{2j}} E_{2j+1,2j} \quad \text{on } \partial L_2^\pm \setminus L_1 \quad (8.20)$$

$$\mathbf{J}_S = I_{d+1} + \sum_{j=2}^{d+1} \sum_{k=1}^{j-1} b_{j,k} e^{-2n(\varphi_k + \dots + \varphi_{j-1})} E_{j,k} \quad \begin{array}{l} \text{on } \partial L_1 \cap L_2 \\ \text{and on } \partial L_2 \cap L_1, \end{array} \quad (8.21)$$

for certain constants $b_{j,k}$ (different constants on $\partial L_1 \cap L_2$ and $\partial L_2 \cap L_1$) whose exact values are not important to us in what follows.

- As $z \rightarrow \infty$ in the sector S_ℓ^\pm , $\ell = 0, \dots, d$,

$$\mathbf{S}(z) = \left(I_{d+1} + O\left(z^{-\frac{2}{d}}\right) \right) \begin{pmatrix} 1 & 0 \\ 0 & \mathbf{A}(z) \end{pmatrix}. \quad (8.22)$$

Proof. The jump matrices (7.33)–(7.35) in the RH problem for \mathbf{T} have a block diagonal form with 2×2 blocks

$$\begin{pmatrix} e^{-2n\varphi_{k,+}} & 1 \\ 0 & e^{-2n\varphi_{k,-}} \end{pmatrix}.$$

The 2×2 block has the standard factorization

$$\begin{pmatrix} 1 & 0 \\ e^{-2n\varphi_{k,-}} & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ e^{-2n\varphi_{k,+}} & 1 \end{pmatrix},$$

since $\varphi_{k,+} = -\varphi_{k,-}$ on Σ_k , see e.g. (7.15)–(7.18). The definition of S in $L_3 \setminus L_2$, see (8.2)–(8.3), has the effect of moving the outer two factors to the jump matrix on the lips of the lenses while keeping the inner factor on the original contour. This accounts for the jump matrices (8.18)–(8.19) on the lips of the lenses, and for the jump matrices (8.14)–(8.16) on the contour Σ_3 outside of the lens L_2 . The jump matrix (7.36) has 2×2 blocks

$$\begin{pmatrix} \omega^{-k} e^{-2n\varphi_{2k,+}} & 1 \\ 0 & \omega^k e^{-2n\varphi_{2k,-}} \end{pmatrix}$$

which has the factorization

$$\begin{pmatrix} 1 & 0 \\ \omega^k e^{-2n\varphi_{2k,-}} & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ \omega^{-k} e^{-2n\varphi_{2k,+}} & 1 \end{pmatrix}.$$

Because of the definition (8.4) of S in $L_2 \setminus L_1$, we obtain (8.17) and (8.20).

The form of the jump matrix (8.21) is immediate from the definitions.

It remains to check the jumps on the parts of Σ_1^* and Σ_2 that lie in the intersection of the lenses. To that end we write

$$\psi_k = -\sum_{j=1}^{k-1} \varphi_j + \sum_{j=k}^d \varphi_j, \quad k = 1, \dots, d+1, \quad (8.23)$$

where the first sum is 0 if $k = 1$ and the last sum is 0 if $k = d + 1$. Also put

$$\Psi = \text{diag}(e^{\psi_1}, e^{\psi_2}, \dots, e^{\psi_{d+1}}).$$

Then (8.5) can be written as

$$\mathbf{S} = \mathbf{T}\Psi^n C^\pm \Psi^{-n}, \quad \text{in } L_1^\pm \cap L_2^\mp.$$

The jump matrix on $\Sigma_1^* \cap L_1 \cap L_2$ therefore is

$$\mathbf{J}_S = \Psi_-^n (C^-)^{-1} \Psi_-^{-n} \mathbf{J}_T \Psi_+^n C^+ \Psi_+^{-n}. \quad (8.24)$$

Recall that $\varphi_{2k-1,+} = -\varphi_{2k-1,-}$ and $\varphi_{2k,+} = \varphi_{2k-1,-} + \varphi_{2k,-} + \varphi_{2k+1,-}$ on Σ_1^* , see (7.15)–(7.18) and (7.28). Then if d is odd, we can check from the definition (8.23) of ψ_k that

$$\begin{aligned} \psi_{2k,+} &= \psi_{2k,-} + \varphi_{1,-} + 2\varphi_{2k-1,-} - \varphi_{d,-} \\ \psi_{2k-1,+} &= \psi_{2k-1,-} + \varphi_{1,-} - 2\varphi_{2k-1,-} - \varphi_{d,-} \\ \psi_{2k-1,+} &= \psi_{2k,-} + \varphi_{1,-} - \varphi_{d,-} \\ \psi_{2k,+} &= \psi_{2k-1,-} + \varphi_{1,-} - \varphi_{d,-}. \end{aligned} \quad (8.25)$$

It follows from these relations and (7.33) that

$$\Psi_-^{-n} \mathbf{J}_T \Psi_+^n = e^{n(\varphi_{1,-} - \varphi_{d,-})} \left(I_{d+1} + \sum_{j=1}^{\lfloor \frac{d}{2} \rfloor} E_{2j-1,2j} \right) \quad (8.26)$$

which is a scalar function times a constant matrix. Then by one of the defining properties (8.7) of C^\pm , we obtain from (8.24) and (8.26)

$$\mathbf{J}_S = e^{n(\varphi_{1,-} - \varphi_{d,-})} \Psi_-^n \text{diag}(i\sigma_2, \dots, i\sigma_2) \Psi_+^{-n}. \quad (8.27)$$

Again using the jump properties (8.25) of ψ_k we then see that (8.27) reduces to (8.14). The argument for d even is similar. The main difference is that the term $-\varphi_{d,-}$ in (8.25) is absent.

The jump matrix \mathbf{J}_S on $\Sigma_2 \cap L_1 \cap L_2$ is calculated in a similar way. Now we use the other property (8.8) that defines the matrices C^\pm .

The asymptotic condition (8.22) is obvious from (7.37) outside the lenses, since $\mathbf{S} = \mathbf{T}$ outside the lenses. Furthermore, it is easy to deduce from (7.21) that the φ functions satisfy

$$\begin{aligned} \varphi_{2k}(z) &= d_{2k,\ell,\pm} z^{\frac{d+1}{d}} + O(\log z), \quad z \in L_2 \cap S_\ell^\pm, \quad k = 1, \dots, \lfloor \frac{d}{2} \rfloor, \\ \varphi_{2k-1}(z) &= d_{2k-1,\ell,\pm} z^{\frac{d+1}{d}} + O(\log z), \quad z \in L_3 \cap S_\ell^\pm, \quad k = 2, \dots, \lfloor \frac{d}{2} \rfloor, \end{aligned} \quad (8.28)$$

as $z \rightarrow \infty$, where $d_{k,\ell,\pm}$ are constants such that $\text{Re } \varphi_{2k}(z) > 0$ ($\text{Re } \varphi_{2k+1}(z) > 0$) for z large enough in L_2 (L_3). Therefore, from (7.37) and the definition (8.3)–(8.4) of \mathbf{S} in the unbounded lenses, we deduce that the asymptotic condition is also valid inside the lenses L_2 and L_3 (although not uniformly up to the sets Σ_2 and Σ_3). \square

Remark 8.4. Note that the jump matrices (8.14), (8.16) and (8.17) are constant. The jump matrix (8.15) is not constant since it contains the term $e^{2n\varphi_1} E_{1,2}$. However, by the discussion at the beginning of this section we have

$$\text{Re } \varphi_1 < 0 \quad \text{on } \Sigma_1 \setminus \Sigma_1^* \quad (8.29)$$

so that the non-constant term is exponentially small if n is large.

The jump matrices (8.18)–(8.21) on the lips of the lenses are exponentially close to the identity matrix, since the lenses were taken so that

$$\operatorname{Re} \varphi_1 > 0 \quad \text{on } \partial L_1 \quad (8.30)$$

$$\operatorname{Re} \varphi_{2k-1} > 0 \quad \text{on } \partial L_3 \quad \text{for } k = 2, \dots, \lfloor \frac{d}{2} \rfloor \quad (8.31)$$

$$\operatorname{Re} \varphi_{2k} > 0 \quad \text{on } \partial L_2 \quad \text{for } k = 1, \dots, \lfloor \frac{d}{2} \rfloor \quad (8.32)$$

$$\operatorname{Re} \varphi_k \geq 0 \quad \text{on } L_1 \cap L_2 \quad \text{for all } k. \quad (8.33)$$

Also taking note of (8.28) we may then conclude that

$$\mathbf{J}_S(z) = I_{d+1} + O\left(e^{-cn|z|^{\frac{d+1}{d}}}\right), \quad z \in (\Sigma_1 \setminus \Sigma_1^*) \cup \partial L_1 \cup \partial L_2 \cup \partial L_3, \quad (8.34)$$

as $n \rightarrow \infty$, and the O -term is uniform for z if we stay away from the endpoints $\omega^\ell x^*$ of Σ_1^* .

Ignoring all exponentially small entries in the jump matrices, we arrive at a new RH problem which we study next.

9 Global parametrix

RH problem 9.1. Find $\mathbf{M} : \mathbb{C} \setminus \Sigma_M \rightarrow \mathbb{C}^{(d+1) \times (d+1)}$ where $\Sigma_M = \Sigma_2 \cup \Sigma_3$ such that:

- \mathbf{M} is analytic in $\mathbb{C} \setminus \Sigma_M$.
- $\mathbf{M}_+ = \mathbf{M}_- \mathbf{J}_M$ on Σ_M , where

$$\mathbf{J}_M = \begin{cases} \operatorname{diag}(i\sigma_2, i\sigma_2, \dots, i\sigma_2, 1) & \text{for } d \text{ even} \\ \operatorname{diag}(i\sigma_2, i\sigma_2, \dots, i\sigma_2) & \text{for } d \text{ odd} \end{cases} \quad \text{on } \Sigma_1^*, \quad (9.1)$$

$$\mathbf{J}_M = \begin{cases} \operatorname{diag}(1, 1, i\sigma_2, \dots, i\sigma_2, 1) & \text{for } d \text{ even} \\ \operatorname{diag}(1, 1, i\sigma_2, \dots, i\sigma_2) & \text{for } d \text{ odd} \end{cases} \quad \text{on } \Sigma_3 \setminus \Sigma_1^*, \quad (9.2)$$

$$\mathbf{J}_M = \begin{cases} \operatorname{diag}(1, i\sigma_2, i\sigma_2, \dots, i\sigma_2) & \text{for } d \text{ even} \\ \operatorname{diag}(1, i\sigma_2, i\sigma_2, \dots, i\sigma_2, -1) & \text{for } d \text{ odd} \end{cases} \quad \text{on } \Sigma_2. \quad (9.3)$$

- \mathbf{M} satisfies the asymptotic condition

$$\mathbf{M}(z) = \left(I_{d+1} + O\left(\frac{1}{z}\right)\right) \begin{pmatrix} 1 & 0 \\ 0 & \mathbf{A}(z) \end{pmatrix} \quad (9.4)$$

as $z \rightarrow \infty$ in the sector S_ℓ^\pm , $\ell = 0, \dots, d$.

- $\mathbf{M}(z) = O((z - \omega^j x^*)^{-1/4})$ as $z \rightarrow \omega^j x^*$, $j = 0, \dots, d$.
- $\mathbf{M}(z)$ remains bounded as $z \rightarrow 0$.

We solve this problem with the help of a meromorphic differential and certain meromorphic functions defined on the Riemann surface \mathcal{R} given in (3.5). Let $\boldsymbol{\eta}$ be the meromorphic differential

that has simple poles at the branch points $\omega^j x^*$, $j = 0, \dots, d$, and a simple pole at ∞_2 (see definition of \mathcal{R}), with

$$\operatorname{Res}(\boldsymbol{\eta}, \omega^j x^*) = -\frac{1}{2}, \quad \operatorname{Res}(\boldsymbol{\eta}, \infty_2) = \frac{d+1}{2}, \quad (9.5)$$

and is holomorphic elsewhere. This meromorphic differential exists and is uniquely determined by these conditions, since \mathcal{R} has genus zero.

We now define

$$u_j(z) = \int_{\infty_1}^z \boldsymbol{\eta}, \quad z \in \mathcal{R}_j, \quad j = 1, \dots, d+1,$$

where ∞_1 is the point at infinity in \mathcal{R}_1 and the path of integration is taken so that it satisfies the following rules:

- 1) The path for $u_1(z)$ stays on the first sheet.
- 2) The path for $u_2(z)$ starts on the first sheet, passes once through Σ_1 and stays on the second sheet. The passage is made via the --side on \mathcal{R}_1 .
- 3) The definition of the path for $u_j(z)$, $3 \leq j \leq d$, is based on the following observation. Suppose that $z \in \mathcal{R}_j$, $j = 3, \dots, d$, and z lies in any of the $2d+2$ sectors S_ℓ^\pm . Then there is only one neighboring sector in the upper sheet \mathcal{R}_{j-1} to which the point z can be connected through a path that crosses the cut connecting the sheets \mathcal{R}_{j-1} and \mathcal{R}_j only once.

We then require that the path for $u_j(z)$, $3 \leq j \leq d$, in its passage from \mathcal{R}_1 to \mathcal{R}_2 , crosses Σ_1 only once through the --side in the upper sheet, and for each $2 \leq k \leq j-1$, the path should cross the cut Σ_k connecting the sheets \mathcal{R}_k and \mathcal{R}_{k+1} only once. The observation above shows that this path is well-defined and clearly any two paths satisfying these requirements will give the same value for $u_j(z)$.

- 4) The path for $u_{d+1}(z)$ is defined as follows. As before, it is required to cross Σ_1 only once through the --side in its passage from \mathcal{R}_1 to \mathcal{R}_2 , and for $2 \leq k \leq d-1$, it is required to cross the cut Σ_k only once in its passage from \mathcal{R}_k to \mathcal{R}_{k+1} . Finally, the path from \mathcal{R}_d to \mathcal{R}_{d+1} should cross Σ_d only once through the --side on the upper sheet.

These rules should be satisfied for any value of d , even or odd. It easily follows from these rules and the residue assumptions on the branch points $\omega^j x^*$ that the following relations hold in the case d odd:

$$u_{1,-} = u_{2,+}, \quad \text{on } \Sigma_1^*, \quad (9.6)$$

$$u_{1,+} = u_{2,-} \pm \pi i, \quad \text{on } \Sigma_1^*, \quad (9.7)$$

$$u_{2j-1,\pm} = u_{2j,\mp}, \quad 2 \leq j \leq \frac{d-1}{2}, \quad \text{on } \Sigma_3, \quad (9.8)$$

$$u_{d+1,+} = u_{d,-}, \quad \text{on } \Sigma_3, \quad (9.9)$$

$$u_{d+1,-} = u_{d,+} \pm \pi i, \quad \text{on } \Sigma_3, \quad (9.10)$$

$$u_{j,+} = u_{j,-}, \quad j = 1, d+1, \quad \text{on } \Sigma_2, \quad (9.11)$$

$$u_{2j,\pm} = u_{2j+1,\mp}, \quad 1 \leq j \leq \frac{d-1}{2}, \quad \text{on } \Sigma_2, \quad (9.12)$$

and in the case d even, the relations (9.6)–(9.7) still hold, and we now have

$$u_{2j-1,\pm} = u_{2j,\mp}, \quad 2 \leq j \leq \frac{d}{2}, \quad \text{on } \Sigma_3, \quad (9.13)$$

$$u_{d+1,+} = u_{d+1,-}, \quad \text{on } \Sigma_3, \quad (9.14)$$

$$u_{1,+} = u_{1,-}, \quad \text{on } \Sigma_2, \quad (9.15)$$

$$u_{2j,\pm} = u_{2j+1,\mp}, \quad 1 \leq j \leq \frac{d}{2} - 1 \quad \text{on } \Sigma_2, \quad (9.16)$$

$$u_{d+1,+} = u_{d,-}, \quad \text{on } \Sigma_2, \quad (9.17)$$

$$u_{d+1,-} = u_{d,+} \pm \pi i, \quad \text{on } \Sigma_2. \quad (9.18)$$

Let us define now the functions $v_j(z) := e^{u_j(z)}$, $j = 1, \dots, d+1$, and set

$$\mathbf{v}(z) := (v_1(z) \ v_2(z) \ \dots \ v_{d+1}(z)).$$

Then the above relations (9.6)–(9.18) give in the case d is odd,

$$\begin{aligned} \mathbf{v}_+(z) &= \mathbf{v}_-(z) \operatorname{diag}(i\sigma_2, \sigma_1, \dots, \sigma_1, i\sigma_2), & z \in \Sigma_1^*, \\ \mathbf{v}_+(z) &= \mathbf{v}_-(z) \operatorname{diag}(1, 1, \sigma_1, \dots, \sigma_1, i\sigma_2), & z \in \Sigma_3 \setminus \Sigma_1^*, \\ \mathbf{v}_+(z) &= \mathbf{v}_-(z) \operatorname{diag}(1, \sigma_1, \dots, \sigma_1, 1), & z \in \Sigma_2, \end{aligned}$$

and in the case d is even,

$$\begin{aligned} \mathbf{v}_+(z) &= \mathbf{v}_-(z) \operatorname{diag}(i\sigma_2, \sigma_1, \dots, \sigma_1, 1), & z \in \Sigma_1^*, \\ \mathbf{v}_+(z) &= \mathbf{v}_-(z) \operatorname{diag}(1, 1, \sigma_1, \dots, \sigma_1, 1), & z \in \Sigma_3 \setminus \Sigma_1^*, \\ \mathbf{v}_+(z) &= \mathbf{v}_-(z) \operatorname{diag}(1, \sigma_1, \dots, \sigma_1, i\sigma_2), & z \in \Sigma_2. \end{aligned}$$

Here $\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $i\sigma_2 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$.

We next introduce the functions

$$\begin{aligned} \widehat{v}_1 &:= v_1, \\ \widehat{v}_{2j} &:= v_{2j}, & 1 \leq j \leq \lfloor \frac{d}{2} \rfloor, \\ \widehat{v}_{2j+1} &:= \begin{cases} -v_{2j+1} & \text{on } \bigcup_{\ell=0}^d S_\ell^+, \\ v_{2j+1} & \text{on } \bigcup_{\ell=0}^d S_\ell^-, \end{cases} & 1 \leq j \leq \lfloor \frac{d-1}{2} \rfloor, \\ \widehat{v}_{d+1} &:= \begin{cases} v_{d+1} & \text{on } \bigcup_{\ell=0}^d S_\ell^+, \\ -v_{d+1} & \text{on } \bigcup_{\ell=0}^d S_\ell^-, \end{cases} & \text{if } d \text{ is odd,} \\ \widehat{v}_{d+1} &:= v_{d+1} & \text{if } d \text{ is even.} \end{aligned}$$

If we let

$$\widehat{\mathbf{v}}(z) := (\widehat{v}_1(z) \ \widehat{v}_2(z) \ \dots \ \widehat{v}_{d+1}(z))$$

then we readily see that $\widehat{\mathbf{v}}$ is analytic in $\mathbb{C} \setminus (\Sigma_2 \cup \Sigma_3)$, and satisfies

$$\widehat{\mathbf{v}}_+ = \widehat{\mathbf{v}}_- \mathbf{J}_M, \quad \text{on } \Sigma_2 \cup \Sigma_3. \quad (9.19)$$

We now explicitly construct the sought function \mathbf{M} in terms of the functions \widehat{v}_j and certain meromorphic functions on \mathcal{R} . Let

$$f^{(0)} \equiv 1, f^{(1)}, f^{(2)}, \dots, f^{(d)},$$

be a basis of the vector space of all meromorphic functions that have a pole of order at most d at ∞_2 and are holomorphic elsewhere in \mathcal{R} . We choose these functions so that $f^{(j)}$ has a pole of order j at ∞_2 . By $f_i^{(j)}$ we denote the restriction of $f^{(j)}$ to the sheet \mathcal{R}_i . Then we set

$$\mathbf{B} := \begin{pmatrix} 1 & 1 & 1 & \cdots & 1 \\ f_1^{(1)} & f_2^{(1)} & f_3^{(1)} & \cdots & f_{d+1}^{(1)} \\ f_1^{(2)} & f_2^{(2)} & f_3^{(2)} & \cdots & f_{d+1}^{(2)} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ f_1^{(d)} & f_2^{(d)} & f_3^{(d)} & \cdots & f_{d+1}^{(d)} \end{pmatrix} \text{diag} (\widehat{v}_1 \quad \widehat{v}_2 \quad \cdots \quad \widehat{v}_{d+1}). \quad (9.20)$$

In virtue of (9.19) we also obtain

$$\mathbf{B}_+ = \mathbf{B}_- \mathbf{J}_M, \quad \text{on } \Sigma_2 \cup \Sigma_3. \quad (9.21)$$

By comparing (9.2)–(9.3) with (5.39)–(5.40) we see that

$$\mathbf{J}_M = \begin{pmatrix} 1 & 0 \\ 0 & \mathbf{J}_A \end{pmatrix} \quad \text{on } (\Sigma_3 \setminus \Sigma_1^*) \cup \Sigma_2.$$

Thus

$$\begin{pmatrix} 1 & 0 \\ 0 & \mathbf{A}_+(z) \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & \mathbf{A}_-(z) \end{pmatrix} \mathbf{J}_M \quad \text{on } (\Sigma_3 \setminus \Sigma_1^*) \cup \Sigma_2. \quad (9.22)$$

From (9.21) and (9.22) we deduce that the function $\mathbf{B}(z) \begin{pmatrix} 1 & 0 \\ 0 & \mathbf{A}^{-1}(z) \end{pmatrix}$ extends to an analytic function in $\mathbb{C} \setminus \Sigma_1^*$, and therefore has a Laurent expansion at infinity.

We have $\widehat{v}_1(z) = 1 + O(1/z)$ as $z \rightarrow \infty$, and in virtue of (9.5), we have $\widehat{v}_j(z) = O(z^{-\frac{d+1}{2d}})$ as $z \rightarrow \infty$ for $j \geq 2$. As $f^{(j)}$ has a pole of order j at ∞_2 , we obtain $f_1^{(j)}(z) = O(1)$ for all j , and $f_i^{(j)}(z) = O(z^{\frac{j}{d}})$ for $i \geq 2$. Thus

$$\mathbf{B}(z) = O(z^{\frac{d-1}{2d}})$$

Also $\mathbf{A}^{-1}(z) = O(z^{\frac{d-1}{2d}})$ as $z \rightarrow \infty$, so that

$$\mathbf{B}(z) \begin{pmatrix} 1 & 0 \\ 0 & \mathbf{A}^{-1}(z) \end{pmatrix} = O(z^{\frac{d-1}{d}}) \quad \text{as } z \rightarrow \infty.$$

Since $\frac{d-1}{d} < 1$ we see that the Laurent expansion does not have any terms with strictly positive powers of z . Hence

$$\mathbf{B}(z) \begin{pmatrix} 1 & 0 \\ 0 & \mathbf{A}^{-1}(z) \end{pmatrix} = C + O\left(\frac{1}{z}\right), \quad z \rightarrow \infty, \quad (9.23)$$

for some constant matrix C .

The matrix C is easily seen to be invertible, as the functions $f^{(j)}$ are linearly independent. Then

$$\mathbf{M}(z) = C^{-1} \mathbf{B}(z) \quad (9.24)$$

satisfies all the conditions in the RH problem for \mathbf{M} .

The $(1, 1)$ entry of \mathbf{M} is of special interest, since it appears in the asymptotic formula (2.11). This entry is given by

$$\mathbf{M}_{1,1}(z) = \exp(u_1(z)) = \exp\left(\int_{\infty_1}^z \boldsymbol{\eta}\right), \quad z \in \mathbb{C} \setminus \Sigma_1^*, \quad (9.25)$$

where η is the meromorphic differential defined above. To see this, observe that according to (9.24) and (9.20),

$$\mathbf{M}_{1,1}(z) = \widehat{v}_1(z) \sum_{i=1}^{d+1} (C^{-1})_{1,i} f_1^{(i-1)}(z) = e^{u_1(z)} \sum_{i=1}^{d+1} (C^{-1})_{1,i} f_1^{(i-1)}(z),$$

where $(C^{-1})_{1,i}$ denotes the $(1, i)$ entry of C^{-1} . Since $\widehat{v}_j(z) = O(z^{-\frac{d+1}{2d}})$ as $z \rightarrow \infty$ for $j \geq 2$, it follows from the expression of $\mathbf{A}^{-1}(z)$ and (9.23) that the entries $(1, i)$, $i \geq 2$, of C are all zero. This implies that $(C^{-1})_{1,i} = 0$ for all $i \geq 2$, hence $\mathbf{M}_{1,1}(z) = c e^{u_1(z)}$ for some constant c . By (9.4) we have $\mathbf{M}_{1,1}(z) = 1 + O(z^{-1})$, so $c = 1$ and (9.25) follows.

10 The final transformation $\mathbf{S} \mapsto \mathbf{R}$ and conclusion of the steepest descent analysis

In this section we introduce the final transformation of the RH problem and give the proof of Theorem 2.3. This final transformation makes use of the global parametrix from the previous section and of a local parametrix $\mathbf{P}^{(\text{Airy})}$ involving Airy functions that is defined on small disks around the endpoints $\omega^\ell x^*$, $\ell = 0, \dots, d$ of the star Σ_1^* . We define disks

$$D(\omega^\ell x^*, \delta) = \{z \in \mathbb{C} \mid |z - \omega^\ell x^*| < \delta\},$$

where $\delta > 0$ is taken sufficiently small. In any case, we want $\delta < \widehat{x} - x^*$ and the disks should be contained in the lense L_3 . Recall that $t_0 < t_{0,\text{crit}}$, and therefore we know by Lemma 3.6 that the density of μ_1^* vanishes as a square root at the endpoints of Σ_1^* . This property and the fact that the RH problem is locally of size 2×2 allow us to construct in a standard way a function $\mathbf{P}^{(\text{Airy})}$ that is the solution to the following problem.

RH problem 10.1. • $\mathbf{P}^{(\text{Airy})}$ is continuous on $\left(\bigcup_{\ell=0}^d \overline{D(\omega^\ell x^*, \delta)}\right) \setminus \Sigma_S$ and is analytic in its interior $\left(\bigcup_{\ell=0}^d D(\omega^\ell x^*, \delta)\right) \setminus \Sigma_S$.

- $\mathbf{P}_+^{(\text{Airy})} = \mathbf{P}_-^{(\text{Airy})} \mathbf{J}_S$ on $\Sigma_S \cap \bigcup_{\ell=0}^d D(\omega^\ell x^*, \delta)$, where \mathbf{J}_S is the jump matrix for \mathbf{S} as given in (8.14), (8.15), (8.18).
- $\mathbf{P}^{(\text{Airy})}$ matches with the global parametrix \mathbf{M} in the sense that

$$\mathbf{P}^{(\text{Airy})}(z) = \mathbf{M}(z)(I_{d+1} + O(n^{-1})), \quad (10.1)$$

uniformly for $z \in \bigcup_{\ell=0}^d \partial D(\omega^\ell x^*, \delta)$.

We omit here the construction of $\mathbf{P}^{(\text{Airy})}$ in terms of Airy functions. The details in the 2×2 case can be found in [8].

We now define the final transformation $\mathbf{S} \mapsto \mathbf{R}$.

Definition 10.2. We define the matrix-valued function $\mathbf{R} : \mathbb{C} \setminus (\Sigma_S \cup \bigcup_{\ell} \partial D(\omega^\ell x^*, \delta)) \rightarrow \mathbb{C}^{(d+1) \times (d+1)}$ by

$$\mathbf{R}(z) := \begin{cases} \mathbf{S}(z) (\mathbf{P}^{(\text{Airy})})^{-1}(z), & \text{in the disks } D(\omega^\ell x^*, \delta), \\ \mathbf{S}(z) \mathbf{M}(z)^{-1}, & \text{outside the disks.} \end{cases} \quad (10.2)$$

Since the jump matrices of \mathbf{S} and \mathbf{M} agree on Σ_1^* , Σ_2 and on $\Sigma_3 \setminus \Sigma_1$, we see from (10.2) that \mathbf{R} has analytic continuation across Σ_2 and $\Sigma_3 \setminus \Sigma_1$ and across the part of Σ_1^* outside the disks. The jump matrices of \mathbf{S} and $\mathbf{P}^{(\text{Airy})}$ agree inside the disks, and therefore \mathbf{R} can be analytically continued inside the disks. The result is that R is analytic in $\mathbb{C} \setminus \Sigma_R$, where Σ_R is a system of contours as shown in Figure 10 for the case $d = 3$. Thus Σ_R consists of the intervals $\omega^\ell[x^* + \delta, \hat{x}]$ for $\ell = 0, \dots, d$, the part of the boundary ∂L_1 of the lens L_1 that is outside of the disks, the full boundaries ∂L_2 and ∂L_3 of the other lenses, and the circles $\partial D(\omega^\ell x^*, \delta)$. These circles are given the positive orientation in Σ_R .




Figure 10: The contour Σ_R in the case $d = 3$.

We immediately observe from (10.2) that \mathbf{R} satisfies the following RH problem.

RH problem 10.3. • The matrix \mathbf{R} is analytic in $\mathbb{C} \setminus \Sigma_R$.

- We have $\mathbf{R}_+ = \mathbf{R}_- \mathbf{J}_R$ on Σ_R , where

$$\mathbf{J}_R(z) = \begin{cases} \mathbf{M}(z) \mathbf{P}^{(\text{Airy})}(z)^{-1}, & \text{for } z \in \bigcup_{\ell=0}^d \partial D(\omega^\ell x^*, \delta), \\ \mathbf{M}_-(z) \mathbf{J}_S(z) \mathbf{M}_+^{-1}(z), & \text{for } z \in \Sigma_R \setminus \left(\bigcup_{\ell=0}^d \partial D(\omega^\ell x^*, \delta) \right). \end{cases} \quad (10.3)$$

- $\mathbf{R}(z) = I_{d+1} + O(z^{-1})$ as $z \rightarrow \infty$.

Note that the asymptotic conditions (8.22) and (9.4) for \mathbf{S} and \mathbf{M} both contain $\begin{pmatrix} 1 & 0 \\ 0 & A(z) \end{pmatrix}$, and this is cancelled in the asymptotic condition for \mathbf{R} . Then one gets $\mathbf{R}(z) = I_{d+1} + O(z^{-\frac{2}{d}})$ as $z \rightarrow \infty$. However, since all jump matrices in the RH problem for \mathbf{R} are exponentially close to the identity matrix for large z , see (10.5), it can be shown that the O -term improves to $O(z^{-1})$.

The important feature of the RH problem associated with \mathbf{R} is that the jump matrix $\mathbf{J}_R(z)$ tends to the identity matrix as $n \rightarrow \infty$, provided that the point $\widehat{x} > x^*$ is chosen so that it lies in the region where $\operatorname{Re} \varphi_1(z) < 0$. Indeed, from (10.1) and (10.3) we get

$$\mathbf{J}_R(z) = I_{d+1} + O(n^{-1}), \quad z \in \bigcup_{\ell=0}^d \partial D(\omega^\ell x^*, \delta), \quad (10.4)$$

and on the remaining parts of Σ_R we have convergence to the identity matrix exponentially fast, specifically

$$\mathbf{J}_R(z) = I_{d+1} + O\left(e^{-cn|z|^{\frac{d+1}{d}}}\right), \quad \text{for } z \text{ elsewhere on } \Sigma_R, \quad (10.5)$$

for some constant $c > 0$. The estimate (10.5) follows from (10.3) and the corresponding estimate (8.34) for $\mathbf{J}_S(z)$, which is valid uniformly on the parts of Σ_R that come from $\Sigma_1 \setminus \Sigma_1^*$, and from the lips of the lenses.

We can now conclude the steepest descent analysis. From (10.4)–(10.5) we deduce several important consequences. First, it follows that the RH problem 10.3 has a unique solution if n is large enough. The solution \mathbf{R} can be expressed as a Neumann series. Since the transformations performed during the steepest descent analysis

$$\mathbf{Y} \mapsto \mathbf{X} \mapsto \mathbf{U} \mapsto \mathbf{T} \mapsto \mathbf{S} \mapsto \mathbf{R} \quad (10.6)$$

are all invertible, this implies that a solution \mathbf{Y} to the original RH problem 4.2 uniquely exists. In particular, the polynomial $P_{n,n}$, which is the $(1, 1)$ entry of \mathbf{Y} , uniquely exists. We also deduce from (10.4)–(10.5) that the solution R to the RH problem 10.3 is itself close to the identity as $n \rightarrow \infty$. In fact, it follows from (10.4) and (10.5) that R satisfies

$$\mathbf{R}(z) = I_{d+1} + O\left(\frac{1}{n(1+|z|)}\right), \quad \text{as } n \rightarrow \infty, \quad (10.7)$$

uniformly for $z \in \mathbb{C} \setminus \Sigma_R$.

10.1 Proof of Theorem 2.3

We unravel the transformations (10.6) in order to express the polynomial $P_{n,n}$ in terms of the matrix-valued function \mathbf{R} and then we use (10.7).

We start from the relation $P_{n,n}(z) = \mathbf{Y}_{1,1}(z)$ that we know from Lemma 4.3. From (5.41), (6.1) and (7.32) we easily obtain

$$P_{n,n}(z) = \mathbf{X}_{1,1}(z) = \mathbf{U}_{1,1}(z) = \mathbf{T}_{1,1}(z) e^{ng_1(z)}, \quad z \in \mathbb{C} \setminus \Sigma_1^*.$$

Next, by definition of \mathbf{S} in Definition 8.1 we know that $\mathbf{S}(z) = \mathbf{T}(z)$ for all $z \in \mathbb{C} \setminus (L_2 \cup L_3)$, and so certainly $\mathbf{S}_{1,1}(z) = \mathbf{T}_{1,1}(z)$ for those z . But from (8.3) and (8.4) we see that $\mathbf{S}_{1,1}(z) = \mathbf{T}_{1,1}(z)$ also for $z \in (L_2 \cup L_3) \setminus L_1$. Hence

$$P_{n,n}(z) = \mathbf{S}_{1,1}(z) e^{ng_1(z)}, \quad z \in \mathbb{C} \setminus L_1. \quad (10.8)$$

On account of (10.2), we have $\mathbf{S} = \mathbf{R}\mathbf{M}$ outside of the disks $D(\omega^\ell x^*, \delta)$, so using (10.7) we obtain

$$\mathbf{S}_{1,1}(z) = (1 + O(n^{-1})) \mathbf{M}_{1,1}(z) + O(n^{-1})$$

for such z . Since $\mathbf{M}_{1,1}(z)$ is an analytic function with no zeros in $\mathbb{C} \setminus \Sigma_1^*$, the above estimate can be rewritten as

$$\mathbf{S}_{1,1}(z) = (1 + O(n^{-1})) \mathbf{M}_{1,1}(z), \quad z \in \mathbb{C} \setminus \left(\bigcup_{\ell=0}^d D(\omega^\ell x^*, \delta) \right), \quad (10.9)$$

uniformly for z in the indicated set.

Finally, from (10.8) and (10.9) we obtain (2.11) uniformly for $z \in \mathbb{C} \setminus \left(L_1 \cup \bigcup_{\ell=0}^d D(\omega^\ell x^*, \delta) \right)$.

Since the lens L_1 and the disks $D(\omega^\ell x^*, \delta)$ can be taken as small as we like, it follows that (2.11) is also valid uniformly for z in compact subsets of $\mathbb{C} \setminus \Sigma_1^*$.

11 Acknowledgements

The first author is supported by KU Leuven Research Grant OT/12/073, the Belgian Interuniversity Attraction Pole P07/18, and the FWO Flanders projects G.0641.11 and G.0934.13.

The second author was supported by a postdoctoral fellowship from the Fund for Scientific Research - Flanders (Belgium) during his stay at the University of Leuven.

The authors are grateful to the participants of the AIM workshop on Vector Equilibrium problems and their Applications to Random Matrix Models in April 2012 for fruitful and inspiring discussions on the normal matrix model.

References

- [1] A.I. Aptekarev, V.A. Kalyagin, J. Van Iseghem, The genetic sum's representation for the moments of a system of Stieltjes functions and its applications, *Constr. Approx.* 16 (2000), 487–524.
- [2] A.I. Aptekarev, V.A. Kalyagin, E.B. Saff, Higher order three-term recurrences and asymptotics of multiple orthogonal polynomials, *Constr. Approx.* 30 (2009), 175–223.
- [3] Y. Ameur, H. Hedenmalm, N. Makarov, Fluctuations of eigenvalues of random normal matrices, *Duke Math. J.* 159 (2011), 31–81.
- [4] Y. Ameur, H. Hedenmalm, N. Makarov, Random normal matrices and Ward identities, to appear in *Ann. Prob.*, preprint arXiv:1109.5941.
- [5] F. Balogh, M. Bertola, S.-Y. Lee, K.D.T-R McLaughlin, Strong asymptotics of the orthogonal polynomial with respect to a measure supported on the plane, *Comm. Pure Appl. Math.* 68 (2015), 112–172.
- [6] B. Beckermann, V. Kalyagin, A. Matos, F. Wielonsky, Equilibrium problems for vector potentials with semidefinite interaction matrices and constrained masses, *Constr. Approx.* 37 (2013), 101–134.
- [7] P.M. Bleher, A.B.J. Kuijlaars, Orthogonal polynomials in the normal matrix model with a cubic potential, *Adv. Math.* 230 (2012), 1272–1321.
- [8] P. Deift, *Orthogonal Polynomials and Random Matrices: a Riemann-Hilbert approach*, Courant Lecture Notes in Mathematics Vol. 3, Amer. Math. Soc., Providence R.I. 1999.

- [9] S. Delvaux, A.B.J. Kuijlaars, A graph-based equilibrium problem for the limiting distribution of non-intersecting Brownian motions at low temperature, *Constr. Approx.* 32 (2010), 467–512.
- [10] S. Delvaux, A. López, High-order three-term recursions, Riemann-Hilbert minors and Nikishin systems on star-like sets, *Constr. Approx.* 37 (2013), 383–453.
- [11] M. Duits, A.B.J. Kuijlaars, Painlevé I asymptotics for orthogonal polynomials with respect to a varying quartic weight, *Nonlinearity* 19 (2006), 2211–2245.
- [12] M. Duits, A.B.J. Kuijlaars, Universality in the two matrix model: a Riemann-Hilbert steepest descent analysis, *Comm. Pure Appl. Math.* 62 (2009), 1076–1153.
- [13] M. Eiermann, R.S. Varga, Zeros and local extreme points of Faber polynomials associated with hypocycloidal domains, *Electron. Trans. Numer. Anal.* 1 (1993), 49–71.
- [14] P. Elbau, Random normal matrices and polynomial curves, Ph.D. thesis, ETH Zürich, arXiv:0707.0425.
- [15] P. Elbau, G. Felder, Density of eigenvalues of random normal matrices, *Comm. Math. Phys.* 259 (2005), 433–450.
- [16] A.S. Fokas, A.R. Its, and A.V. Kitaev, The isomonodromy approach to matrix models in 2D quantum gravity, *Commun. Math. Phys.* 147 (1992), 395–430.
- [17] A. Hardy, A.B.J. Kuijlaars, Weakly admissible vector equilibrium problems, *J. Approx. Theory* 164 (2012), 854–868.
- [18] M.X. He, E.B. Saff, The zeros of Faber polynomials for an m -cusped hypocycloid, *J. Approx. Theory* 78 (1994), 410–432.
- [19] H. Hedenmalm, N. Makarov, Coulomb gas ensembles and Laplacian growth, *Proc. London Math. Soc.* 106 (2013), 859–907.
- [20] A.R. Its and L. A. Takhtajan. Normal matrix models, $\bar{\partial}$ -problem, and orthogonal polynomials on the complex plane, arXiv:0708.3867.
- [21] I. Kostov, I. Krichever, M. Mineev-Weinstein, P. Wiegmann, A. Zabrodin, τ -function for analytic curves, in: “Random Matrices and Their Applications”, MSRI Publications Vol. 40 (eds. P. Bleher, A. Its), Cambridge Univ. Press, Cambridge, 2001, pp. 285–299.
- [22] A.B.J. Kuijlaars, M.Y. Mo, The global parametrix in the Riemann-Hilbert steepest descent analysis for orthogonal polynomials, *Comput. Meth. Funct. Theory* 11 (2011), 161–178.
- [23] A.B.J. Kuijlaars, A. Tovbis, The supercritical regime in the normal matrix model with cubic potential, preprint arXiv:1412.7597.
- [24] S.-Y. Lee, R. Teodorescu, P. Wiegmann, Shocks and finite-time singularities in Hele-Shaw flow, *Physica D* 238 (2009), 1113–1128.
- [25] S.-Y. Lee, R. Teodorescu, P. Wiegmann, Viscous shocks in Hele-Shaw flow and Stokes phenomena of the Painlevé I transcendent, *Physica D* 240 (2011), 1080–1091.
- [26] A. López García, Asymptotics of multiple orthogonal polynomials for a system of two measures supported on a starlike set, *J. Approx. Theory* 163 (2011), 1146–1184.

- [27] M. Mineev-Weinstein, M. Putinar, R. Teodorescu, Random matrices in 2D, Laplacian growth and operator theory, *J. Phys. A: Math. Theor.* 41 (2008), 263001, 74pp.
- [28] E.M. Nikishin and V.N. Sorokin, *Rational approximations and orthogonality*, Translations of Mathematical Monographs vol. 92, American Mathematical Society, Providence, RI, 1991.
- [29] T. Ransford, *Potential Theory in the Complex Plane*, London Mathematical Society Student Texts vol. 28, Cambridge University Press, 1995.
- [30] R. Riser, Universality in Gaussian Random Normal Matrices, arXiv:1312.0068.
- [31] N. Ben Romdhane, On the zeros of d -orthogonal d -symmetric polynomials, *J. Math. Anal. Appl.* 344 (2008), 888–897.
- [32] E.B. Saff, V. Totik, *Logarithmic Potentials with External Fields*, Springer-Verlag, Berlin, 1997.
- [33] C. Sinclair, M. Yattselev, Universality for ensembles of matrices with potential theoretic weights on domains with smooth boundary, *J. Approx. Theory* 164 (2012), 682–708.
- [34] R. Teodorescu, E. Bettelheim, O. Agam, A. Zabrodin, P. Wiegmann, Normal random matrix ensemble as a growth problem, *Nuclear Phys. B* 704 (2005), 407–444.
- [35] W. Van Assche, J. Geronimo, A.B.J. Kuijlaars, Riemann-Hilbert problems for multiple orthogonal polynomials, in: “Special Functions 2000” (J. Bustoz, et al., eds.), Kluwer, Dordrecht, 2001, pp. 23–59.
- [36] P. Wiegmann, A. Zabrodin, Conformal maps and integrable hierarchies, *Comm. Math. Phys.* 213 (2000), 523–538.
- [37] P. Wiegmann, A. Zabrodin, Large scale correlations in normal and general non-hermitian matrix ensembles, *J. Phys. A* 36 (2003), 3411–3424.
- [38] A. Zabrodin, Random matrices and Laplacian growth, Chapter 39 in: “The Oxford Handbook of Random Matrix Theory” (G. Akemann, J. Baik, P. Di Francesco, eds.), Oxford University Press, 2011.