Discovering functional DNA elements using population genomic information: A proof of concept using human mtDNA

Daniel R. Schrider and Andrew D. Kern

Department of Genetics, Rutgers, The State University of New Jersey, Piscataway, NJ 08854

Correspondence: D.R.S. (dan.schrider@rutgers.edu)

ABSTRACT

Identifying the complete set of functional elements within the human genome would be a windfall for multiple areas of biological research including medicine, molecular biology, and evolution. Complete knowledge of function would aid in the prioritization of loci when searching for the genetic bases of disease or adaptive phenotypes. Because mutations that disrupt function are disfavored by natural selection, purifying selection leaves a detectable signature within functional elements; accordingly this signal has been exploited for over a decade through the use of genomic comparisons of distantly related species. While this is so, the functional complement of the genome changes extensively across time and between lineages, therefore, evidence of the current action of purifying selection in humans is essential. Because the removal of deleterious mutations by natural selection also reduces within-species genetic diversity within functional loci, dense population genetic data have the potential to reveal genomic elements that are currently functional. Here we assess the potential of this approach by examining an ultra-deep sample of human mitochondrial genomes (n=16,411). We show that the high density of polymorphism in this dataset precisely delineates regions experiencing purifying selection. Further, we show that the number of segregating alleles at a site is strongly correlated with its divergence across species after accounting for known mutational biases in human mtDNA (ρ =0.51; P<2.2x10-16). These two measures track one another at a remarkably fine scale across many loci-a correlation that is purely the result of natural selection. Our results demonstrate that genetic variation has the potential to reveal exactly which nucleotides in the genome are currently performing important functions and likely to have deleterious fitness effects when mutated. As more complete human genomes are sequenced, similar power to reveal purifying selection may be achievable in the human nuclear genome.

INTRODUCTION

Only 1-2% of human genome lies within protein-coding sequence (Lander et al. 2001). Determining the extent to which the remainder of the genome is functional is crucial to our understanding of human biology. A variety of recently developed experimental techniques have aided in the search of non-coding DNA for functional elements (Dunham et al. 2012), however, on their own these techniques can produce a huge number of false positives (Graur et al. 2013). Searches for the evolutionary signature of purifying selection have therefore proved a more fruitful strategy for identifying functional elements; indeed phylogenetic searches comparing sequences of related species have revealed that roughly 5% of the human genome is constrained by natural selection (Chinwalla et al. 2002; Siepel et al. 2005; Lunter et al. 2006; Birney et al. 2007; Davydov et al. 2010), and similar strategies have been used to predict the phenotypic severity of mutations (Stone and Sidow 2005). While whole-genome comparisons aimed at identifying the footprints of selection are highly effective, they have been used primarily to detect elements under constraint for hundreds of millions of years of evolutionary history (Siepel et al. 2005; Davydov et al. 2010). However, the set of functional elements in the genome experiences considerable turnover (Demuth et al. 2006). Comparative genomic techniques will fail in these instances, and particularly for the supremely interesting cases of human-specific gain (Knowles and McLysaght 2009) and loss of function (Wang et al. 2006).

Surveys of genetic diversity within species, on the other hand, have the potential to identify regions currently experiencing purifying selection and that are therefore functional, as purifying selection will remove genetic diversity from such loci. Unfortunately, genetic variation in the human genome is quite sparse, with a comparison of any two homologous chromosomes uncovering <1 single nucleotide polymorphism (SNP) every kilobase (Lander et al. 2001). Sampling more individuals, however, yields additional polymorphisms, and an ultra-deep sample of mitochondrial variation from 16,411 genomes is available on the MITOMAP database (Ruiz-Pesini et al. 2007). These data are extremely polymorphic, with more than one SNP every other base pair on average. This dataset thus serves as an ideal proving ground for the approach of identifying functional constraint using massive amounts of polymorphism data, which will soon be available for nuclear genomes. Here we show that the density of polymorphism in these data closely tracks divergence at a fine scale, implying that these data can indeed be used to reveal the strength of purifying selection in the human mitochondrial genome at a very high resolution. Our results suggest an enormous potential for population genomic data to uncover functional DNA elements, including those not conserved across species.

RESULTS AND DISCUSSION

We set out to determine the extent to which polymorphism data reveal the strength of purifying selection across the human mitochondrial genome, and downloaded the coordinates of all 8,944 SNPs from MITOMAP (http://www.mitomap.org/MITOMAP). We reasoned that if the density of polymorphism were governed by the amount of purifying selection acting on each site, then SNP density would be correlated with divergence across species, in accordance with expectations under the Neutral model (Kimura 1982). This is indeed what we observe, in the form of a strong correlation between the number of alleles per site and its average negated PhyloP score (Siepel et al. 2006) measuring divergence across vertebrates (Spearman's ρ =0.52; P<2.2x10⁻¹⁶). This correlation is also highly significant when averaging polymorphism and divergence within 10 bp adjacent windows (ρ =0.50; P<2.2x10⁻¹⁶; Figure 1).

While this observation is consistent with purifying selection both removing diversity and constraining divergence at functional elements, such a pattern could also be generated by variation in the spontaneous mutation rate. It has been shown that mutation rate in the mitochondria varies according to the duration for which a given site remains single stranded on the H strand (D_{ssH}) during DNA replication (Reves et al. 1998). We also find evidence for this in the form a significant correlation between divergence at each site and the duration the site is single-stranded on the H strand during replication, though this correlation is far weaker than that shared between polymorphism and divergence ($\rho=0.11$; $P<2.2\times10^{-16}$). Moreover, after correcting for D_{ssH} , the correlation between polymorphism and divergence at individuals sites is essentially unchanged and still highly significant ($\rho=0.49$; $P<2.2\times10^{-16}$). Rather than being driven by a subset of mitochondrial loci, this correlation is significant (at P < 0.05) in 36/37 genes, and is significant in 35/37 genes after correcting for D_{ssH} (Table 1). This correlation is also far stronger at nonsynonymous than synonymous sites ($\rho=0.25$ for second codon position sites; $P<2.2 \times 10^{-16}$; ρ =0.079 for fourfold degenerate sites; P=3.6x10⁻⁴; Figure 2), as expected if purifying selection is a more predominant force at nonsynonymous sites. Finally, the minor allele frequencies of SNPs from mtDB (Ingman and Gyllensten 2006) are correlated with divergence (ρ =0.076; P=4.0x10⁻⁶), even though variation in mutation rate is not expect to affect allele frequencies. Thus purifying

selection uniquely drives patterns of polymorphism in the human mitochondrial genome. This finding supports previous reports that purifying selection is a prominent force in the mitochondrial genome (Rand and Kann 1996; Nielsen and Weinreich 1999; Elson et al. 2004; Stewart et al. 2008), providing further evidence that the hypothesis that drift alone governs mitochondrial evolution (Avise et al. 1987; Palumbi et al. 2001) should be rejected. The patterns we observed are not the result of positive selection, as the fixation of a beneficial mutation through a selective sweep removes all genetic diversity from a non-recombining chromosome (Smith and Haigh 1974). We are thus limited to observing mutations occurring since the most recent sweep.

Having established that patterns of polymorphism across the human mitochondria are largely determined by purifying selection, we sought to determine the resolution at which these data reveal the strength of selection acting on particular sites in the genome. We examined patterns of SNP diversity and divergence in 5 bp sliding windows across each gene in the mitochondrial genome, observing the extent to which the two measures mirror one another on a small scale. In particular, within each window we calculated the average SNP density per base pair and the average probability that the site is not conserved across vertebrates according to phastCons (Siepel et al. 2005). We find that for many loci, tRNA genes in particular, these two measures track one another to a surprising extent (e.g. the phenylalanine and tryptophan tRNA genes shown in Figure 3; the remaining tRNA genes are shown in supplementary figure S1). This result demonstrates that SNP density has the ability to reveal the strength of selection at a surprisingly detailed resolution-on the scale of a few base pairs. We therefore sought to use SNP density to predict function at a fine scale via a hidden Markov model (HMM; Rabiner 1989). Using a similar strategy as phastCons (Siepel et al. 2005), we learned a two-state HMM (constrained vs. unconstrained) where the observation for each site in the genome is the number of alleles at the site. We then used this HMM to predict constrained regions to which we refer as mitoPopCons elements. There is extremely strong overlap between mitoPopCons elements obtained from polymorphism data and phastCons elements predicted from divergence (Pvalue<0.0001; Figure 4; Methods). SNP diversity therefore has the ability to accurately predict function at a fine scale in the human mitochondria.

We have shown that ultra-dense polymorphism data can be used to accurately detect functional nucleotides in the human mitochondrial genome, potentially at the level of the individual base pair, while sidestepping limitations of phylogenetic approaches. This result suggests that as whole-genome sequencing becomes more ubiquitous, it will become possible to perform such high-resolution prediction in the nuclear genome as well. Approaches such as ours will then have an enormous impact on biological research, allowing for the discovery of the complete set of functional elements in the human genome and the degree to which new mutations at each site are deleterious. Such efforts will vastly improve predictions of the phenotypic impact of mutations occurring in humans and will prioritize searches for diseasecausing mutations. This information will also reveal species-specific changes in selective pressures at the resolution of individual nucleotides, greatly improving our understanding of how the functional components of genomes evolve.

METHODS

We converted all coordinates from obtained from MITOMAP and mtDB from the CRS to the human reference genome's (GRCh37) coordinate space by aligning the two mitochondrial sequences using MUSCLE (Edgar 2004). We downloaded phyloP scores for each mitochondrial site from the UCSC Genome Browser Database (Meyer et al. 2013). Because there are no allele frequency data available for MITOMAP SNPs, we used the number of alleles at each site as our measure of diversity. For the correlation of minor allele frequency with divergence, we used frequencies of biallelic SNPs from mtDB. We measured the duration of single-strandedness (D_{ssH}) following Chong and Mueller (Chong and Mueller 2013), using the coordinates of the light-strand origin of replication from MITOMAP. For the D_{ssH} analyses we omitted sites within the control region. To control for the D_{ssH} -divergence correlation, we fit a generalized linear model treating the number of segregating alleles at each site as a Poisson distributed response variable and with D_{ssH} value as the predictive variable using R. We then calculated the correlation between PhyloP scores and the polymorphism residuals. The HMM analysis was performed in MATLAB with the same transition matrix as used to learn the phastCons HMM for the UCSC Genome Browser (Kent et al. 2002). We then used Baum-Welch training to learn the emission parameters only for the constrained and unconstrained states by using large number of pseudocounts for transition matrix to ensure it remained invariant during training. Next, we predicted constrained elements by using the Viterbi algorithm. We tested for significant overlap between our HMM predictions and mammalian phastCons elements from the UCSU Genome Browser by counting the number of base pairs constrained according to both methods, and compared this count to those obtained after permuting the coordinates of our predictions. This was repeated for 10,000 permutations to obtain a P-value.

ACKNOWLEDGEMENTS

D. R. S. was supported by the National Institutes of Health under Ruth L. Kirschstein National Research Service Award F32 GM105231-01, and A. D. K. was supported in part by Rutgers University and National Science Foundation Award MCB-1161367.

REFERENCES

Avise JC, Arnold J, Ball RM, Bermingham E, Lamb T, Neigel JE, Reeb CA and Saunders NC. 1987. Intraspecific phylogeography: the mitochondrial DNA bridge between population genetics and systematics. Annual review of ecology and systematics 18: 489-522.

Birney E, Stamatoyannopoulos JA, Dutta A, et al. 2007. Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project. Nature 447: 799-816.

Chinwalla AT, Cook LL, Delehaunty KD, et al. 2002. Initial sequencing and comparative analysis of the mouse genome. Nature 420: 520-562.

Chong RA and Mueller RL. 2013. Evolution along the mutation gradient in the dynamic mitochondrial genome of salamanders. Genome Biol Evol.

Davydov EV, Goode DL, Sirota M, Cooper GM, Sidow A and Batzoglou S. 2010. Identifying a high fraction of the human genome to be under selective constraint using GERP++. PLoS Comput Biol 6: e1001025.

Demuth JP, De Bie T, Stajich JE, Cristianini N and Hahn MW. 2006. The evolution of mammalian gene families. PLoS ONE 1: e85.

Dunham I, Kundaje A, Aldred SF, et al. 2012. An integrated encyclopedia of DNA elements in the human genome. Nature 489: 57-74.

Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res 32: 1792-1797.

Elson J, Turnbull D and Howell N. 2004. Comparative genomics and the evolution of human mitochondrial DNA: assessing the effects of selection. The American Journal of Human Genetics 74: 229-238.

Graur D, Zheng Y, Price N, Azevedo RB, Zufall RA and Elhaik E. 2013. On the immortality of television sets: "function" in the human genome according to the evolution-free gospel of ENCODE. Genome Biol Evol 5: 578-590.

Ingman M and Gyllensten U. 2006. mtDB: Human Mitochondrial Genome Database, a resource for population genetics and medical sciences. Nucleic Acids Res 34: D749-D751.

Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM and Haussler D. 2002. The human genome browser at UCSC. Genome Res 12: 996-1006.

Kimura M. 1982. The neutral theory as a basis for understanding the mechanism of evolution and variation at the molecular level. In: Kimura M, editor. Molecular Evolution, Protein Polymorphism and the Neutral Theory Springer-Verlag, Berlin. Berlin: Springer-Verlag. p. 3-56.

Knowles DG and McLysaght A. 2009. Recent de novo origin of human protein-coding genes. Genome Res 19: 1752-1759.

Lander ES, Linton LM, Birren B, et al. 2001. Initial sequencing and analysis of the human genome. Nature 409: 860-921.

Lunter G, Ponting CP and Hein J. 2006. Genome-wide identification of human functional DNA using a neutral indel model. PLoS Comput Biol 2: e5.

Meyer LR, Zweig AS, Hinrichs AS, et al. 2013. The UCSC Genome Browser database: extensions and updates 2013. Nucleic Acids Res 41: D64-D69.

Nielsen R and Weinreich DM. 1999. The age of nonsynonymous and synonymous mutations in animal mtDNA and implications for the mildly deleterious theory. Genetics 153: 497-506.

Palumbi SR, Cipriano F and Hare MP. 2001. Predicting nuclear gene coalescence from mitochondrial data: the three-times rule. Evolution 55: 859-868.

Rabiner LR. 1989. A tutorial on hidden Markov models and selected applications in speech recognition. Proceedings of the IEEE 77: 257-286.

Rand DM and Kann LM. 1996. Excess amino acid polymorphism in mitochondrial DNA: contrasts among genes from Drosophila, mice, and humans. Mol Biol Evol 13: 735-748.

Reyes A, Gissi C, Pesole G and Saccone C. 1998. Asymmetrical directional mutation pressure in the mitochondrial genome of mammals. Mol Biol Evol 15: 957-966.

Ruiz-Pesini E, Lott MT, Procaccio V, et al. 2007. An enhanced MITOMAP with a global mtDNA mutational phylogeny. Nucleic Acids Res 35: D823-D828.

Siepel A, Bejerano G, Pedersen JS, et al. 2005. Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes. Genome Res 15: 1034-1050.

Siepel A, Pollard KS and Haussler D editors. Research in Computational Molecular Biology. 2006.

Smith JM and Haigh J. 1974. The hitch-hiking effect of a favourable gene. Genet Res 23: 23-35.

Stewart JB, Freyer C, Elson JL, Wredenberg A, Cansu Z, Trifunovic A and Larsson N-G. 2008. Strong purifying selection in transmission of mammalian mitochondrial DNA. PLoS Biol 6: e10.

Stone EA and Sidow A. 2005. Physicochemical constraint violation by missense substitutions mediates impairment of protein function and disease severity. Genome Res 15: 978-986.

Wang X, Grus WE and Zhang J. 2006. Gene losses during human origins. PLoS Biol 4: e52.



Figure 1: The correlation between polymorphism and divergence in the human mitochondrial genome. The average number of alleles per base pair in 10 bp windows is shown on the x-axis, and divergence as measured by the negated phyloP score is shown on the y-axis.



Figure 2: The correlation between polymorphism and divergence at nonsynonymous and synonymous sites. The number of alleles observed at each site is shown on the x-axis and divergence (negative phyloP score) is shown on the y-axis at (A) 2nd codon position (nonsynonymous) sites and (B) 4-fold degenerate (synonymous) sites. We added noise to the number of alleles in order to reveal the density of sites along the y-axis.



Figure 3: The probability of polymorphism versus the probability of unconstrained evolution across vertebrates. (A) 5bp sliding genomic windows showing SNP density (blue) and one minus the probability of conservation across vertebrates (red) according to PhastCons(Siepel et al. 2005) across the Phenylalanine tRNA gene. (B) The same plot for the Tryptophan tRNA gene.



Figure 4: Comparison of conserved elements called from phylogenetic data and those called from population genetic data. This image form the UCSC Genome Browser shows positions 1-5,000 of the human mitochondrial genome. Conserved elements called from the polymorphism-based HMM (mitoPopCons) appear in blue, while phastCons elements obtained from a comparison of mammalian genomes appear in red. PhastCons conservation probabilities are shown at the bottom in green. Gene locations are shown at the top.

						Spearman's	
		Gene	Gene	Spear		ρ	<i>P</i> -value
Gene	Gene	Start	End	man's		(correcting	(correcting
Name	Description	(hg19)	(hg19)	ρ	<i>P</i> -value	for D_{SSH})	for D_{SSH})
	tRNA						
MT-TF	phenylalanine	579	649	-0.484	1.90E-05	-0.468948	3.71E-05
MT-	12S ribosomal						
RNR1	RNA	650	1603	-0.429	<2.20E-16	-0.4169046	<2.20E-16
MT-TV	tRNA valine	1604	1672	-0.336	0.004783	-0.326022	0.006261
MT-	16S ribosomal						
RNR2	RNA	1673	3230	-0.443	<2.20E-16	-0.4435798	<2.20E-16
	tRNA leucine						
MT-TL1	1	3231	3305	-0.301	0.008593	-0.29837	0.00932
	NADH						
MT-	Dehydrogenase						
ND1	subunit 1	3308	4263	-0.542	<2.20E-16	-0.5149936	<2.20E-16
	tRNA						
MT-TI	isoleucine	4264	4332	-0.269	0.02551	-0.2523745	0.03643
	tRNA						
MT-TQ	glutamine	4330	4401	-0.463	4.19E-05	-0.476385	2.34E-05
	tRNA						
MT-TM	methionine	4403	4470	-0.291	0.01611	-0.2880727	0.01721
	NADH						
MT-	dehydrogenase						
ND2	subunit 2	4471	5512	-0.516	<2.20E-16	-0.461188	<2.20E-16
	tRNA						
MT-TW	tryptophan	5513	5580	-0.509	9.42E-06	-0.4626812	7.11E-05
MT-TA	tRNA alanine	5588	5656	-0.429	0.0002381	-0.4412621	0.0001475
	tRNA						
MT-TN	asparagine	5658	5730	-0.378	0.0009899	-0.3765626	0.001025
MT-TC	tRNA cysteine	5762	5827	-0.596	1.26E-07	-0.5709837	5.55E-07
MT-TY	tRNA tyrosine	5827	5892	-0.352	0.00371	-0.3486535	0.004118
	Cytochrome c						
MT-	oxidase						
CO1	subunit I	5905	7446	-0.633	<2.20E-16	-0.6186776	<2.20E-16
MT-TS1	tRNA serine 1	7447	7515	-0.627	8.31E-09	-0.6107229	2.51E-08
	tRNA aspartic						
MT-TD	acid	7519	7586	-0.365	0.002186	-0.320253	0.007758
	Cytochrome c						
MT-	oxidase						
CO2	subunit II	7587	8270	-0.573	<2.20E-16	-0.5291891	<2.20E-16
MT-TK	tRNA lysine	8296	8365	-0.331	0.005158	-0.2809919	0.01846
MT-	ATP synthase						
ATP8	F0 subunit 8	8367	8573	-0.266	0.0001042	-0.2637231	0.0001233

Table 1: Gene-specific correlations between SNP density and negative phyloP score.

MT-	ATP synthase						
ATP6	F0 subunit 6	8528	9208	-0.389	<2.20E-16	-0.3823366	<2.20E-16
	Cytochrome c						
MT-	oxidase						
CO3	subunit III	9208	9991	-0.538	<2.20E-16	-0.5314281	<2.20E-16
MT-TG	tRNA glycine	9992	10059	-0.396	0.0008191	-0.3404812	0.004497
	NADH						
MT-	dehydrogenase						
ND3	subunit 3	10060	10405	-0.510	<2.20E-16	-0.5000739	<2.20E-16
MT-TR	tRNA arginine	10406	10470	-0.470	7.66E-05	-0.4415206	0.0002317
	NADH						
MT-	dehydrogenase						
ND4L	subunit 4L	10471	10767	-0.491	<2.20E-16	-0.5133884	<2.20E-16
	NADH						
MT-	dehydrogenase						
ND4	subunit 4	10761	12138	-0.561	<2.20E-16	-0.5419698	<2.20E-16
MT-TH	tRNA histidine	12139	12207	-0.277	0.02112	-0.1786413	0.1419
MT-TS2	tRNA serine2	12208	12266	-0.470	0.0001712	-0.489146	8.45E-05
MT-TL2	tRNA leucine2	12267	12337	-0.374	0.001311	-0.3543069	0.002434
	NADH						
MT-	dehydrogenase						
ND5	subunit 5	12338	14149	-0.532	<2.20E-16	-0.5141183	<2.20E-16
	NADH						
MT-	dehydrogenase						
ND6	subunit 6	14150	14674	-0.500	<2.20E-16	-0.4776221	<2.20E-16
	tRNA glutamic						
MT-TE	acid	14675	14743	-0.347	0.003468	-0.3476731	0.003421
MT-							
CYB	Cytochrome b	14748	15888	-0.551	<2.20E-16	-0.5025513	<2.20E-16
	tRNA						
MT-TT	threonine	15889	15954	-0.560	1.00E-06	-0.4354723	0.0002578
MT-TP	tRNA proline	15957	16024	-0.103	0.4044	-0.1870294	0.1267



Supplementary figure S1: Polymorphism versus divergence in 5bp adjacent windows across the 20 remaining mitochondrial tRNA genes. Phenylalanine and Tryptophan are shown in Figure 3.