Kohn-Sham Theory in the Presence of Magnetic Field

Andre Laestadius*

Department of Mathematics, KTH Royal Institute of Technology, Sweden (Dated: June 1, 2021)

Abstract

In the well-known Kohn-Sham theory in Density Functional Theory, a fictitious non-interacting system is introduced that has the same particle density as a system of N electrons subjected to mutual Coulomb repulsion and an external electric field. For a long time, the treatment of the kinetic energy was not correct and the theory was not well-defined for N-representable particle densities. In the work of [Hadjisavvas and Theophilou, Phys. Rev. A, 1984, 30, 2183], a rigorous Kohn-Sham theory for N-representable particle densities was developed using the Levy-Lieb functional. Since a Levy-Lieb-type functional can be defined for Current Density Functional Theory formulated with the paramagnetic current density, we here develop a rigorous N-representable Kohn-Sham approach for interacting electrons in magnetic field. Furthermore, in the one-electron case, criteria for N-representable particle densities to be v-representable are given.

I. INTRODUCTION

In the fundamental paper by Hohenberg and Kohn [1], the theoretical foundation of Density Functional Theory (DFT) was established. The Hohenberg-Kohn theorem states that, for a quantum mechanical system, the particle density ρ determines the scalar potential of the system up to a constant. Subsequently, Kohn and Sham provided an algorithm [2], the so-called Kohn-Sham equations, for computing the density. These equations bear much resemblance to the Hartree-Fock integro-differential equations. The idea of Kohn and Sham was to introduce a fictitious system of non-interacting particles that has the same particle density as the real interacting system. This is achieved by means of the exchange-correlation functional, which accounts for the non-classical two-particle interactions and the residual between the interacting and non-interacting kinetic energy. However, this functional remains unknown.

In the work of Hadjisavvas and Theophilou [3], a mathematically rigorous Kohn-Sham approach was developed. The importance of this work relies on the fact that N-representability can be guaranteed for a proper wavefunction, whereas v-representability cannot. This means, in principle, that any v-representable formalism is unjustified.

In the presence of a magnetic field, no Hohenberg-Kohn theorem exists at the present time (that is valid for any number of electrons). For the formulation of Current Density Functional Theory (CDFT) that uses the paramagnetic current density j^p , it is well-known that the density pair (ρ, j^p) does not determine the scalar potential and vector potential of the system [4]. Counterexamples have been constructed that show that a ground-state can come from two different Hamiltonians [4, 5]. Thus, the particle density ρ and the paramagnetic current density j^p do not fully determine the Hamiltonian. For a many-electron system, neither proof nor counterexample exists so far in the literature for a Hohenberg-Kohn theorem formulated with the total current density j [5, 6]. In the one-electron case, on the other hand, it is possible to give a direct proof that ρ and j determine the scalar and vector potential up to a gauge transformation [5, 6].

However, since the density pair (ρ, j^p) determines the (possibly degenerate) ground-state(s) of the system [5, 7], this work aims at continue the N-representable approach of [3] and develop a rigorous Kohn-Sham approach for CDFT formulated with the paramagnetic current density j^p .

II. CURRENT DENSITY FUNCTIONAL THEORY

We will in this paper consider a system of N interacting electrons subjected to both an electric and a magnetic field. The system's Hamiltonian is given by (in suitable units)

$$H(v,A) = \sum_{k=1}^{N} ((i\nabla_k - A(x_k))^2 + v(x_k)) + \sum_{1 \le k < l \le N} |x_k - x_l|^{-1},$$

where v(x) is the scalar potential and A(x) the vector potential. The magnetic field is computed from $B(x) = \nabla \times A(x)$. Throughout we will assume that the ground-state is non-degenerate, i.e., dim $\ker(e_0 - H(v, A)) = 1$, where e_0 is the lowest eigenvalue of H(v, A).

A. Preliminaries

To begin with, some mathematical concepts needed for the forthcoming discussion are introduced. We first mention some relevant function spaces. If for some $p \in [1, \infty)$ a function f satisfies $\int_{\mathbb{R}^n} |f|^p < \infty$, then f belongs to the normed space $L^p(\mathbb{R}^n)$ with norm $||f||_{L^p(\mathbb{R}^n)} = (\int_{\mathbb{R}^n} |f|^p)^{1/p}$. In the case $p = \infty$, we say $f \in L^\infty(\mathbb{R}^n)$ if

$$||f||_{L^{\infty}(\mathbb{R}^n)} = \operatorname{ess\,sup}\{|f||x \in \mathbb{R}^n\} < \infty.$$

Furthermore, $f \in L^2(\mathbb{R}^n)$ is said to belong to the Hilbert space $\mathcal{H}^1(\mathbb{R}^n)$ if

$$||f||_{\mathcal{H}^{1}(\mathbb{R}^{n})}^{2} = \int_{\mathbb{R}^{n}} |f|^{2} + \int_{\mathbb{R}^{n}} |\nabla f|^{2} < \infty.$$

Let $B_R = \{x \in \mathbb{R}^n | |x| \le R\}$ for R > 0. Then $f \in L^1_{loc}(\mathbb{R}^n)$ whenever $\int_{B_R} |f| < \infty$ for any B_R . For a vector u such that $(u)_l \in L^p$, l = 1, 2, 3, we write $u \in (L^p)^3$.

We say that a sequence $\{\psi_k\} \subset L^p(\mathbb{R}^n)$ converges in $L^p(\mathbb{R}^n)$ -norm to $\psi \in L^p(\mathbb{R}^n)$ if $\int_{\mathbb{R}^n} |\psi_k - \psi|^p \to 0$ as $k \to \infty$, and we write $\psi_k \to \psi$. For the Hilbert space $L^2(\mathbb{R}^n)$, with inner product $(\psi, \phi)_{L^2(\mathbb{R}^n)} = \int_{\mathbb{R}^n} \psi^* \phi$, we say that $\{\psi_k\} \subset L^2(\mathbb{R}^n)$ converges weakly to $\psi \in L^2(\mathbb{R}^n)$ if $(\psi_k, \phi)_{L^2(\mathbb{R}^n)} \to (\psi, \phi)_{L^2(\mathbb{R}^n)}$ as $k \to \infty$ for all $\phi \in L^2(\mathbb{R}^n)$, and we write $\psi_k \to \psi$. For weak convergence in $\mathcal{H}^1(\mathbb{R}^n)$, we require $(\psi_k, \phi)_{\mathcal{H}^1(\mathbb{R}^n)} \to (\psi, \phi)_{\mathcal{H}^1(\mathbb{R}^n)}$ as $k \to \infty$ for all $\phi \in \mathcal{H}^1(\mathbb{R}^n)$, where the inner product of $\mathcal{H}^1(\mathbb{R}^n)$ is given by $(\psi, \phi)_{\mathcal{H}^1(\mathbb{R}^n)} = \int_{\mathbb{R}^n} \psi^* \phi + \int_{\mathbb{R}^n} \nabla \psi^* \cdot \nabla \phi$. Weak convergence on $\mathcal{H}^1(\mathbb{R}^n)$ implies weak convergence in the $L^2(\mathbb{R}^n)$ sense. A functional f is said to be weakly lower semi continuous if $\psi_k \to \psi$ implies $\liminf_{k\to\infty} f(\psi_k) \geq f(\psi)$. In particular, $\liminf_{k\to\infty} ||\psi_k||_{L^2(\mathbb{R}^n)} \geq ||\psi||_{L^2(\mathbb{R}^n)}$ if $\psi_k \to \psi$ weakly in $L^2(\mathbb{R}^n)$.

For a fixed particle number N, define the set of proper wavefunctions to be

$$W_N = \{ \psi \in \mathcal{H}^1(\mathbb{R}^{3N}) | \psi \text{ antisymmetric and } ||\psi||_{L^2(\mathbb{R}^{3N})} = 1 \}$$

and let the ground-state energy of H(v, A) be given by

$$e_0(v, A) = \inf\{\mathcal{E}_{v, A}(\psi) | \psi \in W_N\},$$

where

$$\mathcal{E}_{v,A}(\psi) = \sum_{k=1}^{N} \left(\int_{\mathbb{R}^{3N}} |(i\nabla_k - A(x_k))\psi|^2 + \int_{\mathbb{R}^{3N}} |\psi|^2 v(x_k) \right) + \sum_{1 \le k < l \le N} \int_{\mathbb{R}^{3N}} |\psi|^2 |x_k - x_l|^{-1}.$$

We will define the inner-product $(\psi, H(v, A)\psi)_{L^2}$ as the number $\mathcal{E}_{v,A}(\psi)$ for $\psi \in W_N$, even if $H(v, A)\psi \notin L^2$.

The particle and paramagnetic current density for $\psi \in W_N$ are computed from

$$\rho_{\psi}(x) = N \int_{\mathbb{R}^{3(N-1)}} |\psi(x, x_2, \dots, x_N)|^2 dx_2 \dots dx_N,$$

$$j_{\psi}^p(x) = N \operatorname{Im} \int_{\mathbb{R}^{3(N-1)}} \psi^*(x, x_2, \dots, x_N) \nabla_x \psi(x, x_2, \dots, x_N) dx_2 \dots dx_N,$$

respectively. We will use the notation $\psi \mapsto (\rho, j^p)$ to mean $\rho_{\psi} = \rho$ and $j_{\psi}^p = j^p$. Furthermore, we shall use the notation H_0 for the Hamiltonian H(v, A) when the potential terms are set to zero, i.e.,

$$(\psi, H_0 \psi)_{L^2} = \sum_{k=1}^N \int_{\mathbb{R}^{3N}} |\nabla_k \psi|^2 + \sum_{1 \le k < l \le N} \int_{\mathbb{R}^{3N}} |\psi|^2 |x_k - x_l|^{-1}.$$

Note that

$$\mathcal{E}_{v,A}(\psi) = (\psi, H(v,A)\psi)_{L^2} = (\psi, H_0\psi)_{L^2} + 2\int_{\mathbb{R}^3} j_{\psi}^p \cdot A + \int_{\mathbb{R}^3} \rho_{\psi}(v + |A|^2),$$

which follows from a direct computation.

B. N-representable DFT

A v-representable particle density is a density ρ that satisfies $\rho = \rho_{\psi}$ and where ψ is the ground-state of some H(v). (We will use the notation H(v) = H(v,0) and $e_0(v) = e_0(v,0)$ when not considering magnetic fields.) The set of N-representable particle densities is given by [8]

$$I_N = \left\{ \rho | \rho \ge 0, \int_{\mathbb{R}^3} \rho = N, \rho^{1/2} \in \mathcal{H}^1(\mathbb{R}^3) \right\}.$$

As demonstrated by Englisch and Englisch in [9], not every N-representable particle density is v-representable. For $\rho \in I_N$, the Levy-Lieb functional

$$F_{LL}(\rho) = \inf\{(\psi, H_0\psi)_{L^2} | \psi \in W_N, \psi \mapsto \rho\}$$

is well-defined. As was proven in [8] (Theorem 3.3), there exists a $\psi_0 \in W_N$ such that $F_{LL}(\rho) = (\psi_0, H_0\psi_0)_{L^2}$ and $\rho_{\psi_0} = \rho$. The functional $F_{LL}(\rho)$ extends the Hohenberg-Kohn functional to N-representable densities, and for the ground-state energy we have

$$e_0(v) = \inf \left\{ F_{LL}(\rho) + \int_{\mathbb{R}^3} \rho v | \rho \in I_N \right\}.$$

Note that the number $e_0(v)$ is well-defined for $v \in L^{3/2}(\mathbb{R}^3) + L^{\infty}(\mathbb{R}^3)$ even if H(v) does not have a ground-state. $(\int_{\mathbb{R}^3} \rho v$ is finite for all $\rho \in I_N$, since $I_N \subset L^1(\mathbb{R}^3) \cap L^3(\mathbb{R}^3)$, see [8].)

C. N-representable CDFT

A density pair (ρ, j^p) is said to be v-representable if there exists a ψ that is the groundstate of some Hamiltonian H(v, A) such that $\rho = \rho_{\psi}$ and $j^p = j_{\psi}^p$. We denote this set of densities \mathcal{A}_N , i.e.,

$$\mathcal{A}_N = \{(\rho, j^p) | \text{there exists a } H(v, A) \text{ with ground-state } \psi \text{ such that } \psi \mapsto (\rho, j^p) \}.$$

Now, assume that $H(v_1, A_1)$ and $H(v_2, A_2)$ have the ground-states ψ and ϕ , respectively. Then from Theorem 9 in [5], if $\psi \mapsto (\rho, j^p)$ and $\phi \mapsto (\rho, j^p)$, it follows that $\psi = \text{const. } \phi$. For $(\rho, j^p) \in \mathcal{A}_N$, let ψ_{ρ, j^p} denote the ground-state of some H(v, A) such that $\psi \mapsto (\rho, j^p)$. Then the generalized Hohenberg-Kohn functional

$$F_{HK}(\rho, j^p) = (\psi_{\rho, j^p}, H_0 \psi_{\rho, j^p})_{L^2}$$

is well-defined on A_N . Furthermore (Theorem 2 in [10]),

$$e_0(v,A) = \min \left\{ F_{HK}(\rho, j^p) + 2 \int_{\mathbb{R}^3} j^p \cdot A + \int_{\mathbb{R}^3} \rho(v + |A|^2) \Big| (\rho, j^p) \in \mathcal{A}_N \right\}$$

for $(v, A) \in V_N$, where

$$V_N = \{(v, A) | H(v, A) \text{ has a unique ground-state} \}.$$

However, a $\psi \in W_N$ may be such that $(\rho_{\psi}, j_{\psi}^p) \notin \mathcal{A}_N$. From Proposition 3 in [10], $\psi \in W_N$ implies that $\psi \mapsto (\rho, j^p) \in Y_N$, where

$$Y_N = \left\{ (\rho, j^p) | \rho \ge 0, \int_{\mathbb{R}^3} \rho = N, \rho^{1/2} \in \mathcal{H}^1(\mathbb{R}^3), j^p \in (L^1(\mathbb{R}^3))^3, \int_{\mathbb{R}^3} |j^p|^2 \rho^{-1} < \infty \right\}.$$

The set Y_N is referred to as the set of N-representable density pairs (ρ, j^p) . It is a convex set and $\mathcal{A}_N \subsetneq Y_N$ (Proposition 4 in [10]). For $(\rho, j^p) \in Y_N$, define as in [10]

$$Q(\rho, j^p) = \inf\{(\psi, H_0\psi)_{L^2} | \psi \in W_N, \psi \mapsto (\rho, j^p)\}.$$

The functional $Q(\rho, j^p)$ is the generalization of the Levy-Lieb functional $F_{LL}(\rho)$. It also depends on the paramagnetic current density j^p . The functional $Q(\rho, j^p)$ inherits many properties of $F_{LL}(\rho)$: by Theorem 5 and Theorem 6 in [10], we have (i) $Q(\rho, j^p) = F_{HK}(\rho, j^p)$ for $(\rho, j^p) \in \mathcal{A}_N$, (ii) there exists a $\psi_m \in W_N$ such that $Q(\rho, j^p) = (\psi_m, H_0 \psi_m)_{L^2}$ and where $\psi_m \mapsto (\rho, j^p)$, and (iii)

$$e_0(v,A) = \inf \Big\{ Q(\rho,j^p) + 2 \int_{\mathbb{R}^3} j^p \cdot A + \int_{\mathbb{R}^3} \rho(v+|A|^2) \Big| (\rho,j^p) \in Y_N \Big\}.$$

In [3], $F_{LL}(\rho)$ was used to obtain a rigorous Kohn-Sham theory for N-representable densities. Before generalizing this to CDFT formulated with j^p , we shall discuss the following question raised in [3]: since a $\psi_0 \in W_N$ exists such that $F_{LL}(\rho) = (\psi_0, H_0\psi_0)_{L^2}$ and $\psi_0 \mapsto \rho$, does ψ_0 satisfy any Schrödinger equation, i.e., is there a v(x) such that $H(v)\psi = e\psi$?

III. CHARACTERIZATION OF V-REPRESENTABLE PARTICLE DENSITIES

We start be stating the mentioned result of Lieb (Theorem 3.3 in [8]) for the functional $F_{LL}(\rho)$.

Theorem 1 There exists a ψ_0 in W_N such that for $\rho \in I_N$, $F_{LL}(\rho) = (\psi_0, H_0 \psi_0)_{L^2}$ and $\rho_{\psi_0} = \rho$.

Let $\rho \in I_N$. In light of Theorem 1, if the minimizer ψ_0 would be the ground-state of some Hamiltonian H(v), then ρ would be v-representable. However, since the v-representable densities are a proper subset of the N-representable ones [9], there exists $\rho \in I_N$ such that the corresponding minimizer ψ_0 is not the ground-state of any Hamiltonian H(v). Also note that, if ρ is v-representable, then the minimizer ψ_0 is also the ground-state associated with ρ . This so since if ρ is v-representable, then by the definition of the minimizer ψ_0 , we have

$$(\psi_0, H_0\psi_0)_{L^2} + \int_{\mathbb{R}^3} \rho v = e_0(v)$$

for some v, i.e., ψ_0 is the ground-state of H(v). (A similar result holds for a minimizer of $Q(\rho, j^p)$, see Proposition 5.)

Now, let N = 1. Note the following: $(\psi, H_0 \psi)_{L^2} = \int_{\mathbb{R}^3} |\nabla \psi|^2 dx \ge \int_{\mathbb{R}^3} |\nabla \psi|^2 dx$. Thus, for $F_{LL}(\rho)$, it is enough to minimize over the non-negative functions of W_1 , i.e.,

$$F_{LL}(\rho) = \inf \left\{ \int_{\mathbb{R}^3} |\nabla \psi|^2 dx | \psi \in W_1, \psi \ge 0, \psi^2 = \rho \right\}.$$

We now give criteria when ψ_0 in Theorem 1 is an eigenfunction of some H(v).

Proposition 2 (i) Let N=1 and $\rho \in I_1$ be such that ψ_0 fulfills $\Delta \psi_0 \in L^2(\mathbb{R}^3)$ and $\psi_0 \neq 0$ almost everywhere (a.e.), where $\psi_0 \geq 0$ minimizes $\int_{\mathbb{R}^3} |\nabla \psi|^2$ subject to the constraint $\psi^2 = \rho$. Then there exists a $\phi_0 \in L^2(\mathbb{R}^3)$ and a constant e such that, with $v-e=\phi_0/\rho^{1/2}$, ψ_0 satisfies

$$-\Delta\psi_0 + v\psi_0 = e\psi_0,$$

and where $\int_{\mathbb{R}^3} v |\psi_0|^2 > -\infty$.

(ii) For N=1, there exists $\rho_0 \in I_1$ such that $\Delta \psi_0 \notin L^2(\mathbb{R}^3)$, and $-\Delta \psi_0 + v\psi_0 = 0$ implies $\int_{\mathbb{R}^3} v |\psi_0|^2 = -\infty$.

Proof. By assumption, $\psi_0 > 0$ a.e. and $\psi_0 = \rho^{1/2}$. Now, set $\phi_0 = \Delta \psi_0$, which is in $L^2(\mathbb{R}^3)$. Then with $v - e = \phi_0/\rho^{1/2}$ the conclusion of the first part follows, since

$$\int_{\mathbb{R}^3} v |\psi_0|^2 = \int_{\mathbb{R}^3} \phi_0 \rho^{1/2} + e \ge -||\phi_0||_{L^2} + e.$$

For the second part, set, for small $|x_1|$, $\rho_0(x) = \rho_1(x_1)\tilde{\rho}(x_2, x_3)$, where $\tilde{\rho}(x_2, x_3)$ is regular and $\rho_1(x_1) = (a + b|x_1|^{\varepsilon+1/2})^2$, a, b < 0 and $0 < \varepsilon < 1/2$. Then $\Delta \psi_0 \notin L^2(\mathbb{R}^3)$. Furthermore, $-\Delta \psi_0 + v\psi_0 = 0$ implies $\int_{\mathbb{R}^3} v|\psi_0|^2 = -\infty$. (The density ρ_0 is the counterexample of Englisch and Englisch that shows that not every N-representable density is v-representable, see [9].)

Note that ψ_0 is not proven to be the ground-state of $-\Delta + v$. However, we have

Corollary 3 Let ρ , ψ_0 and ϕ_0 be as in Proposition 2 (i). In addition, assume that $\phi_0 \leq C\rho^{1/2}$ for some constant C and that $\rho^{-1} \in L^1_{loc}(\mathbb{R}^3)$. Then ψ_0 is the ground-state of $-\Delta + v$.

Proof. From Proposition 2, we know that $-\Delta\psi_0 + v\psi_0 = e\psi_0$, where $v = \phi_0/\rho^{1/2} + e$. By Schwarz's inequality, it follows that $v \in L^1_{loc}(\mathbb{R}^3)$. Since v is also bounded above, we have by Corollary 11.9 in [11] that $\psi_0 > 0$ is the ground-state of $-\Delta + v$.

We can thus conclude with the following characterization: if $\rho \in I_1$ satisfies (i) $\rho > 0$ (a.e.), (ii) $\Delta \rho^{1/2} \in L^2(\mathbb{R}^3)$ and bounded above by a constant times $\rho^{1/2}$, and (iii) $\rho^{-1} \in L^1_{loc}$, then ρ is v-representable.

IV. RIGOROUS KOHN-SHAM THEORY FOR CDFT

By means of the Levy-Lieb-type density functional $Q(\rho, j^p)$ we can formulate a rigorous N-representable Kohn-Sham approach for CDFT as that of Ref. [3] for DFT. Now, fix the particle number N. We say that a wavefunction $\phi \in W_N$ is a determinant if there exist N orthonormal one-particle functions f_k such that

$$\phi(x_1,\ldots,x_N) = (N!)^{-1/2} \det[f^k(x_l)]_{k,l}.$$

Let the space of all normalized determinants of finite kinetic energy be denoted W_S , i.e.,

$$W_S = \{ \phi | \phi \text{ is a determinant}, ||\phi||_{L^2(\mathbb{R}^{3N})} = 1, (\phi, K\phi)_{L^2(\mathbb{R}^{3N})} < \infty \},$$

where $K = -\sum_{k=1}^{N} \Delta_k$. Note that, in particular, for a $\phi \in W_S$, we have $\rho_{\phi} = \sum_{k=1}^{N} |f^k|^2$ and

$$(\phi, K\phi)_{L^2(\mathbb{R}^{3N})} = \sum_{k=1}^N \int_{\mathbb{R}^3} |\nabla f^k|^2 dx.$$

Thus, $||\phi||_{L^2(\mathbb{R}^{3N})} = 1$ and $(\phi, K\phi)_{L^2(\mathbb{R}^{3N})} < \infty$ are equivalent to $f^k \in \mathcal{H}^1(\mathbb{R}^3)$ for all k. Also note that a $\psi \in W_N$ is not in general an element of W_S , i.e., $W_S \subsetneq W_N$.

Furthermore, define, for a non-interacting system, the non-interacting Hamiltonian

$$H'(v, A) = \sum_{k=1}^{N} ((i\nabla_k - A(x_k))^2 + v(x_k)).$$

The non-interacting ground-state energy is then given by

$$e_0'(v, A) = \inf \{ \mathcal{E}_{v, A}'(\psi) | \psi \in W_N \},$$

where $\mathcal{E}'_{v,A}(\psi)$ is given by the relation

$$\mathcal{E}'_{v,A}(\psi) + \sum_{1 \le k \le l \le N} \int_{\mathbb{R}^{3N}} |\psi|^2 |x_k - x_l|^{-1} = \mathcal{E}_{v,A}(\psi).$$

This motivates: set, for $(\rho, j^p) \in Y_N$,

$$Q'(\rho, j^p) = \inf\{(\psi, K\psi)_{L^2} | \psi \in W_N, \psi \mapsto (\rho, j^p)\}.$$

For $Q(\rho, j^p)$ and $Q'(\rho, j^p)$ we have the following.

Theorem 4 Fix $(\rho, j^p) \in Y_N$, then (i) there exists a $\psi_m \in W_N$ such that $\psi_m \mapsto (\rho, j^p)$ and $Q(\rho, j^p) = (\psi_m, H_0 \psi_m)_{L^2}$, and (ii) there exists a $\psi'_m \in W_N$ such that $\psi'_m \mapsto (\rho, j^p)$ and $Q'(\rho, j^p) = (\psi'_m, K \psi'_m)_{L^2}$.

Proof. Part (i) above is just Theorem 5 in [10]. However, for (ii), we can use the same proof. For the sake of completeness we include the proof in [10] here applied to $Q'(\rho, j^p)$.

Let $\{\psi^j\}_{j=1}^{\infty}$ be a minimizing sequence, i.e., $\psi^j \in W_N$, $\psi^j \mapsto (\rho, j^p)$ and

$$\lim_{j \to \infty} (\psi^j, K\psi^j)_{L^2} = Q'(\rho, j^p).$$

Since $\{\psi^j\}_{j=1}^{\infty}$ is bounded in $\mathcal{H}^1(\mathbb{R}^{3N})$, by the Banach-Alaoglu theorem there exists a subsequence and a $\psi'_m \in \mathcal{H}^1(\mathbb{R}^{3N})$ such that $\psi^{j_k} \rightharpoonup \psi'_m$ weakly in $\mathcal{H}^1(\mathbb{R}^{3N})$ as $k \to \infty$. Since the functional $\psi \mapsto (\psi, K\psi)_{L^2}$ is weakly lower semi continuous, we know that

$$(\psi'_m, K\psi'_m)_{L^2} \leq Q'(\rho, j^p).$$

However, it remains to prove that $\psi'_m \mapsto (\rho, j^p)$. In the proof of Theorem 3.3 in [8], it is shown that $\psi^{j_k} \to \psi'_m$ in $L^2(\mathbb{R}^{3N})$ and $\psi'_m \mapsto \rho$. Now, let g be the characteristic function of any measurable set in \mathbb{R}^3 . For l = 1, 2, 3 and $k = 1, 2, \ldots$, let

$$I_l(k) = \left| \int_{\mathbb{R}^{3N}} [(\psi^{j_k})^* \partial_l \psi^{j_k} - (\psi'_m)^* \partial_l \psi'_m] g \right|.$$

Then

$$I_{l}(k) \leq \left| \int_{\mathbb{R}^{3N}} (\psi^{j_{k}} - \psi'_{m})^{*} (\partial_{l} \psi^{j_{k}}) g \right| + \left| \int_{\mathbb{R}^{3N}} (\psi'_{m})^{*} (\partial_{l} \psi^{j_{k}} - \partial_{l} \psi'_{m}) g \right|$$

$$\leq ||\psi^{j_{k}} - \psi'_{m}||_{L^{2}} ||(\partial_{l} \psi^{j_{k}}) g||_{L^{2}} + \left| \int_{\mathbb{R}^{3N}} (\psi'_{m} g^{*})^{*} (\partial_{l} \psi^{j_{k}} - \partial_{l} \psi'_{m}) \right|.$$

Thus $I_l(k)$ tends to zero as $k \to \infty$ (because $\psi^{j_k} \to \psi'_m$ in $L^2(\mathbb{R}^{3N})$ -norm and $\psi^{j_k} \rightharpoonup \psi'_m$ weakly in $\mathcal{H}^1(\mathbb{R}^{3N})$ as $k \to \infty$). Since $\psi^{j_k} \mapsto j^p$ for all k, we have $\int_{\mathbb{R}^3} (j^p)_l g = \int_{\mathbb{R}^3} (j^p_{\psi'_m})_l g$, i.e., $j^p_{\psi'_m}(x) = j^p(x)$ a.e.

Proposition 5 Assume that $(\rho, j^p) \in \mathcal{A}_N$, i.e., there exists a H(v, A) with ground-state ψ such that $\psi \mapsto (\rho, j^p)$. Then the minimizer ψ_m is the ground-state of H(v, A).

Proof. Since $\psi \mapsto (\rho, j^p)$, we have $(\psi, H_0 \psi)_{L^2} \ge (\psi_m, H_0 \psi_m)_{L^2}$. The conclusion then follows from

$$e_0(v,A) \le (\psi_m, H(v,A)\psi_m)_{L^2} = (\psi_m, H_0\psi_m)_{L^2} + 2\int_{\mathbb{R}^3} j^p \cdot A + \int_{\mathbb{R}^3} \rho(v+|A|^2)$$

$$\le (\psi, H_0\psi)_{L^2} + 2\int_{\mathbb{R}^3} j^p \cdot A + \int_{\mathbb{R}^3} \rho(v+|A|^2) = (\psi, H(v,A)\psi)_{L^2} = e_0(v,A). \blacksquare$$

Note that when H_0 is replaced by K, $Q'(\rho, j^p)$ is the minimal kinetic energy for $\psi \in W_N$ such that $\rho_{\psi} = \rho$ and $j_{\psi}^p = j^p$. Next we will introduce another kinetic energy density functional.

A. Non-interacting kinetic energy density functional

Set, for $(\rho, j^p) \in Y_N$,

$$T_{\text{det}}(\rho, j^p) = \inf\{(\phi, K\phi)_{L^2} | \phi \in W_S, \phi \mapsto (\rho, j^p)\}.$$

For $(\rho, j^p) \in Y_N$, we remark that the set $\{\phi \in W_S | \phi \mapsto (\rho, j^p)\}$ is not empty, at least when $N \geq 4$. This follows from the determinant construction in [12]. However, for all N, the set $\{\phi \in W_S | \phi \mapsto (\rho, j^p), \nabla \times (j^p/\rho) = 0\}$ is non-empty (see [10, 12]).

We have that $T_{\text{det}}(\rho, j^p) \geq Q'(\rho, j^p)$ on Y_N . Now, let the set of non-interacting v-representable densities be denoted \mathcal{A}'_N ,

$$\mathcal{A}'_N = \{(\rho, j^p) | H'(v, A) \text{ has a unique ground-state}\}.$$

If $(\rho, j^p) \in \mathcal{A}'_N$, by the same argument as in the proof of Proposition 5, we can conclude that ψ'_m is the ground-state of some H'(v, A). Clearly, ψ'_m is in this case a determinant. Thus, $T_{\text{det}}(\rho, j^p) = Q'(\rho, j^p)$ on \mathcal{A}'_N .

An important property of $T_{\text{det}}(\rho, j^p)$ is that the infimum actually is a minimum. For the proof, we need the following:

(i) For $k=1,\ldots,N$, assume that $f_j^k\to f^k$ in L^2 -norm as $j\to\infty$ and for each j, $(f_j^k,f_j^l)_{L^2}=\delta_{kl}$. Then f^1,\ldots,f^N are orthonormal. This so since

$$(f^k, f^l)_{L^2} = \lim_{j \to \infty} (f_j^k, f^l)_{L^2} = \lim_{j \to \infty} [(f_j^k, f^l - f_j^l)_{L^2} + (f_j^k, f_j^l)_{L^2}] = \delta_{kl},$$

where we used that $|(f_j^k, f^l - f_j^l)_{L^2}| \le ||f_j^k||_{L^2}||f^l - f_j^l||_{L^2} \to 0$ as $j \to \infty$.

(ii) If $f_j
ightharpoonup f$ weakly in L^2 as $j
ightharpoonup \infty$ and $||f_j||_{L^2}
ightharpoonup ||f||_{L^2}$ as $j
ightharpoonup \infty$, then $f_j
ightharpoonup f$ in L^2 -norm as $j
ightharpoonup \infty$. (This is an elementary fact and can be checked by expanding $||f_j - f||_{L^2}^2 = (f_j - f, f_j - f)_{L^2}$.)

Theorem 6 Let $(\rho, j^p) \in Y_N$. If N < 4 we also assume $\nabla \times (j^p/\rho) = 0$. Then there exists a determinant ϕ_m such that $\phi_m \mapsto (\rho, j^p)$ and $T_{det}(\rho, j^p) = (\phi_m, K\phi_m)_{L^2}$.

Proof. Fix $(\rho, j^p) \in Y_N$ and let $\{D^j\}_{j=1}^{\infty} \subset W_S$ be a sequence of minimizing determinants, i.e., $D^j \mapsto (\rho, j^p)$ and $\lim_{j \to \infty} (D^j, KD^j)_{L^2} = T_{\text{det}}(\rho, j^p)$. From the proof of Theorem 4, there exists a subsequence D^{j_n} and a $\phi_m \in W_N$ such that $\phi_m \mapsto (\rho, j^p)$,

$$T_{\text{det}}(\rho, j^p) = (\phi_m, K\phi_m)_{L^2}$$

and $D^{j_n} \to \phi_m$ in L^2 -norm. It remains to show that $\phi_m \in W_S$. To meet that end, let

$$D^{j}(x_{1},...,x_{N}) = (N!)^{-1/2} \det[f_{j}^{k}(x_{l})]_{k,l},$$

where for each j the N one-particle functions f_j^k are orthonormal. By the Banach-Alaoglu theorem, there exist N functions f^k such that (for a subsequence) $f_j^k \rightharpoonup f^k$ weakly in L^2 as $j \to \infty$. We furthermore claim that f^1, \ldots, f^N are orthonormal. If we could prove that $f_j^k \to f^k$ in L^2 -norm, it would follow that $(f^k, f^l)_{L^2} = \delta_{kl}$.

We shall prove $f_j^k \to f^k$ by demonstrating that $||f_j^k||_{L^2} \to ||f^k||_{L^2}$. This together with the fact that $f_j^k \to f^k$ weakly in L^2 gives the desired result. Let $\varepsilon > 0$ and choose a characteristic function χ such that $\int_{\mathbb{R}^3} \rho(1-\chi) < \varepsilon$. Since for each j, $D^j \mapsto \rho$, we have for each k,

$$\int_{\mathbb{R}^3} |f_j^k|^2 (1-\chi) \le \sum_{k=1}^N \int_{\mathbb{R}^3} |f_j^k|^2 (1-\chi) = \int_{\mathbb{R}^3} \rho(1-\chi) < \varepsilon.$$

By the Rellich-Kondrachov theorem, we can choose a subsequence such that $\chi f_{j_n}^k \to \chi f^k$ in L^2 -norm. But this implies

$$\int_{\mathbb{R}^3} |f^k|^2 \ge \int_{\mathbb{R}^3} \chi |f^k|^2 = \lim_{n \to \infty} \int_{\mathbb{R}^3} \chi |f_{j_n}^k|^2 \ge 1 - \varepsilon.$$

Conversely, by the lower semi continuity of the L^2 -norm, $1 = \liminf_{j \to \infty} ||f_j^k||_{L^2} \ge ||f^k||_{L^2}$, and we have $||f^k||_{L^2} = 1$.

Returning to the fact that $f_{j_n}^k \rightharpoonup f^k$ weakly in L^2 , we note that $\Pi_{k=1}^N f_{j_n}^k(x_k) \rightharpoonup \Pi_{k=1}^N f^k(x_k)$ weakly in $L^2(\mathbb{R}^{3N})$ (since product-functions are dense in $L^2(\mathbb{R}^{3N})$). But then

$$D^{j_n} \rightharpoonup (N!)^{-1/2} \det[f^k(x_l)]_{k,l},$$

where f^1, \ldots, f^N are orthonormal. However, since $D^{j_n} \to \phi_m$, we have $\phi_m \in W_S$.

B. N-representable Kohn-Sham theory

In the Kohn-Sham approach [2], a non-interacting system is introduced that has the same ground-state density as the fully interacting system. The idea is then to use an element of W_S , i.e., a determinant, to compute the ground-state density. On \mathcal{A}'_N , the (generalized) Kohn-Sham density functional $T_{KS}(\rho, j^p)$ satisfies

$$T_{KS}(\rho, j^p) = T_{\text{det}}(\rho, j^p) = Q'(\rho, j^p).$$

Moreover, T_{KS} defines an exchange-correlation functional $E_{xc}(\rho, j^p)$ on $\mathcal{A}_N \cap \mathcal{A}'_N$ according to

$$E_{xc}(\rho, j^p) = F_{HK}(\rho, j^p) - \frac{1}{2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{\rho(x)\rho(y)}{|x - y|} dx dy - T_{KS}(\rho, j^p).$$

Now, to obtain an N-representable Kohn-Sham scheme, define two functionals on W_S ,

$$\mathcal{G}_K(\phi) = \inf\{(f, Kf)_{L^2} | f \in W_S, f \mapsto (\rho_\phi, j_\phi^p)\},\$$

$$\mathcal{G}_{H_0}(\phi) = \inf\{(f, H_0 f)_{L^2} | f \in W_N, f \mapsto (\rho_\phi, j_\phi^p)\}.$$

Note that, by Theorem 4 and Theorem 6, there exists a $\psi_m \in W_N$ and a $\phi_m \in W_S$ such that $\mathcal{G}_{H_0}(\phi) = (\psi_m, H_0\psi_m)_{L^2}$ and $\mathcal{G}_K(\phi) = (\phi_m, K\phi_m)_{L^2}$ and where $\psi_m, \phi_m \mapsto (\rho_\phi, j_\phi^p)$. Furthermore, we can use the existence of the minimizers ψ_m and ϕ_m and define, for $\phi \in W_S$,

$$\Delta T(\phi) = (\psi_m, K\psi_m)_{L^2} - (\phi_m, K\phi_m)_{L^2},$$

$$E_{xc}^W(\phi) = (\psi_m, \sum_{1 \le k < l \le N} |x_k - x_l|^{-1} \psi_m)_{L^2} - \frac{1}{2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{\rho_\phi(x) \rho_\phi(y)}{|x - y|} dx dy.$$

On W_S , we now introduce the following energy functional

$$\mathcal{G}_{v,A}(\phi) = (\phi, K\phi)_{L^2} + \Delta T(\phi) + 2 \int_{\mathbb{R}^3} j_{\phi}^p \cdot A + \int_{\mathbb{R}^3} \rho_{\phi}(v + |A|^2) + E_{xc}^W(\phi) + \frac{1}{2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{\rho_{\phi}(x)\rho_{\phi}(y)}{|x - y|} dx dy.$$

We then have

Theorem 7 Assume that H(v, A) has a unique ground-state ψ_0 . Let $e_0(v, A)$, ρ_0 and j_0^p denote the ground-state energy, ground-state particle density and ground-state paramagnetic current density, respectively. If N < 4 we assume that $\nabla \times (j_0^p/\rho_0) = 0$. Then

$$e_0(v, A) = \inf \{ \mathcal{G}_{v,A}(\phi) | \phi \in W_S \} = \mathcal{G}_{v,A}(\phi_m)$$

for some $\phi_m \in W_S$. Moreover, $\rho_{\phi_m} = \rho_0$ and $j_{\phi_m}^p = j_0^p$, i.e., the ground-state densities can be computed from the determinant ϕ_m that minimizes $\mathcal{G}_{v,A}$.

Proof. First note, for any $\phi \in W_S$, we have

$$\mathcal{G}_{v,A}(\phi) = (\phi, K\phi)_{L^{2}} + ((\psi_{m}, K\psi_{m})_{L^{2}} - (\phi_{m}, K\phi_{m})_{L^{2}})
+ 2 \int_{\mathbb{R}^{3}} j_{\phi}^{p} \cdot A + \int_{\mathbb{R}^{3}} \rho_{\phi}(v + |A|^{2}) + (\psi_{m}, \sum_{1 \leq k < l \leq N} |x_{k} - x_{l}|^{-1}\psi_{m})_{L^{2}}
\geq (\psi_{m}, (K + \sum_{1 \leq k < l \leq N} |x_{k} - x_{l}|^{-1})\psi_{m})_{L^{2}} + 2 \int_{\mathbb{R}^{3}} j_{\phi}^{p} \cdot A + \int_{\mathbb{R}^{3}} \rho_{\phi}(v + |A|^{2})
= \mathcal{E}_{v,A}(\psi_{m}) \geq e_{0}(v, A),$$

where we used that $(\phi, K\phi)_{L^2} - (\phi_m, K\phi_m)_{L^2} \ge 0$ and $\psi_m \mapsto (\rho_\phi, j_\phi^p)$. In the next step, we want to show that there exists a $\phi_0 \in W_S$ such that $\mathcal{G}_{v,A}(\phi_0) = e_0(v,A)$ and $\phi_0 \mapsto (\rho_0, j_0^p)$.

Let $\phi \in W_S$ be a determinant such that $\phi \mapsto (\rho_0, j_0^p)$ (if N < 4, we need the assumption $\nabla \times (j_0^p/\rho_0) = 0$). By Theorem 6, we then have

$$\mathcal{G}_K(\phi) = T_{\text{det}}(\rho_0, j_0^p) = (\phi_m, K\phi_m)_{L^2},$$

for some $\phi_m \in W_S$. Note that ϕ_m is a determinant such that $\phi_m \mapsto (\rho_0, j_0^p)$ and

$$\mathcal{G}_K(\phi_m) = (\phi_{m,m}, K\phi_{m,m})_{L^2} = (\phi_m, K\phi_m)_{L^2}.$$

Furthermore,

$$\mathcal{G}_{H_0}(\phi_m) = Q(\rho_0, j_0^p) = (\psi_m, H_0 \psi_m)_{L^2},$$

for some $\psi_m \in W_N$, which follows from Theorem 4. Note that $\psi_m \mapsto (\rho_0, j_0^p) = (\rho_{\phi_m}, j_{\phi_m}^p)$. We have,

$$\begin{aligned} e_0(v,A) &= (\psi_m, H(v,A)\psi_m)_{L^2} \\ &= (\psi_m, H_0\psi_m)_{L^2} + 2\int_{\mathbb{R}^3} j_0^p \cdot A + \int_{\mathbb{R}^3} \rho_0(v + |A|^2) \\ &= (\psi_m, K\psi_m)_{L^2} + (\psi_m, \sum_{1 \le k < l \le N} |x_k - x_l|^{-1} \psi_m)_{L^2} + 2\int_{\mathbb{R}^3} j_{\phi_m}^p \cdot A + \int_{\mathbb{R}^3} \rho_{\phi_m}(v + |A|^2), \end{aligned}$$

where the first equality follows from Proposition 5. Since

$$\Delta T(\phi_m) = (\psi_m, K\psi_m)_{L^2} - (\phi_m, K\phi_m)_{L^2}$$

and

$$E_{xc}^{W}(\phi_m) = (\psi_m, \sum_{1 \le k \le l \le N} |x_k - x_l|^{-1} \psi_m)_{L^2} - \frac{1}{2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{\rho_{\phi_m}(x) \rho_{\phi_m}(y)}{|x - y|} dx dy,$$

it follows that

$$e_{0}(v,A) = (\phi_{m}, K\phi_{m})_{L^{2}} + 2\int_{\mathbb{R}^{3}} j_{\phi_{m}}^{p} \cdot A + \int_{\mathbb{R}^{3}} \rho_{\phi_{m}}(v + |A|^{2})$$

$$+ \frac{1}{2} \int_{\mathbb{R}^{3}} \int_{\mathbb{R}^{3}} \frac{\rho_{\phi_{m}}(x)\rho_{\phi_{m}}(y)}{|x - y|} dxdy + E_{xc}^{W}(\phi_{m}) + \Delta T(\phi_{m}) = \mathcal{G}_{v,A}(\phi_{m}). \quad \blacksquare$$

Remarks. (i) Any density pair (ρ, j^p) computed from a $\phi \in W_S$ is N-representable, but not necessarily (non-interacting) v-representable. So Theorem 7 establishes a Kohn-Sham approach for N-representable densities (whereas T_{KS} is only defined on \mathcal{A}'_N).

- (ii) Recall that no Hohenberg-Kohn theorem can exist for CDFT formulated with the paramagnetic current density. On the other hand, since ρ and j^p determine the ground-state, the Hohenberg-Kohn variational principle continues to hold for CDFT formulated with these densities. However, the N-representable Kohn-Sham approach outlined here does not use any variational principle for densities. Instead, the approach relies on the existence of minimizers for certain functionals.
- (iii) If we set $\phi(x_1, \ldots, x_N) = (N!)^{-1/2} \det[f^k(x_l)]_{k,l}$ and define on $(\mathcal{H}^1(\mathbb{R}^3))^N$ the functional

$$\mathcal{E}(f^{1},\dots,f^{N}) = \sum_{k=1}^{N} \int_{\mathbb{R}^{3}} |\nabla f^{k}|^{2} + 2 \sum_{k=1}^{N} \int_{\mathbb{R}^{3}} \operatorname{Im}(\overline{f}^{k} \nabla f^{k}) \cdot A + \sum_{k=1}^{N} \int_{\mathbb{R}^{3}} |f^{k}|^{2} (v + |A|^{2}) + \frac{1}{2} \sum_{k,l=1}^{N} \int_{\mathbb{R}^{3}} \int_{\mathbb{R}^{3}} \frac{|f^{k}(x)|^{2} |f^{l}(y)|^{2}}{|x - y|} dx dy + E_{xc},$$

where $E_{xc} = \Delta T + E_{xc}^W$, we can obtain the usual Kohn-Sham equations by minimizing $\mathcal{E}(f^1, \ldots, f^N)$ subject to the constraint $(f^k, f^l)_{L^2} = \delta_{kl}$.

ACKNOWLEDGMENTS

The author is thankful to Michael Benedicks for useful comments and discussions.

- * andrela@math.kth.se; Department of Mathematics, KTH Royal Institute of Technology, Sweden
- [1] P. Hohenberg and W. Kohn, Phys. Rev. 136, B864 (Nov 1964), http://link.aps.org/doi/10.1103/PhysRev.136.B864
- [2] W. Kohn and L. J. Sham, Phys. Rev. 140, A1133 (Nov 1965), http://link.aps.org/doi/10.1103/PhysRev.140.A1133
- [3] N. Hadjisavvas and A. Theophilou, Phys. Rev. A 30, 2183 (Nov 1984), http://link.aps.org/doi/10.1103/PhysRevA.30.2183
- [4] K. Capelle and G. Vignale, Phys. Rev. B **65**, 113106 (Feb 2002), http://link.aps.org/doi/10.1103/PhysRevB.65.113106
- [5] A. Laestadius and M. Benedicks, To appear in Int. Jour. Quant. Chem.
- [6] E. I. Tellgren, S. Kvaal, E. Sagvolden, U. Ekström, A. M. Teale, and T. Helgaker, Phys. Rev. A 86, 062506 (Dec 2012), http://link.aps.org/doi/10.1103/PhysRevA.86.062506
- [7] G. Vignale and M. Rasolt, Phys. Rev. Lett. 59, 2360 (Nov 1987), http://link.aps.org/doi/10.1103/PhysRevLett.59.2360
- [8] E. H. Lieb, Int. Jour. Quant. Chem. 24, 243 (1983), http://dx.doi.org/10.1002/qua.560240302
- [9] H. Englisch and R. Englisch, Physica A **121**, 253 (1983)
- [10] A. Laestadius, arXiv abs/1404.0825 (2014), http://arxiv.org/abs/1404.0825
- [11] E. H. Lieb and M. Loss, Analysis (American Mathematical Society, Providence, Rhode Island, 2001)
- [12] E. H. Lieb and R. Schrader, Phys. Rev. A 88, 032516 (Sep 2013), http://link.aps.org/doi/10.1103/PhysRevA.88.032516