

COMPLEX HADAMARD MATRICES CONTAINED IN A BOSE–MESNER ALGEBRA

TAKUYA IKUTA AND AKIHIRO MUNEMASA

Dedicated to Chris Godsil on the occasion of his 65th birthday

ABSTRACT. A complex Hadamard matrix is a square matrix H with complex entries of absolute value 1 satisfying $HH^* = nI$, where $*$ stands for the Hermitian transpose and I is the identity matrix of order n . In this paper, we first determine the image of a certain rational map from the d -dimensional complex projective space to $\mathbb{C}^{d(d+1)/2}$. Applying this result with $d = 3$, we give constructions of complex Hadamard matrices, and more generally, type-II matrices, in the Bose–Mesner algebra of a certain 3-class symmetric association scheme. In particular, we recover the complex Hadamard matrices of order 15 found by Ada Chan. We compute the Haagerup sets to show inequivalence of resulting type-II matrices, and determine the Nomura algebras to show that the resulting matrices are not decomposable into generalized tensor products.

1. INTRODUCTION

A complex Hadamard matrix is a square matrix H with complex entries of absolute value 1 satisfying $HH^* = nI$, where $*$ stands for the Hermitian transpose and I is the identity matrix of order n . They are the natural generalization of real Hadamard matrices. Complex Hadamard matrices appear frequently in various branches of mathematics and quantum physics.

A type-II matrix, or an inverse orthogonal matrix, is a square matrix W with nonzero complex entries satisfying $WW^{(-)\top} = nI$, where $W^{(-)}$ denotes the entrywise inverse of W . Obviously, a complex Hadamard matrix is a type-II matrix. Two type-II matrices W_1 and W_2 are said to be equivalent if there exist diagonal matrices D, D' with nonzero

Date: April 21, 2015.

2010 Mathematics Subject Classification. 05E30, 05B34.

Key words and phrases. association scheme, complex Hadamard matrix, type-II matrix.

The work of T.I. was supported by JSPS KAKENHI grant number 25400215, and that of A.M. was supported by JSPS KAKENHI grant number 26400003.

complex diagonal entries, and permutation matrices T, T' , such that $DW_1D' = TW_2T'$ holds.

Complete classifications of complex Hadamard matrices, and of type-II matrices are only available up to order $n = 5$ (see [7, 14, 10]). Although it is shown by Craigen [7] that there are uncountably many equivalence classes of complex Hadamard matrices of order n whenever n is a composite number, some type-II matrices are more closely related to combinatorial objects than the others. Szollosi [16] used design theoretical methods to construct complex Hadamard matrices. Strongly regular graphs were used to construct type-II matrices in [5, 6]. See [15] for a generalization. In this paper, we construct type-II matrices and complex Hadamard matrices in the Bose–Mesner algebra of a certain 3-class symmetric association scheme. In particular, we recover the complex Hadamard matrix of order 15 found in [4].

The method of finding complex Hadamard matrices in the Bose–Mesner algebra of a symmetric association scheme generalizes the classical work of Goethals and Seidel [9]. Assuming that the association scheme is symmetric, the resulting complex Hadamard matrices are symmetric. It turns out that this assumption enables us to consider only the real parts of the entries of a complex Hadamard matrix, since the orthogonality can be expressed in terms of the real parts. Extending this reduction to type-II matrices, we are led to consider a rational map whose inverse is explicitly given in Section 2. In Section 3, we explain why only real parts come into play when we construct complex Hadamard matrices in the Bose–Mesner algebra of a symmetric association scheme. In Section 4, we consider a particular family of 3-class association schemes. This family was found after extensive computer experimentation on the list of 3-class association schemes up to 100 vertices given in [8]. Surprisingly, most other association schemes up to 100 vertices, with the exceptions of amorphic or pseudocyclic schemes, do not admit a complex Hadamard matrix in their Bose–Mesner algebras. In Section 5, we compute the Haagerup set to show inequivalence of type-II matrices constructed in Section 4. In Section 6, we show that the Nomura algebra of each of the type-II matrices constructed in Section 4 has dimension 2. This implies that our matrices are not equivalent to generalized tensor products defined in [11].

All the computer calculations in this paper were performed by Magma [2].

2. THE IMAGE OF A RATIONAL MAP

We define a polynomial in three indeterminates X, Y, Z as follows:

$$g(X, Y, Z) = X^2 + Y^2 + Z^2 - XYZ - 4.$$

Lemma 1.

$$g\left(\frac{X}{Y} + \frac{Y}{X}, \frac{X}{Z} + \frac{Z}{X}, \frac{Z}{Y} + \frac{Y}{Z}\right) = 0.$$

Proof. Straightforward. \square

Lemma 2. *In the rational function field with four indeterminates X, Y, Z and z , the following identity holds:*

$$ww' = 1 + \frac{z^2g + (2zX - zYZ + f)f}{z(zZ - Y)(zY - Z)}, \quad (1)$$

where

$$\begin{aligned} f &= z^2 - zX + 1, \\ g &= g(X, Y, Z), \\ w &= \frac{z^2 - 1}{zZ - Y}, \\ w' &= \frac{z^{-2} - 1}{z^{-1}Z - Y}. \end{aligned}$$

Proof. Straightforward. \square

We define a polynomial in six indeterminates $X_{0,1}, X_{0,2}, X_{0,3}, X_{1,2}, X_{1,3}, X_{2,3}$ as follows:

$$h(X_{0,1}, X_{0,2}, X_{0,3}, X_{1,2}, X_{1,3}, X_{2,3}) = \det \begin{bmatrix} 2 & X_{0,1} & X_{0,2} \\ X_{0,1} & 2 & X_{1,2} \\ X_{0,3} & X_{1,3} & X_{2,3} \end{bmatrix}.$$

Lemma 3. *In the rational function field with four indeterminates X_0, X_1, X_2, X_3 , set*

$$x_{i,j} = \frac{X_i}{X_j} + \frac{X_j}{X_i} \quad (0 \leq i < j \leq 3).$$

Then $h(x_{0,1}, x_{0,2}, x_{0,3}, x_{1,2}, x_{1,3}, x_{2,3}) = 0$.

Proof. Straightforward. \square

For a finite set N and a positive integer k , we denote by $\binom{N}{k}$ the collection of all k -element subsets of N .

Lemma 4. Let $N = \{0, 1, \dots, d\}$, $N_3 = \binom{N}{3}$ and $N_4 = \binom{N}{4}$. Let $a_{i,j}$ ($0 \leq i, j \leq d$, $i \neq j$) be complex numbers satisfying

$$a_{i,j} = a_{j,i} \quad (0 \leq i < j \leq d), \quad (2)$$

$$g(a_{i,j}, a_{j,k}, a_{i,k}) = 0 \quad (\{i, j, k\} \in N_3), \quad (3)$$

$$h(a_{i,j}, a_{i,k}, a_{i,\ell}, a_{j,k}, a_{j,\ell}, a_{k,\ell}) = 0 \quad (\{i, j, k, \ell\} \in N_4). \quad (4)$$

Assume

$$a_{i_0, i_1} \neq \pm 2 \quad \text{for some } i_0, i_1 \text{ with } 0 \leq i_0 < i_1 \leq d. \quad (5)$$

Let w_{i_0}, w_{i_1} be nonzero complex numbers satisfying

$$\frac{w_{i_0}}{w_{i_1}} + \frac{w_{i_1}}{w_{i_0}} = a_{i_0, i_1}. \quad (6)$$

Then for complex numbers w_i ($0 \leq i \leq d$, $i \neq i_0, i_1$), the following are equivalent:

(i) for all i, j with $0 \leq i, j \leq d$ and $i \neq j$,

$$\frac{w_j}{w_i} + \frac{w_i}{w_j} = a_{i,j} \quad (7)$$

(ii) for all i, j with $0 \leq i \leq d$, $i \neq i_0, i_1$,

$$w_i = \frac{w_{i_1}^2 - w_{i_0}^2}{a_{i_1, i} w_{i_1} - a_{i_0, i} w_{i_0}}. \quad (8)$$

Moreover, if one of the two equivalent conditions (i), (ii) is satisfied, $a_{i,j}$ ($0 \leq i < j \leq d$) are all real and

$$-2 < a_{i_0, i_1} < 2, \quad (9)$$

then $|w_i| = |w_j|$ for $0 \leq i < j \leq d$.

Proof. Without loss of generality, we may assume $i_0 = 0$ and $i_1 = 1$. Thus (6) reads

$$\frac{w_0}{w_1} + \frac{w_1}{w_0} = a_{0,1}. \quad (10)$$

Then by (5) and (3) with $\{i, j, k\} = \{0, 1, i\}$, we find

$$a_{0,i}^2 + a_{1,i}^2 - a_{0,1} a_{0,i} a_{1,i} \neq 0. \quad (11)$$

First we need to check that the denominator of (8) is nonzero. Suppose $a_{1,i} w_1 - a_{0,i} w_0 = 0$. If, moreover $a_{1,i} = 0$, then we have $a_{0,i} = 0$, but this contradicts (11). Thus we have $a_{1,i} \neq 0$, and

$$\frac{w_1}{w_0} = \frac{a_{0,i}}{a_{1,i}}.$$

Then by (10), we have

$$\frac{a_{1,i}}{a_{0,i}} + \frac{a_{0,i}}{a_{1,i}} = a_{0,1},$$

but again this contradicts (11). Therefore, w_i is well-defined. Moreover, we claim $w_i \neq 0$. Indeed, $w_i = 0$ would imply $w_1^2 = w_0^2$. Then by (10), we have $a_{0,1} = \pm 2$, which contradicts (5).

Clearly, (10) implies (7) with $(i, j) = (0, 1)$. Suppose $j \geq 2$. Observe that $x = w_j = (w_1^2 - w_0^2)/(a_{1,j}w_1 - a_{0,j}w_0)$ given in (8) is a unique common root of the equations

$$\begin{aligned} x^2 - w_0 a_{0,j} x + w_0^2 &= 0, \\ x^2 - w_1 a_{1,j} x + w_1^2 &= 0. \end{aligned}$$

This can be seen by $g(a_{0,1}, a_{0,j}, a_{1,j}) = 0$. Thus (7) holds if $i = 0$ or 1 . Now assume $\{i, j\} \cap \{0, 1\} = \emptyset$. By (4), we have

$$\begin{aligned} 0 &= w_0^2 w_1^2 w_i w_j h(a_{0,1}, a_{0,i}, a_{0,j}, a_{1,i}, a_{1,j}, a_{i,j}) \\ &= \det \begin{bmatrix} 2w_0^2 & w_0 w_1 a_{0,1} & w_0 w_i a_{0,i} \\ w_0 w_1 a_{0,1} & 2w_1^2 & w_1 w_i a_{1,i} \\ w_0 w_j a_{0,j} & w_1 w_j a_{1,j} & w_i w_j a_{i,j} \end{bmatrix} \\ &= \det \begin{bmatrix} 2w_0^2 & w_0^2 + w_1^2 & w_0^2 + w_i^2 \\ w_0^2 + w_1^2 & 2w_1^2 & w_1^2 + w_i^2 \\ w_0^2 + w_j^2 & w_1^2 + w_j^2 & w_i w_j a_{i,j} \end{bmatrix} \\ &= -(w_1^2 - w_0^2)^2 (w_i w_j a_{i,j} - w_i^2 - w_j^2). \end{aligned}$$

Thus (7) holds.

Conversely, suppose that $\{w_i\}_{i=0}^d$ satisfy (7). Setting $(i, j) = (0, 1)$ in (7) gives (10). Setting $i = 1$ and replacing j by i in (7) gives

$$\frac{w_i}{w_1} + \frac{w_1}{w_i} = a_{1,i}.$$

By (7), we have

$$\begin{aligned} a_{1,i} w_1 w_i &= w_1^2 + w_i^2 \\ &= a_{0,i} w_0 w_i + w_1^2 - w_0^2, \end{aligned}$$

which gives (8).

Finally, assume that $a_{i,j}$ ($0 \leq i < j \leq d$) are all real and that (9) holds for $(i_0, i_1) = (0, 1)$. Then by (6), w_1/w_0 is an imaginary number with absolute value 1. This means

$$\left(\frac{w_1}{w_0}\right)^{-1} = \overline{\left(\frac{w_1}{w_0}\right)}. \quad (12)$$

For $j \geq 2$, set $(X, Y, Z, z) = (a_{0,1}, a_{0,j}, a_{1,j}, w_1/w_0)$ in Lemma 2. It is easy to check that all the denominators in Lemma 2 are nonzero. Moreover,

$$f = 0 \quad (\text{by (6)}), \quad (13)$$

$$g = 0 \quad (\text{by (3)}), \quad (14)$$

$$w = \frac{w_j}{w_0} \quad (\text{by (8)}), \quad (15)$$

and

$$\begin{aligned} \overline{w'} &= \overline{\left(\frac{\left(\frac{w_1}{w_0}\right)^{-2} - 1}{\left(\frac{w_1}{w_0}\right)^{-1} a_{1,j} - a_{0,j}} \right)} \\ &= \frac{\left(\frac{w_1}{w_0}\right)^2 - 1}{\frac{w_1}{w_0} a_{1,j} - a_{0,j}} \quad (\text{by (12)}) \\ &= \frac{w_1^2 - w_0^2}{w_0(a_{1,j}w_1 - a_{0,j}w_0)} \\ &= \frac{w_j}{w_0} \quad (\text{by (8)}) \\ &= w \quad (\text{by (15)}). \quad (16) \end{aligned}$$

Now

$$\begin{aligned} 1 &= ww' \quad (\text{by (1), (13), (14)}) \\ &= |w|^2 \quad (\text{by (16)}) \\ &= \left| \frac{w_j}{w_0} \right|^2 \quad (\text{by (15)}). \end{aligned}$$

Therefore $|w_i| = |w_j|$ for $0 \leq i < j \leq d$. \square

In the following theorem, $\mathbb{C}^{d(d+1)/2}$ will mean the \mathbb{C} -vector space whose coordinates are indexed by ordered pairs (i, j) of integers with $0 \leq i < j \leq d$.

Theorem 1. *Let d, N, N_3, N_4 be as in Lemma 4. Define $\phi : (\mathbb{C}^\times)^{d+1} \rightarrow \mathbb{C}^{d(d+1)/2}$ by*

$$\phi(w_0, \dots, w_d) = \left(\frac{w_i}{w_j} + \frac{w_j}{w_i} \right)_{0 \leq i < j \leq d}.$$

Then the image of ϕ coincides with the zeros of the ideal generated by the polynomials

$$g(X_{i,j}, X_{j,k}, X_{i,k}) \quad (\{i, j, k\} \in N_3), \quad (17)$$

$$h(X_{i,j}, X_{i,k}, X_{i,\ell}, X_{j,k}, X_{j,\ell}, X_{k,\ell}) \quad (\{i, j, k, \ell\} \in N_4), \quad (18)$$

where $X_{i,j} = X_{j,i}$.

Proof. In the polynomial ring $\mathbb{C}[X_{i,j} \mid 0 \leq i < j \leq d]$, let I denote the ideal generated by (17) and (18). By Lemmas 1 and 3, the image of ϕ is contained in the set of zeros of I .

Conversely, let $a = (a_{i,j})_{0 \leq i < j \leq d}$ be a zero of I . If there exists i_0, i_1 with $0 \leq i_0 < i_1 \leq d$ and $a_{i_0, i_1} \neq \pm 2$, then Lemma 4 implies that a is in the image of ϕ . Next suppose that $a_{i,j} \in \{\pm 2\}$ for $0 \leq i < j \leq d$. Then

$$a_{0,1} = \frac{a_{0,1}}{2} + \frac{2}{a_{0,1}}.$$

Also, since $g(a_{0,i}, a_{0,j}, a_{i,j}) = 0$, we have $a_{0,i}a_{0,j}a_{i,j} = 8$. Thus

$$\begin{aligned} a_{i,j} &= \frac{8}{a_{0,i}a_{0,j}} \cdot \frac{a_{0,i}^2}{4} \cdot \frac{a_{0,j}^2}{4} \\ &= \frac{1}{2}a_{0,i}a_{0,j} \\ &= \frac{a_{0,i}}{a_{0,j}} + \frac{a_{0,j}}{a_{0,i}}. \end{aligned}$$

Therefore, $a = \phi(2, a_{0,1}, \dots, a_{0,d})$ is in the image of ϕ . \square

The following lemma will be used in the proof of Theorem 2.

Lemma 5. Let $\mathbb{Q}(X_1, X_2, X_3)$ be the rational function field with three indeterminates X_1, X_2, X_3 . Set $X_0 = 1$ and

$$x_{i,j} = \frac{X_i}{X_j} + \frac{X_j}{X_i} \quad (0 \leq i < j \leq 3).$$

Then

$$\begin{aligned} &(X_1X_2X_3 + 1)(x_{0,1}x_{0,2} + x_{0,3} - x_{1,2}) \\ &= (X_1X_2 + X_3) \left(x_{0,1}x_{0,2}x_{0,3} + 2 - \frac{1}{2}(x_{1,2}x_{0,3} + x_{1,3}x_{0,2} + x_{2,3}x_{0,1}) \right). \end{aligned}$$

Proof. Straightforward. \square

3. TYPE-II MATRICES CONTAINED IN A BOSE–MESNER ALGEBRA

Throughout this section, we let \mathcal{A} denote a symmetric Bose–Mesner algebra with adjacency matrices $A_0 = I, A_1, \dots, A_d$. Let n be the size of the matrices A_i , and we denote by

$$P = (P_{i,j})_{\substack{0 \leq i \leq d \\ 0 \leq j \leq d}}$$

the first eigenmatrix of \mathcal{A} . Then the adjacency matrices are expressed as

$$A_j = \sum_{i=0}^d P_{i,j} E_i \quad (j = 0, 1, \dots, d),$$

where $E_0 = \frac{1}{n}J, E_1, \dots, E_d$ are the primitive idempotents of \mathcal{A} . The second eigenmatrix

$$Q = (Q_{i,j})_{\substack{0 \leq i \leq d \\ 0 \leq j \leq d}}$$

is defined as $Q = nP^{-1}$, so that

$$E_j = \frac{1}{n} \sum_{i=0}^d Q_{i,j} A_i \quad (j = 0, 1, \dots, d)$$

holds. Since $QP = nI$ and $Q_{i,0} = P_{i,0} = 1$ for $i = 0, 1, \dots, d$, we have

$$\sum_{j=1}^d Q_{i,j} = n\delta_{i,0} - 1. \quad (19)$$

Lemma 6. *Let w_0, w_1, \dots, w_d be nonzero complex numbers, and set*

$$W = \sum_{j=0}^d w_j A_j \in \mathcal{A}. \quad (20)$$

Set

$$\beta_k = \sum_{j=0}^d w_j P_{k,j}, \quad \beta'_k = \sum_{j=0}^d w_j^{-1} P_{k,j} \quad (k = 0, 1, \dots, d). \quad (21)$$

Then the following are equivalent.

- (i) W is a type-II matrix,
- (ii) $\beta_k \beta'_k = n$ for all k with $1 \leq k \leq d$.

Proof. Since

$$W = \sum_{k=0}^d \beta_k E_k, \quad W^{(-)} = \sum_{k=0}^d \beta'_k E_k.$$

W is type-II matrix if and only if $\beta_k \beta'_k = n$ for $0 \leq k \leq d$. In particular, (i) implies (ii).

To prove the converse, it suffices to show $\beta_0\beta'_0 = n$. Since W is symmetric, the diagonal entries of $WW^{(-)}$ are all n . Thus

$$\begin{aligned} n^2 &= \text{tr } WW^{(-)} \\ &= \sum_{k=0}^d \beta_k\beta'_k \text{tr } E_k \\ &= \beta_0\beta'_0 + \sum_{k=1}^d n \text{tr } E_k \\ &= \beta_0\beta'_0 + n \text{tr}(I - E_0) \\ &= \beta_0\beta'_0 + n(n - 1), \end{aligned}$$

and hence $\beta_0\beta'_0 = n$. \square

Lemma 7. Let $X_{i,j}$ ($0 \leq i < j \leq d$) be indeterminates and let e_k be the polynomial defined by

$$e_k = \sum_{0 \leq i < j \leq d} P_{k,i}P_{k,j}X_{i,j} + \sum_{i=0}^d P_{k,i}^2 - n \quad (k = 1, \dots, d). \quad (22)$$

Let $a_{i,j}$ ($0 \leq i, j \leq d$, $i \neq j$) and w_i ($0 \leq i \leq d$) be complex numbers. Assume that $w_i \neq 0$ for all i with $0 \leq i \leq d$ and that (7) holds. Then the following statements are equivalent:

- (i) the matrix W given by (20) is a type-II matrix,
- (ii) $(a_{i,j})_{0 \leq i < j \leq d}$ is a common zero of e_k ($k = 1, \dots, d$).

Moreover, if one of the two equivalent conditions (i), (ii) is satisfied, $a_{i,j} \in \mathbb{R}$ for all i, j with $0 \leq i < j \leq d$, and (9) holds for some i_0, i_1 with $0 \leq i_0 < i_1 \leq d$, then W is a scalar multiple of a complex Hadamard matrix.

Proof. Define β_k, β'_k by (21). Then

$$\beta_k\beta'_k = \sum_{0 \leq i < j \leq d} P_{k,i}P_{k,j} \left(\frac{w_i}{w_j} + \frac{w_j}{w_i} \right) + \sum_{i=0}^d P_{k,i}^2 \quad (k = 1, \dots, d). \quad (23)$$

Thus, equivalence of (i) and (ii) follows immediately from Lemma 6.

Moreover, suppose that $a_{i,j} \in \mathbb{R}$ for all i, j with $0 \leq i, j \leq d$, $i \neq j$, and (9) holds for some i_0, i_1 . Then (5) holds. This, together with (7) enables us to use Lemma 4 to conclude that w_0, w_1, \dots, w_d have the same absolute value. Therefore W is a scalar multiple of a complex Hadamard matrix. \square

4. INFINITE FAMILIES OF COMPLEX HADAMARD MATRICES

Let $q \geq 4$ be an integer, and $n = q^2 - 1$. We consider a three-class association scheme $\mathcal{X} = (X, \{R_i\}_{i=0}^3)$ with the first eigenmatrix:

$$P = \begin{bmatrix} 1 & \frac{q^2}{2} - q & \frac{q^2}{2} & q - 2 \\ 1 & \frac{q}{2} & -\frac{q}{2} & -1 \\ 1 & -\frac{q}{2} + 1 & -\frac{q}{2} & q - 2 \\ 1 & -\frac{q}{2} & \frac{q}{2} & -1 \end{bmatrix}. \quad (24)$$

For $q = 2^s$ with an integer $s \geq 2$, there exists a 3-class association scheme with the first eigenmatrix (24) (see [3, 12.1.1]).

Let $\mathcal{M} = \langle A_0, A_1, A_2, A_3 \rangle$ be the Bose–Mesner algebra of \mathcal{X} . Then, \mathcal{X} has two non-trivial fusion schemes. One is an imprimitive scheme $\mathcal{X}_1 = (X, \{R_0, R_1 \cup R_2, R_3\})$ with the first eigenmatrix:

$$P_1 = \begin{bmatrix} 1 & q(q-1) & q-2 \\ 1 & 0 & -1 \\ 1 & -q+1 & q-2 \end{bmatrix}. \quad (25)$$

Another is a primitive scheme $\mathcal{X}_2 = (X, \{R_0, R_1 \cup R_3, R_2\})$ with the first eigenmatrix:

$$P_2 = \begin{bmatrix} 1 & \frac{q^2}{2} - 2 & \frac{q^2}{2} \\ 1 & \frac{q}{2} - 1 & -\frac{q}{2} \\ 1 & -\frac{q}{2} - 1 & \frac{q}{2} \end{bmatrix}. \quad (26)$$

Theorem 2. *Let w_1, w_2, w_3 be nonzero complex numbers. The matrix*

$$W = A_0 + w_1 A_1 + w_2 A_2 + w_3 A_3 \in \mathcal{M} \quad (27)$$

is a type-II matrix if and only if one of the following holds:

(i) $w_1 = w_2 = w_3$, where

$$w_3 + \frac{1}{w_3} + q^2 - 3 = 0,$$

(ii) w_3 is as in (i), and

$$w_1 = w_2 = \frac{-(q-3)w_3 + (q-1)}{q^2 - 2q - 1},$$

(iii)

$$w_1 + \frac{1}{w_1} = \frac{2(q^2 - 6)}{q^2 - 4}, \quad w_2 = -1, \quad w_3 = w_1,$$

(iv)

$$w_1 = w_3 = 1, \quad w_2 + \frac{1}{w_2} = \frac{-2(q^2 - 2)}{q^2},$$

$$(v) \quad w_1 + \frac{1}{w_1} = -\frac{2}{q}, \quad w_2 = \frac{1}{w_1}, \quad w_3 = 1,$$

$$(vi) \quad w_1 + \frac{1}{w_1} = a_{0,1},$$

and

$$w_i = \frac{w_1^2 - 1}{a_{1,i}w_1 - a_{0,i}} \quad (i = 2, 3),$$

where

$$a_{0,1} = \frac{-(q-1)(q-2) + (q+2)r}{2q(q+1)},$$

$$a_{0,2} = \frac{(q+2)(q-1) - (q-2)r}{2q(q-3)},$$

$$a_{0,3} = \frac{5q^2 - 2q - 19 - (q-1)r}{2(q+1)(q-3)},$$

$$a_{1,2} = \frac{2(-q^4 + 2q^3 + 4q^2 - 10q + 1 + (q-1)r)}{q^2(q+1)(q-3)},$$

$$a_{1,3} = -a_{0,2},$$

$$r^2 = (17q-1)(q-1).$$

Note that $w_1w_2 = -w_3$ holds.

Proof. Let $K = \mathbb{Q}(r)$, where r is a square root of $(17q-1)(q-1)$. Note that K is either \mathbb{Q} or its quadratic extension (see Remark 2). Consider the polynomial ring

$$R = K[X_{0,1}, X_{0,2}, X_{0,3}, X_{1,2}, X_{1,3}, X_{2,3}].$$

We assume that W is a type-II matrix. For $i, j \in \{0, 1, 2, 3\}$, define $a_{i,j}$ by (7), where $w_0 = 1$. We write

$$\mathbf{a} = (a_{0,1}, a_{0,2}, a_{0,3}, a_{1,2}, a_{1,3}, a_{2,3})$$

for brevity. Then by Lemmas 1, 3 and 7, \mathbf{a} is a common zero of the polynomials

$$g(X_{i,j}, X_{i,k}, X_{j,k}) \quad (\{i, j, k\} \in \binom{\{0, 1, 2, 3\}}{3}),$$

$$h(X_{i,j}, X_{i,k}, X_{i,l}, X_{j,k}, X_{j,l}, X_{k,l}) \quad (\{i, j, k, l\} = \{0, 1, 2, 3\}),$$

$$e_k \quad (k \in \{1, 2, 3\}).$$

Let \mathcal{I} be the ideal of R generated by these polynomials. Then we can verify that \mathcal{I} contains the polynomial $b_1 b_2 b_3 b_4^+ b_4^-$, where

$$\begin{aligned} b_1 &= X_{1,2} - 2, \\ b_2 &= (q^2 - 4)X_{1,2} + 2q^2 - 12, \\ b_3 &= q^2 X_{1,2} + 2q^2 - 4, \\ b_4^\pm &= q^2(q+1)(q-3)X_{1,2} - 2(-q^4 + 2q^3 + 4q^2 - 10q + 1 \pm (q-1)r). \end{aligned}$$

First assume $a_{1,2}$ is a zero of the polynomial b_1 . This implies $w_1 = w_2$. Let \mathcal{I}_1 denote the ideal generated by \mathcal{I} and b_1 . Then we can verify that \mathcal{I}_1 contains the polynomials $X_{0,3} + q^2 - 3$ and $c_1 c_2$, where

$$\begin{aligned} c_1 &= X_{0,1} + q^2 - 3, \\ c_2 &= (q^2 - 2q - 1)X_{0,1} - (q^3 - 3q^2 - q + 7). \end{aligned}$$

In particular,

$$a_{0,3} = w_3 + \frac{1}{w_3} = -(q^2 - 3). \quad (28)$$

If $a_{0,1}$ is a zero of the polynomial c_1 , then let \mathcal{I}'_1 denote the ideal generated by \mathcal{I}_1 and c_1 . We can verify that \mathcal{I}'_1 contains the polynomial $X_{1,3} - 2$. Hence $w_1 = w_2 = w_3$, and we have Case (i).

If $a_{0,1}$ is a zero of the polynomial c_2 , then

$$a_{0,1} = \frac{q^3 - 3q^2 - q + 7}{q^2 - 2q - 1}. \quad (29)$$

Let \mathcal{I}''_1 denote the ideal generated by \mathcal{I}_1 and c_2 . We can verify that

$$(q^2 - 2q - 1)X_{1,3} + (q^3 - q^2 - q - 3) \in \mathcal{I}''_1.$$

This implies

$$a_{1,3} = -\frac{q^3 - q^2 - q - 3}{q^2 - 2q - 1} \neq \pm 2. \quad (30)$$

Applying Lemma 4 with (28)–(30), we obtain from (8),

$$\begin{aligned} w_1 = w_2 &= \frac{w_3^2 - 1}{a_{1,3}w_3 - a_{0,1}} \\ &= \frac{-(q-3)w_3 + (q-1)}{q^2 - 2q - 1}. \end{aligned}$$

This gives Case (ii).

Next assume $a_{1,2}$ is a zero of the polynomial b_2 . Let \mathcal{I}_2 denote the ideal generated by \mathcal{I} and b_2 . Then we can verify that \mathcal{I}_2 contains the polynomials $X_{0,2} + 2$, $X_{1,3} - 2$, and $(q^2 - 4)X_{0,1} - 2(q^2 - 6)$. Hence $w_2 = -1$, $w_1 = w_3$ and $w_1 + 1/w_1 = 2(q^2 - 6)/(q^2 - 4)$. This gives Case (iii).

Next assume $a_{1,2}$ is a zero of the polynomial b_3 . Let \mathcal{I}_3 denote the ideal generated by \mathcal{I} and b_3 . Then we can verify that \mathcal{I}_3 contains the polynomials $X_{0,3} - 2$ and c_4c_5 , where

$$\begin{aligned} c_4 &= q^2X_{0,2} + 2(q^2 - 2), \\ c_5 &= qX_{0,2} + 2. \end{aligned}$$

In particular, $w_3 = 1$.

If $a_{0,2}$ is a zero of the polynomial c_4 , then

$$a_{0,2} = -\frac{2(q^2 - 2)}{q^2}.$$

Let \mathcal{I}'_3 denote the ideal generated by \mathcal{I} and c_4 . We can verify that \mathcal{I}'_3 contains the polynomial $X_{0,1} - 2$. Hence $w_1 = w_3 = 1$, $w_2 + 1/w_2 + 2(q^2 - 2)/q^2 = 0$, and we have Case (iv).

If $a_{0,2}$ is a zero of the polynomial c_5 , then

$$a_{0,2} = -\frac{2}{q}. \tag{31}$$

Let \mathcal{I}''_3 denote the ideal generated by \mathcal{I}_3 and c_5 . We can verify that \mathcal{I}''_3 contains the polynomials $qX_{0,1} + 2$ and $q^2X_{1,2} + 2(q^2 - 2)$. Thus

$$a_{0,1} = -\frac{2}{q}, \tag{32}$$

$$a_{1,2} = -\frac{2(q^2 - 2)}{q^2}. \tag{33}$$

Since

$$\begin{aligned} w_1w_2 + \frac{1}{w_1w_2} &= \left(w_1 + \frac{1}{w_1}\right) \left(w_2 + \frac{1}{w_2}\right) - \left(\frac{w_2}{w_1} + \frac{w_1}{w_2}\right) \\ &= a_{0,1}a_{0,2} - a_{1,2} \\ &= 2 \end{aligned}$$

by (31)–(33), we have $w_1w_2 = 1$. This gives Case (v).

Next assume $a_{1,2}$ is a zero of the polynomial $b_4^+b_4^-$. Without loss of generality, we may assume $a_{1,2}$ is a zero of the polynomial b_4^+ . Let \mathcal{I}_4 denote the ideal generated by \mathcal{I} and b_4^+ . Then we can verify that \mathcal{I}_4

contains the following polynomials:

$$2q(q+1)X_{0,1} + (q-1)(q-2) - (q-2)r, \quad (34)$$

$$2q(q-3)X_{0,2} - (q+2)(q-1) + (q-2)r, \quad (35)$$

$$2(q+1)(q-3)X_{0,3} - (5q^2 - 2q - 19) + (q-1)r, \quad (36)$$

$$q^2(q+1)(q-3)X_{1,2} - 2(-q^4 + 2q^3 + 4q^2 - 10q + 1 + (q-1)r), \quad (37)$$

$$2q(q-3)X_{1,3} + (q+2)(q-1) - (q-2)r. \quad (38)$$

Since $a_{1,2} \neq \pm 2$, we can apply Lemma 4 to obtain Case (vi) from (8). Moreover, we can verify that \mathcal{I}_4 contains the polynomial

$$X_{0,1}X_{0,2} + X_{0,3} - X_{1,2}.$$

Thus

$$\begin{aligned} 0 &= (w_1w_2w_3 + 1)(a_{0,1}a_{0,2} + a_{0,3} - a_{1,2}) \\ &= (w_1w_2 + w_3)(a_{0,1}a_{0,2}a_{0,3} + 2 - \frac{1}{2}(a_{1,2}a_{0,3} + a_{1,3}a_{0,2} + a_{2,3}a_{0,1})) \end{aligned}$$

by Lemma 5. Since the ideal generated by \mathcal{I}_4 and the polynomial

$$X_{0,1}X_{0,2}X_{0,3} + 2 - \frac{1}{2}(X_{1,2}X_{0,3} + X_{1,3}X_{0,2} + X_{2,3}X_{0,1})$$

is trivial, we conclude $w_1w_2 + w_3 = 0$.

Conversely, assume that w_1, w_2 , and w_3 are given in Theorem 2. Then, we show that the matrix W given in (27) is a type-II matrix. To do this, from Lemma 7 we only check that \mathbf{a} defined by (7) is a zero of the polynomials (22).

First consider Case (i). Then, by (7) we have

$$\mathbf{a} = (-q^2 + 3, -q^2 + 3, -q^2 + 3, 2, 2, 2),$$

and this is a zero of the polynomials (22). Hence W is a type-II matrix.

Next consider Case (ii). This means $w_1 = aw_3 + b$, with

$$\begin{aligned} a &= -\frac{q-3}{q^2-2q-1}, \\ b &= \frac{q-1}{q^2-2q-1}. \end{aligned}$$

We claim

$$\frac{1}{w_1} = a\frac{1}{w_3} + b.$$

Indeed, this can be checked by showing

$$(aw_3 + b) \left(a\frac{1}{w_3} + b \right) = 1$$

using $w_3 + \frac{1}{w_3} = -(q^2 - 3)$. Thus

$$\begin{aligned} a_{0,1} &= w_1 + \frac{1}{w_1} \\ &= \frac{q^3 - 3q^2 - q + 7}{q^2 - 2q - 1}, \\ a_{1,3} &= \frac{w_1}{w_3} + \frac{w_3}{w_1} \\ &= \frac{aw_3 + b}{w_3} + w_3 \left(a \frac{1}{w_3} + b \right) \\ &= 2a + b \left(w_3 + \frac{1}{w_3} \right) \\ &= 2a - b(q^2 - 3) \\ &= \frac{-q^3 + q^2 + q + 3}{q^2 - 2q - 1}. \end{aligned}$$

Now

$$\mathbf{a} = (a_{0,1}, a_{0,1}, -(q^2 - 3), 2, a_{1,3}, a_{1,3})$$

is a zero of the polynomials (22). Hence W is a type-II matrix.

Next consider Case (iii). Then, by (7) we have

$$\mathbf{a} = \left(\frac{2(q^2 - 6)}{q^2 - 4}, -2, \frac{2(q^2 - 6)}{q^2 - 4}, -\frac{2(q^2 - 6)}{q^2 - 4}, 2, -\frac{2(q^2 - 6)}{q^2 - 4} \right),$$

and this is a zero of the polynomials (22). Hence W is a type-II matrix.

Next consider Case (iv). Then, by (7) we have

$$\mathbf{a} = \left(2, \frac{-2(q^2 - 2)}{q^2}, 2, \frac{-2(q^2 - 2)}{q^2}, 2, \frac{-2(q^2 - 2)}{q^2} \right),$$

and this is a zero of the polynomials (22). Hence W is a type-II matrix.

Next consider Case (v). Then, by (7) we have

$$\begin{aligned} a_{1,2} &= a_{0,1}^2 - 2 \\ &= \frac{-2(q^2 - 2)}{q^2}, \end{aligned}$$

and

$$\mathbf{a} = \left(-\frac{2}{q}, -\frac{2}{q}, 2, \frac{-2(q^2 - 2)}{q^2}, -\frac{2}{q}, -\frac{2}{q} \right),$$

and this is a zero of the polynomials (22). Hence W is a type-II matrix.

Lastly consider Case (vi). Then a is a zero of the polynomials (22). Hence W is a type-II matrix. \square

Corollary 1. *Let W be a type-II matrix in Theorem 2. Then, W is a complex Hadamard matrix if and only if W is given in (iii), (iv), (v), or (vi) with $r = \sqrt{(17q-1)(q-1)} > 0$.*

Proof. Since $q \geq 4$, we have $n \geq 15$. Hence, $|w_3| \neq 1$ for Cases (i) and (ii). For Cases (iii), (iv) and (v), it is easy to see that

$$-2 < w_i + \frac{1}{w_i} < 2 \quad \text{for some } i \in \{1, 2, 3\}.$$

Therefore, by Lemma 7 the Cases (iii), (iv) and (v) give complex Hadamard matrices.

Lastly, we consider the Case (vi). If $r > 0$, then we have $0 < a_{0,1} < 2$. In fact, since $r > 3q$, we have $a_{0,1} > 0$. Also,

$$a_{0,1} - 2 = \frac{-8q(q+1)(q-3)^2}{2q(q+1)(5q^2+q+2+(q+2)r)} < 0.$$

Thus, we have $0 < a_{0,1} < 2$. By Lemma 7, W is a complex Hadamard matrix.

If $r < 0$, then

$$a_{0,1} + 2 = \frac{-8q(q+1)(q^2-5)}{2q(q+1)(3q^2+7q-2-(q+2)r)} < 0.$$

This implies $|w_1| \neq 1$. Therefore, W is not a complex Hadamard matrix. \square

Chan [4], found three complex Hadamard matrices on the line graph of the Petersen graph. This is the 3-class association scheme with the first eigenmatrix (24), where $q = 4$, and the three matrices can be described as the matrix W in (27) with w_1, w_2, w_3 given as follows.

$$w_1 = 1, \quad w_2 = \frac{-7 \pm \sqrt{15}i}{8}, \quad w_3 = 1, \quad (39)$$

$$w_1 = \frac{5 \pm \sqrt{11}i}{6}, \quad w_2 = -1, \quad w_3 = w_1, \quad (40)$$

$$w_1 = \frac{-1 \pm \sqrt{15}i}{4}, \quad w_2 = w_1^{-1}, \quad w_3 = 1. \quad (41)$$

The cases (39), (40) and (41) are given by (iv), (iii) and (v), respectively, of Theorem 2. Note that (39) is equivalent to the matrix U_{15} in [16].

The complex Hadamard matrix of order 15 constructed in Theorem 2 (vi) seems to be new. This is obtained by setting $q = 4$ and

$r = \sqrt{201}$, and has coefficients $w_1, w_2, w_3 = -w_1w_2$, where

$$\begin{aligned} w_1 + \frac{1}{w_1} &= a_{0,1}, \\ a_{0,1} &= \frac{3}{20}(\sqrt{201} - 1), \\ w_2 &= \frac{a_{0,1}w_1 - 2}{a_{1,2}w_1 - a_{0,2}}, \\ a_{0,2} &= -\frac{1}{4}(\sqrt{201} - 9), \\ a_{1,2} &= \frac{3\sqrt{201} - 103}{40}. \end{aligned}$$

We have verified using the span condition [13, Proposition 4.1] that, this matrix, as well as the one given by (39) are isolated, while the two matrices given by (40) and (41) do not satisfy the span condition.

Remark 1. None of the complex Hadamard matrices given in Corollary 1 is a Butson–Hadamard matrix, that is, entries are roots of unity. This follows from the fact that $a_{i,j}$ is not an algebraic integer for some i, j . Indeed, for the Case (iii),

$$a_{0,1} = 2 - \frac{4}{(q-2)(q+2)}$$

is not an integer.

As for the Case (iv),

$$a_{0,2} = -2 + \frac{4}{q^2}$$

is not an integer.

As for the Case (v),

$$a_{0,1} = \frac{-2}{q}$$

is not an integer.

Finally, for the Case (vi), first suppose that r is irrational. If $a_{0,1}$ is an algebraic integer, then so is

$$a'_{0,1} = \frac{-(q-1)(q-2) - (q+2)r}{2q(q+1)}$$

But then

$$a_{0,1} + a'_{0,1} = -1 + \frac{4q-2}{q(q+1)}$$

is an integer, which is absurd.

Next suppose that r is rational. Then $a_{0,1}$ is an integer, so we have $a_{0,1} \in \{0, \pm 1, \pm 2\}$. But this does not occur.

Remark 2. Let q be an even positive integer with $q \geq 4$. Then $\sqrt{(q-1)(17q-1)}$ is an integer if and only if

$$q \in \left\{ \frac{1}{34} \left(\operatorname{tr} \left((2177 + 528\sqrt{17})^n (433 + 105\sqrt{17}) \right) + 18 \right) \mid n \in \mathbb{Z} \right\} \\ = \{ \dots, 41210, 10, 26, 110890, 482812730, \dots \},$$

where tr denotes the trace from $\mathbb{Q}(\sqrt{17})$ to \mathbb{Q} , i.e., $\operatorname{tr}(a+b\sqrt{17}) = 2a$ for $a, b \in \mathbb{Q}$. In this case, the entries of the matrix W in Theorem 2(vi) are quadratic irrationals, so there are two pairs of algebraically conjugate matrices arising from two choices for r . If $\sqrt{(17q-1)(q-1)}$ is not an integer, then the four matrices in Theorem 2(vi) are algebraically conjugate to each other.

5. EQUIVALENCE

For a type-II matrix W of order n , the Haagerup set $H(W)$ (see [10]) is defined as

$$H(W) = \left\{ \frac{W_{i_1, j_1} W_{i_2, j_2}}{W_{i_1, j_2} W_{i_2, j_1}} \mid 1 \leq i_1, i_2, j_1, j_2 \leq n \right\}.$$

We also define

$$K(W) = \left\{ w + \frac{1}{w} \mid w \in H(W) \setminus \{1\} \right\}.$$

If W_1 and W_2 are equivalent, then clearly $H(W_1) = H(W_2)$, and hence $K(W_1) = K(W_2)$. In this section, we compute the Haagerup sets of type-II matrices constructed in Theorem 2 to conclude that some of them are inequivalent to others.

We suppose that

$$W = \sum_{i=0}^d w_i A_i$$

is a complex Hadamard matrix, where A_0, \dots, A_d are the adjacency matrices of a symmetric Bose–Mesner algebra of an association scheme $(X, \{R_i\}_{i=0}^d)$, and $w_0 = 1$. Let $H(W)$ be the Haagerup set of W . Then

$$H(W) = \bigcup_{i=1}^4 H_i(W),$$

where

$$H_i(W) = \left\{ \frac{W_{x_1, y_1} W_{x_2, y_2}}{W_{x_2, y_1} W_{x_1, y_2}} \mid x_1, x_2, y_1, y_2 \in X, |\{x_1, x_2, y_1, y_2\}| = i \right\}$$

for $i = 1, 2, 3, 4$. Clearly,

$$\begin{aligned} H_1(W) &= \{1\}, \\ H_2(W) &= \{1\} \cup \{w_i^{\pm 2} \mid i = 1, \dots, d\}. \end{aligned} \quad (42)$$

It should be remarked that, although $H(W)$ is an invariant, none of $H_i(W)$ ($i = 2, 3, 4$) is.

Lemma 8. *If $|X| \geq 3$, then*

$$H_3(W) = \{1\} \cup \left\{ \left(\frac{w_i w_j}{w_k} \right)^{\pm 1} \mid 1 \leq i, j, k \leq d, p_{ij}^k > 0 \right\}.$$

Proof. Let

$$w = \frac{W_{x_1, y_1} W_{x_2, y_2}}{W_{x_2, y_1} W_{x_1, y_2}} \in H_3(W),$$

where $|\{x_1, x_2, y_1, y_2\}| = 3$. If $x_1 = x_2$, then $w = 1$. If $x_1 = y_1$, then $w = (w_i w_j / w_k)^{-1}$, where $(x_2, x_1) \in R_i$, $(x_1, y_2) \in R_j$, $(x_2, y_2) \in R_k$, and $1 \leq i, j, k \leq d$. All other cases can be treated in a similar manner, and we conclude that $H_3(W)$ is contained in the right-hand side. Arguing in the opposite way, we obtain the reverse containment. \square

Lemma 9. *Let Δ be a subset of $\{1, \dots, d\}$. Suppose that there exists $i \in \{1, \dots, d\}$ such that $p_{i_1, j_1}^i > 0$ for any $i_1, j_1 \in \Delta$. Then*

$$H_4(W) \supset \left\{ \frac{w_{i_1} w_{i_2}}{w_{j_1} w_{j_2}} \mid i_1, i_2, j_1, j_2 \in \Delta \right\} \setminus \{1\}.$$

In particular, if there exists $i \in \{1, \dots, d\}$ such that $p_{i_1, j_1}^i > 0$ for any $i_1, j_1 \in \{1, \dots, d\}$, then

$$H_4(W) \setminus \{1\} = \left\{ \frac{w_{i_1} w_{i_2}}{w_{j_1} w_{j_2}} \mid i_1, i_2, j_1, j_2 \in \{1, \dots, d\} \right\} \setminus \{1\}.$$

Proof. Pick $(x_1, x_2) \in R_i$ with $i \neq 0$. For any $i_1, i_2, j_1, j_2 \in \Delta$, there exists y_1 such that $(x_1, y_1) \in R_{i_1}$ and $(x_2, y_1) \in R_{j_1}$. Similarly, there exists y_2 such that $(x_1, y_2) \in R_{j_2}$ and $(x_2, y_2) \in R_{i_2}$. If $(i_1, j_1) \neq (j_2, i_2)$, then $y_1 \neq y_2$. Thus $|\{x_1, x_2, y_1, y_2\}| = 4$, and hence

$$\begin{aligned} H_4(W) &\supset \left\{ \frac{w_{i_1} w_{i_2}}{w_{j_1} w_{j_2}} \mid i_1, i_2, j_1, j_2 \in \Delta, (i_1, j_1) \neq (j_2, i_2) \right\} \\ &\supset \left\{ \frac{w_{i_1} w_{i_2}}{w_{j_1} w_{j_2}} \mid i_1, i_2, j_1, j_2 \in \Delta \right\} \setminus \{1\}. \end{aligned}$$

The second statement follows immediately from the first one, since $H_4(W) \setminus \{1\}$ is contained in the right-hand side by the definition. \square

Lemma 10. *Suppose that there exists $i \in \{1, \dots, d-1\}$ such that $p_{i_1, j_1}^i > 0$ for any $i_1, j_1 \in \{1, \dots, d-1\}$. Moreover, suppose $p_{i, j}^d > 0$ for any $j \in \{1, \dots, d-1\}$. Then*

$$H_4(W) \setminus \{1\} = \left\{ \frac{w_{i_1} w_{i_2}}{w_{j_1} w_{j_2}} \mid i_1, i_2, j_1, j_2 \in \{1, \dots, d\} \right\} \setminus \{1\}.$$

Proof. Clearly, $H_4(W) \setminus \{1\}$ is contained in the right-hand side, so we show the reverse containment. Setting $\Delta = \{1, \dots, d-1\}$ in Lemma 9, we have

$$H_4(W) \setminus \{1\} \supset \left\{ \frac{w_{i_1} w_{i_2}}{w_{j_1} w_{j_2}} \mid i_1, i_2, j_1, j_2 \in \{1, \dots, d-1\} \right\} \setminus \{1\}.$$

It remains to show that

$$\frac{w_{i_1} w_d}{w_{j_1} w_{j_2}} \in H_4(W) \text{ and } \frac{w_d^2}{w_{j_1} w_{j_2}} \in H_4(W) \quad (\forall i_1, j_1, j_2 \in \{1, \dots, d-1\}).$$

Pick $(x_1, x_2) \in R_i$. Then by the assumption, there exists y_1 such that $(x_1, y_1) \in R_{i_1}$ and $(x_2, y_1) \in R_{j_1}$. Similarly, there exists y_2 such that $(x_1, y_2) \in R_{j_2}$ and $(x_2, y_2) \in R_d$. Now

$$\frac{w_{i_1} w_d}{w_{j_1} w_{j_2}} = \frac{W_{x_1, y_1} W_{x_2, y_2}}{W_{x_2, y_1} W_{x_1, y_2}} \in H_4(W).$$

Replacing i_1 by d in the above argument gives

$$\frac{w_d^2}{w_{j_1} w_{j_2}} = \frac{W_{x_1, y_1} W_{x_2, y_2}}{W_{x_2, y_1} W_{x_1, y_2}} \in H_4(W).$$

□

Below, we determine the Haagerup set of the type-II matrices given in Theorem 2. In what follows, let $\mathfrak{X} = (X, \{R_i\}_{i=0}^3)$ be an association scheme with the first eigenmatrix (24), where q is an even positive integer with $q \geq 4$. The intersection numbers of \mathfrak{X} are given by

$$B_1 = \begin{bmatrix} 0 & 1 & 0 & 0 \\ \frac{q^2}{2} - q & \frac{(q-2)^2}{q} & \frac{(q-2)^2}{q} & \frac{q(q-4)}{4} \\ 0 & \frac{q(q-2)}{4} & \frac{q(q-2)}{4} & \frac{q^2}{4} \\ 0 & \frac{q-4}{2} & \frac{q-2}{2} & 0 \end{bmatrix}, \quad (43)$$

$$B_2 = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & \frac{q(q-2)}{4} & \frac{q(q-2)}{4} & \frac{q^2}{4} \\ \frac{q^2}{2} & \frac{q^2}{4} & \frac{q^2}{4} & \frac{q^2}{4} \\ 0 & \frac{q}{2} & \frac{q-2}{2} & 0 \end{bmatrix}, \quad (44)$$

$$B_3 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & \frac{q-4}{2} & \frac{q-2}{2} & 0 \\ 0 & \frac{q}{2} & \frac{q-2}{2} & 0 \\ q-2 & 0 & 0 & q-3 \end{bmatrix}, \quad (45)$$

where B_h has (i, j) -entry p_{hi}^j ($0 \leq i, j \leq 3$).

Lemma 11. *Let $W = I + \sum_{i=1}^3 w_i A_i$ be a type-II matrix belonging to the Bose–Mesner algebra of \mathfrak{X} . Then*

$$\begin{aligned} H(W) = & \{w_i^{\pm 2} \mid i = 1, 2, 3\} \\ & \cup \left\{ \left(\frac{w_{i_1} w_{i_2}}{w_{i_3}} \right)^{\pm 1} \mid 1 \leq i_1, i_2, i_3 \leq 3, p_{i_2, i_3}^{i_1} > 0 \right\} \\ & \cup \left\{ \frac{w_{i_1} w_{i_2}}{w_{j_1} w_{j_2}} \mid i_1, i_2, j_1, j_2 \in \{1, 2, 3\} \right\}. \end{aligned}$$

Proof. Note that $H_3(W)$ is determined by Lemma 8. We can also easily see that the assumptions of Lemma 10 are satisfied by setting $i = 2$. This gives $H_4(W) \setminus \{1\}$. Since $H_1(W) = \{1\} \subset H_4(W)$, we have the desired expression for $H(W)$. \square

Using Lemma 11, we can determine the Haagerup set $H(W)$ for each type-II matrix given in Theorem 2. Note that the description of $H(W)$ in Table 1 is valid for all even $q \geq 4$, even though $p_{11}^3 = 0$ for $q = 4$.

	$H(W) \setminus \{1\}$	$K(W)$
(i)	$\{w_1^{\pm 1}, w_1^{\pm 2}\}$	$-q^2 + 3, q^4 - 6q^2 + 7$
(ii)	$\{w_1^{\pm 1}, w_1^{\pm 2}, w_3^{\pm 1}, w_3^{\pm 2},$ $\left(\frac{w_1^2}{w_3}\right)^{\pm 1}, \left(\frac{w_3}{w_1}\right)^{\pm 1}, \left(\frac{w_3}{w_1}\right)^{\pm 2}\}$	$-q^2 + 3, q^4 - 6q^2 + 7,$ $\frac{q^3 - 3q^2 - q + 7}{q^2 - 2q - 1}, \dots$
(iii)	$\{-1, \pm w_1^{\pm 1}, \pm w_1^{\pm 2}\}$	$-2, \pm \frac{2(q^2 - 6)}{q^2 - 4}, \pm \frac{2(q^4 - 16q^2 + 56)}{(q^2 - 4)^2}$
(iv)	$\{w_2^{\pm 1}, w_2^{\pm 2}\}$	$-\frac{2(q^2 - 2)}{q^2}, \frac{2(q^4 - 8q^2 + 8)}{q^4}$
(v)	$\{w_1^{\pm 1}, w_1^{\pm 2}, w_1^{\pm 3}, w_1^{\pm 4}\}$	$-\frac{2}{q}, \dots$
(vi)	$\{-1, \pm w_1^{\pm 1}, \pm w_2^{\pm 1}, \pm w_1^{\pm 2},$ $\pm w_2^{\pm 2}, (w_1^{\pm 1} w_2^{\pm 1})^2, \pm w_1^{\pm 1} w_2^{\pm 1},$ $\pm (w_1^2 w_2)^{\pm 1}, \pm (w_1 w_2^2)^{\pm 1}\}$	$-2, \frac{-(q-1)(q-2) + (q+2)r}{2q(q+1)}, \dots$

TABLE 1. Haagerup sets

The elements of $H(W)$ given in Table 1 can be found as follows: As for the Case (i), $K(W)$ has two elements

$$w_1 + \frac{1}{w_1} = -q^2 + 3, \quad w_1^2 + \frac{1}{w_1^2} = q^4 - 6q^2 + 7.$$

As for (ii), $K(W)$ contains

$$w_1 + \frac{1}{w_1} = \frac{q^3 - 3q^2 - q + 7}{q^2 - 2q - 1},$$

by (29). The Cases (iii) and (iv) are immediate. Finally, it is clear that $K(W)$ contains $-\frac{2}{q}$ and -2 , in the Cases (v) and (vi), respectively. We do not need the remaining elements of $K(W)$ to prove the following propositions.

Proposition 1. *Let W_1, \dots, W_6 be type-II matrices given in (i)–(vi) of Theorem 2, respectively. Then W_1, \dots, W_6 are pairwise inequivalent.*

Proof. It suffices to show that the sets $K_i = K(W_i)$ ($i = 1, \dots, 6$) are pairwise distinct. Since $-1 \in H(W_i)$ if and only if $i \in \{3, 6\}$, we see that $K_i \neq K_j$ whenever $i \in \{1, 2, 4, 5\}$ and $j \in \{3, 6\}$. Since W_4 and W_5 are complex Hadamard matrices by Corollary 1, we see that K_4 and K_5 are contained in the interval $[-2, 2]$, while neither K_1 nor K_2 is. Thus $K_i \neq K_j$ whenever $i \in \{1, 2\}$ and $j \in \{4, 5\}$. Also, we have $K_1 \neq K_2$ since

$$\frac{q^3 - 3q^2 - q + 7}{q^2 - 2q - 1} \in K_2 \setminus K_1,$$

and $K_4 \neq K_5$ since

$$-\frac{2}{q} \in K_5 \setminus K_4.$$

It can be checked directly that the element

$$a_{0,1} = \frac{-(q-1)(q-2) + (q+2)r}{2q(q+1)} \in K_6$$

does not belong to K_3 . Thus $K_3 \neq K_6$. \square

Proposition 2. *Let W_+ and W_- be type-II matrices given in Theorem 2 (vi) with $r > 0$ and $r < 0$, respectively. Then W_+ and W_- are inequivalent.*

Proof. By Corollary 1, W_+ is a complex Hadamard matrix. Thus every element of $H(W_+)$ has absolute value 1, and hence $K(W_+)$ is contained in the interval $[-2, 2]$. On the other hand, the element

$$a_{0,1} = \frac{-(q-1)(q-2) - (q+2)\sqrt{(17q-1)(q-1)}}{2q(q+1)} \in K(W_-)$$

does not belong to the interval $[-2, 2]$. Thus $K(W_+) \neq K(W_-)$. \square

We were able to use the Haagerup set to distinguish some of the complex Hadamard matrices in Theorem 2. This is because the Haagerup set can be described by the intersection numbers of the association

scheme, and is independent of the isomorphism class. In general, if $q \geq 8$ is a power of 2, there may be many non-isomorphic association schemes with the eigenmatrix (24). We do not know whether complex Hadamard matrices having the same coefficients are equivalent if they belong to Bose–Mesner algebras of non-isomorphic association schemes.

Note that there are two type-II matrices described in Theorem 2(i), since $w_1 = w_2 = w_3$ is either of the two zeros of a quadratic equation. Similarly, there are two type-II matrices in each of (ii)–(v) in Theorem 2. Moreover, there are four type-II matrices in (vi) because there are two choices for r and $a_{0,1}^2 - 4 \neq 0$. The following lemma shows that the two type-II matrices in Theorem 2(i) are inequivalent, and so are those in Theorem 2(ii).

Lemma 12. *Let W and W' be type-II matrices belonging to the Bose–Mesner algebra of an association scheme $\mathcal{X} = (X, \{R_i\}_{i=0}^d)$. Suppose that each of W and W' has $d + 1$ distinct entries, the valencies of \mathcal{X} are pairwise distinct, and $\min\{p_{11}^i \mid 0 < i \leq d\} > \frac{|X|}{2}$. If W and W' are equivalent, then W is a scalar multiple of W' .*

Proof. Suppose that W and W' are equivalent. Then there exist diagonal matrices $D = \text{diag}(d_1, \dots, d_n)$, $D' = \text{diag}(d'_1, \dots, d'_n)$ and permutation matrices $T = (\delta_{\tau(i),j})$, $T' = (\delta_{\tau'(i),j})$ such that

$$DWD' = TW'T'^{-1} \tag{46}$$

holds. Let

$$W = \sum_{i=0}^d w_i A_i,$$

$$W' = \sum_{i=0}^d w'_i A_i.$$

For $x \in X$, define

$$R_1(x) = \{y \in X \mid (x, y) \in R_1\}.$$

Let $x_1, x_2 \in X$ be distinct. Then $|R_1(x_1) \cap R_1(x_2)| > \frac{|X|}{2}$ by the assumption. Similarly, $|R_1(\tau(x_1)) \cap R_1(\tau(x_2))| > \frac{|X|}{2}$, hence

$$|\tau'^{-1}(R_1(\tau(x_1)) \cap R_1(\tau(x_2)))| > \frac{|X|}{2}.$$

Therefore, there exists

$$y \in R_1(x_1) \cap R_1(x_2) \cap \tau'^{-1}(R_1(\tau(x_1)) \cap R_1(\tau(x_2))).$$

Equivalently, y satisfies

$$(x_1, y), (x_2, y), (\tau(x_1), \tau'(y)), (\tau(x_2), \tau'(y)) \in R_1.$$

Comparing (x_1, y) - and (x_2, y) -entries of (46), we find

$$d_{x_1} w_1 d'_y = d_{x_2} w_1 d'_y = w'_1.$$

This implies $d_{x_1} = d_{x_2}$. Since $x_1, x_2 \in X$ were arbitrary, $D = dI$ for some d . Similarly, $D' = d'I$ for some d' . We now have

$$dd'W = TW'T'^{-1}.$$

Since W has $d + 1$ distinct entries, we have

$$\begin{aligned} |X|k_i &= |\{(x, y) \in X \times X \mid (TW'T'^{-1})_{x,y} = dd'w_i\}| \\ &= |\{(x, y) \in X \times X \mid W'_{\tau(x), \tau'(y)} = dd'w_i\}| \\ &= |\{(x, y) \in X \times X \mid W'_{x,y} = dd'w_i\}|. \end{aligned}$$

Since W' has $d + 1$ distinct entries and the valencies are pairwise distinct, we obtain $dd'w_i = w'_i$. Hence $dd'W = W'$. \square

Proposition 3. *Let W be a type-II matrix given in (i) of Theorem 2. Then W and $W^{(-)}$ are inequivalent. The same conclusion holds if W is a type-II matrix given in (ii) of Theorem 2.*

Proof. First, let W be a type-II matrix given in (i) of Theorem 2. Then the matrices W and $W^{(-)}$ belong to the Bose–Mesner algebra of the association scheme $(X, \{R_0, R_1 \cup R_2 \cup R_3\})$. Since W has two distinct entries and this association scheme satisfies the hypothesis of Lemma 12, W and $W^{(-)}$ are inequivalent.

If W is a type-II matrix given in (ii) of Theorem 2, then W and $W^{(-)}$ belong to the Bose–Mesner algebra of the association scheme $(X, \{R_0, R_1 \cup R_2, R_3\})$. Since W has three distinct entries and this association scheme satisfies the hypothesis of Lemma 12, W and $W^{(-)}$ are inequivalent. \square

We do not know whether the two type-II matrices in each of (iii)–(v) in Theorem 2 are equivalent or not, and whether the two type-II matrices in Theorem 2(vi) with a given sign for r are equivalent or not.

6. NOMURA ALGEBRAS

Since $q^2 - 1$ is a composite, there are uncountably many inequivalent complex Hadamard matrices of order $q^2 - 1$ by [7]. Indeed, such matrices can be constructed using generalized tensor products [11]. We show that none of our complex Hadamard matrices is equivalent to a generalized tensor product. This is done by showing that the Nomura

algebra of any of our complex Hadamard matrices has dimension 2. According to [11], the Nomura algebra of the generalized tensor product of type-II matrices is imprimitive, and this is never the case when it has dimension 2.

For a type-II matrix $W \in M_X(\mathbb{C})$ and $a, b \in X$, we define column vectors Y_{ab} by setting

$$(Y_{ab})_x = \frac{W_{xa}}{W_{xb}} \quad (x \in X).$$

The *Nomura algebra* $N(W)$ of W is the algebra of matrices in $M_n(\mathbb{C})$ such that Y_{ab} is an eigenvector for all $a, b \in X$. It is shown in [12, Theorem 1] that the Nomura algebra is a Bose–Mesner algebra. We first show that the Nomura algebra consists of symmetric matrices.

Throughout this section, we let N denote the Nomura algebra of any of the type-II matrices given in Theorem 2.

Lemma 13. *The algebra N is symmetric.*

Proof. Suppose that N is not symmetric. Then by [12, Proposition 6(i)], there exists $(b, c) \in X^2$ with $b \neq c$ such that

$$\sum_{x \in X} \frac{W_{x,b}^2}{W_{x,c}^2} = 0$$

This is equivalent to

$$\sum_{j,k} p_{jk}^i \frac{w_j^2}{w_k^2} = 0$$

for some $i \in \{1, 2, 3\}$. Using the notation (7), we have

$$\begin{aligned} \sum_{j,k} p_{jk}^i \frac{w_j^2}{w_k^2} &= \sum_{j < k} p_{jk}^i \left(\frac{w_j^2}{w_k^2} + \frac{w_k^2}{w_j^2} \right) + \sum_{j=0}^3 p_{jj}^i \\ &= \sum_{j < k} p_{jk}^i \left(\left(\frac{w_j}{w_k} + \frac{w_k}{w_j} \right)^2 - 2 \right) + \sum_{j=0}^3 p_{jj}^i \\ &= \sum_{j < k} p_{jk}^i (a_{j,k}^2 - 2) + \sum_{j=0}^3 p_{jj}^i. \end{aligned} \tag{47}$$

It can be verified by computer that (47) is nonzero for each of the cases (i)–(vi) in Theorem 2. \square

Since N is symmetric, the adjacency matrices of N are the $(0, 1)$ -matrices representing the connected components of the Jones graph defined as follows (see [12, Sect. 3.3]). The *Jones graph* of a type-II

matrix $W \in M_X(\mathbb{C})$ is the graph with vertex set X^2 such that two distinct vertices (a, b) and (c, d) are adjacent whenever $\langle Y_{ab}, Y_{cd} \rangle \neq 0$, where $\langle \cdot, \cdot \rangle$ denotes the ordinary (not Hermitian) scalar product.

Proposition 4. *The algebra N coincides with the linear span of I and J . In particular, none of the type-II matrices given in Theorem 2 is equivalent to a nontrivial generalized tensor product.*

Proof. We aim to show that $\{(x, y) \mid x, y \in X, x \neq y\}$ is a connected component of the Jones graph. Fix x, y, z with $(x, y), (y, z), (z, x) \in R_3$. Then in the Jones graph, (x, y) and (y, x) belong to the same connected component, and (x, z) and (z, x) belong to the same connected component, by Lemma 13.

We claim that (x, y) and (x, z) belong to the same connected component. Indeed, if (x, y) and (x, z) belong to different connected components, then (y, x) and (z, x) belong to different connected components. In particular,

$$\langle Y_{xy}, Y_{xz} \rangle = \langle Y_{yx}, Y_{zx} \rangle = 0.$$

Let

$$c_{i,j,k} = |\{u \in X \mid (x, u) \in R_i, (y, u) \in R_j, (z, u) \in R_k\}|.$$

Then for $j, k \in \{1, 2\}$,

$$c_{1,j,k} + c_{2,j,k} = c_{j,1,k} + c_{j,2,k} = c_{j,k,1} + c_{j,k,2} = p_{jk}^3,$$

and

$$\sum_{i,j,k=0}^3 c_{i,j,k} \frac{w_i^2}{w_j w_k} = \sum_{i,j,k=0}^3 c_{i,j,k} \frac{w_j w_k}{w_i^2} = 0.$$

It can be verified by computer that these conditions give rise to a polynomial equation in q which has no solution in even positive integers. Therefore, we have proved the claim. This implies that, for each equivalence class C of the equivalence relation $R_0 \cup R_3$, $(C \times C) \cap R_3$ is a clique in the Jones graph.

Let C and C' be two distinct equivalence classes of $R_0 \cup R_3$. We claim that, for any $(x, z) \in C \times C'$, there exist $y \in C$ and $y' \in C'$ such that $\langle Y_{xy}, Y_{xz} \rangle \neq 0$ and $\langle Y_{y'z}, Y_{xz} \rangle \neq 0$. Suppose $(x, z) \in R_1$ and $\langle Y_{xy}, Y_{xz} \rangle = 0$ for all $y \in C$. Then

$$\begin{aligned} 0 &= \sum_{y \in C} \langle Y_{xy}, Y_{xz} \rangle \\ &= \sum_{y \in C} \sum_{u \in X} (Y_{xy})_u (Y_{xz})_u \end{aligned}$$

$$\begin{aligned}
 &= \sum_{y \in C} \sum_{u \in X} \frac{W_{xu}^2}{W_{yu} W_{zu}} \\
 &= \sum_{y \in C} \sum_{i,j=0}^3 \sum_{u \in R_i(x) \cap R_j(z)} \frac{W_{xu}^2}{W_{yu} W_{zu}} \\
 &= \sum_{i,j=0}^3 \sum_{u \in R_i(x) \cap R_j(z)} \sum_{k=0}^3 \sum_{y \in C \cap R_k(u)} \frac{w_i^2}{w_k w_j} \\
 &= \sum_{i,j=0}^3 \sum_{u \in R_i(x) \cap R_j(z)} \sum_{k=0}^3 p_{3k}^i \frac{w_i^2}{w_k w_j} \\
 &= \sum_{i,j,k=0}^3 p_{ij}^1 p_{3k}^i \frac{w_i^2}{w_k w_j}.
 \end{aligned}$$

It can be verified by computer that this leads to a polynomial equation in q which has no solution in even positive integers. Similarly, suppose $(x, z) \in R_2$ and $\langle Y_{xy}, Y_{xz} \rangle = 0$ for all $y \in C$. Then

$$\sum_{i,j,k=0}^3 p_{ij}^2 p_{3k}^i \frac{w_i^2}{w_k w_j} = 0,$$

and again this leads to a contradiction. Thus, there exists $y \in C$ such that $\langle Y_{xy}, Y_{xz} \rangle \neq 0$. Switching the role of x and z , we see that there exists $y' \in C'$ such that $\langle Y_{y'z}, Y_{xz} \rangle \neq 0$. Therefore, we have proved the claim.

Since C and C' are arbitrary, the claim shows that, in the Jones graph, R_3 is contained in a connected component, and that every element $(x, z) \in R_1 \cup R_2$ is adjacent to an element of R_3 . Thus, $R_1 \cup R_2 \cup R_3$ is a connected component of the Jones graph. Therefore, $\dim N = 2$. □

Acknowledgements. We are very grateful to Ferenc Szöllősi and Ada Chan for helpful discussions on various parts of the paper. We also thank Doug Leonard for giving us suggestions for Section 2 and Takao Komatsu for consultation on Pell equations.

REFERENCES

[1] E. Bannai and T. Ito, *Algebraic Combinatorics I: Association Schemes*, Benjamin/Cummings, Menlo Park, 1984.
 [2] W. Bosma, J. Cannon, and C. Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput., 24 (1997), 235–265.

- [3] A. E. Brouwer, A. M. Cohen, and A. Neumaier, *Distance-Regular Graphs*, Springer-Verlag, Berlin, Heidelberg, 1989.
- [4] A. Chan, *Complex Hadamard matrices and strongly regular graphs*, arXiv:1102.5601.
- [5] A. Chan and C. Godsil, *Type-II matrices and combinatorial structures*, *Combinatorica*, **30** (2010), 1–24.
- [6] A. Chan and R. Hosoya, *Type-II matrices attached to conference graphs*, *J. Algebraic Combin.* **20** (2004), 341–351.
- [7] R. Craigen, *Equivalence classes of inverse orthogonal and unit Hadamard matrices*, *Bull. Austral. Math. Soc.* **44** (1991), no. 1, 109–115.
- [8] E. van Dam, *Three-class association schemes*, *J. Algebraic Combin.* **10** (1999), 69–107.
- [9] J. M. Goethals and J. J. Seidel, *Strongly regular graphs derived from combinatorial designs*, *Can. J. Math.*, **22**, (1970), 597–614.
- [10] U. Haagerup, *Orthogonal maximal Abelian $*$ -subalgebras of $n \times n$ matrices and cyclic n -roots*, *Operator Algebras and Quantum Field Theory (Rome)*, Cambridge, MA, International Press, (1996), 296–322.
- [11] R. Hosoya and H. Suzuki, *Type II matrices and their Bose-Mesner algebras*, *J. Algebraic Combin.* **17** (2003), 19–37.
- [12] F. Jaeger, M. Matsumoto, and K. Nomura, *Bose-Mesner algebras related to type II matrices and spin models*, *J. Algebraic Combin.* **8** (1998), 39–72.
- [13] R. Nicoara, *A finiteness result for commuting squares of matrix algebras*, *J. Operator Theory* **55** (2006), 101–116.
- [14] K. Nomura, *Type II matrices of size five*, *Graphs Combin.* **15** (1999), 79–92.
- [15] A. D. Sankey, *Type-II matrices in weighted Bose-Mesner algebras of ranks 2 and 3*, *J. Algebraic Combin.* **32** (2010), 133–153.
- [16] F. Szöllösi, *Exotic complex Hadamard matrices and their equivalence*, *Cryptogr. Commun.* **2** (2010), no. 2, 187–198.
- [17] W. Tadej and K. Życzkowski *A concise guide to complex Hadamard matrices*, *Open Syst. Inf. Dyn.* **13** (2006), 133–177.

APPENDIX A. VERIFICATION BY MAGMA

Proof of Lemma 1.

```

F<X,Y,Z>:=FunctionField(Rationals(),3);
g:=X^2+Y^2+Z^2-X*Y*Z-4;
g @ hom<F->F|X/Y+Y/X,X/Z+Z/X,Z/Y+Y/Z> eq 0;

```

Proof of Lemma 2.

```

F4<X,Y,Z,z>:=FunctionField(Rationals(),4);
f:=z^2-z*X+1;
g:=X^2+Y^2+Z^2-X*Y*Z-4;
w:=(z^2-1)/(z*Z-Y);
ww:=(z^(-2)-1)/(z^(-1)*Z-Y);
w*ww eq 1+(z^2*g+(2*z*X-z*Y*Z+f)*f)/(z*(z*Z-Y)*(z*Y-Z));

```

Proof of Lemma 3.

```

F<X0,X1,X2,X3>:=FunctionField(Rationals(),4);
x01:=X0/X1+X1/X0;
x02:=X0/X2+X2/X0;
x03:=X0/X3+X3/X0;
x12:=X1/X2+X2/X1;
x13:=X1/X3+X3/X1;
x23:=X2/X3+X3/X2;
hX:=Matrix(F,3,3,[2,x01,x02, x01,2,x12, x03,x13,x23]);
Determinant(hX) eq 0;

```

Proof of Lemma 4.

```

R<w0,w1,a01,a0j,a1j>:=PolynomialRing(Rationals(),5);
F:=FieldOfFractions(R);
I:=ideal<R|w0^2+w1^2-w0*w1*a01,a01^2+a0j^2+a1j^2-a01*a0j*a1j-4>;
x:=(w1^2-w0^2)/(a1j*w1-a0j*w0);
num:=Numerator(x^2-w0*a0j*x+w0^2);
num in I;

R<w0,w1,wi,wj,aij>:=PolynomialRing(Rationals(),5);
M:=Matrix(R,3,3,[
2*w0^2,w0^2+w1^2,w0^2+wi^2,
w0^2+w1^2,2*w1^2,w1^2+wi^2,
w0^2+wj^2,w1^2+wj^2,wi*wj*aij]);
Determinant(M) eq -(w1^2-w0^2)^2*(wi*wj*aij-wi^2-wj^2);

```

Proof of Lemma 5.

```

F<X1,X2,X3>:=FunctionField(Rationals(),3);
x01:=1/X1+X1;
x02:=1/X2+X2;
x03:=1/X3+X3;
x12:=X1/X2+X2/X1;
x13:=X1/X3+X3/X1;
x23:=X2/X3+X3/X2;
(X1*X2*X3+1)*(x01*x02+x03-x12) eq
(X1*X2+X3)*(x01*x02*x03+2-1/2*(x12*x03+x13*x02+x23*x01));

```

Proof of Theorem 2.

```

d:=3;
d2s:={&cat[[[i,j]:j in [i+1..d]:i in [0..d-1]]];
d2:={Seqset(s):s in d2s};
R:=PolynomialRing(Rationals(),#d2+2);
q:=R. (#d2+2);
r:=R. (#d2+1);
X:=func<i,j|R.Position(d2,{i,j})>;
g:=func<i,j,k|X(i,j)^2+X(i,k)^2+X(j,k)^2-X(i,j)*X(i,k)*X(j,k)-4>;
h:=func<i,j,k,l|(X(k,l)^2-4)*X(i,j)
-X(k,l)*(X(k,i)*X(l,j)+X(k,j)*X(l,i))
+2*(X(k,i)*X(k,j)+X(l,i)*X(l,j))>;

eigenP:=Matrix(R,4,4,[
1,1/2*q^2-q,1/2*q^2,q-2,
1,1/2*q,-1/2*q,-1,
1,-1/2*q+1,-1/2*q,q-2,
1,-1/2*q,1/2*q,-1]);
P:=func<i,j|eigenP[i+1,j+1]>;
n:={&+[P(0,i):i in [0..d]]};
n eq q^2-1;

e:=func<i|-n+&+[P(i,j)^2:j in [0..d]]
+&+[P(i,j[1])*P(i,j[2])*X(j[1],j[2]):j in d2s]>;
s3:={Setseq(x):x in Subsets({0..d},3)};
eq7:={g(i[1],i[2],i[3]):i in s3} cat
[h(0^i,1^i,2^i,3^i):i in Sym({0..d})] cat
[e(i):i in [1..d]] cat
[r^2-(17*q-1)*(q-1)];
I:=ideal<R|eq7>;

```

```

b1:=X(1,2)-2;
b2:=(q^2-4)*X(1,2)+2*q^2-12;
b3:=q^2*X(1,2)+2*q^2-4;
a12D:=q^2*(q+1)*(q-3);
a12Nplus:= 2*(-q^4+2*q^3+4*q^2-10*q+1 +(q-1)*r);
a12Nminus:=2*(-q^4+2*q^3+4*q^2-10*q+1 -(q-1)*r);
b4plus:=a12D*X(1,2)-a12Nplus;
b4minus:=a12D*X(1,2)-a12Nminus;
q*(q-1)*b1*b2*b3*b4plus*b4minus in I;

I1:=ideal<R|I,b1>;
c1:=X(0,1)+n-2;
a01D:=q^2-2*q-1;
a01N:=q^3-3*q^2-q+7;
c2:=a01D*X(0,1)-a01N;
X(0,3)+n-2 in I1;
(q-1)^2*c1*c2 in I1;
I11:=ideal<R|I1,c1>;
(q-1)*(q+1)*(q-2)*(q^2-5)*(X(1,3)-2) in I11;

I12:=ideal<R|I1,c2>;
a13D:=q^2-2*q-1;
a13N:=- (q^3-q^2-q-3);
-a13N+2*a13D eq (q+1)*(q^2-5);
-a13N-2*a13D eq (q-1)^3;
(q-1)*(q+1)*(q-2)*(q^2-5)*(a13D*X(1,3)-a13N) in I12;
a01:=a01N/a01D;
a03:=- (n-2);
a13:=a13N/a13D;
FA<w3>:=PolynomialRing(FieldOfFractions(R));
FA3:=FA/ideal<FA|w3^2+1-a03*w3>;
QA3:=FieldOfFractions(FA3);
w1:=QA3!((w3^2-1)/(a13*w3-a01));
w1 eq QA3!((- (q-3)*w3+(q-1))/(q^2-2*q-1));

I2:=ideal<R|I,b2>;
q*(q-1)*(q+1)*(q-3)*(n-4)*(X(0,2)+2) in I2;
q*(q-1)*(q+1)*(q-3)*(n-4)*(X(1,3)-2) in I2;
q*(q-1)*(q+1)*(q-3)*(n-4)*((n-3)*X(0,1)-2*(n-5)) in I2;

I3:=ideal<R|I,b3>;
c4:=q^2*X(0,2)+2*(n-1);

```

```

c5:=q*X(0,2)+2;
X(0,3)-2 in I3;
(q-1)^2*c4*c5 in I3;

I31:=ideal<R|I3,c4>;
(q-1)*(q+1)*(q-2)*(X(0,1)-2) in I31;

I32:=ideal<R|I3,c5>;
(q-1)*(q+1)*(q-2)*(q*X(0,1)+2) in I32;
q^2*X(1,2)+2*(n-1) in I32;
a02:=-2/q;
a01:=-2/q;
a12:=-2*(n-1)/q^2;
a01*a02-a12 eq 2;

I4:=ideal<R|I,b4plus>;
a01D:=2*q*(q+1);
a01N:=- (q-1)*(q-2)+(q+2)*r;
a01:=a01D*X(0,1)-a01N;
a02D:=2*q*(q-3);
a02N:=(q+2)*(q-1)-(q-2)*r;
a02:=a02D*X(0,2)-a02N;
a03D:=2*(q+1)*(q-3);
a03N:=5*q^2-2*q-19-(q-1)*r;
a03:=a03D*X(0,3)-a03N;
a12D:=q^2*(q+1)*(q-3);
a12N:=2*(-q^4+2*q^3+4*q^2-10*q+1+(q-1)*r);
a12:=a12D*X(1,2)-a12N;
a13D:=a02D;
a13N:=-a02N;
a13:=a13D*X(1,3)-a13N;
a23D:=a01D;
a23N:=-a01N;
a23:=a23D*X(2,3)-a23N;
q*(q+1)*(q-1)^2*(q-3)^2*(q^2-5)*a01 in I4;
q*(q+1)^2*(q-1)^2*(q-3)*(q^2-5)*a02 in I4;
a03 in I4;
a12 in I4;
q*(q+1)^2*(q-1)^2*(q-3)*(q^2-5)*a13 in I4;
q*(q+1)*(q-1)^2*(q-3)^2*(q^2-5)*a23 in I4;
b4pp:=Evaluate(b4plus,[0,0,0,2,0,0,r,q]);
(b4pp+2*(q-1)*r)^2-4*(q-1)^2*(17*q-1)*(q-1) eq

```



```

16*q^2*(q+1)*(q-1)^2*(q-3)*(q^2-5);
b4pm:=Evaluate(b4plus,[0,0,0,-2,0,0,r,q]);
(b4pm+2*(q-1)*r)^2-4*(q-1)^2*(17*q-1)*(q-1) eq
-64*q^2*(q+1)*(q-3);
q^2*(q+1)*(q-1)^2*(q-3)*(q^2-5)*(X(0,1)*X(0,2)+X(0,3)-X(1,2)) in I4;
I41:=ideal<R|I4,X(0,1)*X(0,2)*X(0,3)+2
-1/2*(X(1,2)*X(0,3)+X(1,3)*X(0,2)+X(2,3)*X(0,1))>;
q*(q-1)^2*(q^2-5)^2 in I41;

```

Conversely,

```

Rqr<rr,qq>:=PolynomialRing(Rationals(),2);
F:=FieldOfFractions(Rqr/ideal<Rqr|rr^2-(17*qq-1)*(qq-1)>);
d:=3;
d2s:={cat[[[i,j]:j in [i+1..d]]:i in [0..d-1]]};
d2:={Seqset(s):s in d2s};
q:=F!qq;
r:=F!rr;
R:=PolynomialRing(F,#d2);
X:=func<i,j|R.Position(d2,{i,j})>;
g:=func<i,j,k|X(i,j)^2+X(i,k)^2+X(j,k)^2-X(i,j)*X(i,k)*X(j,k)-4>;
h:=func<i,j,k,l|(X(k,l)^2-4)*X(i,j)
-X(k,l)*(X(k,i)*X(l,j)+X(k,j)*X(l,i))
+2*(X(k,i)*X(k,j)+X(l,i)*X(l,j))>;
eigenP:=Matrix(F,4,4,[
1,1/2*q^2-q,1/2*q^2,q-2,
1,1/2*q,-1/2*q,-1,
1,-1/2*q+1,-1/2*q,q-2,
1,-1/2*q,1/2*q,-1]);
P:=func<i,j|eigenP[i+1,j+1]>;
n:={P(0,i):i in [0..d]};
n eq q^2-1;
e:=func<i|-n+&+[P(i,j)^2:j in [0..d]]
+&+[P(i,j[1])*P(i,j[2])*X(j[1],j[2]):j in d2s]>;
s3:={Setseq(x):x in Subsets({0..d},3)};
eq7rq:={g(i[1],i[2],i[3]):i in s3} cat
[h(0^i,1^i,2^i,3^i):i in Sym({0..d})] cat
[e(i):i in [1..d]];
subs1:=[-n+2,-n+2,-n+2,2,2,2];
&and[Evaluate(f,subs1) eq 0:f in eq7rq];
a:=- (q-3)/(q^2-2*q-1);
b:=(q-1)/(q^2-2*q-1);
a^2+a*b*(-(n-2))+b^2 eq 1;

```

```

a01:=(q^3-3*q^2-q+7)/(q^2-2*q-1);
a*(-(n-2))+2*b eq a01;
a13:=(-q^3+q^2+q+3)/(q^2-2*q-1);
2*a-b*(n-2) eq a13;
subs2:=[a01,a01,-(n-2),2,a13,a13];
&and[Evaluate(f,subs2) eq 0:f in eq7rq];
a01_iii:=2*(n-5)/(n-3);
subs3:=[a01_iii,-2,a01_iii,-a01_iii,2,-a01_iii];
&and[Evaluate(f,subs3) eq 0:f in eq7rq];
a02_iv:=-2*(n-1)/(n+1);
subs4:=[2,a02_iv,2,a02_iv,2,a02_iv];
&and[Evaluate(f,subs4) eq 0:f in eq7rq];
subs5:=[-2/q,-2/q,2,-2*(n-1)/(n+1),-2/q,-2/q];
&and[Evaluate(f,subs5) eq 0:f in eq7rq];
a01D:=2*q*(q+1);
a01N:=- (q-1)*(q-2)+(q+2)*r;
a01:=a01N/a01D;
a02D:=2*q*(q-3);
a02N:=(q+2)*(q-1)-(q-2)*r;
a02:=a02N/a02D;
a03D:=2*(q+1)*(q-3);
a03N:=5*q^2-2*q-19-(q-1)*r;
a03:=a03N/a03D;
a12D:=q^2*(q+1)*(q-3);
a12N:=2*(-q^4+2*q^3+4*q^2-10*q+1+(q-1)*r);
a12:=a12N/a12D;
a13D:=a02D;
a13N:=-a02N;
a13:=a13N/a13D;
a23D:=a01D;
a23N:=-a01N;
a23:=a23N/a23D;
subs6:=[a01,a02,a03,a12,a13,a23];
&and[Evaluate(f,subs6) eq 0:f in eq7rq];

```

Proof of Corollary 1.

```

a01-2 eq -8*q*(q+1)*(q-3)^2/(2*q*(q+1)*(5*q^2+q+2+(q+2)*r));
a01+2 eq -8*q*(q+1)*(q^2-5)/(2*q*(q+1)*(3*q^2+7*q-2-(q+2)*r));

```

Isolation.

```

n:=15;

```

```

A0:=ScalarMatrix(n,1);
J:=Parent(A0)! [1:i in [1..n^2]];
L03:=LineGraph(OddGraph(3));
A1:=AdjacencyMatrix(L03);
A2:=A1^2-A1-4*A0;
A3:=J-A0-A1-A2;
DM:=DistanceMatrix(L03);
DM eq A1+2*A2+3*A3;

hermitianConjugate:=
  func<H|Parent(H)! [ComplexConjugate(x):x in Eltseq(Transpose(H))]>;

complexHadamard:=function(xyz)
  AA:=[ChangeRing(A,Parent(xyz[1])):A in [A1,A2,A3]];
  return A0+xyz[1]*AA[1]+xyz[2]*AA[2]+xyz[3]*AA[3];
end function;

spanCondition:=function(H)
  F:=Parent(H[1,1]);
  MnF:=Parent(H);
  n:=Nrows(H);
  Es:=[MnF|0:i in [1..n]];
  for i in [1..n] do
    Es[i][i,i]:=1;
  end for;
  EsF:=[MnF|e:e in Es];
  Hs:=hermitianConjugate(H);
  Vn:=VectorSpace(F,n^2);
  bracket:=sub<Vn|[Vn|Eltseq(v*Hs*w*H-Hs*w*H*v):v,w in EsF]>;
  return Dimension(bracket) eq n^2-2*n+1;
end function;

F<s>:=QuadraticField(-15);
y:=(-7+s)/8;
H:=complexHadamard([1,y,1]);
H*hermitianConjugate(H) eq n*A0;
spanCondition(H);

F<s>:=QuadraticField(-11);
x:=(5+s)/6;
H:=complexHadamard([x,-1,x]);
H*hermitianConjugate(H) eq n*A0;

```

```

not spanCondition(H);

F<s>:=QuadraticField(-15);
x:=(-1+s)/4;
H:=complexHadamard([x,x^(-1),1]);
H*hermitianConjugate(H) eq n*A0;
not spanCondition(H);

F<s>:=QuadraticField(201);
Z:=(53-3*s)/10;
R<T>:=PolynomialRing(F);
K<z>:=ext<F|T^2-Z*T+1>;
z+1/z eq Z;
x:=1/144*((-5*Z+31)*z-25*Z+155);
xb:=1/144*((-5*Z+31)*z^(-1)-25*Z+155);
y:=1/144*((25*Z-155)*z+5*Z-31);
yb:=1/144*((25*Z-155)*z^(-1)+5*Z-31);
x*xb eq 1;
y*yb eq 1;
H:=complexHadamard([x,y,z]);
H*hermitianConjugate(H) eq n*A0;
spanCondition(H);

```

Remark 1.

```

mustbe0:=func<a01v|(2*q*(q+1)*a01v+(q-1)*(q-2))^2-((q+2)*r)^2>;
mustbe0(0) eq -8*q*(q+1)*(q-1)*(2*q+7);
mustbe0(1) eq -8*q*(q+1)*(q^2+6*q-8);
mustbe0(-1) eq -8*q*(q+1)*(2*q^2+3*q-6);
mustbe0(2) eq 8*q*(q+1)*(q-3)^2;
mustbe0(-2) eq -8*q*(q+1)*(q^2-5);

```

Table 1.

```

HWminus1:=function(w)
  I3:={1..3};
  H3q:={w[i1]*w[i2]/w[i3]:i1,i2,i3 in I3
    |#[i:i in [i1,i2,i3]|i eq 3] ne 2};
  H3q4:={w[i1]*w[i2]/w[i3]:i1,i2,i3 in I3
    |#[i:i in [i1,i2,i3]|i eq 3] ne 2 and {i1,i2,i3} ne {1,3}};
  plus:={w[i]^2:i in I3} join H:H in [H3q,H3q4];
  return {(p join {x^(-1):x in p} join
    {w[i1]*w[i2]/(w[j1]*w[j2]):i1,i2,j1,j2 in I3})
    diff {1}:p in plus};

```

```

end function;
Rw<w1,w2,w3>:=FunctionField(Rationals(),3);
HWminus1([w1,w1,w1]) eq
  {&join{{w1^s,w1^(s*2)}:s in {1,-1}}};
HWminus1([w1,w1,w3]) eq
  {&join{{w^s,w^(s*2)}:s in {1,-1},w in {w1,w3}} join
  &join{{(w1^2/w3)^s,(w3/w1)^s,(w3/w1)^(s*2)}:s in {1,-1}}};
HWminus1([w1,-1,w1]) eq {-1} join
  &join{{s1*w1^s,s1*w1^(s*2)}:s,s1 in {1,-1}}};
HWminus1([1,w2,1]) eq {&join{{w2^s,w2^(s*2)}:s in {1,-1}}};
HWminus1([w1,w1^(-1),1]) eq {{w1^(s*k):s in {1,-1},k in {1..4}}};
HWminus1([w1,w2,-w1*w2]) eq {-1} join
  {s0*w^(s*k):w in {w1,w2},s,s0 in {1,-1},k in {1,2}} join
  &join{{s0*w1^s1*w2^s2,(w1^s1*w2^s2)^2}:s0,s1,s2 in {1,-1}}
  join &join{{s0*(w1^2*w2^(-1))^s,s0*(w1^(-1)*w2^2)^s
  :s,s0 in {1,-1}}};
// (i)
Rq<q>:=FunctionField(Rationals());
(-q^2+3)^2-2 eq q^4-6*q^2+7;
// (ii)
Rw3<w3>:=FunctionField(Rq);
A:=- (q-3)/(q^2-2*q-1);
B:=(q-1)/(q^2-2*q-1);
(A^2+B^2-1)/(A*B) eq q^2-3;
w1:=A*w3+B;
(A/w3+B)-1/w1 eq 1/w1*A*B*(w3+1/w3+(q^2-3));
// (iii)
(2*(q^2-6)/(q^2-4))^2-2 eq 2*(q^4-16*q^2+56)/(q^2-4)^2;
// (iv)
(-2*(q^2-2)/q^2)^2-2 eq 2*(q^4-8*q^2+8)/q^4;

```

Proof of Proposition 1. (iii) $\not\cong$ (vi)

```

Rq<q>:=FunctionField(Rationals());
k3a:=(q^2-6)/(q^2-4);
k3b:=2*(q^4-16*q^2+56)/(q^2-4)^2;
ra1:=(k3a+(q-1)*(q-2)/(2*q*(q+1)))/(q+2);
fac:=Factorization(Numerator(ra1^2-(17*q-1)*(q-1)));
#fac eq 1 and Degree(fac[1][1]) gt 1;
ra2:=(-k3a+(q-1)*(q-2)/(2*q*(q+1)))/(q+2);
fac:=Factorization(Numerator(ra2^2-(17*q-1)*(q-1)));
#fac eq 1 and Degree(fac[1][1]) gt 1;
rb1:=(k3b+(q-1)*(q-2)/(2*q*(q+1)))/(q+2);

```

```

fac:=Factorization(Numerator(rb1^2-(17*q-1)*(q-1)));
#fac eq 1 and Degree(fac[1][1]) gt 1;
rb2:=(-k3b+(q-1)*(q-2)/(2*q*(q+1)))/(q+2);
fac:=Factorization(Numerator(rb2^2-(17*q-1)*(q-1)));
#fac eq 1 and Degree(fac[1][1]) gt 1;

```

(i) $\not\equiv$ (ii)

```

n:=q^2-1;
Numerator((2-n)-(q^3-3*q^2-q+7)/(q^2-2*q-1))
eq -(q-2)*(q+1)*(q^2-5);
Numerator((n^2-4*n+2)-(q^3-3*q^2-q+7)/(q^2-2*q-1))
eq (q-2)*(q+1)^2*(q^3-2*q^2-4*q+7);

```

(iv) $\not\equiv$ (v)

```

Numerator(-2*(n-1)/(n+1)-(-2)/q)
eq -2*(q-2)*(q+1);
Numerator(2*(n^2-6*n+1)/(n+1)^2-(-2)/q)
eq 2*(q-2)*(q+1)*(q^2+2*q-4);

```

Proof of Proposition 2.

```

ra1:=(2+(q-1)*(q-2)/(2*q*(q+1)))/(q+2);
fac:=Factorization(Numerator(ra1^2-(17*q-1)*(q-1)));
#fac eq 1 and Degree(fac[1][1]) gt 1;
ra2:=(-2+(q-1)*(q-2)/(2*q*(q+1)))/(q+2);
fac:=Factorization(Numerator(ra2^2-(17*q-1)*(q-1)));
#fac eq 1 and Degree(fac[1][1]) gt 1;

```

Proof of Lemma 13.

```

R<r,q>:=PolynomialRing(Rationals(),2);
F:=FieldOfFractions(R);
n:=q^2-1;
fr:=r^2-(17*q-1)*(q-1);
B1:=Matrix(R,4,4,[
0,1,0,0,
q^2/2-q,(q-2)^2/4,(q-2)^2/4,q*(q-4)/4,
0,q*(q-2)/4,q*(q-2)/4,q^2/4,
0,(q-4)/2,(q-2)/2,0]);

B2:=Matrix(R,4,4,[
0,0,1,0,
0,q*(q-2)/4,q*(q-2)/4,q^2/4,
q^2/2,q^2/4,q^2/4,q^2/4,
0,q/2,(q-2)/2,0]);

```

```

B3:=Matrix(R,4,4,[
0,0,0,1,
0,(q-4)/2,(q-2)/2,0,
0,q/2,(q-2)/2,0,
q-2,0,0,q-3]);

B0:=ScalarMatrix(4,R!1);
BB:=[B0,B1,B2,B3];
pijk:=func<i,j,k|BB[i+1][j+1,k+1]>;

isSymNbas:=function(ajk)
  aijs:=[[ajk[1],ajk[2],ajk[3]],
        [1,ajk[4],ajk[5]],[1,1,ajk[6]]];
  aij:=func<i,j|aijs[i+1,j]>;
  ff:=[F|&+[pijk(j,k,i)*(aij(j,k)^2-2):j,k in [0..3]|j lt k
        +&+[pijk(j,j,i):j in [0..3]]:i in [1..3]];
  II:=[ideal<R|fr, Numerator(ff[k])>:k in [1..3]];
  return [Basis(EliminationIdeal(II[k],{q})):k in [1..3]];
end function;

// case (i)
subs1:=[-n+2,-n+2,-n+2,2,2,2];
bass:=isSymNbas(subs1);
[Factorization(bass[i][1]):i in [1..3]] eq
[[<q-2,1>,<q-1,1>,<q+1,1>,<q+2,1>]:i in [1..3]];

// case (ii)
a01:=(q^3-3*q^2-q+7)/(q^2-2*q-1);
a13:=(-q^3+q^2+q+3)/(q^2-2*q-1);
subs2:=[a01,a01,-(n-2),2,a13,a13];
bass:=isSymNbas(subs2);
b1:=q^4-10*q^2+4*q+17;
fac:=[[<q-2,1>,<q-1,1>,<q+1,1>,<b1,1>],
      [<q-2,1>,<q-1,1>,<q+1,1>,<b1,1>],
      [<q-2,1>,<q-1,1>,<q+1,1>,<q+2,1>]];
[Factorization(bass[i][1]):i in [1..3]] eq fac;

// case (iii)
a01_iii:=2*(n-5)/(n-3);
subs3:=[a01_iii,-2,a01_iii,-a01_iii,2,-a01_iii];
bass:=isSymNbas(subs3);

```

```

b1:=q^6-13*q^4+28*q^2+64;
b2:=q^4-9*q^2+24;
[Factorization(bass[i][1]):i in [1..3]] eq
  [[<b1,1>],[<b2,1>],[<b1,1>]];

// case (iv)
a02_iv:=-2*(n-1)/(n+1);
subs4:=[2,a02_iv,2,a02_iv,2,a02_iv];
bass:=isSymNbas(subs4);
bass eq [[(q-2)*(q-1)*(q+1)*(q+2)]:i in [1..3]];

// case (v)
subs5:=[-2/q,-2/q,2,-2*(n-1)/(n+1),-2/q,-2/q];
bass:=isSymNbas(subs5);
b1:=(q-2)*(q-1)*(q+1)*(q^2-2*q-4);
b3:=(q-2)*(q-1)*(q+1)*(q+2);
bass eq [[b1],[b1],[b3]];

// case (vi)
a01D:=2*q*(q+1);
a01N:=- (q-1)*(q-2)+(q+2)*r;
a01:=a01N/a01D;
a02D:=2*q*(q-3);
a02N:=(q+2)*(q-1)-(q-2)*r;
a02:=a02N/a02D;
a03D:=2*(q+1)*(q-3);
a03N:=5*q^2-2*q-19-(q-1)*r;
a03:=a03N/a03D;
a12D:=q^2*(q+1)*(q-3);
a12N:=2*(-q^4+2*q^3+4*q^2-10*q+1+(q-1)*r);
a12:=a12N/a12D;
a13:=-a02;
a23:=-a01;
bass:=isSymNbas([a01,a02,a03,a12,a13,a23]);
bass eq [
  [q^3*(q-3)^2*(q-1)*(q+1)^2*
   (q^9-q^8-12*q^7+14*q^6+49*q^5
   +51*q^4-894*q^3-464*q^2+4664*q-272)],
  [q^3*(q-3)^2*(q-2)*(q-1)*(q+1)^2*
   (q^7+3*q^6-4*q^5+2*q^4+57*q^3-q^2-86*q+92)],
  [q^2*(q-3)^2*(q-1)*(q+1)^2*
   (q^8-2*q^7+66*q^5-273*q^4-288*q^3+1344*q^2-288*q+16)]];

```


Proof of Proposition 4.

```

R<c111,c112,c121,c122,c211,c212,c221,c222,
  w1,w2case4,w3case2,r,q>
:=PolynomialRing(Rationals(),13);
varname:=[[i,j,k]:i,j,k in [1,2]];
c:=func<i,j,k|R.Position(varname,[i,j,k])>;
fr:=r^2-(17*q-1)*(q-1);
w0:=1;

B1:=Matrix(R,4,4,[
0,1,0,0,
q^2/2-q,(q-2)^2/4,(q-2)^2/4,q*(q-4)/4,
0,q*(q-2)/4,q*(q-2)/4,q^2/4,
0,(q-4)/2,(q-2)/2,0]);

B2:=Matrix(R,4,4,[
0,0,1,0,
0,q*(q-2)/4,q*(q-2)/4,q^2/4,
q^2/2,q^2/4,q^2/4,q^2/4,
0,q/2,(q-2)/2,0]);

B3:=Matrix(R,4,4,[
0,0,0,1,
0,(q-4)/2,(q-2)/2,0,
0,q/2,(q-2)/2,0,
q-2,0,0,q-3]);

B0:=ScalarMatrix(4,R!1);
BB:=[B0,B1,B2,B3];
pijk:=func<i,j,k|BB[i+1][j+1,k+1]>;

cijk:=function(i,j,k)
  if 0 in {i,j,k} then
    if [i,j,k] in {[0,3,3],[3,0,3],[3,3,0]} then
      return 1;
    else
      return 0;
    end if;
  elif 3 in {i,j,k} then
    if {3} eq {i,j,k} then
      return pijk(3,3,3)-1;

```

```

else
  return 0;
end if;
else
  return c(i,j,k);
end if;
end function;

fx:=[cijjk(1,j,k)+cijjk(2,j,k)-pijk(j,k,3):j,k in [1,2]];
fy:=[cijjk(j,1,k)+cijjk(j,2,k)-pijk(j,k,3):j,k in [1,2]];
fz:=[cijjk(j,k,1)+cijjk(j,k,2)-pijk(j,k,3):j,k in [1,2]];
fxyz:=fx cat fy cat fz;

test:=function(fa,ww)
  alpha:=func<i|ww[i+1]>;
  ff:=%+[cijjk(i,j,k)*alpha(i)^2/(alpha(j)*alpha(k))
  :i,j,k in [0..3]];
  gg:=%+[cijjk(i,j,k)*alpha(j)*alpha(k)/alpha(i)^2
  :i,j,k in [0..3]];
  I:=ideal<R|[fr,fa,Numerator(ff),Numerator(gg)] cat fxyz>;
  return Basis(EliminationIdeal(I,{q}));
end function;

test2:=function(fa,ww)
  alpha:=func<i|ww[i+1]>;
  t1:=%+[pijk(i,j,1)*pijk(3,k,i)*alpha(i)^2/(alpha(j)*alpha(k))
  :i,j,k in [0..3]];
  t2:=%+[pijk(i,j,2)*pijk(3,k,i)*alpha(i)^2/(alpha(j)*alpha(k))
  :i,j,k in [0..3]];
  I1:=ideal<R|[fr,fa,Numerator(t1)]>;
  I2:=ideal<R|[fr,fa,Numerator(t2)]>;
  return [Basis(EliminationIdeal(I1,{q})),
  Basis(EliminationIdeal(I2,{q}))];
end function;

// case (i)
fa1:=w1^2+(q^2-3)*w1+1;
ww:=[w0,w1,w1,w1];
test(fa1,ww) eq [(q+1)*(q-1)*(q^2-5)];
test2(fa1,ww) eq [[(q-2)*(q+1)*(q-1)*(q^2-5)]:i in [1,2]];

// case (ii)

```

```

fa3:=w3case2^2+(q^2-3)*w3case2+1;
w2:=(-(q-3)*w3case2+q-1)/(q^2-2*q-1);
ww:=[w0,w2,w2,w3case2];
test(fa3,ww) eq [(q+1)*(q-1)*(q^2-5)];
test2(fa3,ww) eq
  [[(q-2)*(q+1)*(q-1)*(q^2-5)*(q^2-2*q-1)^2]:i in [1,2]];

// case (iii)
fa1:=R!((q^2-4)*(w1^2-2*(q^2-6)/(q^2-4)*w1+1));
ww:=[w0,w1,-1,w1];
test(fa1,ww) eq [(q^2+4)*(q^2-5)];
test2(fa1,ww) eq
  [[(q-2)*(q+1)*(q^2-5)*(q^5-5*q^4+12*q^3-24*q^2+64)]:i in [1,2]];

// case (iv)
fa2:=R!(q^2*(w2case4^2+2*(q^2-2)/q^2*w2case4+1));
ww:=[w0,1,w2case4,1];
test(fa2,ww) eq [q^2*(q+1)*(q-1)];
test2(fa2,ww) eq [[q^2*(q-2)*(q+1)*(q-1)]:i in [1,2]];

// case (v)
fa1:=R!(q*(w1^2+2/q*w1+1));
ww:=[w0,w1,1/w1,1];
test(fa1,ww) eq [q^2*(q+1)*(q-1)];
test2(fa1,ww) eq [[q*(q-2)*(q+1)*(q-1)]:i in [1,2]];

// case (vi)
a01D:=2*q*(q+1);
a01N:=-(q-1)*(q-2)+(q+2)*r;
a01:=a01N/a01D;
a02D:=2*q*(q-3);
a02N:=(q+2)*(q-1)-(q-2)*r;
a02:=a02N/a02D;
a03D:=2*(q+1)*(q-3);
a03N:=5*q^2-2*q-19-(q-1)*r;
a03:=a03N/a03D;
a12D:=q^2*(q+1)*(q-3);
a12N:=2*(-q^4+2*q^3+4*q^2-10*q+1+(q-1)*r);
a12:=a12N/a12D;
a13:=-a02;
a23:=-a01;
fa1:=a01D*w1^2-a01N*w1+a01D;

```

```
w2:=(a01*w1-2)/(a12*w1-a02);  
w3:=(a01*w1-2)/(a13*w1-a03);  
ww:=[w0,w1,w2,w3];  
test(fa1,ww) eq [q^12*(q-3)^12*(q-1)*(q+1)^6*(q-1/9)*(q^2-5)^3];  
test2(fa1,ww) eq [[q^9*(q-3)^12*(q-1)^2*(q+1)^7*(q^2-5)^4*  
(q^4+6/19*q^3-48/19*q^2+8/19*q+16/19)]:i in [1,2]];
```

APPENDIX B. DETAILED PROOF OF REMARK 2

Lemma 14. *Let d be a square-free positive integer, and let u, v be positive integers satisfying $u^2 - dv^2 = 1$ and $u > 1$. Let a be a positive integer. If x and y are positive integers satisfying $x^2 - dy^2 = a$ and $x > \sqrt{a(u+1)}/2$, then there exist integers x_0, y_0, n with $0 < x_0 \leq \sqrt{a(u+1)}/2$ such that*

$$x + \sqrt{d}y = (u + \sqrt{d}v)^n(x_0 + \sqrt{d}y_0). \quad (48)$$

Proof. Suppose that the statement is false, and choose the minimal counterexample $x > \sqrt{a(u+1)}/2$ with $x^2 - dy^2 = a$, not expressible in the form (48). Then $2x^2 > a(u+1)$, so $\frac{a}{x^2} < \frac{2}{u+1}$. Thus

$$\begin{aligned} u^2x^2 &> (u^2 - 1)(x^2 - a) = d^2v^2y^2 \\ &= (u - 1)^2x^2\frac{u+1}{u-1}\left(1 - \frac{a}{x^2}\right) \\ &> (u - 1)^2x^2\frac{u+1}{u-1}\left(1 - \frac{2}{u+1}\right) \\ &= (u - 1)^2x^2. \end{aligned}$$

This implies $ux > dvy > (u - 1)x$, or equivalently,

$$x > ux - dvy > 0.$$

Set

$$\begin{aligned} x_1 &= ux - dvy, \\ y_1 &= |uy - vx|. \end{aligned}$$

Then $x > x_1 > 0$ and $y_1 \geq 0$.

$$\begin{aligned} x_1 \pm \sqrt{d}y_1 &= ux - dvy \pm \sqrt{d}(uy - vx) \\ &= (u \mp \sqrt{d}v)(x \pm \sqrt{d}y), \end{aligned} \quad (49)$$

and so

$$\begin{aligned} x_1^2 - dy_1^2 &= (u^2 - dv^2)(x - dy)^2 \\ &= a. \end{aligned}$$

Note that if $y_1 = 0$, then $x_1 = \sqrt{a}$, so

$$x_1 \leq \sqrt{a(u+1)}/2. \quad (50)$$

Since x is the minimal counterexample, we have either (50), or there exist integers x_0, y_0, n with $0 < x_0 \leq \sqrt{a(u+1)}/2$ such that

$$x_1 + \sqrt{d}y_1 = (u + \sqrt{d}v)^n(x_0 + \sqrt{d}y_0). \quad (51)$$

In the former case, (49) implies

$$x + \sqrt{d}y = (u + \sqrt{d}v)(x_1 + \sqrt{d}y_1),$$

so (48) holds by setting $(x_0, y_0, n) = (x_1, y_1, 1)$.

In the latter case, (49) and (51) imply

$$x + \sqrt{d}y = (u + \sqrt{d}v)^{n+1}(x_0 + \sqrt{d}y_0),$$

so (48) again holds.

In either case, we obtain a contradiction to the the assumption that x is a counterexample. \square

Example 1. Since the Pell equation

$$u^2 - 17v^2 = 1$$

has a solution $(u, v) = (33, 8)$, Lemma 14 implies that all solutions of the equation

$$x^2 - 17y^2 = 64 \tag{52}$$

can be expressed by those with $x < \sqrt{64(33+1)/2} = 8\sqrt{17}$ and powers of $33 + 8\sqrt{17}$. Observe that the non-negative solutions (x, y) of (52) with $x < 8\sqrt{17}$ are

$$\{(8, 0), (9, 1), (26, 6)\}.$$

Now Lemma 14 implies

$$\begin{aligned} & \{x + \sqrt{17}y \mid x, y \in \mathbb{Z}, x, y > 0, x^2 - 17y^2 = 64\} \\ & \subset \{(33 + 8\sqrt{17})^n \cdot 8 \mid n \in \mathbb{Z}\} \cup \{(33 + 8\sqrt{17})^n(9 + \sqrt{17}) \mid n \in \mathbb{Z}\} \\ & \quad \cup \{(33 + 8\sqrt{17})^n(26 + 6\sqrt{17}) \mid n \in \mathbb{Z}\} \end{aligned}$$

`33^2-17*8^2 eq 1;`

`[v:v in [1..8]|IsSquare(17*v^2+1)] eq [8];`

`[x:x in [1..Floor(8*Sqrt(17))]|IsSquare((x^2-64)/17)]`

`eq [8,9,26];`

Lemma 15. *Let q be an even positive integer with $q \geq 4$. Then $\sqrt{(q-1)(17q-1)}$ is an integer if and only if*

$$\begin{aligned} q & \in \left\{ \frac{1}{34}(\text{tr}((2177 + 528\sqrt{17})^n(433 + 105\sqrt{17})) + 18) \mid n \in \mathbb{Z} \right\} \tag{53} \\ & = \{ \dots, 41210, 10, 26, 110890, 482812730, \dots \}. \end{aligned}$$

Proof. Setting $x = 17q - 9$ and $y = \sqrt{(q-1)(17q-1)}$, we have

$$\begin{aligned} x^2 - 17y^2 & = (17q - 9)^2 - 17r^2 \\ & = 17^2q^2 - 2 \cdot 17 \cdot 9q + 81 - 17(q-1)(17q-1) \\ & = 64. \end{aligned}$$

Thus, by Example 1, we have

$$\begin{aligned} x + \sqrt{17}y &\in \{(33 + 8\sqrt{17})^n \cdot 8 \mid n \in \mathbb{Z}\} \\ &\cup \{(33 + 8\sqrt{17})^n(9 + \sqrt{17}) \mid n \in \mathbb{Z}\} \\ &\cup \{(33 + 8\sqrt{17})^n(26 + 6\sqrt{17}) \mid n \in \mathbb{Z}\} \end{aligned}$$

Since q is even, we have $x \equiv -9 \pmod{34}$. On the other hand, we have

$$\begin{aligned} (33 + 8\sqrt{17})^n \cdot 8 = a + b\sqrt{17} &\implies a \equiv 8(-1)^n \pmod{34}, \\ (33 + 8\sqrt{17})^n(9 + \sqrt{17}) = a + b\sqrt{17} &\implies a \equiv 9(-1)^n \pmod{34}, \\ (33 + 8\sqrt{17})^n(26 + 6\sqrt{17}) = a + b\sqrt{17} &\implies a \equiv 8(-1)^{n+1} \pmod{34}, \end{aligned}$$

Thus

$$\begin{aligned} x + \sqrt{17}y &\in \{(33 + 8\sqrt{17})^{2n+1}(9 + \sqrt{17}) \mid n \in \mathbb{Z}\} \\ &= \{(2177 + 528\sqrt{17})^n(433 + 105\sqrt{17}) \mid n \in \mathbb{Z}\} \end{aligned} \quad (54)$$

Thus

$$x \in \left\{ \frac{1}{2} \operatorname{tr}((2177 + 528\sqrt{17})^n(433 + 105\sqrt{17})) \mid n \in \mathbb{Z} \right\} \quad (55)$$

Since $q = (x + 9)/17$, we obtain (53).

Conversely, if (53) holds, then (55) holds, and hence (54) holds for some integer y . Moreover, (54) implies $x^2 - 17y^2 = 64$, and hence $y^2 = (17q - 1)(q - 1)$. \square

```
K<s>:=QuadraticField(17);
(33+8*s)^2 eq 2177+528*s;
(33+8*s)*(9+s) eq 433+105*s;
[1/34*(Trace((2177+528*s)^n*(433+105*s))+18):n in [-2..2]]
eq [41210,10,26,110890,482812730];
```

When (53) holds, $a_{0,1}$ is rational. This means that w_1 is at most quadratic. The following computation suggests that w_1 is always quadratic.

```
K<s>:=QuadraticField(17);
qs:=[1/34*(Trace((2177+528*s)^n*(433+105*s))+18):n in [-1000..1000]];
w1rational:=function(q)
t,r:=IsSquare((17*q-1)*(q-1));
if t then
a01:=(-(q-1)*(q-2)+(q+2)*r)/(2*q*(q+1));
return IsSquare(a01^2-4);
else
return false;
```

```
end if;  
end function;  
&and[not D(q):q in qs];
```

KOBE GAKUIN UNIVERSITY, KOBE, 650-8586, JAPAN
E-mail address: ikuta@law.kobegakuin.ac.jp

TOHOKU UNIVERSITY, SENDAI, 980-8579, JAPAN
E-mail address: munemasa@math.is.tohoku.ac.jp