

Search for CP violation in $D^0 \rightarrow h^+h^-$ decays at CDF

SABATO LEO

*University of Illinois at Urbana-Champaign, 61801- Urbana, IL, USA,
for the CDF Collaboration.*

I report on a measurement of CP -violating asymmetries (A_Γ) between effective lifetimes of D^0 or \bar{D}^0 in fully reconstructed $D^0 \rightarrow K^+K^-$ and $D^0 \rightarrow \pi^+\pi^-$ decays collected in $p\bar{p}$ collisions by the Collider Detector at Fermilab experiment. The full CDF data set corresponding to 9.7 fb^{-1} of integrated luminosity is used. The flavor of the charm meson at production is determined by exploiting the strong-interaction decay $D^{*+} \rightarrow D^0\pi^+$, while the contamination from mesons originated in b -hadron decays is evaluated and subtracted from the sample. Signal yields as functions of the observed decay-time distributions are extracted from maximum likelihood fits and used to measure the asymmetries. The results, $A_\Gamma(K^+K^-) = (-1.9 \pm 1.5 \text{ (stat)} \pm 0.4 \text{ (syst)}) \times 10^{-3}$ and $A_\Gamma(\pi^+\pi^-) = (-0.1 \pm 1.8 \text{ (stat)} \pm 0.3 \text{ (syst)}) \times 10^{-3}$, and their combination, $A_\Gamma = (-1.2 \pm 1.2) \times 10^{-3}$, are consistent with the SM predictions and other experimental determinations.

PRESENTED AT

8th International Workshop on the CKM Unitarity Triangle
(CKM 2014), Vienna, Austria, September 8-12, 2014.

1 Introduction

In charm transitions, the standard model (SM) predicts CP -violating effects not exceeding $O(10^{-2})$ [1]. Indeed, no CP -violating effects have been firmly established yet in charm dynamics [2].

The decay-time-dependent rate asymmetries of charm mesons two body hadron decays ($D \rightarrow h^+h^-$, $h = K, \pi$),

$$\mathcal{A}_{CP}(t) = \frac{d\Gamma(D^0 \rightarrow h^+h^-)/dt - d\Gamma(\bar{D}^0 \rightarrow h^+h^-)/dt}{d\Gamma(D^0 \rightarrow h^+h^-)/dt + d\Gamma(\bar{D}^0 \rightarrow h^+h^-)/dt}, \quad (1)$$

probe non-SM physics contributions in the *oscillation* and *penguin* transition amplitudes which could be affected by non-SM contributions that enhance the magnitude of the observed CP violation w.r.t. SM expectation. The asymmetry $\mathcal{A}_{CP}(t)$ includes both direct and indirect CP violation effects. The slow oscillations rate [2] of charm mesons allows approximating Eq. (1) as [3],

$$\mathcal{A}_{CP}(t) \approx \mathcal{A}_{CP}^{\text{dir}}(h^+h^-) - \frac{\langle t \rangle}{\tau} A_{\Gamma}(h^+h^-) \text{ with } A_{\Gamma} = \frac{\hat{\tau}(\bar{D}^0 \rightarrow h^+h^-) - \hat{\tau}(D^0 \rightarrow h^+h^-)}{\hat{\tau}(\bar{D}^0 \rightarrow h^+h^-) + \hat{\tau}(D^0 \rightarrow h^+h^-)} \quad (2)$$

where $\langle t \rangle$ is the sample mean of decay time, τ is the CP -averaged D lifetime [2], $\mathcal{A}_{CP}^{\text{dir}}$ is related to direct CP violation, and A_{Γ} is the asymmetry between the *effective* lifetimes $\hat{\tau}$ of D^0 and \bar{D}^0 and is mostly due to indirect CP violation. Recent measurements of A_{Γ} [4] showed consistency with CP symmetry with $O(10^{-3})$ uncertainties. However, additional determinations with comparable precision may improve the knowledge of CP violation in the charm sector. In this note I report a measurement of A_{Γ} using fully reconstructed $D^0 \rightarrow K^+K^-$ and $D^0 \rightarrow \pi^+\pi^-$ decays collected in $p\bar{p}$ collisions by the Collider Detector at Fermilab experiment. The full CDF data set corresponding to 9.7 fb^{-1} of integrated luminosity is used.

2 Selection and reconstruction

Online data selection is based on pairs of charged particles displaced from the $p\bar{p}$ collision point. Offline, a D candidate is reconstructed using two oppositely charged tracks fit to a common decay vertex. A charged particle with $p_T > 400 \text{ MeV}/c$ is associated with each D candidate to form $D^{*\pm}$ candidates. Constraining the $D^{*\pm}$ decay vertex to lie on the beam-line results in a 25% improvement in $D^{*\pm}$ mass resolution w.r.t Ref. [3]. Ref. [3] details the offline selection. The h^+h^- mass of selected candidates is required to be within about $24 \text{ MeV}/c^2$ of the known D^0 mass, m_{D^0} [2], to separate $D \rightarrow K^+K^-$ and $D \rightarrow \pi^+\pi^-$ samples. Final selected samples contain 6.1×10^5 $D^0 \rightarrow K^+K^-$, 6.3×10^5 $\bar{D}^0 \rightarrow K^+K^-$, 2.9×10^5 $D^0 \rightarrow \pi^+\pi^-$,




Figure 1: Distributions of $D\pi^\pm$ mass with fit results overlaid for the $D \rightarrow K^+K^-$ sample (a). Distributions of D^0 impact parameter with fit results overlaid for background-subtracted $D^{*+} \rightarrow D^0(\rightarrow \pi^+\pi^-)\pi_s^+$ decays restricted to the decay-time bin $6.16 < t/\tau < 20$ (b).

and $3.0 \times 10^5 \bar{D}^0 \rightarrow \pi^+\pi^-$ signal events. The main backgrounds are real D^0 decays associated with random pions or random combinations of three tracks (combinatorics) for the $\pi^+\pi^-$ sample, while the K^+K^- sample is also polluted by misreconstructed multibody charm meson decays (i.e. $D^0 \rightarrow h^-\pi^+\pi^0$ and $D^0 \rightarrow h^-\ell^+\nu_\ell$, where ℓ is a muon or an electron), Figure. 1.

3 Determination of the asymmetry

The flavor-conserving strong-interaction processes $D^{*+} \rightarrow D^0\pi^+$ and $D^{*-} \rightarrow \bar{D}^0\pi^-$ allow identification of the initial flavor through the charge of the low-momentum π meson (soft pion, π_s). D^0 or \bar{D}^0 subsamples are thus divided in equally populated 30 bins of decay time between 0.15τ and 20τ . In each bin, the average decay-time $\langle t \rangle$ is determined from a sample of about $13 \times 10^6 D^{*\pm} \rightarrow D(\rightarrow K^\mp\pi^\pm)\pi_s^\pm$ signal decays. Signal and background yields in the signal region are determined in each decay-time bin, and for each flavor, through χ^2 fits of the $D\pi_s^\pm$ mass distribution. The functional form of the signal shapes is determined from simulation [3], with parameters tuned in the sample of $D \rightarrow K^\mp\pi^\pm$ decays, independently for each D flavor and decay-time bin. The resulting signal-to-background proportions are used to construct signal-only distributions of the D impact parameter (IP). In each bin and for each flavor background-subtracted IP distributions are formed by subtracting IP distributions of background candidates, sampled in the $2.015 < M(D\pi^\pm) < 2.020 \text{ GeV}/c^2$

region for the $\pi^+\pi^-$ sample, from IP distributions of signal candidates which have $M(D\pi_s^\pm)$ within $2.4 \text{ MeV}/c^2$ of the known D^{*+} mass [2]. Contamination from multi-body decays in the K^+K^- sample is taken into account using as background candidates in the sideband $m_{D^0} - 64 \text{ MeV}/c^2 < M(K^+K^-) < m_{D^0} - 40 \text{ MeV}/c^2$ and with $M(D\pi_s^\pm)$ within $2.4 \text{ MeV}/c^2$ of the known $D^{*\pm}$ mass. A χ^2 fit of these signal-only IP distributions identifies $D^{*\pm}$ mesons from b -hadron decays (*secondary*) and determines the yields of charm (N_{D^0}) and anticharm ($N_{\bar{D}^0}$) mesons directly produced in the $p\bar{p}$ collision (*primary*). Double-Gaussian models are used for both the primary and secondary components. The parameters of the primary component are derived from a fit of candidates in the first decay-time bin ($t/\tau < 1.18$), where any bias from the $\mathcal{O}(\%)$ secondary contamination is negligible, and fixed in all fits. The parameters of the secondary component are determined by the fit independently for each decay-time bin, Figure 1. The yields are then combined into the asymmetry $A = (N_{D^0} - N_{\bar{D}^0})/(N_{D^0} + N_{\bar{D}^0})$, which is fit with the linear function in Eq. (2). The slope of the function, which yields A_Γ , is extracted using a χ^2 fit. The fit is shown in Fig. 2 and yields $A_\Gamma(K^+K^-) = (-1.9 \pm 1.5 \text{ (stat)}) \times 10^{-3}$ and $A_\Gamma(\pi^+\pi^-) = (-0.1 \pm 1.8 \text{ (stat)}) \times 10^{-3}$. In both samples, we observe a few percent value for $A(0)$, due to the known detector-induced asymmetry in the soft-pion reconstruction efficiency [3]. The independence of instrumental asymmetries from decay time is demonstrated by the analysis of $D \rightarrow K^\mp\pi^\pm$ decays, where no indirect CP violation occurs and instrumental asymmetries are larger; an asymmetry compatible with zero is found, $(-0.5 \pm 0.3) \times 10^{-3}$.

For the $\pi^+\pi^-$ analysis, the dominant systematic uncertainty of 0.028% arises from the choice of the impact-parameter shape of the secondary component whereas for the K^+K^- sample this effect only contributes 0.013%. The choice of the background sideband has a dominant effect in the K^+K^- analysis (0.038%) and a minor impact (0.010%) on the $\pi^+\pi^-$ result.

4 Conclusions

A measurement of the difference in effective lifetime between anticharm and charm mesons reconstructed in $D^0 \rightarrow K^+K^-$ and $D^0 \rightarrow \pi^+\pi^-$ decays using the full CDF data set is reported. The final results,

$$\begin{aligned} A_\Gamma(\pi^+\pi^-) &= (-0.1 \pm 1.8 \text{ (stat)} \pm 0.3 \text{ (syst)}) \times 10^{-3}, \\ A_\Gamma(K^+K^-) &= (-1.9 \pm 1.5 \text{ (stat)} \pm 0.4 \text{ (syst)}) \times 10^{-3}, \end{aligned} \quad (3)$$

are consistent with CP symmetry and combined to yield $A_\Gamma = (-1.2 \pm 1.2) \times 10^{-3}$ [5]. The results are also consistent with the current best results [4], have the second best precisions, and contribute to improve the global constraints on indirect CP violation in charm meson dynamics.




Figure 2: Effective lifetime asymmetries as functions of decay time for the (a) $D \rightarrow K^+K^-$ and (b) $D \rightarrow \pi^+\pi^-$ samples. Results of fits not allowing for (red dotted line) and allowing for (blue solid line) CP violation are overlaid.

References

- [1] S. Bianco, F. L. Fabbri, D. Benson, and I. I. Bigi, Riv. Nuovo Cim. **26N7**, 1 (2003);
- [2] Y. Amhis *et al.* (Heavy Flavor Averaging Group), arXiv:1207.1158 and online update at <http://www.slac.stanford.edu/xorg/hfag>; K.A. Olive *et al.* (Particle Data Group), Chin. Phys. C **38**, 090001 (2014).
- [3] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. D **85**, 012009 (2012); A. Di Canto, Ph.D. thesis, University of Pisa, 2011, FERMILAB-THESIS-2011-29; T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **109**, 111801 (2012).
- [4] M. Staric *et al.* (Belle Collaboration), Phys. Rev. Lett. **98**, 211803 (2007) and preliminary update in arXiv:1212.3478; J. P. Lees *et al.* (BaBar Collaboration), Phys. Rev. D **87**, 012004 (2013); R. Aaij *et al.* (LHCb Collaboration), Phys. Rev. Lett. **112**, 041801 (2014).
- [5] T. A. Aaltonen *et al.* [CDF Collaboration], arXiv:1410.5435 [hep-ex].