

Spin-structures and proper group actions

Peter Hochs* and Varghese Mathai†

October 12, 2018

Abstract

We generalise Atiyah and Hirzebruch's vanishing theorem for actions by compact groups on compact Spin-manifolds to possibly non-compact groups acting properly and cocompactly on possibly non-compact Spin-manifolds. As corollaries, we obtain some vanishing results for an \hat{A} -type genus.

Contents

1	Results and applications	2
2	Spin-structures on slices	5
3	Proofs of the results	7

Introduction

In 1970, Atiyah and Hirzebruch [2] proved the following remarkable result.

Theorem 1. *Let N be a compact, connected even dimensional manifold and K be a compact connected Lie group acting smoothly and non-trivially on N . Suppose also that N has a K -invariant Spin structure. Then the equivariant index of the Dirac operator on N vanishes in the representation ring of K ,*

$$\text{index}_K(\not{D}_N) = 0 \quad \in R(K). \quad (0.1)$$

In particular, the \hat{A} -genus of N is zero.

*University of Adelaide, peter.hochs@adelaide.edu.au

†University of Adelaide, mathai.varghese@adelaide.edu.au

Their result then inspired many, especially Witten [13] who studied two-dimensional quantum field theories and the index of the Dirac operator on free loop space LN , relating it to the rigidity of certain Dirac-type operators on N and the elliptic genus, which was proved in [4, 11].

Our goal in this note is to extend Theorem 1 to the non-compact setting. The result is Theorem 2, which can be stated in an equivalent way as Theorem 3. One consequence is a result related to rigidity, Corollary 4. These results involve K-theory and K-homology of C^* -algebras, but they have purely differential geometric consequences, as noted in Corollary 5.

Acknowledgements

The first author was supported by a Marie Curie fellowship from the European Union. The second author thanks the Australian Research Council for support via the ARC Discovery Project grant DP130103924.

1 Results and applications

Let M be a manifold, on which a connected Lie group G acts properly and isometrically. Suppose that the action is cocompact, i.e. M/G is compact, and that M has a G -equivariant Spin-structure. Let

$$\text{index}_G(\not{D}_M) \in K_\bullet(C_r^*G)$$

be the equivariant index of the associated Spin-Dirac operator. Here $K_\bullet(C_r^*G)$ is the K-theory of the reduced group C^* -algebra of G , and index_G denotes the analytic assembly map used in the Baum–Connes conjecture [3, 9]. If G is compact, then $K_0(C_r^*G) = R(G)$ and $K_1(C_r^*G) = \{0\}$, and the analytic assembly map is the usual equivariant index.

Let $K < G$ be a maximal compact subgroup, and suppose G/K has a G -equivariant Spin-structure. This is true for a double cover of G , as pointed out in Section 2.

The action by G on M will be called *properly trivial* if all stabilisers are maximal compact subgroups of G . For a proper action, the stabilisers cannot be larger. The action is called *properly nontrivial* if it is not properly trivial.

For any manifold X , we write \hat{A}_X for the \hat{A} -class of X . If X is compact, then we denote its \hat{A} -genus by

$$\hat{A}(X) := \int_X \hat{A}_X.$$

Atiyah and Hirzebruch's Theorem 1 generalises as follows.

Theorem 2. *As above, let G be a connected Lie group, with maximal compact subgroup K , such that G/K has a G -equivariant Spin-structure. Suppose that G acts properly and cocompactly on a manifold M , and that M has a G -equivariant Spin-structure. If the action is properly nontrivial, then*

$$\text{index}_G(\not\partial_M) = 0.$$

This result can be restated in an equivalent form as follows.

Theorem 3. *Consider the setting of Theorem 2. One has $\text{index}_G(\not\partial_M) \neq 0$ if and only if there is a compact Spin-manifold N with $\hat{A}(N) \neq 0$, and a G -equivariant diffeomorphism*

$$M \cong G/K \times N,$$

where G acts trivially on N .

A first consequence of Theorem 2 is a rigidity-type result. This involves the Dirac induction map

$$\text{D-Ind}_K^G : R(K) \rightarrow K_\bullet(C_r^*G). \quad (1.1)$$

Here $R(K)$ is the representation ring of K . This map is an isomorphism of Abelian groups by the Connes–Kasparov conjecture, which was proved for (almost) connected groups in [5].

Corollary 4. *In the setting of Theorem 2, one has*

$$\text{index}_G(\not\partial_M) \in \mathbb{Z} \cdot \text{D-Ind}_K^G[\mathbb{C}],$$

where $[\mathbb{C}] \in R(K)$ is the class of the trivial representation.

Theorems 2 and 3 and Corollary 4 have consequences within differential geometry, not involving K -theory and K -homology. Let $c \in C_c^\infty(M)$ be a cutoff function, that is to say a non-negative function satisfying

$$\int_G c(g^{-1}m)dg = 1$$

for all $m \in M$, for a fixed left Haar measure dg on G . The averaged \hat{A} -genus of the action by G on M is

$$\hat{A}_c(M) := \int_M c \hat{A}_M.$$

It was shown to be independent of c in [12].

Corollary 5. *Consider the setting of Theorem 2.*

1. *If the action is properly nontrivial, then $\hat{A}_c(M) = 0$.*
2. *If $\hat{A}_c(M) \neq 0$, then there is a compact Spin-manifold N with $\hat{A}(N) \neq 0$, and a G -equivariant diffeomorphism*

$$M \cong G/K \times N,$$

where G acts trivially on N .

3. *The averaged \hat{A} -genus $\hat{A}_c(M)$ is an integer multiple of $\hat{A}_c(G/K)$.*

Remark 6. There are many group actions that satisfy the hypotheses of Theorem 2. Indeed, if K acts on a compact Spin-manifold N as in Theorem 1, then Theorem 2 applies to the action by G on the fibred product $G \times_K N$, as we will see. If $K = S^1$, then it is proved in the theorem in Section 2.3 in [2] that any compact oriented manifold X with $\hat{A}(X) = 0$ has the property that mX (for some $m \in \mathbb{N}$) is oriented cobordant to a compact Spin manifold N which has a non-trivial S^1 -action on each of its components. Then the action by K on N satisfies the hypotheses of Theorem 1, so that the action by G on $G \times_K N$ satisfies the conditions of Theorem 2.

Remark 7. In the setting of Theorems 2 and 3, there is a proper, equivariant map $p : M \rightarrow G/K$ (see Theorem 8 below). The map p_* induced on K -homology relates the equivariant indices on M and G/K by the diagram

$$\begin{array}{ccc} K_\bullet^G(M) & \xrightarrow{\text{index}_G} & K_\bullet(C_r^*G). \\ p_* \downarrow & \nearrow \cong & \\ K_\bullet^G(G/K) & \xrightarrow{\text{index}_G} & \end{array}$$

Since the Baum–Connes conjecture is true for connected groups by Theorem 1.1 in [5], the equivariant index on G/K defines an isomorphism

$$K_\bullet^G(G/K) \cong K_\bullet(C_r^*G).$$

Hence $p_*[\partial_M] = 0$ if and only if $\text{index}_G(\partial_M) = 0$, so the (non)vanishing of $\text{index}_G(\partial_M) = 0$ in Theorems 2 and 3 can be replaced by the (non)vanishing of $p_*[\partial_M] = 0$. This removes K -theory and the assembly map from these results.

2 Spin-structures on slices

We begin by recalling the smooth version of Abels' slice theorem for proper group actions. Let M be a smooth manifold, and let G be a connected Lie group acting properly on M . Let $K < G$ be maximal compact.

Theorem 8 (p. 2 of [1]). *There is a smooth, K -invariant submanifold $N \subset M$, such that the map $[g, n] \mapsto gn$ is a G -equivariant diffeomorphism*

$$G \times_K N \cong M \quad (2.1)$$

Here the left hand side is the quotient of $G \times N$ by the action by K given by

$$k \cdot (g, n) = (gk^{-1}, kn),$$

for $k \in K$, $g \in G$ and $n \in N$.

We call (2.1) an associated Abels fibration of M , as it is a fibre bundle over G/K with fibre N . From now on, fix N as in Theorem 8.

The fixed point set N^K of the action by K on N is related to the action by G on M in the following way.

Lemma 9. *One has*

$$M_{(K)} = G \cdot N^K \cong G/K \times N^K,$$

where $M_{(K)}$ is the set of points in M with stabilisers conjugate to K .

Proof. Let $m \in M_{(K)}$, and write $m = [g, n]$ for $g \in G$ and $n \in N$, under the correspondence (2.1). Then $G_m = gK_n g^{-1}$. So G_m is conjugate to K if and only if K_n is. Since $K_n < K$, it is conjugate to K precisely if it equals K . \square

Now fix a K -invariant inner product on the Lie algebra \mathfrak{g} of G , and let $\mathfrak{p} \subset \mathfrak{g}$ be the orthogonal complement to the Lie algebra \mathfrak{k} of K . Suppose $\text{Ad} : K \rightarrow \text{SO}(\mathfrak{p})$ lifts to

$$\widetilde{\text{Ad}} : K \rightarrow \text{Spin}(\mathfrak{p}). \quad (2.2)$$

This is always possible if one replaces G by a double cover. Indeed, consider the diagram

$$\begin{array}{ccc} \widetilde{K} & \xrightarrow{\widetilde{\text{Ad}}} & \text{Spin}(\mathfrak{p}) \\ \pi_K \downarrow & & \downarrow \pi \\ K & \xrightarrow{\text{Ad}} & \text{SO}(\mathfrak{p}), \end{array}$$

where

$$\begin{aligned}\tilde{K} &:= \{(k, a) \in K \times \text{Spin}(\mathfrak{p}); \text{Ad}(k) = \pi(a)\}; \\ \pi_K(k, a) &:= k; \\ \widetilde{\text{Ad}}(k, a) &:= a,\end{aligned}$$

for $k \in K$ and $a \in \text{Spin}(\mathfrak{p})$. Then for all $k \in K$,

$$\pi_K^{-1}(k) \cong \pi^{-1}(\text{Ad}(k)) \cong \mathbb{Z}_2,$$

so π_K is a double covering map. Since G/K is contractible, \tilde{K} is the maximal compact subgroup of a double cover of G .

Suppose M has a G -equivariant Spin-structure $P_M \rightarrow M$. In Section 3.2 of [6] and Section 3.2 of [8], an induction procedure of equivariant Spin^c -structures from N to M is described, which we will denote by Ind_N^M here. We will use the fact that any G -equivariant Spin-structure on M can be obtained via this induction procedure. (See also Proposition 3.10 in [8].)

Lemma 10. *There is a K -equivariant Spin-structure $P_N \rightarrow N$ such that*

$$P_M = \text{Ind}_N^M(P_N).$$

Proof. Let $\mathfrak{p}_N \rightarrow N$ be the trivial vector bundle $N \times \mathfrak{p} \rightarrow N$, with the diagonal K -action. It has the K -equivariant Spin-structure

$$N \times \text{Spin}(\mathfrak{p}) \rightarrow N$$

on \mathfrak{p}_N , where K acts diagonally on $N \times \text{Spin}(\mathfrak{p})$ via the lift (2.2) of the adjoint action by K on \mathfrak{p} . Since

$$TM = G \times_K (TN \oplus \mathfrak{p}_N)$$

(see Proposition 2.1 and Lemma 2.2 in [6]), the restriction $P_M|_N$ is a K -equivariant Spin-structure on

$$TM|_N = TN \times \mathfrak{p}_N.$$

Together with the Spin^c -structure on \mathfrak{p}_N , this determines a K -equivariant Spin-structure on TN by the two-out-of-three lemma. Explicitly, if \mathcal{S}_M is the spinor bundle associated to P_M , and $\Delta_{\mathfrak{p}}$ is the standard representation of $\text{Spin}(\mathfrak{p})$, then the spinor bundle \mathcal{S}_N associated to P_N is determined by

$$\mathcal{S}_M|_N = \mathcal{S}_N \otimes \Delta_{\mathfrak{p}}.$$

The Clifford action $c_{\mathcal{S}_N}$ by TN on \mathcal{S}_N is determined by the property that for all $n \in N$, $v \in T_n N$ and $X \in \mathfrak{p}$,

$$c_{\mathcal{S}_M|_N}(v, X) = \begin{cases} c_{\mathcal{S}_N}(v) \otimes 1_{\Delta_{\mathfrak{p}}} + \varepsilon_{\mathcal{S}_N} \otimes c_{\mathfrak{p}}(X) & \text{if } \dim G/K \text{ is even;} \\ c_{\mathcal{S}_N}(v) \otimes 1_{\Delta_{\mathfrak{p}}} + \varepsilon_{\mathcal{S}_N} \otimes \varepsilon_{\Delta_{\mathfrak{p}}} c_{\mathfrak{p}}(X) & \text{if } \dim G/K \text{ is odd.} \end{cases}$$

Here $\varepsilon_{\mathcal{S}_N}$ and $\varepsilon_{\Delta_{\mathfrak{p}}}$ are the grading operators on \mathcal{S}_N and $\Delta_{\mathfrak{p}}$, respectively. See also Section 3.1 in [10].

In Lemma 3.9 of [8], it is shown that

$$P_M = \text{Ind}_N^M(P_N).$$

□

Remark 11. We have only considered the principle bundle part $P_M \rightarrow M$ of a Spin-structure on M , not the isomorphism

$$P_M \times_{\text{Spin}(\dim M)} \mathbb{R}^{\dim M} \cong TM.$$

This isomorphism determines the Riemannian metric on M induced by the Spin-structure. The index of \not{D}_M is independent of this metric, however, so Lemma 10 is enough for our purposes.

3 Proofs of the results

The *quantisation commutes with induction* techniques of [6, 7], adapted to the Spin-setting, allow us to deduce Theorems 2 and 3 from Atiyah and Hirzebruch's Theorem 1. This is based on the fact that the Dirac induction map (1.1) relates the equivariant indices of the Spin-Dirac operators \not{D}_N on N and \not{D}_M on M , associated to the Spin-structures P_N and P_M , respectively, to each other. (See also Theorem 5.7 in [8].)

Proposition 12. *One has*

$$\text{D-Ind}_K^G(\text{index}_K(\not{D}_N)) = \text{index}_G(\not{D}_M) \in K_{\bullet}(C_r^*G).$$

Proof. Let $K_{\bullet}^K(N)$ and $K_{\bullet}^G(M)$ be the equivariant K-homology groups [3] of N and M , respectively. In Theorem 4.6 in [6] and Theorem 4.5 in [7], a map

$$K\text{-Ind}_K^G : K_{\bullet}^K(N) \rightarrow K_{\bullet}^G(M)$$

is constructed, such that the following diagram commutes:

$$\begin{array}{ccc}
K_{\bullet}^G(M) & \xrightarrow{\text{index}_G} & K_{\bullet}(C_r^*G) \\
\uparrow K\text{-Ind}_K^G & & \uparrow D\text{-Ind}_K^G \\
K_{\bullet}^K(N) & \xrightarrow{\text{index}_K} & R(K).
\end{array}$$

In Section 6 of [6], it is shown that the K-homology class of a Spin^c -Dirac operator on N , associated to a connection ∇^N on the determinant line bundle of a Spin^c -structure, is mapped to the class of a Spin^c -Dirac operator on M associated to a connection ∇^M induced by ∇^N on the determinant line bundle of the induced Spin^c -structure, by the map $K\text{-Ind}_K^G$. In the Spin -setting, both connections ∇^N and ∇^M are trivial connections on trivial line bundles. Hence one gets

$$K\text{-Ind}_K^G[\partial_N] = [\partial_M],$$

and the result follows. \square

Proof of Theorems 2 and 3. Consider the setting of Theorem 2, let $N \subset M$ be as in Theorem 8. Consider a K -equivariant Spin -structure on N as in Lemma 10. By Proposition 12, we have

$$\text{index}_G(\partial_M) = D\text{-Ind}_K^G(\text{index}_K(\partial_N)). \quad (3.1)$$

The stabiliser of a point $m \in M$ is a maximal compact subgroup of G if and only if $m \in M_{(K)}$. Hence, by Lemma 9, the condition on the stabilisers of the action by G on M is equivalent to the action by K on N being nontrivial. So Theorem 1 implies that

$$\text{index}_K(\partial_N) = 0,$$

and Theorem 2 follows.

To prove Theorem 3, we note that by injectivity of Dirac induction, the equality (3.1) and Theorem 1 imply the equivalences

$$\begin{aligned}
\text{index}_G(\partial_M) \neq 0 & \Leftrightarrow \text{index}_K(\partial_N) \neq 0 \\
& \Leftrightarrow K \text{ acts trivially on } N \text{ and } \hat{A}(N) \neq 0 \\
& \Leftrightarrow M \cong G/K \times N \text{ and } \hat{A}(N) \neq 0.
\end{aligned}$$

\square

Remark 13. In the proof of Theorem 1 in [2], the fixed point set of the action plays an important role. Because proper actions by noncompact groups do not have fixed points, the authors think it is unlikely that the arguments in [2] generalise directly to a proof of Theorems 2 and 3. This is one reason to apply the induction argument used here.

It remains to prove Corollaries 4 and 5.

Proof of Corollary 4. If $\text{index}_G(\partial_M) = 0$, the result holds trivially. If $\text{index}_G(\partial_M)$ is nonzero, then we have the decomposition $M \cong G/K \times N$ of Theorem 3. By multiplicativity of the analytic assembly map (see Theorem 5.2 in [6]), we then conclude that

$$\text{index}_G(\partial_M) = \text{index}(\partial_N) \text{index}_G(\partial_{G/K}) = \text{index}(\partial_N) \text{D-Ind}_K^G[\mathbb{C}]. \quad (3.2)$$

□

Proof of Corollary 5. Let $\tau : C_r^*G \rightarrow \mathbb{C}$ be the von Neumann trace, determined by

$$\tau(R(f)^*R(f)) = \int_G |f(g)|^2 dg,$$

for $f \in L^1(G) \cap L^2(G)$, where R denotes the right regular representation. This induces a morphism $\tau_* : K_\bullet(C_r^*G) \rightarrow \mathbb{R}$. Wang showed in Theorem 6.12 in [12] that

$$\hat{A}_c(M) = \tau_*(\text{index}(\partial_M)).$$

So $\text{index}(\partial_M) = 0$ implies $\hat{A}_c(M) = 0$. Furthermore,

$$\tau_*(\text{D-Ind}_K^G[\mathbb{C}]) = \hat{A}_c(G/K).$$

Hence Theorem 2, Theorem 3 and Corollary 4 imply Corollary 5. □

References

- [1] H. Abels, *Parallelizability of proper actions, global K-slices and maximal compact subgroups*, Math. Ann. 212 (1974), 1–19.
- [2] M. Atiyah, F. Hirzebruch, *Spin-manifolds and group actions*, 1970 Essays on Topology and Related Topics (Mémoires dédiés Georges de Rham) Springer, New York, 18–28.

- [3] P. Baum, A. Connes, N. Higson, *Classifying space for proper actions and K-theory of group C*-algebras*, C*-algebras: 1943-1993 (San Antonio, TX, 1993), Contemp. Math., 167, Amer. Math. Soc., Providence, RI (1994), 240–291.
- [4] R. Bott and C. Taubes, *On the rigidity theorem of Witten*, J. Amer. Math. Soc., 2 (1989), 137-186.
- [5] J. Chabert, S. Echterhoff and R. Nest, *The Connes–Kasparov conjecture for almost connected groups and for linear p-adic groups*, Publ. Math. Inst. Hautes Études Sci. 97 (2003), 239–278.
- [6] P. Hochs, *Quantisation commutes with reduction at discrete series representations of semisimple groups*, Adv. Math. 222 (2009), no. 3, 862–919.
- [7] P. Hochs, *Quantisation of presymplectic manifolds, K-theory and group representations*, Proc. Amer. Math. Soc. (to appear), 17 pages, arXiv:1211.0107.
- [8] P. Hochs and V. Mathai, *Quantising proper actions on Spin^c -manifolds*, 62 pages, arXiv:1408.0085.
- [9] G. Kasparov, *Equivariant KK-theory and the Novikov conjecture*. Invent. Math. **91** (1988), no. 1, 147-201.
- [10] R.J. Plymen, *Strong Morita equivalence, spinors and symplectic spinors*, J. Operator Theory 16 (1986), no. 2, 305–324.
- [11] C. Taubes, *S^1 actions and elliptic genera*, Comm. Math. Phys. 122 (1989), 455-526.
- [12] H. Wang, *L^2 -index formula for proper cocompact group actions*. J. Noncommut. Geom. 8 (2014), no.2, 393–432.
- [13] E. Witten, *Elliptic genera and quantum field theory*, Comm. Math. Phys. 109 (1987), no. 4, 525–536.