

The history of the universe is an elliptic curve

Robert Coquereaux

Centre de Physique Théorique (CPT),

Aix Marseille Université, Université de Toulon, CNRS, CPT, UMR 7332, 13288 Marseille, France

Abstract

Friedmann-Lemaître equations with contributions coming from matter, curvature, cosmological constant, and radiation, when written in terms of conformal time u rather than in terms of cosmic time t , can be solved explicitly in terms of standard Weierstrass elliptic functions. The spatial scale factor, the temperature, the densities, the Hubble function, and almost all quantities of cosmological interest (with the exception of t itself) are elliptic functions of u , in particular they are bi-periodic with respect to a lattice of the complex plane, when one takes u complex. After recalling the basics of the theory, we use these explicit expressions, as well as the experimental constraints on the present values of density parameters (we choose for the curvature density a small value in agreement with experimental bounds) to display the evolution of the main cosmological quantities for one real period $2\omega_r$ of conformal time (the cosmic time t “never ends” but it goes to infinity for a finite value $u_f < 2\omega_r$ of u). A given history of the universe, specified by the measured values of present-day densities, is associated with a lattice in the complex plane, or with an elliptic curve, and therefore with two Weierstrass invariants g_2, g_3 . Using the same experimental data we calculate the values of these invariants, as well as the associated modular parameter and the corresponding Klein j -invariant. If one takes the flat case $k = 0$, the lattice is only defined up to homotheties, and if one, moreover, neglects the radiation contribution, the j -invariant vanishes and the corresponding modular parameter τ can be chosen in one corner of the standard fundamental domain of the modular group (equianharmonic case: $\tau = \exp(2i\pi/3)$). Several exact – i.e., non-numerical – results of independent interest are obtained in that case.

1 Introduction

Friedmann-Lemaître equation is studied in many places (articles, books, encyclopedias, etc.). The subject is hardly new and it is known that the general solution to this differential equation involves elliptic integrals [8], but few people tried to obtain explicit expressions. To our knowledge, the first article where such an analysis is performed in a detailed manner, keeping all the contributions coming from matter, curvature, cosmological constant, and radiation, is [2]. Explicit formulae for all range of values of the reduced cosmological constant, and assuming¹ $k = +1$ (the so-called “closed case”), are given in the same reference. This work was quickly followed by [5], where the same analysis was performed for the flat and open cases. Fifteen years later or so (see [3] and the lectures [4]), the same analysis was reconsidered, in the light of experiments showing that the cosmological constant was probably not zero, after all. . . A little bit more than thirty years after the first paper we find it useful to return to this problem since the range of cosmological parameters specifying our universe history has been made more precise thanks to recent experiments, and since the elliptic functions (essentially Weierstrass elliptic functions) that allow one to give explicit expressions for the quantities of interest have been made available, with a very good precision, in most computer packages. About this last point it seems that many people still prefer to perform simplifying assumptions or use techniques of numerical integration to describe the evolution of quantities of cosmological interest in terms of cosmic time. We think that using exact expressions is highly preferable, not only for conceptual reasons, but also for practical reasons: some features of the solutions (presence of local extrema, inflection points, behavior near the singularities, etc.) are almost obvious if one uses exact expressions, because of the well-known properties of the involved functions, but may be sometimes difficult to detect numerically. If one is interested in the evolution of the solutions in terms of conformal time, i.e., “before” the Big Bang, or “after” the end of cosmic time (infinity), the use of such expressions is, of course, required.

As it is clear from the very definition of elliptic functions, cosmological solutions of Friedmann equations are periodic in conformal time, they are even bi-periodic if the latter is allowed to take complex values. This is by no means in contradiction with the fact that many solutions (in particular the one that seems to be dictated by experiments) describe a never-ending universe starting with a Big Bang, since the “never-ending” qualifier refers to the cosmic time variable, not to the conformal time variable. Rather than trying to study the evolution of the spatial scale factor $a(t)$ as a function of cosmic time t , the starting point of the method of resolution is to notice that it is much better to give a parametric representation $(a(u), t(u))$ of $a(t)$, because $a(u)$, like most functions of cosmological interest (not the function $t(u)$), turns out to be an elliptic function of u , the conformal time. Actually, the resolution of Friedmann equations for the quantity $1/a$, the inverse of the scale factor, is almost immediate; this quantity is essentially equal to the temperature \tilde{T} of the Cosmic Microwave Blackbody (CMB) radiation (warning: in our paper the variable T will denote another quantity that differs from \tilde{T} by a scale factor).

The present article is supposed to be self-contained and does not require from the reader any familiarity with Friedmann equations. We nevertheless still refer to the article [2] for a general study of their analytic solutions, for all possible values of parameters, but here we shall restrict our attention to those solutions that are compatible with the recent experiments. In particular the

¹This assumption was then made by a majority of people, but at the same time the cosmological constant was not fashioned and was almost always assumed to be zero in the cosmology community.

evolution of density parameters and of other quantities of cosmological interest, that are displayed in sec. 4.2 as functions of conformal time, are determined by using the experimental results for the present-day values of density parameters and Hubble constant. Section 2 is certainly standard, but it serves the purpose of specifying our notations. At the beginning of section 3, we remind the reader that the main features of the evolution of the temperature, as a function of conformal time, can be simply obtained by studying the classical motion of point in a potential. Then we give explicit analytical solutions, with or without radiation (the latter case is of course simpler but hides interesting physical phenomena that show up in a neighborhood of $u = 0$). In the same section we define the Weierstrass invariants of the universe. In section 4 we use the values of present-day densities coming from experiment (in particular from the Planck collaboration) to calculate and display the behavior of most quantities of cosmological interest in terms of conformal time. This is done by assuming a small (positive) value for the present value of the curvature density, compatible with the experimental bounds. This section ends with a table giving the values of various quantities at several important dates in the (conformal) history of the universe – the fact, for instance, that the curvature density function has an extremum located in our past is a phenomenon that is often overlooked. Special features of the case $k = 0$ (flat case) are discussed in section 5. In that particular situation, the experimental data only allows one to determine up to scale the lattice in the complex plane associated with the universe history. In other words, the corresponding elliptic curve is only known up to isomorphism, but several exact results (values of periods, value of conformal time when $t \rightarrow \infty$, etc.) can nevertheless also be obtained in that case. Section 6 contains miscellaneous comments. Several properties of elliptic functions in relation with lattices, tori, elliptic curves, and modular considerations, are given in the appendix.

As it was already mentioned, the general method of resolution of Friedmann equation in terms of elliptic functions was discussed in [2], but a good part of the discussion relating cosmology to modular considerations is only described in the present paper. In this respect, let us summarize some of the properties that will be discussed (see also the abstract): to each universe history determined by the measurement of cosmological parameters one can associate an elliptic curve, or, equivalently, a lattice in the complex plane, or a torus with a complex structure. If $k = \pm 1$, the measurement of all the densities (summing up to 1) specifies in particular the Weierstrass invariants g_2 and g_3 . The period parallelogram can be chosen as a rhombus, symmetric with respect to the real axis. Conversely, the history of the universe, described by the evolution of the temperature as a function of conformal time is fully specified by the Weierstrass invariants, together with today's date (the value of the conformal time “now”), and a dimensionful quantity defining the centimeter, for instance the Hubble constant. If $k = 0$, the lattice is only determined up to a complex homothety and the elliptic curve only up to isomorphism (the invariants g_2, g_3 are obtained up to scale). If one furthermore neglects the effect of radiation, the curve is equianharmonic, which corresponds to a modular parameter equal to $exp(2i\pi/3)$ sitting in the corner of the boundary of the standard fundamental domain for the modular group. In all cases one can determine the value of the Klein j -invariant.

2 Friedmann-Lemaître equations and elliptic functions

2.1 Natural variables for the Friedmann-Lemaître equations

It is usual to model space-time as a four-dimensional smooth manifold endowed with a pseudo-riemannian structure specified by a metric. Assuming that our space-time neighborhood is, in first approximation, homogeneous and isotropic, this metric is described, in local coordinates (t, χ, θ, ϕ) , i.e., in the domain of a chart, by the line element ds^2 . There are three possibilities ($k = \pm 1$ or 0):

$$ds^2 = -dt^2 + a(t)^2 [d\chi^2 + s(\chi)^2 (d\theta^2 + \sin^2 \theta d\phi^2)]$$

with $s(\chi) = \sin(\chi)$ if $k = +1$, $s(\chi) = \sinh(\chi)$ if $k = -1$, and $s(\chi) = \chi$ if $k = 0$.

In conventional cosmology one assumes that the average local energy described by a rank-two tensor T , the energy-momentum tensor, splits into three parts (G denotes the Newton constant):

- Vacuum contribution $\rho_{vac} = \frac{\Lambda}{8\pi G}$ where Λ is the cosmological constant,
- Radiation contribution ρ_{rad} such that $a^4(t)\rho_{rad}(t) = const. = \frac{3}{8\pi G}C_r$,
- Averaged matter contribution ρ_{mat} such that $a^3(t)\rho_{mat}(t) = const. = \frac{3}{8\pi G}C_m$,

Einstein's equations read $E_{\mu\nu} = 8\pi G T_{\mu\nu}$ where E is the Einstein tensor. The cosmological term described by Λ is sometimes written explicitly on the lhs of those equations but here it is included as a part of T itself, on the rhs. The quantities Λ , C_r and C_m are constant, as well as ρ_{vac} , but ρ_{rad} and ρ_{mat} are, a priori, time-dependent quantities. Assuming a Levi-Civita connection (no torsion), the Einstein tensor is determined from the metric alone, and Einstein's equations imply that the evolution of $a(t)$ is governed by the Friedmann equation:

$$\frac{1}{a^2} \left(\frac{da}{dt} \right)^2 = \frac{C_r}{a^4} + \frac{C_m}{a^3} - \frac{k}{a^2} + \frac{\Lambda}{3}. \quad (1)$$

Lemaître [8] did, long ago, an analytic study of the solutions of Friedmann equations, with a cosmological constant; his discussion, made in terms of a , the scale factor, and t , the cosmic time, involves elliptic integrals. In order to discuss these equations it is however convenient [2] to use another set of variables, i.e., to introduce a *conformal time* u , and a dimensionless *reduced temperature* $T(u)$, defined as:

$$du = \frac{dt}{a} \quad \text{and} \quad T(u) = \frac{1}{\Lambda_c^{1/2} a(u)} \quad \text{with} \quad \sqrt{\Lambda_c} = \frac{2}{3C_m} \quad (2)$$

The variable u is natural, both geometrically (it gives three-dimensional geodesic distances) and analytically, as we shall see below, since it is only when we express the cosmological quantities of interest in terms of $T(u)$ that these quantities can be written themselves as elliptic functions with respect to a particular lattice.

Intuitively, this change of variables replaces the scale factor a , that measures the “size” of the spatial universe, by its inverse, a quantity proportional to the temperature of the cosmic microwave

radiation (see later). In a Big Bang cosmology, T is infinite at the Big Bang and decreases as the universe expands. This change of variables replaces the cosmic time t by a parameter u , the conformal time, that measures geodesic distances in dimensionless units. The previous equations define u only up to an additive constant; in a Big Bang cosmology, it is natural to set $u = 0$ at the Big Bang, so that u gives the dimensionless distance along the trajectory of a photon that would have been emitted at the Big Bang. This is a perfectly natural way of measuring “time”. Rather than describing the dynamics of the universe by a single function $a(t)$ of one variable, we therefore use the parametric equations $(T(u), t(u))$. The advantage is that the differential equation for $T(u)$ is very simple – see below – and can be integrated immediately in terms of standard elliptic functions. Very often – and in particular, as we shall see later, for the cosmological solutions of physical interest – the function $t(u)$ approaches a logarithmic singularity when u approaches a finite limit u_f . In other words, when the cosmic time goes to infinity, the universe expands for ever (in terms of t) and cools down ($T \sim 1/a \rightarrow 0$) but the conformal time goes to u_f . So, even if the universe is spatially closed (let us say that it is S^3), an observer will never see the back of his head if the cosmology is such that u_f is finite and smaller than 2π , even if this observer waits for an infinite (cosmic) time.

In this article we use the natural system of units for which $\hbar = c = 1$. All quantities are therefore homogenous with L^p , for some integer p , where L is a length (the cm, say). In particular the quantities C_r , C_m and Λ have dimensions L^2, L, L^{-2} respectively, a has dimension L and t has dimension L^{-1} . The Newton constant G is homogeneous with L^2 . Notice that T and u are *dimensionless*.

With these new variables, Friedmann equation becomes

$$(dT/du)^2 = \alpha T^4 + \frac{2}{3}T^3 - kT^2 + \frac{\lambda}{3} \quad (3)$$

where

$$\lambda = \frac{\Lambda}{\Lambda_c} \quad \text{and} \quad \alpha = C_r \Lambda_c \quad (4)$$

are two constant dimensionless parameters: the *reduced cosmological constant* and the *reduced radiation parameter*. We shall see later how these parameters can be extracted from the more standard densities used to describe experimental results.

The next section will deal with analytic solutions, but the direct link with the theory of elliptic functions should be already clear from the fact that, with our parametrization, the RHS of eq 3 is a polynomial of degree four.

2.2 Hubble function and density “parameters”

As a rule we shall add an upper or lower index o to denote the present-day value of the cosmological quantities. For instance t_o is the age of the universe (using cosmic time), u_o its age in terms of conformal time, a_o is the present value of the scale factor, etc.

The *Hubble function* describing the rate of expansion is defined, as usual, by $H = \frac{d \text{Log}(a)}{dt}$ and can be written, in terms of the reduced temperature $T(u)$ as

$$H(u) = -\Lambda_c^{1/2} \frac{dT}{du} \quad (5)$$

In natural units, most quantities used in cosmology are dimensionless, but the Hubble function is a dimensionful quantity homogenous to an inverse length, so we may consider that the present value of $H_o = H(u_0)$, or rather of $(H_o)^{-1}$, defines what the centimeter is (today).

Notice that

$$H^2 = \alpha \Lambda_c T^4 + 2/3 \Lambda_c T^3 - k \Lambda_c T^2 + \Lambda/3 \quad (6)$$

Multiplying this equation by $1/H^2$, one obtains the famous relation² :

$$1 = \Omega_r + \Omega_m + \Omega_K + \Omega_\Lambda \quad (7)$$

with

$$\Omega_K = -kT^2 \frac{\Lambda_c}{H^2}, \quad \Omega_m = \frac{2}{3} T^3 \frac{\Lambda_c}{H^2}, \quad \Omega_\Lambda = \frac{\lambda}{3} \frac{\Lambda_c}{H^2}, \quad \Omega_r = \alpha T^4 \frac{\Lambda_c}{H^2} \quad (8)$$

Equivalently,

$$\Omega_K = -\frac{k}{a^2 H^2}, \quad \Omega_m = \frac{C_m}{a^3 H^2}, \quad \Omega_\Lambda = \frac{\Lambda}{3H^2}, \quad \Omega_r = \frac{C_r}{a^4 H^2} \quad (9)$$

We remind the reader that those quantities³, although often called “density parameters”, are function of time (u or t). For this reason it is usual to introduce the notations Ω_m^o , Ω_r^o , Ω_K^o and Ω_Λ^o to denote their present-day values. Obviously, Ω_m and Ω_r are positive but Ω_Λ and Ω_K could be of both signs. It is quite common to call $\Omega = \Omega_m + \Omega_r + \Omega_\Lambda$ the “total density parameter”. Since $-\Omega_K = \Omega - 1$ and since, for historical reasons, Ω_K is negative when $k = +1$ and vice-versa, one sees that $k = +1, 0$ or -1 , respectively, if $\Omega > 1, = 1$, or < 1 .

The *constant* parameters Λ , k , C_m and C_r entering the original Friedmann equation are expressed as follows in terms of the *time dependent* density parameters:

$$\Lambda = 3H^2 \Omega_\Lambda, \quad k = -\text{sign}(\Omega_K), \quad C_m = \frac{\Omega_m}{|\Omega_K|^{3/2} H}, \quad C_r = \frac{\Omega_r}{|\Omega_K|^2 H^2} \quad (10)$$

In the closed case ($k = +1$), rather than C_m or Λ_c , one often uses the mass M

$$M = 2\pi^2 a^3 \rho_m = \frac{\pi}{2G\sqrt{\Lambda_c}} = \frac{3\pi}{4G} C_m \quad (11)$$

² Ω_r^o being non zero, but very small, it is sometimes dropped from equation 7, and because Ω_K^o is experimentally compatible with 0, it is also often dropped from many presentations. It is interesting to remember that thirty years ago, it was another parameter (namely Ω_Λ) that was often forgotten from many presentations.

³ The densities $\Omega_r(u)$, $\Omega_m(u)$, $\Omega_\Lambda(u)$ were respectively called $\alpha_S(\tau)$, $\Omega(\tau)$ and $\lambda_S(\tau)$ in ref. [2].

to parametrize the matter contents of the universe.

If $k = \pm 1$, the *constant* parameters α and λ , introduced in equation (4) can be expressed as follows in terms of the *time dependent* density parameters:

$$\alpha = \frac{4}{9} \frac{\Omega_r |\Omega_K|}{\Omega_m^2} \quad \lambda = \frac{27}{4} \frac{\Omega_\Lambda \Omega_m^2}{|\Omega_K^3|} \quad (12)$$

If $k = 0$, the measurements (now) of Ω_r , Ω_m and Ω_Λ give no information on the individual values of α and λ , but they are nevertheless related — at all times — by the relation:

$$\alpha^3 \lambda = \frac{16}{27} \frac{\Omega_r^3 \Omega_\Lambda}{\Omega_m^4} \quad (13)$$

2.3 Temperature, units and dimensions

The reason for calling T a dimensionless “reduced temperature” is that it is proportional to the temperature \tilde{T} of the black body radiation. Indeed, $\rho_{rad} = 4\sigma\tilde{T}^4$ where σ is the Stefan-Boltzmann constant. Since $\rho_{rad} = \frac{3}{8\pi G} \alpha \Lambda_c T^4$, one finds

$$\tilde{T}^4 = \frac{3}{8\pi G} \frac{\alpha \Lambda_c}{4\sigma} T^4 \quad (14)$$

With $\hbar = c = 1$, the value of the Stefan-Boltzmann constant is $\sigma = \pi^2 k_B^4 / 60 = 59.8 \text{ cm}^{-4}$, where k_B is the Boltzmann constant. Notice that \tilde{T} is in degrees Kelvin, hence dimensionless, but the Boltzmann constant k_B (as $k_B \tilde{T}$) is an energy, hence homogenous to an inverse length, as it should. We call T_o and \tilde{T}_o the *present* values of the time-dependent quantities T and \tilde{T} .

As already recalled, in this system of units all quantities are either dimensionless or have a dimension which is some power of a length ($[cm]$). We gather the relevant information as follows:

$$t \sim a \sim C_m \sim [cm], \quad G \sim C_r \sim [cm^2], \quad H \sim k_B \sim \text{energy} \sim [cm^{-1}]$$

$$\Lambda \sim \Lambda_c \sim [cm^{-2}], \quad \rho_{vac} \sim \rho_{rad} \sim \rho_m \sim \sigma \sim [cm^{-4}]$$

Finally we list the dimensionless quantities:

$$\Omega_m \sim \Omega_\Lambda \sim \Omega_K \sim \Omega_r \sim q \sim T \sim \tilde{T} \sim u \sim \alpha \sim \lambda \sim [cm^0 = 1]$$

Remember that $k, \alpha, \lambda, \Lambda, \Lambda_c$ are constant parameters.




Figure 1: Potential for the associated mechanical system. Case $\alpha \neq 0$

3 Solutions

3.1 Qualitative Behavior of Solutions

Friedmann equation in $\{u, T\}$ variables (eq 3) can also be written

$$\left(\frac{dT}{du}\right)^2 + V_{\alpha,k}(T) = \frac{\lambda}{3} \quad \text{with} \quad V_{\alpha,k}(T) = -\alpha T^4 - \frac{2}{3}T^3 + kT^2 \quad (15)$$

This is the equation of a one-dimensional mechanical system with “coordinate” T , potential $V_{\alpha,k}(T)$ (displayed in Fig 1) and total energy $\lambda/3$. We shall also set

$$Q_{\alpha,k}(T) = -V_{\alpha,k}(T) + \lambda/3 \quad (16)$$

When no confusion arises we write $V = V_{\alpha,k}$, $Q = Q_{\alpha,k}$. Formally, $du = dT/\sqrt{Q}$.

The kinetic energy being non negative, the associated mechanical system describes a horizontal line in the $(V(T), T)$ plane but never penetrates under the curve $V_{\alpha,k}(T)$ –this would correspond to u imaginary. The length of the vertical line segment between a point belonging to the curve and a point with same value of T but belonging to the horizontal line $\lambda/3$ (on which the associated mechanical system moves) is a measure of $(dT/du)^2$.

For a given value of α , the radiation parameter, and for $k = \pm 1$, the curve $V_{\alpha,k}(T)$ has typically two bumps (two local maxima). For $k = 1$ (closed universe), the right maximum occurs for a positive




Figure 2: Potential for the associated mechanical system. Case $\alpha = 0$

value of T and $V(T)$ whereas, for $k = -1$ (open case), this maximum is shifted to $T = 0$ and $V(T) = 0$. For $k = 0$ (flat case), the right maximum disappears and we are left with an inflection point at $T = 0, V(T) = 0$. If $\alpha = 0$, the curve $V(T)$ becomes a cubic (Fig. 2) and the LHS maximum disappears: it moves to $-\infty$ as α goes to 0.

Warning: figure 1 gives only the qualitative features of the curve $V_{\alpha,k}(T)$. Indeed, for reasonable values of α and λ , i.e., values compatible with experimental constraints, the vertical coordinate of the left maximum should be at least 1000 times higher than the vertical coordinate of the right maximum. Let us call $\lambda_{\pm}/3$ the ordinates of the non-zero extrema of the potential $V_{\alpha,k}(T)$ (see fig. 1). For small values of α , the values of these extrema (maxima if $k = +1$) are given by $\lambda_- \simeq k(1 - 3\alpha k)$ and $\lambda_+ \simeq \frac{1}{16\alpha^3}(1 + 12k\alpha)$; the former goes to k , and the latter to infinity when α goes to 0. Actually, if $k = 0$, one has exactly $\lambda_- = 0$ and $\lambda_+ = 1/(16\alpha^3)$.

All the recent experimental results (see sec. 4.1) seem to agree on the fact that Λ is non-zero and positive. As a consequence, λ is also positive. This will be assumed in the rest of this paper. At the end of sec. 4.1 we will show, using the experimental values (or bounds) on the densities, that if $k = 1$, then $\lambda_- < \lambda < \lambda_+$, and if $k = 0$ or -1 , then $0 < \lambda < \lambda_+$. Actually, if $k = 0$ the right maximum of the potential disappears: ($\lambda_- = 0$) and if $k = -1$, the (right) maximum becomes a (left) minimum and moves to the non-physical region

In other words, the associated mechanical system moves along an horizontal line like the one displayed in fig. 1, with an ordinate located between the two extrema of the potential $V_{\alpha,k}(T)$. Typically, a given universe starts from the right of the picture (an infinite T corresponding to the Big Bang) and moves to the left until it reaches the vertical axis ($T = 0$). This takes place in a finite conformal time u_f but corresponds to a cosmic time t going to infinity, so that, in the universe in which we live, ‘‘History’’ stops there. However, the solution can be continued for $T < 0$ (a negative radius a) until the system bumps against $V(T)$ and goes back to right infinity; the system then jumps to left infinity, follows the same horizontal line (but now from left to right) till it bumps against $V(T)$ again, and comes back. This round trip of the associated mechanical system is done in a (conformal) time $2\omega_r$ – a period of the corresponding elliptic function. If $k = 1$ there is an




Figure 3: Qualitative behavior of $T(u)$. Typical plots for cases (i) ($k = 1, \alpha = 0$), assuming $\lambda \geq 1$, (ii) ($k = 0, \alpha = 0$), assuming $\lambda \geq 0$, (iii) ($k = -1, \alpha = 0$), and case (iv) $\alpha \neq 0$, assuming $\lambda_- \leq \lambda \leq \lambda_+$. The function $T(u)$ is doubly periodic for complex u , and periodic, with period $2\omega_r$, if the conformal time u is real (graphs on the left). Only the decreasing positive branch of $T(u)$, in the first period, is “physical” i.e., describes the history of our universe (graphs on the right). In the first three cases the inflection point occurs respectively for $T > 0$, $T = 0$, and $T < 0$. We have the same type of behavior when $\alpha \neq 0$ but the curve also develops a negative connected branch (see details in fig. 4).

inflection point for $T(u)$ coming from the existence of a positive right maximum for the curve $V(T)$: the expansion speeds up anyway, but there is a time u_I for which the rate of expansion vanishes. If $k = 0$ the right maximum disappears and there is no inflection point for $T(u)$. In the case $k = -1$ the inflection point of $T(u)$ moves to the non-physical region.

Since the radius a is proportional to $1/T$, the discussion in terms of a is of course different: the system starts with $a = 0$ (Big Bang), and expands forever; as $\lambda > 1$ the expansion speeds up in all three cases $k = \pm 1, k = 0$. Let us stress the fact that only the first part of the motion of the associated mechanical system (from right-infinity to the intersection with the vertical axis) is physically relevant for the history of the universe in which we live.

The above elementary discussion shows immediately that, as a function of the conformal time u , the behavior of the reduced temperature $T(u)$ is described by fig 3. Notice that, if $\alpha = 0$ the connected negative branch(es) of this curve disappears. Call u_f the first positive zero of T , i.e., the first positive value of u for which $T(u_f) = 0$. Only that part of the curve corresponding to the interval $0 \leq u \leq u_f$ matters, for the universe in which we live.

3.2 Notations for special values of the conformal time

The end of time. Call $u_f = \int_0^\infty dT/\sqrt{Q(T)}$. The conformal time reaches u_f when the reduced temperature T becomes equal to 0. This corresponds to an infinite value of cosmic time. The value u_f is given by the previous integral but one can also determine numerically its value by looking at the first zero of the function $T(u)$ along the real axis. We call u_f “the end of time”, but this is only a shortening for “the value of the conformal time at the end of cosmic time”, since, after that, the conformal time goes on....




Figure 4: Qualitative behavior of $T(u)$, radiation being taken into account. Special values of the temperature (extrema T_a, T_b) and of the conformal time ($u_f, u_g, u_f + u_g, 2\omega_r$) are displayed on the figure. The width of the connected negative branch (between the two asymptotes) is $2\delta_c = 2\omega_r - (u_f + u_g)$, it goes to 0 when the effect of radiation is neglected, at the same time the connected negative part of the curve is shifted towards $-\infty$ i.e., the smallest (negative) extremum T_a becomes very large. Experimentally, the contribution of radiation is small, which means that $2\delta_c$ is very small compared to $2\omega_r$. The physical branch of the history of the universe is the interval $[0, u_f]$.

After the end of time: the negative history. If $\alpha = 0$ (no radiation), and for given λ , call T_M the solution of the equation $V(T) = \lambda$. If $\alpha \neq 0$, call $T_a \leq T_b$ the two solutions of the same equation (see graphs 1 and 2). In the first case, call $\delta_M = \int_{T_M}^0 dT/\sqrt{Q(T)}$. In the second case, call $\delta_M = \int_{T_b}^0 dT/\sqrt{Q(T)}$. As a function of conformal time, the reduced temperature T has a (negative) minimum T_M (if $\alpha = 0$), or T_b (if $\alpha \neq 0$) which is obtained for $u_M = u_f + \delta_M$. This occurs after the end of time. After the end of time u_f , the reduced temperature $T(u)$ becomes negative and continues to decrease until it reaches T_M (or T_b) then it starts to increase and vanishes again when $u = u_g$, with $u_g = u_f + 2\delta_M$.

After the negative history: a new beginning. When $u > u_g$, the temperature is again positive, it starts from zero and increases to infinity (big crunch), this happens when $u = u_f + u_g = 2u_M$. Although it could be a perfectly allowed region, it is not the branch of the universe in which we live. If the chosen cosmological model uses $\alpha = 0$, the value $2u_M = 2 \int_{T_M}^{\infty} dT/\sqrt{Q(T)}$ coincides with $2\omega_r$, the real period. Otherwise (i.e., if we take $\alpha \neq 0$), the curve $T(u)$ develops a negative connected branch for u between $2u_M$ and the period $2\omega_r$: *the radiation conformal shift*⁴ $2\delta_c = 2\omega_r - 2u_M$, that vanishes if $\alpha = 0$, measures the size of this unphysical branch; as we see from graph 1, it can be obtained numerically as the integral $\delta_c = \int_{-\infty}^{T_a} dT/\sqrt{Q(T)}$.

The complex history. Under the potential, i.e., for $T_a < T < T_b$ (using $\alpha \neq 0$), or for $-\infty < T < T_M$ (using $\alpha = 0$), the polynomial $Q(T)$ is negative and the conformal time is purely imaginary. As a function of the complex argument u , the function $T(u)$ is doubly periodic in the complex plane. We

⁴in ref [2] the number $2\delta_c$ was called u_c

have already determined one (real) half-period ω_r , by integrating $dT/\sqrt{Q(T)}$ along the branches $\int_{-\infty}^{T_a} + \int_{T_b}^{\infty}$, integration between iT_a and iT_b gives us the (complex) half-period $i\omega_i$. So, in terms of real quantities, we have: $\omega_i = \int_{T_a}^{T_b} dT/\sqrt{|Q(T)|}$. If one uses $\alpha = 0$, we just replace T_a by $-\infty$.

Let us summarize the previous discussion (see also fig. 4), by the following :

$$u_f = \int_0^{\infty} \frac{dT}{\sqrt{Q(T)}}, \quad \delta_M = \int_{T_b}^0 \frac{dT}{\sqrt{Q(T)}}, \quad \delta_c = \int_{-\infty}^{T_a} \frac{dT}{\sqrt{Q(T)}}, \quad \omega_i = \int_{T_a}^{T_b} \frac{dT}{\sqrt{|Q(T)|}} \quad (17)$$

$$u_M = u_f + \delta_M, \quad u_g = u_f + 2\delta_M, \quad 2\omega_r = u_f + u_g + 2\delta_c = 2u_M + 2\delta_c.$$

If one uses $\alpha = 0$, one sets $T_a \rightarrow \infty$, $T_b = T_M$, and therefore $\delta_c = 0$, $\omega_r = u_M$.

The conformal time at the inflection point u_I . From the associated mechanical system, see figs 1 or 2, we see that when the reduced cosmological constant λ is such that $\lambda_- < \lambda < \lambda_+$ (case $\alpha \neq 0$), or is such that $1 < \lambda$ (case $\alpha = 0$), the curve $T(u)$ has an inflection point between 0 and u_f , and another one (after the negative history) when $u > u_g$. This occurs both for models with $k = 1$ and $k = -1$, but this point is “physical”, i.e., it occurs for a positive temperature, only if $k = 1$. Its position u_I can be determined by solving $T''(u) = 0$. Rather than solving this equation, one may notice that, assuming $k = +1$, the value $T(u_I)$ is also equal to T_+ , the positive real value of T for which the potential $V(T)$ is maximal (T_+ is α -dependent, but $T_+ = 1$ if $\alpha = 0$), therefore, u_I can be found by solving the equation $T(u_I) = T_+$, if $\alpha \neq 0$, or the equation $T(u_I) = 1$, if $\alpha = 0$. Notice that if $k = 0$ the second derivative of T vanishes when $T = 0$, i.e., when $u = u_f$ (end of time), but does not change sign: there is no inflection point in the physical region.

The conformal time now u_0 . The above special values of the conformal time depend on the parameters used to construct the cosmological model (a universe history). In contradistinction, the conformal time u_0 just specifies the date “today”, i.e., where we are on the curve describing a universe history. This value u_0 is defined by the equation $T(u_0) = T_o$ where T_o is in principle taken from experiment (see equation 14).

Assuming $\alpha = 0$ to simplify, we have obviously $0 < u_f < u_M = \omega_r < u_g < u_g + u_f = 2\omega_r$. If $T_I > 0$, i.e., if $k = 1$, it is of course physically interesting to know if u_I is smaller or larger than u_0 . Using the experimental results on the present day values of the densities we will see that u_I is very close to u_f and still in our future, in other words $u_o < u_I < u_f$. If $k = 0$ this inflection point disappears.

Periods and Weierstrass invariants. We saw how to determine two periods $2\omega_r$ and $2\omega_i$ by integration (see above). As it is discussed in sections 3.3.1 and 3.3.2, as well as in the Appendix, another way to encode the lattice with respect to which the cosmological quantities are elliptic (in particular, doubly periodic) is to introduce the Weierstrass invariants g_2, g_3 . We shall come back to it, but let us only mention now that several mathematical computer packages offer facilities to convert Weierstrass invariants (g_2, g_3) into complex half-periods (ω_1, ω_2) , and vice versa, but the reader should remember that, given the two invariants, the choice of a base in the corresponding lattice of the complex plane is not unique (cf. Appendix), and the result returned for the periods by a computer package will depend upon the package and may even depend upon the program version! Calling $\Delta = g_2^3 - 27g_3^2 = 2^{-4}3^{-3}\lambda(\lambda - \lambda_+)(\lambda - \lambda_-)$ the modular determinant, and assuming $\Delta < 0$ as it seems to be experimentally the case (see sec. 4.1), we can choose⁵ for elementary periodicity

⁵At the moment, with Mathematica version 10, [13] the two half-periods returned by the command $\{\omega_1, \omega_2\} = \text{WeierstrassHalfPeriods}[\{g_2, g_3\}]$ are such that $\omega_r + i\omega_i = 2\omega_2$ and $-i\omega_i = \omega_1$.

cell the rhombus $\{0, \omega_r - i\omega_i, \omega_r + i\omega_i, 2\omega_r\}$, where the numbers ω_r and ω_i are real; in particular its diagonal $\{0, 2\omega_r\}$ is real, and as a function of the real variable u , the number $2\omega_r$ is the fundamental period.

3.3 Analytical solutions

3.3.1 The case without radiation ($\alpha = 0$)

For a solution starting with a Big Bang the contribution of the term αT^4 to the Friedmann equation is only important when the universe is very young. As the discussion is in any case easier in that case, we shall first assume $\alpha = 0$. The RHS of eq 3 is then a cubic polynomial, and setting $T = 6y + k/2$ brings this equation to the form

$$(dy/du)^2 = 4y^3 - g_2y - g_3 \quad (18)$$

where the parameters g_2 and g_3 , called the Weierstrass invariants, are given by

$$g_2 = k^2/12 \quad \text{and} \quad g_3 = \frac{1}{6^3}(k - 2\lambda) \quad (19)$$

The analytic solution, for the reduced temperature T , is immediate: as a function of u , it is the (scaled and shifted) Weierstrass elliptic function⁶ $y = \mathcal{P}$ corresponding to the invariants g_2 and g_3 .

$$T(u) = 6\mathcal{P}(u; g_2, g_3) + \frac{k}{2} \quad (20)$$

Since $\mathcal{P}(u) = \frac{1}{u^2} + O(u^2)$ as $u \rightarrow 0$, i.e., near the Big Bang, we see that $T(u) \sim \frac{6}{u^2} + k/2$ ($u \rightarrow 0$) but, on physical grounds, one should not use this approximation of $T(u)$ for small u since one cannot neglect the effect of radiation (the α term) near the Big Bang.

The end of time⁷ occurs at the first zero, u_f , of $T(u)$. Calling u_g the next zero, we have $u_f + u_g = 2\omega_r$ where the RHS is the period along the real axis.

Since T vanishes for u equal to u_f and $u_g = 2\omega_r - u_f$, the elliptic function $1/T$ has a pole for these two values, with a behavior dictated by the dominant term $\sqrt{\lambda/3}$ in the RHS of eq 3, so that we can write it immediately⁸ in terms of the Weierstrass function $\zeta(u)$. We obtain in this way an alternative explicit expression for the reduced temperature $T(u)$ (remember that $T(u) = 1/(a(u)\sqrt{\Lambda_c})$):

$$\frac{1}{T(u)} = \sqrt{\frac{3}{\lambda}} \quad \{\zeta(u - u_g) - \zeta(u - u_f) + \zeta(u_g) - \zeta(u_f)\} \quad (21)$$

The dedicated reader can check that the following provides still another expression for the same function $T(u)$. In this formula, $\sigma(u) = \sigma(u; g_2, g_3)$ denotes the Weierstrass σ function of the same

⁶See sec. 6.

⁷Physically, the cosmic time $t(u)$ develops a logarithmic singularity when $u \rightarrow u_f$, this is indeed “the end of time”.

⁸See the last paragraph of the appendix.

lattice. Rather than using the information that we have about the the poles and principal parts of $1/T$, it uses the fact that we know its poles and zeroes.

$$T(u) = T_M \times \frac{\sigma^2\left(\frac{u_f + u_g}{2}\right)}{\sigma^2\left(\frac{u_f - u_g}{2}\right)} \times \frac{\sigma(u - u_f) \sigma(u - u_g)}{\sigma(u) \sigma(u - u_f - u_g)} \quad (22)$$

Here $T_M = T_b = T(\omega_r)$, see figs 1,2, is the minimum value of $T(u)$; it can be determined from $dT/du = 0$, i.e., as $T_M = 6e_2 + \frac{1}{2}$ where e_2 is the real cubic root of the polynomial $4y^3 - g_2y - g_3$. This is numerically easy to find but it can also be given in closed form: $e_2 = a_+ + a_-$, where $a_{\pm} = -\frac{1}{2}(-g_3 \pm \sqrt{-\Delta/27})^{1/3}$, with $\Delta = g_2^3 - 27g_3^2$.

Eq 20 looks simpler than eqs 21 or 22 but the latter is numerically as convenient as the first and, as we shall see, generalize straightforwardly to the $\alpha \neq 0$ case.

The cosmic time. Integrating eq 2 we obtain

$$\sqrt{\frac{\Lambda}{3}} t(u) = \ln\left[\frac{\sigma(u_f + u)}{\sigma(u_f - u)}\right] - 2u \zeta(u_f) \quad (23)$$

This function is not elliptic, it has a logarithmic singularity at $u = u_f$. In a neighborhood of u_f , to the left, we have $\sqrt{\frac{\Lambda}{3}} t(u) \sim -\ln(u_f - u)$. Between u_f and u_g the cosmic time is not real.

3.3.2 The case with radiation ($\alpha \neq 0$)

From the analytic point of view, what happens is that the two poles of $T(u)$ become distinct, each one of them is therefore of first order, since $T(u)$ is still elliptic of order 2. The RHS of eq 3 is now a quartic polynomial $Q(T) = -V(T) + \lambda/3$, but such a polynomial can be brought to a cubic form by a simple change of variables. Let T_j be any one of the (possibly complex) roots of the equation $Q(T) = 0$, then the fractional linear transformation

$$y = \frac{Q'(T_j)}{4} \frac{1}{T - T_j} + \frac{Q''(T_j)}{24} \quad (24)$$

brings eq (3) to the same form as before (eq 18), but the invariants g_2 and g_3 are now given by

$$g_2 = \frac{k^2}{12} + \frac{\alpha\lambda}{3} \quad \text{and} \quad g_3 = \frac{1}{6^3}(k - 2\lambda) - \frac{\alpha\lambda k}{18} \quad (25)$$

Again, we have $y = \mathcal{P}(u; g_2, g_3)$ and one can solve eq 24 for T in terms of y to get an explicit expression for the function $T(u)$. However, in the present case, it is much simpler to express T in terms of the Weierstrass zeta function $\zeta = \zeta(u; g_2, g_3)$. The obtained expression is exactly the same as the one (eq 21) obtained in the previous section, with the difference that the invariants g_2, g_3 are now given by eq 25, and that $u_f + u_g \neq 2\omega_r$ since we have $u_f + u_g + 2\delta_c = 2\omega_r$ with $2\delta_c = 2 \int_{-\infty}^{T_a} dT/\sqrt{Q(T)} \neq 0$. With $\alpha \neq 0$, and assuming⁹ $\lambda_- \leq \lambda \leq \lambda_+$, the polynomial $Q(T)$ has two real roots $T_a \leq T_b < 0$ (we had only one, $T_M = T_b$, in the case $\alpha = 0$ and $\lambda > 1$).

⁹This corresponds to the experimental situation, see our discussion in sec. 4.1.

The cosmic time. Equation (23) is modified as follows:

$$\sqrt{\frac{\Lambda}{3}} t(u) = \ln\left[\frac{\sigma(u_f)\sigma(u-u_g)}{\sigma(u_g)\sigma(u-u_f)}\right] + u(\zeta(u_g) - \zeta(u_f)) \quad (26)$$

Compared to the case $\alpha = 0$, modifications in the interval $0 \leq u \leq u_f$ occur for very small u .

4 Experimental constraints and evolution of cosmological quantities with conformal time

The experimental values given for Ω_m^o and Ω_Λ^o in table 2 of [12] assume $\Omega_K^o = 0$ (the spatially flat “base Λ CDM model”). This very special value of the curvature density is certainly compatible with present-day experiments but, using observations of the CMB together with the results coming from detection of gravitational lensing, it seems (formula 67b of [12]) that one can only constrain Ω_K^o to percent level precision: $100 \Omega_K^o = -1.0_{-1.9}^{+1.8}$. This small value found for Ω_K^o is often described in the literature by sentences like “the universe is spatially flat”. From the analytical and geometrical points of view, the flat case $k = 0$ (that implies $\Omega_K = 0$ at all times since densities have constant sign and $k = -\text{sign}(\Omega_K)$) is however very special, see section 5. For conceptual—and philosophical—reasons, some people may find cases $k = 0$ and $k = -1$ a bit unpleasant, see in particular the discussion in pages 748, 749 of [9], while some other people, also for conceptual—and philosophical—reasons, prefer to take $k = 0$. In general we do not assume spatial flatness, and in the next three subsections we shall use values of Ω_m^o and Ω_Λ^o that allow for a non-zero curvature density Ω_K^o within the experimental bounds. For definiteness, and since the latter is only constrained to be smaller than a few percents (not such a small number), we shall take $\Omega_K^o = -0.01$, so $k = +1$ (closed case). Subsection (5) will be devoted to the case $k = 0$.

4.1 Experimental present values for density parameters, CMB temperature, and Hubble function

Newton constant and Stefan-Boltzmann constants. With $\hbar = c = 1$, the value of the Stefan-Boltzmann constant is $\sigma = \pi^2 k_B^4 / 60 = 59.8 \text{ cm}^{-4}$, where k_B is the Boltzmann constant and the Newton constant (sometimes called “Planck area” in those units) is $G = 2.61 \cdot 10^{-66} \text{ cm}^2$.

CMB temperature. The present day value of the CMB temperature given by [12] is $\tilde{T}_o = 2.7255 \text{ K}$.

Hubble parameter. The present value of the Hubble “constant” (i.e., now), given by [12] is $H_o = 100 h \text{ km sec}^{-1} \text{ Mpc}^{-1}$, with $h \simeq 0.688 \pm 0.008$. Its mean experimental value, in natural units, is therefore $H_o = (7.437 \pm 0.13) \cdot 10^{-29} \text{ cm}^{-1}$. In the following we shall take h (later called h_o) equal to 0.688.

Radiation density Ω_r . From eq. 14 and using the present day experimental value of the CMB temperature \tilde{T}_o , one finds $\alpha \Lambda_c T_o^4 = 5.23 \cdot 10^{-63} \text{ cm}^{-2}$. From eq. 8 for Ω_r , and using the experimental value for H_o , one finds $\Omega_r^o = (5.45 \pm 0.20) \cdot 10^{-5}$. This value is therefore obtained from the measurements of \tilde{T}_o and H_o . Taking into account the effect of all relativistic particles (massless neutrino)

is not expected to change this value much. The term Ω_r^o , which is well determined but very small compared to the other terms, is often dropped from eq 7 written $1 = \Omega_r^o + \Omega_K^o + \Omega_\Lambda^o + \Omega_m^o$. It is certainly legitimate to perform this approximation at present times, but of course not at all times and certainly not at the beginning of the expansion.

Curvature density (cf. discussion at the beginning of this section). In contradistinction to the latter, the present day curvature density Ω_K^o is not well determined, but it seems to be also quite small. For this reason it may look legitimate to drop the term Ω_K^o from equation eq. 7. Doing so is certainly valid at present times, but setting $k = 0$ (which implies $\Omega_K = 0$ at all times) is a strong and disputable hypothesis on the topology and the dynamics of the mathematical model chosen for our universe. In any case we want to study and display the *evolution* of Ω_K as a function of u . We take $\Omega_K^o = -0.01$, a value compatible with the experimental bounds. This implies¹⁰ $k = +1$.

Matter density. According to the recent measurements (see in particular [12]), Ω_m^o is about 0.3. The value quoted by [7] for CMB + WMAP + BAO is $\Omega_m^o = 0.293 \pm 0.010$. For definiteness, we shall take $\Omega_m^o = 0.293$ in the following numerical experiments. One could for instance assume a contribution of 0.047 from baryons and 0.246 from dark matter but the fact that matter density seems to be dominated by its so-called “dark matter” component (compared to its baryonic component) is irrelevant for the present analysis since both contribute in the same way to Friedmann equations.

Vacuum density (often called “dark energy density”). The experimental constraints — see those given by [12], in particular fig. 25 of [12] — are $\Omega_\Lambda^o = 0.707 \pm 0.010$. We shall take $\Omega_\Lambda^o = 0.717$.

From equation 13 and using the experimental values of Ω_r^o , Ω_m^o and Ω_Λ^o , one finds $\lambda\alpha^3 \simeq 8.20 \cdot 10^{-12}$. This value is independent of the hypothesis made about the curvature density (in particular it also holds if $k = 0$).

If we assume $k = \pm 1$, the equations 12 make sense and give individual values — or constraints — for the parameters α and λ . The sum of the four densities should be 1 at all times, but since they enter multiplicatively in the expressions of α and λ , one obtains reasonable bounds by keeping Ω_K^o as a free variable in eqs 12. Doing so leads to $\alpha \simeq 2.70 \cdot 10^{-4} |\Omega_K^o|$ and $\lambda \simeq 0.415 / |\Omega_K^o|^3$. If we assume that the present day value of the curvature density is bounded, in norm, by 10^{-2} , we find $\alpha < 2.7 \cdot 10^{-6}$ and $\lambda > 4.1 \cdot 10^5$. As already written, we shall use $|\Omega_K^o| = 10^{-2}$ in the following numerical experiments.

Warning: if we assume $k = 0$, so that Ω_K is strictly zero at all times, we cannot use eqs 12 to provide individual values (or bounds) for α or λ . The only constraint that we have is on the value of $\lambda\alpha^3$.

We called $\lambda_\pm/3$ the ordinates of the left and right maxima of the potential $V_{\alpha,k}(T)$ of the associated mechanical system (see fig. 1 in sec. 3.1).

If $k = 0$, one has $\lambda_+ = \frac{1}{16\alpha^3}$ and $\lambda_- = 0$. The numerical value obtained previously for the product $\lambda\alpha^3$ is clearly much smaller than $\frac{1}{16}$. Therefore $\lambda \ll \lambda_+$. On the other hand, experimentally, Λ and therefore λ , are positive. So we have the constraint $\lambda_- = 0 < \lambda \ll \lambda_+$.

¹⁰The spatial universe is then, topologically, a sphere $S^3 \sim SU(2)$ or a quotient of the latter by a (discrete) binary polyhedral subgroup of $SU(2)$.

If $k = +1$, from the experimental bounds on Ω_K^o , we conclude, as before, that the value of α is itself very small, so that we can use for λ_{\pm} the approximations $\lambda_- \simeq (1 - 3\alpha)$ and $\lambda_+ \simeq \frac{1}{16\alpha^3}(1 + 12\alpha)$ given in 3.1, or even $\lambda_- \simeq 1$ and $16\lambda_+\alpha^3 \simeq 1$, but λ is at the same time much larger than 1 and such that $\lambda\alpha^3 \simeq 8.20 \cdot 10^{-12}$. As a result: $\lambda_- < \lambda \ll \lambda_+$.

The case $k = -1$ can be discussed similarly, but here λ_- is negative and corresponds to a minimum of the potential; it is also associated with an inflection point of the curve $T(u)$ but this occurs when T is negative, i.e., in the unphysical region (after the end of time).

In all cases, the constraint $\lambda_- < \lambda < \lambda_+$ tells us (see fig. 1) that, as a function of conformal time u , the temperature will decrease from plus infinity to 0, for some value u_f (the end of time), it will then become negative until it reaches a minimum (the associated mobile bumps against the potential since $\lambda \ll \lambda_+$), and goes back to (plus) infinity. This branch is followed, since $\alpha \neq 0$, by a round trip of the mobile in the non-physical region (a negative branch of the curve $T(u)$). In the case $k = 1$ the curve $T(u)$ has an inflection point in the physical region; this inflection point disappears if $k = 0$, and moves to the unphysical region if $k = -1$, but in the three cases, given the experimental constraints on the densities and the corresponding bounds on λ , the overall features of the temperature, as a function of the conformal time, are the same.

4.2 Evolution of the cosmological quantities as functions of conformal time

In the cases $k = \pm 1$, from the independent measurements of H_o , Ω_m^o , Ω_Λ^o and \tilde{T}_o , one can determine the evolution of all cosmological quantities in terms of the conformal time u . Here we take $k = +1$. The various quantities are obtained, in turn, as follows. Eq. 14, using \tilde{T}_o , give the product $\alpha\Lambda_c T_o^4$. Then eq. 8, with H_o , gives Ω_r^o . The densities Ω_m^o and Ω_K^o , together with eq. 7 gives Ω_K^o . The parameters α and λ are then obtained from eq. 12. Λ comes from Ω_Λ^o and H_o , using eq. 8. This gives Λ_c since λ is known, and also T_o , since $\alpha\Lambda_c T_o^4$ is known. Eqs. 25 give the two invariants g_2 and g_3 . The Weierstrass functions are then perfectly determined. The different cosmological quantities of interest, as functions of u , are finally given by eqs. 5, 21, together with 16, 17 (or 20 if radiation is neglected), and 8.

Temperature \tilde{T}

As a function of conformal time, the evolution of the usual temperature \tilde{T} is displayed in fig. 5. Remember that it is obtained by multiplying T by the known (and constant) scaling factor given by eq. 14. As already mentioned several times, this function is periodic in the complex plane, and in particular along the real axis, but the physical branch (i.e., the history of our universe) is only given by the interval $[0, u_f]$, the value u_f being the “end of time”, which corresponds to $t \rightarrow \infty$ in terms of cosmic time. With $k = 1$, i.e., in a universe with the topology of S^3 (or a discrete quotient of the latter), the fact that $u_f \leq 2\pi$ means that an observer looking forward will never be able to see the back of his head, even if he waits for a very long time.




Figure 5: Evolution of the universe temperature \tilde{T} as a function of conformal time. Cases: (i) In the interval $[0, u_f]$, (ii) Around the present value u_o , (iii) For a real period $[0, u_f + u_g \lesssim 2\omega_r]$. The physical branch is the interval $[0, u_f]$ where the curve intersects the real axis (first zero): u_f is the end of time. The curve possesses a negative branch in the interval $[u_f + u_g, 2\omega_r]$ which is not displayed (see figs 3, 4 and 13).




Figure 6: Evolution of the radiation density $\Omega_r(u)$ as a function of conformal time. Cases: (i) Near the Big Bang. (ii) In the interval $[0, u_f]$, the physical branch. (iii) Around the present value u_o . (iv) For a real period $[0, u_f + u_g \lesssim 2\omega_r]$. In the tiny interval $[u_f + u_g, 2\omega_r]$ the curve has two infinite positive branches that are not displayed.

Radiation, matter, vacuum, curvature densities $\Omega_r, \Omega_m, \Omega_\Lambda, \Omega_K$ and $\Omega = \Omega_m + \Omega_\Lambda + \Omega_r$

The curves displaying the behavior of those densities, as functions of conformal time, are respectively given in figures 6, 7, 8, 9, 10. We already commented about the fact that we choose (arbitrarily) the value $\Omega_K^o = -0.01$ (so $k = +1$), which is non-zero, but compatible with the experimental bounds, to perform the calculations leading to the following plots. The values obtained for the conformal time u of the various points of interest (extrema, inflection point, present day value, end of time, etc.), as well as the values obtained for the modular parameters (Weierstrass invariants, periods, j-function, etc), are quite sensitive to the choice of Ω_K^o ; however, the overall features of the curves are rather stable, given the present values of the other densities.




Figure 7: Evolution of the matter density $\Omega_m(u)$ as a function of conformal time. Cases: (i) Near the Big Bang. (ii) In the interval $[0, u_f]$, the physical branch. (iii) Around the present value u_o . (iv) For a real period $[0, u_f + u_g \lesssim 2\omega_r]$. In the tiny interval $[u_f + u_g, 2\omega_r]$ the curve has two infinite negative branches that are not displayed.




Figure 8: Evolution of the vacuum density $\Omega_\Lambda(u)$ as a function of conformal time. Cases: (i) In the interval $[0, u_f]$, the physical branch. (ii) Around the present value u_o . (iii) For a real period $[0, u_f + u_g \lesssim 2\omega_r]$. In the tiny interval $[u_f + u_g, 2\omega_r]$ the curve has two infinite positive branches that are not displayed.




Figure 9: Evolution of the curvature density $\Omega_K(u)$ as a function of conformal time. Cases: (i) In the interval $[0, u_f]$, the physical branch. (ii) Around the present value u_o . (iii) For a real period $[0, u_f + u_g \lesssim 2\omega_r]$. In the tiny interval $[u_f + u_g, 2\omega_r]$ the curve has two infinite negative branches that are not displayed.




Figure 10: Evolution of the sum $\Omega(u) = \Omega_m(u) + \Omega_\Lambda(u) + \Omega_r(u)$ as a function of conformal time. Cases: (i) In the interval $[0, u_f]$, the physical branch. (ii) Around the present value u_o . (iii) For a real period $[0, u_f + u_g \lesssim 2\omega_r]$. In the tiny interval $[u_f + u_g, 2\omega_r]$ the curve is not displayed.




Figure 11: Evolution of the Hubble “constant” $H(u)$ as a function of conformal time. We plot $h(u)$, with $H(u) = 100 h(u) \text{ km s}^{-1} \text{ Mpc}^{-1}$. Cases: (i) In the interval $[0, u_f]$, the physical branch. (ii) Around the present value u_o . (iii) For a real period $[0, u_f + u_g \lesssim 2\omega_r]$. The branch of the curve in the tiny interval $[u_f + u_g, 2\omega_r]$ is not displayed.

Hubble function H and deceleration function q

Rather than plotting the Hubble function $H(u) \times cm$, we give in fig. 11 the evolution of the h coefficient as a function of conformal time. As usual, $H = (100 h 10^5 \text{ cm}) / (s \text{ Mpc})$, so $H \times cm = 1.0810076 10^{-28} h$, with $h_o = 0.688$, the present value of h .

Another useful quantity is the *deceleration function* — a misnomer¹¹ since we seem to live (now) in an accelerating expanding universe: $q = -a(da/dt)^{-2}(d^2a/dt^2) = \Omega_m/2 - \Omega_\Lambda + \Omega_r$. It can be expressed in terms of $T(u)$ as follows:

$$q = \frac{-\frac{\lambda}{3} + \frac{1}{3}T^3 + \alpha T^4}{\frac{\lambda}{3} - kT^2 + \frac{2}{3}T^3 + \alpha T^4}$$

The behavior of this function $q(u)$ is given in fig. 12.

¹¹In the old books it is not uncommon to find q defined as $\Omega_m/2$ since it was then fashioned to set to zero both the vacuum and the radiative contributions.




Figure 14: Evolution of the spatial scale factor $a(u)/cm$ as a function of conformal time, (i) in the interval $[0, u_f]$, the physical branch. (ii) For a real period $[0, u_f + u_g \lesssim 2\omega_r]$. In the tiny interval $[u_f + u_g, 2\omega_r]$ the curve is not displayed. (iii) Evolution of the cosmic time $t(u)/cm$ as a function of conformal time.

Near the Big Bang, we have $\sqrt{\Lambda/3} a(u) = (\mathcal{P}(u_f) - \mathcal{P}(u_g))u + \frac{1}{2}(\mathcal{P}'(u_g) - \mathcal{P}'(u_f))u^2 + O(u)^3$ and $\sqrt{\Lambda/3} t(u) = \frac{1}{2}(\mathcal{P}(u_f) - \mathcal{P}(u_g))u^2 + \frac{1}{6}(\mathcal{P}'(u_g) - \mathcal{P}'(u_f))u^3 + O(u)^4$ so, $a(t) \sim \sqrt{2} (\frac{\Lambda}{3})^{-1/4} (\mathcal{P}(u_f) - \mathcal{P}(u_g))^{1/2} t^{1/2}$.

If radiation is neglected, the first term in the Taylor expansions of $a(u)$ and $t(u)$ vanish, since in that case $\mathcal{P}(u_f) = \mathcal{P}(u_g) = -k/12$, and we have instead $a(t) \sim (\frac{\Lambda}{3})^{-1/6} 3^{1/3} (\frac{3}{2})^{1/3} (\mathcal{P}'(u_g) - \mathcal{P}'(u_f))^{1/3} t^{2/3}$. We recover, in both cases, the known results for the power-law behavior of $a(t)$ which can also be derived directly by keeping only the leading term in the differential equation for the function $a(t)$.

4.3 Tables

The experimental inputs are in boldface. $\Omega_r(u_o)$ is exactly deduced from the measurements of the temperature $\tilde{T}[u_o]$ and the Hubble constant H_o . Remember that we choose for $\Omega_K[u_o]$ an arbitrary non-zero negative value compatible with the experimental bounds (so $k = +1$). We recall the notations used for special values of the conformal time u : 0 (Big Bang), u_o (present value), u_f (first zero of T , i.e., end of cosmic time: $t \rightarrow \infty$), u_g (second zero of T), $2\omega_r$ (real period of T), u_I (inflection point for T), u_k (value of u for which Ω_K as an extremum in the interval $[0, u_f]$). The other quantities are either standard in cosmology (their definitions have been recalled in the text), or they are related to modular considerations: the Weierstrass invariants g_2, g_3 have been defined in sections 3.3.1, 3.3.2, see also the end of sec. 3.2 for the half-periods (ω_1, ω_2) , and we refer to the appendix for the other quantities (modular discriminant Δ , modular parameter τ and Klein invariant j).

$$\begin{array}{llll}
 H_o * cm = \mathbf{7.44 10^{-29}} & a_o/cm = 1.31 10^{28} & t_o/cm = 1.31 10^{28} & H_o * t_o = 0.973 \\
 \Lambda * cm^2 = 1.19 10^{-56} & \Lambda_c * cm^2 = 2.90 10^{-62} & \lambda = 409049. & \alpha = 2.72 10^{-6} \\
 \rho_m[u_o] * cm^4 = 7.41 10^7 & \rho_{vac} * cm^4 = 1.81 10^8 & C_m/cm = 3.91 10^{30} & q_o = -0.57
 \end{array}$$

$u[0]=0$	$u_k=0.272$	$u_o=0.330$	$u_I=0.441$	$u_f=0.444$
$\tilde{T}[0]=\infty$	$\tilde{T}[u_k]=4.626$	$\tilde{T}[u_o]=\mathbf{2.725}$	$\tilde{T}[u_I]=0.062$	$\tilde{T}[u_f]=0$
$T[0]=\infty$	$T[u_k]=74.22$	$T[u_o]=42.72$	$T[u_I]=1$	$T[u_f]=0$
$\Omega_m[0]=0^a$	$\Omega_m[u_k]=0.675$	$\Omega_m[u_o]=\mathbf{0.293}$	$\Omega_m[u_I]=4.89 \cdot 10^{-6}$	$\Omega_m[u_f]=0$
$\Omega_\Lambda[0]=0$	$\Omega_\Lambda[u_k]=0.338$	$\Omega_\Lambda[u_o]=\mathbf{0.717}$	$\Omega_\Lambda[u_I]=1.$	$\Omega_\Lambda[u_f]=1.$
$\Omega_K[0]=0$	$\Omega_K[u_k]=-0.014$	$\Omega_K[u_o]=\mathbf{-0.01}$	$\Omega_K[u_I]=-7.33 \cdot 10^{-6}$	$\Omega_K[u_f]=0$
$\Omega_r[0]=1^b$	$\Omega_r[u_k]=2.043 \cdot 10^{-4}$	$\Omega_r[u_o]=\mathbf{5.21992 \cdot 10^{-5}}$	$\Omega_r[u_I]=1.992 \cdot 10^{-11}$	$\Omega_r[u_f]=0$
$a[0]=0$	$a[u_k]/cm=7.90 \cdot 10^{28}$	$a[u_o]/cm=1.34 \cdot 10^{29}$	$a[u_I]/cm=5.86 \cdot 10^{30}$	$a[u_f]/cm=\infty$
$t[0]=0$	$t[u_k]/cm=7.00 \cdot 10^{27}$	$t[u_o]/cm=1.31 \cdot 10^{28}$	$t[u_I]/cm=7.22 \cdot 10^{28}$	$t[u_f]/cm=\infty$
		$g_2=0.454$	$g_3=-3787.$	
	$\omega_1=-0.387i$	$\omega_2=0.336+0.194i$	$\omega_r=0.671$	$u_g=0.889$
	$\tau=-0.50+0.86i$	$j=-2.412 \cdot 10^{-10}$	$\Delta=-3.87 \cdot 10^8$	

4.4 Comments

Very quickly after the Big Bang, taking into account the radiation term in Friedmann equations, i.e., using eq. 21 together with eq. 17 rather than using only eq. 20, only changes, in the previous table, the third decimal digit in the values of all quantities of interest, at least in the “physical branch” of the history ($u < u_f$). However the influence of this term is very important when $u \rightarrow 0$. See in particular the values of the densities at the point $u = 0$. Analytically the curve $T(u)$ also develops a connected negative branch, between $u_f + u_g$ and $2\omega_r$, but this is physically irrelevant since it occurs long after the end of time u_f . If radiation is neglected ($\alpha = 0$), the quartic term in Friedmann equations disappears, and $2\omega_r = u_f + u_g$.

The (negative) curvature density Ω_K has a minimum for a value u_k in the physical branch, see fig. 9 and the table in sec. 4.3, showing that $0 < u_k < u_o$, so this happened in our past. It is incorrect to think or claim that the experimental smallness of Ω_K^o implies that it was even smaller in the past: indeed, although Ω_K goes to 0 at the Big Bang, it has – actually it had – an extremum for $u = u_k$.

If one takes radiation into account, as one should, one finds that the matter density Ω_m is zero at the Big Bang, and quickly increases — see fig. 7 — to reach a maximum almost equal to 1 because the other densities are negligible in this region. Using the same experimental values as before, the maximum is reached for a value of conformal time $u_m \simeq 0.10526$. After that, Ω_m stabilizes for a while and starts to slowly decrease ($\Omega_m^o \simeq 0.293$ nowadays) to reach 0 at the end of time. The sharp increase of Ω_m at the Big Bang is already encoded in the Friedmann equations (with radiation, of course). Notice that we do not introduce any inflation mechanism in this paper. Near $u = 0_+$, this density behaves as $\frac{2u}{3\sqrt{\alpha}}$. If one incorrectly forgets the radiative term contribution, the matter density behaves instead as $1 + \frac{u^2}{4}$ when $u \rightarrow 0$; notice that the former expression does not go continuously to the latter when $\alpha \rightarrow 0$.

^aIf one does not include the radiation density in the Friedmann equations, i.e., if one sets $\Omega_r = 0$ at all times, one finds $\Omega_m[0] = 1$ instead of 0, in agreement with the fact that, in that case, $\Omega_m[0] + \Omega_\Lambda[0] + \Omega_K[0] = 1$.

^bThis is in agreement with the fact that $\Omega_m[0] + \Omega_\Lambda[0] + \Omega_K[0] + \Omega_r[0] = 1$, see previous footnote.

With the given experimental values, one finds $0 < u_m < u_k < u_o < u_I < u_f$. The inflection point u_I of $T(u)$ is therefore still in our future, but it is located very near the end of time u_f , and is hard to see on the figures.

The table gives $u_f = 0.444$, therefore, looking forward, you cannot hope to see the back of your head (assuming $k = 1$) since $u_f < 2\pi$, even if you wait a very long time. However, as already mentioned, this value is quite sensitive to the (arbitrary) chosen input for Ω_K^o , so, there is still hope!

In contradistinction with other quantities (like $g_2, g_3, \lambda, \alpha$, etc.), the numerical values found for t_o , the cosmic time, now, and a_o , the cosmic scale factor, now, are quite stable with respect to the choice of Ω_K^o . In terms of cosmic time, here are a few important dates in the history of the universe: $t(0) = 0$, $t(u_m) = 4.58 \cdot 10^8 \text{ yr}$, $t(u_k) = 7.40 \cdot 10^9 \text{ yr}$, $t(u_o) = t_o = 1.38 \cdot 10^{10} \text{ yr}$, $t(u_I) = 7.63 \cdot 10^{10} \text{ yr}$, $t(u_f) = \infty$.

One should remember that the function T , introduced in the first section, differs from the usual temperature \tilde{T} by a multiplicative dimensionless factor $(\frac{3}{8\pi G} \frac{\alpha \Lambda_c}{4\sigma})^{1/4}$, see eq. 14. Using the values of the previous table, this factor is about $6.23 \cdot 10^{-2}$. It is quite sensitive to the input chosen for Ω_K .

Analytically, the main qualitative feature that distinguishes the case $\alpha \neq 0$ from the case $\alpha = 0$ is that, in the former, the two poles of $T(u)$ are distinct, consequently, each of them is of first order.

Because of the smallness of the j -invariant, the modular parameter τ appears to be numerically very close to $\exp(2i\pi/3)$, but it cannot be equal to this special value since $k = 1$ and $\alpha\lambda > 0$ imply $g_2 \neq 0$ (see eq. 25), so j , given by eq. 35, is strictly non-zero. We shall return to this in sec. 5.2.

5 Some specific features of the flat case $k = 0$

If one assumes $k = 0$ from the very beginning, the Friedmann equation simplifies (the quadratic term vanishes), and becomes $(\frac{dT}{du})^2 = \alpha T^4 + \frac{2}{3}T^3 + \frac{\lambda}{3}$, with, as usual, $\lambda = \frac{\Lambda}{\Lambda_c}$, $\sqrt{\Lambda_c} = \frac{2}{3C_m}$ and $\alpha = C_r \Lambda_c$. We have immediately $\frac{\Omega_m}{\Omega_\Lambda} = \frac{2}{\lambda}T^3$, $\frac{\Omega_r}{\Omega_m} = \frac{3}{2\alpha}T$ but this information, taken at conformal time u_o , is insufficient to obtain separately T_o , α or λ . In other words, given the values of H_o , \tilde{T}_o , Ω_m^o , Ω_Λ^o , and Ω_r^o (the sum of the last three being set to 1), one cannot obtain explicit values for all the cosmological quantities appearing in the table given in section 4.3, for instance one cannot obtain the values of λ , of α , or of the conformal time “now” i.e., u_o : one extra piece of data, that could be for instance Λ_c , is missing. However, as we shall see below, the experimental data singles out a well-defined isomorphism class of elliptic curves.

5.1 General study

If we don't neglect radiation, the invariants are given by equations 25 where we set $k = 0$. So

$$g_2 = \alpha \lambda / 3, \quad \text{and} \quad g_3 = -2\lambda / 6^3 \tag{27}$$

From equations 8, one immediately obtains:

$$\frac{27}{16} \lambda \alpha^3 = \frac{\Omega_r^3 \Omega_\Lambda}{\Omega_m^4} \quad (28)$$

As $\lambda \alpha^3$ does not depend upon the value of the conformal time, it can be evaluated by replacing the densities by their present-day values. Notice that this quantity is positive, although rather small; indeed, using the same experimental values as in the previous section, one finds $\lambda \alpha^3 \simeq 8.20 \cdot 10^{-12}$.

We cannot, in the case $k = 0$, obtain separately the values of λ or α from the measurement of densities. However, since the product $\lambda \alpha^3$ is fixed, we can investigate what happens under scaling. The value of the cosmological constant Λ being determined, changing the parameter Λ_c (or, equivalently the “mass” of the universe) amounts to change the value of their ratio, λ . We therefore set $\alpha' = \alpha s^2$ and $\lambda' = \lambda/s^6$, where s is an arbitrary parameter; in this way $\lambda' \alpha'^3 = \lambda \alpha^3$ is kept fixed. From the expressions of the Weierstrass invariants g_2 and g_3 , we find immediately $g'_2 = g_2/s^4$ and $g'_3 = g_3/s^6$. Setting $\Delta = g_2^3 - 27g_3^2$, $j = g_2^3/\Delta$ and using the same definitions for Δ' and j' in terms of the prime invariants g'_2, g'_3 , we notice that $\Delta' = g_2'^3 - 27g_3'^2 = (g_2^3 - 27g_3^2)/s^{12} = \Delta/s^{12}$ so that $j' = j$. The quantity Δ is known as the modular discriminant, and j as the Klein invariant (see the Appendix for details). In other words, if one assumes $k = 0$, the measurement of Ω_m^o , Ω_Λ^o , and Ω_r^o (the sum of the three being set to 1), H_o and \tilde{T}_o does not specify a single lattice in the complex plane, or the corresponding field of elliptic functions, but a class of homothetic lattices. Equivalently, this experimental data does not single out an elliptic curve but an isomorphism class of elliptic curves specified by the j -invariant. If one introduces rescaled conformal time parameters $u' = s u$ (with the same rescaling for periods, of course), the invariants g_2, g_3 , and the discriminant Δ , are rescaled as previously, in agreement with the general results recalled in the Appendix.

In terms of cosmological parameters the modular discriminant Δ is readily evaluated, one finds $\Delta = \frac{\lambda^2}{2 \times 6^3} (-1 + 16\lambda\alpha^3)$, so that $\Delta < 0 \Leftrightarrow \lambda\alpha^3 < 1/16$, a condition that obviously holds experimentally, since $\lambda\alpha^3 \simeq 8.20 \cdot 10^{-12}$. Δ is therefore negative. From the expression of Weierstrass invariants g_2, g_3 , one obtains the Klein invariant

$$j = 1 + \frac{1}{-1 + 16\lambda\alpha^3} = 1 + \frac{1}{-1 + \frac{256}{27} \frac{\Omega_\Lambda \Omega_r^3}{\Omega_m^4}} \quad (29)$$

Using the same experimental values as in the previous section, namely $\Omega_m^o = 0.293$, $\Omega_\Lambda^o = 0.717$, and $\Omega_r^o = 5.20 \cdot 10^{-5}$ one finds $j \simeq -7.50 \cdot 10^{-14}$.

A value of j being given, there are several methods to find a modular parameter τ , in the upper half-plane, such that $j(\tau) = j$. One method to determine this parameter (defined only up to a modular transformation) is recalled in the Appendix, but in our case, the numerical precision on the – very small – value of j is insufficient to distinguish τ from the boundary point $exp(2i\pi/3)$ where j is exactly 0, the latter situation occurring when the radiation contribution to Friedmann equation is neglected (next subsection).

The expressions giving the function $T(u)$, the densities, etc, in terms of Weierstrass functions, are the same as in the previous section, but for the fact that the invariants g_2, g_3 are slightly simpler since $k=0$. The overall features of the curves showing the evolution of cosmological quantities of interest are also similar, but for the fact Ω_K is now strictly zero, and that there is no inflection point in the physical region for the curve $T(u)$, as it is a priori clear by looking at the behavior of the associated mechanical system for $k = 0$, in fig. 1 and 2.

5.2 Neglecting radiation

k being set to 0, let us assume moreover that we neglect the radiative contribution (i.e., we also set $\alpha = 0$). This is of course only an approximation, as the density Ω_r , as measured today, and although small, is clearly non-zero. Nevertheless, this limit case allows one to obtain several interesting exact results. Setting $T = 6y$ in the Friedmann equation gives: $\frac{dy^2}{du} = 4y^3 + \frac{\lambda}{108}$. The solution is immediate: y is the Weierstrass elliptic function, with invariants $g_2 = 0$ and $g_3 = -\lambda/108$.

$$T(u) = 6 \mathcal{P}(u; g_2 = 0, g_3 = -\frac{\lambda}{108})$$

The end of time u_f , i.e., the first (real) positive zero of T will exist only if $g_3 < 0$, i.e., iff $\lambda > 0$, otherwise it is complex. Since, experimentally (nowadays), the cosmological constant Λ is positive, we have also $\lambda > 0$ and u_f exists. The determination of the zeros of the Weierstrass \mathcal{P} function, in one periodicity cell, is in general a difficult problem — see the article [6] where the problem is solved in full generality – but in the present case, one does not need to use these techniques. Indeed, it is not too difficult, using elliptic integrals, to solve the particular equation $\mathcal{P}(u, 0, 4) = 0$, i.e., for $g_2 = 0$, $g_3 = 4$. One finds the solution $u = \frac{i}{6} B(\frac{1}{6}, \frac{1}{3})$, where B is the Euler beta function. Since, by scaling (see Appendix), $\mathcal{P}(u; g_2, g_3) = s^2 \mathcal{P}(u; g_2/s^4, g_3/s^6)$, we have $\mathcal{P}(u; 0, g_3) = s^2 \mathcal{P}(u; 0, g_3/s^6)$. If $g_3/s^6 = 4$, the solution is known (cf supra), so we take $s = (g_3/4)^{1/6}$, and $\mathcal{P}(u; 0, g_3)$ will vanish for a value $u_f = \frac{i}{6} B(\frac{1}{6}, \frac{1}{3}) \frac{1}{(g_3/4)^{1/6}}$. We summarize the discussion as follows:

Consider the equation $\mathcal{P}(u; 0, g) = 0$. If $g > 0$ there is no real solution. If $g < 0$, there are two real zeroes (u_f and u_g) in each periodicity cell, $u_f + u_g$ being the smallest real period, and the position of the first is:

$$u_f = \frac{\Gamma(1/6) \Gamma(1/3)}{2^{2/3} 3 (-g)^{1/6} \sqrt{\pi}}$$

Here Γ is the Euler Gamma-function. In cosmology, with $k = 0$ and $\lambda > 0$, one has $g_3 = -\lambda/108$. Therefore the end of time u_f that solves the equation $T(u_f) = 6 \mathcal{P}(u_f; 0, g_3) = 0$ is given by :

$$u_f = \frac{2^{2/3} \sqrt{3} \Gamma(1/3) \Gamma(7/6)}{\lambda^{1/6} \sqrt{\pi}} \quad (30)$$

In the theory of elliptic functions, the case $g_2 = 0$ is often referred to as the *equianharmonic case*. This name¹² already appears in Abramowitz and Stegun [1], where it refers to the case $g_2 = 0$, $g_3 = 1$; the case where $g_2 = 0$ and g_3 is an arbitrary complex number can be obtained from the latter by scaling and it is therefore justified to keep the same terminology. For $g_2 = 0$, $g_3 = 1$, the equianharmonic half-periods are known [1, 10]: $(\frac{\Gamma(1/3)^3}{4\pi}, e^{i\pi/3} \frac{\Gamma(1/3)^3}{4\pi})$. The quantity called ω_r in this paper (half the smallest real period for the temperature, expressed as a function of conformal time) is then immediately obtained by scaling, one gets:

$$\omega_r = \frac{3 \Gamma(1/3)^3}{2 \cdot 2^{2/3} \lambda^{1/6} \pi} \quad (31)$$

This universe is therefore very special, not only geometrically, since we have set $k = 0$, but also because of its dynamics. Indeed, when the first Weierstrass invariant vanishes ($g_2 = 0$), the Klein j -invariant vanishes and the modular parameter (ratio of the periods) of the associated elliptic curve

¹²which means “equally anharmonic” since it is related to a geometrical configuration where the anharmonic ratio (cross ratio) is a cube root or a sixth root of unity.

(or of the corresponding lattice, see our discussion in the Appendix) can be chosen at the corner¹³ ($e^{2i\pi/3}$) of the boundary of the fundamental domain for the full modular group. The discussion carried out in the previous subsection holds: when $k = 0$ the available experimental data does not single out a specific elliptic curve but an isomorphism class of elliptic curves specified by the j -invariant. If moreover one takes $\alpha = 0$, like in this section, the j -invariant is zero (equianharmonic case).

6 Miscellaneous (slightly eccentric) comments

The elliptic curve associated with a given cosmology: is the universe special ? The flat case ($k = 0$) with no radiation pressure ($\alpha = 0 \Leftrightarrow \Omega_r(u) = 0$) is clearly special since the associated modular parameter sits in a corner, $exp(2i\pi/3)$, of the fundamental domain (equianharmonic case). Whether or not $k = 0$ is a debatable issue: it seems to be experimentally clear, nowadays, that Ω_K^o is “small”, but Ω_K is a time-dependent function and one should not set it to zero at all times unless one can argue, on some theoretical grounds, that the constant k should be 0, rather than 1 or -1 (see the comments in sec. 4). Assuming $k = 0$ is certainly convenient from an experimental point of view and may be to the liking of partisans of the inflation mechanism (although widely accepted, one should remember that strong criticisms have been voiced against inflation). On the other hand, considering that space is flat, or asymptotically flat, and that it is filled in first approximation with an ideal fluid whose density goes to zero at infinity, may be perceived by some people as a retrograde idea going against the evolution of the scientific concepts that were at the roots of General Relativity and Friedmann equations. Any kind of hypothesis made on k will also trigger epistemological or philosophical debates, for instance: should one believe that the universe “exists” by itself, independently of our more or less elaborated fits?

The no-radiation pressure hypothesis is of different nature, as it is clearly only an approximation: although small, Ω_r^o is non-zero and it is well-measured. Whether or not $k = 0$, one comment that one can make is the following: for the Weierstrass \mathcal{P} function to have real values —the CMB temperature is real— on the real axis (conformal time), one needs the associated lattice to be identical with its conjugate, equivalently g_2 and g_3 should be real; this is automatic (see eqs. 25) in the framework of Friedmann equations, then the j -invariant is real as well, so the modular parameter τ , taken in the standard fundamental domain, should belong either to the boundary of the domain, or to the half-axis iy , with $y > 1$, since these geodesic arcs are precisely those subsets where j takes real values. It seems unreasonable to extend this kind of discussion further... Elliptic curves can be “special” for a variety of arithmetical reasons, but it is clear that human experiments, in cosmology, do not give exact results in the sense of arithmetics!

Aeons. The solutions of Friedmann equations, written as differential equations with respect to conformal time u , and almost all quantities of cosmological interest related to these solutions, are periodic functions of u , and are actually bi-periodic (elliptic functions) when the variable u is extended to the complex plane. They automatically describe a denumerable infinity of identical cosmologies (depending upon the choice of the parameters, only one interval within each real period may describe a possible physical history). For a wide range of parameters — including the values

¹³Choosing $e^{2i\pi/3}$ or $e^{i\pi/3}$ is irrelevant: both values are equivalent under the modular transformation $\tau \mapsto \tau + 1$.




Figure 15: The temperature T as a function of the complex conformal time: (i) The real part of $T(u)$ for an argument running in a domain containing several complex periods, (ii) Plot of the parametric region defined by $X = \text{Re}(T(u))$, $Y = \text{Re}(T(u))$, with $u = x + iy$ running in the same set as in (i).

that are experimentally measured — the temperature function, an elliptic function of u , goes to zero when the conformal time reaches a finite value u_f , corresponding, in terms of cosmic time, to $t \rightarrow \infty$. This happens, of course, in each period. In other words, Friedmann equations, from the very beginning, describe an infinite sequence of identical FLRW spacetimes, each one lasting an infinite number of years, in terms of cosmic time. This observation (that was explicitly made in ref. [2]) may inspire science-fiction writers and may trigger deep philosophical or metaphysical thoughts about the notion of reality or eternity, but it does not convey much physical meaning: a periodic function is a function on a circle, a bi-periodic function is a function on a torus, and each period of $T(u)$ describe the “same” universe. The fact of perceiving those universes as distinct does not mean much... unless, of course, one can introduce a way to make them (slightly) different, and a way to communicate —i.e., to send information— from one period to the next. In a slightly different framework, such a connection was proposed recently in [11] and popularized under the denomination “conformal cyclic cosmology”, the name “aeon” was then chosen to denote each member of the infinite sequence of spacetimes. Along that same line of thought, and taking into account the fact that all the functions of interest are bi-periodic, one could go one step further and talk about “complex aeons” when considering the doubly periodic lattice of identical (or almost identical) spacetimes obtained by allowing the conformal time to be complex: see fig. 15. We did not try, in the present article, to suggest any perturbation to the background provided by homogeneous and isotropic cosmologies, or to suggest a mechanism that could connect the different aeons that are rooted in the structure of Friedmann-Lemaître differential equations.

Appendix: a short memo on elliptic functions and elliptic curves

Elliptic functions.

Given two independent vectors of the plane, i.e., equivalently two non-zero complex numbers ω_1 and ω_2 such that ω_2/ω_1 is not real, one can build the lattice L generated by $2\omega_1$ and $2\omega_2$ as the infinite set

$\{m2\omega_1 + n2\omega_2\}$, where m, n are arbitrary integers. The two complex numbers $2\omega_1$ and $2\omega_2$ are called¹⁴ the *periods* of the lattice, and $\{\omega_1, \omega_2\}$ are the *half-periods*. An elliptic function with respect to L is a non-constant meromorphic function of the complex variable u that is periodic with respect to the given lattice, i.e., doubly periodic.

The Weierstrass elliptic function¹⁵ \mathcal{P} associated with the lattice L is defined as

$$\mathcal{P}(u) = \frac{1}{u^2} + \sum_{\omega \in L - \{0\}} \left(\frac{1}{(u - \omega)^2} - \frac{1}{\omega^2} \right) \quad (32)$$

From this definition it is clear that \mathcal{P} is periodic with respect to L (i.e., doubly periodic), that it is even, that it has a pole of order two at the origin (and at all other vertices of L), that it is meromorphic, and that

$$\mathcal{P}'(u)^2 = 4\mathcal{P}(u)^3 - g_2\mathcal{P}(u) - g_3 \quad (33)$$

where $g_2 = 60 \sum_{\omega \in L - \{0\}} \frac{1}{\omega^4}$, and $g_3 = 140 \sum_{\omega \in L - \{0\}} \frac{1}{\omega^6}$. These two (complex in general) numbers are called the Weierstrass invariants of the lattice L . Near the origin, $\mathcal{P}(u) \sim \frac{1}{u^2} + \frac{g_2}{20}u^2 + \frac{g_3}{28}u^4 + \dots$

The Weierstrass elliptic function \mathcal{P} is a function of the complex variable u , but it depends on the chosen lattice L . In order to make this dependence explicit one writes $\mathcal{P}(u; L)$ although it is quite standard to denote the same function by $\mathcal{P}(u; g_2, g_3)$ or $\mathcal{P}(u|\omega_1, \omega_2)$.

Modular considerations

A lattice L can be specified by a pair of half-periods (ω_1, ω_2) , but given any 2×2 matrix with integer coefficients a, b, c, d and determinant 1 (i.e., an element of the infinite discrete group $SL(2, \mathbb{Z})$), one obtains a new pair of periods for the same lattice L by setting $\begin{pmatrix} \omega'_1 \\ \omega'_2 \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix}$. The invariants g_2, g_3 only depend on the lattice: the invariants calculated from any two pairs of periods of the same lattice are the same. Conversely, given g_2 and g_3 , one can determine a pair of periods for the lattice L , but the previous comment shows that this pair is not uniquely determined.

Rather than choosing different pairs of periods for the same lattice, one can also rescale the lattice itself by an arbitrary (real or complex) non-zero number s , and in particular replace the pair of half-periods (ω_1, ω_2) by $(s\omega_1, s\omega_2)$. One gets another lattice, homothetic –by definition– with the first. The lattice is different, its Weierstrass invariants are different (one finds $g_2(s\omega_1, s\omega_2) = s^{-4}g_2(\omega_1, \omega_2)$ and $g_3(s\omega_1, s\omega_2) = s^{-6}g_3(\omega_1, \omega_2)$) and the corresponding \mathcal{P} -function is also different, but the former is simply related to the latter: one has the homogeneity relation: $\mathcal{P}(su|s\omega_1, s\omega_2) = s^{-2}\mathcal{P}(u|\omega_1, \omega_2)$. In particular, choosing $s = 1/\omega_1$, one obtains:

$$\mathcal{P}(u|\omega_1, \omega_2) = \frac{1}{\omega_1^2} \mathcal{P}\left(\frac{u}{\omega_1} \middle| 1, \tau = \frac{\omega_2}{\omega_1}\right) \quad (34)$$

Using an homothety with scale factor $1/\omega_1$, one can replace the basis $\{\omega_1, \omega_2\}$ by $\{1, \tau\}$ with $\tau = \omega_2/\omega_1$. It does not cost anything to assume that $\tau = \omega_2/\omega_1$ belongs to the upper half-plane (if it does not, just permute the two periods).

The value of $\tau = \omega_2/\omega_1$, called the modular parameter, clearly does not change if one only rescales the periods; however if one uses another set of periods $\{\omega'_1, \omega'_2\}$ for the same lattice (rescaled or not), one will

¹⁴About half the planet calls (ω_1, ω_2) what we call $(2\omega_1, 2\omega_2)$.

¹⁵Not to be confused with the pathological Weierstrass function that is continuous everywhere and differentiable nowhere.




Figure 16: Standard fundamental domain for the modular group, and its neighbors. Each point of the domain specifies a lattice in the complex plane, up to homothety. The boundary point $e^{2i\pi/3}$ describes a flat universe without radiation (equianharmonic elliptic curve).

have in general $\tau' \neq \tau$. As the periods are only defined up to the action of $SL(2, \mathbb{Z})$, the same is true for τ . In other words, one can find four integers a, b, c, d , with $ad - bc = 1$, such that $\tau' = (a\tau + b)/(c\tau + d)$. Notice that τ' also belongs to the upper half-plane. Conversely, one proves that if τ and τ' are related in this way, they are associated with the same lattice or with two homothetic lattices.

The j -invariant, or Klein invariant¹⁶, of a lattice is defined as the complex number

$$j(L) = \frac{g_2(L)^3}{\Delta} \quad \text{where} \quad \Delta = g_2(L)^3 - 27g_3(L)^2 \quad (35)$$

From the homogeneity transformations of g_2 and g_3 one sees that $j(L)$ is invariant under lattice rescaling. Actually, a detailed analysis leads to the following result: Given L and L' , two lattices in \mathbb{C} , then $j(L) = j(L')$ if and only if L and L' are homothetic. As a function of τ , the meromorphic function j is invariant with respect to the action of the modular group, and one has the following famous expansion (Fourier series), usually written in terms of $q = e^{2i\pi\tau}$, where all the coefficients are positive integers:

$$1728 j(\tau) = 1/q + 744 + 196884q + 21493760q^2 + \dots \quad (36)$$

An arbitrary complex number j being given, there are several ways to invert the Klein function, i.e., to find a value of τ (up to a modular transformation) solving the equation $j(\tau) = j$. This can be done for instance as follows: solve the cubic equation $256(1-x)^3/x^2 = 1728j$ in x , then write $x = \ell(1-\ell)$ and solve this quadratic equation in ℓ . One obtains finally $\tau = i \frac{{}_2F_1(1/2, 1/2, 1, 1-\ell)}{{}_2F_1(1/2, 1/2, 1, \ell)}$ where ${}_2F_1$ is the Gauss hypergeometric function. The first step of the method amounts to solve a sextic equation for ℓ and there are a priori six possible choices¹⁷ for the solution (six values for a cross-ratio), but τ , up to a modular transformation, will not depend upon this choice.

Elliptic curves

Choose a lattice L . If we set $x = \mathcal{P}(z)$ and $y = \mathcal{P}'(z)$, we have $y^2 = 4x^3 - g_2x - g_3$. Using homogenous coordinates $x = X/U$, $y = Y/U$, one gets the equation of a cubic in the complex projective plane: $Y^2U = 4X^3 - g_2XU^2 - g_3U^3$. The homothetic lattice sL determines an isomorphic cubic (use the homogeneity relations for $g_2, g_3, \mathcal{P}, \mathcal{P}'$, and set $x = \tilde{X}/U$, $y = \tilde{Y}/U$, with $\tilde{X} = X/s^3$ and $\tilde{Y} = Y/s^3$).

¹⁶It is convenient to set $J(\tau) = 1728j(\tau)$ but it is not uncommon to see J called j (or vice-versa) in the literature.

¹⁷The modular function ℓ , as a function of τ is usually called $\lambda(\tau)$ in the literature, but we changed the notation for $\ell(\tau)$, as λ already refers to the reduced cosmological constant.

An elliptic curve is a projective variety isomorphic to a non-singular curve of degree 3 in the complex projective plane, together with a distinguished point. Every such cubic can be brought to the Weierstrass form (i.e., to the previous form). This (complex) curve is parametrized by setting $x = \mathcal{P}(z)$ and $y = \mathcal{P}'(z)$, for some lattice specified by g_2 and g_3 .

A function (like \mathcal{P}) defined on the complex plane \mathbb{C} and periodic w.r.t. a lattice L is, by definition, a function on \mathbb{C}/L . The latter manifold is, topologically, and by construction, a torus (it is obtained by identifying the opposite sides of a period parallelogram). All 2-tori are diffeomorphic as real manifolds, but a torus constructed as previously is also a complex manifold because the choice of a lattice specifies a complex structure (two proportional lattices determine the same complex structure). Up to isomorphism, an elliptic curve can therefore also be defined as a 2-torus endowed with a complex structure.

Elliptic functions (continuation).

Taking into account equation (33), any elliptic function, for a given lattice, can be written as a rational function of \mathcal{P} and its first derivative \mathcal{P}' . In particular, any rational function of an elliptic function f is also elliptic (with respect to the same lattice) but its order will coincide with the order of f only if the transformation is fractional linear, like the transformations (20) or (24). It is known, since Liouville, that if a is an arbitrary complex number (including infinity), the number of solutions of the equation $f(u) = a$, called the *order* of f , is independent of a , if multiplicities are properly counted. The order of an elliptic function is at least two and the Weierstrass \mathcal{P} function of a lattice can be defined as *the* elliptic function of order 2 that has a pole of order 2 at the origin and is such that $1/u^2 - \mathcal{P}(u)$ vanishes at $u = 0$.

The Weierstrass ζ function is the primitive of $-\mathcal{P}$ which is such that $\zeta(u) - 1/u$ vanishes at the origin. The odd meromorphic function ζ has a pole of first order at all the vertices of the lattice defined by the invariants (g_2, g_3) . Warning: ζ is not periodic (hence not elliptic !) but it is quasi-periodic: if $a \neq b$ then $\zeta(u - a) - \zeta(u - b)$ is elliptic of order 2 with poles at $u = a$ and $u = b$.

The Weierstrass σ function is defined as an entire function that vanishes at $u = 0$ and whose logarithmic derivative is $\zeta(u)$. It is not elliptic, but if a_1, a_2, b_1, b_2 are complex numbers such that $a_1 + a_2 = b_1 + b_2$, then $\sigma(u - a_1)\sigma(u - a_2)/\sigma(u - b_1)\sigma(u - b_2)$ is elliptic of order 2 with poles at b_1, b_2 and zeroes at a_1, a_2 .

Acknowledgments

This text is based, in parts, on talks or lectures given by the author in Gordon Mac Kay Laboratory (Harvard Univ., 1981), CERN (Geneva, 2000), CBPF (Rio de Janeiro, 2002), LUTH (Meudon, 2013), CPT (Marseille, 2013) and IAFE (Univ. Buenos Aires, 2014). Hospitality of these research centers is greatly acknowledged.

References

- [1] M. Abramowitz and I.A. Stegun, eds. (1972), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, New York: Dover Publications, ISBN 978-0-486-61272-0.
- [2] R. Coquereaux and A. Grossmann, *Analytic Discussion of Spatially Closed Friedman Universes with Cosmological Constant and Radiation Pressure*, Ann. of Phys. **143**, No. **2** (1982), 296-356. Ann. of Phys. Erratum **170** (1986) 490.
- [3] R. Coquereaux and A. Grossmann, *Large scale geometry and evolution of a universe with radiation pressure and cosmological constant.*, CERN-TH/2000-180, arXiv:0101369
- [4] R. Coquereaux and A. Grossmann, *Evolução de um universo com pressão de radiação e constante cosmológica*, abridged version of [3] in portuguese, Anais IV Escola do CBPF, Rio de Janeiro, 15 a 26 de julho de 2002, Ligia MCS Rodrigues et al, editores, CBPF. Ao Livro Técnico, Rio de Janeiro (2003).
<http://www.cpt.univ-mrs.fr/%7Ecoque/FriedmannRio.pdf>
- [5] M. Dabrowski and J. Stelmach, *Analytic Solutions of Friedman Equation for Spatially Opened Universes with Cosmological, Constant and Radiation Pressure*, Ann. of Phys. 166 (1986), 422-442.
- [6] M. Eichler and D.Zagier, *On the zeros of the Weierstrass \mathcal{P} -function*, Math. Ann. 258, 399-407 (1982).
- [7] O. Lahav and A.R. Liddle, *The Cosmological Parameters 2014*, arXiv:1401.1389
- [8] G. Lemaître, *L'univers en expansion*, Ann. Soc. Sci. Bruxelles **A 53** (1933), 51.
- [9] C.W. Misner, K.S. Thorne, and J.A. Wheeler, *Gravitation*, Gravitation, San Francisco.
- [10] NIST, *Digital Library of Mathematical Functions*, <http://dlmf.nist.gov>
- [11] R. Penrose *Before the Big Bang: An Outrageous New Perspective and its Implications for Particle Physics*. Proceedings of the EPAC 2006, Edinburgh, Scotland: 27592762 (2006).
- [12] Planck Collaboration, *Planck 2013 results. XVI. Cosmological parameters* , arXiv:1303.5076
- [13] Wolfram Research, Inc., Mathematica, Version 10.0, Champaign, IL (2012).