Comprehensive Time-Series Regression Models Using Gretl—U.S. GDP and Government Consumption Expenditures & Gross Investment from 1980 to 2013¹

Juehui Shi Department of Operations Management and Strategy University at Buffalo, The State University of New York (SUNY)

Reference

Cottrell, A., and Lucchetti, R. (2018) "Gretl User's Guide," URL http://ricardo.ecn.wfu.edu/pub//gretl/manual/en/gretl-guide.pdf, Online retrieved 04/05/2018.

Abstract: Using Gretl, I apply ARMA, Vector ARMA, VAR, state-space model with a Kalman filter, transfer-function and intervention models, unit root tests, cointegration test, volatility models (ARCH, GARCH, ARCH-M, GARCH-M, Taylor-Schwert GARCH, GJR, TARCH, NARCH, APARCH, EGARCH) to analyze quarterly time series of GDP and Government Consumption Expenditures & Gross Investment (GCEGI) from 1980 to 2013. The article is organized in three sections: (I) Definition; (II) Regression Models; (III) Discussion [Summary of Major Findings and Their Managerial Implications, Comparison of Empirical Results, Contributions to Literature, Limitations and Future Research, Gretl Scripts]. Additionally, I discovered a unique interaction between GDP and GCEGI in both the short-run and the long-run and provided policy makers with some suggestions. For example in the short run, GDP responded positively and very significantly (0.00248) to GCEGI, while GCEGI reacted positively but not too significantly (0.08051) to GDP. In the long run, current GDP responded negatively and permanently (0.09229) to a shock in past GCEGI, while current GCEGI reacted negatively yet temporarily (0.29821) to a shock in past GDP. Therefore, policy makers should not adjust current GCEGI based merely on the condition of current and past GDP. Although increasing GCEGI does help GDP in the short-term, significantly abrupt increase in GCEGI might not be good to the long-term health of GDP. Instead, a balanced, sustainable, and economically viable solution is recommended, so that the short-term benefits to the current economy from increasing GCEGI often largely secured by the long-term loan outweigh or at least equal to the negative effect to the future economy from the long-term debt incurred by the loan. Finally, I found that non-normally distributed volatility models generally perform better than normally distributed ones. More specifically, TARCH-GED performs the best in the group of non-normally distributed, while GARCH-M does the best in the group of normally distributed.

¹Copy and Distribution of this paper is strictly prohibited without original author's approval. Cite the work appropriately. Work is completed under the supervision of my advisor Dr. Winston T. Lin.

SECTION I: DEFINITON

(1) Nonstationarity

Phenomena: In order to use statistical tests for an ARMA model selection, we need to estimate the underlying stochastic process, which can be derived by the mean, variance, and covariance of the sample data. However, these quantities are only meaningful if they are independent of time. If that is the case, the series is stationary. If not, then it is nonstationary. The most commonly considered is covariance stationarity with those following conditions: $E(y_t) = m$; $E[(y_t - m)^2] = \gamma_0 < \infty$; $E[(y_t - m)(y_{t-i} - m)] = \gamma_i$, where t=1,...,T and i=...,-2, -1, 0, 1, 2, ... The nonstationary series do not meet all or either one of the conditions.

Methodology: Univariate ARIMA for example ARIMA(1, 1, 1): $\Delta y_t = \mu + \varphi \Delta y_{t-1} + \varepsilon_t - \theta \varepsilon_{t-1}$, where the nonstationary time series y_t is differenced once to obtain stationary series $\Delta y_t(y_t - y_{t-1})$. Sometimes, it is necessary to differentiate the data more than once to obtain a stationary series. The process can be also denoted as $ARIMA(p, d, q) = \varphi_p(L)\Delta^d y_t = \mu + \theta_q(L)\varepsilon_t$, where d is the order of the differencing operator $\Delta^d = (1 - L)^d$. For example, if d=1, $\Delta_{y_t}^1 = (1 - L)^1 y_t = y_t - y_{t-1}$. $ARIMA(1, 1, 1) = \varphi_1(L)\Delta^1 y_t = \mu + \theta_1(L)\varepsilon_t$. Additionally, the most widely used test for non-stationarity is the Augmented Dickey-Fuller (ADF) unit root test. If the tested data has unit root, then the series is non-stationary.

Supply chain implication: Often, we observe some kind of trending behavior within wholesaler or retailer's sales data, for example the sales in the end of the sample are higher (lower) than the initial sales. Since the stationary data requires the mean level is independent of time (which implies that the average sales level at the beginning of the sample period is equal to the average sales level at the end of the sample), it is normal for supply chain time series to be nonstationary.

(2) Seasonality

Phenomena: Seasonality exists when there is a highly fluctuating pattern such as seasonal sales or advertising pattern in the retailing business. Seasonal processes can be identified from the ACF and PACF functions, similarly to the nonseasonal ARIMA processes, except that the patterns occur at lags s, 2s, 3s, ... instead of 1, 2, 3, ... In addition, seasonal unit root tests have been developed to detect the order of seasonal integration (Franses 1998 and Hylleberg 1992).

Methodology: Univariate ARIMA with (i) purely deterministic seasonal processes, (ii) stationary seasonal processes, and (iii) integrated seasonal processes (Maddala and Kim 1998, p. 363). For example (i) can be model as $y_t = \mu + \sum_{s=1}^{S-1} b_s d_{st} + \varepsilon_t$ where S is the maximum number of seasons (12 for monthly data, 4 for quarterly data), d_{st} is a dummy variable taking the value one in season s and zero otherwise, and b_s is a parameter. (ii) ARMA type if seasonal fluctuations in sales levels or random shocks die out over time in a seasonal way. (iii) integrated, when nonstationary seasonal patterns exist. It can be models as $ARIMA(P, D, Q)_s$, where s is the seasonal lag. For instance, $ARIMA(1, 0, 1)_{12}$: $y_t = \mu + \varphi y_{t-12} + \varepsilon_t - \theta \varepsilon_{t-12}$.

Supply chain implication: Retailer could have much higher advertising expenditures and sales figures in Spring than in Summer. In fact, many time series in supply chain display such seasonal patterns caused by managerial decisions, weather conditions, events, holidays, etc. Additionally, manufacturer could face similar fluctuation for raw material shortage due to adverse weather impact or unfavorable market condition, which also inevitably affect wholesaler's procurement and sales.

(3) Changing volatility

Phenomena: Volatility exists when the variance of the dependent variable changes over time. The volatility can also change over time. For instance the U.S. stock returns index (NASDAQ) experiences a relatively sedate period from 1992 to 1996. Then, stock returns become much more volatile until early 2004. Volatility increases again at the end of 2009.

Methodology: ARCH, GARCH, ARCH-M, EGARCH and other ARCH variants or VAR and impulse response functions.

Supply chain implication: Volatility widely exists in the supply chain where manufacturers require to procure precious metal (silver, gold, platinum) as raw materials whose price are usually volatile over time, due to uncertain supply and demand.

(4) Dynamics

Phenomena: Dynamics exists when the time series exhibit a deterministic trend which implies that the level is not constant, but can be perfectly predicted if the underlying deterministic function is known. The linear time trend is the most commonly used function for such purpose: $y_t = \mu + \beta t + \varepsilon_t$, t=1, ..., T, where the long run behavior of y_t is perfectly determined by the series' individual growth path βt . In the long run the series always returns to its individual growth path βt . Therefore, such series is often called Trend Stationary (TS), because it is stationary around a trend.

Methodology: We can incorporate the deterministic trend into ARMA model. For example: $dynamic AR(1): y_t = \mu + \beta t + \varphi y_{t-1} + \varepsilon_t, or ARMA(1, 1): y_t = \mu + \beta t + \varphi y_{t-1} - \theta \varepsilon_{t-1} + \varepsilon_t.$

Supply chain implication: Wholesaler's demand could be dynamic but predictable with a pattern, so is manufacturer.

(5) Randomness

Phenomena: Randomness exists when the time series exhibit a stochastic trend which implies that the variation is systematic but hardly predictable, because every temporary deviation may change the long-run performance of the series. Such phenomenon is also called "shock persistence". A simple example of such series is named as the Random Walk (RW) process: $y_t = \beta + y_{t-1} + \varepsilon_t$.

Methodology: ARMA with a stochastic trend: $y_t = \beta + \sum_{i=1}^p \varphi_p y_{t-p} + \sum_{i=1}^q \theta_q \varepsilon_{t-q} + \varepsilon_t$. A simple case of AR(1): $y_t = \beta + \varphi_1 y_{t-1} + \varepsilon_t$, or ARMA(1, 1): $y_t = \beta + \varphi_1 y_{t-1} - \theta_1 \varepsilon_{t-1} + \varepsilon_t$.

Supply chain implication: Retailer's demand is unpredictable, but rather stochastic due to the nature of retailing business.

(4) + (5) Dynamics + Randomness

Phenomena: A deterministic trend and a stochastic trend can coexist in a very complex situation, for example: $y_t = \mu + \beta t + \sum_{i=1}^t \varepsilon_{t-i} + \varepsilon_t$, where $\mu + \beta t$ is the deterministic trend, $\sum_{i=1}^t \varepsilon_{t-i}$ is the stochastic trend, and ε_t is the noise term.

Methodology: ARMA with a dynamic and stochastic trend: $y_t = \mu + \beta t + \sum_{i=1}^p \varphi_p y_{t-p} + \sum_{i=1}^q \theta_q \varepsilon_{t-q} + \varepsilon_t$. A simple case of AR(1): $y_t = \mu + \beta t + \varphi_1 y_{t-1} + \varepsilon_t$, or ARMA(1, 1): $y_t = \mu + \beta t + \varphi_1 y_{t-1} - \theta_1 \varepsilon_{t-1} + \varepsilon_t$.

Supply chain implication: Consumer's demand is changing all the time, sometime predictable but other time not, therefore the combination of both dynamics and stochastic works better.

(6) Nonnormality

Phenomena: Nonnormality exists if residuals of the time series are not normally distributed. We have to utilize GARCH or its variants integrating nonnormal distributions such as student's t or generalized error to explain the endogenous variables.

Methodology: ARCH, ARCH-M, GARCH, GARCH-M, EGARCH and other GARCH variants with nonnormal residual distributions such as t, GED, skewed t, skewed-GED.

Supply chain implication: Error distributions within time series, especially of volatile data, are nonnormal. Sales or price series within the supply chain can often be quite heteroscedastic with high volatility over time.

(7) Nonlinearity

Phenomena: Sometimes, the observed series are not only influenced by its own lagged past value but also by other determinant exogenous variables. For example, one of the main fields of interest in marketing is the determination of the effect of marketing actions (advertising campaign, pricing) on sales fluctuation.

Methodology: ARMAX, transfer function and intervention model.

Supply chain implication: Since retailer side of the supply chain frequently involves marketing decisions such as pricing and advertising expenditures, sales time series can be effectively modeled as ARMAX by incorporating pricing and advertising as exogenous determining variables. Such model provides the researchers with important insights of how pricing and advertising decisions contribute to demand changes within the supply chain. Moreover, pricing also heavily influences demand on the side of both manufacturer and wholesaler. We can even further study the joint cause and effect of pricing-demand from downstream (manufacturer→wholesaler→retailer) to upstream (retailer→wholesaler→manufacturer).

SECTION II: REGRESSION MODELS

Data source: GDP¹, Government Consumption Expenditures & Gross Investment²

Annotations: 1 = GDPC1Q, 2 = GCEC1Q

(1) The Box-Jenkins Univariate ARIMA Approach:

Note: Maximum lag for the correlogram (ACF and PACF) is set to 20. I denote observations GDPC1Q as Y_{1t} and GCEC1Q as Y_{2t} .

Data collected:

- a) Use for model construction (Sample Range): 1980 Q1-2006 Q1, Billions of Chained 2005 Dollars, Quarterly, Seasonally Adjusted Annual Rate.
- b) Use for forecast performance assessment (Forecast Range): 2006 Q2-2013 Q1


Sample size for each observation: 105.

I. Identification

In this section, I draw the graphs of ACF and PACF for all the observed variables, including GDPC1Q and GCEC1Q, which help me to initially identify fitted models.

LAG	ACF		PACF		Q-stat.	[p-value]
1 2 3 4 5	0.9725 0.9445 0.9157 0.8873 0.8591	* * * * * * * * * * * *	0.9725 -0.0250 -0.0280 -0.0062 -0.0123	* * *	102.1765 199.4772 291.8301 379.4130 462.3396	[0.000] [0.000] [0.000] [0.000]
6 7 8 9 10	0.8308 0.8027 0.7739 0.7443 0.7145	* * * * * * * * * * * *	-0.0187 -0.0103 -0.0280 -0.0319 -0.0187		540.6633 614.5253 683.8957 748.7268 809.1079	[0.000] [0.000] [0.000] [0.000] [0.000]
11 12 13 14 15 16 17 18	0.6846 0.6555 0.6268 0.5986 0.5702 0.5423 0.5148 0.4880	* * * * * * * * * * * * * * * * * * *	-0.0205 -0.0032 -0.0089 -0.0109 -0.0194 -0.0093 -0.0108 -0.0108		865.1257 917.0296 965.0109 1009.2447 1049.8350 1086.9642 1120.8026 1151.5546	[0.000] [0.000] [0.000] [0.000] [0.000] [0.000] [0.000]
19 20	0.4610 0.4332	* * *	-0.0213 -0.0352		1179.3186 1204.1175	[0.000] [0.000]


Autocorrelation function for GDPC1Q



The ACF graph for GDPC1Q dies out slowly (exponentially decaying), with one spike in PACF that cuts off after lag 1. Data is nonstationary. Therefore, the first model for GDPC1Q (Y_{1t}) is initially identified as ARIM(1, 1, 0), ARIMA(0, 1, 1), or ARIMA (1, 1, 1).

Autocorrelation function for GCEC1Q

LAG	ACF		PACF		Q-stat.	[p-value]
1	0.9697	* * *	0.9697	* * *	101.5772	[0.000]
2	0.9399	* * *	-0.0063		197.9379	[0.000]
3	0.9071	* * *	-0.0651		288.5790	[0.000]
4	0.8738	* * *	-0.0281		373.5108	[0.000]
5	0.8400	* * *	-0.0229		452.7871	[0.000]
6	0.8053	* * *	-0.0339		526.3749	[0.000]
7	0.7681	* * *	-0.0600		594.0054	[0.000]
8	0.7310	* * *	-0.0185		655.8961	[0.000]
9	0.6931	* * *	-0.0308		712.1199	[0.000]
10	0.6549	* * *	-0.0286		762.8375	[0.000]
11	0.6161	* * *	-0.0309		808.2024	[0.000]
12	0.5781	* * *	-0.0083		848.5792	[0.000]
13	0.5415	* * *	0.0006		884.3816	[0.000]
14	0.5043	* * *	-0.0325		915.7805	[0.000]
15	0.4682	* * *	-0.0080		943.1473	[0.000]
16	0.4308	* * *	-0.0461		966.5724	[0.000]
17	0.3946	* * *	-0.0062		986.4527	[0.000]
18	0.3611	* * *	0.0199		1003.2914	[0.000]
19	0.3294	* * *	0.0052		1017.4698	[0.000]
20	0.2986	* * *	-0.0133		1029.2550	[0.000]



The ACF graph for GCEC1Q dies out slowly (exponentially decaying), with one spike in PACF that cuts off after lag 1. Data is nonstationary. Therefore, the second model for GCEC1Q (Y_{2t}) is initially identified as ARIMA(1, 1, 0), ARIMA(0, 1, 1), or ARIMA (1, 1, 1).

II & III. Estimation and Diagnostic Checking

In this section, I fit the following ARIMA models into collected data (GDPC1Q and GCEC1Q), for example ARIMA (1, 0, 0); (1, 1, 0); (0, 0, 1); (0, 1, 1); (1, 0, 1); (1, 1, 1). I will rule out any ARIMA model, which either does not have significant parameters or pass the initial normality test with significantly low p-value that rejects the null of normal distribution. Additionally, the model must meet those calculation requirements, such as convergence criterion and finite initial value in the objective function. Finally, I might need to increase the AR or MA order to 2 if necessary to obtain normally-distributed residual.

If the tested model meets the preliminary requirements for significant parameters and residual normality distribution with successful calculation, the estimated models are suitable to be tested further against autocorrelation and heteroskedasticity.

For diagnostic checking, I will perform additional tests on all the models passing the normality test, for instance Ljung-Box Q test for autocorrelation and ARCH test for heteroskedasticity. The selection technique for the best fitted model is to have the best overall white noise, which is normally distributed, independently distributed (NO autocorrelation), and homoskedastic (NO ARCH effect).

	Normality (χ^2)	Autocorrelation (Ljung-Box	ARCH Effect (LM)
	Test	Q) Test (lag order 4)	Test (lag order 4)
Model 1			
GDPC1Q (Y_{1t})			
ARIMA (1, 1, 0)	6.406 (0.04063)	9.64775 (0.02181)	11.4214 (0.0222148)
ARIMA (0, 0, 1)	3.548 (0.16963)	343.18 (4.471e-074)	99.6012 (1.19602e-
			020)
ARIMA (0, 1, 1)	6.943 (0.03107)	14.2602 (0.002572)	8.3121 (0.0807918)
ARIMA (1, 1, 1)	4.976 (0.08308)	3.64901 (0.1613)	12.384 (0.0147128)
ARIMA (2, 1, 0)	5.343 (0.06915)	1.05835 (0.5891)	15.8064 (0.00329019)
ARIMA (0, 1, 2)	7.162 (0.02784)	2.50263 (0.2861)	13.6701 (0.00842606)
Model 2			
GCEC1Q (Y_{2t})			
ARIMA (0, 0, 1)	0.791 (0.67336)	317.396 (1.708e-068)	97.9932 (2.63013e-
			020)
ARIMA (2, 1, 1)	0.491 (0.78234)	3.69189 (0.05468)	3.0714 (0.545949)
ARIMA (1, 1, 2)	0.411 (0.81410)	2.13669 (0.1438)	2.68573 (0.611714)

Below is the summarized testing results:

P value for the testing statistics is in the parenthesis. 0.02 is used as significance level for all the hypothesis testing.

Null hypothesis for ARCH Effect (LM) test is that no ARCH effect is present.

Feasible models are highlighted in red, which need to be further screened for the best appropriate model.

Ideally, studied models should pass all three tests. But ARCH model can be used to correct the heteroskedasticity if the model has passed both normality and autocorrelation tests. ARIMA (1, 1, 0) is appropriate for model 1, because it has passed all the tests. The reason for not selecting ARIMA (2, 1, 0) or ARIMA (0, 1, 2) is that they did not pass the ARCH effect test, which also have more parameters than ARIMA (1, 1, 0). However, ARIMA (1, 1, 1) would be appropriate if I choose 0.01 as significance level for all the hypothesis testing. Therefore, I will estimate both ARIMA (1, 1, 0) and ARIMA (1, 1, 1) model, then decide which one I will utilize as model 1 based on their forecasting performance and overall the goodness of fit.

Additionally, I choose ARIMA (1, 1, 2) for model 2, because it has passed all the tests with the lowest test statistics (highest p value) for each test.

Below are the estimated results of ARIMA (1, 1, 0)/ARIMA (1, 1, 1) for model 1 GDPC1Q (Y_{1t}) and ARIMA (1, 1, 2) for model 2 GCEC1Q (Y_{2t}) .

Model 1.1 GDPC1Q (Y_{1t}): ARIMA (1, 1, 0), using observations 1980:1-2006:1 (T = 105)						
Dependent variable: (1-L) GDPC1Q						
	Standard e	errors bas	ed on l	Hessian		
	Coefficient	Std. Err	or	Z	p-value	
const	66.9945	7.9377	'8	8.4400	< 0.00001	***
phi_1	0.327894	0.09317	'87	3.5190	0.00043	***
Mean dependent var	66.78	3000	SD de	ependent var	58	36696
Mean of innovations				f innovations		91789
Log-likelihood	-569.6	585	Akaike	e criterion	114	45.317
Schwarz criterion	1153	.279	Hanna	n-Quinn	114	48.543
	Real	Imagin	ary	Modulus	Frequency	
AR						
Root 1	3.0498	0.	0000	3.0498	0.000	0

For Model 1.1:

$$\Delta^{1d} Y_{1t} = c_1 + \varphi_1 \Delta^{1d} Y_{1,t-1} + \varepsilon_{1t}$$
(1.1)

Where c_1 is a constant, ε_{1t} is white noise

$$\Delta^{1d}Y_{1t} = Y_{1t} - Y_{1,t-1} \tag{1.2}$$


Combine (1.1) and (1.2), we have: $Y_{1t} - Y_{1,t-1} = c_1 + \varphi_1 (Y_{1,t-1} - Y_{1,t-2}) + \varepsilon_{1t}$

Replace the symbol with estimated parameters, we have:

$$Y_{1t} - Y_{1,t-1} = 66.9945 + 0.327894 (Y_{1,t-1} - Y_{1,t-2}) + \varepsilon_{1t}$$
(1.3)

$$\therefore Y_{1t} = 66.9945 + 1.327894Y_{1,t-1} - 0.327894Y_{1,t-2} + \varepsilon_{1t}$$
(1.4)

More simply, let first difference of GDPC1Q be d_GDPC1Q; $Y_{d_{GDPC1Q}}$ be $\Delta^{1d}Y_1$, then:



Actual and fitted GDPC1Q fitted actual

Model 1.1 estimation range: 1980:1 - 2006:1 Standard error of residuals = 54.9179

	GDPC1Q	fitted	residual	
1980:1	5903.40	5951.49	-48.0923	
1980:2	5782.40	5954.62	-172.223	*
1980:3	5771.70	5787.75	-16.0508	
1980:4	5878.40	5813.22	65.1825	
1981:1	6000.60	5958.41	42.1878	
1981:2	5952.70	6085.69	-132.995	
1981:3	6025.00	5982.02	42.9802	
1981:4	5950.00	6093.73	-143.733	*
1982:1	5852.30	5970.43	-118.134	
1982:2	5884.00	5865.29	18.7093	
1982:3	5861.40	5939.42	-78.0202	
1982:4	5866.00	5899.02	-33.0155	
1983:1	5938.90	5912.53	26.3657	
1983:2	6072.40	6007.83	64.5706	
1983:3	6192.20	6161.20	31.0002	
1983:4	6320.20	6276.51	43.6924	

1984:1	6442.80	6407.20	35.6036
1984:2	6554.00	6528.03	25.9743
1984:3	6617.70	6635.49	-17.7877
1984:4	6671.60	6683.61	-12.0128
1985:1	6734.50	6734.30	0.200579
1985:2	6791.50	6800.15	-8.65047
1985:3	6897.60	6855.22	42.3841
1985:4	6950.00	6977.42	-27.4155
1986:1	7016.80	7012.21	4.59242
1986:2	7045.00	7083.73	-38.7293
1986:3	7112.90	7099.27	13.6275
1986:4	7147.30	7180.19	-32.8899
1987:1	7186.90	7203.61	-16.7055
1987:2	7263.30	7244.91	18.3895
1987:3	7326.30	7333.38	-7.07703
1987:4	7451.70	7391.98	59.7167
1988:1	7490.20	7537.84	-47.6438
1988:2	7586.40	7547.85	38.5501
1988:3	7625.60	7662.97	-37.3693
1988:4	7727.40	7683.48	43.9206
1989:1	7799.90	7805.81	-5.90554
1989:2	7858.30	7868.70	-10.3982
1989:3	7920.60	7922.47	-1.87494
1989:4	7937.90	7986.05	-48.1537
1990:1	8020.80	7988.60	32.2015
1990:2	8052.70	8093.01	-40.3083
1990:3	8052.60	8108.19	-55.5858
1990:4	7982.00	8097.59	-115.593
1991:1	7943.40	8003.88	-60.4766
1991:2	7997.00	7975.77	21.2308
1991:3	8030.70	8059.60	-28.9011
1991:4	8062.20	8086.78	-24.5760
1992:1	8150.70	8117.55	33.1454
1992:2	8237.30	8224.74	12.5555
1992:3	8322.30	8310.72	11.5785
1992:4	8409.80	8395.20	14.6031
1993:1	8425.30	8483.52	-58.2167
1993:2	8479.20	8475.41	3.79170
1993:3	8523.80	8541.90	-18.0994
1993:4	8636.40	8583.45	52.9500
1994:1	8720.50	8718.35	2.15321
1994:2	8839.80	8793.10	46.6982
1994:3	8896.70	8923.94	-27.2437
1994:4	8995.50	8960.38	35.1169
1995:1	9017.60	9072.92	-55.3219
1995:2	9037.00	9069.87	-32.8724

1995:3	9112.90	9088.39	24.5129	
1995:4	9176.40	9182.81	-6.41309	
1996:1	9239.30	9242.25	-2.94720	
1996:2	9399.00	9304.95	94.0495	
1996:3	9480.80	9496.39	-15.5906	
1996:4	9584.30	9552.65	31.6523	
1997:1	9658.00	9663.26	-5.26296	
1997:2	9801.20	9727.19	74.0083	
1997:3	9924.20	9893.18	31.0197	
1997:4	10000.3	10009.6	-9.25688	
1998:1	10094.8	10070.3	24.5213	
1998:2	10185.6	10170.8	14.7881	
1998:3	10320.0	10260.4	59.6013	
1998:4	10498.6	10409.1	89.5051	
1999:1	10592.1	10602.2	-10.0878	
1999:2	10674.9	10667.8	7.11598	
1999:3	10810.7	10747.1	63.6244	
1999:4	11004.8	10900.3	104.546	
2000:1	11033.6	11113.5	-79.8701	
2000:2	11248.8	11088.1	160.731	*
2000:3	11258.3	11364.4	-106.089	
2000:4	11325.0	11306.4	18.5591	
2001:1	11287.8	11391.9	-104.096	
2001:2	11361.7	11320.6	41.0717	
2001:3	11330.4	11431.0	-100.557	
2001:4	11370.0	11365.2	4.83713	
2002:1	11467.1	11428.0	39.0895	
2002:2	11528.1	11544.0	-15.8644	
2002:3	11586.6	11593.1	-6.52747	
2002:4	11590.6	11650.8	-60.2077	
2003:1	11638.9	11636.9	1.96248	
2003:2	11737.5	11699.8	37.7368	
2003:3	11930.7	11814.9	115.844	
2003:4	12038.6	12039.1	-0.475032	
2004:1	12117.9	12119.0	-1.10569	
2004:2	12195.9	12188.9	6.97208	
2004:3	12286.7	12266.5	20.1983	
2004:4	12387.2	12361.5	25.7013	
2005:1	12515.0	12465.2	49.8207	
2005:2	12570.7	12601.9	-31.2308	
2005:3	12670.5	12634.0	36.5104	
2005:4	12735.6	12748.2	-12.6497	
2006:1	12896.4	12802.0	94.4282	

Note: * denotes a residual in excess of 2.5 standard errors

Log-likelihood

1142.394

Model 1.2 GDPC1Q (Y_{1t}): ARIMA (1, 1, 1), using observations 1980:1-2006:1 (T = 105) Dependent variable: (1-L) GDPC1Q Standard errors based on Hessian Std. Error Coefficient *p*-value Z*** Const 66.2629 11.1287 5.9542 < 0.00001 *** phi 1 0.759918 0.139983 5.4286 < 0.00001 theta 1 -0.479305 0.180154 -2.6605 0.00780 *** Mean dependent var 66.78000 S.D. dependent var 58.36696 Mean of innovations 0.894814 S.D. of innovations 53.61904

Schwarz criterion	1153.	010 Hanna	n-Quinn	1146.695	
	Real	Imaginary	Modulus	Frequency	
AR Root 1	1.3159	0.0000	1.3159	0.0000	
MA Root 1	2.0864	0.0000	2.0864	0.0000	

-567.1968

For Model 1.2:

$$\Delta^{1d} Y_{1t} = c_1 + \varphi_1 \Delta^{1d} Y_{1,t-1} + \varepsilon_{1t} + \theta_1 \varepsilon_{1,t-1}$$
(1.6)

Akaike criterion

Where c_1 is a constant, ε_{1t} is white noise

$$\Delta^{1d} Y_{1t} = Y_{1t} - Y_{1,t-1} \tag{1.7}$$

Combine (1.6) and (1.7), we have: $Y_{1t} - Y_{1,t-1} = c_1 + \varphi_1 (Y_{1,t-1} - Y_{1,t-2}) + \varepsilon_{1t} + \theta_1 \varepsilon_{1,t-1}$


Replace the symbol with estimated parameters, we have:

$$Y_{1t} - Y_{1,t-1} = 66.2629 + 0.759918 (Y_{1,t-1} - Y_{1,t-2}) + \varepsilon_{1t} - 0.479305\varepsilon_{1,t-1}$$
(1.8)

$$\therefore Y_{1t} = 66.2629 + 1.759918Y_{1,t-1} - 0.759918Y_{1,t-2} + \varepsilon_{1t} - 0.479305\varepsilon_{1,t-1}$$
(1.9)

More simply, let first difference of GDPC1Q be d_GDPC1Q; $Y_{d_{GDPC1Q}}$ be $\Delta^{1d}Y_1$, then:

$$\Delta^{1d}Y_{1t} = 66.2629 + 0.759918\Delta^{1d}Y_{1,t-1} + \varepsilon_{1t} - 0.479305\varepsilon_{1,t-1}$$
(1.10)



lag

Regression residuals (= observed - fitted GDPC1Q)

Actual and fitted GDPC1Q fitted actual -

Model 1.2 estimation range: 1980:1 - 2006:1 Standard error of residuals = 53.619

	GDPC1Q	fitted	residual	
1980:1	5903.40	5950.76	-47.3624	
1980:2	5782.40	5952.81	-170.406	*
1980:3	5771.70	5785.19	-13.4899	
1980:4	5878.40	5785.89	92.5081	
1981:1	6000.60	5931.13	69.4672	
1981:2	5952.70	6076.09	-123.388	
1981:3	6025.00	5991.34	33.6567	
1981:4	5950.00	6079.72	-129.719	
1982:1	5852.30	5971.09	-118.789	
1982:2	5884.00	5850.90	33.0992	
1982:3	5861.40	5908.13	-46.7333	
1982:4	5866.00	5882.53	-16.5338	
1983:1	5938.90	5893.33	45.5711	
1983:2	6072.40	5988.36	84.0360	
1983:3	6192.20	6149.48	42.7214	
1983:4	6320.20	6278.67	41.5300	

1984:1	6442.80	6413.47	29.3277
1984:2	6554.00	6537.82	16.1826
1984:3	6617.70	6646.65	-28.9549
1984:4	6671.60	6695.89	-24.2934
1985:1	6734.50	6740.11	-5.61192
1985:2	6791.50	6800.90	-9.39708
1985:3	6897.60	6855.23	42.3722
1985:4	6950.00	6973.83	-23.8265
1986:1	7016.80	7017.15	-0.348276
1986:2	7045.00	7083.64	-38.6379
1986:3	7112.90	7100.86	12.0426
1986:4	7147.30	7174.63	-27.3348
1987:1	7186.90	7202.45	-15.5513
1987:2	7263.30	7240.36	22.9450
1987:3	7326.30	7326.27	0.0314818
1987:4	7451.70	7390.07	61.6318
1988:1	7490.20	7533.36	-43.1617
1988:2	7586.40	7556.05	30.3471
1988:3	7625.60	7660.87	-35.2670
1988:4	7727.40	7688.20	39.1992
1989:1	7799.90	7801.88	-1.97971
1989:2	7858.30	7871.85	-13.5513
1989:3	7920.60	7925.08	-4.48285
1989:4	7937.90	7986.00	-48.1000
1990:1	8020.80	7990.01	30.7904
1990:2	8052.70	8084.95	-32.2476
1990:3	8052.60	8108.31	-55.7063
1990:4	7982.00	8095.13	-113.133
1991:1	7943.40	7998.48	-55.0834
1991:2	7997.00	7956.38	40.6226
1991:3	8030.70	8034.17	-3.46948
1991:4	8062.20	8073.88	-11.6806
1992:1	8150.70	8107.64	43.0555
1992:2	8237.30	8213.22	24.0756
1992:3	8322.30	8307.48	14.8222
1992:4	8409.80	8395.70	14.1029
1993:1	8425.30	8485.44	-60.1416
1993:2	8479.20	8481.81	-2.61333
1993:3	8523.80	8537.32	-13.5206
1993:4	8636.40	8580.08	56.3187
1994:1	8720.50	8710.88	9.61867
1994:2	8839.80	8795.71	44.0928
1994:3	8896.70	8925.23	-28.5327
1994:4	8995.50	8969.52	25.9764
1995:1	9017.60	9074.04	-56.4377
1995:2	9037.00	9077.35	-40.3535
=			

1995:3	9112.90	9086.99	25.9075	
1995:4	9176.40	9174.07	2.33140	
1996:1	9239.30	9239.45	-0.145765	
1996:2	9399.00	9303.08	95.9229	
1996:3	9480.80	9490.29	-9.49098	
1996:4	9584.30	9563.42	20.8813	
1997:1	9658.00	9668.85	-10.8514	
1997:2	9801.20	9735.12	66.0845	
1997:3	9924.20	9894.25	29.9460	
1997:4	10000.3	10019.2	-18.9250	
1998:1	10094.8	10083.1	11.6910	
1998:2	10185.6	10176.9	8.68292	
1998:3	10320.0	10266.3	53.6528	
1998:4	10498.6	10412.3	86.2747	
1999:1	10592.1	10608.9	-16.7778	
1999:2	10674.9	10687.1	-12.2024	
1999:3	10810.7	10759.6	51.1218	
1999:4	11004.8	10905.3	99.4977	
2000:1	11033.6	11120.5	-86.9187	
2000:2	11248.8	11113.1	135.745	*
2000:3	11258.3	11363.2	-104.879	
2000:4	11325.0	11331.7	-6.69674	
2001:1	11287.8	11394.8	-107.005	
2001:2	11361.7	11326.7	34.9726	
2001:3	11330.4	11417.0	-86.6038	
2001:4	11370.0	11364.0	5.96732	
2002:1	11467.1	11413.1	53.9589	
2002:2	11528.1	11530.9	-2.83368	
2002:3	11586.6	11591.7	-5.12162	
2002:4	11590.6	11649.4	-58.8184	
2003:1	11638.9	11637.7	1.15991	
2003:2	11737.5	11691.0	46.5435	
2003:3	11930.7	11806.0	124.672	
2003:4	12038.6	12033.7	4.93147	
2004:1	12117.9	12134.1	-16.2398	
2004:2	12195.9	12201.9	-5.95371	
2004:3	12286.7	12273.9	12.7644	
2004:4	12387.2	12365.5	21.7091	
2005:1	12515.0	12469.1	45.9251	
2005:2	12570.7	12606.0	-35.3138	
2005:3	12670.5	12645.9	24.6381	
2005:4	12735.6	12750.4	-14.8390	
2006:1	12896.4	12808.1	88.3085	

Note: * denotes a residual in excess of 2.5 standard errors

Model 1.3 GCEC1Q (*Y*_{2t}): ARIMA (1, 1, 2), using observations 1980:1-2006:1 (T = 105)

Dependent variable: (1-L) GCEC1Q Standard errors based on Hessian								
	Coefficient Std. Error z p-value							
const	9.7675	2.54067	3.8445	0.00012	***			
phi 1	0.811342	0.12216	6.6416	< 0.00001	***			
theta 1	-0.943283	0.141859	-6.6494	< 0.00001	***			
theta_2	0.273345	0.11415	2.3946	0.01664	**			
Mean dependent var	10.03		dependent var		92138			
Mean of innovations	0.127	798 S.D.	S.D. of innovations		15.25431			
Log-likelihood	-435.2		Akaike criterion).4622			
Schwarz criterion	893.7320 Hanna		nan-Quinn	885	5.8394			
	Real	Imaginary	Modulus	Frequency				
AR								
Root 1	1.2325	0.000) 1.2325	0.000	0			
MA								
Root 1	1.7254	-0.8254	1.9127	-0.071	0			
Root 2	1.7254	0.8254	1.9127	0.071	0			
Model 1 2.								

For Model 1.3:

$$\Delta^{1d} Y_{2t} = c_2 + \varphi_1 \Delta^{1d} Y_{2,t-1} + \varepsilon_{2t} + \theta_1 \varepsilon_{2,t-1} + \theta_2 \varepsilon_{2,t-2}$$
(1.11)

Where c_2 is a constant, ε_{2t} is white noise

$$\Delta^{1d} Y_{2t} = Y_{2t} - Y_{2,t-1} \tag{1.12}$$

Combine (1.11) and (1.12): $Y_{2t} - Y_{2,t-1} = c_2 + \varphi_1 (Y_{2,t-1} - Y_{2,t-2}) + \varepsilon_{2t} + \theta_1 \varepsilon_{2,t-1} + \theta_2 \varepsilon_{2,t-2}$


Replace the symbol with estimated parameters, we have:

$$Y_{2t} - Y_{2,t-1} = 9.7675 + 0.811342 (Y_{2,t-1} - Y_{2,t-2}) + \varepsilon_{2t} - 0.943283\varepsilon_{2,t-1}$$
(1.13)
+ 0.273345\varepsilon_{2,t-2}


$$Y_{2t} = 9.7675 + 1.811342Y_{2,t-1} - 0.811342Y_{2,t-2} + \varepsilon_{2t} - 0.943283\varepsilon_{2,t-1}$$
(1.14)
+ 0.273345\varepsilon_{2,t-2}

More simply, let first difference of GCEC1Q be d_GCEC1Q; $Y_{d_{GCEC1Q}}$ be $\Delta^{1d}Y_2$, then:


$$\Delta^{1d} Y_{2t} = 9.7675 + 0.811342 \Delta^{1d} Y_{2,t-1} + \varepsilon_{2t} - 0.943283 \varepsilon_{2,t-1}$$
(1.15)
+ 0.273345 \varepsilon_{2,t-2}



Regression residuals (= observed - fitted GCEC1Q)



Actual and fitted GCEC1Q



Model 1.3 estimation range: 1980:1 - 2006:1 Standard error of residuals = 15.2543

	GCEC1Q	fitted	residual
1980:1	1365.40	1353.57	11.8325
1980:2	1369.70	1374.22	-4.51706
1980:3	1350.80	1381.94	-31.1393
1980:4	1349.40	1364.51	-15.1100
1981:1	1367.30	1356.14	11.1595
1981:2	1370.40	1369.12	1.27825
1981:3	1367.30	1376.59	-9.28578
1981:4	1379.90	1375.74	4.16444
1982:1	1378.50	1385.50	-6.99946
1982:2	1386.50	1386.95	-0.447576
1982:3	1396.00	1393.34	2.65761
1982:4	1420.10	1402.92	17.1788
1983:1	1430.80	1426.02	4.78191
1983:2	1443.00	1441.51	1.49089

1983:3	1468.00	1454.64	13.3581
1983:4	1443.20	1477.93	-34.7333
1984:1	1457.80	1461.34	-3.53614
1984:2	1489.20	1465.33	23.8703
1984:3	1500.20	1493.04	7.16417
1984:4	1532.30	1510.73	21.5655
1985:1	1549.90	1541.80	8.09730
1985:2	1584.70	1564.28	20.4209
1985:3	1625.80	1597.73	28.0719
1985:4	1635.50	1640.09	-4.59111
1986:1	1653.20	1657.22	-4.01676
1986:2	1688.30	1671.94	16.3625
1986:3	1726.60	1702.09	24.5116
1986:4	1716.60	1740.87	-24.2683
1987:1	1723.70	1739.92	-16.2213
1987:2	1734.60	1739.97	-5.37092
1987:3	1734.60	1745.92	-11.3186
1987:4	1755.60	1745.65	9.94874
1988:1	1747.10	1762.00	-14.9025
1988:2	1751.70	1758.82	-7.12305
1988:3	1750.70	1759.92	-9.22040
1988:4	1786.20	1758.48	27.7182
1989:1	1775.20	1788.18	-12.9789
1989:2	1802.80	1787.94	14.8627
1989:3	1819.70	1809.47	10.2316
1989:4	1829.40	1829.67	-0.265702
1990:1	1857.60	1842.16	15.4399
1990:2	1860.40	1867.69	-7.28578
1990:2	1859.80	1875.61	-15.8074
1990:4	1878.30	1874.08	4.22474
1991:1	1885.90	1886.85	-0.946534
1991:2	1892.50	1895.96	-3.45658
1991:2	1883.50	1902.70	-19.1994
1991:4	1875.60	1895.21	-19.6062
1992:1	1889.90	1884.28	5.62072
1992:2	1887.60	1892.68	-5.08371
1992:2	1897.30	1893.91	3.39160
1992:4	1897.90	1902.42	-4.52389
1993:1	1877.90	1905.42	-27.5239
1993:2	1876.50	1888.24	-11.7421
1993:3	1874.60	1880.76	-6.15944
1993:4	1883.90	1877.50	6.39839
1994:1	1859.90	1885.57	-25.6690
1994.1	1867.70	1868.23	-0.532646
1994.2	1900.50	1869.36	31.1429
1994:3	1884.10	1809.30	-15.3326
1774.4	1004.10	10/2.40	-15.5520

1995:1	1891.60	1895.61	-4.01243	
1995:2	1897.90	1899.12	-1.22155	
1995:3	1893.70	1904.91	-11.2097	
1995:4	1872.50	1902.38	-29.8750	
1996:1	1884.50	1882.26	2.24122	
1996:2	1911.60	1885.80	25.8015	
1996:3	1909.70	1911.70	-2.00461	
1996:4	1925.90	1918.94	6.95521	
1997:1	1929.40	1933.78	-4.37778	
1997:2	1946.00	1940.11	5.88693	
1997:3	1948.20	1954.56	-6.36131	
1997:4	1951.50	1959.44	-7.93735	
1998:1	1939.70	1961.77	-22.0685	
1998:2	1981.90	1950.62	31.2839	
1998:3	2000.20	1982.44	17.7606	
1998:4	2018.10	2008.69	9.41165	
1999:1	2028.40	2030.44	-2.04266	
1999:2	2036.90	2043.10	-6.19898	
1999:3	2063.30	2050.93	12.3718	
1999:4	2095.90	2073.20	22.7025	
2000:1	2078.70	2106.16	-27.4594	
2000:2	2106.40	2098.70	7.70476	
2000:3	2099.80	2115.94	-16.1432	
2000:4	2106.20	2113.62	-7.42154	
2001:1	2137.30	2115.82	21.4768	
2001:2	2181.70	2142.09	39.6118	*
2001:3	2177.80	2188.07	-10.2717	
2001:4	2216.40	2197.00	19.4047	
2002:1	2250.40	2228.45	21.9513	
2002:2	2272.00	2264.43	7.57376	
2002:3	2290.40	2290.22	0.176208	
2002:4	2305.70	2309.08	-3.37545	
2003:1	2300.90	2323.19	-22.2884	
2003:2	2335.10	2318.95	16.1501	
2003:3	2342.00	2343.36	-1.36407	
2003:4	2343.70	2355.14	-11.4422	
2004:1	2354.90	2357.34	-2.44240	
2004:2	2363.50	2365.01	-1.50594	
2004:3	2372.10	2373.07	-0.973166	
2004:4	2357.60	2381.43	-23.8266	
2005:1	2359.90	2369.89	-9.98745	
2005:2	2362.40	2366.52	-4.11691	
2005:3	2383.90	2367.42	16.4755	
2005:4	2373.40	2386.52	-13.1201	
2006:1	2397.10	2383.60	13.4969	

Note: * denotes a residual in excess of 2.5 standard errors


IV. Forecast

In this section, I use already defined models to forecast GDPC1Q and GCEC1Q from 2006 Q2 to 2013 Q1. I will also perform the forecasting accuracy assessment at the end of this session, comparing forecast with the actual data.

For Model 1.1 GDPC1Q ARIMA (1, 1, 0):

Obs	GDPC1Q	prediction	std. error	95% interval
2006:2	12948.7	12994.2	54.9179	(12886.5, 13101.8)
2006:3	12950.4	13071.2	91.2910	(12892.3, 13250.2)
2006:4	13038.4	13141.5	120.616	(12905.1, 13377.9)
2007:1	13056.1	13209.6	145.159	(12925.1, 13494.1)
2007:2	13173.6	13277.0	166.425	(12950.8, 13603.2)
2007:3	13269.8	13344.1	185.357	(12980.8, 13707.4)
2007:4	13326.0	13411.1	202.555	(13014.1, 13808.1)
2008:1	13266.8	13478.1	218.411	(13050.0, 13906.2)
2008:2	13310.5	13545.1	233.193	(13088.1, 14002.2)
2008:3	13186.9	13612.1	247.094	(13127.8, 14096.4)
2008:4	12883.5	13679.1	260.254	(13169.0, 14189.2)
2009:1	12711.0	13746.1	272.779	(13211.5, 14280.7)
2009:2	12701.0	13813.1	284.754	(13255.0, 14371.2)
2009:3	12746.7	13880.1	296.246	(13299.5, 14460.7)
2009:4	12873.1	13947.1	307.308	(13344.8, 14549.4)
2010:1	12947.6	14014.1	317.985	(13390.8, 14637.3)
2010:2	13019.6	14081.1	328.316	(13437.6, 14724.6)
2010:3	13103.5	14148.1	338.331	(13484.9, 14811.2)
2010:4	13181.2	14215.1	348.058	(13532.9, 14897.2)
2011:1	13183.8	14282.1	357.521	(13581.3, 14982.8)
2011:2	13264.7	14349.0	366.739	(13630.3, 15067.8)
2011:3	13306.9	14416.0	375.731	(13679.6, 15152.5)
2011:4	13441.0	14483.0	384.513	(13729.4, 15236.7)
2012:1	13506.4	14550.0	393.099	(13779.6, 15320.5)
2012:2	13548.5	14617.0	401.502	(13830.1, 15404.0)
2012:3	13652.5	14684.0	409.732	(13881.0, 15487.1)
2012:4	13665.4	14751.0	417.800	(13932.1, 15569.9)
2013:1	13750.1	14818.0	425.715	(13983.6, 15652.4)

For 95% confidence intervals, z(0.025) = 1.96



Overall, we can see that ARIMA (1, 1, 0) provides a good fit for GDPC1Q data. It gives a fairly accurate forecasting. However, although forecasts from 2006 Q2 to 2008 Q3 are within the 95% percent interval, the graph shows that the red line of actual data has gradually moving out of the confidence interval starting from 2007 Q4 and climbing back up toward the interval from 2009 Q1. Such trend exactly coincides with the way of how the economy has evolved since great recession of 2008. But, the weakness of ARIMA model is that it could not predict such trend but

rather assume the same pattern from 1980 Q1-2006 Q1, that is, the ARIMA model is not good for volatility analysis, but GARCH is. Finally, GDP does strive back after a deep dip in 2009 Q1.

For Model 1.2 GDPC1Q ARIMA (1, 1, 1):


For 95% confidence intervals, z(0.025) = 1.96

Obs	GDPC1Q	prediction	std. error	95% interval
2006:2	12948.7	12992.2	53.6190	(12887.1, 13097.3)
2006:2	12950.4	13080.9	87.1201	(12910.1, 13057.5)
2006:4	13038.4	13164.2	118.346	(12932.2, 13396.1)
2000:4	13056.1	13243.4	147.950	(12953.4, 13533.4)
2007:1	13173.6	13319.5	176.035	(12974.5, 13664.5)
2007:2	13269.8	13393.2	202.658	(12996.0, 13790.4)
2007:3	13209.0	13465.2	202.030	(12990.0, 13790.4) (13018.5, 13911.8)
2007:4	13266.8	13535.8	251.807	(13042.2, 14029.3)
2008:2	13200.0	13605.3	274.514	(13067.3, 14143.3)
2008:2	13186.9	13674.1	296.104	(13093.7, 14254.4)
2008:4	12883.5	13742.2	316.672	(13121.6, 14362.9)
2009:1	12711.0	13809.9	336.307	(13150.8, 14469.1)
2009:2	12701.0	13877.3	355.092	(13181.3, 14573.3)
2009:3	12746.7	13944.4	373.104	(13213.1, 14675.6)
2009:4	12873.1	14011.3	390.409	(13246.1, 14776.5)
2010:1	12947.6	14078.0	407.071	(13280.2, 14875.9)
2010:2	13019.6	14144.6	423.143	(13315.3, 14974.0)
2010:3	13103.5	14211.2	438.676	(13351.4, 15071.0)
2010:4	13181.2	14277.7	453.714	(13388.4, 15166.9)
2011:1	13183.8	14344.1	468.296	(13426.2, 15261.9)
2011:2	13264.7	14410.5	482.456	(13464.9, 15356.1)
2011:3	13306.9	14476.8	496.228	(13504.2, 15449.4)
2011:4	13441.0	14543.2	509.638	(13544.3, 15542.0)
2012:1	13506.4	14609.5	522.712	(13585.0, 15634.0)
2012:2	13548.5	14675.8	535.473	(13626.3, 15725.3)
2012:3	13652.5	14742.1	547.941	(13668.1, 15816.0)
2012:4	13665.4	14808.4	560.135	(13710.5, 15906.2)
2013:1	13750.1	14874.6	572.071	(13753.4, 15995.9)

Forecast evaluation statistics

Mean Error	-787.19
Mean Squared Error	8.2292e+005
Root Mean Squared Erro	or 907.15
Mean Absolute Error	787.19
Mean Percentage Error	-5.9698
Mean Absolute Percent	tage Error 5.9698
Theil's U	9.2447
Bias proportion, UM	0.75302

Regression proportion, UR0.18362Disturbance proportion, UD0.063366



The above forecasting graph indicates that ARIMA (1, 1, 1) overall provides a good fit for model 1. Compared with ARIMA (1, 1, 0), it has wider confidence interval (green area). We can see that the actual data from 2013 Q1 is actually touching the interval, however there still is a gap between the red line and the interval in the ARIMA (1, 1, 0) forecast. The reason for the green interval not capturing the red line is because of the unexpected 2008 recession. Hence, I think that GARCH might have the advantage over ARIMA, since it has the ability and room for volatility forecasting. It means that GARCH would allow us to forecast data, which is expected to have large variations. I think that is the purpose for heteroskedasticity to exist within such data.


Based on the forecasting performance, testing and estimating results, it is very difficult to choose between ARIMA (1, 1, 0) and ARIMA (1, 1, 1). But the residual ACF and PACF of ARIMA (1, 1, 1) provides a slightly better result than that of ARIMA (1, 1, 0). We can see that ACF and

PACF of ARIMA (1, 1, 0) has a spike crossing the ± 0.2 boundary at lag 2, while nothing is crossing the boundary in ACF and PACF of ARIMA (1, 1, 1). It means that ARIMA (1, 1, 1) is slightly more stable than ARIMA (1, 1, 0), which probably provides better forecasting results. I also want to point out that although MAPE of ARIMA (1, 1, 0) is slightly better than ARIMA (1, 1, 1) (5.5643% compared with 5.9698%), it does not mean that ARIMA (1, 1, 0) is better than ARIMA (1, 1, 1), because of the stability issue mentioned above. Plus, there are only two residuals of ARIMA (1, 1, 1) in excess of 2.5 standard errors, compared with three of ARIMA (1, 1, 0). Therefore, I will fit GDPC1Q data with ARIMA (1, 1, 1).

For Model 1.3 GCEC1Q ARIMA (1, 1, 2):

For 95% confidence intervals, z(0.025) = 1.96

Oha	CCECIO	nradiation	atd arman	95% interval
Obs 2006:2	GCEC1Q 2399.10	prediction 2401.85	std. error 15.2543	(2371.96, 2431.75)
2006:2	2399.10	2401.83	20.1999	(2371.65, 2450.83)
2006:4	2409.40	2420.70	25.6319	(2370.47, 2470.94)
2007:1	2406.70	2430.22	31.2272	(2369.02, 2491.43)
2007:2	2426.80	2439.79	36.8190	(2367.62, 2511.95)
2007:3	2447.90	2449.39	42.3182	(2366.45, 2532.33)
2007:4	2455.30	2459.02	47.6769	(2365.58, 2552.47)
2008:1	2473.90	2468.68	52.8708	(2365.06, 2572.31)
2008:2	2484.50	2478.36	57.8891	(2364.90, 2591.82)
2008:3	2510.70	2488.06	62.7296	(2365.11, 2611.01)
2008:4	2520.50	2497.77	67.3951	(2365.68, 2629.86)
2009:1	2531.60	2507.49	71.8915	(2366.59, 2648.40)
2009:2	2590.40	2517.22	76.2265	(2367.82, 2666.62)
2009:3	2614.30	2526.96	80.4084	(2369.36, 2684.56)
2009:4	2621.10	2536.70	84.4462	(2371.19, 2702.21)
2010:1	2600.40	2546.45	88.3487	(2373.29, 2719.61)
2010:2	2618.70	2556.20	92.1243	(2375.64, 2736.76)
2010:3	2616.70	2565.95	95.7812	(2378.22, 2753.68)
2010:4	2587.40	2575.71	99.3273	(2381.03, 2770.39)
2011:1	2540.70	2585.47	102.770	(2384.04, 2786.89)
2011:2	2535.40	2595.23	106.115	(2387.25, 2803.21)
2011:3	2516.60	2604.99	109.370	(2390.63, 2819.35)
2011:4	2502.70	2614.75	112.539	(2394.18, 2835.33)
2012:1	2483.70	2624.52	115.629	(2397.89, 2851.15)
2012:2	2479.40	2634.28	118.644	(2401.74, 2866.82)
2012:3	2503.10	2644.05	121.590	(2405.73, 2882.36)
2012:4	2458.10	2653.81	124.468	(2409.86, 2897.77)
2013:1	2427.10	2663.58	127.285	(2414.10, 2913.05)
	,			(=



Root Mean Squared Error87.855Mean Absolute Error62.244Mean Percentage Error-1.092Mean Absolute Percentage Error2.4814Theil's U3.8702Bias proportion, UM0.088923Regression proportion, UR0.35153Disturbance proportion, UD0.55955

The forecast red line is within the 95% confidence interval, which is a good sign for accurate forecasting. And overall, we can find that the ARIMA (1, 1, 2) fits well for the GCEC1Q dataset. We do see that there is a sharp reduction in Government Consumption Expenditures & Gross Investment from 2010 Q2. Compared with the above forecast graph of GDPC1Q, we can see the

confidence interval for GCEC1Q is much wider, meaning that the government spending each year has larger variations than GDP does and it becomes much more volatile.

......

(2) The Tiao-Box (1981) Multiple Time Series (Multivariate VARMA) Approach:

1980Q1-2006Q1 as the sample range; GDPC1Q and GCEC1Q as the observations.

Since we use ARIMA (1, 1, 1) for GDPC1Q and ARIMA (1, 1, 2) for GCEC1Q, we will first fit the sample data in either VARIMA (1, 1, 1) or VARIMA (1, 1, 2), then comparing the results and selecting the best one as our final model. VARIMA (1, 1, 0) is essentially VAR (1), which will be discussed in section 3.

Recall that we write univariate ARIMA (1, 1, 1) of GDPC1Q as:

$$\Delta^{1d} Y_t = \mu_0 + \varphi_1 \Delta^{1d} Y_{t-1} + \varepsilon_t + \theta_1 \varepsilon_{t-1}$$

And univariate ARIMA (1, 1, 2) of GCEC1Q as:

$$\Delta^{1d}Y_t = \mu_0 + \varphi_1 \Delta^{1d}Y_{t-1} + \varepsilon_t + \theta_1 \varepsilon_{t-1} + \theta_2 \varepsilon_{t-2}$$

Now, VARIMA (1, 1, 1) for both GDPC1Q (Y_1) and GCEC1Q (Y_2) can be written as:

$$\begin{bmatrix} \Delta^{1d} Y_{1t} \\ \Delta^{1d} Y_{2t} \end{bmatrix} = \begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix} + \begin{bmatrix} \varphi_{1,1} & \varphi_{1,2} \\ \varphi_{2,1} & \varphi_{2,2} \end{bmatrix} \begin{bmatrix} \Delta^{1d} Y_{1,t-1} \\ \Delta^{1d} Y_{2,t-1} \end{bmatrix} + \begin{bmatrix} \varepsilon_{1,t} \\ \varepsilon_{2,t} \end{bmatrix} + \begin{bmatrix} \theta_{1,1} & \theta_{1,2} \\ \theta_{2,1} & \theta_{2,2} \end{bmatrix} \begin{bmatrix} \varepsilon_{1,t-1} \\ \varepsilon_{2,t-1} \end{bmatrix}$$
(2.1)

(2.1) can also be written as:

$$\Delta^{1d}Y_{1t} = \mu_1 + \varphi_{1,1}\Delta^{1d}Y_{1,t-1} + \varphi_{1,2}\Delta^{1d}Y_{2,t-1} + \varepsilon_{1,t} + \theta_{1,1}\varepsilon_{1,t-1} + \theta_{1,2}\varepsilon_{2,t-1}$$
(2.2)
$$\Delta^{1d}Y_{2t} = \mu_2 + \varphi_{2,1}\Delta^{1d}Y_{1,t-1} + \varphi_{2,2}\Delta^{1d}Y_{2,t-1} + \varepsilon_{2,t} + \theta_{2,1}\varepsilon_{1,t-1} + \theta_{2,2}\varepsilon_{2,t-1}$$

```
gretl version 1.9.14
Current session: 2014-01-17 21:03
? series x = diff(GDPC1Q)
Generated series x (ID 11)
? series y = diff(GCEC1Q)
Generated series y (ID 12)
? series a = uhat1
Generated series a (ID 13)
? series b = uhat2
Generated series b (ID 14)
? system
? equation x const x(-1) y(-1) a(-1) b(-1)
```

? equation y const x(-1) y(-1) a(-1) b(-1)? end system ? estimate \$system method=ols Equation system, Ordinary Least Squares Equation 1: OLS, using observations 1980:2-2006:1 (T = 104) Dependent variable: x coefficient std. error t-ratio p-value _____ const82.2102129.9470.63260.5284x_10.7511790.2367253.1730.0020***y_1-6.4344712.6694-0.50790.6127a_1-0.4601450.259285-1.7750.0790*b_15.9178312.65800.46750.6412 Mean dependent var 67.24038 S.D. dependent var 58.45773 Sum squared resid 294663.4 S.E. of regression 54.55638 0.162846 Adjusted R-squared 0.129022 R-squared Equation 2: OLS, using observations 1980:2-2006:1 (T = 104) Dependent variable: y coefficient std. error t-ratio p-value _____ const8.5912538.16380.22510.8224x_10.06014900.06952350.86520.3890y_1-0.2619463.72085-0.070400.9440a_1-0.1058780.0761489-1.3900.1675b_10.2795123.717510.075190.9402 Mean dependent var 9.920192 S.D. dependent var 15.95743 Sum squared resid 25415.58 S.E. of regression 16.02258 R-squared 0.030971 Adjusted R-squared -0.008182 Cross-equation VCV for residuals (correlations above the diagonal) 2833.3 (0.246) 204.94 244.38 $\log determinant = 13.3853$ Breusch-Pagan test for diagonal covariance matrix: Chi-square(1) = 6.30863 [0.0120]Now, we can write (2.1) and (2.2) as: $\begin{bmatrix} \Delta^{1d} Y_{1t} \\ \Delta^{1d} Y_{2t} \end{bmatrix} = \begin{bmatrix} 82.2102 \\ 8.59125 \end{bmatrix} + \begin{bmatrix} 0.751179 & -6.43447 \\ 0.060149 & -0.261946 \end{bmatrix} \begin{bmatrix} \Delta^{1d} Y_{1,t-1} \\ \Delta^{1d} Y_{2,t-1} \end{bmatrix} + \begin{bmatrix} \varepsilon_{1,t} \\ \varepsilon_{2,t} \end{bmatrix}$ (2.1') $+\begin{bmatrix}-0.460145 & 5.91783\\-0.105878 & 0.279512\end{bmatrix}\begin{bmatrix}\varepsilon_{1,t-1}\\\varepsilon_{2,t-1}\end{bmatrix}$

$$\Delta^{1d} Y_{1t} = 82.2102 + 0.751179 \Delta^{1d} Y_{1,t-1} - 6.43447 \Delta^{1d} Y_{2,t-1} + \varepsilon_{1,t}$$
(2.2')
- 0.460145\varepsilon_{1,t-1} + 5.91783\varepsilon_{2,t-1}

$$\Delta^{1d} Y_{2t} = 8.59125 + 0.060149 \Delta^{1d} Y_{1,t-1} - 0.261946 \Delta^{1d} Y_{2,t-1} + \varepsilon_{2,t}$$
$$- 0.105878 \varepsilon_{1,t-1} + 0.279512 \varepsilon_{2,t-1}$$

Notes: error terms uhat1 and uhat2 are from ARIMA (1, 1, 1) of GDPC1Q and ARIMA (1, 1, 1) of GCEC1Q respectively.

In comparison, VARIMA (1, 1, 2) for both GDPC1Q (Y_1) and GCEC1Q (Y_2) can be written as:

$$\begin{bmatrix} \Delta^{1d} Y_{1t} \\ \Delta^{1d} Y_{2t} \end{bmatrix} = \begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix} + \begin{bmatrix} \varphi_{1,1} & \varphi_{1,2} \\ \varphi_{2,1} & \varphi_{2,2} \end{bmatrix} \begin{bmatrix} \Delta^{1d} Y_{1,t-1} \\ \Delta^{1d} Y_{2,t-1} \end{bmatrix} + \begin{bmatrix} \varepsilon_{1,t} \\ \varepsilon_{2,t} \end{bmatrix} + \begin{bmatrix} \theta_{1,1} & \theta_{1,2} \\ \theta_{2,1} & \theta_{2,2} \end{bmatrix} \begin{bmatrix} \varepsilon_{1,t-1} \\ \varepsilon_{2,t-1} \end{bmatrix}$$
(2.3)
$$+ \begin{bmatrix} \vartheta_{1,1} & \vartheta_{1,2} \\ \vartheta_{2,1} & \vartheta_{2,2} \end{bmatrix} \begin{bmatrix} \varepsilon_{1,t-2} \\ \varepsilon_{2,t-2} \end{bmatrix}$$

(2.3) can also be written as:

$$\Delta^{1d}Y_{1t} = \mu_1 + \varphi_{1,1}\Delta^{1d}Y_{1,t-1} + \varphi_{1,2}\Delta^{1d}Y_{2,t-1} + \varepsilon_{1,t} + \theta_{1,1}\varepsilon_{1,t-1} + \theta_{1,2}\varepsilon_{2,t-1}$$
(2.4)
+ $\vartheta_{1,1}\varepsilon_{1,t-2} + \vartheta_{1,2}\varepsilon_{2,t-2}$

$$\Delta^{1d} Y_{2t} = \mu_2 + \varphi_{2,1} \Delta^{1d} Y_{1,t-1} + \varphi_{2,2} \Delta^{1d} Y_{2,t-1} + \varepsilon_{2,t} + \theta_{2,1} \varepsilon_{1,t-1} + \theta_{2,2} \varepsilon_{2,t-1} + \theta_{2,1} \varepsilon_{1,t-2} + \theta_{2,2} \varepsilon_{2,t-2}$$

```
gret1 version 1.9.14
Current session: 2014-01-17 21:27
? series x = diff(GDPC1Q)
Generated series x (ID 11)
? series y = diff(GCEC1Q)
Generated series y (ID 12)
? series a = uhat1
Generated series a (ID 13)
? series b = uhat2
Generated series b (ID 14)
? system
? equation x const x(-1) y(-1) a(-1) b(-1) a(-2) b(-2)
? equation y const x(-1) y(-1) a(-1) b(-1) a(-2) b(-2)
? end system
? estimate $system method=ols
```

Equation system, Ordinary Least Squares

Equation 1: OLS, using observations 1980:3-2006:1 (T = 103) Dependent variable: $\mathbf x$

	coefficient	std. error	t-ratio	p-value	
const	46.7843	21.0741	2.220	0.0288 **	
x_1	0.369566	0.267391	1.382	0.1701	
y_1	-0.219727	1.18141	-0.1860	0.8528	
a 1	-0.113708	0.283210	-0.4015	0.6889	

-0.1326011.22455-0.10830.91400.2165160.1240601.7450.0841*-0.1043260.379611-0.27480.7840 b 1 a 2 b_2 Mean dependent var 69.06796 S.D. dependent var 55.67800 Sum squared resid 263046.7 S.E. of regression 52.34568 R-squared 0.168111 Adjusted R-squared 0.116118 Equation 2: OLS, using observations 1980:3-2006:1 (T = 103) Dependent variable: y coefficient std. error t-ratio p-value _____ const0.1566036.333610.024730.9803x_10.005823490.08036180.072470.9424y_10.9505560.3550622.6770.0087a_1-0.03941230.0851159-0.46300.6444b_1-1.004330.368026-2.7290.0076a_20.01118160.03728500.29990.7649b_20.1655780.1140891.4510.1500 Mean dependent var 9.974757 S.D. dependent var 16.02571 Sum squared resid 23759.55 S.E. of regression 15.73198 R-squared 0.093008 Adjusted R-squared 0.036321 Cross-equation VCV for residuals (correlations above the diagonal) 2553.9 (0.264)202.99 230.68 $\log determinant = 13.2139$ Breusch-Pagan test for diagonal covariance matrix: Chi-square(1) = 7.20435 [0.0073]

Now, we can write (2.3) and (2.4) as:

$$\begin{split} \begin{bmatrix} \Delta^{1d} Y_{1t} \\ \Delta^{1d} Y_{2t} \end{bmatrix} &= \begin{bmatrix} 46.7843 \\ 0.156603 \end{bmatrix} + \begin{bmatrix} 0.369566 & -0.219727 \\ 0.00582349 & 0.950556 \end{bmatrix} \begin{bmatrix} \Delta^{1d} Y_{1,t-1} \\ \Delta^{1d} Y_{2,t-1} \end{bmatrix} + \begin{bmatrix} \varepsilon_{1,t} \\ \varepsilon_{2,t} \end{bmatrix} & (2.3') \\ &+ \begin{bmatrix} -0.113708 & -0.132601 \\ -0.0394123 & -1.00433 \end{bmatrix} \begin{bmatrix} \varepsilon_{1,t-1} \\ \varepsilon_{2,t-1} \end{bmatrix} \\ &+ \begin{bmatrix} 0.216516 & -0.104326 \\ 0.0111816 & 0.165578 \end{bmatrix} \begin{bmatrix} \varepsilon_{1,t-2} \\ \varepsilon_{2,t-2} \end{bmatrix} \\ \Delta^{1d} Y_{1t} &= 46.7843 + 0.369566\Delta^{1d} Y_{1,t-1} - 0.219727\Delta^{1d} Y_{2,t-1} + \varepsilon_{1,t} \\ &- 0.113708\varepsilon_{1,t-1} - 0.132601\varepsilon_{2,t-1} + 0.216516\varepsilon_{1,t-2} \end{bmatrix}$$

 $-0.104326\varepsilon_{2,t-2}$

$$\begin{split} \Delta^{1d}Y_{2t} &= 0.156603 + 0.00582349 \Delta^{1d}Y_{1,t-1} + 0.950556 \Delta^{1d}Y_{2,t-1} + \varepsilon_{2,t} \\ &\quad - 0.0394123\varepsilon_{1,t-1} - 1.00433\varepsilon_{2,t-1} + 0.0111816\varepsilon_{1,t-2} \\ &\quad + 0.165578\varepsilon_{2,t-2} \end{split}$$

Notes: error terms uhat1 and uhat2 are from ARIMA (1, 1, 2) of GDPC1Q and ARIMA (1, 1, 2) of GCEC1Q respectively.

Essentially, Multivariate VARMA (2.1) and (2.3) can be transformed back to Univariate ARMA, if:


$$\begin{bmatrix} \varphi_{1,1} & \varphi_{1,2} \\ \varphi_{2,1} & \varphi_{2,2} \end{bmatrix} = \begin{bmatrix} \varphi_{1,1} & 0 \\ 0 & \varphi_{2,2} \end{bmatrix} \text{ and } \begin{bmatrix} \theta_{1,1} & \theta_{1,2} \\ \theta_{2,1} & \theta_{2,2} \end{bmatrix} = \begin{bmatrix} \theta_{1,1} & 0 \\ 0 & \theta_{2,2} \end{bmatrix} \text{ and } \begin{bmatrix} \vartheta_{1,1} & \vartheta_{1,2} \\ \vartheta_{2,1} & \vartheta_{2,2} \end{bmatrix} = \begin{bmatrix} \vartheta_{1,1} & 0 \\ 0 & \vartheta_{2,2} \end{bmatrix}$$

We conclude that Multivariate VARMA is more generalized than Univariate ARMA.

The findings are not surprising at all that both VARIMA (1, 1, 1) and VARIMA (1, 1, 2) are not good at explaining the relationship between GDPC1Q and GCEC1Q and between GCEC1Q and GDPC1Q, either showing the statistical significance for such relationships. Thereby, we resort to VAR (1) analysis in the following section 3, because VAR (1) shares the same component AR with both ARIMA (1, 1, 1) of GDPC1Q and ARIMA (1, 1, 2) of GCEC1Q, in comparison VARIMA (1, 1, 1) does not share the same MA component with ARIMA (1, 1, 2) and neither does VARIMA (1, 1, 2) share the same MA component with ARIMA (1, 1, 1), which might be the reason that VAR (1) is more widely used in the field compared to VARMA that however is more generalized than VAR, because of which we can consider both Univariate ARMA, VAR and VMA as special cases of VARMA.

(3) The VAR (p) Model and Impulse-Response Function:

1980Q1-2006Q1 as sample range; GDPC1Q and GCEC1Q as endogenous variables.



lag

I. Identification:

Since the above correlogram indicates nonstationarity, we have to difference GDPC1Q and GCEC1Q for at least once, denoting them as d GDPC1Q and d GCEC1Q.

VAR sys	stem, maximum	lag order	8		
The asterisks below indicate the best (that is, minimized) values of the respective information criteria, AIC = Akaike criterion, BIC = Schwarz Bayesian criterion and HQC = Hannan-Quinn criterion.					
lags	loglik	p(LR)	AIC	BIC	HQC
1 2 3 4 5 6 7 8	-914.45020 -910.05198 -904.12837 -901.71218 -900.28794 -899.41488 -897.98592 -889.94431	0.06639 0.01852 0.30493 0.58349 0.78232 0.58188 0.00291	18.978355 18.970144 18.930482* 18.963138 19.016246 19.080719 19.133730 19.050398	19.137615* 19.235578 19.302090 19.440919 19.600201 19.770848 19.930033 19.952874	19.042752* 19.077472 19.080742 19.156329 19.252369 19.359773 19.455716 19.415315

The above analysis tells us to identify the model as VAR (1), because both BIC and HQC are significant at lag 1.

II. Estimation:

VAR system, lag order 1 OLS estimates, observations 1980:2-2006:1 (T = 104) Log-likelihood = -993.60761 Determinant of covariance matrix = 681522.47AIC = 19.2232BIC = 19.3758HQC = 19.2850Portmanteau test: LB(26) = 109.69, df = 100 [0.2385]

Equation 1: d GDPC1Q

	Coefficient	Std. Er	rror	t-ratio	p-value	
const	49.3131	8.48	76	5.8100	< 0.00001	***
d_GDPC1Q_1	0.360705	0.0954	613	3.7785	0.00027	***
d_GCEC1Q_1	-0.589345	0.346	763	-1.6996	0.09229	*
Mean dependent var	67.2	4038	S.D. 0	lependent var	58.	45773
Sum squared resid	3058	327.1	S.E. o	of regression	55.	02718
R-squared	0.13	1130	Adjus	sted R-squared	0.1	13925
F(2, 101)	7.62	1464	P-valu	ue(F)	0.0	00826
rho	-0.09	6450	Durbi	n-Watson	2.0	78413

F-tests of zero restrictions: All lags of d_GDPC1Q F(1, 101) = 14.277 [0.0003]All lags of d_GCEC1Q F(1, 101) = 2.8885 [0.0923]

Equation 2: d_GCEC1Q

	Coefficient	Std. Er	rror	t-ratio	p-value	
const	11.8267	2.471	76	4.7847	< 0.0000	***
d GDPC1Q 1	-0.0290698	0.0278	002	-1.0457	0.29821	
d_GCEC1Q_1	0.000855439	0.1009	984	0.0085	0.99326	
Mean dependent va	r 9.920)192	S.D. (dependent var	1	5.95743
Sum squared resid	2593	6.91	S.E. 0	of regression	1	6.02501
R-squared	0.01	1093	Adjus	sted R-squared	-0	.008489
F(2, 101)	0.566	5501	P-val	ue(F)	0	.569301
rho	0.014	4121	Durb	in-Watson	1	.962761
	F-tests	of zero	restric	tions:		
All la	gs of d_GDPC1	Q F(1	l, 101)	= 1.0934 [0.2	.982]	
All lags	of d_GCEC1Q	F(1, 1	101) =	7.1758e-005 [0).9933]	

Let $y_{d_GDPC1Q,t}$ be $\Delta^{1d}y_{1,t} = y_{1t} - y_{1,t-1}$ and $y_{d_GCEC1Q,t}$ be $\Delta^{1d}y_{2,t} = y_{2t} - y_{2,t-1}$, then we have VAR (1) model in matrix form:

$$\begin{bmatrix} \Delta^{1d} y_{1,t} \\ \Delta^{1d} y_{2,t} \end{bmatrix} = \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} + \begin{bmatrix} A_{1,1} & A_{1,2} \\ A_{2,1} & A_{2,2} \end{bmatrix} \begin{bmatrix} \Delta^{1d} y_{1,t-1} \\ \Delta^{1d} y_{2,t-1} \end{bmatrix} + \begin{bmatrix} e_{1,t} \\ e_{2,t} \end{bmatrix}$$
(3.1)

Equation (3.1) can also be written as:

$$\Delta^{1d} y_{1,t} = c_1 + A_{1,1} \Delta^{1d} y_{1,t-1} + A_{1,2} \Delta^{1d} y_{2,t-1} + e_{1,t}$$

$$\Delta^{1d} y_{2,t} = c_2 + A_{2,1} \Delta^{1d} y_{1,t-1} + A_{2,2} \Delta^{1d} y_{2,t-1} + e_{2,t}$$
(3.2)

Inputting the estimated parameters into equation (3.1) and (3.2), we have:

$$\begin{bmatrix} \Delta^{1d} y_{1,t} \\ \Delta^{1d} y_{2,t} \end{bmatrix} = \begin{bmatrix} 49.3131 \\ 11.8267 \end{bmatrix} + \begin{bmatrix} 0.360705 & -0.589345 \\ -0.0290698 & 0.000855439 \end{bmatrix} \begin{bmatrix} \Delta^{1d} y_{1,t-1} \\ \Delta^{1d} y_{2,t-1} \end{bmatrix} + \begin{bmatrix} e_{1,t} \\ e_{2,t} \end{bmatrix}$$
(3.3)
$$\Delta^{1d} y_{1,t} = 49.3131 + 0.360705\Delta^{1d} y_{1,t-1} - 0.589345\Delta^{1d} y_{2,t-1} + e_{1,t}$$
(3.4)
$$\Delta^{1d} y_{2,t} = 11.8267 - 0.0290698\Delta^{1d} y_{1,t-1} + 0.000855439\Delta^{1d} y_{2,t-1} + e_{2,t}$$

Impulse Responses:

2	17.085	-1.5728
3	7.0897	-0.49801
4	2.8508	-0.20652
5	1.15	-0.083049
6	0.46376	-0.033502
7	0.18702	-0.01351
8	0.075423	-0.0054483
9	0.030416	-0.0021972
10	0.012266	-0.00088608
11	0.0049467	-0.00035734
12	0.0019949	-0.00014411
13	0.0008045	-5.8115e-005
14	0.00032444	-2.3436e-005
15	0.00013084	-9.4513e-006
16	5.2764e-005	-3.8115e-006
17	2.1279e-005	-1.5371e-006
18	8.5812e-006	-6.1988e-007
19	3.4606e-006	-2.4998e-007
20	1.3956e-006	-1.0081e-007

Responses to a one-standard error shock in d_GCEC1Q


period	d_GDPC1Q	d_GCEC1Q
1	0	15.224
2	-8.972	0.013023
3	-3.2439	0.26082
4	-1.3238	0.094523
5	-0.53321	0.038564
6	-0.21506	0.015533
7	-0.086728	0.006265
8	-0.034976	0.0025265
9	-0.014105	0.0010189
10	-0.0056882	0.0004109
11	-0.0022939	0.00016571
12	-0.00092509	6.6825e-005
13	-0.00037307	2.6949e-005
14	-0.00015045	1.0868e-005
15	-6.0673e-005	4.3828e-006
16	-2.4468e-005	1.7675e-006
17	-9.8674e-006	7.1279e-007
18	-3.9793e-006	2.8745e-007
19	-1.6048e-006	1.1592e-007
20	-6.4717e-007	4.6749e-008


Decomposition of variance for d_GDPC1Q period std. error d_GDPC1Q d_GCEC1Q

54.2277	100.0000	0.0000
57.5591	97.5703	2.4297
58.0848	97.3022	2.6978
58.1697	97.2583	2.7417
58.1836	97.2512	2.7488
58.1858	97.2500	2.7500
58.1862	97.2498	2.7502
58.1862	97.2498	2.7502
58.1862	97.2498	2.7502
58.1862	97.2498	2.7502
58.1862	97.2498	2.7502
58.1862	97.2498	2.7502
58.1862	97.2498	2.7502
58.1862	97.2498	2.7502
58.1862	97.2498	2.7502
58.1862	97.2498	2.7502
58.1862	97.2498	2.7502
58.1862	97.2498	2.7502
58.1862	97.2498	2.7502
58.1862	97.2498	2.7502
	58.0848 58.1697 58.1836 58.1858 58.1862 58.1862 58.1862 58.1862 58.1862 58.1862 58.1862 58.1862 58.1862 58.1862 58.1862 58.1862 58.1862 58.1862 58.1862 58.1862 58.1862 58.1862	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$

Decomposition of variance for d_GCEC1Q


period	std. error	d GDPC1Q	d GCEC1Q
1	15.7922	7.0707	92.9293
2	15.8703	7.9834	92.0166
3	15.8803	8.0717	91.9283
4	15.8819	8.0870	91.9130
5	15.8822	8.0895	91.9105
6	15.8822	8.0899	91.9101
7	15.8822	8.0899	91.9101
8	15.8822	8.0899	91.9101
9	15.8822	8.0899	91.9101
10	15.8822	8.0899	91.9101
11	15.8822	8.0899	91.9101
12	15.8822	8.0899	91.9101
13	15.8822	8.0899	91.9101
14	15.8822	8.0899	91.9101
15	15.8822	8.0899	91.9101
16	15.8822	8.0899	91.9101
17	15.8822	8.0899	91.9101
18	15.8822	8.0899	91.9101
19	15.8822	8.0899	91.9101
20	15.8822	8.0899	91.9101






response of d_GDPC1Q to a shock in d_GCEC1Q, with bootstrap confidence interval

The above analysis tells us that d_GDPC1Q reacts positively to a shock in d_GCEC1Q (both decrease) between period 1 and 2, positively (both increase) from 2 to 3, and negatively (d_GCEC1Q decrease, d_GDPC1Q increase) from 3 to 15. But, starting from 15, the shock of d_GCEC1Q fades away. The overall sign of impulse response is negative (except for period 1), which conforms to the sign of coefficient (-0.589345) of $y_{2,t-1}$ in equation 1 of (3.4). It implies that a shock in government consumption expenditure and gross investment can either decrease or increase GDP temporarily, however GDP will react negatively to such shock in the long run. Hence, the results suggest that government should keep a steady rather than volatile fiscal policy. Since the shock of GCEC on GDP fades out after certain period of time, influencing GDP merely through government expenditure is not optimal in the long run. Further researches are needed for a more comprehensive and effective approach. Final equilibrium is reached when government expenditure shock has little to no impact on GDP.




In contrast, d_GCEC1Q reacts positively to a shock in d_GCEC1Q (both decrease) between period 1 and 2, negatively from 2 to 3 (d_GDPC1Q decrease, d_GCEC1Q increase), and negatively from 3 to 15 (d_GDPC1Q decrease, d_GCEC1Q increase). However, a shock in d_GDPC1Q after 15 is fading out. The overall sign of impulse response is negative (except for period 1), which corresponds to the sign of coefficient (-0.0290698) of $y_{1,t-1}$ in equation 2 of (3.4). The results suggest that GDP does influence government spending negatively in the long run, but its negative impact force gradually reduces over time. Interestingly, government spending response negatively to GDP shock from period 2 to 3. One theorizes that government try to stimulate weak economy through spending, but unexpectedly continuous expenditure will hurt GDP in the long term. But equilibrium is reached when GDP shock has little to no effect on fiscal policy.


In comparison, one finds that a shock of government spending has more profound influence on GDP than a shock of GDP on expenditure, because the absolute value of coefficient of $y_{2,t-1}$ in equation 1 is larger than that of $y_{1,t-1}$ in equation 2 (0.589345 > 0.0290698). The response impulse graphs also hints that response of d_GDPC1Q to a shock in d_GCEC1Q has a deeper valley (curve) than response of d_GCEC1Q to d_GDPC1Q.




VAR residuals



From the VAR residuals time plot, we can see that residuals of equation $y_{2,t}(d_GCEC1Q)$ has narrower bandwidth than $y_{1,t}(d_GDPC1Q)$. The VAR inverse roots chart shows that VAR (1) is a stable system with both roots within the unit circle.



VAR inverse roots in relation to the unit circle



From the above correlograms, we can find that uhat1 (residual from VAR system, equation 1) appears slightly more stable than uhat2 (residual from VAR system, equation 2), because uhat2 is over the upper bound 0.2 at lag 3.

III. Diagnostic Checking:

1) Autocorrelation test:

```
Equation 1:
Ljung-Box Q' = 8.5046 with p-value = P(Chi-square(4) > 8.5046) = 0.0747
Equation 2:
Ljung-Box Q' = 10.2071 with p-value = P(Chi-square(4) > 10.2071) = 0.0371
```

Autocorrelation test passes for both equation 1 and 2 at p-value of 0.0747 and 0.0371 respectively. The null hypothesis of no autocorrelation cannot be rejected.

2) ARCH test:

Test for ARCI	H of order 4			
Equation 1:	coefficient	std. error	t-ratio	p-value
alpha(0) alpha(1)	1472.71 0.110710	618.846 0.102541	2.380 1.080	0.0193 ** 0.2830

ARCH test passes for both equation 1 and 2 with p-value of 0.0610456 and 0.681783 respectively. The null hypothesis of no ARCH effect cannot be rejected.

3) Multivariate normality test:

```
Residual correlation matrix, C (2 x 2)

1.0000 0.26591

0.26591 1.0000

Eigenvalues of C

0.734092

1.26591

Doornik-Hansen test

Chi-square(4) = 7.23579 [0.1239]
```

Doornik-Hansen test shows p value of 0.1239. The null hypothesis of normality cannot be rejected at any significant level.

4) Cointegration test (lag order 4):


```
Step 1: testing for a unit root in d_GDPClQ
Augmented Dickey-Fuller test for d_GDPClQ
including 3 lags of (1-L)d_GDPClQ
(max was 4, criterion modified AIC)
sample size 100
unit-root null hypothesis: a = 1
test with constant
model: (1-L)y = b0 + (a-1)*y(-1) + ... + e
1st-order autocorrelation coeff. for e: 0.041
lagged differences: F(3, 95) = 4.212 [0.0076]
estimated value of (a - 1): -0.444352
```

test statistic: tau c(1) = -3.33419asymptotic p-value 0.01345 Step 2: testing for a unit root in d GCEC1Q Augmented Dickey-Fuller test for d GCEC1Q including 4 lags of (1-L)d GCEC1Q (max was 4, criterion modified AIC) sample size 100 unit-root null hypothesis: a = 1 test with constant model: $(1-L)y = b0 + (a-1)*y(-1) + \dots + e$ 1st-order autocorrelation coeff. for e: -0.003 lagged differences: F(4, 94) = 2.567 [0.0431]estimated value of (a - 1): -0.566423test statistic: tau c(1) = -2.98579asymptotic p-value 0.03625 Step 3: cointegrating regression Cointegrating regression -OLS, using observations 1980:1-2006:1 (T = 105) Dependent variable: d GDPC1Q coefficient std. error t-ratio p-value _____ const59.24156.630338.9351.72e-014***d_GCEC1Q0.7514920.3535462.1260.0359** Mean dependent var 66.78000 S.D. dependent var 58.36696 Sum squared resid339408.8S.E. of regression57.40410R-squared0.042022Adjusted R-squared0.032721Log-likelihood-573.2411Akaike criterion1150.482Schwarz griterion1155.700Warpan Ouipp1152.623
 Schwarz criterion
 1155.790
 Hannan-Quinn
 1152.633

 rho
 0.383388
 Durbin-Watson
 1.218974
 Step 4: testing for a unit root in uhat Augmented Dickey-Fuller test for uhat including one lag of (1-L)uhat (max was 4, criterion modified AIC) sample size 100 unit-root null hypothesis: a = 1 model: $(1-L)y = (a-1)*y(-1) + \ldots + e$ 1st-order autocorrelation coeff. for e: 0.038 estimated value of (a - 1): -0.442103test statistic: tau c(2) = -4.05898asymptotic p-value 0.005862 There is evidence for a cointegrating relationship if: (a) The unit-root hypothesis is not rejected for the individual variables. (b) The unit-root hypothesis is rejected for the residuals (uhat) from the cointegrating regression.

The cointegration test results indicate that (a) The unit-root hypothesis is rejected for d_GDPC1Q and d_GCEC1Q; (b) The unit-root hypothesis is rejected for the residuals (uhat) from the cointegrating regression. Thereby, cointegration is not present for variables d_GDPC1Q

and d_GCEC1Q. VAR(1) is adequate and VECM is not needed. Additionally, residuals plot for both equations has relatively stable trend. Based on the results of diagnostic checking, the fitted model is appropriate.



IV. Forecast:


For 95% confidence intervals, t(101, 0.025) = 1.984

Obs	d GDPC1Q	prediction	std. error	95% interval
2006:2	undefined	93.3470	54.2277	(-14.2262, 200.920)
2006:3	undefined	78.7567	57.5591	(-35.4251, 192.939)
2006:4	undefined	72.3467	58.0848	(-42.8779, 187.571)
2007:1	undefined	69.7836	58.1697	(-45.6096, 185.177)
2007:2	undefined	68.7490	58.1836	(-46.6715, 184.170)
2007:3	undefined	68.3318	58.1858	(-47.0931, 183.757)
2007:4	undefined	68.1636	58.1862	(-47.2621, 183.589)
2008:1	undefined	68.0958	58.1862	(-47.3301, 183.522)
2008:2	undefined	68.0684	58.1862	(-47.3574, 183.494)
2008:3	undefined	68.0574	58.1862	(-47.3685, 183.483)
2008:4	undefined	68.0529	58.1862	(-47.3729, 183.479)
2009:1	undefined	68.0511	58.1862	(-47.3747, 183.477)
2009:2	undefined	68.0504	58.1862	(-47.3754, 183.476)
2009:3	undefined	68.0501	58.1862	(-47.3757, 183.476)
2009:4	undefined	68.0500	58.1862	(-47.3759, 183.476)
2010:1	undefined	68.0499	58.1862	(-47.3759, 183.476)

2010:2	undefined	68.0499	58.1862	(-47.3759, 183.476)
2010:3	undefined	68.0499	58.1862	(-47.3759, 183.476)
2010:4	undefined	68.0499	58.1862	(-47.3759, 183.476)
2011:1	undefined	68.0499	58.1862	(-47.3759, 183.476)
2011:2	undefined	68.0499	58.1862	(-47.3759, 183.476)
2011:3	undefined	68.0499	58.1862	(-47.3759, 183.476)
2011:4	undefined	68.0499	58.1862	(-47.3759, 183.476)
2012:1	undefined	68.0499	58.1862	(-47.3759, 183.476)
2012:2	undefined	68.0499	58.1862	(-47.3759, 183.476)
2012:3	undefined	68.0499	58.1862	(-47.3759, 183.476)
2012:4	undefined	68.0499	58.1862	(-47.3759, 183.476)
2013:1	undefined	68.0499	58.1862	(-47.3759, 183.476)

For 95% confidence intervals, t(101, 0.025) = 1.984

Obs	d GCEC1Q	prediction	std. error	95% interval
2006:2	undefined	7.17258	15.7922	(-24.1549, 38.5000)
2006:3	undefined	9.11928	15.8703	(-22.3632, 40.6017)
2006:4	undefined	9.54508	15.8803	(-21.9571, 41.0473)
2007:1	undefined	9.73179	15.8819	(-21.7736, 41.2372)
2007:2	undefined	9.80646	15.8822	(-21.6995, 41.3124)
2007:3	undefined	9.83659	15.8822	(-21.6694, 41.3426)
2007:4	undefined	9.84875	15.8822	(-21.6573, 41.3548)
2008:1	undefined	9.85365	15.8822	(-21.6524, 41.3597)
2008:2	undefined	9.85562	15.8822	(-21.6504, 41.3617)
2008:3	undefined	9.85642	15.8822	(-21.6496, 41.3625)
2008:4	undefined	9.85674	15.8822	(-21.6493, 41.3628)
2009:1	undefined	9.85687	15.8822	(-21.6492, 41.3629)
2009:2	undefined	9.85692	15.8822	(-21.6491, 41.3630)
2009:3	undefined	9.85695	15.8822	(-21.6491, 41.3630)
2009:4	undefined	9.85695	15.8822	(-21.6491, 41.3630)
2010:1	undefined	9.85696	15.8822	(-21.6491, 41.3630)
2010:2	undefined	9.85696	15.8822	(-21.6491, 41.3630)
2010:3	undefined	9.85696	15.8822	(-21.6491, 41.3630)
2010:4	undefined	9.85696	15.8822	(-21.6491, 41.3630)
2011:1	undefined	9.85696	15.8822	(-21.6491, 41.3630)
2011:2	undefined	9.85696	15.8822	(-21.6491, 41.3630)
2011:3	undefined	9.85696	15.8822	(-21.6491, 41.3630)
2011:4	undefined	9.85696	15.8822	(-21.6491, 41.3630)
2012:1	undefined	9.85696	15.8822	(-21.6491, 41.3630)
2012:2	undefined	9.85696	15.8822	(-21.6491, 41.3630)
2012:3	undefined	9.85696	15.8822	(-21.6491, 41.3630)
2012:4	undefined	9.85696	15.8822	(-21.6491, 41.3630)
2013:1	undefined	9.85696	15.8822	(-21.6491, 41.3630)



VAR (1) forecast of d_GDPC1Q shows downward trend, compared with upward of d_GCEC1Q. Interestingly, both forecasts are moving exactly the opposite direction at the initial stage, but only the decreasing rate of d_GDPC1Q is much larger than the increasing of d_GCEC1Q. In one's opinion, empirical cause and effect is very evident for those two relations, GDP and Government. It is fruitful for future researchers to explore such phenomenon, critically important to the fiscal policy maker.

(4) A State-Space Model with A Kalman Filter (ARMA estimation via Kalman example)

For Model 1.2 GDPC1Q ARIMA (1, 1, 1):

Recall that:

First difference of GDPC1Q (*I*, 1)=d_GDPC1Q and $Y_{d_GDPC1Q} = \Delta^{1d}Y_1 = Y_1 - Y_{1,t-1}$

$$\Delta^{1d}Y_{1t} = c_1 + \varphi_1 \Delta^{1d}Y_{1,t-1} + \varepsilon_{1t} + \theta_1 \varepsilon_{1,t-1}$$

$$\Delta^{1d} Y_{1t} = 66.2629 + 0.759918 \Delta^{1d} Y_{1,t-1} + \varepsilon_{1t} - 0.479305 \varepsilon_{1,t-1}$$

Based on James Hamilton's with slight variations, a state-space model can be written as:

$$\xi_{t+1} = \mathbf{F}_t \xi_t + \mathbf{v}_t \tag{4.1}$$

$$y_t = A'_t \mathbf{x}_t + \mathbf{H}'_t \boldsymbol{\xi}_t + \mathbf{w}_t \tag{4.2}$$

Where (4.1) is the state transition equation and (4.2) is the observation or measurement equation. The state vector, ξ_t , is $(r \times 1)$ and the vector of observables, y_t , is $(n \times 1)$; x_t is a $(k \times 1)$ vector of exogenous variables. The $(r \times 1)$ vector v_t and the $(n \times 1)$ vector w_t are assumed to be vector white noise:

$$E(\mathbf{v}_t \mathbf{v}'_s) = \mathbf{Q}_t \text{ for } t = s, \text{ otherwise } 0$$
$$E(\mathbf{w}_t \mathbf{w}'_s) = \mathbf{R}_t \text{ for } t = s, \text{ otherwise } 0$$

The Kalman filter provides a very efficient yet accurate recursive way to compute the likelihood of ARMA models. ARMA (1, 1) can be written as:

$$y_t = \phi y_{t-1} + \varepsilon_t + \theta \varepsilon_{t-1} \tag{4.3}$$

In order for equation (4.3) to be transformed into state-space form, we can define a latent process $\xi_t = (1 - \phi L)^{-1} \varepsilon_t$. Then, the observation equation (4.3) can be rewritten as:

$$y_t = \xi_t + \theta \xi_{t-1} \tag{4.4}$$

Construct the following two-equation system:

$$\xi_t = \phi \xi_{t-1} + \varepsilon_t$$

$$\xi_{t-1} = \xi_{t-1} + 0 \xi_{t-2} + 0$$
(4.5)

And the state transition equation—(4.5) in matrix form will be:

$$\begin{bmatrix} \xi_t \\ \xi_{t-1} \end{bmatrix} = \begin{bmatrix} \phi & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} \xi_{t-1} \\ \xi_{t-2} \end{bmatrix} + \begin{bmatrix} \varepsilon_t \\ 0 \end{bmatrix}$$
(4.6)

In comparison, for ARIMA (1, 1, 1), we just need to modify (4.3) as:

$$\Delta^{1d} y_t = \phi \Delta^{1d} y_{t-1} + \varepsilon_t + \theta \varepsilon_{t-1}$$
(4.7)

Modify (4.4) as:

$$\Delta^{1d} y_t = \xi_t + \theta \xi_{t-1} \tag{4.8}$$

Equation (4.5) and (4.6) are the same. We first build the filter, then use Maximum Likelihood (ML) to estimate the parameters of ARIMA (1, 1, 1) with the sample range of 1980 Q1-2006 Q1. The numerical results are demonstrated below:

(gretl scripts)

```
series y = diff(GDPC1Q)
/* parameter initalization */
phi = 0
theta = 0
sigma = 1
/* Kalman filter setup */
matrix H = {1; theta}
matrix F = {phi, 0; 1, 0}
matrix Q = {sigma^2, 0; 0, 0}
```

```
kalman
  obsy y
  obsymat H
  statemat F
  statevar Q
end kalman
/* maximum likelihood estimation */
mle logl = ERR ? NA : $kalman llt
  H[2] = theta
  F[1,1] = phi
  Q[1,1] = sigma^2
  ERR = kfilter()
  params phi theta sigma
end mle -h
(gretl results)
Using numerical derivatives
Tolerance = 1.81899e-012
Function evaluations: 120
Evaluations of gradient: 50
Model 1: ML, using observations 1980:1-2006:1 (T = 105)
logl = ERR ? NA : $kalman llt
Standard errors based on Hessian
       estimate std. error z
                                p-value
 _____
 phi
        0.975190 0.0252938 38.55 0.0000 ***
        -1.43617 0.231869 -6.194 5.87e-010 ***
 theta
 sigma
         38.3644 6.70031
                              5.726 1.03e-08 ***
Log-likelihood -570.6483 Akaike criterion
                                            1147.297
Schwarz criterion 1155.258 Hannan-Quinn
                                             1150.523
Also, we are able to get the results on the following scalar H, F, Q:
```

$$H = \begin{bmatrix} 1 \\ -1.43617 \end{bmatrix}; F = \begin{bmatrix} 0.97519 & 0 \\ 1 & 0 \end{bmatrix}; Q = \begin{bmatrix} 1471.79253 & 0 \\ 0 & 0 \end{bmatrix}$$

In conclusion, we can write:

$$\xi_{t+1} = \begin{bmatrix} 0.97519 & 0\\ 1 & 0 \end{bmatrix} \xi_t + v_t \tag{4.1'}$$

$$y_t = \begin{bmatrix} 1\\ -1.43617 \end{bmatrix} \xi_t \tag{4.2'}$$

where
$$E(\mathbf{v}_t \mathbf{v}'_s) = \begin{bmatrix} 1471.79253 & 0\\ 0 & 0 \end{bmatrix}$$
 for $t = s$, otherwise 0

And,

$$\Delta^{1d} y_t = 0.975190 \,\Delta^{1d} y_{t-1} + \varepsilon_t - 1.43617 \,\varepsilon_{t-1} \tag{4.7'}$$

$$\Delta^{1d} y_t = \xi_t - 1.43617\xi_{t-1} \tag{4.8'}$$

$$\begin{bmatrix} \xi_t \\ \xi_{t-1} \end{bmatrix} = \begin{bmatrix} 0.975190 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} \xi_{t-1} \\ \xi_{t-2} \end{bmatrix} + \begin{bmatrix} \varepsilon_t \\ 0 \end{bmatrix}$$
(4.6')

We find that ARIMA (1, 1, 1) of GDPC1Q without a constant has the same positive sign on φ and negative sign on θ , plus significant parameters, compared to ARIMA (1, 1, 1) with a constant.

(5) Transfer-Function and Intervention Models (VAR and ARMAX)

Sample range: 1980Q1-2006Q1

*Times series models with exogenous variables are also known as *transfer function models*.

VAR (1) modeled in section 3 is one kind of the intervention model. Recall that:

Let $y_{d_GDPC1Q,t}$ be $\Delta^{1d}y_{1,t} = y_{1t} - y_{1,t-1}$ and $y_{d_GCEC1Q,t}$ be $\Delta^{1d}y_{2,t} = y_{2t} - y_{2,t-1}$, then we have VAR (1) model in matrix form:

$$\Delta^{1d} y_{1,t} = 49.3131 + 0.360705 \Delta^{1d} y_{1,t-1} - 0.589345 \Delta^{1d} y_{2,t-1} + e_{1,t}$$
(3.4)
$$\Delta^{1d} y_{2,t} = 11.8267 - 0.0290698 \Delta^{1d} y_{1,t-1} + 0.000855439 \Delta^{1d} y_{2,t-1} + e_{2,t}$$

In the above model, the intervention government expenditure $(\Delta^{1d} y_{2,t-1})$ negatively affects GDP $(\Delta^{1d} y_{1,t})$ permanently and in return, $\Delta^{1d} y_{2,t}$ reacts negatively to a shock in $\Delta^{1d} y_{1,t-1}$ temporarily. Detailed impulse response analysis can be found in section 3.

Another kind of the intervention model is to add a regressor in Model 1 d_GDPC1Q_ARIMA (1, 1, 1) or Model 2 d_GCEC1Q_ARIMA (1, 1, 2), namely ARMAX. Recall that:

$$\Delta^{1d}Y_{1t} = c_1 + \varphi_1 \Delta^{1d}Y_{1,t-1} + \varepsilon_{1t} + \theta_1 \varepsilon_{1,t-1}$$
(1.16)

$$\Delta^{1d} Y_{2t} = c_2 + \varphi_1 \Delta^{1d} Y_{2,t-1} + \varepsilon_{2t} + \theta_1 \varepsilon_{2,t-1} + \theta_2 \varepsilon_{2,t-2}$$
(1.11)

Now modify equation (1.6) and (1.11) as follows:

$$\Delta^{1d} y_{1t} = c_1 + \varphi_1 \Delta^{1d} y_{1,t-1} + \xi_1 \Delta^{1d} y_{2t} + \varepsilon_{1t} + \theta_1 \varepsilon_{1,t-1}$$
(5.1)

$$\Delta^{1d} y_{2t} = c_2 + \varphi_1 \Delta^{1d} y_{2,t-1} + \xi_1 \Delta^{1d} y_{1t} + \varepsilon_{2t} + \theta_1 \varepsilon_{2,t-1} + \theta_2 \varepsilon_{2,t-2}$$
(5.2)

Equation (5.1) can be then estimated:

Model 5.1: ARMAX, using observations 1980:1-2006:1 (T = 105) Dependent variable: d_GDPC1Q Standard errors based on Hessian							
Coefficient Std. Error z p-value							
const	57.0849	11.2986	5.0524	< 0.00001	***		
phi_1	0.720079	0.137865	5.2231	< 0.00001	***		
theta_1	-0.383438	0.174517	-2.1971	0.02801	**		
d_GCEC1Q	0.922759	0.304911	3.0263	0.00248	***		
Mean dependent var	66.78		dependent var	58.36696			
Mean of innovations	0.777		S.D. of innovations		48378		
Log-likelihood	-562.9		Akaike criterion		1135.887		
Schwarz criterion	1149.	157 Hann	an-Quinn	1141.264			
	Real	Imaginary	Modulus	Frequency			
AR							
Root 1	1.3887	0.0000	1.3887	0.000	0		
MA							
Root 1	2.6080	0.0000	2.6080	0.000	0		

$$\Delta^{1d} y_{1t} = 57.0849 + 0.720079 \Delta^{1d} y_{1,t-1} + 0.922759 \Delta^{1d} y_{2t} + \varepsilon_{1t}$$
(5.3)
- 0.383438 \varepsilon_{1,t-1}

Frequency distribution for uhat11, obs 133-237 number of bins = 11, mean = 0.77744, sd = 52.613

interval	midpt	frequency	rel.	cum.	
< -143.79	-158.24	1	0.95%	0.95%	
-143.79114.89	-129.34	4	3.81%	4.76%	*
-114.8985.988	-100.44	2	1.90%	6.67%	
-85.98857.086	-71.537	6	5.71%	12.38% '	* *
-57.08628.185	-42.636	13	12.38%	24.76%	* * * *

-28.185 -	0.71708	-13.734	26	24.76%	49.52%	*******
0.71708 -	29.619	15.168	22	20.95%	70.48%	* * * * * * *
29.619 -	58.521	44.070	21	20.00%	90.48%	* * * * * * *
58.521 -	87.422	72.972	7	6.67%	97.14%	* *
87.422 -	116.32	101.87	1	0.95%	98.10%	
>=	116.32	130.78	2	1.90%	100.00%	

Test for null hypothesis of normal distribution: Chi-square(2) = 5.861 with p-value 0.05338

Equation (5.3) passes the normality test for the null hypothesis cannot be rejected with p-value of

0.05338.

Test for autocorrelation up to order 4

Ljung-Box Q' = 1.58435, with p-value = P(Chi-square(2) > 1.58435) = 0.4529 Model 5.1 passes the Ljung-Box autocorrelation test as the null hypothesis cannot be rejected

with p-value of 0.4529.


Test for ARCH of order 4


	coefficient	std. error	t-ratio	p-value		
alpha(0) alpha(1) alpha(2) alpha(3) alpha(4)	1295.21 0.0370509 0.229126 0.0790132 0.0875882	538.403 0.101975 0.101382 0.0885744 0.0889496	2.406 0.3633 2.260 0.8921 0.9847	0.0181 ** 0.7172 0.0261 ** 0.3746 0.3272		
Null hypothesis: no ARCH effect is present Test statistic: LM = 9.78853						

with p-value = P(Chi-square(4) > 9.78853) = 0.0441445

Model 5.1 has no ARCH effect because of p-value 0.0441445.

Regression residuals (= observed - fitted d_GDPC1Q)





Actual and fitted d_GDPC1Q versus d_GCEC1Q

The above graphs show a positive relationship between d_GDPC1Q and residual. Also, there seems to be a positive relationship between d_GDPC1Q and d_GCEC1Q concurrently, corresponding to d_GCEC1Q coefficient of 0.922759 with p-value 0.00248, which implies that GDP reacts very positively to changes in government spending at the same spatial time. This is exactly the way of how economic stimulus package works. But its long term effect on GDP is different as discussed in the VAR analysis.

Equation (5.2) can also be estimated:


Model 5	5.2: ARMAX, usin	g observations	1980:1-2006	(T = 105)			
Dependent variable: d GCEC1Q							
Standard errors based on Hessian							
	Coefficient	Std. Error	Z	p-value			
const	6.6791	3.01259	2.2171	0.02662	**		
phi_1	0.777622	0.132586	5.8650	< 0.00001	***		
theta_1	-0.862196	0.159308	-5.4121	< 0.00001	***		
theta_2	0.239026	0.127766	1.8708	0.06137	*		
d_GDPC1Q	0.0483137	0.0276438	1.7477	0.08051	*		

Mean dependent var	10.03	143	S.D. de	ependent var	15.92138
Mean of innovations	0.021	340	S.D. of	finnovations	15.03032
Log-likelihood	-433.6	534	Akaike	e criterion	879.3068
Schwarz criterion	895.2	305	305 Hannan-Quinn		885.7594
	ת 1	T			5
	Real	Imagi	nary	Modulus	Frequency
AR					
Root 1	1.2860		0.0000	1.2860	0.0000
MA					
Root 1	1.8036	_(0.9648	2.0454	-0.0782
Root 2	1.8036		0.9648	2.0454	0.0782


 $\Delta^{1d} y_{2t} = 6.6791 + 0.777622 \Delta^{1d} y_{2,t-1} + 0.0483137 \Delta^{1d} y_{1t} + \varepsilon_{2t}$ (5.4) - 0.862196\varepsilon_{2,t-1} + 0.239026\varepsilon_{2,t-2}

with p-value = P(Chi-square(4) > 2.56683) = 0.63271

Diagnostic checks out O.K. for Model 5.2. The model passes all the tests, with normally distribution residuals, without autocorrelation and ARCH effect.







Regression residuals (= observed - fitted d_GCEC1Q)

d_GCEC1Q shows positive relationship with residual. There's no obvious pattern how d_GCEC1Q reacts to d_GDPC1Q concurrently. But it appears that the fitted values is clustering around coordinates (70, 9). This conforms to the numerical results that d_GDPC1Q has little positive impact or significance (coefficient 0.0483137 with p-value 0.08051) on d_GCEC1Q, which also implies that the response of government expenditure to changes in GDP is minimal concurrently. But its long term impact on government spending is different as shown in the VAR analysis.

Equation (5.1) and (5.2) can be also modified as (5.5) and (5.6) respectively and estimated jointly as equation (5.7):

$$\Delta^{1d} y_{1t} = c_1 + \varphi_{1,1} \Delta^{1d} y_{1,t-1} + \xi_{1,1} \Delta^{1d} y_{2t} + \varepsilon_{1t} + \theta_{1,1} \varepsilon_{1,t-1}$$
(5.5)

$$\Delta^{1d} y_{2t} = c_2 + \varphi_{2,1} \Delta^{1d} y_{2,t-1} + \xi_{2,1} \Delta^{1d} y_{1t} + \varepsilon_{2t} + \theta_{2,1} \varepsilon_{2,t-1}$$
(5.6)

$$\begin{bmatrix} \Delta^{1d} y_{1t} \\ \Delta^{1d} y_{2t} \end{bmatrix} = \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} + \begin{bmatrix} \varphi_{1,1} & \varphi_{1,2} \\ \varphi_{2,1} & \varphi_{2,2} \end{bmatrix} \begin{bmatrix} \Delta^{1d} y_{1,t-1} \\ \Delta^{1d} y_{2,t-1} \end{bmatrix} + \begin{bmatrix} \xi_{1,1} & \xi_{1,2} \\ \xi_{2,1} & \xi_{2,2} \end{bmatrix} \begin{bmatrix} \Delta^{1d} y_{2t} \\ \Delta^{1d} y_{1t} \end{bmatrix} + \begin{bmatrix} \varepsilon_{1t} \\ \varepsilon_{2t} \end{bmatrix}$$

$$+ \begin{bmatrix} \theta_{1,1} & \theta_{1,2} \\ \theta_{2,1} & \theta_{2,2} \end{bmatrix} \begin{bmatrix} \varepsilon_{1,t-1} \\ \varepsilon_{2,t-1} \end{bmatrix}$$
Parameters to be estimated:
$$\begin{bmatrix} c_1 \\ c_2 \end{bmatrix}, \begin{bmatrix} \varphi_{1,1} & \varphi_{1,2} \\ \varphi_{2,1} & \varphi_{2,2} \end{bmatrix}, \begin{bmatrix} \xi_{1,1} & \xi_{1,2} \\ \xi_{2,1} & \xi_{2,2} \end{bmatrix}, \sigma^2 \left(\begin{bmatrix} \varepsilon_{1t} \\ \varepsilon_{2t} \end{bmatrix} \right), \begin{bmatrix} \theta_{1,1} & \theta_{1,2} \\ \theta_{2,1} & \theta_{2,2} \end{bmatrix}$$

Equation (5.7) can be considered Multivariate VARMAX process, which is a special case of B-J Multivariate VARMA.

.....

(6) Unit Root Tests (Augmented Dickey-Fuller test)

Sample range 1980Q1-2006Q1; Lag order 12 for ADF test.

For GDPC1Q Data:

```
Augmented Dickey-Fuller test for GDPC1Q
including 2 lags of (1-L)GDPC1Q
(max was 12, criterion modified AIC)
sample size 105
unit-root null hypothesis: a = 1
test with constant
model: (1-L)y = b0 + (a-1)*y(-1) + ... + e
1st-order autocorrelation coeff. for e: 0.020
lagged differences: F(2, 101) = 6.669 [0.0019]
estimated value of (a - 1): 0.0045083
test statistic: tau_c(1) = 1.68139
asymptotic p-value 0.9997
```

The test results indicate that GDPC1Q has unit root with significantly high asymptotic p-value 0.9997. This tells us that Model 1 data is non-stationary and we need to difference it at least once.

For GCEC1Q Data:

```
Augmented Dickey-Fuller test for GCEC1Q
including one lag of (1-L)GCEC1Q
(max was 12, criterion modified AIC)
sample size 105
unit-root null hypothesis: a = 1
test with constant
model: (1-L)y = b0 + (a-1)*y(-1) + ... + e
1st-order autocorrelation coeff. for e: 0.002
estimated value of (a - 1): -0.000652051
test statistic: tau_c(1) = -0.124135
asymptotic p-value 0.9451
```

The test results show that Model 2 GCEC1Q has unit root with relatively high asymptotic p-

value 0.9451, which means that Model 2 data is non-stationary and we have to difference it at

least once.

(7) Cointegration Test (Engle-Granger test) and Error Correction Model (if Contegration is *Present*)

Sample range 1980Q1-2006Q1; Variables GDPC1Q and GCEC1Q; Lag order 4.

Step 1: testing for a unit root in GDPC1Q Augmented Dickey-Fuller test for GDPC1Q including 2 lags of (1-L)GDPC1Q (max was 4, criterion modified AIC) sample size 100 unit-root null hypothesis: a = 1 test with constant model: $(1-L)y = b0 + (a-1)*y(-1) + \dots + e$ 1st-order autocorrelation coeff. for e: 0.062 lagged differences: F(2, 96) = 8.226 [0.0005]estimated value of (a - 1): 0.00430732 test statistic: tau c(1) = 1.66308asymptotic p-value 0.9996 Step 2: testing for a unit root in GCEC1Q Augmented Dickey-Fuller test for GCEC1Q including one lag of (1-L)GCEC10 (max was 4, criterion modified AIC) sample size 100 unit-root null hypothesis: a = 1

```
test with constant
   model: (1-L)y = b0 + (a-1)*y(-1) + \dots + e
   1st-order autocorrelation coeff. for e: 0.006
   estimated value of (a - 1): -0.00249536
   test statistic: tau c(1) = -0.439981
   asymptotic p-value 0.9
Step 3: cointegrating regression
Cointegrating regression -
OLS, using observations 1980:1-2006:1 (T = 105)
Dependent variable: GDPC1Q
             coefficient std. error t-ratio p-value
  _____
  const -4023.80 312.349 -12.88 3.38e-023 ***
  GCEC1Q
            6.88632 0.165023 41.73 2.38e-066 ***
Mean dependent var 8846.749 S.D. dependent var 2128.680
Sum squared resid 26317802 S.E. of regression 505.4826
R-squared0.944154Adjusted R-squared0.943611Log-likelihood-801.6578Akaike criterion1607.316Schwarz criterion1612.624Hannan-Quinn1609.467rho0.972847Durbin-Watson0.049978
Step 4: testing for a unit root in uhat
Augmented Dickey-Fuller test for uhat
including 3 lags of (1-L)uhat
(max was 4, criterion modified AIC)
sample size 100
unit-root null hypothesis: a = 1
   model: (1-L)y = (a-1)*y(-1) + \ldots + e
   1st-order autocorrelation coeff. for e: -0.026
   lagged differences: F(3, 96) = 4.967 [0.0030]
   estimated value of (a - 1): -0.0468461
   test statistic: tau c(2) = -2.16499
   asymptotic p-value 0.4417
There is evidence for a cointegrating relationship if:
(a) The unit-root hypothesis is not rejected for the individual variables.
(b) The unit-root hypothesis is rejected for the residuals (uhat) from the
    cointegrating regression.
Test results with constant indicate that variables GDPC1Q and GCEC1Q are not cointegrated,
because a) The unit-root hypothesis is not rejected for the individual variables; b) However, the
cointegrating regression has unit root with 0.4417 p-value, therefore the unit-root hypothesis is
NOT rejected for the residuals (uhat) from the cointegrating regression. The estimated
cointegrating system is shown in step 3.
```

(8) Volatility Tests by the ARCH, GARCH, ARCH-M, GARCH-M, EGARCH and other GARCH variants

Sample (full) range: 1947Q1-2013Q1

In order to obtain comparable results (AIC, BIC, HQC), we use 100*ldiff(GDPC1Q) [100 times log first difference of GDPC1Q] for all ARCH, GARCH, ARCH-M, GARCH-M, TS-GARCH, GJR, TARCH, NARCH, APARCH, EGARCH. Let treated endogenous variable 100*ldiff(GDPC1Q) be y_t .

Model 8.1: ARCH (1)

I. Identification:

LM tests show that ARIMA (1, 1, 1) of GDPC1Q has ARCH effect with p-value of 0.0147128, however ARIMA (1, 1, 2) of GCEC1Q does not has such effect with p-value of 0.611714. Therefore, dependent variable GDPC1Q will be tested for volatility.

II. Estimation and Diagnostic Checking:

Model 8.1: ARCH (1), using observations 1947:2-2013:1 (T = 264)								
	Dependent variable: 100*ldiff(GDPC1Q)							
	Standard	errors ba	ased on	Hessian				
	Coefficient	Std. E	rror	Z	p-value			
const	0.798974	0.058	7239	13.6056	< 0.00001	***		
alpha(0)	0.632159	0.0764	4556	8.2683	< 0.00001	***		
alpha(1)	0.36451	0.109	357	3.3332	0.00086	***		
Mean dependent var	0.77	6391	S.D. 0	dependent var	0.9	83733		
Log-likelihood	-357.	1958	Akaik	criterion	722	2.3915		
Schwarz criterion	736.	6953	Hanna	an-Quinn	728	8.1392		

Unconditional error variance = 0.994759

$$y_t = 0.798974 + e_t \tag{8.1}$$

$$e_t | I_{t-1} \sim N(0, h_t)$$
 (8.2)

$$h_t = 0.632159 + 0.36451e_{t-1}^2 \tag{8.3}$$

Frequency distribution for uhat11, obs 2-265 number of bins = 17, mean = -0.0225837, sd = 0.987495


interval	midpt	frequency	rel.	cum.
< -3.3342	-3.5438	1	0.38%	0.38%
-3.33422.9151	-3.1247	1	0.38%	0.76%
-2.91512.4960	-2.7055	1	0.38%	1.14%
-2.49602.0768	-2.2864	5	1.89%	3.03%

-2.0768 -	-1.6577	-1.8673	9	3.41%	6.44%	*
-1.6577 -	-1.2385	-1.4481	9	3.41%	9.85%	*
-1.2385 -	-0.81940	-1.0290	14	5.30%	15.15%	*
-0.81940 -	-0.40027	-0.60984	39	14.77%	29.92%	* * * * *
-0.40027 -	0.018873	-0.19070	67	25.38%	55.30%	*******
0.018873 -	0.43801	0.22844	44	16.67%	71.97%	* * * * * *
0.43801 -	0.85715	0.64758	27	10.23%	82.20%	* * *
0.85715 -	1.2763	1.0667	28	10.61%	92.80%	* * *
1.2763 -	1.6954	1.4859	10	3.79%	96.59%	*
1.6954 -	2.1146	1.9050	4	1.52%	98.11%	
2.1146 -	2.5337	2.3241	2	0.76%	98.86%	
2.5337 -	2.9528	2.7433	0	0.00%	98.86%	
>=	2.9528	3.1624	3	1.14%	100.00%	
Test for nul	l hypothes	is of norm	al distribu	tion:		
Chi-square(2	= 19.654	with p-va	lue 0.00005			

Normality (Doornik-Hansen) test of H_0 is rejected with extremely small p-value 0.00005.

Residual definitely is not normally distributed.

III. Analysis:



The residual plot shows approximately 33 GDP spikes outside of the blue upper and lower boundaries. There are large spikes occurring around the year of 1949, 1950, 1951, 1953, 1954,

1955, 1958, 1961, 1978, 1980, 1982, 1991, and 2009. From a historian point of view, the analysis accurately catches behaviors of the economy during major recessions, for example 1981-1982, 1990-1991, 2001, especially the great recession 2008. More interestingly, the analysis also sends out ex-ante and ex-post warning signals to policy makers. For example, 1980, just before 1981-1982 recession shows volatility (huge downward spike) in GDP followed by 1983, 1984. Additionally, 1988, 2000, and 2007 exhibit instable economic situation just before several major recessions. In conclusion, proper volatility analysis can help the government not only monitor economy but also prevent major recessions through proper economic policy adjustment.

Model 8.2: GARCH (1, 1)

Model 8.2: GARCH (1, 1), using observations 1947:2-2013:1 (T = 264)						
	Dependent va	riable: 1	l00*ldi	ff(GDPC1Q)		
	Standard	errors b	ased on	Hessian		
	Coefficient	Std. E	Error	Z	p-value	
const	0.820552	0.055	6729	14.7388	< 0.00001	***
alpha(0)	0.0385705	0.02	223	1.7351	0.08273	*
alpha(1)	0.255331	0.074	9903	3.4048	0.00066	***
beta(1)	0.734709	0.060	7463	12.0947	< 0.00001	***
Mean dependent var Log-likelihood	0.77 -346.			lependent var te criterion		83733 8.7286
Schwarz criterion		6084		an-Quinn).9132

Unconditional error variance = 3.87253

The GARCH (1, 1) model can be expressed as:


$$y_t = 0.820552 + e_t \tag{8.4}$$

$$e_t | I_{t-1} \sim N(0, h_t)$$
 (8.5)

$$h_t = 0.0385705 + 0.255331e_{t-1}^2 + 0.734709h_{t-1}$$
(8.6)

GARCH (1, 1) shows definite nonnormality with p value of 0.00005, same as ARCH (1).

III. Analysis:



The residual plot of GARCH (1, 1) is almost identical to ARCH (1), but slightly different from it. For instance, GARCH (1, 1)'s upper and lower boundaries are more adaptive and volatile than ARCH (1)'s, therefore GARCH includes all the traits of ARCH, but also catches what ARCH misses. In other words, GARCH is more generalized than ARCH and thus it reflects volatility more accurately.

Model 8.3: ARCH-M (1)

```
(gretl scripts)
series y = 100*1diff(GDPC1Q)
scalar mu = 0.0
scalar delta = 0.04
scalar alpha = 0.4
scalar beta = 0.0
scalar theta = 0.0001
```

```
mle ll = -0.5*(\log(h) + (e^2)/h)
 series h = var(y)
 series e = y - theta*h
 series h = delta + alpha*(e(-1))^2
 params theta delta alpha
end mle
(gretl script results)
Using numerical derivatives
Tolerance = 1.81899e-012
Function evaluations: 214
Evaluations of gradient: 57
Model 19: ML, using observations 1947:3-2013:1 (T = 263)
ll = -0.5*(log(h) + (e^2)/h)
Standard errors based on Outer Products matrix
       estimate std. error z
                                p-value
 _____
        0.829080 0.0566959 14.62 2.00e-048 ***
 theta
        0.632092 0.0602920 10.48 1.02e-025 ***
 delta
         0.365035 0.102237 3.570 0.0004 ***
 alpha
Log-likelihood -114.1423 Akaike criterion
                                             234.2847
Schwarz criterion 245.0011 Hannan-Quinn
                                               238.5914
                                     y_t = \theta h_t + e_t
                                    h_t = \delta + \alpha_1 e_{t-1}^2
```

Input the parameters into equation 8.7 and 8.8, we get:

$$y_t = 0.82908h_t + e_t \tag{8.9}$$

(8.7)

(8.8)

$$h_t = 0.632092 + 0.365035e_{t-1}^2 \tag{8.10}$$

Model 8.4: GARCH-M (1, 1)

(gretl scripts) series y = 100*ldiff(GDPC1Q) scalar mu = 0.0 scalar delta = 0.04

(8.11)

(8.12)

(8.13)

(8.14)

scalar alpha = 0.4scalar beta = 0.5scalar theta = 0.0001mle ll = $-0.5*(\log(h) + (e^2)/h)$ series h = var(y)series e = y - theta*h series $h = delta + alpha*(e(-1))^2 + beta*h(-1)$ params theta delta alpha beta end mle (gretl script results) Model 13: ML, using observations 1947:2-2013:1 (T = 264) $ll = -0.5*(log(h) + (e^2)/h)$ Standard errors based on Outer Products matrix estimate std. error z p-value 0.847852 0.0563744 15.04 4.04e-051 *** theta 0.0386463 0.0211140 1.830 0.0672 * delta 0.255461 0.0558469 4.574 4.78e-06 *** alpha 0.734511 0.0412182 17.82 4.94e-071 *** beta Log-likelihood -104.2686 Akaike criterion 216.5371 Schwarz criterion 230.8409 Hannan-Quinn 222.2848 $y_t = \theta h_t + e_t$ $h_t = \delta + \alpha_1 e_{t-1}^2 + \beta_1 h_{t-1}$ Input the parameters into equation 8.11 and 8.12, we get: $y_t = 0.847852h_t + e_t$ $h_t = 0.0386463 + 0.255461e_{t-1}^2 + 0.734511h_{t-1}$ Model 8.5: Taylor-Schwert GARCH Model: Taylor/Schwert's GARCH(1,1) (Normal) Dependent variable: y

Conditional mean equation

	coefficient	std. error	Z	p-value	
const	0.836373	0.00279419	299.3	0.0000	* * *

Sample: 1947:2-2013:1 (T = 264), VCV method: Hessian

Conditional	variance	equation
Conditional	variance	equatior

	coefficient	std. error	Z	p-value	
omega alpha beta	0.0662126 0.252868 0.738908	0.0345328 0.0588503 0.0643768	1.917 4.297 11.48	0.0552 1.73e-05 1.70e-030	* * * * * * *
Llik: BIC:	-345.48651 713.27682		98.97303 04.72073		

Model 8.6: GJR (The Glosten Jagannathan Runkle GARCH)

Model: GJR(1,1) [Glosten et al.] (Normal)*
Dependent variable: y
Sample: 1947:2-2013:1 (T = 264), VCV method: OPG

Conditional mean equation

	coefficient	std. error	Z	p-value
const	0.850098	0.0518322	16.40	1.88e-060 ***

Conditional variance equation

	coefficient	std. error	Z	p-value	
omega	0.0858183	0.0455942	1.882	0.0598	*
alpha	0.341302	0.100939	3.381	0.0007	* * *
gamma	0.313410	0.114741	2.731	0.0063	* * *
beta	0.592662	0.0887584	6.677	2.43e-011	* * *

(alt. parametrization)

	coefficient	std. error	z z	p-value	
delta alpha gamma beta	0.0858183 0.160892 0.427870 0.592662	0.0455942 0.0863810 0.150627 0.0887584	1.882 1.863 2.841 6.677	0.0598 0.0625 0.0045 2.43e-011	* * * * *
Llik: BIC:	-344.38603 716.65180		98.77205 05.95668		

Model 8.7: TARCH

Model: TARCH(1,1) [Zakoian] (Normal)
Dependent variable: y
Sample: 1947:2-2013:1 (T = 264), VCV method: Hessian

Conditional mean equation

	coefficient	std. error	Z	p-value	
const	0.830636	0.0490712	16.93	2.84e-064	* * *

Conditional variance equation

coefficient std. error z p-value

 omega
 0.107087
 0.0500759
 2.138
 0.0325
 **

 alpha
 0.284205
 0.0661465
 4.297
 1.73e-05

 gamma
 0.363679
 0.130245
 2.792
 0.0052

 beta
 0.664701
 0.0864832
 7.686
 1.52e-014

 Llik:
 -341.62854
 AIC:
 693.25708
 BIC:
 711.13683
 HQC:
 700.44171

Model 8.8: NARCH

Model: NARCH(1,1) [Higgins and Bera] (Normal)
Dependent variable: y
Sample: 1947:2-2013:1 (T = 264), VCV method: Hessian

Conditional mean equation

coefficient std. error z p-value const 0.843591 3.95938e-05 2.131e+04 0.0000 ***

Conditional variance equation

	coefficient	std. error	c z	p-value	
omega alpha beta delta	0.0811524 0.156068 0.779823 0.246768	0.0349671 0.0728736 0.0701164 0.282203	2.321 2.142 11.12 0.8744	0.0203 0.0322 9.82e-029 0.3819	* * * * * * *
Llik: BIC:	-345.06132 718.00239		00.12265 07.30727		

Model 8.9: APARCH

Model: APARCH(1,1) [Ding] (Normal)
Dependent variable: y
Sample: 1947:2-2013:1 (T = 264), VCV method: Hessian

Conditional mean equation

	coefficient	std. error	Z	p-value	
const	0.829406	0.0553688	14.98	9.97e-051	* * *

Conditional variance equation

	coefficient	std. erro	or z	p-value	
omega alpha gamma beta delta	0.112231 0.247851 0.387464 0.687873 0.690535	0.0510380 0.0790212 0.142873 0.0880695 0.419295	3.137 2.712	0.0279 0.0017 0.0067 5.69e-015 0.0996	* * * * * * * * *
Llik: BIC:	-341.37515 716.20599		694.75029 703.37184		

Model 8.10: EGARCH (1, 1)

Model: EGARCH(1,1) [Nelson] (Normal)
Dependent variable: y
Sample: 1947:2-2013:1 (T = 264), VCV method: Hessian

Conditional mean equation

	coefficient	std. error	Z	p-value	
const	0.836103	0.0500604	16.70	1.27e-062 **	* *

Conditional variance equation

	coefficient	std. erro	r z	p-value	
omega alpha gamma beta	-0.464418 0.541685 -0.196858 0.843356	0.0937466 0.110848 0.0756444 0.0583697	4.887 -2.602	7.27e-07 1.03e-06 0.0093 2.56e-047	* * * * * * * * *
Llik: BIC:	-340.43681 708.75337		690.87362 698.05825		

Goodness-of-Fit Comparison

Model Criteria	AIC	BIC	HQC
ARCH (1)*	722.3915	736.6953	728.1392
GARCH (1, 1)*	703.7286	721.6084	710.9132
ARCH-M (1)*	234.2847	245.0011	238.5914
GARCH-M (1, 1)*	216.5371	230.8409	222.2848
TS-GARCH (1, 1)*	698.97303	713.27682	704.72073
GJR (1, 1)*	698.77205	716.65180	705.95668
TARCH (1, 1)*	693.25708	711.13683	700.44171
NARCH (1, 1)	700.12265	718.00239	707.30727
APARCH (1, 1)*	694.75029	716.20599	703.37184
EGARCH (1, 1)*	690.87362	708.75337	698.05825

Residuals are normally distributed.

* denotes that parameters (all the coefficients) are significant.

Discussion of the Best Volatility Model

From the above goodness of fit comparison chart, we learn that GARCH-M (1, 1) definitely has the lowest AIC, BIC, HQC. For a more comprehensive comparison, we now assume that the normality assumption of residual distribution is violated and those distributions (t, GED—

Generalized Error Distribution, skewed t, skewed GED) will be used to construct comparable models. The general technique for obtaining the best volatility model is to experiment with models of nonnormal distribution that results in the lowest possible overall fitness criterion, which we compare with those of normal distribution with the lowest criterion. More specifically, the algorithm of obtaining the best volatility model is to first perform self-comparison within the models of nonnormal distribution; second perform cross-comparison between the models of nonnormal distribution and those of normal distribution. (The reason to first perform self-comparison within the models of nonnormal distribution, as results shown in the following table). For example, first compare fitness criterion among t, GED, Skewed t, Skewed GED, get the lowest result; second compare the model of nonnormal distribution with the lowest fitness criterion with its counterpart of normal distribution, get the lowest result as the best fitted model. Please see the following experiment results:

	AIC	BIC	HQC
<u>ARCH (1)</u>			
t*	712.58939	726.89319	718.33709
GED*	706.94010	721.24390	712.68780
Skewed t	714.58926	732.46901	721.77389
Skewed GED	N/A	N/A	N/A
<u>GARCH (1, 1)</u>			
t	694.76479	712.64453	701.94941
GED	691.90814	709.78789	699.09277
Skewed t	696.76243	718.21812	705.38398
Skewed GED	693.86115	715.31684	702.48270
<u>TS-GARCH (1, 1)</u>			
t*	693.75727	711.63702	700.94190
GED	691.12038	709.00012	698.30500
Skewed t	695.72381	717.17951	704.34536
Skewed GED	692.97543	714.43113	701.59698
<u>GJR (1, 1)</u>			

t*	694.17832	715.63401	702.79987
GED	691.54820	713.00390	700.16975
Skewed t	695.53312	720.56476	705.59159
Skewed GED	692.19620	717.22785	702.25468
<u>TARCH (1, 1)</u>			
t	692.16754	713.62323	700.78909
GED*	689.33261	710.78830	697.95416
Skewed t	693.37396	718.40561	703.43244
Skewed GED	689.77211	714.80376	699.83059
<u>NARCH (1, 1)</u>			
t	695.74212	717.19781	704.36367
GED	693.08054	714.53624	701.70209
Skewed t	697.72246	722.75410	707.78094
Skewed GED	694.97241	720.00406	705.03089
<u>APARCH (1, 1)</u>			
t	694.15953	719.19118	704.21801
GED*	691.33184	716.36349	701.39032
Skewed t	695.35481	723.96241	706.85022
Skewed GED	691.70526	720.31285	703.20066
<u>EGARCH (1, 1)</u>			
t	690.32769	711.78338	698.94924
GED	N/A	N/A	N/A
Skewed t	690.99794	716.02958	701.05642
Skewed GED	N/A	N/A	N/A

For the purpose of comparison, models with nonnormal distribution use the same covariance estimator as those with normal distribution. Results seem not to vary significantly from using different covariance estimation (Sandwich, Hessian, OPG).

For the purpose of comparable results, models with nonnormal distribution (ARCH, GARCH, TS-GARCH, GJR, TARCH, NARCH, APARCH, EGARCH) use 100*ldiff(GDPC1Q).

N/A means that calculation error occurs for associated distribution to estimate the corresponding model.

* means that parameters (all the coefficients) are significant.

From the above experiment results, we can find that GED (General Error Distribution) works the best for all the models of nonnormal distribution with the lowest overall fitness criterion. TARCH (1, 1) GED is the best model within the nonnormal group. APARCH (1, 1) GED is the second. TS-GARCH (1, 1) t and GJR (1, 1) t stands the third for the former has slightly lower overall fitness criterion than the latter. On the other hand, within the group of normal distribution, we find that GARCH-M (1, 1) and ARCH-M (1, 1) are very different from the rest with the lowest AIC, BIC, HQC in the 200 range. GARCH-M is slightly better than ARCH-M. In the 700 range, EGARCH is the first; TARCH and APARCH are the second (TARCH is slightly better than APARCH); TS-GARCH and GJR are the third (TS-GARCH is slightly better than GJR). Therefore, we conclude that GARCH-M and ARCH-M models are the best for volatility test of GDPC1Q. TARCH (1, 1) GED is the second best. EGARCH (1, 1) takes the third.

SECTION III: DISCUSSION

1) Summary of Major Research Findings and Their Managerial Implications

GDP—GDPC1Q (Y_{1t}), Government Expenditure—GCEC1Q (Y_{2t})

Sample range:1980Q1—2006Q1 for (i)—(vii); Sample range:1947Q1-2013Q1 for (viii)

(i) Univariate ARIMA

$$\Delta^{1d} Y_{1t} = 66.2629 + 0.759918 \Delta^{1d} Y_{1,t-1} + \varepsilon_{1t} - 0.479305 \varepsilon_{1,t-1}$$
(1.10)

$$\Delta^{1d} Y_{2t} = 9.7675 + 0.811342 \Delta^{1d} Y_{2,t-1} + \varepsilon_{2t} - 0.943283 \varepsilon_{2,t-1}$$
(1.15)
+ 0.273345 \varepsilon_{2,t-2}

Through Univariate ARMA analysis, we find that each observable value for GDP and Government Expenditure of current year are significantly positively associated with the value of past year, meaning that the future trend of both GDP and Government Spending can be accurately predicted via their past pattern, which is critical for policy makers to monitor and control the direction of overall economy. The forecasting results show a promising bouncing back trend of both GDP and Government Expenditure.

(ii) Multivariate ARIMA

The VARIMA (1, 1, 1) and VARIMA (1, 1, 2) analysis does not give us any useful insights regarding the interrelationship between GDP and Government Expenditure and between GDP because of statistically insignificant corresponding

coefficients. However, we do find that both GDPC1Q and GCEC1Q from current year reacts positively to those from previous year, which is consistent with the findings from the univariate ARIMA analysis.

(iii) VAR and Impulse Response Function

$$\begin{bmatrix} \Delta^{1d} y_{1,t} \\ \Delta^{1d} y_{2,t} \end{bmatrix} = \begin{bmatrix} 49.3131 \\ 11.8267 \end{bmatrix} + \begin{bmatrix} 0.360705 & -0.589345 \\ -0.0290698 & 0.000855439 \end{bmatrix} \begin{bmatrix} \Delta^{1d} y_{1,t-1} \\ \Delta^{1d} y_{2,t-1} \end{bmatrix} + \begin{bmatrix} e_{1,t} \\ e_{2,t} \end{bmatrix}$$
(3.3)

$$\Delta^{1d} y_{1,t} = 49.3131 + 0.360705 \Delta^{1d} y_{1,t-1} - 0.589345 \Delta^{1d} y_{2,t-1} + e_{1,t}$$
(3.4)
$$\Delta^{1d} y_{2,t} = 11.8267 - 0.0290698 \Delta^{1d} y_{1,t-1} + 0.000855439 \Delta^{1d} y_{2,t-1} + e_{2,t}$$

Through VAR and impulse response, we find that the negative response of GDP from current year to a shock of Government Expenditure from last year is statistically significant, while the negative response of Government Spending from current year to a shock of GDP from last year is statistically insignificant. For example, a column of insignificant coefficients means that the empirical shocks of the corresponding variable have only exhibited temporary effects on the variables of the system, whereas a column of significant coefficients indicates permanent effects on the system (Katarina Juselius "The Cointegrated VAR Model: Methodology and Applications"). Based on Juselius's finding, we conclude the VAR analysis that a shock of lastyear Government Expenditure on current-year GDP is not only negative but also permanent, while a shock of last-year GDP on current-year Government Expenditure is not necessarily negative but also temporary implies that substantially large government expenditure from the past will negatively impact GDP in the long-run, while temporary increase of government spending alone in the future might not necessarily be the only answer to a weak economy from the past. Thereby, we suggest that policy makers should be cautious about boosting economy via increasing government expenditure alone, rather government spending should be adjusted accordingly with consideration of GDP not only in the short-run but in the long-run as well. (iv) State-Space Model of ARMA via the Kalman Filter and Maximum Likelihood Estimation

$$\xi_{t+1} = \begin{bmatrix} 0.97519 & 0\\ 1 & 0 \end{bmatrix} \xi_t + v_t \tag{4.1'}$$

$$y_t = \begin{bmatrix} 1\\ -1.43617 \end{bmatrix} \xi_t \tag{4.2'}$$

where
$$E(\mathbf{v}_t \mathbf{v}'_s) = \begin{bmatrix} 1471.79253 & 0\\ 0 & 0 \end{bmatrix}$$
 for $t = s$, otherwise 0

And,

$$\Delta^{1d} y_t = 0.975190 \,\Delta^{1d} y_{t-1} + \varepsilon_t - 1.43617 \,\varepsilon_{t-1} \tag{4.7'}$$

$$\Delta^{1d} y_t = \xi_t - 1.43617\xi_{t-1} \tag{4.8'}$$

$$\begin{bmatrix} \xi_t \\ \xi_{t-1} \end{bmatrix} = \begin{bmatrix} 0.975190 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} \xi_{t-1} \\ \xi_{t-2} \end{bmatrix} + \begin{bmatrix} \varepsilon_t \\ 0 \end{bmatrix}$$
(4.6')

We find that ARIMA (1, 1, 1) of GDPC1Q without a constant has the same positive sign on φ and negative sign on θ , plus significant parameters, compared to ARIMA (1, 1, 1) with a constant, meaning that there is a statistically significant positive relationship between last year GDP and current year GDP. This provides policy makers with a very accurate yet powerful tool to monitor, control and predict the direction of overall economy.

(v) Transfer Function and Intervention Model—ARMAX

$$\Delta^{1d} y_{1t} = 57.0849 + 0.720079 \Delta^{1d} y_{1,t-1} + 0.922759 \Delta^{1d} y_{2t} + \varepsilon_{1t}$$
(5.3)
- 0.383438 \varepsilon_{1,t-1}

$$\Delta^{1d} y_{2t} = 6.6791 + 0.777622 \Delta^{1d} y_{2,t-1} + 0.0483137 \Delta^{1d} y_{1t} + \varepsilon_{2t}$$
(5.4)
- 0.862196\varepsilon_{2,t-1} + 0.239026\varepsilon_{2,t-2}

Through ARMAX results, we find that both concurrent relationship between Government Expenditure and GDP and between GDP and Government Expenditure are statistically significantly positive, meaning that by increasing current-year Government Spending, policy makers can boost current-year GDP, while strong current-year GDP stimulates increasing current-year Government Expenditure in return, which is consistent with the classic Keynesian theory that increasing government investment in infrastructure can boost the economy in the short run.

(vi) Unit Root Test (ADF Test)

The ADF tests show that sample data of both GDPC1Q and GCEC1Q are not stationary, meaning that we have to difference the sample data at least for once in order for its suitability for further ARMA analysis.

(vii) Cointegration Test (Engle-Granger test)

The Engle-Granger tests show no sign of conintegration between GDPC1Q and GCEC1Q, meaning that it is not necessary for us to use error-corrected VAR (VEC).

(viii) Volatility Tests via ARCH, GARCH, ARCH-M, GARCH-M, EGARCH, and other ARCH GARCH variants

We find that the classic GARCH-M performs the best for GDPC1Q data assuming normal distribution, while TARCH (1, 1) GED performs the best for GDPC1Q data assuming non-normal distribution. Overall, models with non-normal distribution have lower AIC, BIC, HQC than models with normal distribution, excluding ARCH-M and GARCH-M. The results explicitly suggest the best volatility model to policy makers and practitioners for analyzing GDP with either normal or non-normal distribution.

2) Comparison of Empirical Results

(i) Univariate vs. Multivariate VARMA

 $\begin{aligned} & ARIMA (1, 1, 1) \ GDPC1Q: \\ & \Delta^{1d}Y_{1t} = 66.2629 + 0.759918\Delta^{1d}Y_{1,t-1} + \varepsilon_{1t} - 0.479305\varepsilon_{1,t-1} \\ & ARIMA (1, 1, 2) \ GCEC1Q: \\ & \Delta^{1d}Y_{2t} = 9.7675 + 0.811342\Delta^{1d}Y_{2,t-1} + \varepsilon_{2t} - 0.943283\varepsilon_{2,t-1} \end{aligned}$ (1.15)

$$+ 0.273345\varepsilon_{2,t-2}$$

$$VARIMA (1, 1, 1) GDPC1Q & GCEC1Q: \begin{bmatrix} \Delta^{1d}Y_{1t} \\ \Delta^{1d}Y_{2t} \end{bmatrix} = \begin{bmatrix} 82.2102 \\ 8.59125 \end{bmatrix} + \begin{bmatrix} 0.751179 & -6.43447 \\ 0.060149 & -0.261946 \end{bmatrix} \begin{bmatrix} \Delta^{1d}Y_{1,t-1} \\ \Delta^{1d}Y_{2,t-1} \end{bmatrix} + \begin{bmatrix} \varepsilon_{1,t} \\ \varepsilon_{2,t} \end{bmatrix} + \begin{bmatrix} -0.460145 & 5.91783 \\ -0.105878 & 0.279512 \end{bmatrix} \begin{bmatrix} \varepsilon_{1,t-1} \\ \varepsilon_{2,t-1} \end{bmatrix}$$

$$\Delta^{1d}Y_{1t} = 82.2102 + 0.751179\Delta^{1d}Y_{1,t-1} - 6.43447\Delta^{1d}Y_{2,t-1} + \varepsilon_{1,t} - 0.460145\varepsilon_{1,t-1} + 5.91783\varepsilon_{2,t-1}$$

$$\Delta^{1d}Y_{1t} = 82.2102 + 0.751179\Delta^{1d}Y_{1,t-1} - 6.43447\Delta^{1d}Y_{2,t-1} + \varepsilon_{1,t} - 0.460145\varepsilon_{1,t-1} + 5.91783\varepsilon_{2,t-1}$$

$$\Delta^{1d}Y_{1t} = 82.2102 + 0.000140\Delta^{1d}Y_{1,t-1} - 6.43447\Delta^{1d}Y_{2,t-1} + \varepsilon_{1,t} - 0.460145\varepsilon_{1,t-1} + 5.91783\varepsilon_{2,t-1}$$

$$\Delta^{1d} Y_{2t} = 8.59125 + 0.060149 \Delta^{1d} Y_{1,t-1} - 0.261946 \Delta^{1d} Y_{2,t-1} + \varepsilon_{2,t} - 0.105878 \varepsilon_{1,t-1} + 0.279512 \varepsilon_{2,t-1}$$

VARIMA (1, 1, 2) GDPC1Q & GCEC1Q:

$$\begin{bmatrix} \Delta^{1d} Y_{1t} \\ \Delta^{1d} Y_{2t} \end{bmatrix} = \begin{bmatrix} 46.7843 \\ 0.156603 \end{bmatrix} + \begin{bmatrix} 0.369566 & -0.219727 \\ 0.00582349 & 0.950556 \end{bmatrix} \begin{bmatrix} \Delta^{1d} Y_{1,t-1} \\ \Delta^{1d} Y_{2,t-1} \end{bmatrix} + \begin{bmatrix} \varepsilon_{1,t} \\ \varepsilon_{2,t} \end{bmatrix}$$

$$+ \begin{bmatrix} -0.113708 & -0.132601 \\ -0.0394123 & -1.00433 \end{bmatrix} \begin{bmatrix} \varepsilon_{1,t-1} \\ \varepsilon_{2,t-1} \end{bmatrix}$$

$$+ \begin{bmatrix} 0.216516 & -0.104326 \\ 0.0111816 & 0.165578 \end{bmatrix} \begin{bmatrix} \varepsilon_{1,t-2} \\ \varepsilon_{2,t-2} \end{bmatrix}$$

$$(2.3')$$

$$\begin{split} \Delta^{1d}Y_{1t} &= 46.7843 + 0.369566\Delta^{1d}Y_{1,t-1} - 0.219727\Delta^{1d}Y_{2,t-1} + \varepsilon_{1,t} \\ &\quad -0.113708\varepsilon_{1,t-1} - 0.132601\varepsilon_{2,t-1} + 0.216516\varepsilon_{1,t-2} \\ &\quad -0.104326\varepsilon_{2,t-2} \end{split} \tag{2.4'}$$

$$\Delta^{1d}Y_{2t} &= 0.156603 + 0.00582349\Delta^{1d}Y_{1,t-1} + 0.950556\Delta^{1d}Y_{2,t-1} + \varepsilon_{2,t} \\ &\quad -0.0394123\varepsilon_{1,t-1} - 1.00433\varepsilon_{2,t-1} + 0.0111816\varepsilon_{1,t-2} \end{split}$$

 $+ 0.165578\varepsilon_{2,t-2}$

VARI (1, 1):

$$\begin{bmatrix} \Delta^{1d} y_{1,t} \\ \Delta^{1d} y_{2,t} \end{bmatrix} = \begin{bmatrix} 49.3131 \\ 11.8267 \end{bmatrix} + \begin{bmatrix} 0.360705 & -0.589345 \\ -0.0290698 & 0.000855439 \end{bmatrix} \begin{bmatrix} \Delta^{1d} y_{1,t-1} \\ \Delta^{1d} y_{2,t-1} \end{bmatrix} + \begin{bmatrix} e_{1,t} \\ e_{2,t} \end{bmatrix}$$
(3.3)

$$\Delta^{1d} y_{1,t} = 49.3131 + 0.360705 \Delta^{1d} y_{1,t-1} - 0.589345 \Delta^{1d} y_{2,t-1} + e_{1,t}$$
(3.4)
$$\Delta^{1d} y_{2,t} = 11.8267 - 0.0290698 \Delta^{1d} y_{1,t-1} + 0.000855439 \Delta^{1d} y_{2,t-1} + e_{2,t}$$

The coefficient value of $\Delta^{1d} Y_{1,t-1}$ from the first equation in (2.2') is remarkably similar to its counterpart in (1.10) (0.751179 compared with 0.759918), plus both coefficients having the same positive sign. In contrast, the coefficient value of $\Delta^{1d}Y_{2,t-1}$ from the second equation in (2.4') is very close to its counterpart in (1.15) (0.950556 compared with 0.811342), plus both coefficients having the same positive sign. Additionally, the coefficient value of $\varepsilon_{1,t-1}$ in the first equation of (2.2') is close to its counterpart in (1.10) (-0.460145 compared with -0.479305), plus both coefficients having the same negative sign. In contrast, the coefficient value of $\varepsilon_{2,t-1}$ in the second equation in (2.4') is very close to its counterpart in (1.15) (-1.00433 compared with -0.943283), plus both coefficients having the same negative sign. We find the results are very interesting, because the first equation of multivariate VARIMA (1, 1, 1) is much better than the second equation in the system as to the overall parameter's accuracy and sign from the first equation close to that of univariate ARIMA (1, 1, 1), in comparison the second equation of multivariate VARIMA (1, 1, 2) is much better than the first equation in the system as to the overall parameter's accuracy and sign from the second equation close to that of univariate ARIMA (1, 1, 2). We also see much difference in both value and sign of the parameters between first and second equation in the system (2.2') and (2.4').

(ii) ARMAX vs. VAR

ARMAX analysis shows us that there are positive relationships between GDP and Government Spending and between Government Spending and GDP, both concurrently in the short-run. In comparison, VAR results indicate that current GDP responses negatively to a shock in past Government Spending permanently in the long-run, while current Government Spending not necessarily yet temporarily reacts negatively to a shock in past GDP. The combined findings suggest that policy makers should not adjust current Government Expenditure based merely on the condition and performance of past economy (GDP), because the negative relationship between current Government Spending and past GDP is statistically insignificant, thus such relationship is only temporary. Instead, policy makers should focus on both the short-term benefits of spending to the current economy and the long-term effects of expenditure on the future economy, thereby finding a balanced yet economically viable solution to both short-term and long-term propensity. Specifically, decisions on government expenditure should not be limited by the current economic condition, and abrupt increase in government spending alone (without monetary policy—interest rate—adjustment) in the short-term might not be good to the long-term health of the economy. Fundamentally, for a more balanced approach, we theorize and predict that the short-term benefits to the current economy from increasing government expenditure often largely secured by the long-term loan should outweigh the negative effects to the future economy from the long-term debt incurred by the loan, or at least equal to.

(iii) VARMA vs. VAR

We are unable to obtain meaningful information from VARMA analysis on GDP and Government Expenditure, because the associated parameters are statistically insignificant due to the technical difficulties mentioned on the bottom of page 34. In comparison, VAR does give us a lot insights as elaborated in the above title (ii).

(iv) ARCH, GARCH, ARCH-M, GARCH-M, TS-GARCH, GJR, TARCH, NARCH, APARCH, EGARCH; Normal vs. Non-normal distribution

We find that models with non-normal distribution have overall smaller AIC, BIC, HQC, compared with models with normal distribution, excluding ARCH-M and GARCH-M, suggesting that non-normally distributed volatility models generally perform better than those normally distributed. Looking into each group of normally and non-normally distributed, we find that TARCH (1, 1) GED performs the best in the group of non-normally distributed, while GARCH-M does the best in the group of normally distributed.

3) Contributions to Literature

The contribution of this research to the current economic literature is profound in the following ways: 1) the combined analysis of ARMAX and VAR offers better insights on the cause and

effect between GDP and Government Expenditure not only in the short-run but in the long-run as well. VAR also builds a bridge enabling us to study how short-term effects long-term, interchangeably between GDP and Government Spending and between Government Spending and GDP. Thereby, the findings are practically significant to policy makers as shown in section 9; 2) theories and predictions derived from the ARMAX and VAR analysis and results are not only theoretically novel but also empirically significant, as shown in section 10, regarding the current economic debates on the topic between short-term stimulus spending and long-term debt; 3) we have empirically demonstrated that volatility models with non-normal distribution overall perform better than models distributed normally; 4) Kalman filter is both theoretically and empirically demonstrated to generate better estimations; 5) we are able to show similarity and difference between univariate ARMA and multivariate VARMA both theoretically and empirically.

4) Limitations and Future Research

Model limitations

(i) Unit root test: arbitrary lag order.

(ii) VAR: point estimate.

(iii) Multivariate VARMA/VAR: not including heteroskedasticity component like GARCH.

(iv) GARCH: univariate. (Multivariate GARCH has been discussed and used in the literature.)

Data limitations

Economic analysis is only limited to the U.S. data. Extension to include the foreign country data along with their comparison to the U.S. might provide further insights to the effectiveness and influence of fiscal policy on GDP.

Future research

Theoretical extension1: ARMAX extends to Multivariate VARMAX.

Theoretical extension2: VAR extends to VARCH or VGARCH.

Theoretical extension3: Consider combining GARCH with Multivariate ARMA. For example,

GARCH + Multivariate VARMA = Multivariate VGAR(MA)CH. The idea is to incorporate the component of moving average to the Multivariate GARCH model.

Empirical extension: Use VAR and ARMAX to first study the individual effect of each monetary policy (interest rate) and fiscal policy (government spending) on GDP; then study the simultaneous effects of monetary and fiscal policy on GDP; and finally compare the results of

step one and two. The goal is to not only examine application of the classic Keynesian theory in the short-term, but more importantly explore the long-term effects of both monetary policy and fiscal policy on GDP as well.

```
5) Gretl Scripts for System Equation (VAR, VARMA, etc.)
```

VAR (1)

```
series x = diff(GDPC1Q)
series y = diff(GCEC1Q)
system
    equation x const x(-1) y(-1)
    equation y const x(-1) y(-1)
end system
estimate $system method=ols
```

Notes: The developed VAR gretl system procedure can also be used similarly in other

multivariate AR analysis (such as VARX), with slight modification.

VARMA (1, 1)

```
series x = diff(GDPC1Q)
series y = diff(GCEC1Q)
series a = uhat1
series b = uhat2
system
    equation x const x(-1) y(-1) a(-1) b(-1)
    equation y const x(-1) y(-1) a(-1) b(-1)
end system
estimate $system method=ols
```

Notes for the VARMA gretl procedures:

Step 1: Run ARIMA (1, 1, 1) for GDPC1Q once and run ARIMA (1, 1, 1) for GCEC1Q once

Step 2: Save to the dataset each of the residuals (uhat1 and uhat2) from each univariate ARIMA

(1, 1, 1) of GDPC1Q and GCEC1Q respectively.

Step 3: Input the above VARMA (1, 1) gretl scripts and get the results.

Notes: The developed VARMA procedures can be used similarly in other multivariate ARMA

analysis (such as VARMAX) as well, with slight modification.

VARMA (1, 2)

series x = diff(GDPC1Q) series y = diff(GCEC1Q) series a = uhat1 series b = uhat2 system equation x const x(-1) y(-1) a(-1) b(-1) a(-2) b(-2) equation y const x(-1) y(-1) a(-1) b(-1) a(-2) b(-2) end system estimate \$system method=ols

Disclaimer: This document and its associated work, including but not limited to methodology, script, coding, ideas, insights, discoveries, implications, theoretical and empirical extensions are all copyright protected. Copy and distribution of this paper without original author's permission is strictly prohibited. Please cite the work appropriately. Since this is not a peer reviewed article, author will not be responsible for any errors.