

Hypergeometric Hodge modules

Thomas Reichelt and Christian Sevenheck

April 16, 2020

Abstract

We consider mixed Hodge module structures on GKZ-hypergeometric differential systems. We show that the Hodge filtration on these \mathcal{D} -modules is given by the order filtration, up to a suitable shift. As an application, we prove a conjecture on the existence of non-commutative Hodge structures on the reduced quantum \mathcal{D} -module of a nef complete intersection inside a toric variety.

Contents

1	Introduction	2
2	A guiding example	5
3	GKZ-systems and the Fourier-Laplace transformation	15
3.1	GKZ-systems and strict resolutions	15
3.2	Fourier-Laplace transformed GKZ-systems	17
4	Hodge filtration on torus embeddings	19
4.1	V-filtration	20
4.2	Compatibility of filtrations	24
4.3	Calculation of the Hodge filtration	30
5	Integral transforms of torus embeddings	36
5.1	Hypergeometric modules, Gauß-Manin systems and the Radon transformation	36
5.2	Integral transforms of twisted structure sheaves	39
5.3	Calculation in charts	42
5.4	A Koszul complex	48
5.5	\mathcal{R} -modules	51
5.6	A quasi-isomorphism	54
5.7	Computation of the direct image	56
5.8	Local cohomology of semi-group rings	58
5.9	Statement and proof of the main theorem	63
5.10	Duality	65
5.11	Hodge structures on affine hypersurfaces of tori	67
6	Landau-Ginzburg models and non-commutative Hodge structures	71

2010 *Mathematics Subject Classification*. 14F10 (primary), 32C38 (secondary)

Keywords: GKZ systems, Hodge modules, Hodge filtration, Radon transformation, non-commutative Hodge structures
The authors are supported by the DFG grants RE 3567/1-1 and SE 1114/5-1.

1 Introduction

In a series of papers Gel'fand, Graev, Kapranov and Zelevinskiĭ [GfGZ87], [GfZK89] introduced a system of differential equations which generalize the classical differential systems satisfied by the hypergeometric functions of Gauß, Appell, Bessel and others. These generalized systems are nowadays called GKZ-systems. The initial data of a GKZ system consists of a $d \times n$ integer matrix and a parameter vector β . Although the definition of a GKZ system has a combinatorial flavor it was early realized that at least for non-resonant parameter vectors β GKZ-systems come from geometry [GKZ90], i.e. they are isomorphic to a direct image of some twisted structure sheaf on an algebraic variety. In [Rei14], the first-named author has shown that certain GKZ-systems actually carry a much richer structure, namely, they underlie *mixed Hodge modules* in the sense of M. Saito (see [Sai90]). One of the main goals of this paper is the explicit calculation of the corresponding Hodge filtration on these modules.

An important application of GKZ systems is mirror symmetry for weak Fano complete intersections in toric varieties. We have shown in our previous papers [RS15, RS17] how to express variants of the mirror correspondence as an equivalence of differential systems of “GKZ-type”. However, an important point was left open in these articles: The mirror statements given there actually involve differential systems (i.e., holonomic \mathcal{D} -modules) with some additional data, sometimes called *lattices*. These are constructed by a variant of the Fourier-Laplace transformation from regular holonomic *filtered* \mathcal{D} -modules. The filtration in question is the Hodge filtration on these modules, but a concrete description of it is missing in [RS15, RS17]. As a consequence, the most important Hodge theoretic property of the differential system entering in the mirror correspondence was formulated only as a conjecture in [RS17] (conjecture 6.15): the so-called *reduced quantum \mathcal{D} -module*, which governs certain Gromov-Witten invariants of nef complete intersections in toric varieties conjecturally underlies a *variation of non-commutative Hodge structures*. We prove this conjecture here (see Theorem 6.6), it appears as a consequence of the main result of the present paper, which determines the Hodge filtration on the GKZ-systems. More precisely, as GKZ-systems are defined as cyclic quotients of the Weyl algebra, we obtain (Theorem 5.35) that this Hodge filtration is given by the filtration induced from the order of differential operators up to a suitable shift. In some sense, this finishes the Hodge theoretic study of mirror symmetry for this class of varieties since we can now express the mirror correspondence as an isomorphism of non-commutative Hodge structures, which are the correct generalization of ordinary Hodge structures in the case where the underlying differential equations acquire irregular singularities, as it is the case for the quantum \mathcal{D} -module of weak Fano varieties (in contrast to the Calabi-Yau case).

Another application of our main result, which can be found in the two papers [CnDS19] and [CnDRS19], is the calculation of the so-called irregular Hodge filtration on certain 1-dimensional classical hypergeometric modules. The irregular Hodge filtration has been introduced by C. Sabbah (see [Sab18]) in order to attach Hodge-type numerical invariants (namely dimensions of graded parts of a filtration) to differential systems acquiring irregular singularities. In geometric situations, like those where regular functions on quasi-projective manifolds are studied as Landau-Ginzburg models of certain quantum cohomology theories, the irregular Hodge filtration has a concrete description using certain logarithmic de Rham complexes, as has been shown by Esnault, Sabbah and Yu ([ESY17]), see also the discussion in [KKP17]). Classical hypergeometric systems are also the most prominent example of *rigid* \mathcal{D} -modules (see [Kat90]), so the computation of these invariants for them is of particular interest. It turns out that confluent classical hypergeometric modules (these are precisely those with irregular singularities) are obtained from GKZ-systems by a dimensional reduction and a Fourier-Laplace transformation. Using our result (i.e., Theorem 5.35), one can explicitly describe the irregular Hodge filtration (and give closed formulas for irregular Hodge numbers) of certain such systems (see [CnDS19, Theorem 4.7] and [CnDRS19, Theorem 5.9] for more details).

Let us give a short overview on the content of this article and the precise statements of the main results. Notice that section 2 below provides a detailed description of these results and parts of their proofs for rather simple example, which is related to the quantum \mathcal{D} -module of \mathbb{P}^1 . We advise the reader to go through this example in order to understand the strategy of the proof in the general case in the main body of this article.

The main result of this paper is obtained in two major steps, which occupy section 4 resp. sec-

tion 5. First we study embeddings of tori into affine spaces given by a monomial map $h_A : T = (\mathbb{C}^*)^d \hookrightarrow \mathbb{C}^n; (t_1, \dots, t_d) \mapsto (\underline{t}^{a_1}, \dots, \underline{t}^{a_n})$, where $\underline{t}^{a_i} = \prod_{k=1}^d t_k^{a_{ki}}$ and where the matrix of columns $A = (a_i)_{i=1, \dots, n} \in M(d \times n, \mathbb{Z})$ satisfies certain combinatorial properties related to the geometry of the semi-group ring $\mathbb{C}[\text{NA}]$. Consider the twisted structure sheaf $\mathcal{O}_T^\beta := \mathcal{D}_T / \mathcal{D}_T(\partial_{t_1} t_1 + \beta_1, \dots, \partial_{t_d} t_d + \beta_d)$. It was shown in [SW09] that the direct image $h_{A+} \mathcal{O}_T^\beta$ has an explicit description as a Fourier-Laplace transformed GKZ-system $\check{\mathcal{M}}_A^\beta$ (cf. Definition 3.5) in case the parameter vector β is not strongly resonant (cf. Definition 3.8). We consider the corresponding direct image $\mathcal{H}^0(h_{A*} {}^p \mathbb{C}_T^{H, \beta})$ in the category of complex mixed Hodge modules, and calculate its Hodge filtration (cf. Theorem 4.17) in case that the parameter vector β lies in the set of admissible parameters \mathfrak{A}_A (cf. Formula (15)). More precisely, this first result can be stated as follows.

Theorem (Theorem 4.17 below). *For $\beta \in \mathfrak{A}_A$ the Hodge filtration on $\check{\mathcal{M}}_A^\beta$ is equal to the order filtration shifted by $n - d$, i.e.*

$$F_{p+(n-d)}^H \check{\mathcal{M}}_A^\beta = F_p^{\text{ord}} \check{\mathcal{M}}_A^\beta.$$

If the matrix A we started with satisfies a homogeneity property, then the underlying \mathcal{D} -module of this mixed Hodge module is a (monodromic) Fourier-Laplace transformation of the GKZ-system we are interested in. It should be noticed that Theorem 4.17 is of independent interest, its statement is related to the description of the Hodge filtration on various cohomology groups associated with singular toric varieties. We plan to discuss this question in a subsequent work. The main point in Theorem 4.17 is to determine the canonical V -filtration on the direct image module along the boundary divisor $\overline{\text{im}}(h_A) \setminus \text{im}(h_A)$, i.e., the calculation of some Bernstein polynomials.

The second step, carried out in section 5 consists in studying the behavior of a twisted structure sheaf on a torus under a certain integral transformation which generalizes the Radon transformation in [Rei14]. It is well-known (see [Bry86] and [DE03]) that there is a close relation between the Fourier-Laplace transformation and the Radon transformation for holonomic \mathcal{D} -modules, however, the former does not a priori preserve the category of mixed Hodge modules whereas the latter does. This fact is one of the main points in the proof of the existence of a mixed Hodge module structure on GKZ-systems in [Rei14]. We calculate the behaviour of the Hodge filtration under the various functors entering into the integral transformation functor, an essential tool for these calculations is the so called *Euler-Koszul-complex* (or some variants of it) as introduced in [MMW05]. We finally get the following statement for the Hodge filtration on the GKZ system \mathcal{M}_A^β (cf. Theorem 5.35). We call a matrix homogenous if all of its columns lie in an affine hyperplane. Moreover, an integer matrix is called normal, if the semi-group generated by its columns is the intersection of the cone generated by these columns with the lattice generated by them (see Formula (11) below).

Theorem (Theorem 5.35 below). *Let \tilde{A} be a homogeneous, normal $(d+1) \times (n+1)$ integer matrix, $\tilde{\beta} \in \mathfrak{A}_{\tilde{A}}$ and $\beta_0 \in (-1, 0]$. Then the GKZ-system $\mathcal{M}_{\tilde{A}}^{\tilde{\beta}}$ carries the structure of a mixed Hodge module whose Hodge filtration is given by the shifted order filtration, i.e.*

$$(\mathcal{M}_{\tilde{A}}^{\tilde{\beta}}, F_{\bullet}^H) \simeq (\mathcal{M}_{\tilde{A}}^{\tilde{\beta}}, F_{\bullet+d}^{\text{ord}}).$$

The second last part of section 5 deals with the Hodge module structure on the holonomic dual GKZ-system (which, under the assumptions on the initial data, is also a GKZ-system). The last subsection of section 5 explains how one can deduce from our main result the computation of Batyrev (see [Bat93]) of the Hodge filtration on the relative cohomology of smooth affine hypersurfaces in algebraic tori.

Finally, in section 6 we explain the above mentioned conjecture from [RS17] and show how its proof can be deduced from our main result (cf. Theorem 6.6). To formulate this result, consider a smooth projective toric variety X_Σ and a split globally generated vector bundle \mathcal{E} satisfying some positivity assumptions (ensuring, in particular, that the subvariety $Y := s^{-1}(0)$ for a generic section $s \in \Gamma(X_\Sigma, \mathcal{E})$ is smooth weak Fano). Then we have the reduced quantum \mathcal{D} -module $\overline{\text{QDM}}(X_\Sigma, \mathcal{E})$ (see [RS17, Section 4.1]) and the mirror map Mir (see Theorem 6.5 below as well as [RS17, Theorem 6.9] and the references therein). We get the following result.

Theorem (Theorem 6.6 below). *Let X_Σ be a k -dimensional smooth, toric variety, let $\mathcal{L}_1, \dots, \mathcal{L}_l$ be globally generated line bundles such that $-K_{X_\Sigma} - \mathcal{E}$ is nef, where $\mathcal{E} = \bigoplus_{j=1}^l \mathcal{L}_j$, \mathcal{L}_j being ample for*

$j = 1, \dots, l$. Then the smooth $\mathcal{R}_{\mathbb{C}_z \times \mathcal{KM}^\circ}$ -module $(\text{id}_{\mathbb{C}_z} \times \text{Mir})^* \overline{\text{QDM}}(X_\Sigma, \mathcal{E})$ underlies a variation of pure polarized non-commutative Hodge structures.

While we were working on this paper, a preprint of T. Mochizuki ([Moc15b]) appeared where [RS17, Conjecture 6.15] is shown with rather different methods. The arguments in his paper work entirely in the category of mixed twistor modules, using the full strength of [Moc15a]. It seems possible from his approach to obtain our main result by considering equivariant twistor modules (see [Moc15b, Section 6.4]). However, this treatment of the regular case in loc.cit. (i.e., the case of mixed Hodge modules) seems to be done in a more restrictive setup, in two ways: First of all, there is an extra assumption ([Moc15b, Assumption 6.51]) excluding the calculation of the Hodge filtration on GKZ-systems for examples coming from local mirror symmetry (however, the mirror symmetry consequence, i.e., the proof of [RS17, Conjecture 6.15] is covered by [Moc15b]). On the other hand, there is no discussion on GKZ-systems (or its twistor versions) for non-zero parameter vectors β in loc.cit. in contrast to our main result (i.e., in contrast to Theorem 5.35, which holds for non-zero parameters satisfying a natural combinatorial condition). Besides, we feel that the calculation of the Hodge filtration on GKZ-systems which only uses properties of the category of mixed Hodge modules (not passing via twistor modules) may be of independent interest.

To finish this introduction, we introduce some notation and conventions used throughout the paper. Let X be a smooth algebraic variety over \mathbb{C} of dimension d_X . We denote by $M(\mathcal{D}_X)$ the abelian category of algebraic left \mathcal{D}_X -modules on X and denote the abelian subcategory of (regular) holonomic \mathcal{D}_X -modules by $M_h(\mathcal{D}_X)$ (resp. $(M_{rh}(\mathcal{D}_X))$). The full triangulated subcategory in $D^b(\mathcal{D}_X)$, consisting of objects with (regular) holonomic cohomology, is denoted by $D_h^b(\mathcal{D}_X)$ (resp. $D_{rh}^b(\mathcal{D}_X)$).

Let $f : X \rightarrow Y$ be a map between smooth algebraic varieties. Let $M \in D^b(\mathcal{D}_X)$ and $N \in D^b(\mathcal{D}_Y)$, then we denote by

$$f_+ M := Rf_*(\mathcal{D}_{Y \leftarrow X} \overset{L}{\otimes} M) \quad \text{resp.} \quad f^+ M := \mathcal{D}_{X \rightarrow Y} \overset{L}{\otimes} f^{-1} M[d_X - d_Y]$$

the direct resp. inverse image for \mathcal{D} -modules. Recall that the functors f_+, f^+ preserve (regular) holonomicity (see e.g., [HTT08, Theorem 3.2.3]). We denote by $\mathbb{D} : D_h^b(\mathcal{D}_X) \rightarrow (D_h^b(\mathcal{D}_X))^{opp}$ the holonomic duality functor. Recall that for a single holonomic \mathcal{D}_X -module M , the holonomic dual is also a single holonomic \mathcal{D}_X -module ([HTT08, Proposition 3.2.1]) and that holonomic duality preserves regular holonomicity ([HTT08, Theorem 6.1.10]).

For a morphism $f : X \rightarrow Y$ between smooth algebraic varieties we additionally define the functors $f_\dagger := \mathbb{D} \circ f_+ \circ \mathbb{D}$ and $f^\dagger := \mathbb{D} \circ f^+ \circ \mathbb{D}$.

Let $MF(\mathcal{D}_X)$ be the category of filtered \mathcal{D}_X -modules (M, F) where the ascending filtration F_\bullet satisfies

1. $F_p M = 0$ for $p \ll 0$
2. $\bigcup_p F_p M = M$
3. $(F_p \mathcal{D}_X) F_q M \subset F_{p+q} M$ for $p \in \mathbb{Z}_{\geq 0}, q \in \mathbb{Z}$

where $F_\bullet \mathcal{D}_X$ is the filtration by the order of the differential operator.

We denote by $\text{MHM}(X)$ the abelian category of algebraic mixed Hodge modules and by $D^b\text{MHM}(X)$ the corresponding bounded derived category. The forgetful functor to the bounded derived category of regular holonomic \mathcal{D} -modules is denoted by

$$D_{\text{mod}} : D^b\text{MHM}(X) \longrightarrow D_{rh}^b(\mathcal{D}_X).$$

For each morphism $f : X \rightarrow Y$ between complex algebraic varieties, there are induced functors

$$f_*, f! : D^b\text{MHM}(X) \longrightarrow D^b\text{MHM}(Y)$$

and

$$f^*, f^\dagger : D^b\text{MHM}(Y) \rightarrow D^b\text{MHM}(X),$$

which are interchanged by \mathbb{D} . The functors $f_*, f_!, f^*, f^!$ lift the analogous functors $f_+, f_\dagger, f^\dagger, f^+$ on $D_{rh}^b(\mathcal{D}_X)$. Let \mathbb{Q}_{pt}^H be the unique mixed Hodge structure with $\mathrm{Gr}_i^W = \mathrm{Gr}_i^F = 0$ for $i \neq 0$ and underlying vector space \mathbb{Q} . Denote by $a_X : X \rightarrow \{pt\}$ the map to the point and set

$$\mathbb{Q}_X^H := a_X^* \mathbb{Q}_{pt}^H.$$

The shifted object ${}^p\mathbb{Q}_X^H := \mathbb{Q}_X^H[d_X]$ lies in $\mathrm{MHM}(X)$ and is equal to $(\mathcal{O}_X, F, \mathbb{Q}_X[d_X], W)$ with $\mathrm{Gr}_p^F = 0$ for $p \neq 0$ and $\mathrm{Gr}_i^W = 0$ for $i \neq d_X$. We have $\mathbb{D}\mathbb{Q}_X^H \simeq a_X^! \mathbb{Q}_{pt}^H$ and, since X is smooth, the isomorphism

$$\mathbb{D}\mathbb{Q}_X^H \simeq \mathbb{Q}_X^H(d_X)[2d_X]. \quad (1)$$

Here (d_X) denotes the Tate twist (see e.g., [Sai90, page 257]).

We also have to consider the category $\mathrm{MHM}(X, \mathbb{C})$ of complex mixed Hodge modules which can be defined as follows (see [DS13, Definition 3.2.1]): First note that one can naturally extend the notion of a \mathbb{Q} -mixed Hodge module (i.e., an object of $\mathrm{MHM}(X)$) to \mathbb{R} -mixed Hodge module, due to the work of Mochizuki on mixed twistor modules (see in particular [Moc15a, Section 13.5], where the notion of a K -mixed Hodge module is considered, K being any subfield of \mathbb{R}). Then we say that a filtered \mathcal{D}_X -module (M, F_\bullet) underlies a complex mixed Hodge module if it is a direct summand of a filtered \mathcal{D}_X -module that underlies an \mathbb{R} -mixed Hodge module. Many properties of the category of \mathbb{R} -mixed Hodge modules carry over to $\mathrm{MHM}(X, \mathbb{C})$ since they are stable by direct summands.

Let $T = (\mathbb{C}^*)^d$ be a torus with coordinates t_1, \dots, t_d and $\beta \in \mathbb{R}^d$. We denote by \mathcal{O}_T^β the \mathcal{D}_T -module

$$\mathcal{O}_T^\beta = \mathcal{D}_T / ((\partial_{t_i} t_i + \beta_i)_{i=1, \dots, d})$$

and by $\mathbb{C}_T^{H, \beta}$ the complex Hodge module $(\mathcal{O}_T^\beta, F, W)$ with $\mathrm{Gr}_p^F \mathbb{C}_T^{H, \beta} = 0$ for $p \neq 0$ and $\mathrm{Gr}_i^W \mathbb{C}_T^{H, \beta} = 0$ for $i \neq d$. Finally we set ${}^p\mathbb{C}_T^{H, \beta} := \mathbb{C}_T^{H, \beta}[d_T]$.

Acknowledgments: We thank Takuro Mochizuki, Claude Sabbah and Uli Walther for many stimulating discussions on the content of this paper.

2 A guiding example

In this section, we intend to discuss a particular example, related to the quantum differential equation of \mathbb{P}^1 , where most of the techniques used in the main body of the paper can be written down quite explicitly. We hope that this section will help the reader to find his way through the technical difficulties of the paper.

Let \tilde{A} be the following 2×3 -matrix with integer entries:

$$\begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & -1 \end{pmatrix}.$$

As explained in Definition 3.1 below, any $d \times n$ -integer matrix A together with a vector $\beta \in \mathbb{C}^d$ defines a cyclic $\mathcal{D}_{\mathbb{C}^n}$ -module called a GKZ-system and denoted by \mathcal{M}_A^β . For the above matrix, this system for the vector $\beta = 0$ is given by

$$\mathcal{M}_{\tilde{A}}^0 = \mathcal{D}_{\mathbb{C}^3} / (\partial_{\lambda_0}^2 - \partial_{\lambda_1} \partial_{\lambda_2}, \lambda_0 \partial_{\lambda_0} + \lambda_1 \partial_{\lambda_1} + \lambda_2 \partial_{\lambda_2}, \lambda_1 \partial_{\lambda_1} - \lambda_2 \partial_{\lambda_2})$$

It is well known that $\mathcal{M}_{\tilde{A}}^0$ is holonomic (see [Ado94]) and regular ([SW08]). Moreover, it follows from [Rei14, Theorem 3.5] that $\mathcal{M}_{\tilde{A}}^0$ underlies a mixed Hodge module ${}^H\mathcal{M}_{\tilde{A}}^0 \in \mathrm{MHM}(\mathbb{C}^3)$. The purpose of this introductory section is to explain and partly prove the following statement, which is a very special case of the main theorem of this paper (Theorem 5.35).

Theorem 2.1. *We have an isomorphism of filtered $\mathcal{D}_{\mathbb{C}^3}$ -modules*

$$(\mathcal{M}_{\tilde{A}}^0, F_\bullet^H) \simeq (\mathcal{M}_{\tilde{A}}^0, F_{\bullet+1}^{ord}),$$

where F_\bullet^H denotes the filtration such that the filtered module $(\mathcal{M}_{\tilde{A}}^0, F_\bullet^H)$ underlies the mixed Hodge module ${}^H\mathcal{M}_{\tilde{A}}^0$ and where F_\bullet^{ord} is the filtration induced on $\mathcal{M}_{\tilde{A}}^0$ by the filtration on $\mathcal{D}_{\mathbb{C}^3}$ by orders of differential operators.

The proof of this theorem will be done in several steps that parallel the main steps of the proof of Theorem 5.35. The two major simplifications are that we are dealing with the very special matrix \tilde{A} , whereas in Theorem 5.35 any $(d+1) \times (n+1)$ -matrix \tilde{A} satisfying some combinatorial conditions is considered, and moreover we restrict to the parameter value $\beta = 0$ (compare with the general definition of GKZ-systems $\mathcal{M}_{\tilde{A}}^{\beta}$ in Definition 3.1 resp. with the Definition 3.5 for the sheaf $\tilde{\mathcal{M}}_{\tilde{A}}^{\beta}$). This avoids considering some rather involved combinatorial condition relating \tilde{A} and β (see the definition of the set $\mathfrak{A}_{\tilde{A}}$ in Equation (15) below).

We first consider the morphism

$$\begin{aligned} h_{\tilde{A}} : (\mathbb{C}^*)^2 &\longrightarrow \mathbb{C}^3 =: W \\ (t_0, t_1) &\longmapsto (t_0, t_0 \cdot t_1, t_0 \cdot t_1^{-1}) =: (w_0, w_1, w_2) \end{aligned}$$

where the exponents of the monomials in the components of the map are exactly the columns of \tilde{A} . As explained in more detail in Theorem 3.9 and Proposition 3.11 below, it follows from a result of Schulze and Walther ([SW09]) that we have the following isomorphism of \mathcal{D}_W -modules

$$\begin{aligned} h_{\tilde{A},+} \mathcal{O}_{(\mathbb{C}^*)^2} &\cong h_{\tilde{A},+} \mathcal{D}_{(\mathbb{C}^*)^2} / (\partial_{t_0} t_0 + 2, \partial_{t_1} t_1 - 1) \\ &\stackrel{!}{\cong} \mathcal{D}_W / (w_0^2 - w_1 w_2, \partial_{w_0} w_0 + \partial_{w_1} w_1 + \partial_{w_2} w_2, \partial_{w_1} w_1 - \partial_{w_2} w_2) =: \tilde{\mathcal{M}}_{\tilde{A}}^0. \end{aligned}$$

The main point in this result is to show that (left) multiplication by w_0 is invertible on $\tilde{\mathcal{M}}_{\tilde{A}}^0$, then the isomorphism follows since the map $h_{\tilde{A}}$ can be decomposed as

$$h_{\tilde{A}} : (\mathbb{C}^*)^2 \xrightarrow{h_{\tilde{A},1}} \mathbb{C}^* \times \mathbb{C}^2 \xrightarrow{h_{\tilde{A},2}} W$$

where both $h_{\tilde{A},1}$ and $h_{\tilde{A},2}$ are embeddings and where $h_{\tilde{A},1}$ sends (t_0, t_1) to $(t_0, t_0 \cdot t_1, t_0 \cdot t_1^{-1})$ and is closed (this follows since the map $\mathbb{C}^* \rightarrow \mathbb{C}^2, t \mapsto (t, t^{-1})$ is a closed embedding), whereas $h_{\tilde{A},2}$ is the canonical open embedding of $\mathbb{C}^* \times \mathbb{C}^2$ into W .

A second consequence of Proposition 3.11 below is that $\tilde{\mathcal{M}}_{\tilde{A}}^0$ underlies a mixed Hodge module on W , namely the object $h_{\tilde{A},*} {}^p\mathcal{Q}_T^H \in \text{MHM}(W)$. A first step to prove Theorem 2.1 above is to compute the Hodge filtration on $\tilde{\mathcal{M}}_{\tilde{A}}^0$. This task can be divided into two steps: First we have to compute the Hodge filtration on the module ${}^*\tilde{\mathcal{M}}_{\tilde{A}}^0 := h_{\tilde{A},1,+} \mathcal{O}_{(\mathbb{C}^*)^2}$ which underlies the mixed Hodge module $h_{\tilde{A},1,*} {}^p\mathcal{Q}_T^H \in \text{MHM}(\mathbb{C}^* \times \mathbb{C}^2)$. This is done via a rather direct argument since $h_{\tilde{A},1}$ is closed. The second step, which is more delicate, is to obtain from this the Hodge filtration on $h_{\tilde{A},2,+} {}^*\tilde{\mathcal{M}}_{\tilde{A}}^0 = \tilde{\mathcal{M}}_{\tilde{A}}^0$.

Here we are faced with the fundamental problem of extending a mixed Hodge module from the complement of a (smooth) divisor to the total space. While this operation is easily understood at the level of \mathcal{D} -modules, one cannot simply use the direct image functors for \mathcal{O} -modules to calculate the extension of the filtration steps of the Hodge filtration since those are by definition \mathcal{O} -coherent, a property that is lost under direct images of open embeddings. As is explained in more detail at the beginning of section 4.3, this problem is solved by intersecting the direct image with the canonical V -filtration along the divisor in question. In order to compute the Hodge filtration on $h_{\tilde{A},2,+} {}^*\tilde{\mathcal{M}}_{\tilde{A}}^0$, we thus have to calculate this V -filtration along $w_1 = 0$.

Notice that in the main body of the text, we follow a slightly different strategy, due to the fact that the factorization of $h_{\tilde{A}}$ used above into a closed and an open embedding may look different depending on the shape of the matrix \tilde{A} . In general, one can always consider the factorization into a map between tori, which is a closed embedding (which would be the map from $(\mathbb{C}^*)^2$ to $(\mathbb{C}^*)^3$ in the above example) followed by the canonical open embedding from the torus into affine space. The latter, however, is the extension over a normal crossing divisor, which is not smooth. In order to apply the techniques sketched above, one has to compose further with a graph embedding with respect to the equation of the normal crossing divisor (see diagram (23) and the arguments following it).

The first (easy) step of the calculation of the Hodge filtration on $\tilde{\mathcal{M}}_{\tilde{A}}^0$ can be formulated as follows.

Lemma 2.2 (Compare Lemma 4.14 below for the general case). *The direct image $\mathcal{H}^0 h_{\tilde{A},1,+} \mathcal{O}_{(\mathbb{C}^*)^2}$ is isomorphic to the cyclic $\mathcal{D}_{\mathbb{C}^* \times \mathbb{C}^2}$ -module*

$$\mathcal{D}_{\mathbb{C}^* \times \mathbb{C}^2} / (w_0^2 - w_1 w_2, \partial_{w_0} w_0 + \partial_{w_1} w_1 + \partial_{w_2} w_2, \partial_{w_1} w_1 - \partial_{w_2} w_2)$$

and the Hodge filtration on this module is given, under this isomorphism, by the induced order filtration, shifted by one, i.e., we have

$$F_p^H \mathcal{H}^0 h_{\tilde{A},1,+} \mathcal{O}_{(\mathbb{C}^*)^2} = F_{p-1}^{ord} [\mathcal{D}_{\mathbb{C}^* \times \mathbb{C}^2} / (w_0^2 - w_1 w_2, \partial_{w_0} w_0 + \partial_{w_1} w_1 + \partial_{w_2} w_2, \partial_{w_1} w_1 - \partial_{w_2} w_2)]$$

Proof. We can factor $h_{\tilde{A},1}$ further as $h_{\tilde{A},1} = \tilde{h}_{\tilde{A},1} \circ \tilde{h}_{\tilde{A},1}$, where

$$\begin{aligned} \tilde{h}_{\tilde{A},1} : (\mathbb{C}^*)^2 &\longmapsto (\mathbb{C}^*)^3 \\ (t_0, t_1) &\longmapsto (t_0, t_0 \cdot t_1, t_0 \cdot t_1^{-1}) \end{aligned}$$

and where $\tilde{h}_{\tilde{A},1} : (\mathbb{C}^*)^3 \hookrightarrow \mathbb{C}^* \times \mathbb{C}^2$ is the canonical open embedding. Then since $h_{\tilde{A},1}$ is a closed embedding, we know that the support of ${}^* \mathcal{M}_{\tilde{A}}^0$ is disjoint from the divisor $(\mathbb{C}^* \times \mathbb{C}^2) \setminus (\mathbb{C}^*)^3$, which implies that $F_p^H {}^* \mathcal{M}_{\tilde{A}}^0 = \tilde{h}_{\tilde{A},1,*} F_p^H \tilde{h}_{\tilde{A},1,+} \mathcal{O}_{(\mathbb{C}^*)^2}$. It therefore suffices to determine the filtration steps $F_p^H \tilde{h}_{\tilde{A},1,+} \mathcal{O}_{(\mathbb{C}^*)^2}$, or, more precisely, to show that

$$F_p^H \tilde{h}_{\tilde{A},1,+} \mathcal{O}_{(\mathbb{C}^*)^2} = F_{p-1}^{ord} [\mathcal{D}_{(\mathbb{C}^*)^3} / (w_0^2 - w_1 w_2, \partial_{w_0} w_0 + \partial_{w_1} w_1 + \partial_{w_2} w_2, \partial_{w_1} w_1 - \partial_{w_2} w_2)]$$

Consider the coordinate change

$$\begin{aligned} \phi : (\mathbb{C}^*)^3 &\longrightarrow (\mathbb{C}^*)^3 \\ (w_0, w_1, w_2) &\longmapsto (w_0, w_1/w_0, w_1 w_2/w_0^2) =: (u_0, u_1, u_2) \end{aligned}$$

so that $(\phi \circ \tilde{h}_{\tilde{A},1})(t_0, t_1) = (t_0, t_1, 1)$ and

$$(\phi \circ \tilde{h}_{\tilde{A},1})_+ \mathcal{O}_{(\mathbb{C}^*)^2} = \mathcal{D}_{(\mathbb{C}^*)^3} / (u_2 - 1, \partial_{u_0} u_0 + 2, \partial_{u_1} u_1) = (\mathcal{D}_{(\mathbb{C}^*)^2} / (\partial_{u_0} u_0 + 2, \partial_{u_1} u_1)) [\partial_{u_2}].$$

According to [Sai93, Formula (1.8.6)], we have

$$F_{p+1}^H (\phi \circ \tilde{h}_{\tilde{A},1})_+ \mathcal{O}_{(\mathbb{C}^*)^2} = \sum_{p_1+p_2=p} F_{p_1}^H (\mathcal{D}_{(\mathbb{C}^*)^2} / (\partial_{u_0} u_0 + 2, \partial_{u_1} u_1)) \partial_{u_2}^{p_2}$$

Since $F_{p_1}^H (\mathcal{D}_{(\mathbb{C}^*)^3} / (\partial_{u_0} u_0, \partial_{u_1} u_1)) = F_{p_1}^{ord} (\mathcal{D}_{(\mathbb{C}^*)^3} / (\partial_{u_0} u_0, \partial_{u_1} u_1))$, we obtain from the above formula that

$$F_p^H (\phi \circ \tilde{h}_{\tilde{A},1})_+ \mathcal{O}_{(\mathbb{C}^*)^2} = F_{p-1}^{ord} (\mathcal{D}_{(\mathbb{C}^*)^3} / (u_2 - 1, \partial_{u_0} u_0, \partial_{u_1} u_1)).$$

Since ϕ is invertible, we obtain that the Hodge filtration on $\phi_+^{-1}((\phi \circ \tilde{h}_{\tilde{A},1})_+ \mathcal{O}_{(\mathbb{C}^*)^2}) = \tilde{h}_{\tilde{A},1,+} \mathcal{O}_{(\mathbb{C}^*)^2}$ is the order filtration on $\mathcal{D}_{(\mathbb{C}^*)^3} / (w_0^2 - w_1 w_2, \partial_{w_0} w_0 + \partial_{w_1} w_1 + \partial_{w_2} w_2, \partial_{w_1} w_1 - \partial_{w_2} w_2)$, shifted by one. As discussed above, the closure of the support of $\tilde{h}_{\tilde{A},1,+} \mathcal{O}_{(\mathbb{C}^*)^2}$ in $\mathbb{C}^* \times \mathbb{C}^2$ lies entirely in the torus $(\mathbb{C}^*)^3$, therefore, we obtain

$$F_p^H h_{\tilde{A},1,+} \mathcal{O}_{(\mathbb{C}^*)^2} = F_{p-1}^{ord} [\mathcal{D}_{\mathbb{C}^* \times \mathbb{C}^2} / (w_0^2 - w_1 w_2, \partial_{w_0} w_0 + \partial_{w_1} w_1 + \partial_{w_2} w_2, \partial_{w_1} w_1 - \partial_{w_2} w_2)],$$

as required. \square

The next step is to compute the Hodge filtration of the open direct image $h_{\tilde{A},2,+} {}^* \check{\mathcal{M}}_{\tilde{A}}^0 = \check{\mathcal{M}}_{\tilde{A}}^0$. As mentioned above, this needs information on the canonical V -filtration of the module $\check{\mathcal{M}}_{\tilde{A}}^0$ with respect to the smooth divisor $\{w_0 = 0\}$. More precisely, we have the following important formula (see Formula (22) below, as well as [Sai93, Proposition 4.2.]):

$$F_p^H \check{\mathcal{M}}_{\tilde{A}}^0 = \sum_{i \geq 0} \partial_{w_0}^i \left(V^0 \check{\mathcal{M}}_{\tilde{A}}^0 \cap h_{\tilde{A},2,*} F_{p-i}^H {}^* \check{\mathcal{M}}_{\tilde{A}}^0 \right) \quad (2)$$

Hence we need to determine the object $V^0 \check{\mathcal{M}}_A^0$. For what follows, it is more convenient to work out everything at the level of global sections. Since all modules we are considering here are defined on affine spaces, this is obviously sufficient.

We refer to [MM04] for details on the V -filtration. For what follows in this introduction, we only need that $V^0 D_{\mathbb{C}^3} = \mathbb{C}[w_0, w_1, w_2] \langle w_0 \partial_{w_0}, \partial_{w_1}, \partial_{w_2} \rangle$ and the following characterization of the canonical V -filtration of the holonomic module $\check{M}_A^0 = \Gamma(\mathbb{C}^3, \check{\mathcal{M}}_A^0)$, copied from [MM04, Definition 4.3-3, Proposition 4.3-9] (notice again that we work at the level of global sections): For any $m \in \check{M}_A^0$ we consider its Bernstein-Sato polynomial $b_m(x) \in \mathbb{C}[x]$, which is the unique monic polynomial of smallest degree satisfying the functional equation $b_m(\partial_{w_0} w_0) m \in w_0 \cdot V^0(D_{\mathbb{C}^3}) m$. The set of roots of $b_m(x)$ is denoted by $\text{ord}(m)$. Then we have

$$V^\alpha \check{M}_A^0 := \left\{ m \in \check{M}_A^0 \mid \text{ord}(m) \subset [\alpha, \infty) \right\}.$$

Our first step is to compute the Bernstein-Sato polynomial for the class $[1] \in \check{M}_A^0$. Here we have the following result.

Proposition 2.3 (Compare Lemma 4.4 below for the general case). *Consider the class of $1 \in D_{\mathbb{C}^3}$ in the quotient \check{M}_A^0 , denoted by $[1]$. Then we have $b_{[1]}(s) = s^2$.*

Proof. It is sufficient to find a functional equation in $D_{\mathbb{C}^3}$ of the form $(\partial_{w_0} w_0)^2 = w_0 \cdot P + \check{I}_A^0$, where

$$\check{I}_A^0 := (w_0^2 - w_1 w_2, \partial_{w_0} w_0 + \partial_{w_1} w_1 + \partial_{w_2} w_2, \partial_{w_1} w_1 - \partial_{w_2} w_2)$$

and where $P \in \mathbb{C}[w_0, w_1, w_2] \langle w_0 \partial_{w_0}, \partial_{w_1}, \partial_{w_2} \rangle$. We will show that

$$(-\partial_{w_1} w_1 - \partial_{w_2} w_2)^2 \in w_0 \cdot \mathbb{C}[w_1, w_2] \langle \partial_{w_1}, \partial_{w_2} \rangle + \check{I}_A^0, \quad (3)$$

which suffices to conclude. We have

$$\begin{aligned} (-\partial_{w_1} w_1 - \partial_{w_2} w_2)^2 &= 2\partial_{w_1} \partial_{w_2} w_1 w_2 + (\partial_{w_1} w_1)^2 + (\partial_{w_2} w_2)^2 \\ &\equiv 2w_0^2 \cdot \partial_{w_1} \partial_{w_2} + (\partial_{w_1} w_1)^2 + (\partial_{w_2} w_2)^2 \pmod{\check{I}_A^0} \\ &= 2w_0^2 \cdot \partial_{w_1} \partial_{w_2} + (\partial_{w_1} w_1 - \partial_{w_2} w_2)^2 + 2\partial_{w_1} \partial_{w_2} w_1 w_2 \\ &\equiv w_0^2 \cdot 4 \cdot \partial_{w_1} \partial_{w_2} \pmod{\check{I}_A^0}, \end{aligned}$$

which shows Formula (3). □

Notice that the above calculation can be extended to any matrix \tilde{A} of the form

$$\tilde{A} = \begin{pmatrix} 1 & 1 & \dots & 1 \\ 0 & a_{11} & \dots & a_{1n} \\ \vdots & \vdots & & \vdots \\ 0 & a_{d1} & \dots & a_{dn} \end{pmatrix}$$

where the columns of the matrix

$$A = \begin{pmatrix} a_{11} & \dots & a_{1n} \\ \vdots & & \vdots \\ a_{n1} & \dots & a_{nn} \end{pmatrix}$$

are the primitive integral generators of the fan of a smooth projective toric Fano manifold. Then one has to consider the classical cohomology algebra of this manifold, which admits a toric description (see, e.g. [Ful93, Section 5.2]), and the functional equation (i.e. the analogue of Formula 3) can be deduced from the relations in this algebra. Notice also that this is in fact an argument which is a much simplified version of the one used to prove Lemma 4.4 below (actually, the proof of this lemma relies on the main result of the separate paper [RSW18], which is based on general arguments from toric algebra, such as Euler-Koszul complexes, toric modules etc.). Lemma 4.4 is also more general in the sense that the matrix considered there is not necessarily defined by the rays of a smooth toric variety.

We have the following consequence of the above calculation which gives complete control on the integer part of the canonical V -filtration on \check{M} .

Corollary 2.4 (Compare Proposition 4.8 below for the general case). *Denote by $V_{ind}^\bullet \check{M}_A^0$ the filtration induced on \check{M}_A^0 by the V -filtration (with respect to w_0) on $D_{\mathbb{C}^3}$. Then for all $k \in \mathbb{Z}$, we have $V_{ind}^k \check{M}_A^0 = V^k \check{M}_A^0$.*

Proof. The proof is more or less similar to the general case in Proposition 4.8 below. In the case $k \geq 0$ any element $[P] \in V_{ind}^k \check{M}_A^0$ has an expression

$$[P] = \left[\sum_{i=0}^l w_0^k (w_0 \partial_{w_0})^i \cdot P_i \right] + [R]$$

where $P_i \in \mathbb{C}[w_1, w_2] \langle \partial_{w_1}, \partial_{w_2} \rangle$ and where $[R] \in V_{ind}^{k+1} \check{M}_A^0$. On the other hand, if $k > 0$, we can always write an element $[P] \in V_{ind}^{-k} \check{M}_A^0$ as

$$[P] = \left[\sum_{i=0}^l \partial_{w_1}^k (w_1 \partial_{w_1})^i \cdot P_i \right] + [R].$$

where P_i and $[R]$ are as above. One easily deduces (see the calculations in the proof of Proposition 4.8 below) from the functional equation $(\partial_{w_0} w_0)^2 [1] \in V_{ind}^1 \check{M}_A^0$ proved in Proposition 2.3 above that we have

$$\begin{aligned} (\partial_{w_0} w_0 - k)^2 [P] &\in V_{ind}^{k+1} \check{M}_A^0 && \text{for } k \geq 0 \text{ and } [P] \in V_{ind}^k \check{M}_A^0 \\ (\partial_{w_0} w_0 + k)^2 [P] &\in V_{ind}^{-k+1} \check{M}_A^0 && \text{for } k > 0 \text{ and } [P] \in V_{ind}^{-k} \check{M}_A^0 \end{aligned}$$

General considerations on the canonical V -filtration (see, e.g., [MM04, section 4.2 and 4.3] and the argument in the proof of Proposition 4.8) then imply that $V_{ind}^k \check{M}_A^0 = V^k \check{M}_A^0$. \square

Finally, we arrive at the following first main step toward the proof of theorem 2.1.

Proposition 2.5 (Compare Theorem 4.17 below for the general case). *Let \tilde{A} and $h_{\tilde{A}} : (\mathbb{C}^*)^2 \rightarrow \mathbb{C}^3 = W$ be as above, then for any $k \in \mathbb{Z}$ we have the following isomorphism of \mathcal{O}_W -modules*

$$F_p^H h_{\tilde{A},+} \mathcal{O}_{(\mathbb{C}^*)^2} \cong F_{p-1}^{ord} [\mathcal{D}_W / (w_0^2 - w_1 w_2, \partial_{w_0} w_0 + \partial_{w_1} w_1 + \partial_{w_2} w_2, \partial_{w_1} w_1 - \partial_{w_2} w_2)]$$

Proof. The main tool to obtain a description of the Hodge filtration is Formula (2) from above (see [Sai93, Proposition 4.2.]) which at the level of global sections reads

$$F_p^H \check{M}_A^0 = \sum_{i \geq 0} \partial_{w_0}^i \left(V^0 \check{M}_A^0 \cap F_{p-i}^H {}^* \check{M}_A^0 \right). \quad (4)$$

Recall from Lemma 2.2 that for all $l \in \mathbb{Z}$ we have

$$F_l^H {}^* \check{M}_A^0 = F_{l-1}^{ord} \check{M}_A^0.$$

In particular, since $F_p^{ord} \check{M}_A^0 = 0$ for all $p < 0$, we have $F_p^H \check{M}_A^0 = 0$ for all $p < 1$. On the other hand, we have seen in Proposition 2.3 that $[1] \in V^0 \check{M}_A^0$. Obviously, we have $[1] \in F_0^{ord} \check{M}_A^0$, which implies that $[1] \in F_1^H \check{M}_A^0$. Since \check{M}_A^0 is a cyclic D_W -module and since both filtrations F^H and F^{ord} on it are good filtrations, we obtain the inclusion

$$F_{p-1}^{ord} \check{M}_A^0 \subset F_p^H \check{M}_A^0$$

for all $p \in \mathbb{Z}$. It remains to show the reverse inclusion $F_p^H \check{M}_A^0 \subset F_{p-1}^{ord} \check{M}_A^0$. Using formula (4) as well as Corollary 2.4 and Lemma 2.2 this amounts to

$$\sum_{i \geq 0} \partial_{w_0}^i \left(V_{ind}^0 \check{M}_A^0 \cap F_{p-i}^{ord} \check{M}_A^0 \right) \subset F_p^{ord} \check{M}_A^0.$$

We obviously have $\partial_{w_0}^i F_{p-i}^{ord} \check{M}_A^0 \subset F_p^{ord} \check{M}_A^0$ for all i so it only remains to show that

$$V_{ind}^0 \check{M}_A^0 \cap F_l^{ord} \check{M}_A^0 \subset F_l^{ord} \check{M}_A^0$$

for all $l \in \mathbb{Z}$. Consider any class $[P] \in V_{ind}^0 \check{M}_A^0 \cap F_l^{ord} \check{M}_A^0$. Since we have ${}^* \check{M}_A^0 = \check{M}_A^0[w_0^{-1}]$, we can write

$$P = w_0^{-k} P_k + w_0^{-k+1} P_{-k+1} + \dots,$$

where $P_i \in \mathbb{C}[w_1, w_2] \langle \partial_{w_0}, \partial_{w_1}, \partial_{w_2} \rangle$. It follows that $w_0^k \cdot [P] \in V_{ind}^k \check{M}_A^0 \cap F_l^{ord} \check{M}_A^0$. We thus have to prove

$$V_{ind}^k \check{M}_A^0 \cap F_l^{ord} \check{M}_A^0 \subset w_0^k F_l^{ord} \check{M}_A^0.$$

for all $k, l \in \mathbb{Z}$. Take any class $[Q] \in V_{ind}^k \check{M}_A^0 \cap F_l^{ord} \check{M}_A^0$, then suppose that we can find a representative $Q \in V^k D_W \cap F_l D_W$ of $[Q]$. This means that $Q = w_0^k \cdot \tilde{Q}$, with $\tilde{Q} \in F_l D_W$, as required. Hence we obtain $[Q] \in w_0^k F_l^{ord} \check{M}_A^0$. It thus remains to show the existence of such a representative $Q \in V^k D_W \cap F_l D_W$, and this is exactly the content of the next lemma. \square

Lemma 2.6 (Compare Proposition 4.9 below for the general case). *Let A be as above, then for all $k, l \in \mathbb{Z}$, the morphism*

$$V^k D_W \cap F_l D_W \longrightarrow V_{ind}^k \check{M}_A^0 \cap F_l^{ord} \check{M}_A^0$$

is surjective.

Proof. The proof relies on the theory of Gröbner bases in the Weyl algebra. We will not give any definition here, but we refer to subsection 4.1 for details about monomial orders and Gröbner bases in the non-commutative setup.

Consider any class $m \in V_{ind}^k \check{M}_A^0 \cap F_l^{ord} \check{M}_A^0$. Then we can find $P \in F_l D_W$ and $Q \in V^k D_W$ such that $[P] = [Q] = m$, that is, $P = Q + i$ for some

$$i \in \check{I}_A^0 = (w_0^2 - w_1 w_2, \partial_{w_0} w_0 + \partial_{w_1} w_1 + \partial_{w_2} w_2, \partial_{w_1} w_1 - \partial_{w_2} w_2).$$

We chose a minimal $r \in \mathbb{N}$ with $Q \in F_r D_W$. If $r \leq l$, we are done since then $Q \in V^k D_W \cap F_l D_W$ is the preimage of m we are looking for. Hence suppose $r > l$. It is easy to see that then $i \in F_r D_W$ and the class of i in $\text{Gr}_r^F D_W$ is non-zero. For any operator $R \in D_W$, write

$$\sigma(R) \in \text{Gr}_\bullet^F D_W = \mathbb{C}[w_0, w_1, w_2, \xi_0, \xi_1, \xi_2] =: \mathbb{C}[\underline{w}, \underline{\xi}]$$

for its symbol. The three generators of \check{I}_A^0 from above form a Gröbner basis of this ideal with respect to the partial ordering given by the weight vector $(0, 1)$ (i.e. where w_i has weight 0 and ∂_{w_i} has weight 1), notice that this weight vector induces the filtration F_\bullet on D_W . This can be directly shown by a Macaulay2 calculation, the corresponding general result is Corollary 4.13 below, where we treat a slightly different situation however (we add a column to our matrix which is the sum of all other columns, this corresponds to composing with the graph embedding of the equation of a normal crossing divisor, see also the remarks before Lemma 4.14 above).

We can therefore conclude that there is an expression

$$\sigma(i) = \tilde{i}_1 \cdot (w_0^2 - w_1 w_2) + \tilde{i}_2 \cdot (\xi_0 w_0 + \xi_1 w_1 + \xi_2 w_2) + \tilde{i}_3 \cdot (\xi_1 w_1 - \xi_2 w_2).$$

with $\tilde{i}_1, \tilde{i}_2, \tilde{i}_3 \in \mathbb{C}[\underline{w}, \underline{\xi}]$. Let $i_1, i_2, i_3 \in D_W$ be the normally ordered operators obtained from $\tilde{i}_1, \tilde{i}_2, \tilde{i}_3$ by replacing ξ_k by ∂_{w_k} . Then we define

$$i' := i_1(w_0^2 - w_1 w_2) + i_2(\partial_{w_0} w_0 + \partial_{w_1} w_1 + \partial_{w_2} w_2) + i_3(\partial_{w_1} w_1 - \partial_{w_2} w_2) \in \check{I}_A^0$$

and clearly $i' \in F_l D_W$. However, we also have $i' \in V^k D_W$, since it can again be shown by a direct computation that the three polynomials $w_0^2 - w_1 w_2, \xi_0 w_0 + \xi_1 w_1 + \xi_2 w_2, \xi_1 w_1 - \xi_2 w_2$ form a Gröbner basis of the ideal they generate with respect to a partial ordering given by the weight vector $(-1, 0, 0, 1, 0, 0)$ (i.e. the weight of w_0 is -1 , the weight of ξ_0 is 1 and all other weights are zero). Notice again that this weight vector yields the filtration induced from $V^\bullet D_W$ on $\mathbb{C}[\underline{w}, \underline{\xi}]$. The corresponding general result (for the case of the extended matrix with one added column) is found in Corollary 4.13, 2. below.

Summarizing, we obtain that the operator $Q - i'$ satisfies

1. $[Q - i'] = [Q] = m$.

2. $Q - i' \in V^k D_W$
3. $Q - i' \in F_{l-1} D_W$ (this follows from $\sigma(Q) = \sigma(i) = \sigma(i')$).

Hence we see by descending induction on l that we can construct an operator $Q' \in F_l D_W \cap V^k D_W$ such that $[Q'] = m$. This shows the statement. \square

We have finished the proof of Proposition 2.5 above, which roughly summarizes the content of section 4 in the main body of the paper for our particular example. We now turn to the statements corresponding to section 5 below (for our example), that is, we are going to complete the proof of Theorem 2.1. Recall that the GKZ system $\mathcal{M}_{\tilde{A}}^0$ can be described as a Fourier-Laplace transform of a torus embedding

$$\mathcal{M}_{\tilde{A}}^0 \simeq \text{FL}(h_{\tilde{A},+} \mathcal{O}_{(\mathbb{C}^*)^2})$$

Since the matrix \tilde{A} is homogeneous (i.e. $(1, \dots, 1)$ is in its row span) the \mathcal{D} -module $\mathcal{M}_{\tilde{A}}^0$ has a different presentation involving only (proper) direct image functors and inverse image functors but excluding the use of the Fourier-Laplace transformation (see [Rei14, Proposition 2.7(iii)]). Let us recall some ingredients of this construction in the present situation. Consider the torus embedding

$$\begin{aligned} g : \mathbb{C}^* &\longrightarrow \mathbb{P}^2 \\ t &\mapsto (1 : t : t^{-1}), \end{aligned}$$

then $\mathcal{M}_{\tilde{A}}^0$ can be described by a Radon type transform of the \mathcal{D} -module $g_+ \mathcal{O}_{\mathbb{C}^*}$. More precisely, we have a commutative diagram

$$\begin{array}{ccccc} & & U & & \\ & \swarrow \pi_1^U & \downarrow j_U & \searrow \pi_2^U & \\ \mathbb{P}^2 & \xleftarrow{\pi_1} & \mathbb{P}^2 \times \mathbb{C}^3 & \xrightarrow{\pi_2} & \mathbb{C}^3 \end{array}$$

where U is the complement of the universal hyperplane in $\mathbb{P}^2 \times \mathbb{C}^3$, i.e. $U := \{\lambda_0 w_0 + \lambda_1 w_1 + \lambda_2 w_2 \neq 0\}$. The GKZ-system $\mathcal{M}_{\tilde{A}}^0$ is now given by

$$\mathcal{M}_{\tilde{A}}^0 \simeq \mathcal{R}_c^\circ(g_+ \mathcal{O}_{\mathbb{C}^*}) \simeq \pi_{2,+}^U \pi_1^{U,\dagger} g_+ \mathcal{O}_{\mathbb{C}^*} \simeq \pi_{2,+} \left(\pi_1^\dagger g_+ \mathcal{O}_T \otimes j_{U,\dagger} \mathcal{O}_U \right)$$

where the last isomorphism follows from the projection formula.

Since $\mathcal{O}_{\mathbb{C}^*}$ carries a trivial Hodge module structure and since the category of algebraic mixed Hodge modules is stable under the (proper) direct image functor and the (exceptional) inverse image functor this induces a mixed Hodge module structure on $\mathcal{M}_{\tilde{A}}^0$.

For technical reasons that mainly occur when dealing with the case $\beta \neq 0$ as we do in the main body of this paper, we will pursue a slightly different approach.

Consider the map $F : \mathbb{C}^* \times \mathbb{C}^3 \rightarrow \mathbb{C}$ given by the Laurent polynomial $\lambda_0 + \lambda_1 t + \lambda_2 t^{-1}$ and let $j : \mathbb{C}^* \rightarrow \mathbb{C}$ the canonical embedding. Denote by p resp. q the projection from $\mathbb{C}^* \times \mathbb{C}^3$ to the second resp. first factor. We consider the integral transform of $\mathcal{O}_{\mathbb{C}^*}$ from \mathbb{C}^* to \mathbb{C}^3 with kernel $F^\dagger j_! \mathcal{O}_{\mathbb{C}^*}$ and prove in Proposition 5.4 that this integral transform is isomorphic to the GKZ-system $\mathcal{M}_{\tilde{A}}^0$ (In the more general case of a non-zero $\tilde{\beta} = (\beta_0, \beta)$ we would start with $\mathcal{O}_{\mathbb{C}^*}^{\tilde{\beta}}$ and use the kernel $F^\dagger j_! \mathcal{O}_{\mathbb{C}^*}^{\beta_0}$).

Since this integral transformation preserves the category of mixed Hodge modules we define a Hodge module structure on the GKZ system by

$${}^H \mathcal{M}_{\tilde{A}}^0 := \mathcal{H}^6(p_*(q^{*p} \mathbb{C}_{\mathbb{C}^*} \otimes F^* j_!^p \mathbb{C}_{\mathbb{C}^*})) \quad (5)$$

In proposition 5.5 we prove that this approach coincides with the Radon transform for integer β_0 .

In order to compute the Hodge filtration on (5) explicitly we have to consider a partial compactification of $\mathbb{C}^* \times \mathbb{C}^3$, since the projection $p : \mathbb{C}^* \times \mathbb{C}^2$ is not proper. For this we use the locally closed embedding $g : \mathbb{C}^* \rightarrow \mathbb{P}^2$. As an intermediate step we compute the Hodge filtration on the mixed Hodge module

$${}^H \mathcal{N} := (g \times id)_*(q^{*p} \mathbb{C}_{\mathbb{C}^*} \otimes F^* j_!^p \mathbb{C}_{\mathbb{C}^*})$$

The space $\mathbb{P}^2 \times \mathbb{C}^3$ is covered by the three charts $W_u = \{w_u \neq 0\}$ for $u = 0, 1, 2$. The map g factors over each chart and is given by

$$\begin{aligned} g_0 : \mathbb{C}^* &\longrightarrow W_0 & g_1 : \mathbb{C}^* &\longrightarrow W_1 & g_2 : \mathbb{C}^* &\longrightarrow W_2 \\ t &\mapsto (t, t^{-1}) & t &\mapsto (t^{-1}, t^{-2}) & t &\mapsto (t, t^2) \end{aligned}$$

We obtain in Formula (41) below that the restriction of ${}^H\mathcal{N}$ to W_u can be written as a direct product

$${}^H\mathcal{N}_u = \mathcal{H}^0(p g_{u*} \mathbb{C}_{\mathbb{C}^*}^H) \boxtimes \mathcal{H}^2(pr_u j_!^p \mathbb{C}_{\mathbb{C}^*}^H)$$

where $pr_u : \mathbb{C}^3 \rightarrow \mathbb{C}$ is the projection to the $(u+1)$ -th factor. The Hodge filtration on the first factor can be computed by using Theorem 4.17, the computation of the second factor is straightforward (cf. Remark 4.18) (we check the assumption of Theorem 4.17 in Lemma 5.8).

Define the matrices

$$A_0^s := \begin{pmatrix} 0 & 0 & 1 & 1 & 1 \\ 1 & -1 & 0 & 1 & -1 \end{pmatrix} \quad A_1^s := \begin{pmatrix} 0 & 0 & 1 & 1 & 1 \\ -1 & -2 & 0 & 1 & -1 \end{pmatrix} \quad A_2^s := \begin{pmatrix} 0 & 0 & 1 & 1 & 1 \\ 1 & 2 & 0 & 1 & -1 \end{pmatrix}$$

We show in Lemma 5.13 that the \mathcal{D} -module underlying ${}^H\mathcal{N}_u$ is isomorphic to a partial Laplace transformation of $\mathcal{M}_{A_u^s}^0$ in the w -variables. More precisely we have

$$\mathcal{N}_0 = \mathcal{D}_{\mathbb{C}^2 \times \mathbb{C}^3} / \overset{(\vee)}{\mathcal{I}}_{A_0^s}, \quad \mathcal{N}_1 = \mathcal{D}_{\mathbb{C}^2 \times \mathbb{C}^3} / \overset{(\vee)}{\mathcal{I}}_{A_1^s}, \quad \mathcal{N}_2 = \mathcal{D}_{\mathbb{C}^2 \times \mathbb{C}^3} / \overset{(\vee)}{\mathcal{I}}_{A_2^s}$$

where $\overset{(\vee)}{\mathcal{I}}_{A_0^s}$ is generated by Euler operators

$$\overset{(\vee)}{E}_0^0 := \lambda_0 \partial_{\lambda_0} + \lambda_1 \partial_{\lambda_1} + \lambda_2 \partial_{\lambda_2}, \quad \overset{(\vee)}{E}_1^0 := -w_{10} \partial_{w_{10}} + w_{20} \partial_{w_{20}} + \lambda_1 \partial_{\lambda_1} - \lambda_2 \partial_{\lambda_2}$$

and the box operators

$$\overset{(\vee)}{\square}_{(1,1,0,0,0)} := w_{10} w_{20} - 1, \quad \overset{(\vee)}{\square}_{(0,0,2,-1,-1)} := \partial_{\lambda_0}^2 - \partial_{\lambda_1} \partial_{\lambda_2}, \quad \overset{(\vee)}{\square}_{(0,1,-1,1,0)} := w_{20} \partial_{\lambda_1} - \partial_{\lambda_0}$$

The ideal $\overset{(\vee)}{\mathcal{I}}_{A_1^s}$ is generated by Euler operators

$$\overset{(\vee)}{E}_0^1 := \lambda_0 \partial_{\lambda_0} + \lambda_1 \partial_{\lambda_1} + \lambda_2 \partial_{\lambda_2}, \quad \overset{(\vee)}{E}_1^1 := w_{01} \partial_{w_{01}} + 2w_{21} \partial_{w_{21}} + \lambda_1 \partial_{\lambda_1} - \lambda_2 \partial_{\lambda_2}$$

and box operators

$$\overset{(\vee)}{\square}_{(2,-1,0,0,0)} := w_{01}^2 - w_{21}, \quad \overset{(\vee)}{\square}_{(0,0,2,-1,-1)} := \partial_{\lambda_0}^2 - \partial_{\lambda_1} \partial_{\lambda_2}, \quad \overset{(\vee)}{\square}_{(1,0,-1,1,0)} := w_{01} \partial_{\lambda_1} - \partial_{\lambda_0}$$

The ideal $\overset{(\vee)}{\mathcal{I}}_{A_2^s}$ is generated by Euler operators

$$\overset{(\vee)}{E}_0^2 := \lambda_0 \partial_{\lambda_0} + \lambda_1 \partial_{\lambda_1} + \lambda_2 \partial_{\lambda_2}, \quad \overset{(\vee)}{E}_1^2 := -w_{02} \partial_{w_{01}} - 2w_{12} \partial_{w_{21}} + \lambda_1 \partial_{\lambda_1} - \lambda_2 \partial_{\lambda_2}.$$

and box operators

$$\overset{(\vee)}{\square}_{(2,-1,0,0,0)} := w_{02}^2 - w_{12}, \quad \overset{(\vee)}{\square}_{(0,0,2,-1,-1)} := \partial_{\lambda_0}^2 - \partial_{\lambda_1} \partial_{\lambda_2}, \quad \overset{(\vee)}{\square}_{(1,0,1,-1,0)} := w_{02} \partial_{\lambda_0} - \partial_{\lambda_1}$$

The Hodge filtration on these systems is given by $F_{p+1}^H \mathcal{N}_u = F_p^{ord} \mathcal{D}_{W_u \times \mathbb{C}^3} / \overset{(\vee)}{\mathcal{I}}_{A_u^s}$.

Using the fact that \mathcal{N}_u is a partial Fourier-Laplace transform of $\mathcal{M}_{A_u^s}$ we use the results of section 3.1 to construct a strict resolution of $(\mathcal{N}_u, F_{\bullet}^H)$ at the level of global sections which is given by the Euler-Koszul complex

$$\overset{(\vee)}{K}_u^{\bullet} := D_{W_u \times \mathbb{C}^3} / \overset{(\vee)}{J}_{A_u^s} \xrightarrow{\cdot (-\overset{(\vee)}{E}_1^u, \overset{(\vee)}{E}_0^u)^t} (D_{W_u \times \mathbb{C}^3} / \overset{(\vee)}{J}_{A_u^s})^2 \xrightarrow{\cdot (\overset{(\vee)}{E}_0^u, \overset{(\vee)}{E}_1^u)} D_{W_u \times \mathbb{C}^3} / \overset{(\vee)}{J}_{A_u^s}$$

where the left ideal $J_{A_u^s}$ is generated by the "box-type" generators from above. It remains to compute the projection of $(\mathcal{N}, F_\bullet^H)$ under the map $\mathbb{P}^2 \times \mathbb{C}^3 \rightarrow \mathbb{C}^3$. In order to do this we lift the filtered D -modules (N_u, F_\bullet^H) as well as their strictly filtered resolution $(\overset{(\vee)}{K}_u^\bullet, F_\bullet)$ to the category of $R_{\mathcal{W}_u \times \mathcal{C}^3}$ -modules where $\mathcal{W}_u \times \mathcal{C}^3 := \mathbb{C}_z \times W_u \times \mathbb{C}^3$ and

$$R_{\mathcal{W}_u \times \mathcal{C}^3} = \mathbb{C}[z, (w_{iu})_{i \neq u}, \lambda_0, \lambda_1, \lambda_2] \langle (z \partial_{w_{iu}})_{i \neq u}, z \partial_{\lambda_0}, z \partial_{\lambda_1}, z \partial_{\lambda_2} \rangle$$

This is done by the Rees construction, i.e. we associate to the filtered $D_{W_u \times \mathbb{C}^3}$ -modules (N_u, F_\bullet^H) resp. $(\overset{(\vee)}{K}_u^\bullet, F_\bullet)$ the $R_{\mathcal{W}_u \times \mathcal{C}^3}$ -modules

$$N_u := R_F N_u = \bigoplus_{p \in \mathbb{Z}} F_p N_u z^p, \quad K_u^\bullet := R_F \overset{(\vee)}{K}_u^\bullet$$

and similarly for the filtered $\mathcal{D}_{\mathbb{P}^2 \times \mathbb{C}^3}$ -module $(\mathcal{N}, F_\bullet^H)$ to which we associate $\mathcal{N} := \mathcal{O}_{\mathcal{P} \times \mathcal{C}^3} \otimes_{\mathcal{O}_{\mathbb{P}^2 \times \mathbb{C}^3}[z]} R_{F^H} \mathcal{N}$, where $\mathcal{P} \times \mathcal{C}^3 := \mathbb{C}_z \times \mathbb{P}^2 \times \mathbb{C}^3$.

Instead of computing the projection of the filtered \mathcal{D} -module $(\mathcal{N}, F_\bullet^H)$ we compute the projection of the \mathcal{R} -module \mathcal{N} . This is given by

$$\pi_{2+} \mathcal{N} \simeq R\pi_{2*} DR_{\mathcal{P} \times \mathcal{C}^3 / \mathcal{C}^3}(\mathcal{N}),$$

where this time π_2 denotes the map $\mathcal{P} \times \mathcal{C}^3 \rightarrow \mathcal{C}^3$.

Since this is hard to compute directly we construct a resolution \mathcal{K}^\bullet from the local resolutions K_u^\bullet and get the double complex $\Omega_{\mathcal{P} \times \mathcal{C}^3 / \mathcal{C}^3}^{\bullet+2} \otimes \mathcal{K}^\bullet$:

$$\begin{array}{ccccc} \mathcal{K}^0 & \xrightarrow{II d^{1,0}} & \Omega_{\mathcal{P} \times \mathcal{C}^3 / \mathcal{C}^3}^1 \otimes \mathcal{K}^0 & \xrightarrow{II d^{2,0}} & \Omega_{\mathcal{P} \times \mathcal{C}^3 / \mathcal{C}^3}^2 \otimes \mathcal{K}^0 \\ \uparrow Id^{0,0} & & \uparrow Id^{1,0} & & \uparrow Id^{2,0} \\ \mathcal{K}^{-1} & \xrightarrow{II d^{1,-1}} & \Omega_{\mathcal{P} \times \mathcal{C}^3 / \mathcal{C}^3}^1 \otimes \mathcal{K}^{-1} & \xrightarrow{II d^{2,-1}} & \Omega_{\mathcal{P} \times \mathcal{C}^3 / \mathcal{C}^3}^2 \otimes \mathcal{K}^{-1} \\ \uparrow Id^{0,-1} & & \uparrow Id^{1,-1} & & \uparrow Id^{2,-1} \\ \mathcal{K}^{-2} & \xrightarrow{II d^{1,-2}} & \Omega_{\mathcal{P} \times \mathcal{C}^3 / \mathcal{C}^3}^1 \otimes \mathcal{K}^{-2} & \xrightarrow{II d^{2,-2}} & \Omega_{\mathcal{P} \times \mathcal{C}^3 / \mathcal{C}^3}^2 \otimes \mathcal{K}^{-2} \end{array}$$

This double complex gives rise to two spectral sequences: The first one is given by first taking cohomology in the vertical direction. This gives the IE_1 -page where only the $IE_1^{0,q}$ -terms are non-zero and are isomorphic to $\Omega_{\mathcal{P} \times \mathcal{C}^3 / \mathcal{C}^3}^{q+2} \otimes \mathcal{N}$. If we consider the second spectral sequence and take cohomology in the horizontal direction we get the $II E_1$ -page. Here $II E_1^{p,q} = 0$ for $q \neq 0$ and we set $\mathcal{L}^\bullet := II E_1^{p,0}$. Since both spectral sequences degenerate at the second page we get a quasi-isomorphism $\Omega_{\mathcal{P} \times \mathcal{C}^3 / \mathcal{C}^3}^{\bullet+2} \otimes \mathcal{N} \simeq \mathcal{L}^\bullet$ (cf. Proposition 5.20).

In order to get an explicit representation of \mathcal{L}^\bullet we introduce a sheaf of rings \mathcal{S} on $\mathcal{P} \times \mathcal{C}^3$ and an ideal $\mathcal{J} \subset \mathcal{S}$ which are locally given by

$$\Gamma(\mathcal{W}_u \times \mathcal{C}^3, \mathcal{S}) := S_{\mathcal{W}_u \times \mathcal{C}^3} := \mathbb{C}[z, \lambda_0, \lambda_1, \lambda_2, (w_{iu})_{i \neq u}] \langle z \partial_{\lambda_0}, z \partial_{\lambda_1}, z \partial_{\lambda_2} \rangle$$

resp.

$$\Gamma(\mathcal{W}_u \times \mathcal{C}^3, \mathcal{J}) = J_{A_u^s}$$

where $J_{A_u^s}$ is the left ideal in $S_{\mathcal{W}_u \times \mathcal{C}^3}$ generated by the corresponding box operators. Define the following Euler operators

$$\tilde{E}_0 = \lambda_0 z \partial_{\lambda_0} + \lambda_1 z \partial_{\lambda_1} + \lambda_2 z \partial_{\lambda_2}, \quad \tilde{E}_1 = \lambda_1 z \partial_{\lambda_1} - \lambda_2 z \partial_{\lambda_2}$$

we get the following quasi-isomorphism

$$\mathcal{L}^\bullet \simeq Kos^\bullet(z^{-1} \mathcal{S} / \mathcal{J}, (\tilde{E}_k)_{k=0,1})$$

hence we get $\pi_{2+}\mathcal{N} \simeq R\pi_{2*} \left(\text{Kos}^\bullet(z^{-1}\mathcal{S}/\mathcal{I}, (\tilde{E}_k)_{k=0,1}) \right)$.

Since $\mathcal{C}^3 = \mathbb{C}_z \times \mathbb{C}^3$ is affine it is enough to compute the global sections of $\pi_{2+}\mathcal{N}$ which are given by

$$R\Gamma \left(\text{Kos}^\bullet(z^{-1}\mathcal{S}/\mathcal{I}, (\tilde{E}_k)_{k=0,1}) \right).$$

We will show that each term in the Koszul complex is Γ -acyclic which boils down to the fact that \mathcal{S}/\mathcal{I} is Γ -acyclic.

Define the matrix

$$A^s := \begin{pmatrix} 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 \\ 0 & -1 & -1 & 0 & -1 & -1 \end{pmatrix}$$

and the ring

$$S := \mathbb{C}[z, w_0, w_1, w_2, \lambda_0, \lambda_1, \lambda_2] \langle z\partial_{\lambda_0}, z\partial_{\lambda_1}, z\partial_{\lambda_2} \rangle$$

Let J_{A^s} be the left ideal in S generated by

$$w_0^2 - w_1w_2, \quad \partial_{\lambda_0}^2 - \partial_{\lambda_1}\partial_{\lambda_2}, \quad w_1\partial_{\lambda_1} - w_0\partial_{\lambda_0}, \quad w_2\partial_{\lambda_2} - w_0\partial_{\lambda_0}$$

(these are box operators with respect to the matrix A^s). The associated sheaf $\widetilde{S/J_{A^s}}$ is isomorphic to \mathcal{S}/\mathcal{I} . The associated graded module is defined by

$$\Gamma_*(\mathcal{S}/\mathcal{I}) := \bigoplus_{a \in \mathbb{Z}} \Gamma(\mathcal{P} \times \mathcal{C}^3, (\mathcal{S}/\mathcal{I})(a))$$

The difference between S/J_{A^s} and the associated graded module is measured by local cohomology modules of S/J_{A^s} , more precisely we have (cf. Proposition 5.22)

$$0 \longrightarrow H_{(\underline{w})}^0(S/J_{A^s}) \longrightarrow S/J_{A^s} \longrightarrow \Gamma_*(\mathcal{S}/\mathcal{I}) \longrightarrow H_{(\underline{w})}^1(S/J_{A^s}) \longrightarrow 0$$

and

$$\bigoplus_{a \in \mathbb{Z}} H^i(\mathcal{P} \times \mathcal{C}^3, (\mathcal{S}/\mathcal{I})(a)) \simeq H_{(\underline{w})}^{i+1}(S/J_{A^s}) \quad (6)$$

where (\underline{w}) is the ideal in $\mathbb{C}[z, w_0, w_1, w_2, \lambda_0, \lambda_1, \lambda_2]$ generated w_0, w_1, w_2 . Notice that all terms involved carry a natural \mathbb{Z} -grading by setting $\deg(w_i) = 1$ and $\deg(\lambda_i) = \deg(\partial_{\lambda_i}) = 0$ for $i = 0, 1, 2$. The generators of J_{A^s} lie in the commutative subring $T := \mathbb{C}[w_0, w_1, w_2, \partial_{\lambda_0}, \partial_{\lambda_1}, \partial_{\lambda_2}] \subset S$. We denote by K_{A^s} the corresponding ideal in T . It is easily seen that the ring T/K_{A^s} is isomorphic to the semi-group ring $\mathbb{C}[\mathbb{N}A^s]$.

We prove that the local cohomology modules turning up in Formula (6) above can be rewritten as follows (cf. Lemma 5.24):

$$H_{(\underline{w})}^k(S/J_{A^s}) \simeq S \otimes_T H_I^k(\mathbb{C}[\mathbb{N}A^s])$$

where the ideal $I \subset \mathbb{C}[\mathbb{N}A^s]$ is generated by w_0, w_1, w_2 . Hence, we have reduced the problem to a well-known subject in commutative algebra, since the local cohomology groups $H_I^k(\mathbb{C}[\mathbb{N}A^s])$ can be explicitly computed by the so-called Ishida complex. Let σ be the face which is generated by the first three columns of A^s (the columns which correspond to variables w_0, w_1, w_2). For a face $\tau \subseteq \sigma$ we define the localization $\mathbb{C}[\mathbb{N}A^s]_\tau := \mathbb{C}[\mathbb{N}A^s + \mathbb{Z}(A^s \cap \tau)]$. Put

$$L_\tau^k := \bigoplus_{\substack{\tau \subset \sigma \\ \dim \tau = k}} \mathbb{C}[\mathbb{N}A^s]_\tau$$

The Ishida complex therefore takes the form

$$L_\sigma^\bullet : 0 \longrightarrow L_\sigma^0 \longrightarrow L_\sigma^1 \longrightarrow L_\sigma^2 \longrightarrow 0$$

We prove in Proposition 5.25 that $H_I^k(\mathbb{C}[\mathbb{N}A^s]) \simeq H^k(L_\sigma^\bullet)$. Finally we show in Corollary 5.31 that $H^k(L_\sigma^\bullet) = 0$ for $k \neq 2$ (so we have local cohomology only in the top degree) and that the \mathbb{Z} -degrees in

$H^2(L_\sigma^\bullet)$ are purely negative. We refer the reader to Example 5.27 for more details in this particular case.

We can therefore conclude that

$$S/J_{A^s} \simeq \Gamma_*(\mathcal{S}/\mathcal{J}) \quad \text{and} \quad H^i(\mathcal{P} \times \mathcal{C}^3, \mathcal{S}/\mathcal{J}) = 0 \quad \text{for all } i \geq 1$$

Putting things together we conclude that the global sections of $\pi_{2+}\mathcal{N}$ are given by $\Gamma(\text{Kos}^\bullet(z^{-1}\mathcal{S}/\mathcal{J}, (\tilde{E}_k)_{k=0,1}))$. The latter one can be easily computed and gives

$$\Gamma\mathcal{H}^0\pi_{2+}\mathcal{N} = z^{-1}R_{\mathcal{C}^3}/I_A^\lambda \quad \text{and} \quad \Gamma\mathcal{H}^i\pi_{2+}\mathcal{N} = 0 \quad \text{for } i \geq 1$$

where $R_{\mathcal{C}^3} := \mathbb{C}[z, \lambda_0, \lambda_1, \lambda_2]\langle z\partial_{\lambda_0}, z\partial_{\lambda_1}, z\partial_{\lambda_2} \rangle$ and the left ideal I_A^λ is generated by the box operator $(z\partial_{\lambda_0})^2 - (z\partial_{\lambda_1})(z\partial_{\lambda_2})$ and the Euler operators \tilde{E}_0, \tilde{E}_1 . But this shows that

$$(\mathcal{M}_A^0, F_\bullet^H) \simeq (\mathcal{M}_A^0, F_{\bullet+1}^{ord})$$

which is the statement of Theorem 2.1 resp. that of Theorem 5.35 below in the general case.

3 GKZ-systems and the Fourier-Laplace transformation

We start by introducing GKZ-systems as well as their Fourier-Laplace transformed versions. Throughout the whole paper, we let W be a finite-dimensional vector space over \mathbb{C} and denote by V its dual vector space. We will fix coordinates w_1, \dots, w_n on W and dual coordinates $\lambda_1, \dots, \lambda_n$ on V .

3.1 GKZ-systems and strict resolutions

Given a $d \times n$ integer matrix $A = (a_{ki})$ we denote by $\underline{a}_1, \dots, \underline{a}_n$ its columns. We define

$$\mathbb{N}A := \sum_{i=1}^n \mathbb{N}\underline{a}_i$$

and similarly for $\mathbb{Z}A$ and $\mathbb{R}_{\geq 0}A$. Throughout the paper we assume that the matrix A satisfies

$$\mathbb{Z}A = \mathbb{Z}^d.$$

Definition 3.1. *Let $A = (a_{ki})$ be a $d \times n$ integer matrix with $\mathbb{Z}A = \mathbb{Z}^d$ and $\beta = (\beta_1, \dots, \beta_d) \in \mathbb{C}^d$. Write \mathbb{L}_A for the \mathbb{Z} -module of integer relations among the columns of A and write \mathcal{D}_V for the sheaf of rings of differential operators on V . Define*

$$\mathcal{M}_A^\beta := \mathcal{D}_V/\mathcal{I}_A,$$

where \mathcal{I}_A is the sheaf of left ideals generated by

$$\square_{\underline{l}} := \prod_{i:l_i < 0} \partial_{\lambda_i}^{-l_i} - \prod_{i:l_i > 0} \partial_{\lambda_i}^{l_i}$$

for all $\underline{l} \in \mathbb{L}_A$ and

$$E_k - \beta_k := \sum_{i=1}^n a_{ki} \lambda_i \partial_{\lambda_i} - \beta_k$$

for $i = 1, \dots, d$.

Since GKZ-systems are defined on the affine space $V \cong \mathbb{C}^n$, we will often work with the D -modules of global sections $M_A^\beta := \Gamma(\mathbb{C}^n, \mathcal{M}_A^\beta)$ rather than with the sheaves themselves.

We will now discuss filtrations on GKZ-systems given by a weight vector $(u, v) \in \mathbb{Z}^{2n}$. This weight vector induces an increasing filtration on D_V given by

$$F_p^{(u,v)} D_V = \left\{ \sum_{\substack{\sum_i u_i \gamma_i + v_i \delta_i \leq p \\ \text{finite}}} c_{\gamma\delta} \lambda^\gamma \partial_\lambda^\delta \mid \gamma, \delta \in \mathbb{Z}_{\geq 0}^n \right\}$$

where we set $\lambda^\gamma := \prod_{i=1}^n \lambda_i^{\gamma_i}$ etc. . For an element $P = \sum c_{\gamma\delta} \lambda^\gamma \partial_\lambda^\delta$ we define $ord_{(u,v)}(P) := \max\{\sum_i u_i \gamma_i + v_i \delta_i \mid c_{\gamma\delta} \neq 0\}$. The associated graded ring $\text{Gr}_\bullet^{(u,v)} D_V$ is given by $\bigoplus_p F_p^{(u,v)} D_V / F_{p-1}^{(u,v)} D_V$.

In order to construct a strictly filtered resolution of \mathcal{M}_A^β , we use the theory of Euler-Koszul complexes as introduced in [MMW05]. We will work at the level of global sections. We briefly recall the definition of the Euler-Koszul complex $(K^\bullet, E - \beta)$ from [MMW05, Definition 4.2] (where it is called $\mathcal{K}_\bullet(E - \beta; \mathbb{C}[\mathbb{N}A])$) and placed in positive homological degrees). Its terms are given by

$$K^{-l} = \bigoplus_{0 \leq i_1 < \dots < i_l \leq l} (D_V / J_A) e_{i_1 \dots i_l},$$

where the left ideal $J_A \subset \mathbb{C}[\partial] := \mathbb{C}[\partial_{\lambda_1}, \dots, \partial_{\lambda_n}]$ is generated by

$$\square_{\underline{l}} := \prod_{i:l_i < 0} \partial_{\lambda_i}^{-l_i} - \prod_{i:l_i > 0} \partial_{\lambda_i}^{l_i}, \quad \forall \underline{l} \in \mathbb{L}_A$$

A simple computation using the fact that $\sum_{i=1}^n l_i a_{ki} = 0$ shows that the maps

$$\begin{aligned} D_V / D_V J_A &\longrightarrow D_V / D_V J_A \\ P &\mapsto P \cdot (E_k - \beta_k) \quad \text{for } k = 1, \dots, d \end{aligned} \quad (7)$$

are well defined. Moreover, we have $[E_{k_1} - \beta_{k_1}, E_{k_2} - \beta_{k_2}] = 0$ for $k_1, k_2 \in \{1, \dots, d\}$, and hence we can build the Koszul complex

$$(K^\bullet, E - \beta) = (\dots \xrightarrow{d_{-2}} K^{-1} \xrightarrow{d_{-1}} K^0 \rightarrow 0) := \text{Kos}(D_V / D_V J_A, (E_k - \beta_k)_{k=1, \dots, d}).$$

with D_V -linear differential

$$d_{-l}(e_{i_1 \dots i_l}) := \sum_{k=1}^l (-1)^{l-1} (E_{i_k} - \beta_{i_k}) e_{i_1 \dots \hat{i}_k \dots i_l}$$

If we assume that the semigroup $\mathbb{N}A$ satisfies

$$\mathbb{N}A = \mathbb{Z}^d \cap \mathbb{R}_{\geq 0} A$$

then by a classical result due to Hochster ([Hoc72, theorem 1]) it follows that the semigroup ring $\mathbb{C}[\mathbb{N}A]$ is Cohen-Macaulay. It was shown in [MMW05, Remark 6.4] that in this case $(K^\bullet, E - \beta)$ is a resolution of M_A^β for all $\beta \in \mathbb{C}^d$.

Notice that the filtration $F_\bullet^{(u,v)}$ on D_V induces a filtration on $D_V / D_V J_A$ which we denote by the same symbol. We define the following filtration on each term of the Koszul complex $(K^\bullet, E - \beta)$:

$$F_p^{(u,v)} K^{-l} := \bigoplus_{0 \leq i_1 < \dots < i_l \leq l} F_{p - \sum_{k=1}^l c_{i_k}}^{(u,v)} (D_V / D_V J_A) e_{i_1 \dots i_l},$$

where $c_i = ord_{(u,v)}(E_i - \beta_i)$. This shows that the complex $((K^\bullet, E - \beta), F_\bullet^{(u,v)})$ is filtered, i.e. that the differential d respects the filtration

$$d_{-l}(F_p^{(u,v)} K^{-l}) \subset F_p^{(u,v)} d_{-l}(K^{-l}) := \text{im}(d_{-l}) \cap F_p^{(u,v)} K^{-l+1}.$$

We recall the following well-known criterion for a complex to be strictly filtered, which means

$$d_{-l}(F_p^{(u,v)} K^{-l}) = F_p^{(u,v)} d_{-l}(K^{-l}) = \text{im}(d_{-l}) \cap F_p^{(u,v)} K^{-l+1}$$

Lemma 3.2. *Let*

$$0 \longrightarrow (M_1, F) \xrightarrow{d_1} \dots \xrightarrow{d_{n-1}} (M_n, F) \rightarrow 0$$

be a sequence of filtered D -modules with bounded below filtration. The following properties are equivalent.

1. The map d_k is strict.
2. $H^k(F_p M_\bullet) \simeq F_p H^k(M_\bullet)$ for all p .
3. $H^k(\text{gr}_p^F M_\bullet) \simeq \text{Gr}_p^F H^k(M_\bullet)$ for all p .

Remark 3.3. Suppose that we have $\mathbf{NA} = \mathbb{Z}^d \cap \mathbb{R}_{\geq 0} A$, then in order to prove that the filtered complex $((K^\bullet, E - \beta), F_\bullet^{(u,v)})$ is strict it is enough to show that $H^{-l}(\text{Gr}_\bullet^{(u,v)} K^\bullet) = 0$ for $l > 1$ and $H^0(\text{Gr}_\bullet^{(u,v)} K^\bullet) = \text{Gr}_\bullet^{(u,v)} M_A^\beta$, since we already know that $H^{-l}(K^\bullet) = 0$ for $l > 1$ and $H^0(K^\bullet) = M_A^\beta$.

3.2 Fourier-Laplace transformed GKZ-systems

Let as before W be a n -dimensional vector space over \mathbb{C} and denote by V its dual vector space. Let X be a smooth algebraic variety and $E = X \times W$ be a trivial vector bundle and $E' := X \times V$ its dual. We write $\langle, \rangle : W \times V \rightarrow \mathbb{C}$ for the canonical pairing which extends to a function $\langle, \rangle : E \times E' \rightarrow \mathbb{C}$.

Definition 3.4. Define $\mathcal{L} := \mathcal{O}_{E \times_X E'} e^{-\langle, \rangle}$ which is by definition the free rank one module with differential given by the product rule. Denote by $p_1 : E \times_X E' \rightarrow E$, $p_2 : E \times_X E' \rightarrow E'$ the canonical projections. For $\mathcal{M} \in D_h^b(\mathcal{D}_E)$ the Fourier-Laplace transformation is then defined by

$$\text{FL}_X(\mathcal{M}) := p_{2+}(p_1^+ \mathcal{M} \otimes^L \mathcal{L})[-n]$$

Definition 3.5. Let $A = (a_{ki})$ be a $d \times n$ integer matrix. Let $\beta \in \mathbb{C}^d$. Write \mathbb{L}_A for the \mathbb{Z} -module of relations among the columns of A and write \mathcal{D}_W for the sheaf of rings of algebraic differential operators on W . Define

$$\check{\mathcal{M}}_A^\beta := \mathcal{D}_W / ((\check{\square}_{\underline{m}})_{\underline{m} \in \mathbb{L}_A}, (\check{E}_k + \beta_k)_{k=1, \dots, d}),$$

where

$$\begin{aligned} \check{E}_k &:= \sum_{i=1}^n a_{ki} \partial_{w_i} w_i \quad \text{for } k = 1, \dots, d \\ \check{\square}_{\underline{m} \in \mathbb{L}_A} &:= \prod_{m_i > 0} w_i^{m_i} - \prod_{m_i < 0} w_i^{-m_i}. \end{aligned} \tag{8}$$

Again we will often work with the D_W -module of global sections

$$\check{M}_A^\beta := \Gamma(W, \check{\mathcal{M}}_A^\beta)$$

of the \mathcal{D}_W -module $\check{\mathcal{M}}_A^\beta$. Sometimes we will be interested in the case $\beta = 0$ and will write

$$\check{\mathcal{M}}_A := \check{\mathcal{M}}_A^0 \quad \text{and} \quad M_A := \Gamma(W, \check{\mathcal{M}}_A).$$

Remark 3.6. Notice that $\check{\mathcal{M}}_A^\beta$ is just a Fourier-Laplace transformation (in all variables) of the GKZ-system \mathcal{M}_A^β (cf. Definition 3.1).

The semigroup ring associated with the matrix A is

$$\mathbb{C}[\mathbf{NA}] \simeq \mathbb{C}[\underline{w}] / ((\check{\square}_{\underline{m}})_{\underline{m} \in \mathbb{L}_A}),$$

where $\mathbb{C}[\underline{w}]$ is the commutative ring $\mathbb{C}[w_1, \dots, w_n]$ and the isomorphism follows from [MS05, Theorem 7.3]. The rings $\mathbb{C}[\underline{w}]$ and $\mathbb{C}[\mathbf{NA}]$ are naturally \mathbb{Z}^d -graded if we define $\deg(w_j) = \underline{a}_j$ for $j = 1, \dots, n$. This is compatible with the \mathbb{Z}^d -grading of the Weyl algebra D_W given by $\deg(\partial_{w_j}) = -\underline{a}_j$ and $\deg(w_j) = \underline{a}_j$.

Definition 3.7 ([MMW05, Definition 5.2]). Let N be a finitely generated \mathbb{Z}^d -graded $\mathbb{C}[\underline{w}]$ -module. An element $\alpha \in \mathbb{Z}^d$ is called a true degree of N if N_α is non-zero. A vector $\alpha \in \mathbb{C}^d$ is called a quasi-degree of N , written $\alpha \in \text{qdeg}(N)$, if α lies in the complex Zariski closure $\text{qdeg}(N)$ of the true degrees of N via the natural embedding $\mathbb{Z}^d \hookrightarrow \mathbb{C}^d$.

Schulze and Walther now define the following set of parameters:

Definition 3.8 ([SW09]). *The set*

$$sRes(A) := \bigcup_{j=1}^n sRes_j(A),$$

where

$$sRes_j(A) := \{\beta \in \mathbb{C}^d \mid \beta \in -(\mathbb{N} + 1)\underline{a}_j + qdeg(\mathbb{C}[\mathbb{N}A]/(w_j))\}$$

is called the set of strongly resonant parameters of A .

The matrix A is called pointed if 0 is the only unit in $\mathbb{N}A$. The matrix A gives rise to a map from a torus $T = (\mathbb{C}^*)^d$ with coordinates (t_1, \dots, t_d) into the affine space $W = \mathbb{C}^n$ with coordinates w_1, \dots, w_n :

$$\begin{aligned} h_A : T &\longrightarrow W \\ (t_1, \dots, t_d) &\mapsto (\underline{t}^{\underline{a}_1}, \dots, \underline{t}^{\underline{a}_n}), \end{aligned}$$

where $\underline{t}^{\underline{a}_i} := \prod_{k=1}^d t_k^{a_{ki}}$. Notice that the map h_A is affine and a locally closed embedding, hence the direct image functor for \mathcal{D}_T -modules $(h_A)_+$ is exact.

For a pointed matrix A Schulze and Walther computed the direct image of the twisted structure sheaf

$$\mathcal{O}_T^\beta := \mathcal{D}_T / \mathcal{D}_T \cdot (\partial_{t_1} t_1 + \beta_1, \dots, \partial_{t_d} t_d + \beta_d)$$

under the morphism h_A .

Theorem 3.9 ([SW09] Theorem 3.6, Corollary 3.7). *Let A a pointed $(d \times n)$ integer matrix satisfying $\mathbb{Z}A = \mathbb{Z}^d$, then the following statements are equivalent*

1. $\beta \notin sRes(A)$.
2. $\tilde{\mathcal{M}}_A^\beta \simeq (h_A)_+ \mathcal{O}_T^\beta$.
3. Left multiplication with w_i is invertible on \tilde{M}_A^β for $i = 1, \dots, n$.

Notice that Schulze and Walther [SW09] use the GKZ-system \mathcal{M}_A^β and the convention $deg(\partial_{\lambda_j}) = \underline{a}_j$. We will use $\tilde{\mathcal{M}}_A^\beta$ and $deg(w_j) = \underline{a}_j$ instead.

The aim of section is to generalize the implication 1. \Rightarrow 2. to the case of a non-pointed matrix A . For this we set $\underline{a}_0 := 0$. We will associate to the matrix A the homogenized $(d + 1 \times n + 1)$ matrix \tilde{A} with columns $\tilde{\underline{a}}_i := (1, \underline{a}_i)$ for $i = 0, \dots, n$. Notice that $\mathbb{Z}\tilde{A} = \mathbb{Z}^{d+1}$ holds and that the matrix \tilde{A} is pointed in any case. Consider now the augmented map

$$\begin{aligned} h_{\tilde{A}} : \tilde{T} &\longrightarrow \tilde{W} \\ (t_0, \dots, t_d) &\mapsto (t_0 \underline{t}^{\underline{a}_0}, t_0 \underline{t}^{\underline{a}_1}, \dots, t_0 \underline{t}^{\underline{a}_n}), \end{aligned} \tag{9}$$

where $\tilde{T} = (\mathbb{C}^*)^{d+1}$ and $\tilde{W} = \mathbb{C}^{n+1}$ with coordinates w_0, \dots, w_n . Let \tilde{W}_0 be the subvariety of \tilde{W} given by $w_0 \neq 0$ and denote by $k_0 : \tilde{W}_0 \rightarrow \tilde{W}$ the canonical embedding. The map $h_{\tilde{A}}$ factors through \tilde{W}_0 which gives rise to a map h_0 with $h_{\tilde{A}} = k_0 \circ h_0$. We get the following commutative diagram

$$\begin{array}{ccccc} & & h_{\tilde{A}} & & \\ & & \curvearrowright & & \\ \tilde{T} & \xrightarrow{h_0} & \tilde{W}_0 & \xrightarrow{k_0} & \tilde{W} \\ \downarrow \pi & & \downarrow \pi_0 & & \\ T & \xrightarrow{h_A} & W & & \end{array} \tag{10}$$

where π is the projection which forgets the first coordinate and π_0 is given by

$$\begin{aligned} \pi_0 : \tilde{W}_0 &\longrightarrow W \\ (w_0, w_1, \dots, w_n) &\mapsto (w_1/w_0, \dots, w_n/w_0). \end{aligned}$$

Lemma 3.10. *For each $\beta_0 \in \mathbb{Z}$ we have an isomorphism:*

$$\mathcal{H}^0 \left(h_{A+} \mathcal{O}_T^\beta \right) \simeq \mathcal{H}^0 \left(\pi_{0+} k_0^+ \left(h_{\tilde{A}+} \mathcal{O}_{\tilde{T}}^{(\beta_0, \beta)} \right) \right).$$

Proof. We show the claim by using the following isomorphisms

$$\begin{aligned} \mathcal{H}^0 h_{A+} \mathcal{O}_T^\beta &\simeq \mathcal{H}^0 h_{A+} \mathcal{H}^0 \pi_+ \mathcal{O}_{\tilde{T}}^{(\beta_0, \beta)} \simeq \mathcal{H}^0 h_{A+} \pi_+ \mathcal{O}_{\tilde{T}}^{(\beta_0, \beta)} \simeq \mathcal{H}^0 \pi_{0+} h_{0+} \mathcal{O}_{\tilde{T}}^{(\beta_0, \beta)} \\ &\simeq \mathcal{H}^0 \pi_{0+} k_0^+ k_{0+} h_{0+} \mathcal{O}_{\tilde{T}}^{(\beta_0, \beta)} \simeq \mathcal{H}^0 \pi_{0+} k_0^+ h_{\tilde{A}+} \mathcal{O}_{\tilde{T}}^{(\beta_0, \beta)}. \end{aligned}$$

The first isomorphism follows from the fact that π is a projection with fiber \mathbb{C}^* , the second isomorphism follows from the exactness of $(h_A)_+$ and the fourth from the fact that $k_0^+(k_0)_+ \simeq id_{\tilde{W}_0}$. \square

The following proposition is the generalization of Theorem 3.9 to the non-pointed case.

Proposition 3.11. *Let $A = (a_{ki})$ be a $d \times n$ integer matrix satisfying $\mathbb{Z}A = \mathbb{Z}^d$ and let $\beta \in \mathbb{C}^d$ with $\beta \notin sRes(A)$, then $\mathcal{H}^0 \left((h_A)_+ \mathcal{O}_T^\beta \right)$ is isomorphic to $\check{\mathcal{M}}_A^\beta$.*

Proof. The proof relies on Lemma 3.10 and the theorem of Schulze and Walther in the pointed case. Notice that we can find a $\beta_0 \in \mathbb{Z}$ with $\beta_0 \gg 0$ such that $(\beta_0, \beta) \notin sRes(\tilde{A})$ by [Rei14, Lemma 1.16] (in loc. cit. the statement is formulated for $\beta \in \mathbb{Q}^d$ but the proof carries over almost word for word in this more general case).

Consider the following isomorphism on \tilde{W}_0 :

$$\begin{aligned} f : \tilde{W}_0 &\longrightarrow \mathbb{C}_{w_0}^* \times W \\ (w_0, \dots, w_n) &\mapsto ((w_0, w_1/w_0, \dots, w_n/w_0) \end{aligned}$$

together with the canonical projection $p : W \times \mathbb{C}_{w_0}^* \rightarrow W$ which forgets the first coordinate. This factors $\pi_0 = p \circ f$, which gives (using Lemma 3.10 above)

$$\begin{aligned} \mathcal{H}^0 \left((h_A)_+ \mathcal{O}_T^\beta \right) &\simeq \mathcal{H}^0 \left((\pi_0)_+ \left((h_{\tilde{A}+} \mathcal{O}_{\tilde{T}}^{(\beta_0, \beta)}) \right)_{|\tilde{W}_0} \right) \simeq \mathcal{H}^0 \left(p_+ f_+ \left((h_{\tilde{A}+} \mathcal{O}_{\tilde{T}}^{(\beta_0, \beta)}) \right)_{|\tilde{W}_0} \right) \\ &\simeq \mathcal{H}^0 \left(p_+ f_+ (\check{\mathcal{M}}_{\tilde{A}}^{(\beta_0, \beta)})_{|\tilde{W}_0} \right). \end{aligned}$$

The \mathcal{D} -module $\mathcal{H}^0 f_+ (\check{\mathcal{M}}_{\tilde{A}}^{(\beta_0, \beta)})_{|\tilde{W}_0}$ is isomorphic to $\mathcal{D}_{W \times \mathbb{C}_{w_0}^*} / \mathcal{I}'_0$ where \mathcal{I}'_0 is generated by

$$\check{\square}_{\underline{m} \in \mathbb{L}_A} = \prod_{i: m_i > 0} w_i^{m_i} - \prod_{i: m_i < 0} w_i^{-m_i}$$

and

$$Z_0 = \partial_{w_0} w_0 + \beta_0 \quad \text{and} \quad E_k = \sum_{i=1}^n a_{ki} \partial_{w_i} w_i + \beta_k$$

Hence $\mathcal{H}^0 f_+ (\check{\mathcal{M}}_{\tilde{A}}^{(\beta_0, \beta)})_{|\tilde{W}_0}$ is isomorphic to $\check{\mathcal{M}}_A^\beta \boxtimes \mathcal{D}_{\mathbb{C}_{w_0}^*} / (\partial_{w_0} w_0 + \beta_0)$ as a \mathcal{D} -module. We therefore have

$$\mathcal{H}^0 \left(p_+ f_+ (\check{\mathcal{M}}_{\tilde{A}}^{(\beta_0, \beta)})_{|\tilde{W}_0} \right) \simeq \mathcal{H}^0 \left(p_+ \mathcal{H}^0 f_+ (\check{\mathcal{M}}_{\tilde{A}}^{(\beta_0, \beta)})_{|\tilde{W}_0} \right) \simeq \mathcal{H}^0 p_+ \left(\check{\mathcal{M}}_A^\beta \boxtimes \mathcal{D}_{\mathbb{C}_{w_0}^*} / (\partial_{w_0} w_0 + \beta_0) \right) \simeq \check{\mathcal{M}}_A^\beta.$$

\square

4 Hodge filtration on torus embeddings

The aim of this section is to compute explicitly the Hodge filtration of $(h_A)_+ \mathcal{O}_T^\beta$ as a mixed Hodge module for certain values of β (cf. Theorem 4.17). We will use this result in section 5 where the behavior of mixed Hodge modules obtained by such torus embeddings under the twisted Radon transformation is studied.

4.1 V-filtration

As above let A be a $d \times n$ integer matrix s.t. $\mathbb{Z}A = \mathbb{Z}^d$. In this section we additionally assume that the matrix A satisfies the following conditions:

$$\mathbb{N}A = \mathbb{Z}^d \cap \mathbb{R}_{\geq 0}A \quad \text{and} \quad \mathbb{N}A \neq \mathbb{Z}^d \quad (11)$$

where $\mathbb{R}_{\geq 0}A$ is the cone generated by the columns of A . As already noticed above, the first condition is equivalent to the fact that the semigroup ring $\mathbb{C}[\mathbb{N}A]$ is normal (see, e.g., [BH93, Section 6.1]). We will again consider the locally closed embedding

$$\begin{aligned} h_A : T &\longrightarrow W \\ (t_1, \dots, t_d) &\mapsto (\underline{t}^{a_1}, \dots, \underline{t}^{a_n}). \end{aligned}$$

Put $D := \{w_1 \cdot \dots \cdot w_n = 0\} \subset W$, $W^* := W \setminus D$, and consider the decomposition $h_A = l_A \circ k_A$, where

$$\begin{aligned} k_A : T &\longrightarrow W^* \\ (t_1, \dots, t_d) &\mapsto (\underline{t}^{a_1}, \dots, \underline{t}^{a_n}). \end{aligned}$$

and where $l_A : W^* \rightarrow W$ is the canonical open embedding.

Lemma 4.1. *The morphism $k_A : T \rightarrow W^*$ is a closed embedding.*

Proof. This is clear, as the image of k_A is precisely the vanishing locus of $(\check{\square}_m)_{m \in \mathbb{L}_A} \subset \Gamma(W^*, \mathcal{O}_{W^*})$. \square

The aim of this subsection is to compute parts of the canonical (descending) V -filtration of $\check{\mathcal{M}}_A^\beta \simeq h_{A+} \mathcal{O}_T^\beta$ (or Kashiwara-Malgrange filtration) along the normal crossing divisor D for certain values of β .

We review very briefly some facts about the V -filtration for differential modules. Let $X = \text{Spec}(R)$ be a smooth affine variety and $Y = \text{div}(t)$ be a smooth reduced principal divisor. Denote by $I = (t)$ the corresponding ideal. Let as before $D_X = \Gamma(X, \mathcal{D}_X)$ be the ring of algebraic differential operators on X , then the V -filtration on D_X is defined by

$$V^k D_X = \{P \in D_X \mid P I^j \subset I^{j+k} \text{ for any } j \in \mathbb{Z}\},$$

where $I^j = R$ for $j \leq 0$. One has

$$\begin{aligned} V^k D_X &= t^k V^0 D_X, \\ V^{-k} D_X &= \sum_{0 \leq j \leq k} \partial_t^j V^0 D_X. \end{aligned}$$

Choose a total ordering $<$ on \mathbb{C} such that, for any $\alpha, \beta \in \mathbb{C}$, the following conditions hold:

1. $\alpha < \alpha + 1$,
2. $\alpha < \beta$ if and only if $\alpha + 1 < \beta + 1$,
3. $\alpha < \beta + m$ for some $m \in \mathbb{Z}$.

We recall the definition of the canonical V -filtration (see, e.g., [Sai93, Section 1]).

Definition 4.2. *Let N be a coherent D_X -module. The canonical V -filtration (or Kashiwara-Malgrange filtration) is an exhaustive filtration on N indexed discretely by \mathbb{C} with total order as above and is uniquely determined by the following conditions*

1. $(V^k D_X)(V^\alpha N) \subset V^{\alpha+k} N$ for all k, α
2. $V^\alpha N$ is coherent over $V^0 D_X$ for any α
3. $t(V^\alpha N) = V^{\alpha+1} N$ for $\alpha \gg 0$
4. the action of $\partial_t t - \alpha$ on $\text{Gr}_V^\alpha N = V^\alpha N / V^{>\alpha} N$ is nilpotent

where $V^{>\alpha}N := \bigcup_{\beta>\alpha} V^\beta$.

The canonical V -filtration is unique if it exists. Its existence is guaranteed if N is D_X -holonomic.

We reduce the computation of the V -filtration on \check{M}_A^β along the possibly singular divisor D to the computation of a V -filtration along a smooth divisor by considering the following graph embedding:

$$\begin{aligned} i_g : W &\longrightarrow W \times \mathbb{C}_t \\ (w_1, \dots, w_n) &\mapsto (w_1, \dots, w_n, w_1 \cdots w_n). \end{aligned}$$

Instead of computing the V -filtration on \check{M}_A^β , we will compute it on $\Gamma(W \times \mathbb{C}_t, \mathcal{H}^0(i_{g+}\check{\mathcal{M}}_A^\beta))$ along $t = 0$ (notice that i_g is an affine embedding hence i_{g+} is exact). In order to compute the direct image we consider the composed map

$$\begin{aligned} i_g \circ h_A : T &\longrightarrow W \times \mathbb{C}_t \\ (t_1, \dots, t_d) &\mapsto (\underline{t}^{\underline{a}_1}, \dots, \underline{t}^{\underline{a}_n}, \underline{t}^{\underline{a}_1 + \dots + \underline{a}_n}). \end{aligned} \quad (12)$$

Notice that the matrix A' , which is built from the columns $\underline{a}_1, \dots, \underline{a}_n, \underline{a}_1 + \dots + \underline{a}_n$, gives a saturated semigroup $\mathbb{N}A' = \mathbb{N}A$. Hence we can apply again Proposition 3.11 to compute

$$\check{\mathcal{M}}_{A'}^\beta \simeq \mathcal{H}^0 i_{g+} \check{\mathcal{M}}_A^\beta \simeq \mathcal{H}^0(i_g \circ h_A)_+ \mathcal{O}_T^\beta.$$

This means that $\mathcal{H}^0 i_{g+} \check{\mathcal{M}}_A^\beta$ is a cyclic $\mathcal{D}_{W \times \mathbb{C}_t}$ -module $\mathcal{D}_{W \times \mathbb{C}_t} / \mathcal{I}'$, where \mathcal{I}' is generated by

$$\check{E}'_k := \sum_{i=1}^n a_{ki} \partial_{w_i} w_i + c_k \partial_t t + \beta_k \quad \text{for } k = 1, \dots, d, \quad (13)$$

where $c_k = a_{k1} + \dots + a_{kn}$ is the k -th component of $c \in \mathbb{Z}^d$ and

$$\check{\square}_{\underline{m} \in \mathbb{L}_{A'}} := \begin{cases} \prod_{m_i > 0} w_i^{m_i} t^{m_{n+1}} - \prod_{m_i < 0} w_i^{-m_i} & \text{for } m_{n+1} \geq 0 \\ \prod_{m_i > 0} w_i^{m_i} - \prod_{m_i < 0} w_i^{-m_i} t^{-m_{n+1}} & \text{for } m_{n+1} < 0 \end{cases} \quad (14)$$

where $\mathbb{L}_{A'}$ is the \mathbb{Z} -module of relations among the columns of A' .

We are going to use the following characterization of the canonical V -filtration along $t = 0$.

Proposition 4.3. [MM04, Definition 4.3-3, Proposition 4.3-9] Let $n \in N$ and set $E := \partial_t$. The Bernstein-Sato polynomial of n is the unitary polynomial of smallest degree, satisfying

$$b(E)n \in V^1(D_X)n.$$

We denote it by $b_n(x) \in \mathbb{C}[x]$ and denote the set of roots of $b_n(x)$ by $\text{ord}(n)$. The canonical V -filtration on N is then given by

$$V^\alpha N = \{n \in N \mid \text{ord}(n) \subset [\alpha, \infty)\}.$$

We will use this characterization to compute the canonical V -filtration on \check{M}_A^β along $t = 0$ for certain $\beta \in \mathbb{R}^d$.

Let $\underline{c} := \underline{a}_1 + \dots + \underline{a}_n$. For all facets F of $\mathbb{R}_{\geq 0}A' = \mathbb{R}_{\geq 0}A$ let $0 \neq \underline{n}_F \in \mathbb{Z}^d$ be the uniquely determined primitive, inward-pointing, normal vector of F , i.e. \underline{n}_F satisfies $\langle \underline{n}_F, F \rangle = 0$, $\langle \underline{n}_F, \mathbb{N}A \rangle \subset \mathbb{Z}_{\geq 0}$ and $\lambda \cdot \underline{n}_F \notin \mathbb{Z}^d$ for $\lambda \in [0, 1)$ (where $\langle \cdot, \cdot \rangle$ is the Euclidean pairing). Set

$$e_F := \langle \underline{n}_F, \underline{c} \rangle \in \mathbb{Z}_{\geq 0}.$$

We show that e_F is always positive. We have $\underline{c} \neq 0$ since otherwise $0 = -\underline{a}_1 - \dots - \underline{a}_n \in \mathbb{N}A$ and therefore $-\underline{a}_i \in \mathbb{N}A$ for all $i \in \{1, \dots, n\}$ which contradicts the assumption $\mathbb{N}A \neq \mathbb{Z}^d$. Furthermore \underline{c} lies in the interior of $\mathbb{R}_{\geq 0}A'$. In order to see this assume to the contrary that \underline{c} lies on some facet F of $\mathbb{R}_{\geq 0}A'$. Then $\langle \underline{n}_F, \underline{c} \rangle = 0$ holds. For $\underline{a}_i \notin F$ we have on the one hand $\underline{c} - \underline{a}_i \in \mathbb{N}A$ and on the other hand $\langle \underline{c} - \underline{a}_i, \underline{n}_F \rangle < 0$ which is a contradiction. Hence \underline{c} is in the interior of $\mathbb{R}_{\geq 0}A$, which shows $e_F \in \mathbb{Z}_{> 0}$.

We define the following set of admissible parameters β :

$$\mathfrak{A}_A := \bigcap_{F: F \text{ facet}} \left\{ \mathbb{R} \cdot F - \left[0, \frac{1}{e_F}\right] \cdot \underline{c} \right\} \quad (15)$$

Lemma 4.4. *Suppose as above that $\mathbb{N}A = \mathbb{Z}^d \cap \mathbb{R}_{\geq 0}A$. Consider the cyclic $D_{W \times \mathbb{C}_t}$ -module $\check{M}_{A'}^\beta$, and its generator $[1] \in \check{M}_{A'}$. Then we have $\text{ord}([1]) \subset [0, 1)$ if $\beta \in \mathfrak{A}_A$.*

Proof. It was shown in [RSW18, Theorem 3.5] that the roots of $b_{[1]}(x)$ for $[1] \in \check{M}_{A'}^\beta$ are contained in the set $\{\epsilon \in \mathbb{C} \mid \epsilon \cdot \underline{c} \in \text{qdeg}(\mathbb{C}[\mathbb{N}A']/(t)) - \beta\}$ which is discrete since \underline{c} lies in the interior of $\mathbb{R}_{\geq 0}A' = \mathbb{R}_{\geq 0}A$ and $\text{qdeg}(\mathbb{C}[\mathbb{N}A']/(t))$ is a finite union of parallel translates of the complex span of faces of $\mathbb{R}_{\geq 0}A'$ (cf. [MMW05]). We will now compute an estimate of the quasi-degrees $\text{qdeg}(\mathbb{C}[\mathbb{N}A']/(t))$. For this we remark that $0 = [P] \in \mathbb{C}[\mathbb{N}A']/(t)$ for $P \in \mathbb{C}[\mathbb{N}A']$ iff $\exists P' \in \mathbb{C}[\mathbb{N}A']$ with $P = P' \cdot t$. In this case we have $\text{deg}(P) \in \mathbb{N}A + \underline{c}$.

Set $L_F := \{\frac{k}{e_F} \cdot \underline{c} + \mathbb{C} \cdot F \mid k = 0, \dots, e_F - 1\}$. Then $L = \bigcup_{F:\text{facet}} L_F$ is Zariski closed and we will show that the set $\text{deg}(\mathbb{C}[\mathbb{N}A']/(t))$ is contained in L . Let $P \in \mathbb{C}[\mathbb{N}A']$ with $0 \neq [P] \in \mathbb{C}[\mathbb{N}A']/(t)$ and set $\underline{p} := \text{deg}(P) \in \mathbb{N}A$. Since $-\underline{c} \notin \mathbb{R}_{\geq 0}A$ there exist a facet F and some $\lambda \in [0, 1)$ such that $\underline{p} - \lambda \underline{c} \in F$, i.e. $\underline{p} = \lambda \underline{c} + \underline{f}$ for some $\underline{f} \in F$. We have $\lambda \cdot e_F = \langle \lambda \underline{c} + \underline{f}, \underline{n}_F \rangle = \langle \underline{p}, \underline{n}_F \rangle \in \mathbb{Z}_{\geq 0}$. Hence $\underline{p} \in L_F \subset L$.

Since $\text{qdeg}(\mathbb{C}[\mathbb{N}A']/(t))$ is by definition the Zariski closure of $\text{deg}(\mathbb{C}[\mathbb{N}A']/(t))$ the former set is contained in L . In particular this shows that the roots of $b_{[1]}(x)$ are contained in the set $\{\epsilon \in \mathbb{C} \mid \epsilon \cdot \underline{c} \in L - \beta\}$. Since L is a union of hypersurfaces which are defined over \mathbb{R} , $\underline{c} \in \mathbb{Z}^d$ and $\beta \in \mathbb{R}^d$, this set is equal to $\{\epsilon \in \mathbb{R} \mid \epsilon \cdot \underline{c} \in L - \beta\}$. Hence for $\beta \in \bigcap_{F:\text{facet}} \{\mathbb{R} \cdot F - [0, \frac{1}{e_F}) \cdot \underline{c}\}$ we can guarantee that the roots of $b_{[1]}(x)$ are contained in $[0, 1)$. \square

We will prove a basic lemma on the set \mathfrak{A}_A which will be of importance later.

Lemma 4.5. *Suppose $\mathbb{N}A = \mathbb{Z}^d \cap \mathbb{R}_{\geq 0}$. Then $\mathfrak{A}_A \cap \text{sRes}(A) = \emptyset$.*

Proof. Recall that $\text{sRes}(A) = \bigcup_{j=1}^n \text{sRes}_j(A) = \bigcup_{j=1}^n -(\mathbb{N} + 1)\underline{a}_j + \text{qdeg}(\mathbb{C}[\mathbb{N}A]/((w_j)))$. Therefore it is enough to show that

$$\mathfrak{A}_A \cap \left\{ -(\mathbb{N} + 1)\underline{a}_j + \text{qdeg}(\mathbb{C}[\mathbb{N}A]/((w_j))) \right\} = \emptyset \quad (16)$$

holds. The following estimate of the quasi-degrees of $\mathbb{C}[\mathbb{N}A]/((w_j))$ can be shown similarly as in the proof of the lemma above

$$\text{qdeg}(\mathbb{C}[\mathbb{N}A]/((w_j))) \subset L_j := \bigcup_{F:\underline{a}_j \notin F} \left\{ \frac{k}{e_{F,j}} \cdot \underline{a}_j + \mathbb{C} \cdot F \mid k = 0, \dots, e_{F,j} - 1 \right\}$$

where $e_{F,j} := \langle \underline{n}_F, \underline{a}_j \rangle$. Hence it is enough to show that for each $j \in \{1, \dots, n\}$ and each facet F with $\underline{a}_j \notin F$ the following holds

$$\left\{ \mathbb{R} \cdot F - [0, \frac{1}{e_F}) \cdot \underline{c} \right\} \cap \left\{ -(\mathbb{N} + 1)\underline{a}_j + \bigcup_{k=0}^{e_{F,j}-1} \frac{k}{e_{F,j}} \cdot \underline{a}_j + \mathbb{R} \cdot F \right\} = \emptyset \quad (17)$$

Since F has codimension one in \mathbb{R}^d and $\underline{a}_j, \underline{c} \notin \mathbb{R} \cdot F$ we can write $\underline{c} = \lambda \underline{a}_j + f$ for some $f \in \mathbb{R} \cdot F$. We get $e_F = \lambda e_{F,j}$. We conclude that (17) is equivalent to


$$\left\{ \mathbb{R} \cdot F - [0, \frac{1}{e_{F,j}}) \cdot \underline{a}_j \right\} \cap \left\{ -(\mathbb{N} + 1)\underline{a}_j + \bigcup_{k=0}^{e_{F,j}-1} \frac{k}{e_{F,j}} \cdot \underline{a}_j + \mathbb{R} \cdot F \right\} = \emptyset.$$

But this holds since $(-\frac{1}{e_{F,j}}, 0] \cap \{-(\mathbb{N} + 1) + \{0, \frac{1}{e_{F,j}}, \dots, \frac{e_{F,j}-1}{e_{F,j}}\}\} = \emptyset$. \square

Example 4.6. *The sets $\text{sRes}(A)$ and \mathfrak{A}_A for the matrix*

$$A = \begin{pmatrix} -1 & 0 & 1 & 2 \\ 1 & 1 & 1 & 1 \end{pmatrix}$$

are sketched below.



We give another estimate of the set \mathfrak{A}_A :

Lemma 4.7. *Suppose $\mathbb{N}A = \mathbb{Z}^d \cap \mathbb{R}_{\geq 0}A$. Let $\underline{b} \in \mathbb{Z}^d \cap \text{int}(\mathbb{R}_{\geq 0}A)$, then $\mathfrak{A}_A \subset -\underline{b} + \text{int}(\mathbb{R}_{\geq 0}A)$.*

Proof. Let F be a face of $\mathbb{R}_{\geq 0}A$. Since F has codimension and $\underline{b}, \underline{c} \notin \mathbb{R}F$ we can write $\underline{c} = \lambda \underline{b} + f$ for some $f \in \mathbb{R} \cdot F$. If we set $e_{F, \underline{b}} := \langle \underline{n}_F, \underline{b} \rangle \in \mathbb{Z}_{>0}$ we get $e_F = \lambda e_{F, \underline{b}}$. hence

$$\{\mathbb{R} \cdot F - [0, \frac{1}{e_F}] \cdot \underline{c}\} = \{\mathbb{R} \cdot F - [0, \frac{1}{e_{F, \underline{b}}}] \cdot \underline{b}\} \subset \{\mathbb{R} \cdot F - [0, 1) \underline{b}\} \subset \{\mathbb{R} \cdot F - \underline{b} + (0, \infty) \underline{b}\} \subset -\underline{b} + \{\mathbb{R} \cdot F + (0, \infty) \underline{n}_F\}$$

hence

$$\mathfrak{A} = \bigcap_F \{\mathbb{R} \cdot F - [0, \frac{1}{e_F}] \cdot \underline{c}\} \subset -\underline{b} + \text{int}(\mathbb{R}_{\geq 0}A)$$

□

Next we draw a consequence for the canonical V -filtration with respect to $t = 0$ on $\mathcal{H}^0 i_{g+} \check{\mathcal{M}}_A^\beta$. We will not compute all of its filtration steps, but those corresponding to integer indices, which is sufficient for our purpose. For this consider the induced V -filtration on $\check{M}_{A'}^\beta = \Gamma(W \times \mathbb{C}_t, \mathcal{H}^0 i_{g+} \check{\mathcal{M}}_A^\beta)$

$$V_{ind}^k \check{M}_{A'}^\beta := \{[P] \in \check{M}_{A'}^\beta \mid P \in V^k D_{W \times \mathbb{C}_t}\}.$$

It is readily checked that $V_{ind}^k \check{M}_{A'}^\beta$ is a good V -filtration on $\check{M}_{A'}^\beta$. As $\check{M}_{A'}$ is holonomic, hence specializable along any smooth hypersurface, it admits a Bernstein polynomial $b_{V_{ind}}(x)$ in the sense of [MM04, Définition 4.2-3]. On the other hand, for any section $\sigma : \mathbb{C}/\mathbb{Z} \rightarrow \mathbb{C}$ of the canonical projection $\mathbb{C} \rightarrow \mathbb{C}/\mathbb{Z}$, there is a *unique* good (\mathbb{Z} -indexed) V -filtration $V_{\sigma}^\bullet \check{M}_{A'}^\beta$ on $\check{M}_{A'}^\beta$, such that the roots of $b_{V_{\sigma}}(x)$ lie in $\text{Im}(\sigma)$ (see loc.cit., Proposition 4.2-6). From this we deduce the following result, which describes the integral part of the canonical V -filtration on $\check{M}_{A'}^\beta$.

Proposition 4.8. *If $\mathbb{N}A = \mathbb{Z}^d \cap \mathbb{R}_{\geq 0}A$ and $\beta \in \mathfrak{A}_A$, then for any $k \in \mathbb{Z}$, we have the following equality*

$$V^k \check{M}_{A'}^\beta = V_{ind}^k \check{M}_{A'}^\beta.$$

Proof. Recall (see [MM04, Proposition 4.3-5]) that we have $V^{\alpha+k} \check{M}_{A'}^\beta = V_{\sigma_\alpha}^k \check{M}_{A'}^\beta$ for any $\alpha \in \mathbb{C}, k \in \mathbb{Z}$, where $\sigma_\alpha : \mathbb{C}/\mathbb{Z} \rightarrow \mathbb{C}$ is the section of $\mathbb{C} \rightarrow \mathbb{C}/\mathbb{Z}$ with image equal to $[\alpha, \alpha + 1)$. Hence, in order to prove the proposition it is enough to show that $V_{\sigma_0}^k \check{M}_{A'}^\beta = V_{ind}^k \check{M}_{A'}^\beta$. Using loc. cit., Proposition 4.2-6 it remains to show that the roots of the Bernstein polynomial $b_{V_{ind}}(x)$ are contained in $[0, 1)$.

An element $[P]$ of $V_{ind}^k \check{M}_{A'}^\beta$ for $k \geq 0$ can be written as

$$[P] = \left[\sum_{i=0}^l t^k (\partial_t t)^i P_i \right] + [R],$$

where $[R] \in V_{ind}^{k+1} \check{M}_{A'}^\beta$ and $P_i \in \mathbb{C}[w_1, \dots, w_n] \langle \partial_{w_1}, \dots, \partial_{w_n} \rangle$. We have

$$\begin{aligned} b_{[1]}(\partial_t t - k) \cdot [P] &= \left[\sum_{i=0}^l t^k (\partial_t t)^i P_i \cdot b_{[1]}(\partial_t t) \right] + b_{[1]}(\partial_t t - k) \cdot [R] \\ &= \sum_{i=0}^l t^k (\partial_t t)^i P_i \cdot b_{[1]}(\partial_t t) \cdot [1] + b_{[1]}(\partial_t t - k) \cdot [R]. \end{aligned}$$

But $\sum_{i=0}^l t^k (\partial_t t)^i P_i \cdot b_{[1]}(\partial_t t) \cdot [1] \in V_{ind}^{k+1} \check{M}_{A'}^\beta$ because $\sum_{i=0}^l t^k (\partial_t t)^i P_i \in V^k D$ and $b_{[1]}(\partial_t t) \cdot [1] \in V_{ind}^1 \check{M}_{A'}^\beta$. Therefore

$$b_{[1]}(\partial_t t - k) \cdot [P] \in V_{ind}^{k+1} \check{M}_{A'}^\beta.$$

Now let $[P] \in V_{ind}^{-k} \check{M}_{A'}^\beta$ with $k > 0$. It can be written as

$$[P] = \left[\sum_{i=0}^l \partial_t^k (\partial_t t)^i P_i \right] + [R],$$

where $[R] \in V_{ind}^{k+1} \check{M}_{A'}^\beta$. By a similar argument we have

$$b_{[1]}(\partial_t t + k) \cdot [P] \in V_{ind}^{-k+1} \check{M}_{A'}^\beta.$$

This shows $b_{V_{ind}}(x) \mid b_{[1]}(x)$. Because of Lemma 4.4 the roots of $b_{V_{ind}}(x)$ are contained in $[0, 1)$, the claim follows. \square

4.2 Compatibility of filtrations

In this subsection we are going to show a compatibility result between two filtrations on the $D_{W'}$ -module $\check{M}_{A'}^\beta$ (recall that the $d \times (n+1)$ -matrix A' has columns $\underline{a}_1, \dots, \underline{a}_n, \underline{a}_1 + \dots + \underline{a}_n$). Let, as before, F^{ord} be the filtration induced on $M_{A'}^\beta$ by the filtration $F \bullet D_{W'}$ by orders of differential operators. Moreover, let $V \bullet D_{W'}$ be the V -filtration on $D_{W'}$ with respect to the coordinate w_{n+1} , and denote as before by $V_{ind} \bullet M_{A'}^\beta$ the induced filtration on $M_{A'}^\beta$. Then the main result of this subsection can be stated as follows.

Proposition 4.9. *Let A be a $d \times n$ -integer matrix and suppose that $\mathbb{N}A = \mathbb{Z}^d \cap \mathbb{R}_{\geq 0}A$, $\mathbb{N}A \neq \mathbb{Z}^d$. Let $A' := (\underline{a}_1, \dots, \underline{a}_n, \underline{a}_1 + \dots + \underline{a}_n)$ and consider the left $D_{W'}$ -module*

$$\check{M}_{A'}^\beta = D_{W'} / \left((\check{\square}_{\underline{m}})_{\underline{m} \in \mathbb{L}_{A'}} + (\check{E}'_k + \beta_k)_{k=1, \dots, d} \right).$$

Then the map

$$V^k D_{W'} \cap F_p D_{W'} \longrightarrow V_{ind}^k \check{M}_{A'}^\beta \cap F_p^{ord} \check{M}_{A'}^\beta.$$

is surjective.

The proof of this result will occupy this entire section. Before going into it, let us comment on how this result will enter in the calculation of the Hodge filtration on $\check{M}_{A'}^\beta$. As will be explained in more detail at the beginning of section 4.3, we consider the mixed Hodge module $h_{A*}^p \mathbb{C}_T^{\beta, H}$ with underlying D_W -module $h_{A+} \mathcal{O}_T^\beta$. If A and β satisfy the assumptions of Proposition 3.11, then this D_W -module is $\check{M}_{A'}^\beta$. In order to compute the Hodge filtration on its module of global sections $\check{M}_{A'}^\beta$, we will first consider the module $\check{M}_{A'}^\beta = \Gamma(W', h_{A'+}^\beta \mathcal{O}_T^\beta)$ and compute the Hodge filtration on it. We will use the fact that the embedding $h_{A'} : T \hookrightarrow W'$ can be factored as

$$T \longrightarrow W \times \mathbb{C}_t^* \xrightarrow{j} W' = W \times \mathbb{C}_t,$$

where the first morphism is a *closed* embedding, and the second one is the canonical open embedding of $W \times \mathbb{C}_t^*$ into W' . Then the main tool to compute the Hodge filtration on $\check{M}_{A'}^\beta$ is the following formula of Saito (see formula (22) below). Let $(\mathcal{M}, F_\bullet^H)$ be any filtered $\mathcal{D}_{W \times \mathbb{C}_t^*}$ -module underlying a complex mixed Hodge module in $\text{MHM}(W \times \mathbb{C}_t^*, \mathbb{C})$. Then the direct image $j_+ \mathcal{M}$ underlies a complex mixed Hodge module on W' , and its Hodge filtration is given as

$$F_p^H j_+ \mathcal{M} = \sum_{i \geq 0} \partial_t^i (V^0 j_+ \mathcal{M} \cap j_* (F_q^H \mathcal{M})),$$

where $V \bullet j_+ \mathcal{M}$ denotes the canonical V -filtration on $j_+ \mathcal{M}$ with respect to the divisor $\{t = 0\}$. We are going to apply this formula for the case where \mathcal{M} is the direct image of \mathcal{O}_T^β under the map $T \rightarrow W \times \mathbb{C}_t^*$ (so that $j_+ \mathcal{M} = \check{M}_{A'}^\beta$). Since this map is a closed immersion, we can explicitly calculate the Hodge

filtration on this direct image, i.e., it is given as the shifted order filtration for a cyclic presentation. Moreover, if β satisfies the assumptions of Proposition 4.8, then we have $V^k \tilde{M}_{A'}^\beta = V_{ind}^k \tilde{M}_{A'}^\beta$ for all $k \in \mathbb{Z}$ (in particular, for $k = 0$), so that we have to compute the intersection of the order filtration on $M_{A'}^\beta$ with the *induced* V -filtration on that module. As we will see below in section 4.3, this is possible since these two filtrations satisfy the compatibility statement of the above proposition. Its proof relies on the very specific structure of the hypergeometric ideal $\tilde{I}_{A'} = ((\tilde{\square}_m)_{m \in \mathbb{L}_{A'}} + (\tilde{E}_k + \beta_k)_{k=1, \dots, d}) \subset D_{W'}$ and uses non-commutative Gröbner basis techniques, a good reference for results needed is [SST00]. We will recall the main definitions for the reader's convenience.

To simplify the notation, we rename the coordinate t on W' to be w_{n+1} , that is $\Gamma(W', \mathcal{O}_{W'}) = \mathbb{C}[w_1, \dots, w_n, w_{n+1}]$. We work in the Weyl algebra $D_{W'} = \mathbb{C}[w_1, \dots, w_{n+1}] \langle \partial_{w_1}, \dots, \partial_{w_{n+1}} \rangle$. Any operator $P \in D_{W'}$ has the so-called normally ordered expression $P = \sum_{(\gamma, \delta)} c_{\gamma\delta} w^\gamma \partial_w^\delta \in D_{W'}$, where the sum runs over all pairs (γ, δ) in some finite subset of $\mathbb{N}^{2(n+1)}$.

First we define partial orders on the set of monomials in $D_{W'}$ resp. $\mathbb{C}[w] := \mathbb{C}[w_1, \dots, w_{n+1}]$ resp. $\mathbb{C}[w, \xi] := \mathbb{C}[w_1, \dots, w_{n+1}, \xi_1, \dots, \xi_{n+1}]$ by choosing the weight vectors $(u, v) \in \mathbb{Z}^{2(n+1)}$ with $u_i + v_i \geq 0$ resp. $u \in \mathbb{Z}^{n+1}$. This means that the variables w_i have weight u_i and the partial differentials ∂_{w_i} resp. ξ_i have weight v_i . The associated partial order in $D_{W'}$ is defined as follows: If for two monomials $w^\gamma \partial_w^\delta, w^c \partial_w^d$ we have $\sum_i u_i c_i + v_i d_i < \sum_i u_i \gamma_i + v_i \delta_i$, then by definition $w^\gamma \partial_w^\delta$ is larger than $w^c \partial_w^d$, we write $w^c \partial_w^d \prec_{(u,v)} w^\gamma \partial_w^\delta$ and similarly for $\mathbb{C}[w]$ and $\mathbb{C}[w, \xi]$. The weight vector (u, v) induces an increasing resp. decreasing filtrations on $D_{W'}$ given by

$$F_p^{(u,v)} D_{W'} = \left\{ \sum_{\sum_i u_i \gamma_i + v_i \delta_i \leq p} c_{\gamma\delta} w^\gamma \partial_w^\delta \right\} \quad \text{resp.} \quad F_{(u,v)}^p D_{W'} = \left\{ \sum_{\sum_i u_i \gamma_i + v_i \delta_i \geq p} c_{\gamma\delta} w^\gamma \partial_w^\delta \right\}$$

We define the graded ring $\text{Gr}_\bullet^{(u,v)} D_{W'} := \bigoplus_p F_p^{(u,v)} D_{W'} / F_{p-1}^{(u,v)} D_{W'}$ associated with the weight (u, v) . Notice that for $(u, v) = (0, \dots, 0, 1, \dots, 1)$ (i.e. the w_i have weight 0 and the ∂_{w_i} have weight 1) the ascending filtration $F_\bullet^{(u,v)} D_{W'}$ is the order filtration $F_\bullet D_{W'}$ and for $(u, v) = (0, \dots, 0, -1, 0, \dots, 0, 1)$ the descending filtration $F_{(u,v)}^\bullet D_{W'}$ is the V -filtration with respect to w_{n+1} .

We get well-defined maps

$$\begin{aligned} in_{(u,v)} : D_{W'} &\longrightarrow \text{Gr}_\bullet^{(u,v)} D_{W'} = \mathbb{C}[w, \xi] \\ P = \sum_{\gamma, \delta} c_{\gamma\delta} w^\gamma \partial_w^\delta &\longmapsto in_{(u,v)}(P) := \sum_{\sum_i u_i \gamma_i + v_i \delta_i = m} c_{\gamma\delta} w^\gamma \xi^\delta \end{aligned}$$

where $m := \text{ord}_{(u,v)}(P) := \max\{\sum_i u_i \gamma_i + v_i \delta_i \mid c_{\gamma\delta} \neq 0\}$ and

$$\begin{aligned} in_u : \mathbb{C}[w] &\longrightarrow \text{Gr}_\bullet^u \mathbb{C}[w] = \mathbb{C}[w] \\ Q = \sum_\gamma c_\gamma w^\gamma &\longmapsto in_u(Q) := \sum_{\sum_i u_i \gamma_i = m} c_\gamma w^\gamma \end{aligned}$$

where $m = \max\{\sum_u u_i \gamma_i \mid c_\gamma \neq 0\}$ and

$$\begin{aligned} in_{(u,v)} : \mathbb{C}[w, \xi] &\longrightarrow \text{Gr}_\bullet^{(u,v)} \mathbb{C}[w, \xi] = \mathbb{C}[w, \xi] \\ R = \sum_{\gamma, \delta} c_{\gamma\delta} w^\gamma \xi^\delta &\longmapsto in_{(u,v)}(R) := \sum_{\sum_i u_i \gamma_i + v_i \delta_i = m} c_{\gamma\delta} w^\gamma \xi^\delta \end{aligned}$$

where $m := \text{ord}_{(u,v)}(R) := \max\{\sum_i u_i \gamma_i + v_i \delta_i \mid c_{\gamma\delta} \neq 0\}$. Notice that, in constrast to the case of a total ordering, the initial terms $\text{in}_{(u,v)}$ resp. in_u are not monomials.

Let $I' \subset D_{W'}$ be a left ideal. The set $\text{in}_{(u,v)}(I')$ is an ideal in $\text{Gr}_{\bullet}^{(u,v)} D_{W'}$, and is called initial ideal of I' with respect to the weight vector (u, v) . A finite subset G of $D_{W'}$ is a Gröbner basis of I' with respect to (u, v) if I' is generated by G and $\text{in}_{(u,v)}(I')$ is generated by $\text{in}_{(u,v)}(G)$. Similarly, let $J' \subset \mathbb{C}[w]$ resp. $K' \subset \mathbb{C}[w, \xi]$ be an ideal. The set $\text{in}_u(J')$ resp. $\text{in}_{(u,v)}(K')$ is an ideal in $\text{Gr}_{\bullet}^u \mathbb{C}[w]$ resp. $\text{Gr}_{\bullet}^{(u,v)} \mathbb{C}[w, \xi]$ and is called initial ideal of J' resp. K' with respect to the weight vector u resp. (u, v) . The definition of a Gröbner basis is parallel to the definition above.

Let $\check{I}_{A'} = \left((\check{\square}_m)_{m \in \mathbb{L}_{A'}} + (\check{E}'_k + \beta_k)_{k=1, \dots, d} \right)$ be the hypergeometric ideal. The fake initial ideal $\text{fin}_{(u,v)}(\check{I}_{A'})$ is the following ideal in $\text{Gr}_{(u,v)} D_{W'}$:

$$\text{fin}_{(u,v)}(\check{I}_{A'}) := \text{Gr}_{(u,v)} D_{W'} \cdot \text{in}_u(\check{J}_{A'}) + \sum_{k=1}^d \text{Gr}_{(u,v)} D_{W'} \cdot \text{in}_{(u,v)}(\check{E}'_k + \beta_k)$$

where $\check{J}_{A'} \subset \mathbb{C}[w]$ is the ideal generated by $(\check{\square}_m)_{m \in \mathbb{L}_{A'}}$.

Consider the Koszul complex

$$\dots \xrightarrow{d_{-2}} K^{-1}(\text{Gr}_{\bullet}^{(u,v)}(D_{W'}/D_{W'}\check{J}_{A'})) \xrightarrow{d_{-1}} K^0(\text{Gr}_{\bullet}^{(u,v)}(D_{W'}/D_{W'}\check{J}_{A'})) \longrightarrow 0$$

where

$$K^{-p}(\text{Gr}_{\bullet}^{(u,v)}(D_{W'}/D_{W'}\check{J}_{A'})) = \bigoplus_{1 \leq i_1 < \dots < i_p \leq s+1} \text{Gr}_{\bullet}^{(u,v)}(D_{W'}/D_{W'}\check{J}_{A'}) e_{i_1 \dots i_p}$$

and

$$d_{-p}(e_{i_1 \dots i_p}) = \sum_{k=1}^p (-1)^{k-1} \text{in}_{(u,v)}(\check{E}'_k + \beta_k) e_{i_1 \dots \widehat{i}_k \dots i_p}$$

The following statement is an easy adaption of [SST00, Theorem 4.3.5]

Proposition 4.10. *If the cohomology $H^{-1}(K^{\bullet}(\text{Gr}_{\bullet}^{(u,v)}(D_{W'}/D_{W'}\check{J}_{A'})))$ vanishes, then the initial ideal satisfies $\text{in}_{(u,v)}(\check{I}_{A'}) = \text{fin}_{(u,v)}(\check{I}_{A'})$.*

Proof. After a Fourier-Laplace transform $w_i \rightarrow \partial_{x_i}$ and $\partial_{w_i} \rightarrow -x_i$ the proof carries over word for word from loc. cit. (Notice that in Chapter 4 of loc. cit. the homogeneity of A' assumed, however the proof of this statement does not need this requirement). \square

Recall that A' is a matrix built from the matrix A by adding a column which is the sum over all columns of A . Let $\check{J}_A \subset \mathbb{C}[w_1, \dots, w_n]$ be the ideal generated by $(\check{\square}_l)_{l \in \mathbb{L}_A}$. We choose generators $g_1, \dots, g_{\ell-1}$ of \check{J}_A . Notice that $g_1, \dots, g_{\ell-1}, g_{\ell} := w_{n+1} - w_1 \cdot \dots \cdot w_n$ is a basis of $\check{J}_{A'} \subset \mathbb{C}[w_1, \dots, w_{n+1}]$.

Lemma 4.11. *The elements g_1, \dots, g_{ℓ} form a Gröbner basis of $\check{J}_{A'}$ with respect to the weight vector $(0, \dots, 0, -e)$ with $e > 0$.*

Proof. We have already seen that g_1, \dots, g_{ℓ} is a basis of $\check{J}_{A'}$. It remains to prove that $\text{in}_{(0, \dots, 0, -e)}(g_1) = g_1, \dots, \text{in}_{(0, \dots, 0, -e)}(g_{\ell-1}) = g_{\ell-1}, \text{in}_{(0, \dots, 0, -e)}(g_{\ell}) = w_1 \cdot \dots \cdot w_n$ is a basis of $\text{in}_{(0, \dots, 0, -e)}(\check{J}_{A'})$. Let

$$x = \sum_{i=1}^{\ell} x_i g_i \tag{18}$$

and $-e \cdot N := \max\{\text{ord}_{(0, \dots, 0, -e)}(x_i g_i) \mid i = 1, \dots, \ell\}$. Assume that $\text{ord}_{(0, \dots, 0, -e)}(x) < -e \cdot N$, then the maximal w_{n+1} -degree component of the equation (18) is given by

$$0 = \sum_{i=1}^{\ell-1} w_{n+1}^N p_i g_i + w_{n+1}^N p_{\ell} \cdot (w_1 \cdot \dots \cdot w_n)$$

for polynomials $p_i \in \mathbb{C}[w_1, \dots, w_n]$. Since $\check{J}_A = (g_1, \dots, g_{\ell-1})$ is a prime ideal and $w_1 \cdots w_n \notin \check{J}_A$ we conclude that $p_\ell \in \check{J}_A$. Hence there exist polynomial $q_i \in \mathbb{C}[w_1, \dots, w_n]$ such that $p_\ell = \sum_{i=1}^{\ell-1} q_i g_i$. We get

$$x = \sum_{i=1}^{\ell} x_i g_i - \sum_{i=1}^{\ell-1} w_{n+1}^N p_i g_i - w_{n+1}^N p_\ell \cdot g_\ell + w_{n+1}^{N+1} \left(\sum_{i=1}^{\ell-1} q_i g_i \right) = \sum_{i=1}^{\ell} x'_i g_i$$

for $x'_i \in \mathbb{C}[w_1, \dots, w_{n+1}]$ with $\max\{\text{ord}_{(0, \dots, 0, -e)}(x'_i g_i) \mid i = 1, \dots, \ell\} < -e \cdot N$. By induction we can reduce to the case $\text{ord}_{(0, \dots, 0, -e)}(x) = -e \cdot N$. In this case we get for the maximal w_{n+1} -degree component

$$\text{in}_{(0, \dots, 0, -e)}(x) = \sum w_{n+1}^N p'_i g_i + w_{n+1}^N p'_\ell \cdot (w_1 \cdots w_n) = \sum w_{n+1}^N p'_i \text{in}_{(0, \dots, 0, -e)}(g_i) + w_{n+1}^N p'_\ell \text{in}_{(0, \dots, 0, -e)}(g_\ell)$$

for polynomials $p'_i \in \mathbb{C}[w_1, \dots, w_n]$. This shows the claim. \square

Proposition 4.12. *Let A be a $d \times n$ integer matrix such that $\mathbb{N}A = \mathbb{R}_{\geq 0}A \cap \mathbb{Z}^d$ and $\mathbb{N}A \neq \mathbb{Z}^d$. Let A' be the matrix built from A by adding a column which is the sum over all columns of A . Then*

$$\text{fin}_{(u,v)}(\check{I}_{A'}) = \text{in}_{(u,v)}(\check{I}_{A'})$$

if

1. $(u, v) = (0, 0, \dots, 0, 1, 1, \dots, 1)$
2. $(u, v) = (0, \dots, 0, -e, 1, \dots, 1, 1 + e)$ for $0 < e < 1$.

Proof. The first case was proven in [SST00, Corollary 4.36] for homogeneous A . In order to prove the statement for $(u, v) = (0, 0, \dots, 0, 1, 1, \dots, 1)$ in the general case we first observe that $\text{Gr}_{(u,v)}(D_{W'}/\check{J}_{A'})$ is isomorphic to

$$\mathbb{C}[\xi_1, \dots, \xi_{n+1}] \otimes_{\mathbb{C}} \mathbb{C}[\mathbb{N}A']$$

which is Cohen-Macaulay by the assumption $\mathbb{N}A = \mathbb{R}_{\geq 0}A \cap \mathbb{Z}^d$ and the fact that $\mathbb{N}A = \mathbb{N}A'$ as well as $\mathbb{R}_{\geq 0}A = \mathbb{R}_{\geq 0}A'$. It follows from [BZGM15, Theorem 1.2] that the elements $\text{in}_{(u,v)}(\check{E}_k + \beta_k)$ are part of a system of parameters in $\mathbb{C}[\xi_1, \dots, \xi_{n+1}] \otimes_{\mathbb{C}} \mathbb{C}[\mathbb{N}A']$ and since this ring is Cohen-Macaulay they also form a regular sequence. Therefore $H^{-1}(K^\bullet(\text{Gr}_{(u,v)}(D_{W'}/D_{W'}\check{J}_{A'}))) = 0$ and the claim follows from Proposition 4.10.

We prove the second claim. Since $\mathbb{N}A \neq \mathbb{Z}^d$ holds the last column of A' , which is the sum of the columns of A , is non-zero (this was shown above Lemma 4.4). Hence we can assume (by elementary row manipulations of A' , which do not change the ideal $\check{I}_{A'}$) that the last column of A' is zero except for the entry in the first row. Set

$$\check{e}_k := \sum_{i=1}^n a_{ki} w_i \xi_i \quad \text{for } k = 1, \dots, d.$$

We will use the generators $g_1, \dots, g_{\ell-1}$ of \check{J}_A from Lemma 4.11. It follows from [BZGM15, Theorem 1.2] that $\check{e}_1, \dots, \check{e}_d$ is part of a system of parameters for

$$\begin{aligned} \mathbb{C}[\xi_1, \dots, \xi_n] \otimes_{\mathbb{C}} \mathbb{C}[\mathbb{N}A] &\simeq \mathbb{C}[\xi_1, \dots, \xi_n, w_1, \dots, w_n] / \mathbb{C}[\xi_1, \dots, \xi_n, w_1, \dots, w_n] \check{J}_A \\ &\simeq \mathbb{C}[\xi_1, \dots, \xi_n, w_1, \dots, w_n] / (g_1, \dots, g_{\ell-1}) \end{aligned}$$

where $\mathbb{C}[\mathbb{N}A]$ has Krull dimension d . Therefore

$$\mathbb{C}[\xi_1, \dots, \xi_n, w_1, \dots, w_n] / (g_1, \dots, g_{\ell-1}, \check{e}_1, \dots, \check{e}_r)$$

has Krull dimension n .

We will show that the Krull dimension of

$$\mathbb{C}[\xi_1, \dots, \xi_n, w_1, \dots, w_n] / (g_1, \dots, g_{\ell-1}, w_1 \cdots w_n, \check{e}_2, \dots, \check{e}_r) \tag{19}$$

is also n (notice that we omitted \check{e}_1). The variety corresponding to $\mathbb{C}[\xi_1, \dots, \xi_n] \otimes_{\mathbb{C}} \mathbb{C}[\mathbb{N}A]$ is

$$\mathbb{C}^n \times X_A \subset \mathbb{C}^n \times \mathbb{C}^n$$

where $X_A := \text{Spec } \mathbb{C}[\mathbb{N}A]$. The toric variety X_A is a finite disjoint union of torus orbits where the big dense torus lies in $\{w_1 \cdots w_n \neq 0\}$ and the smaller dimensional tori lie in $\{w_1 \cdots w_n = 0\}$. Hence

$$\mathbb{C}[\xi_1, \dots, \xi_n, w_1, \dots, w_n] / (g_1, \dots, g_{\ell-1}, w_1 \cdots w_n) \quad (20)$$

has Krull dimension $n + d - 1$. The torus orbits of X_A correspond to the faces of the cone $\mathbb{R}_{\geq 0}A$ where the big dense torus corresponds to $\mathbb{R}_{\geq 0}A$ itself. For a face $\tau \subsetneq \mathbb{R}_{\geq 0}A$ the torus orbit $\text{Orb}(\tau)$ is given by $\text{Orb}(\tau) = X_A \cap (\mathbb{C}^*)_{\tau}^*$ where $(\mathbb{C}^*)_{\tau}^* = \{w \in \mathbb{C}^n \mid w_i = 0 \text{ for } a_i \notin \tau, w_j \neq 0 \text{ for } a_j \in \tau\}$. Hence it suffices to prove that $(\mathbb{C}^n \times \text{Orb}(\tau)) \cap V((\check{e}_2, \dots, \check{e}_d)$ has dimension n , where $V((\check{e}_2, \dots, \check{e}_d)$ is the vanishing locus of the ideal generated by $\check{e}_2, \dots, \check{e}_d$. Set $\mathbb{C}_{\xi}^{\tau} = \{\xi \in \mathbb{C}^n \mid \xi_i = 0 \text{ for } a_i \notin \tau\}$. It is enough to show that $\mathbb{C}_{\xi}^{\tau} \times \text{Orb}(\tau) \cap V((\check{e}_2^{\tau}, \dots, \check{e}_d^{\tau}))$ has dimension at most $\#\{i \mid a_i \in \tau\}$, where $\check{e}_k^{\tau} := \sum_{i: a_i \in \tau} a_{ki} w_i \xi_i$. The codimension of $V((\check{e}_2^{\tau}, \dots, \check{e}_d^{\tau}))$ is $\dim(\tau)$ since $(1, 0, \dots, 0) = \frac{1}{c}(a_1 + \dots + a_n)$ (for a suitable $c \in \mathbb{Z} \setminus \{0\}$) lies in the interior of $\mathbb{R}_{\geq 0}A$, hence not in τ and therefore the matrix $(a_{ki})_{k \geq 2, i: a_i \in \tau}$ has rank $\dim(\tau)$. By [BZGM15, Lemma 1.1] the intersection of $\mathbb{C}_{\xi}^{\tau} \times \text{Orb}(\tau)$ with $V((\check{e}_2^{\tau}, \dots, \check{e}_d^{\tau}))$ is transverse. Since the codimension of $V((\check{e}_2^{\tau}, \dots, \check{e}_d^{\tau}))$ is $\dim \text{Orb}(\tau) = \dim(\tau)$ the intersection has dimension $\#\{i \mid a_i \in \tau\}$. This shows that the Krull dimension of (19) is n .

Let

$$\begin{aligned} \check{e}'_1 &:= \check{e}_1 + \left(\sum_{i=1}^n a_{1i} \right) x_{n+1} \xi_{n+1} = in_{(0, \dots, 0, -e, 1, \dots, 1, 1+e)}(\check{E}'_1 + \beta_1) \\ \check{e}'_k &:= \check{e}_k = in_{(0, \dots, 0, -e, 1, \dots, 1, 1+e)}(\check{E}'_k + \beta_k) \quad \text{for } k = 2, \dots, d. \end{aligned}$$

and

$$\begin{aligned} \bar{g}_i &:= in_{(0, \dots, 0, -e, 1, \dots, 1, 1+e)}(g_i) = g_i \quad \text{for } i = 1, \dots, \ell - 1 \\ \bar{g}_{\ell} &:= in_{(0, \dots, 0, -e, 1, \dots, 1, 1+e)}(g_{\ell}) = w_1 \cdots w_n \end{aligned}$$

□

Since $\bar{g}_1, \dots, \bar{g}_{\ell}$ and $\check{e}'_2, \dots, \check{e}'_d$ are independent of w_{n+1}, ξ_{n+1} and $\check{e}'_1 = \check{e}_1 + (\sum_{i=1}^n a_{1i}) x_{n+1} \xi_{n+1}$ is (for degree reasons) a non-zero-divisor on

$$\mathbb{C}[w_{n+1}, \xi_{n+1}] \otimes_{\mathbb{C}} \mathbb{C}[\xi_1, \dots, \xi_n, w_1, \dots, w_n] / (\bar{g}_1, \dots, \bar{g}_{\ell}, \check{e}_2, \dots, \check{e}_d),$$

one easily sees that

$$\mathbb{C}[w_1, \dots, w_{n+1}, \xi_1, \dots, \xi_{n+1}] / (\bar{g}_1, \dots, \bar{g}_{\ell}, \check{e}'_1, \dots, \check{e}'_d)$$

has Krull dimension $n + 1$. It follows from (20) that $\mathbb{C}[w_1, \dots, w_{n+1}, \xi_1, \dots, \xi_{n+1}] / (\bar{g}_1, \dots, \bar{g}_{\ell})$ has Krull dimension $(n + d + 1)$, hence $\check{e}'_1, \dots, \check{e}'_d$ is part of a system of parameters. By the assumption on A the ring

$$\mathbb{C}[w_1, \dots, w_{n+1}, \xi_1, \dots, \xi_{n+1}] / (\bar{g}_1, \dots, \bar{g}_{\ell-1}) \simeq \mathbb{C}[w_{n+1}, \xi_1, \dots, \xi_{n+1}] \otimes_{\mathbb{C}} \mathbb{C}[\mathbb{N}A]$$

is Cohen-Macaulay. Since $\bar{g}_{\ell} = w_1 \cdots w_n$ is not a zero-divisor in the ring above (because $\mathbb{C}[\mathbb{N}A]$ has no non-zero zero-divisors), we see that the ring

$$\mathbb{C}[w_1, \dots, w_{n+1}, \xi_1, \dots, \xi_{n+1}] / (\bar{g}_1, \dots, \bar{g}_{\ell})$$

is also Cohen-Macaulay and therefore $\check{e}'_1, \dots, \check{e}'_d$ is a regular sequence in $\mathbb{C}[w_1, \dots, w_{n+1}, \xi_1, \dots, \xi_{n+1}] / (\bar{g}_1, \dots, \bar{g}_{\ell})$.

Since

$$\text{Gr}_{(0, \dots, 0, -e, 1, \dots, 1, 1+e)}(D_{W'} / D_{W'} \check{J}_{A'}) \simeq \mathbb{C}[w_1, \dots, w_{n+1}, \xi_1, \dots, \xi_{n+1}] / (\bar{g}_1, \dots, \bar{g}_{\ell})$$

and $\check{e}'_k = in_{(0, \dots, 0, -e, 1, \dots, 1, 1+e)}(\check{E}'_k + \beta_k)$ for $k = 1, \dots, d$, we have

$$H^{-1}(K^{\bullet}(\text{Gr}_{(0, \dots, 0, -e, 1, \dots, 1, 1+e)}(D_{W'} / D_{W'} \check{J}_{A'}))) = 0.$$

Using again Proposition 4.10, this shows the second claim.

Corollary 4.13. *Let $g_1, \dots, g_\ell \in \mathbb{C}[w_1, \dots, w_{n+1}]$ be the generators of $\check{J}_{A'}$ defined above Lemma 4.11.*

1. *The $(g_i)_{i=1, \dots, \ell}$ together with $(\check{E}'_k + \beta_k)_{k=1, \dots, d}$ form a Gröbner basis of $\check{I}_{A'}$ with respect to the weight vector $(u, v) = (0, \dots, 0, 1, \dots, 1)$.*
2. *Let $(u, v) = (0, \dots, 0, 1, \dots, 1)$ and set $\tilde{g}_i := in_{(u, v)}(g_i)$ and $\check{E}'_k = in_{(u, v)}(\check{E}'_k + \beta_k)$. The elements $(\tilde{g}_i)_{i=1, \dots, \ell}$ and $(\check{E}'_k)_{k=1, \dots, d}$ form a Gröbner-basis of*

$$\begin{aligned} in_{(u, v)}(\check{I}_{A'}) &= in_{(u, v)}((g_1, \dots, g_\ell, \check{E}'_1 + \beta_1, \dots, \check{E}'_d + \beta_d)) \\ &= (\tilde{g}_1, \dots, \tilde{g}_\ell, \check{E}'_1, \dots, \check{E}'_d) \subset \mathbb{C}[w_1, \dots, w_{n+1}, \xi_1, \dots, \xi_{n+1}] \end{aligned}$$

with respect to the weight vector $(0, \dots, 0, -1, 0, \dots, 0, 1)$ (i.e. w_{n+1} has weight -1 and ξ_{n+1} has weight $+1$).

Proof. 1.) The set $(g_i)_{i=1, \dots, \ell}$ is a Gröbner basis for $\check{J}_{A'}$. Therefore the elements $in_u(g_i) = g_i$ generate $in_u(\check{J}_{A'}) = \check{J}_{A'}$. The elements $(g_i)_{i=1, \dots, \ell}$ and $(\check{E}_k + \beta_k)_{k=1, \dots, d}$ generate $\check{I}_{A'}$ and the elements $(in_{u, v}(g_i))_{i=1, \dots, \ell}$ and $(in_{(u, v)}(\check{E}_k + \beta_k))_{k=1, \dots, d}$ generate $fin_{(u, v)}$ by definition. The claim follows now from Proposition 4.12 1. .

2.) It follows from the first point that the $\tilde{g}_i = in_{(u, v)}(g_i)$ and the $\check{E}'_k = in_{(u, v)}(\check{E}'_k + \beta_k)$ generate $in_{(u, v)}(\check{I}_{A'})$. We have to show that the $in_{(0, \dots, 0, -1, 0, \dots, 0, 1)}(\tilde{g}_i)$ for $i = 1, \dots, \ell$ and the $in_{(0, \dots, 0, -1, 0, \dots, 0, 1)}(\check{E}'_k)$ generate $in_{(0, \dots, 0, -1, 0, \dots, 0, 1)}(in_{(u, v)}(\check{I}_{A'}))$. But this follows from (cf. [SST00, Lemma 2.1.6 (2)])

$$\begin{aligned} in_{(0, \dots, 0, -1, 0, \dots, 0, 1)}(\tilde{g}_i) &= in_{(0, \dots, 0, -e, 1, \dots, 1, 1+e)}(g_i) && \text{for } i = 1, \dots, \ell \\ in_{(0, \dots, 0, -1, 0, \dots, 0, 1)}(\check{E}'_k) &= in_{(0, \dots, 0, -e, 1, \dots, 1, 1+e)}(\check{E}'_k + \beta_k) && \text{for } k = 1, \dots, d \\ in_{(0, \dots, 0, -1, 0, \dots, 0, 1)}(in_{(u, v)}(\check{I}_{A'})) &= in_{(0, \dots, 0, -e, 1, \dots, 1, 1+e)}(\check{I}_{A'}) \end{aligned}$$

for $0 < e \ll 1$ and Proposition 4.12 2. . □

The second notion we are going to introduce relates the order filtration F_\bullet on $D_{W'}$ with the V -filtration that already occurred in the last subsection. Here we consider the descending V -filtration on $D_{W'}$ with respect to $w_{n+1} = 0$, which we denote again by $V^\bullet D_{W'}$. We have

$$V^0 D_{W'} = \sum_{i, k \geq 0} (\partial_{w_{n+1}} w_{n+1})^i (w_{n+1})^k P_i$$

for $P_i \in \mathbb{C}[w_1, \dots, w_n] \langle \partial_{w_1}, \dots, \partial_{w_n} \rangle$ and

$$V^k D_{W'} = w_{n+1}^k V^0 D_{W'} \quad \text{and} \quad V^{-k} D_{W'} = \sum_{j \geq 0} \partial_{w_{n+1}}^j V^0 D_{W'} \quad (21)$$

for $k > 0$.

Recall the left ideal $\check{I}_{A'} \subset D_{W'}$ and the left $D_{W'}$ -modules $\check{M}_{A'}^\beta := D_{W'} / \check{I}_{A'}$ from above. We define filtrations V_{ind}^\bullet and F_\bullet^{ord} on $\check{M}_{A'}^\beta$ by:

$$V_{ind}^k \check{M}_{A'}^\beta := \frac{V^k D_{W'} + \check{I}_{A'}}{\check{I}_{A'}} \quad \text{and} \quad F_p^{ord} \check{M}_{A'}^\beta := \frac{F_p D_{W'} + \check{I}_{A'}}{\check{I}_{A'}}.$$

We are now ready to prove the main result of this subsection.

Proof of Proposition 4.9. Let $m \in V_{ind}^k \check{M}_{A'}^\beta \cap F_p^{ord} \check{M}_{A'}^\beta$. We can find $P, Q \in D_{W'}$ such that $P \in F_p D_{W'}$, $Q \in V^k D_{W'}$ and $[P] = m = [Q]$, i.e. $P = Q - i$ for some $i \in \check{I}_{A'}$. We have to find a Q' with $Q' \in V^k D_{W'} \cap F_p D_{W'}$ with $P = Q' - i'$ for $i' \in I$. We will construct this element Q' by decreasing induction on the order of Q by killing its leading term in each step. For this we will use the special Gröbner basis of $\check{I}_{A'}$ which we constructed in Corollary 4.13 above.

Recall that the weight vector $(u, v) := (0, \dots, 0, 1, \dots, 1)$ induces the order filtration $F_{\bullet}^{(u,v)} = F_{\bullet}^{ord}$ on $D_{W'}$. If $R \in D_{W'}$ and $k := ord_{(u,v)}(R)$ we define the symbol of R by $\sigma_k(R) = in_{(u,v)}(R)$ and set $\sigma_q(R) = 0$ for $q \neq k$. We define a second weight vector $(u', v') := (0, \dots, 0, -1, 0, \dots, 0, 1)$ which induces the descending V -filtration from (21) on $D_{W'}$. The V -filtration and F -filtration also induce filtrations \tilde{V} and \tilde{F} on $\mathrm{Gr}_{\bullet}^{(u,v)} D_{W'} = \mathrm{Gr}_{\bullet}^F D_{W'} = \mathbb{C}[w_1, \dots, w_{n+1}, \xi_1, \dots, \xi_{n+1}]$.

Let $t_Q := ord_{(u,v)} Q$, $t_i := ord_{(u,v)} i$ and set $t := \max(t_Q, t_i)$. Obviously we have $t \geq p$. If $t = p$ we are done. Hence, we assume $t > p$, thus we have

$$0 = \sigma_t(P) = \sigma_t(Q - i)$$

and therefore $t = t_Q = t_i$ which implies $\sigma_t(Q) = \sigma_t(i) \neq 0$. Set $k_Q := ord_{(u',v')}(\sigma_t(Q))$, then we have $\sigma_t(i) = \sigma_t(Q) \in \tilde{V}^{k_Q}$.

Recall from Corollary 4.13 that $\tilde{I}_{A'}$ is generated by $\{G_1, \dots, G_m\} := \{g_1, \dots, g_\ell, \check{E}'_1 + \beta_1, \dots, \check{E}'_d + \beta_d\}$ and these elements form a Gröbner basis with respect to weight vector (u, v) and their initial forms

$$\{\tilde{G}_1, \dots, \tilde{G}_m\} := \{in_{(u,v)}(G_1), \dots, in_{(u,v)}(G_m)\}$$

are a Gröbner basis of $in_{(u,v)}(\tilde{I}_{A'})$ with respect to the weight vector (u', v') . Therefore we can write

$$\sigma_t(i) = \sum_{l=1}^m \tilde{i}_l \tilde{G}_l$$

with $\tilde{i}_l \in \mathbb{C}[w_1, \dots, w_{n+1}, \xi_1, \dots, \xi_{n+1}]$. Using a commutative version of [SST00, Theorem 1.2.10] we can assume that $\tilde{i}_l \in \tilde{V}^{k_Q - k_l}$ where $k_l = ord_{(u',v')}(\tilde{G}_l)$. Since the elements \tilde{G}_l are homogeneous with respect to the variables ξ_1, \dots, ξ_{n+1} we can also assume that $\tilde{i}_l \in \tilde{F}_{t-t_l} \mathrm{Gr}_{\bullet}^F D_{W'}$ where $t_l = ord_{(u,v)}(\tilde{G}_l)$. Let $i_l \in D_{W'}$ be the normally ordered element which we obtain from \tilde{i}_l by replacing ξ_i with ∂_{w_i} . One sees easily that $i_l G_l \in F_t D_{W'} \cap V^{k_Q} D_{W'}$. Therefore the element $i' := \sum_{l=1}^m i_l G_l$ has the following two properties

$$\sigma_t(i') = \sigma_t(i) = \sigma_t(Q) \quad \text{and} \quad i' \in V^{k_Q} D_{W'}$$

where the second property follows from $ord_{(u',v')}(G_l) = ord_{(u',v')}(\tilde{G}_l)$. We therefore have

$$P = Q - i' - (i - i')$$

with $Q - i' \in F_{t-1} D_{W'} \cap V^{k_Q} D_{W'}$. Since obviously we have $k \leq k_Q$, we conclude that $Q - i' \in F_{t-1} D_{W'} \cap V^k D_{W'}$. The claim now follows by descending induction on the order t . \square

4.3 Calculation of the Hodge filtration

In this subsection we want to compute the Hodge filtration on the mixed Hodge module

$$\mathcal{H}^0(h_{A*} {}^p \mathcal{C}_T^{\beta,H}),$$

recall from section 1 that ${}^p \mathcal{C}_T^{\beta,H} = \mathbb{C}_T^{\beta,H}[d] \in \mathrm{MHM}(T)$. Also recall that ${}^p \mathcal{C}_T^{\beta,H}$ has the underlying filtered \mathcal{D} -module $(\mathcal{O}_T^\beta, F_{\bullet}^H \mathcal{O}_T^\beta)$, where the Hodge filtration is given by

$$F_p^H \mathcal{O}_T^\beta = \begin{cases} \mathcal{O}_T^\beta & \text{for } p \geq 0 \\ 0 & \text{else.} \end{cases}$$

We will use several different presentations of \mathcal{O}_T^β as a \mathcal{D}_T -module, namely, for each $\alpha = (\alpha_k)_{k=1, \dots, d} \in \mathbb{Z}^d$ we have a \mathcal{D}_T -linear isomorphism

$$\Gamma(T, \mathcal{O}_T^\beta) \simeq D_T / (\partial_{t_k} t_k + \beta_k + \alpha_k)_{k=1, \dots, d}$$

such that the Hodge filtration is simply the order filtration on the right hand side.

As we have seen in Lemma 4.1, the morphism h_A can be decomposed into the closed embedding $k_A : T \rightarrow W^* = W \setminus D$ and the canonical open embedding $l_A : W^* \rightarrow W$. We have to determine the Hodge filtration on the direct image modules for both mappings. The former is (after some coordinate change) a rather direct calculation, and will be carried out in Lemma 4.14 below. However, understanding the behaviour of the Hodge filtration under the direct image of an open embedding of the complement of a divisor (like the map l_A) is more subtle and at the heart of the theory of mixed Hodge modules (see, e.g., [Sai90, Section (2.b)]). More precisely, since the steps of the Hodge filtration of a mixed Hodge module are coherent modules over the structure sheaf of the underlying variety, the usual direct image functors are not suitable for the case of an open embedding as they do not preserve coherence. In order to circumvent this difficulty, one uses the canonical V -filtration along the boundary divisor, as computed in subsection 4.1 above. Let us give an overview of the strategy to be used below. The actual calculation will be finished only in Theorem 4.17, the main step being Proposition 4.16.

We will need the following formula (copied from [Sai93, Proposition 4.2.]) which describes the extension of a mixed Hodge module over a smooth hypersurface. Let X be a smooth variety, let t, x_1, \dots, x_n be local coordinates on X and $j : Y \hookrightarrow X$ be a smooth hypersurface given by $t = 0$. Let ${}^H\mathcal{M}$ be a mixed Hodge module on $X \setminus Y$ with underlying filtered \mathcal{D} -module $(\mathcal{M}, F_\bullet^H \mathcal{M})$, then

$$F_p^H \mathcal{H}^0 j_+ \mathcal{M} = \sum_{i \geq 0} \partial_t^i F_{p-i}^H V^0 \mathcal{H}^0 j_+ \mathcal{M}, \quad \text{where} \quad F_q^H V^0 \mathcal{H}^0 j_+ \mathcal{M} := V^0 \mathcal{H}^0 j_+ \mathcal{M} \cap j_*(F_q^H \mathcal{M}), \quad (22)$$

here $V^0 \mathcal{H}^0 j_+ \mathcal{M}$ is the canonical V -filtration on the \mathcal{D} -module $\mathcal{H}^0 j_+ \mathcal{M} \simeq j_* \mathcal{M}$, as introduced in Definition 4.2.

If Y is a non-smooth hypersurface locally given by $g = 0$, we consider (locally) the graph embedding

$$\begin{aligned} i_g : X &\longrightarrow X \times \mathbb{C}_t \\ x &\mapsto (x, g(x)) \end{aligned}$$

together with its restriction $i_g^\circ : X \setminus Y \rightarrow X \times \mathbb{C}_t^*$. Notice that i_g° is a closed embedding. Given a mixed Hodge module \mathcal{M} on $X \setminus Y$ we proceed as follows. We first extend the Hodge filtration of $(i_g^\circ)_+ \mathcal{M}$ over the smooth divisor given by $\{t = 0\}$ as explained above. Afterward, we restrict the mixed Hodge module which we obtained to the smooth divisor given by $\{t = g\}$.

After these general remarks, we come back to the situation of the torus embedding $h_A : T \rightarrow W$ described at the beginning of this section. Consider the following commutative diagram

$$\begin{array}{ccccc} & & h_A & & \\ & \curvearrowright & & \searrow & \\ T & \xrightarrow{k_A} & W^* & \xrightarrow{l_A} & W & \xrightarrow{i_g} & W \times \mathbb{C}_t \\ & & & \searrow^{i_g^\circ} & & \uparrow^{j_t} & \\ & & & & & W \times \mathbb{C}_t^* & \end{array} \quad (23)$$

where $W^* := W \setminus D = W \setminus \{w_1 \dots w_n = 0\} \simeq (\mathbb{C}^*)^n$ and where i_g is the graph embedding

$$\begin{aligned} i_g : W &\longrightarrow W \times \mathbb{C}_t \\ w &\mapsto (w, w_1 \dots w_n) \end{aligned} \quad (24)$$

associated with the function $g : W \rightarrow \mathbb{C}_t, w \mapsto w_1 \dots w_n$. Notice that $i_g \circ l_A$ factors over $W \times \mathbb{C}_t^*$. We have the following isomorphisms

$$i_{g+h_A} \mathcal{O}_T^\beta \simeq i_{g+l_A} k_{A+} \mathcal{O}_T^\beta \simeq j_{t+} i_{g+}^\circ k_{A+} \mathcal{O}_T^\beta.$$

Lemma 4.14. *The direct image $\mathcal{H}^0 k_{A+} \mathcal{O}_T^\beta$ is isomorphic to the cyclic \mathcal{D}_{W^*} -module*

$${}^* \check{\mathcal{M}}_{A'}^\beta := \mathcal{D}_{W^*} / \check{\mathcal{I}}_\beta^*$$

where $\check{\mathcal{I}}_\beta^*$ is the left ideal generated by $(\check{E}_k + \beta_k)_{k=1, \dots, d}$ for $\beta = (\beta_k)_{k=1, \dots, d} \in \mathbb{R}^d$ and $(\check{\square}_m)_{m \in \mathbb{L}_A}$. Furthermore, the Hodge filtration on $\check{\mathcal{M}}_{A'}$ is equal to the induced order filtration, shifted by $n - d$, i.e.

$$F_p^H {}^* \check{\mathcal{M}}_{A'} = F_{p-(n-d)}^{ord} \mathcal{D}_{W^*} / \check{\mathcal{I}}_\beta^*.$$

Proof. We factor the map k_A from above in the following way. Let $A = C \cdot E \cdot F$ be the Smith normal form of A , i.e. $C = (c_{pq}) \in GL(d, \mathbb{Z})$, $F = (f_{uv}) \in GL(n, \mathbb{Z})$ and $E = (I_d, 0_{d, n-d})$. This gives rise to the maps

$$\begin{aligned} k_C : T &\longrightarrow T \\ (t_1, \dots, t_d) &\mapsto (\tilde{t}_1, \dots, \tilde{t}_d) = (\underline{t}^{c_1}, \dots, \underline{t}^{c_d}) \\ k_E : T &\longrightarrow (\mathbb{C}^*)^n \\ (\tilde{t}_1, \dots, \tilde{t}_d) &\mapsto (\tilde{w}_1, \dots, \tilde{w}_n) = (\tilde{t}_1, \dots, \tilde{t}_d, 1, \dots, 1) \\ k_F : (\mathbb{C}^*)^n &\longrightarrow W^* \\ (\tilde{w}_1, \dots, \tilde{w}_n) &\mapsto (w_1, \dots, w_n) = (\underline{\tilde{w}}^f, \dots, \underline{\tilde{w}}^f). \end{aligned}$$

For $\gamma \in \mathbb{Z}^d$ we have

$$k_{A+} \mathcal{O}_T^\gamma \simeq (k_F \circ k_E \circ k_C)_+ \mathcal{O}_T^\gamma \simeq k_{F+} k_{E+} k_{C+} \mathcal{O}_T^\gamma.$$

Since all maps and spaces involved are affine, we will work at the level of global sections. We have $\Gamma(T, \mathcal{O}_T^\gamma) = D_T / (\partial_{t_k} t_k + \gamma_k)_{k=1, \dots, d} = D_T / (t_k \partial_{t_k} + \gamma_k + 1)_{k=1, \dots, d}$. Notice that the Hodge filtration in this presentation is simply the order filtration. Since k_C is a change of coordinates we have $\Gamma(T, \mathcal{H}^0(k_{C+} \mathcal{O}_T^\gamma)) \simeq D_T / (\sum_{i=1}^d c_{ki} \tilde{t}_i \partial_{\tilde{t}_i} + \gamma_k + 1)_{k=1, \dots, d}$ and again the Hodge filtration is equal to the order filtration. We now calculate $\Gamma((\mathbb{C}^*)^n, k_{E+} k_{C+} \mathcal{O}_T^\gamma)$. We have

$$\begin{aligned} \Gamma((\mathbb{C}^*)^n, \mathcal{H}^0 k_{E+} k_{C+} \mathcal{O}_T^\gamma) &\simeq \Gamma(T, \mathcal{H}^0 k_{C+} \mathcal{O}_T^\gamma) [\partial_{\tilde{w}_{d+1}}, \dots, \partial_{\tilde{w}_n}] \\ &\simeq D_{(\mathbb{C}^*)^n} / \left(\sum_{i=1}^d c_{ki} \tilde{w}_i \partial_{\tilde{w}_i} + \gamma_k + 1 \right)_{k=1, \dots, d}, (\tilde{w}_i - 1)_{i=d+1, \dots, n}. \end{aligned} \quad (25)$$

The Hodge filtration is (cf. [Sai93, Formula (1.8.6)])

$$\begin{aligned} F_{p+(n-d)}^H \left(\Gamma((\mathbb{C}^*)^n, \mathcal{H}^0 k_{E+} k_{C+} \mathcal{O}_T^\gamma) \right) &= \sum_{p_1+p_2=p} F_{p_1}^H \Gamma(T, \mathcal{H}^0 k_{C+} \mathcal{O}_T^\gamma) \otimes \underline{\partial}^{p_2} \\ &= \sum_{p_1+p_2=p} F_{p_1}^{ord} \Gamma(T, \mathcal{H}^0 k_{C+} \mathcal{O}_T^\gamma) \otimes \underline{\partial}^{p_2} = F_p^{ord} \left(\Gamma((\mathbb{C}^*)^n, \mathcal{H}^0 k_{E+} k_{C+} \mathcal{O}_T^\gamma) \right). \end{aligned} \quad (26)$$

Hence we see that the Hodge filtration on the presentation (25) shifted by $(n-d)$ is equal to the order filtration, i.e. $F_{p+(n-d)}^H = F_p^{ord}$.

The map k_F is again a change of coordinates, so we have

$$\begin{aligned} \Gamma((\mathbb{C}^*)^n, \mathcal{H}^0 k_{F+} k_{E+} k_{C+} \mathcal{O}_T^\gamma) &\simeq D_{(\mathbb{C}^*)^n} / \left(\left(\sum_{j=1}^n a_{kj} w_j \partial_{w_j} + \gamma_k + 1 \right)_{k=1, \dots, d}, (\underline{w}^{\underline{m}_i} - 1)_{i=d+1, \dots, n} \right) \\ &\simeq D_{(\mathbb{C}^*)^n} / \left(\left(\sum_{j=1}^n a_{kj} w_j \partial_{w_j} + \gamma_k + 1 \right)_{k=1, \dots, d}, (\check{\square}_{\underline{m}})_{\underline{m} \in \mathbb{L}_A} \right), \end{aligned} \quad (27)$$

where \underline{m}_i are the columns of the inverse matrix $M = F^{-1}$. The first isomorphism follows from the equality $A = C \cdot E \cdot F$. The second isomorphism follows from the fact that an element $\underline{m} \in \mathbb{Z}^n$ is a relation between the columns of A if and only if it is a relation between the columns of $E \cdot F$. So the Hodge filtration on the presentation (27) shifted by $(n-d)$ is again the order filtration. We have

$$\sum_{j=1}^n a_{kj} w_j \partial_{w_j} + \gamma_k + 1 = \sum_{j=1}^n a_{kj} \partial_{w_j} w_j - \sum_{j=1}^n a_{kj} + \gamma_k + 1.$$

Setting $\gamma_k := \sum_{j=1}^n a_{kj} + \beta_k - 1$, shows that

$$*\check{\mathcal{M}}_{A'}^\beta = \mathcal{H}^0 k_{A+} \mathcal{O}_T^{\sum_{j=1}^n a_j + \beta - 1} \simeq \mathcal{H}^0 k_{A+} \mathcal{O}_T^\beta \quad (28)$$

where the last isomorphism is given by right multiplication with $\underline{t}^{-\sum_{j=1}^n a_j + 1}$ (here $\mathbf{1} := (1, \dots, 1) \in \mathbb{Z}^d$). \square

The next step is to compute the Hodge filtration of $h_{A+}\mathcal{O}_T^\beta \simeq l_{A+}k_{A+}\mathcal{O}_T^\beta$ from that of $k_{A+}\mathcal{O}_T^\beta$. As the map l_A is an open embedding of the complement of a normal crossing divisor, we need to consider the graph embedding embedding i_g° with respect to the function $g = w_1 \cdots w_n$. We proceed as described at the beginning of this section, i.e., we first extend the module $i_{g+}^\circ k_{A+}\mathcal{O}_T^\beta$ over the smooth divisor $\{t = 0\}$.

Lemma 4.15. *The direct image $i_{g+}^\circ k_{A+}\mathcal{O}_T^\beta$ is isomorphic to the cyclic $\mathcal{D}_{W \times \mathbb{C}_t^*}$ -module $\mathcal{D}_{W \times \mathbb{C}_t^*}/\mathcal{I}^\circ$ where \mathcal{I}° is the left ideal generated by $(\check{E}'_k + \beta_k)_{k=1, \dots, d}$ for $\beta = (\beta_k)_{k=1, \dots, d} \in \mathbb{R}^d$ and $(\check{\square}_m)_{m \in \mathbb{L}_{A'}}$. Recall that the vector fields \check{E}'_k have been defined in formula (13) as $\check{E}'_k := \sum_{i=1}^n b_{ki} \partial_{w_i} + c_k \partial_t$ for $k = 1, \dots, d$. Furthermore, the Hodge filtration shifted by $n - d + 1$ is equal to the induced order filtration, i.e., we have*

$$F_p^H i_{g+}^\circ k_{A+}\mathcal{O}_T^\beta \simeq F_{p-(n-d+1)}^{ord} \mathcal{D}_{W \times \mathbb{C}_t^*} / \mathcal{I}^\circ.$$

Proof. We define

$$\widetilde{W} := (W^* \times \mathbb{C}_t^*) \setminus \{\tilde{t} + g(\underline{w}) = 0\}$$

and factor the map i_g° in the following way. Set

$$\begin{aligned} l_1 : W^* &\longrightarrow \widetilde{W} \\ &\underline{w} \mapsto (\underline{w}, 0) \\ l_2 : \widetilde{W} &\longrightarrow W^* \times \mathbb{C}_t^* \\ &(\underline{w}, \tilde{t}) \mapsto (\underline{w}, \tilde{t} + g(\underline{w})) \end{aligned}$$

and let $l_3 : W^* \times \mathbb{C}_t^* \rightarrow W \times \mathbb{C}_t^*$ be the canonical inclusion. We have $i_g^\circ = l_3 \circ l_2 \circ l_1$. For the convenience of the reader, let us summarize these maps in the following diagram

$$\begin{array}{ccccc} W^* & \xrightarrow[\text{open embedding}]{l_A} & W & \xrightarrow{i_g} & W \times \mathbb{C}_t \\ \downarrow \text{closed } l_1 & \searrow i_g^\circ & & & \uparrow \text{open embedding} \\ \widetilde{W} & \xrightarrow[l_2]{\simeq} & W^* \times \mathbb{C}_t^* & \xrightarrow[l_3]{\text{closed on support}} & W \times \mathbb{C}_t^* \\ & & & & \uparrow \text{open embedding} \\ & & & & W \times \mathbb{C}_t \end{array}$$

Notice again that all spaces involved are affine, hence we will work with the modules of global sections. Since l_1 is just the inclusion of a coordinate hyperplane we have

$$\Gamma(\widetilde{W}, \mathcal{H}^0 l_{1+} k_{A+} \mathcal{O}_T^\beta) \simeq \Gamma(W^*, \mathcal{H}^0 k_{A+} \mathcal{O}_T^\beta)[\partial_{\tilde{t}}].$$

The Hodge filtration is given by

$$\Gamma(\widetilde{W}, F_{p+1}^H(\mathcal{H}^0 l_{1+} k_{A+} \mathcal{O}_T^\beta)) \simeq \sum_{p_1+p_2=p} \Gamma(W^*, F_{p_1}^H \mathcal{H}^0 k_{A+} \mathcal{O}_T^\beta) \otimes \partial_{\tilde{t}}^{p_2}. \quad (29)$$

Notice that $\Gamma(\widetilde{W}, \mathcal{H}^0 l_{1+} k_{A+} \mathcal{O}_T^\beta) \simeq D_{\widetilde{W}}/I_1^\circ$ where I_1° is the left ideal generated by $(\check{E}'_k + \beta_k)_{k=1, \dots, d}$, $(\check{\square}_m)_{m \in \mathbb{L}_A}$ and \tilde{t} .

Under this isomorphism the Hodge filtration on $\Gamma(\widetilde{W}, \mathcal{H}^0 l_{1+} k_{A+} \mathcal{O}_T^\beta)$ shifted by $(n - d) + 1$ is equal to the order filtration by Lemma 4.14 and (29). The map l_2 is just a change of coordinates, hence under the substitutions $\tilde{t} \mapsto t = \tilde{t} + g(\underline{w})$, $w_i \partial_{w_i} \mapsto w_i \partial_{w_i} + g(\underline{w}) \partial_t \equiv w_i \partial_{w_i} + \partial_t t$ for $i = 1, \dots, n$ and by using the presentation of $\mathcal{H}^0(k_{A+}\mathcal{O}_T^\beta)$ as acyclic \mathcal{D} -module, we get that

$$\begin{aligned} \Gamma(W^* \times \mathbb{C}_t^*, \mathcal{H}^0 l_{2+} l_{1+} k_{A+} \mathcal{O}_T^\beta) &\simeq D_{W^* \times \mathbb{C}_t^*} / ((\check{E}'_k + \beta_k)_{k=1, \dots, d} + (\check{\square}_m)_{m \in \mathbb{L}_A} + (t - w_1 \cdots w_n)) \\ &\simeq D_{W^* \times \mathbb{C}_t^*} / ((\check{E}'_k + \beta_k)_{k=1, \dots, d} + (\check{\square}_m)_{m \in \mathbb{L}_{A'}}), \end{aligned} \quad (30)$$

where \check{E}'_k was defined in formula (13) and A' is the matrix defined just before that formula. Notice that the Hodge filtration shifted by $(n - d) + 1$ is again equal to the order filtration.

Since the support of $\mathcal{H}^0 l_{2+l_1+k_{A+}} \mathcal{O}_T^\beta$ lies in the subvariety $\{t = g(\underline{w})\}$, the closure of the support in $W \times \mathbb{C}_t^*$ does not meet $D \times \mathbb{C}_t^*$. We conclude that

$$\begin{aligned} \Gamma(W \times \mathbb{C}_t^*, \mathcal{H}^0 i_{g+}^\circ k_{A+} \mathcal{O}_T^\beta) &\simeq \Gamma(W \times \mathbb{C}_t^*, \mathcal{H}^0 l_{3+l_2+l_1+k_{A+}} \mathcal{O}_T^\beta) \\ &\simeq \Gamma(W^* \times \mathbb{C}_t^*, \mathcal{H}^0 l_{2+l_1+k_{A+}} \mathcal{O}_T^\beta) \\ &\simeq D_{W^* \times \mathbb{C}_t^*} / ((\check{E}'_k + \beta_k)_{k=1, \dots, d} + (\check{\square}_m)_{m \in \mathbb{L}_{A'}}) \\ &\simeq D_{W \times \mathbb{C}_t^*} / ((E'_k + \beta_k)_{k=1, \dots, d} + (\check{\square}_m)_{m \in \mathbb{L}_{A'}}). \end{aligned}$$

The Hodge filtration is then simply extended by using the following formula

$$F_p^H \mathcal{H}^0 i_{g+}^\circ k_{A+} \mathcal{O}_T^\beta \simeq F_p^H \mathcal{H}^0 l_{3+l_2+l_1+k_{A+}} \mathcal{O}_T^\beta \simeq l_{3*} F_p^H \mathcal{H}^0 l_{2+l_1+k_{A+}} \mathcal{O}_T^\beta.$$

□

Proposition 4.16. *Let $\beta \in \mathbb{R}^d \setminus sRes(A)$. The direct image $\mathcal{H}^0 j_{t+} i_{g+}^\circ k_{A+} \mathcal{O}_T^\beta$ is isomorphic to the quotient $\check{\mathcal{M}}_{A'}^\beta = \mathcal{D}_{W \times \mathbb{C}_t} / \mathcal{I}'$, where \mathcal{I}' is the left ideal generated by $(\check{E}'_k + \beta_k)_{k=1, \dots, d}$ and $(\check{\square}_m)_{m \in \mathbb{L}_{A'}}$. Furthermore, if $\beta \in \mathfrak{A}_A$, then the Hodge filtration on $\mathcal{D}_{W \times \mathbb{C}_t} / \mathcal{I}'$ shifted by $(n-d)+1$ is equal to the induced order filtration, that is,*

$$F_p^H \mathcal{H}^0 j_{t+} i_{g+}^\circ k_{A+} \mathcal{O}_T^\beta = F_{p-(n-d+1)}^{ord} \mathcal{D}_{W \times \mathbb{C}_t} / \mathcal{I}'.$$

Proof. First recall that we have an isomorphism $\mathcal{H}^0 j_{t+} i_{g+}^\circ k_{A+} \mathcal{O}_T^\beta \simeq \mathcal{H}^0 i_{g+} h_{A+} \mathcal{O}_T^\beta$. The composed map $i_g \circ h_A$ is a torus embedding given by the matrix A' . Hence, we have an isomorphism $\mathcal{H}^0 j_{t+} i_{g+}^\circ k_{A+} \mathcal{O}_T^\beta \simeq \check{\mathcal{M}}_{A'}^\beta$, for $\beta \in \mathbb{R}^d$ and $\beta \notin sRes(A') = sRes(A)$. This shows the first claim.

For the second statement, suppose that $\beta \in \mathfrak{A}_A$. The formula for extending the Hodge filtration over the smooth divisor $\{t = 0\}$ is

$$F_p^H \check{\mathcal{M}}_{A'}^\beta = \sum_{i \geq 0} \partial_t^i (V^0 \check{\mathcal{M}}_{A'}^\beta \cap j_{t*} j_t^{-1} F_{p-i}^H \check{\mathcal{M}}_{A'}^\beta). \quad (31)$$

At the level of global sections the adjunction morphism $\check{\mathcal{M}}_{A'}^\beta \rightarrow j_{t+} j_t^+ \check{\mathcal{M}}_{A'}^\beta$ is given by the inclusion $\check{\mathcal{M}}_{A'}^\beta \rightarrow {}^* \check{\mathcal{M}}_{A'}^\beta$, where ${}^* \check{\mathcal{M}}_{A'}^\beta$ is the $D_{W \times \mathbb{C}_t}$ -module from Lemma 4.15 seen as a $D_{W \times \mathbb{C}_t}$ -module. Hence, at the level of global sections, formula (31) becomes

$$F_p^H \check{\mathcal{M}}_{A'}^\beta = \sum_{i \geq 0} \partial_t^i (V^0 \check{\mathcal{M}}_{A'}^\beta \cap F_{p-i}^H {}^* \check{\mathcal{M}}_{A'}^\beta).$$

Since we have $F_{p+(n-d+1)}^H {}^* \check{\mathcal{M}}_{A'}^\beta = F_p^{ord} {}^* \check{\mathcal{M}}_{A'}^\beta$, by the same lemma, we conclude that $F_{n-d}^H \check{\mathcal{M}}_{A'}^\beta = 0$. The element $1 \in \check{\mathcal{M}}_{A'}^\beta$ is in $V^0 \check{\mathcal{M}}_{A'}^\beta$ by Proposition 4.8 and $1 \in F_{(n-d)+1}^H {}^* \check{\mathcal{M}}_{A'}^\beta = F_0^{ord} {}^* \check{\mathcal{M}}_{A'}^\beta$, and therefore $1 \in F_{(n-d)+1}^H \check{\mathcal{M}}_{A'}^\beta$. Notice that both $(\check{\mathcal{M}}_{A'}^\beta, F_\bullet^H)$ and $(\check{\mathcal{M}}_{A'}^\beta, F_\bullet^{ord})$ are cyclic, well-filtered $D_{W \times \mathbb{C}_t}$ -modules (see e.g. [Sai88, Section 2.1.1]), therefore we can conclude

$$F_p^{ord} \check{\mathcal{M}}_{A'}^\beta \subset F_{p+(n-d+1)}^H \check{\mathcal{M}}_{A'}^\beta.$$

In order to show the reverse inclusion, we have to show

$$F_p^{ord} \check{\mathcal{M}}_{A'}^\beta \supset F_{p+(n-d+1)}^H \check{\mathcal{M}}_{A'}^\beta = \sum_{i \geq 0} \partial_t^i (V^0 \check{\mathcal{M}}_{A'}^\beta \cap F_{p+(n-d+1)-i}^H {}^* \check{\mathcal{M}}_{A'}^\beta) = \sum_{i \geq 0} \partial_t^i (V_{ind}^0 \check{\mathcal{M}}_{A'}^\beta \cap F_{p-i}^{ord} {}^* \check{\mathcal{M}}_{A'}^\beta)$$

for all $p \geq 0$, where the last equality follows from Proposition 4.8 and Lemma 4.15. Since we have

$$F_p^{ord} \check{\mathcal{M}}_{A'}^\beta \supset \partial_t^i F_{p-i}^{ord} \check{\mathcal{M}}_{A'}^\beta \text{ for } p \geq 0 \text{ and } 0 \leq i \leq p$$

it remains to show

$$F_{p-i}^{ord} \check{\mathcal{M}}_{A'}^\beta \supset V_{ind}^0 \check{\mathcal{M}}_{A'}^\beta \cap F_{p-i}^{ord} {}^* \check{\mathcal{M}}_{A'}^\beta \text{ for } p \geq 0 \text{ and } 0 \leq i \leq p$$

resp.

$$F_p^{ord} \check{M}_{A'}^\beta \supset V_{ind}^0 \check{M}_{A'}^\beta \cap F_p^{ord*} \check{M}_{A'}^\beta \text{ for } p \geq 0.$$

Now let $[P] \in V_{ind}^0 \check{M}_{A'}^\beta \cap F_p^{ord*} \check{M}_{A'}^\beta$, then $P \in D_{W \times \mathbb{C}_t}$ can be written as

$$P = t^{-k} P_k + t^{-k+1} P_{k-1} + \dots$$

with $P_i \in \mathbb{C}[w_1, \dots, w_n] \langle \partial_t, \partial_{w_1}, \dots, \partial_{w_n} \rangle$ and $P_k \neq 0$. Since $t^k \cdot [P] \in V_{ind}^k \check{M}_{A'}^\beta \cap F_p^{ord} \check{M}_{A'}^\beta$ it is enough to prove

$$t^k F_p^{ord} \check{M}_{A'}^\beta \supset V_{ind}^k \check{M}_{A'}^\beta \cap F_p^{ord} \check{M}_{A'}^\beta \text{ for } p \geq 0.$$

Given an element $[Q] \in V_{ind}^k \check{M}_{A'}^\beta \cap F_p^{ord} \check{M}_{A'}^\beta$, we can find, using Proposition 4.9, a $Q' \in V^k D_{W \times \mathbb{C}_t} \cap F_p D_{W \times \mathbb{C}_t}$ with $[Q] = [Q']$. But this element Q' can be written as a linear combination of monomials $t^{l_0} w_1^{l_1} \dots w_n^{l_n} \partial_t^{p_0} \partial_{w_1}^{p_1} \dots \partial_{w_n}^{p_n}$ with $p_0 + \dots + p_n \leq p$ and $l_0 - p_0 \geq k$, hence $[Q'] \in t^k F_p^{ord} \check{M}_{A'}^\beta$. This shows the statement in the case $\mathbb{N}A \neq \mathbb{Z}^d$ (recall that $\mathbb{N}A \neq \mathbb{Z}^d$ was a condition used in Proposition 4.9).

In the case $\mathbb{N}A = \mathbb{Z}^d$ the support of $\check{M}_{A'}^\beta$ is disjoint from the divisor $\{t = 0\}$, hence the extension of the Hodge filtration is simply given by $F_p^H \check{M}_{A'}^\beta = j_{t*} F_p^{H*} \check{M}_{A'}^\beta$. Since j_t is an open embedding we have $\check{M}_{A'}^\beta = {}^* \check{M}_{A'}^\beta$ and therefore also $F_p^H \check{M}_{A'}^\beta = F_p^H {}^* \check{M}_{A'}^\beta$. This shows the claim. \square

Now we want to deduce the Hodge filtration on $h_{A+} \mathcal{O}_T^\beta$ from the proposition above.

Theorem 4.17. *Let $\mathbb{N}A = \mathbb{Z}^d \cap \mathbb{R}_{\geq 0} A$ and $\beta \in \mathbb{R}^d \setminus sRes(A)$. The direct image $h_{A+} \mathcal{O}_T^\beta$ is isomorphic to the cyclic D_W -module $\check{\mathcal{M}}_A^\beta := D_W / \check{\mathcal{I}}$, where $\check{\mathcal{I}}$ is the left ideal generated by $(\check{E}_k + \beta_k)_{k=1, \dots, d}$ and $(\check{\square}_m)_{m \in \mathbb{L}_A}$. For $\beta \in \mathfrak{A}_A$ the Hodge filtration on $\check{\mathcal{M}}_A^\beta$ is equal to the order filtration shifted by $n - d$, i.e.*

$$F_{p+(n-d)}^H \check{\mathcal{M}}_A^\beta = F_p^{ord} \check{\mathcal{M}}_A^\beta.$$

Proof. Recall that we have $j_t \circ i_g^\circ \circ k_A = i_g \circ h_A$ where i_f is the graph embedding from (24). The map i_f can be factored by

$$\begin{aligned} i_0 : W &\longrightarrow W \times \mathbb{C}_{\tilde{t}} \\ \underline{w} &\mapsto (\underline{w}, 0) \\ l_g : W \times \mathbb{C}_{\tilde{t}} &\longrightarrow W \times \mathbb{C}_t \\ (\underline{w}, \tilde{t}) &\mapsto (\underline{w}, \tilde{t} + g(\underline{w})). \end{aligned}$$

Once again, we summarize the relevant maps in the following diagram.

$$\begin{array}{ccccc} & & W \times \mathbb{C}_{\tilde{t}} & & \\ & & \uparrow \text{closed } i_0 & \searrow \cong & \\ & & W & \xrightarrow{l_g} & W \times \mathbb{C}_t \\ & \swarrow l & \xrightarrow{i_g} & \text{closed} & \\ W^* & \xrightarrow{l} & W & \xrightarrow{i_g} & W \times \mathbb{C}_t \\ & \searrow i_g^\circ & & & \uparrow \text{open embedding} \\ & & & & \text{complement of smooth divisor} \\ & & & & W \times \mathbb{C}_t^* \\ \text{closed } l_1 \downarrow & & & & \uparrow j_t \\ \widetilde{W} & \xrightarrow{l_2} & W^* \times \mathbb{C}_t^* & \xrightarrow{l_3} & W \times \mathbb{C}_t^* \\ & \cong & & \text{closed on support} & \end{array}$$

We first compute $\mathcal{H}^0(l_g^{-1})_+ \check{\mathcal{M}}_{A'}^\beta$, with its corresponding Hodge filtration. Since $(l_g)^{-1}$ is just a coordinate change we get similarly to formula (30)

$$\Gamma(W \times \mathbb{C}_{\tilde{t}}, \mathcal{H}^0(l_g^{-1})_+ \check{\mathcal{M}}_{A'}^\beta) \simeq D_{W \times \mathbb{C}_{\tilde{t}}} / ((\check{E}_k + \beta_k)_{k=1, \dots, d}, (\check{\square}_m)_{m \in \mathbb{L}_A}, (\tilde{t})), \quad (32)$$

where the Hodge filtration on the right hand side is the induced order filtration shifted by $(n - d + 1)$. Notice that the right hand side of (32) is simply $\check{M}_A^\beta[\partial_{\tilde{t}}]$, hence the Hodge filtration on

$$\check{M}_A^\beta = \Gamma(W, \check{\mathcal{M}}_{A'}^\beta) = \Gamma(W \times \mathbb{C}_{\tilde{t}}, \mathcal{H}^0 i_0^+(l_g^{-1})_+ \check{\mathcal{M}}_{A'}^\beta)$$

is simply the order filtration shifted by $(n - d)$ by [Sai88, Proposition 3.2.2 (iii)]. \square

Remark 4.18. Let $\mathcal{O}_{\mathbb{C}^*}^\beta = \mathcal{D}_{\mathbb{C}^*}/(\partial_t t + \beta)$ and $j_0 : \mathbb{C}^* \rightarrow \mathbb{C}$ be the inclusion. If $\beta \notin \{-1, -2, -3, \dots\}$ then

$$j_{0+}\mathcal{O}_{\mathbb{C}^*}^\beta \simeq \mathcal{D}_{\mathbb{C}}/(\partial_t t + \beta)$$

as well as

$$j_{0\dagger}\mathcal{O}_{\mathbb{C}^*}^{-\beta-1} \simeq \mathbb{D}j_{0+}\mathbb{D}\mathcal{O}_{\mathbb{C}^*}^\beta \simeq \mathbb{D}j_{0+}\mathcal{D}_{\mathbb{C}^*}/(\partial_t t + \beta) \simeq \mathbb{D}\mathcal{D}_{\mathbb{C}}/(\partial_t t + \beta) \simeq \mathcal{D}_{\mathbb{C}}/(t\partial_t - \beta)$$

If, additionally, $\beta \in (-1, 0]$ holds then by Theorem 4.17

$$(j_{0+}\mathcal{O}_{\mathbb{C}^*}^\beta, F_{\bullet}^H) \simeq (\mathcal{D}_{\mathbb{C}}/(\partial_t t + \beta), F_{\bullet}^{ord})$$

In order to compute the Hodge filtration on $j_{0\dagger}\mathcal{O}_{\mathbb{C}^*}^{-\beta-1}$ we use that $(\mathbb{D}\mathcal{O}_{\mathbb{C}^*}^{-\beta-1}, F_p^H) = (\mathcal{D}_{\mathbb{C}^*}/(\partial_t t + \beta), F_{\bullet-1}^{ord})$ and therefore $(j_{0\dagger}\mathbb{D}\mathcal{O}_{\mathbb{C}^*}^{-\beta-1}, F_p^H) = (\mathcal{D}_{\mathbb{C}}/(\partial_t t + \beta), F_{\bullet-1}^{ord})$ holds, since we assumed $\beta \in (-1, 0]$. We use the filtered resolution

$$(\mathcal{D}_{\mathbb{C}^*}, F_{\bullet-2}^{ord}) \xrightarrow{\partial_t t + \beta} (\mathcal{D}_{\mathbb{C}^*}, F_{\bullet-1}^{ord})$$

to resolve $(\mathcal{D}_{\mathbb{C}}/(\partial_t t + \beta), F_{\bullet-1}^{ord})$ and apply $\mathcal{H}om(-, (\mathcal{D}_{\mathbb{C}}, F_{\bullet-2}^{ord}) \otimes \omega_X^\vee)$ to compute the dual of $(\mathcal{D}_{\mathbb{C}}/(\partial_t t + \beta), F_{\bullet-1}^{ord})$ (cf. [Sai94, page 55] for the choice of filtration on $\mathcal{D}_{\mathbb{C}} \otimes \omega_X^\vee$). This gives

$$(j_{0\dagger}\mathcal{O}_{\mathbb{C}^*}^{-\beta-1}, F_p^H) \simeq (\mathcal{D}_{\mathbb{C}}/(t\partial_t - \beta), F_p^{ord})$$

for $\beta \in (-1, 0]$.

5 Integral transforms of torus embeddings

In this section we investigate how the Hodge filtration of $\mathbb{C}_T^{H,\beta}$ behaves under a certain integral transformation, which depends on a matrix \tilde{A} and a parameter β_0 . We show that the outcome of this transform is isomorphic to the GKZ-system $\mathcal{M}_{\tilde{A}}^{(\beta_0, \beta)}$ (cf. Proposition 5.4). In the special case $\beta_0 \in \mathbb{Z}$ we will show that the integral transformation mentioned above is isomorphic to the Radon transform of a torus embedding.

The Radon transformation has been extensively used in the previous papers [Rei14] and [RS17] in order to study hyperplane sections of toric varieties, more precisely, fibres of Laurent polynomials and their compactifications.

In the first subsection below, we give a brief reminder of how certain GKZ-systems can be constructed using the Radon transformation. The second subsection introduces an integral transform that is able to produce all GKZ-systems $\mathcal{M}_{\tilde{A}}^{\tilde{\beta}}$ with $\tilde{\beta} \notin sRes(\tilde{A})$ and homogeneous \tilde{A} .

The next subsections (until 5.9) constitute the main part of this section, where we study in detail how the various functors entering in the definition of this integral transformation act on the twisted structure sheaf. One can roughly divide the construction in two parts: In subsection 5.3 we calculate the push-forward of a tensor product between the twisted structure sheaf and a kernel to a partial compactification. Then one has to study the projection to the parameter space (i.e., the space on which the GKZ-system is defined). The calculation of the behavior of the filtration steps is non-trivial, as the higher direct images of these filtration steps, being coherent \mathcal{O} -modules, do not, a priori, vanish. However, we can show that this is actually the case in the current situation. We formulate this result in the language of \mathcal{R} -modules (i.e., using the Rees construction for filtered \mathcal{D} -modules), and make extensive use of (variants of) the Euler-Koszul complex of hypergeometric modules. All these intermediate steps are contained in the subsections 5.4 until 5.9. A very important technical result is the calculation of some local cohomology groups of a certain semi-group ring, contained in subsection 5.8. The culminating point is then Theorem 5.35 which gives a precise description of the Hodge filtration on certain GKZ-systems.

5.1 Hypergeometric modules, Gauß-Manin systems and the Radon transformation

Here we give a brief reminder on the relationship between GKZ-hypergeometric systems, Gauß-Manin systems of families of Laurent polynomials as developed in [Rei14].

As in [RS15, RS17], we will consider a homogenization of the above systems. Namely, given the matrix $A = (a_{ki})$, we consider the system $\mathcal{M}_{\tilde{A}}^{\tilde{\beta}}$, where \tilde{A} is the $(d+1) \times (n+1)$ integer matrix

$$\tilde{A} := (\tilde{a}_0, \dots, \tilde{a}_n) := \begin{pmatrix} 1 & 1 & \dots & 1 \\ 0 & a_{11} & \dots & a_{1n} \\ \vdots & \vdots & & \vdots \\ 0 & a_{d1} & \dots & a_{dn} \end{pmatrix} \quad (33)$$

and $\tilde{\beta} \in \mathbb{C}^{d+1}$.

In order to show that such a homogenized GKZ-system comes from geometry we have to review briefly the so-called Radon transformation for \mathcal{D} -modules which was introduced by Brylinski [Bry86] and variants were later added by d'Agnolo and Eastwood [DE03].

Let W be the dual vector space of V with coordinates w_0, \dots, w_n , and let $\lambda_0, \dots, \lambda_n$ be coordinates for V . We will denote by $Z \subset \mathbb{P}(W) \times V$ the universal hyperplane given by $Z := \{\sum_{i=0}^n \lambda_i w_i = 0\}$ and denote its complement by $U := (\mathbb{P}(W) \times V) \setminus Z$. Consider the following diagram

$$\begin{array}{ccccc} & & U & & \\ & \swarrow \pi_1^U & \downarrow j_U & \searrow \pi_2^U & \\ \mathbb{P}(W) & \xleftarrow{\pi_1} & \mathbb{P}(W) \times V & \xrightarrow{\pi_2} & V \\ & \swarrow \pi_1^Z & \downarrow i_Z & \searrow \pi_2^Z & \\ & & Z & & \end{array} \quad (34)$$

We will use in the sequel several variants of the so-called Radon transformation in the derived category of mixed Hodge modules. These are functors from $D^b\text{MHM}(\mathbb{P}(W))$ to $D^b\text{MHM}(\mathcal{D}_V)$ given by

$$\begin{aligned} {}^*\mathcal{R}(M) &:= \pi_{2*}^Z(\pi_1^Z)^*M \simeq \pi_{2*}i_{Z*}i_Z^*\pi_1^*M, \\ {}^!\mathcal{R}(M) &:= \mathbb{D} \circ {}^*\mathcal{R} \circ \mathbb{D}(M) \simeq \pi_{2*}^Z(\pi_1^Z)^!M \simeq \pi_{2*}i_{Z*}i_Z^!\pi_1^!M \\ {}^*\mathcal{R}_{cst}(M) &:= \pi_{2*}\pi_1^*M. \\ {}^!\mathcal{R}_{cst}(M) &:= \mathbb{D} \circ {}^*\mathcal{R}_{cst} \circ \mathbb{D}(M) \simeq \pi_{2*}\pi_1^!M. \\ {}^*\mathcal{R}_c^\circ(M) &:= \pi_{2!}^U(\pi_1^U)^*(M) \simeq \pi_{2*}j_{U!}j_U^*\pi_1^*(M) \\ {}^!\mathcal{R}^\circ(M) &:= \mathbb{D} \circ {}^*\mathcal{R}_c^\circ \circ \mathbb{D}(M) \simeq \pi_{2*}^U(\pi_1^U)^!(M) \simeq \pi_{2*}j_{U*}j_U^!\pi_1^!(M). \end{aligned}$$

The adjunction triangle corresponding to the open embedding j_U and the closed embedding i_Z gives rise to the following triangles of Radon transformations

$${}^!\mathcal{R}(M) \longrightarrow {}^!\mathcal{R}_{cst}(M) \longrightarrow {}^!\mathcal{R}^\circ(M) \xrightarrow{+1}, \quad (35)$$

$${}^*\mathcal{R}_c^\circ(M) \longrightarrow {}^*\mathcal{R}_{cst}(M) \longrightarrow {}^*\mathcal{R}(M) \xrightarrow{+1}, \quad (36)$$

where the second triangle is dual to the first.

We now introduce a family of Laurent polynomials defined on $T \times \Lambda := (\mathbb{C}^*)^d \times \mathbb{C}^n$ using the columns of the matrix A , more precisely, we put

$$\begin{aligned} \varphi_A : T \times \Lambda &\longrightarrow V = \mathbb{C}_{\lambda_0} \times \Lambda, \\ (t_1, \dots, t_d, \lambda_1, \dots, \lambda_n) &\mapsto \left(-\sum_{i=1}^n \lambda_i t_i^{a_i}, \lambda_1, \dots, \lambda_n\right). \end{aligned} \quad (37)$$

The following theorem of [Rei14] constructs a morphism between the Gauß-Manin system $\mathcal{H}^0(\varphi_{A,+}\mathcal{O}_{T \times \Lambda})$ resp. its proper version $\mathcal{H}^0(\varphi_{A,\dagger}\mathcal{O}_{T \times \Lambda})$ and certain GKZ-hypergeometric systems and identify both with a corresponding Radon transform.

For this we apply the triangle (35) to $M = g_! \mathbb{D}^p \mathbb{Q}_T^H$ and the triangle (36) to $M = g_*^p \mathbb{Q}_T^H$, where the map g was defined by

$$g : T \longrightarrow \mathbb{P}(W)$$

$$(t_1, \dots, t_d) \mapsto (1 : \underline{t}^{a_1} : \dots : \underline{t}^{a_n}).$$

Theorem 5.1. [Rei14, Lemma 1.11, Proposition 3.4] Let $A = (\underline{a}_1, \dots, \underline{a}_n) \in M(d \times n, \mathbb{Z})$ and $\tilde{A} = (\underline{a}_0, \underline{a}_1, \dots, \underline{a}_n) \in M((d+1) \times (n+1), \mathbb{Z})$ be as above and assume that \tilde{A} satisfies

1. $\mathbb{Z}\tilde{A} = \mathbb{Z}^{d+1}$
2. $\mathbb{N}\tilde{A} = \mathbb{R}_{\geq 0}\tilde{A} \cap \mathbb{Z}^{d+1}$

Then for every $\tilde{\beta} \in \mathbb{N}\tilde{A}$ and every $\tilde{\beta}' \in \text{int}(\mathbb{N}\tilde{A})$, we have that

$$\mathcal{M}_{\tilde{A}}^{\tilde{\beta}} \simeq \text{DMod}(\mathcal{H}^{n+1}(*\mathcal{R}_c^\circ(g_*^p \mathbb{Q}_T^H))) \quad \text{and} \quad \mathcal{M}_{\tilde{A}}^{-\tilde{\beta}'} \simeq \text{DMod}(\mathcal{H}^{-n-1}({}^! \mathcal{R}^\circ(g_! \mathbb{D}^p \mathbb{Q}_T^H)))$$

If we define

$${}^H \mathcal{M}_{\tilde{A}}^{\tilde{\beta}} := \mathcal{H}^{n+1}(*\mathcal{R}_c^\circ(g_*^p \mathbb{Q}_T^H)) \quad \text{and} \quad {}^H \mathcal{M}_{\tilde{A}}^{-\tilde{\beta}'} := \mathcal{H}^{-n-1}({}^! \mathcal{R}^\circ(g_! \mathbb{D}^p \mathbb{Q}_T^H))$$

the following sequences of mixed Hodge modules are exact and dual to each other:

$$\begin{array}{ccccccc}
H^{d-1}(T, \mathbb{C}) \otimes {}^p \mathbb{Q}_V^H & \mathcal{H}^0(\varphi_{A*} {}^p \mathbb{Q}_{T \times \Lambda}^H) & {}^H \mathcal{M}_{\tilde{A}}^{\tilde{\beta}} & H^d(T, \mathbb{C}) \otimes {}^p \mathbb{Q}_V^H \\
\uparrow \simeq & \uparrow \simeq & \uparrow \simeq & \uparrow \simeq \\
0 \longrightarrow \mathcal{H}^n(*\mathcal{R}_{cst}(g_*^p \mathbb{Q}_T^H)) \longrightarrow \mathcal{H}^n(*\mathcal{R}(g_*^p \mathbb{Q}_T^H)) \longrightarrow \mathcal{H}^{n+1}(*\mathcal{R}_c^\circ(g_*^p \mathbb{Q}_T^H)) \longrightarrow \mathcal{H}^{n+1}(*\mathcal{R}_{cst}(g_*^p \mathbb{Q}_T^H)) \longrightarrow 0 \\
\\
0 \leftarrow \mathcal{H}^{-n}({}^! \mathcal{R}_{cst}(g_! \mathbb{D}^p \mathbb{Q}_T^H)) \leftarrow \mathcal{H}^{-n}({}^! \mathcal{R}(g_! \mathbb{D}^p \mathbb{Q}_T^H)) \leftarrow \mathcal{H}^{-n-1}({}^! \mathcal{R}^\circ(g_! \mathbb{D}^p \mathbb{Q}_T^H)) \leftarrow \mathcal{H}^{-n-1}({}^! \mathcal{R}_{cst}(g_! \mathbb{D}^p \mathbb{Q}_T^H)) \leftarrow 0. \\
\downarrow \simeq & \downarrow \simeq & \downarrow \simeq & \downarrow \simeq \\
H_{d+1}(T, \mathbb{C}) \otimes \mathbb{D}^p \mathbb{Q}_V^H & \mathcal{H}^0(\varphi_{A!} \mathbb{D}^p \mathbb{Q}_{T \times \Lambda}^H) & {}^H \mathcal{M}_{\tilde{A}}^{-\tilde{\beta}'} & H_d(T, \mathbb{C}) \otimes \mathbb{D}^p \mathbb{Q}_V^H
\end{array}$$

Proposition 5.2. Let $\tilde{\beta} \in \mathbb{N}\tilde{A}$ and $\tilde{\beta}' \in \text{int}(\mathbb{N}\tilde{A})$. There exists a natural morphism of mixed Hodge modules between ${}^H \mathcal{M}_{\tilde{A}}^{-\tilde{\beta}'}$ and ${}^H \mathcal{M}_{\tilde{A}}^{\tilde{\beta}}$, which is (up to multiplication with a non-zero constant) given on the underlying \mathcal{D}_V -modules by

$$\begin{aligned}
\mathcal{M}_{\tilde{A}}^{-\tilde{\beta}'} &\longrightarrow \mathcal{M}_{\tilde{A}}^{\tilde{\beta}} \\
P &\mapsto P \cdot \partial^{\tilde{\beta} + \tilde{\beta}'},
\end{aligned}$$

where $\partial^{\tilde{\beta} + \tilde{\beta}'}$:= $\prod_{i=0}^n \partial_{\lambda_i}^{k_i}$ for any $\underline{k} = (k_0, \dots, k_n)$ with $\tilde{A} \cdot \underline{k} = \tilde{\beta} + \tilde{\beta}'$.

Proof. First notice that there is a natural morphism of mixed Hodge modules

$$\mathcal{H}^0(\varphi_{A!} \mathbb{D}^p \mathbb{Q}_{T \times \Lambda}^H)(-d-n) \longrightarrow \mathcal{H}^0(\varphi_{A*} {}^p \mathbb{Q}_{T \times \Lambda}^H)$$

which is induced by the morphism $\mathbb{D}^p \mathbb{Q}_{T \times \Lambda}^H(-d-n) \rightarrow {}^p \mathbb{Q}_{T \times \Lambda}^H$. Using the isomorphisms in the second column, this gives a morphism

$$\mathcal{H}^{-n}({}^! \mathcal{R}(g_! \mathbb{D}^p \mathbb{Q}_T^H))(-n-d) \longrightarrow \mathcal{H}^n(*\mathcal{R}(g_*^p \mathbb{Q}_T^H)).$$

Now we can concatenate this with the following morphisms

$$\begin{array}{ccc} \mathcal{H}^n(*\mathcal{R}(g_*^p \mathbb{Q}_T^H)) & \longrightarrow & \mathcal{H}^{n+1}(*\mathcal{R}_c^\circ(g_*^p \mathbb{Q}_T^H)) \\ \uparrow & & \uparrow \text{---} \\ \mathcal{H}^{-n}({}^l\mathcal{R}(g_! \mathbb{D}^p \mathbb{Q}_T^H))(-n-d) & \longleftarrow & \mathcal{H}^{-n-1}({}^l\mathcal{R}^\circ(g_! \mathbb{D}^p \mathbb{Q}_T^H))(-n-d). \end{array}$$

This gives the desired morphism of mixed Hodge modules between ${}^H\mathcal{M}_{\tilde{A}}^{-\tilde{\beta}'}$ ($-d-n$) and ${}^H\mathcal{M}_{\tilde{A}}^{\tilde{\beta}}$. Then it follows from [RS17, Lemma 2.12] that the corresponding morphism of the underlying \mathcal{D}_V -modules is (up to multiplication by a non-zero constant) right multiplication with $\partial^{\tilde{\beta}+\tilde{\beta}'}$. \square

We will now prove a partial generalization of Theorem 5.1 for non-integer β .

Proposition 5.3. *With the notation as above, let $\tilde{\beta} = (\beta_0, \beta) \in (\mathbb{Z} \times \mathbb{R}^d) \setminus \text{sRes}(\tilde{A})$, then we have the following isomorphism*

$$\text{DMod}(\mathcal{H}^{n+1}(*\mathcal{R}_c^\circ(g_*^p \mathbb{C}_T^{\beta, H}))) \simeq \mathcal{M}_{\tilde{A}}^{\tilde{\beta}}$$

This induces the structure of a complex mixed Hodge module on $\mathcal{M}_{\tilde{A}}^{\tilde{\beta}}$ which we call ${}^H\mathcal{M}_{\tilde{A}}^{\tilde{\beta}}$.

Proof. Consider the following commutative diagram with cartesian square

$$\begin{array}{ccc} & & W \\ & \nearrow h_{\tilde{A}} & \uparrow \bar{j} \\ \mathbb{C}^* \times T & \xrightarrow{\tilde{h}} & W \setminus \{0\} \\ \pi_T \downarrow & & \downarrow \pi \\ T & \xrightarrow{g} & \mathbb{P}(W) \end{array}$$

where π_T is the projection to the first factor. We have

$$\bar{j}_+ \pi^+ g_+ \mathcal{O}_T^\beta \simeq \bar{j}_+ \tilde{h}_+ \pi_T^+ \mathcal{O}_T^\beta \simeq h_{\tilde{A}^+} \mathcal{O}_{\mathbb{C}^* \times T}^{(0, \beta)}[1] \simeq h_+ \mathcal{O}_{\mathbb{C}^* \times T}^{(\beta_0, \beta)}[1]$$

for every $\beta_0 \in \mathbb{Z}$. Let $*\mathcal{R}_c^\circ : D_{rh}^b(\mathcal{D}_X) \rightarrow D_{rh}^b(\mathcal{D}_X)$ be the corresponding functor for \mathcal{D} -modules which is given by $M \mapsto \pi_{2+} j_U \dagger j_U^\dagger \pi_1^\dagger M$. We have the following isomorphism

$$*\mathcal{R}_c^\circ(g_+ \mathcal{O}_T^\beta)[-n-1] \simeq \text{FL}(\bar{j}_+ \pi^+ g_+ \mathcal{O}_T^\beta[-1]) \simeq \text{FL}(h_{\tilde{A}^+} \mathcal{O}_{\mathbb{C}^* \times T}^{(\beta_0, \beta)}) \simeq \mathcal{M}_{\tilde{A}}^{\tilde{\beta}}$$

where the first isomorphism follows from [DE03, Proposition 1]. Notice that the various shifts, occurring in the formulas above, stem from a different (shifted) definition of the (exceptional) inverse image for \mathcal{D} -modules in loc. cit. \square

5.2 Integral transforms of twisted structure sheaves

Unfortunately, the Radon transformation produces only GKZ-systems with $\beta_0 \in \mathbb{Z}$, as we can see in Proposition 5.3. To remedy this fact, we introduce a integral transformation which takes care of that by twisting with a kernel which depends on β_0 .

Let $T = (\mathbb{C}^*)^d$ resp. $\tilde{T} := (\mathbb{C}^*)^{d+1}$ be tori with coordinates t_1, \dots, t_d resp. t_0, \dots, t_d , $W = V = \mathbb{C}^{n+1}$ with coordinates w_0, \dots, w_n resp. $\lambda_0, \dots, \lambda_n$ and consider the torus embedding with respect to the matrix \tilde{A}

$$\begin{aligned} h &:= h_{\tilde{A}} : \tilde{T} \longrightarrow W \\ (t_0, \dots, t_d) &\mapsto (t_0, t_0 \underline{t}^{\underline{a}_1}, \dots, t_0 \underline{t}^{\underline{a}_n}) \end{aligned}$$

If $\tilde{\beta} \notin sRes(\tilde{A})$ the GKZ-system $\mathcal{M}_{\tilde{A}}^{\tilde{\beta}}$ is given by $FL(h_+ \mathcal{O}_{\tilde{T}}^{\tilde{\beta}})$. Consider the maps

$$\begin{array}{ccccc} \mathbb{C}^* & \xrightarrow{j} & \mathbb{C} & \xleftarrow{F} & T \times V & \xrightarrow{p} & V \\ & & & & \downarrow q & & \\ & & & & T & & \end{array}$$

where F is given by $(t_1, \dots, t_d, \lambda_0, \dots, \lambda_n) \mapsto \lambda_0 + \sum_{i=1}^n \lambda_i t_i^{a_i}$, where p_1 resp. p_2 is the projection to the first resp. second factor and where j is the inclusion.

Proposition 5.3 showed that a GKZ-system with integer β_0 can be expressed by a Radon transformation, generalizing a result in [Rei14]. The Radon transformation ${}^* \mathcal{R}_c^\circ$ can be seen as an integral transform from \mathbb{P}^n to \mathbb{C}^{n+1} with kernel $j_{U!} {}^p \mathcal{C}_T^H$. Now, in order to construct GKZ-systems with general β_0 we could twist the kernel $j_{U!} {}^p \mathcal{C}_U^H$ which means instead of using the constant module ${}^p \mathcal{C}_U^H$ on U we could use a rank one local system on U with monodormy $e^{2\pi i \beta_0}$ (notice that U has fundamental group isomorphic to \mathbb{Z}). However, due to computational reason we use a slightly different approach. Instead of embedding the torus T in \mathbb{P}^n and considering an integral transform from \mathbb{P}^n to \mathbb{C}^{n+1} we directly define an integral transformation from T to V with a kernel depending on β_0 and A (the matrix A is encoded in the map F). We prove in Proposition 5.4 that the outcome of this integral transformation applied to the \mathcal{D}_T -module \mathcal{O}_T^β is indeed the GKZ-system $\mathcal{M}_{\tilde{A}}^{\tilde{\beta}}$. Finally, we prove in Proposition 5.5 that this approach is compatible with the original approach using the Radon transform from Proposition 5.3.

The following proposition is a variant of a theorem of d'Agnolo and Eastwood [DE03]. It compares the Fourier-Laplace transform of the twisted structure sheaf under a torus embedding with an integral transform of the twisted structure on a smaller torus. The latter description is favorable since it naturally equips the GKZ-system with the structure of a mixed Hodge module.

Proposition 5.4. *Let $\tilde{\beta} = (\beta_0, \beta) \notin sRes(\tilde{A})$ then*

$$\mathcal{M}_{\tilde{A}}^{\tilde{\beta}} \simeq FL(h_+ \mathcal{O}_{\tilde{T}}^{\tilde{\beta}}) \simeq \mathcal{H}^{2n+d+1}(p_+(q^\dagger \mathcal{O}_T^\beta \otimes_{\mathcal{O}} F^\dagger(j_! \mathcal{O}_{\mathbb{C}^*}^{-\beta_0-1})))$$

Proof. Notice that the morphism $h : \tilde{T} \rightarrow \mathbb{C}^{n+1}$ factors as

$$\tilde{T} \xrightarrow{\tilde{j}} \mathbb{C} \times T \xrightarrow{k} W \tag{38}$$

where \tilde{j} is the canonical embedding and k is given by $(t_0, \dots, t_d) \mapsto (t_0, t_0 t_1^{a_1}, \dots, t_0 t_n^{a_n})$. Consider the diagram

$$\begin{array}{ccccc} \mathbb{C} & \xleftarrow{l} & \mathbb{C} \times T & \xrightarrow{k} & W \\ \uparrow p_1 & & \uparrow p_{12} & & \uparrow q_1 \\ \mathbb{C} \times \mathbb{C} & \xleftarrow{id_{\mathbb{C}} \times F} & \mathbb{C} \times T \times V & \xrightarrow{k \times id_{\mathbb{C}^{n+1}}} & W \times V \\ \downarrow p_2 & & \downarrow p_{13} & & \downarrow q_2 \\ \mathbb{C} & \xleftarrow{F} & T \times V & \xrightarrow{p} & V \end{array} \quad \begin{array}{ccc} \mathbb{C} \times T & & \\ \uparrow p_{12} & \searrow f & \\ \mathbb{C} \times T \times V & \xrightarrow{g} & T \\ \downarrow p_{13} & \nearrow q & \\ T \times V & & \end{array}$$

where p_{ij} are the projections to the factors i and j , the maps l, q, p_1, q_1 are the projections to the first and the maps f, g, p, p_2, q_2 , are the projection to the second factor.

We have that $\tilde{j}_+ \mathcal{O}_{\tilde{T}}^{\tilde{\beta}} \simeq j_+ \mathcal{O}_{\mathbb{C}^*}^{\beta_0} \boxtimes \mathcal{O}_T^{\beta} \simeq l^+ j_+ \mathcal{O}_{\mathbb{C}^*}^{\beta_0}[-d] \otimes f^+ \mathcal{O}_T^{\beta}[-1]$ hence we get the following isomorphisms

$$\begin{aligned}
& \mathrm{FL}(h_+ \mathcal{O}_{\tilde{T}}^{\tilde{\beta}}) \\
& \simeq \mathrm{FL}(k_+ \tilde{j}_+ \mathcal{O}_{\tilde{T}}^{\tilde{\beta}}), && \text{factorization of } h \text{ (38)} \\
& \simeq q_{2,+}(q_1^+ k_+ \tilde{j}_+ \mathcal{O}_{\tilde{T}}^{\tilde{\beta}} \otimes \mathcal{L})[-n-1], \\
& \simeq q_{2,+}(q_1^+ k_+(l^+ j_+ \mathcal{O}_{\mathbb{C}^*}^{\beta_0} \otimes f^+ \mathcal{O}_T^{\beta}) \otimes \mathcal{L})[-n-d-2], && \text{use } \mathcal{O}_{\tilde{T}}^{\tilde{\beta}} \simeq \mathcal{O}_{\mathbb{C}^*}^{\beta_0} \boxtimes \mathcal{O}_T^{\beta} \\
& \simeq q_{2,+}((k \times id)_+ p_{12}^+(l^+ j_+ \mathcal{O}_{\mathbb{C}^*}^{\beta_0} \otimes f^+ \mathcal{O}_T^{\beta}) \otimes \mathcal{L})[-n-d-2], && \text{base change} \\
& \simeq q_{2,+}(k \times id)_+(p_{12}^+(l^+ j_+ \mathcal{O}_{\mathbb{C}^*}^{\beta_0} \otimes f^+ \mathcal{O}_T^{\beta}) \otimes (k \times id)^+ \mathcal{L})[-2d-2], && \text{projection formula} \\
& \simeq q_{2,+}(k \times id)_+((id \times F)^+ p_1^+ j_+ \mathcal{O}_{\mathbb{C}^*}^{\beta_0} \otimes g^+ \mathcal{O}_T^{\beta} \otimes (k \times id)^+ \mathcal{L})[-n-2d-3], && g = f \circ p_{12}, \\
& && l \circ p_{12} = p_1 \circ (id \times F) \\
& && p \circ p_{13} = q_2 \circ (k \times id) \\
& \simeq p_+ p_{13,+}((id \times F)^+ p_1^+ j_+ \mathcal{O}_{\mathbb{C}^*}^{\beta_0} \otimes g^+ \mathcal{O}_T^{\beta} \otimes (k \times id)^+ \mathcal{L})[-n-2d-3], \\
& \simeq p_+ p_{13,+}((id \times F)^+ p_1^+ j_+ \mathcal{O}_{\mathbb{C}^*}^{\beta_0} \otimes g^+ \mathcal{O}_T^{\beta} \otimes (id \times F)^+ \mathcal{L}_1)[-3n-2d-3], \\
& \simeq p_+ p_{13,+}(g^+ \mathcal{O}_T^{\beta} \otimes (id \times F)^+(p_1^+ j_+ \mathcal{O}_{\mathbb{C}^*}^{\beta_0} \otimes \mathcal{L}_1))[-2n-d-3], \\
& \simeq p_+ p_{13,+}(p_{13}^+ q^+ \mathcal{O}_T^{\beta} \otimes (id \times F)^+(p_1^+ j_+ \mathcal{O}_{\mathbb{C}^*}^{\beta_0} \otimes \mathcal{L}_1))[-2n-d-3], && g = q \circ p_{13} \\
& \simeq p_+(q^+ \mathcal{O}_T^{\beta} \otimes p_{13,+}(id \times F)^+(p_1^+ j_+ \mathcal{O}_{\mathbb{C}^*}^{\beta_0} \otimes \mathcal{L}_1))[-2n-d-2], && \text{projection formula} \\
& \simeq p_+(q^+ \mathcal{O}_T^{\beta} \otimes F^+ p_{2,+}(p_1^+ j_+ \mathcal{O}_{\mathbb{C}^*}^{\beta_0} \otimes \mathcal{L}_1))[-2n-d-2], && \text{base change} \\
& \simeq p_+(q^+ \mathcal{O}_T^{\beta} \otimes F^+ j_{\dagger} \mathcal{O}_{\mathbb{C}^*}^{-\beta_0-1})[-2n-d-1], && \mathrm{FL}(j_+ \mathcal{O}_{\mathbb{C}^*}^{\beta_0}) \simeq j_{\dagger} \mathcal{O}_{\mathbb{C}^*}^{-\beta_0-1} \\
& \simeq p_+(q^{\dagger} \mathcal{O}_T^{\beta} \otimes F^{\dagger} j_{\dagger} \mathcal{O}_{\mathbb{C}^*}^{-\beta_0-1})[2n+d+1], && q^+ \simeq q^{\dagger}[2n+2], \\
& && F^+ \simeq F^{\dagger}[2n+2d]
\end{aligned}$$

□

The isomorphism $\mathcal{M}_{\tilde{A}}^{\tilde{\beta}} \simeq \mathcal{H}^{2n+d+1}(p_+(q^{\dagger} \mathcal{O}_T^{\beta} \otimes F^{\dagger} j_{\dagger} \mathcal{O}_{\mathbb{C}^*}^{-\beta_0-1}))$ which holds for $\tilde{\beta} \in \mathbb{R}^{d+1} \setminus sRes(\tilde{A})$ endows the GKZ-system with the structure of a complex mixed Hodge module. We define the mixed Hodge module structure by

$$H\mathcal{M}_{\tilde{A}}^{\tilde{\beta}} := \mathcal{H}^{2n+d+1}(p_*(q^* p \mathcal{C}_T^{\beta} \otimes F^* j^* p \mathcal{C}_{\mathbb{C}^*}^{-\beta_0-1})) \quad (39)$$

We now check that the Hodge module structure on $\mathcal{M}_{\tilde{A}}^{\tilde{\beta}}$ induced by the definition (39) coincides with the one of Proposition 5.3 in the case $\beta_0 \in \mathbb{Z}$.

Proposition 5.5. *If $\beta_0 \in \mathbb{Z}$ and $(\beta_0, \beta) \notin sRes(\tilde{A})$ then there is an isomorphism*

$${}^* \mathcal{R}_c^{\circ}(g_* p \mathcal{C}_T^{H,\beta})[n+1] \simeq p_*(q^* p \mathcal{C}_T^{\beta} \otimes F^* j^* p \mathcal{C}_{\mathbb{C}^*}^{H,-\beta_0-1})[2n+d+1]$$

Proof. Consider the following commutative diagram whose squares are cartesian

$$\begin{array}{ccccc}
\mathbb{C}^* & \xleftarrow{\tilde{F}} & U_0 & \xrightarrow{\quad} & U \\
\downarrow j & & \downarrow k_U & & \downarrow j_U \\
\mathbb{C} & \xleftarrow{\tilde{F}} & W_0 \times V & \xrightarrow{j_0 \times id} & \mathbb{P}(W) \times V \xrightarrow{\pi_2} V \\
\uparrow F & \nearrow g_0 \times id & \downarrow \tilde{q} & & \downarrow \pi_1 \\
T \times V & & W_0 & \xrightarrow{j_0} & \mathbb{P}(W) \\
\downarrow q & \nearrow g_0 & & \nearrow g & \\
T & & & &
\end{array}$$

where $W_0 := \mathbb{P}(W) \setminus \{w_0 \neq 0\} = \mathbb{C}^n$ with coordinates w_1, \dots, w_n and $U_0 = U \cap W_0$. We denote by p, p_0, π_1 the projections to the first factor and by \tilde{F} the map $(w_1, \dots, w_n, \lambda_0, \lambda_n) \mapsto \lambda_0 + \sum_{i=1}^n \lambda_i w_i$. We consider the coordinate change ϕ_0 defined by

$$\tilde{t}_k = t_k \text{ for } k = 1, \dots, d, \quad \tilde{\lambda}_0 = \lambda_0 + \sum_{i=1}^n \lambda_i \underline{t}^{a_i} = F \quad \text{and} \quad \tilde{\lambda}_i = \lambda_i \text{ for } i = 1, \dots, n$$

on $T \times V$ and the coordinate change ψ_0 defined by

$$\tilde{w}_j = w_j \text{ for } j = 1, \dots, n, \quad \tilde{\lambda}_0 = \lambda_0 + \sum_{i=1}^n \lambda_i w_j = \tilde{F} \quad \text{and} \quad \tilde{\lambda}_i = \lambda_i \text{ for } i = 1, \dots, n$$

on $W_0 \times V$. Notice that with respect to these coordinates the maps F and \tilde{F} are given by the coordinate function $\tilde{\lambda}_0$. Let $pr : V \rightarrow \mathbb{C}$ be the projection $(\tilde{\lambda}_0, \dots, \tilde{\lambda}_n) \mapsto \tilde{\lambda}_0$. We also have $\psi_0 \circ (g_0 \times id) \circ \varphi_0^{-1} = g_0 \times id$ and the map q factors as $\pi_2 \circ j_0 \circ (g \times id)$. Hence we get

$$\begin{aligned} & p_*(q^* {}^p \mathbf{C}_T^{H,\beta} \otimes F^* j_! {}^p \mathbf{C}_{\mathbb{C}^*}^{H,-\beta_0-1})[2n+d+1] \\ \simeq & p_*(q^* {}^p \mathbf{C}_T^{H,\beta} \otimes F^* j_! {}^p \mathbf{C}_{\mathbb{C}^*}^H)[2n+d+1] && \text{use } \beta_0 \in \mathbb{Z} \\ \simeq & p_*({}^p \mathbf{C}_T^{H,\beta} \boxtimes pr^* j_! {}^p \mathbf{C}_{\mathbb{C}^*}^H)[n+1] && F \text{ is a projection after coordinate change} \\ \simeq & (\pi_2 \circ (j_0 \times id) \circ (g_0 \times id))_* ({}^p \mathbf{C}_T^{H,\beta} \boxtimes pr^* j_! {}^p \mathbf{C}_{\mathbb{C}^*}^H)[n+1] && p = \pi_2 \circ (j_0 \times id) \circ (g_0 \times id) \\ \simeq & (\pi_2 \circ (j_0 \times id))_* (g_{0*} {}^p \mathbf{C}_T^{H,\beta} \boxtimes pr^* j_! {}^p \mathbf{C}_{\mathbb{C}^*}^H)[n+1] \\ \simeq & (\pi_2 \circ (j_0 \times id))_* (\tilde{q}^* g_{0*} {}^p \mathbf{C}_T^\beta \otimes \tilde{F}^* j_! {}^p \mathbf{C}_{\mathbb{C}^*}^H)[n+1] && \tilde{F} \text{ is a projection after coordinate change} \\ \simeq & (\pi_2 \circ (j_0 \times id))_* (\tilde{q}^* g_{0*} {}^p \mathbf{C}_T^{H,\beta} \otimes k_{U!} \dot{F}^* {}^p \mathbf{C}_{\mathbb{C}^*}^H)[n+1] && \text{base change, } \dot{F} \text{ and } \tilde{F} \text{ smooth} \\ \simeq & (\pi_2 \circ (j_0 \times id))_* (\tilde{q}^* g_{0*} {}^p \mathbf{C}_T^{H,\beta} \otimes k_{U!} {}^p \mathbf{C}_{U_0}^H)[n+1] \\ \simeq & (\pi_2 \circ (j_0 \times id))_* ((j_0 \times id)^! \pi_1^* g_* {}^p \mathbf{C}_T^{H,\beta} \otimes k_{U!} {}^p \mathbf{C}_{U_0}^H)[n+1] && g = j_0 \circ g_0 \\ \simeq & \pi_{2*} (\pi_1^* g_* {}^p \mathbf{C}_T^{H,\beta} \otimes (j_0 \times id)_* k_{U!} {}^p \mathbf{C}_{U_0}^H)[n+1] && \text{projection formula} \\ \stackrel{(*)}{\simeq} & \pi_{2*} (\pi_1^* g_* {}^p \mathbf{C}_T^{H,\beta} \otimes j_{U!} {}^p \mathbf{C}_U^H)[n+1] \\ \simeq & \pi_{2*} j_{U!} j_U^* \pi_1^* g_* {}^p \mathbf{C}_T^{H,\beta} [n+1] && \text{projection formula} \\ \simeq & {}^* \mathcal{R}_c^\circ (g_* {}^p \mathbf{C}_T^{H,\beta}) [n+1] \end{aligned}$$

where the isomorphism $(*)$ follows from the fact that $\pi_1^* g_* {}^p \mathbf{C}_T^\beta$ is localized along the divisor $\mathbb{P}(W) \setminus W_0$. \square

5.3 Calculation in charts

We saw Subsection 5.2 that the GKZ-system $\mathcal{M}_A^{\tilde{\beta}}$ can be expressed by an integral transformation from T to V with kernel $F^* j_! {}^p \mathbf{C}_{\mathbb{C}^*}^{-\beta_0-1}$. As a first step we compute the Hodge filtration of an intermediate step in this integral transformation, namely $q^* {}^p \mathbf{C}_T^{H,\beta_u} \otimes F_u^* j_! {}^p \mathbf{C}_{\mathbb{C}^*}^{H,-\beta_0-1}[2n+d+1]$ (cf. Lemma 5.7). We do this by giving different presentations of the kernel (cf. Lemma 5.6) and by using several adapted coordinate systems on $T \times V$ indexed by a variable u which goes from 0 to n . The reason for using these adapted coordinate systems (and not just one) is the fact that we can rewrite the intermediate step as a direct product whose factors are easy to compute. Since the projection $p : T \times V \rightarrow V$ is not proper, we have to extend the intermediate step to a partial compactification. Concretely we are using the factorization

$$T \times V \xrightarrow{g \times id} \mathbb{P}(W) \times V \xrightarrow{\pi_2} V$$

of the projection q . Our goal in this section is to compute the underlying \mathcal{D} -module \mathcal{N} of

$${}^H \mathcal{N} := \mathcal{H}^{2n+d+1} (g \times id)_* (q^* {}^p \mathbf{C}_T^{H,\beta} \otimes F^* j_! {}^p \mathbf{C}_{\mathbb{C}^*}^{H,-\beta_0-1}) \quad (40)$$

together with its Hodge filtration on affine charts W_u of $\mathbb{P}(W) \times V$. The different adapted coordinate systems are now used to compute the direct image of the intermediate step under the embedding $T \times V \rightarrow W_u \times V$ which is simply the restriction of ${}^H\mathcal{N}$ to the affine chart W_u . It turns out that the underlying \mathcal{D} -module of this direct image is a direct product of a torus embedding with respect to a matrix A_u and another rather simple module (cf. equation (41)). We use Theorem 4.17 to compute the Hodge filtration on the first factor in Proposition 5.9, the Hodge filtration on the second factor was computed in Remark 4.18 .

We define the map

$$F_u : T \times V \longrightarrow \mathbb{C}$$

$$(t_1, \dots, t_d, \lambda_0, \dots, \lambda_n) \mapsto \lambda_u + \sum_{\substack{i=0 \\ i \neq u}} \lambda_i \underline{t}^{a_i - a_u} = \left(\lambda_0 + \sum_{i=1} \lambda_i \underline{t}^{a_i} \right) \cdot \underline{t}^{-a_u}$$

(notice that $F_0 = F$). We need the following result

Lemma 5.6. *There is an isomorphism*

$$F^* j_!^p \mathbb{C}_{\mathbb{C}^*}^{H, -\beta_0 - 1} \simeq F_u^* j_!^p \mathbb{C}_{\mathbb{C}^*}^{H, -\beta_0 - 1}$$

for $u = 0, \dots, n$.

Proof. For $u \in \{0, \dots, n\}$ and $G := (\mathbb{C}^*)^d$ consider the action

$$\mu_u : G \times T \times V \longrightarrow T \times V$$

$$(g_1, \dots, g_d, t_1, \dots, t_d, \lambda_0, \dots, \lambda_n) \mapsto (t_1, \dots, t_d, \lambda_0 \underline{g}^{-a_u}, \dots, \lambda_n \underline{g}^{-a_u})$$

and the action $G \times \mathbb{C} \rightarrow \mathbb{C}$ given by $(g_1, \dots, g_d, t) \mapsto \underline{g}^{-a_u} \cdot t$. It is easy to see that the map $F : T \times V \rightarrow \mathbb{C}$ is equivariant with respect to this action. Let $i : T \rightarrow G \times T$ the embedding $(t_1, \dots, t_d) \mapsto (t_1, \dots, t_d, t_1, \dots, t_d)$. Since $j_!^p \mathbb{C}_{\mathbb{C}^*}^{H, -\beta_0 - 1}$ is a G -equivariant mixed Hodge module, the module $F^! j_!^p \mathbb{C}_{\mathbb{C}^*}^{H, -\beta_0 - 1}$ is also G -equivariant. Let $p : G \times T \times V \rightarrow T \times V$ the projection. We have isomorphisms

$$F^* j_!^p \mathbb{C}_{\mathbb{C}^*}^{H, -\beta_0 - 1} \simeq i^* p_2^* F^* j_!^p \mathbb{C}_{\mathbb{C}^*}^{H, -\beta_0 - 1} \simeq i^* \mu_u^* F^* j_!^p \mathbb{C}_{\mathbb{C}^*}^{H, -\beta_0 - 1} \simeq F_u^* j_!^p \mathbb{C}_{\mathbb{C}^*}^{H, -\beta_0 - 1}$$

where the second isomorphism follows from the G -equivariance of $F^* j_!^p \mathbb{C}_T^{H, -\beta_0 - 1}$. \square

We define a coordinate change ϕ_u by

$$\tilde{t}_k = t_k, \quad (\tilde{\lambda}_i)_{i \neq u} = (\lambda_i)_{i \neq u} \quad \text{and} \quad \tilde{\lambda}_u = \lambda_u + \sum_{i \neq u} \lambda_i \underline{t}^{a_i - a_u}$$

Denote by $C_u \in GL(d+1, \mathbb{Z})$ the matrix

$$C_u := \begin{pmatrix} 1 & & & & \\ -a_{1u} & 1 & & & \\ \vdots & & \ddots & & \\ -a_{du} & & & & 1 \end{pmatrix}.$$

and define for $\tilde{\beta} = (\beta_0, \beta) \in \mathbb{Z}^{d+1}$:

$$\tilde{\beta}^u := (\beta_0^u, \beta^u) := C_u \cdot \tilde{\beta}$$

Notice that $\tilde{\beta}^0 = \tilde{\beta}$ since $a_0 = 0$.

Lemma 5.7.

1. With respect to the coordinates defined by ϕ_u the complex $q^* p \mathbb{C}_T^{H, \beta^u} \otimes F_u^* j_! p \mathbb{C}_{\mathbb{C}^*}^{H, -\beta_0 - 1} [2n + d + 1]$ is isomorphic to

$$p \mathbb{C}_T^{H, \beta^u} \boxtimes pr_u^* j_! p \mathbb{C}_{\mathbb{C}^*}^{H, -\beta_0 - 1} [n]$$

where $pr_u : V \rightarrow \mathbb{C}$ is the projection $(\tilde{\lambda}_0, \dots, \tilde{\lambda}_n) \mapsto \tilde{\lambda}_u$. In particular we have

$$\mathcal{H}^k \left(q^* p \mathbb{C}_T^{H, \beta^u} \otimes F_u^* j_! p \mathbb{C}_{\mathbb{C}^*}^{H, -\beta_0 - 1} \right) = 0 \quad \text{for } k \neq 2n + d + 1$$

and the underlying \mathcal{D} -module of $\mathcal{H}^{2n+d+1} \left(q^* p \mathbb{C}_T^{H, \beta^u} \otimes F_u^* j_! p \mathbb{C}_{\mathbb{C}^*}^{H, -\beta_0 - 1} \right)$ is given by the exterior product

$$\mathcal{D}_T / \left((\partial_{\tilde{t}_k} \tilde{t}_k + \beta_k^u)_{k=1, \dots, d} \right) \boxtimes \mathcal{D}_V / \left((\partial_{\tilde{\lambda}_i})_{i \neq u}, \tilde{\lambda}_u \partial_{\tilde{\lambda}_u} - \beta_0 \right).$$

2. For $\alpha \in \mathbb{Z}^d$ and $u_1, u_2 \in \{0, \dots, n\}$ the map

$$\mathcal{H}^{2n+d+1} \left(q^* p \mathbb{C}_T^{H, \beta^{u_1}} \otimes F_{u_1}^* j_! p \mathbb{C}_{\mathbb{C}^*}^{H, -\beta_0 - 1} \right) \longrightarrow \mathcal{H}^{2n+d+1} \left(q^* p \mathbb{C}_T^{H, \beta^{u_2} + \alpha} \otimes F_{u_2}^* j_! p \mathbb{C}_{\mathbb{C}^*}^{H, -\beta_0 - 1} \right),$$

given by right multiplication with \underline{t}^α at the level of $\mathcal{D}_{V \times T}$ -modules, is an isomorphism.

Proof. Notice that the map F_u is just the projection $((\tilde{t}_k)_{k=1, \dots, d}, (\tilde{\lambda}_i)_{i \neq u}, \tilde{\lambda}_u) \mapsto \tilde{\lambda}_u$ with respect to the new coordinates. This gives

$$q^* p \mathbb{C}_T^{H, \beta^u} \otimes F_u^* j_! p \mathbb{C}_{\mathbb{C}^*}^{H, -\beta_0 - 1} [2n + d + 1] \simeq p \mathbb{C}_T^{H, \beta^u} \boxtimes pr_u^* j_! p \mathbb{C}_{\mathbb{C}^*}^{H, -\beta_0 - 1} [n]$$

(the shifts can be seen by noticing that $q^*[n+1]$, $F^*[n+d]$ and $pr_u^*[n]$ are exact. The rest is clear. For the second point we define coordinates Let $((\tilde{t}_k)_{k=1, \dots, d}, (\tilde{\lambda}_i)_{i=0, \dots, n})$ and $((\bar{t}_k)_{k=1, \dots, d}, (\bar{\lambda}_i)_{i=0, \dots, n})$ correspond to the maps ϕ_{u_1} and ϕ_{u_2} , respectively. The coordinate change $\phi_{u_2} \circ \phi_{u_1}^{-1}$ is given by

$$\bar{t}_k = \tilde{t}_k, \quad \bar{\lambda}_{u_1} = \tilde{\lambda}_{u_1} - \sum_{i \neq u_1} \tilde{\lambda}_i \underline{t}^{a_i - a_{u_1}}, \quad \bar{\lambda}_{u_2} = \tilde{\lambda}_{u_1} \underline{t}^{a_{u_1} - a_{u_2}}, \quad \bar{\lambda}_i = \tilde{\lambda}_i$$

for $k = 1, \dots, d$ and $i \neq u_1, u_2$. We get the following transformations:

$$\begin{aligned} \partial_{\tilde{\lambda}_i} &\mapsto \partial_{\bar{\lambda}_i} - \underline{t}^{a_i - a_{u_1}} \partial_{\bar{\lambda}_{u_1}} \equiv \partial_{\bar{\lambda}_i} \\ \tilde{\lambda}_{u_1} \partial_{\tilde{\lambda}_{u_1}} - \beta_0 &\mapsto \bar{\lambda}_{u_2} \underline{t}^{a_{u_2} - a_{u_1}} \partial_{\bar{\lambda}_{u_1}} + \bar{\lambda}_{u_2} \partial_{\bar{\lambda}_{u_2}} - \beta_0 \equiv \bar{\lambda}_{u_2} \partial_{\bar{\lambda}_{u_2}} - \beta_0 \\ \partial_{\tilde{\lambda}_{u_2}} &\mapsto - \underline{t}^{a_{u_2} - a_{u_1}} \partial_{\bar{\lambda}_{u_1}} \\ \partial_{\tilde{t}_k} \tilde{t}_k + \beta_k^{u_1} &\mapsto \partial_{\bar{t}_k} \bar{t}_k - \sum_{i \neq u_1} (a_{ki} - a_{ku_1}) \tilde{\lambda}_i \underline{t}^{a_i - a_{u_1}} \partial_{\bar{\lambda}_{u_1}} + (a_{ku_1} - a_{ku_2}) \bar{\lambda}_{u_2} \partial_{\bar{\lambda}_{u_2}} + \beta_k^{u_1} \\ &\equiv \partial_{\bar{t}_k} \bar{t}_k + (a_{ku_1} - a_{ku_2}) \bar{\lambda}_{u_2} \partial_{\bar{\lambda}_{u_2}} + \beta_k^{u_1} \\ &\equiv \partial_{\bar{t}_k} \bar{t}_k + (a_{ku_1} - a_{ku_2}) \beta_0 + \beta_k^{u_1} \\ &= \partial_{\bar{t}_k} \bar{t}_k + \beta_k^{u_2} \end{aligned}$$

where \equiv means equality modulo the ideal generated by the operators on the left hand side. This shows that

$$\mathcal{D}_V / \left((\partial_{\tilde{\lambda}_i})_{i \neq u_1}, \tilde{\lambda}_{u_1} \partial_{\tilde{\lambda}_{u_1}} - \beta_0 \right) \boxtimes \mathcal{D}_T / \left((\partial_{\tilde{t}_k} \tilde{t}_k + \beta_k^{u_1})_{k=1, \dots, d} \right)$$

is actually equal to

$$\mathcal{D}_V / \left((\partial_{\bar{\lambda}_i})_{i \neq u_2}, \bar{\lambda}_{u_2} \partial_{\bar{\lambda}_{u_2}} - \beta_0 \right) \boxtimes \mathcal{D}_T / \left((\partial_{\bar{t}_k} \bar{t}_k + \beta_k^{u_2})_{k=1, \dots, d} \right)$$

after the change of coordinates $\phi_{u_2} \circ \phi_{u_1}^{-1}$. It is then easy to see that the map

$$\mathcal{D}_{V \times T} / \left((\partial_{\tilde{\lambda}_i})_{i \neq u_2}, \tilde{\lambda}_{u_2} \partial_{\tilde{\lambda}_{u_2}} - \beta_0, (\partial_{\tilde{t}_k} \tilde{t}_k + \beta_k^{u_2})_{k=1, \dots, d} \right) \rightarrow \mathcal{D}_V / \left((\partial_{\bar{\lambda}_i})_{i \neq u_2}, \bar{\lambda}_{u_2} \partial_{\bar{\lambda}_{u_2}} - \beta_0, (\partial_{\bar{t}_k} \bar{t}_k + \beta_k^{u_2} + \alpha_k)_{k=1, \dots, d} \right)$$

is given by right multiplication with \underline{t}^α . This shows the second claim. \square

Let $(w_0 : \dots : w_n)$ be the homogeneous coordinates on $\mathbb{P}(W)$ and denote by $j_u : W_u \hookrightarrow \mathbb{P}(W)$ the chart $w_u \neq 0$ with coordinates $w_{iu} := \frac{w_i}{w_u}$ for $i \neq u$. The map g factors over the chart W_u and gives rise to the map

$$g_u : T \longrightarrow W_u \\ (t_1, \dots, t_n) \mapsto (t_1^{a_0 - a_u}, \dots, t_n^{a_n - a_u}).$$

We define the maps

$$\tilde{F}_u : W_u \times V \longrightarrow \mathbb{C} \\ (w_{iu})_{i \neq u} \mapsto \lambda_u + \sum_{\substack{i=0 \\ i \neq u}}^n \lambda_i w_{iu}$$

As mentioned above we would like to compute the restriction of \mathcal{N} to the affine chart $W_u \times V$. For $u = 0, \dots, n$ we set

$$\begin{aligned} {}^H\mathcal{N}_u &:= {}^H\mathcal{N}|_{W_u \times V} \simeq \mathcal{H}^{2n+d+1}(g_u \times id)_*(q^* {}^p\mathbb{C}_T^{H, \beta^u} \otimes F_u^* j_u^! {}^p\mathbb{C}_{\mathbb{C}^*}^{H, -\beta_0 - 1}) \\ &\simeq \mathcal{H}^n(g_u \times id)_* \left({}^p\mathbb{C}_T^{H, \beta^u} \boxtimes pr_u^* j_u^! {}^p\mathbb{C}_{\mathbb{C}^*}^{H, -\beta_0 - 1} \right) \\ &\simeq \mathcal{H}^n \left({}^p g_{u*} \mathbb{C}_T^{H, \beta^u} \boxtimes pr_u^* j_u^! {}^p\mathbb{C}_{\mathbb{C}^*}^{H, -\beta_0 - 1} \right) \\ &\simeq \mathcal{H}^0({}^p g_{u*} \mathbb{C}_T^{H, \beta^u}) \boxtimes \mathcal{H}^n(pr_u^* j_u^! {}^p\mathbb{C}_{\mathbb{C}^*}^{H, -\beta_0 - 1}) \end{aligned} \quad (41)$$

We now apply the main result of section 4 in order to compute the module $g_{u+} \mathcal{O}_T^{\beta^u}$ together with its corresponding Hodge filtration.

Notice that the embedding g_u is given by the $d \times n$ -matrix $A_u = (a_{ki}^u)$ with columns $(\underline{a}_i - \underline{a}_u)$ for $i \in \{0, \dots, n\} \setminus \{u\}$. We need to check whether the matrices A_u satisfy the conditions in Theorem 4.17. Recall that $\tilde{\beta}^u := (\beta_0^u, \beta^u) := C_u \cdot \tilde{\beta}$.

Lemma 5.8. *Assume that $\mathbb{Z}\tilde{A} = \mathbb{Z}^{d+1}$ and $\mathbb{N}\tilde{A} = \mathbb{Z}^{d+1} \cap \mathbb{R}_{\geq 0}\tilde{A}$, then the matrices A_u satisfy the conditions,*

1. $\mathbb{Z}A_u = \mathbb{Z}^d$
2. $\mathbb{N}A_u = \mathbb{Z}^d \cap \mathbb{R}_{\geq 0}A_u$
3. if $\tilde{\beta} \in \mathfrak{A}_{\tilde{A}}$ then $\beta^u \in \mathfrak{A}_{A_u}$

Proof. Denote by \tilde{A}_u the $(d+1) \times (n+1)$ -matrix with columns $(1, \underline{a}_i - \underline{a}_u)$ for $i \in \{0, \dots, n\}$. We will first show the two properties for the matrix \tilde{A}_u . Notice that we have $C_u \cdot \tilde{A} = C_u \cdot \tilde{A}_0 = \tilde{A}_u$. Since C_u is a linear, invertible map we get $C_u(\mathbb{Z}\tilde{A}) = \tilde{A}_u$, $C_u(\mathbb{N}\tilde{A}) = \tilde{A}_u$ and $C_u(\mathbb{R}_{\geq 0}\tilde{A}) = \mathbb{R}_{\geq 0}\tilde{A}_u$. Therefore the two properties hold for \tilde{A}_u if and only if they hold for \tilde{A} .

Denote by $p : \mathbb{Z}^{d+1} \rightarrow \mathbb{Z}^d$ the projection to the last d -coordinates. Since p maps $(1, \underline{a}_i - \underline{a}_u)$ to $\underline{a}_i - \underline{a}_u$ it is easy to see that the first two properties also hold for A_u .

It follows easily from the definition that $\tilde{\beta} \in \mathfrak{A}_{\tilde{A}}$ if and only if $\tilde{\beta}^u \in \mathfrak{A}_{\tilde{A}_u}$. Hence it is enough to show that $\tilde{\beta} = (\beta_0, \beta) \in \mathfrak{A}_{\tilde{A}}$ implies $\beta \in \mathfrak{A}_A$. We notice first that there is a 1-1 correspondence between facets of $\mathbb{R}_{\geq 0}A$ and facets of $\mathbb{R}_{\geq 0}\tilde{A}$ containing $\tilde{a}_0 = (1, 0, \dots, 0)$ given by

$$F \leftrightarrow \tilde{F} = F + \mathbb{R}_{\geq 0} \cdot (1, 0, \dots, 0)$$

If n_F is a primitive, inward-pointing normal vector of a facet F of $\mathbb{R}_{\geq 0}A$, the vector $n_{\tilde{F}} := (0, n_F)$ is a primitive, inward-pointing normal vector of the corresponding facet \tilde{F} of $\mathbb{R}_{\geq 0}\tilde{A}$. Since $\tilde{c} := \sum_{i=0}^n \tilde{a}_i = (n+1, \underline{c})$, we have $e_{\tilde{F}} = \langle n_{\tilde{F}}, \tilde{c} \rangle = \langle (0, n_F), (n+1, \underline{c}) \rangle = \langle n_F, \underline{c} \rangle = e_F$. We get by definition 15

$$\tilde{\beta} \in \mathfrak{A}_{\tilde{A}} = \bigcap_{\tilde{F} \text{ facet}} \{\mathbb{R} \cdot \tilde{F} - [0, e_{\tilde{F}}] \cdot \tilde{c}\} \subset \bigcap_{\substack{\tilde{F} \text{ facet} \\ \tilde{a}_0 \in \tilde{F}}} \{\mathbb{R} \cdot \tilde{F} - [0, e_{\tilde{F}}] \cdot \tilde{c}\} \Rightarrow \beta \in \bigcap_{F \text{ facet}} \{\mathbb{R} \cdot F - [0, e_F] \cdot \underline{c}\} = \mathfrak{A}_A.$$

□

Denote by \mathbb{L}_{A_u} the \mathbb{Z} -module of relations among the columns of A_u . In order to calculate the direct image of \mathcal{O}_T^β under the map g_u , we use Theorem 4.17 where A_u takes the role of the matrix B in loc.cit.

Proposition 5.9. *Consider the \mathcal{D}_{W_u} -module $\check{\mathcal{M}}_{A_u}^{\beta^u}$ as defined in Definition 3.5, that is, $\check{\mathcal{M}}_{A_u}^{\beta^u} = \mathcal{D}_{W_u}/\check{\mathcal{I}}_{A_u}^{\beta^u}$ where the left ideal $\check{\mathcal{I}}_{A_u}^{\beta^u}$ is generated by*

$$\check{\square}_{\underline{m} \in \mathbb{L}_{A_u}} = \prod_{i \neq u: m_i > 0} w_{iu}^{m_i} - \prod_{i \neq u: m_i < 0} w_{iu}^{-m_i}$$

and the Euler vector fields:

$$\begin{aligned} \check{E}_k^u + \beta_k^u &= \sum_{i \neq u} a_{ki}^u \partial_{w_{iu}} w_{iu} + \beta_k^u \\ &= \sum_{i \neq u} (a_{ki} - a_{ku}) \partial_{w_{iu}} w_{iu} + \beta_k^u. \end{aligned}$$

Then the direct image $g_{u+} \mathcal{O}_T^{\beta^u}$ is isomorphic to $\check{\mathcal{M}}_{A_u}^{\beta^u}$. Moreover, the Hodge filtration on $\check{\mathcal{M}}_{A_u}^{\beta^u}$ is the order filtration shifted by $(n-d)$, i.e.

$$F_{p+(n-d)}^H \check{\mathcal{M}}_{A_u}^{\beta^u} = F_p^{\text{ord}} \check{\mathcal{M}}_{A_u}^{\beta^u}.$$

Proof. The statement follows from Theorem 4.17 and Lemma 5.8. □

We now want to compute how the \mathcal{D} -modules $g_{u+} \mathcal{O}_T^{\beta^u}$ glue on their common domain of definition. Let $u_1, u_2 \in \{0, \dots, n\}$ and denote by $W_{u_1 u_2}$ the intersection $W_{u_1} \cap W_{u_2}$. We fix $u_1, u_2 \in \{0, \dots, n\}$ with $u_1 < u_2$. We have the following change of coordinates between the charts W_{u_1} and W_{u_2}

$$w_{iu_1} = w_{iu_2} w_{u_1 u_2}^{-1} \quad \text{for } i \neq u_2 \quad \text{and} \quad w_{u_2 u_1} = w_{u_1 u_2}^{-1}$$

which gives the following transformation rules for vector fields:

$$w_{iu_1} \partial_{w_{iu_1}} = w_{iu_2} \partial_{w_{iu_2}} \quad \text{for } i \neq u_2 \quad w_{u_2 u_1} \partial_{w_{u_2 u_1}} = - \sum_{i \neq u_2} w_{iu_2} \partial_{w_{iu_2}}. \quad (42)$$

These transformation rules define an algebra isomorphism

$$\iota_{u_1 u_2} : D_{W_{u_1}}[w_{u_2 u_1}^{-1}] \longrightarrow D_{W_{u_2}}[w_{u_1 u_2}^{-1}].$$

The module of global sections $\Gamma(W_{u_1 u_2}, g_{u_1+} \mathcal{O}_T^{\beta^{u_1}})$ can be expressed as the quotient $D_{W_{u_1}}[w_{u_2 u_1}^{-1}] / \check{I}_{A_{u_1}}^{\beta^{u_1}}$, where $\check{I}_{A_{u_1}}^{\beta^{u_1}} \subset D_{W_{u_1}}[w_{u_2 u_1}^{-1}] := \mathbb{C}[(w_{iu_1})_{i \neq u_1}][w_{u_2 u_1}^{-1}] \otimes_{\mathbb{C}[(w_{iu_1})_{i \neq u_1}]} D_{W_{u_1}}$ is the left ideal generated by

1. $\check{E}_k^{u_1} + \beta_k^{u_1} = \sum_{\substack{i=0 \\ i \neq u_1}}^n a_{ki}^{u_1} \partial_{w_{iu_1}} w_{iu_1} + \beta_k^{u_1} = \sum_{\substack{i=0 \\ i \neq u_1}}^n a_{ki}^{u_1} w_{iu_1} \partial_{w_{iu_1}} + \sum_{\substack{i=0 \\ i \neq u_1}}^n a_{ki}^{u_1} + \beta_k^{u_1} \quad k = 1, \dots, d$
2. $\check{\square}_{\underline{m}} = \prod_{\substack{m_i > 0 \\ i \neq u_1}} w_{iu_1}^{m_i} - \prod_{\substack{m_i < 0 \\ i \neq u_1}} w_{iu_1}^{-m_i} \quad \underline{m} \in \mathbb{L}_{A_{u_1}}.$

Let $\gamma^{u_1} := \sum_{i \neq u_1} a_{ki}^{u_1}$, then $\check{I}_{A_{u_1}}^{\beta^{u_1} - \gamma^{u_1}} \subset D_{W_{u_1}}[w_{u_2 u_1}^{-1}]$ is the left ideal generated by

1. $\sum_{\substack{i=0 \\ i \neq u_1}}^n a_{ki}^{u_1} w_{iu_1} \partial_{w_{iu_1}} + \beta_k^{u_1} \quad k = 1, \dots, d$
2. $\square_l = \prod_{\substack{l_i > 0 \\ i \neq u_1}} w_{iu_1}^{l_i} - \prod_{\substack{l_i < 0 \\ i \neq u_1}} w_{iu_1}^{-l_i} \quad l \in \mathbb{L}_{A_{u_1}}$

We get the following isomorphism of $D_{W_{u_1}}[w_{u_2 u_1}^{-1}]$ -modules

$$\begin{aligned} D_{W_{u_1}}[w_{u_2 u_1}^{-1}]/\dot{I}_{A_{u_1}}^{\beta^{u_1}} &\longrightarrow D_{W_{u_1}}[w_{u_2 u_1}^{-1}]/\dot{I}_{A_{u_1}}^{\beta^{u_1}-\gamma^{u_1}} \\ 1 &\mapsto \prod_{i \neq u_1} w_{i u_1}^{-1} \end{aligned} \quad (43)$$

which is the image of the isomorphism

$$\begin{aligned} \mathcal{O}_T^{\gamma^{u_1}+\beta_k^{u_1}-1} &\longrightarrow \mathcal{O}_T^{\beta_k^{u_1}-1} \\ 1 &\mapsto \underline{t}^{-\gamma^{u_1}} = \underline{t}^{-\sum_{i \neq u_1} (a_i - a_{u_1})} \end{aligned}$$

under the functor g_{u_1+} (cf. equation (28)). One obtains the same results for the chart W_{u_2} by exchanging u_1 and u_2 above. Using the transformation rules (42), we can identify $D_{W_{u_1}}[w_{u_2 u_1}^{-1}]$ with $D_{W_{u_2}}[w_{u_1 u_2}^{-1}]$, which gives a well-defined map

$$\iota_{u_1 u_2} : D_{W_{u_1}}[w_{u_2 u_1}^{-1}]/\dot{I}_{A_{u_1}}^{\beta^{u_1}-\gamma^{u_1}} \longrightarrow D_{W_{u_2}}[w_{u_1 u_2}^{-1}]/\dot{I}_{A_{u_2}}^{\beta^{u_2}-\gamma^{u_2}}.$$

We can now give an explicit expression for the gluing map between the various charts of the module $g_+ \mathcal{O}_T^\beta$.

Lemma 5.10. *The isomorphism between $g_{u_1+} \mathcal{O}_T^{\beta^{u_1}}$ and $g_{u_2+} \mathcal{O}_T^{\beta^{u_2}}$ on their common domain of definition $W_{u_1 u_2} = W_{u_1} \cap W_{u_2}$ is given by*

$$\begin{aligned} \Gamma(W_{u_1 u_2}, g_{u_1+} \mathcal{O}_T) &\simeq D_{W_{u_1}}[w_{u_2 u_1}^{-1}]/\dot{I}_{A_{u_1}}^{\beta^{u_1}} \longrightarrow D_{W_{u_2}}[w_{u_1 u_2}^{-1}]/\dot{I}_{A_{u_2}}^{\beta^{u_2}} \simeq \Gamma(W_{u_1 u_2}, g_{u_2+} \mathcal{O}_T) \\ P &\longmapsto \iota_{u_1 u_2}(P) w_{u_1 u_2}^{n+1}. \end{aligned}$$

Proof. This follows easily from the discussion above by concatenating the three maps

$$D_{W_{u_1}}[w_{u_2 u_1}^{-1}]/\dot{I}_{A_{u_1}}^{\beta^{u_1}} \longrightarrow D_{W_{u_1}}[w_{u_2 u_1}^{-1}]/\dot{I}_{A_{u_1}}^{\beta^{u_1}-\gamma^{u_1}} \xrightarrow{\iota} D_{W_{u_2}}[w_{u_1 u_2}^{-1}]/\dot{I}_{A_{u_2}}^{\beta^{u_2}-\gamma^{u_2}} \longrightarrow D_{W_{u_2}}[w_{u_1 u_2}^{-1}]/\dot{I}_{A_{u_2}}^{\beta^{u_2}}$$

and by using the simple computation

$$\left(\prod_{i \neq u_2} w_{i u_2} \right) \cdot \iota \left(\prod_{i \neq u_1} w_{i u_1}^{-1} \right) = w_{u_1 u_2}^{n+1}$$

□

Consider the following change of coordinates θ_u on $W_u \times V$.

$$\tilde{\lambda}_u = \lambda_u + \sum_{\substack{j=0 \\ j \neq u}}^n \lambda_j w_{j u} \quad , \quad \tilde{\lambda}_i = \lambda_i \quad \text{and} \quad \tilde{w}_{i u} = w_{i u} \quad (44)$$

for $i = 0, \dots, n$ and $i \neq u$. Notice that $\theta_u^{-1} \circ (g_u \times id) \circ \phi_u^- g_u \times id$ and \tilde{F}_u is just the projection $((\tilde{w}_{i u})_{i \neq u}, \tilde{\lambda}_0, \dots, \tilde{\lambda}_n) \mapsto \tilde{\lambda}_u$.

Proposition 5.11. *Consider the original coordinates $((w_{i u})_{i \neq u}, (\lambda_0, \dots, \lambda_n))$ of $W_u \times V$. Then there is an isomorphism of $\mathcal{D}_{W_u \times V}$ -modules $\mathcal{N}_u \simeq \mathcal{D}_{W_u \times V} / \mathcal{K}_{A_u}^{\beta^{u_1}}$, where $\mathcal{K}_{A_u}^{\beta^{u_1}}$ is the left $\mathcal{D}_{W_u \times V}$ -ideal generated by the following classes of operators*

1. $\sum_{\substack{i=0 \\ i \neq u}}^n a_{ki}^u \partial_{w_{i u}} w_{i u} - \sum_{i=1}^n a_{ki} \lambda_i \partial_{\lambda_i} + \beta_k$
2. $\check{\square}_{\underline{m}} = \prod_{\substack{m_i > 0 \\ i \neq u}} w_{i u}^{m_i} - \prod_{\substack{m_i < 0 \\ i \neq u}} w_{i u}^{-m_i} \quad \underline{m} \in \mathbb{L}_{A_u}$
3. $\partial_{\lambda_i} - w_{i u} \partial_{\lambda_u} \quad \text{for } i = 0, \dots, n \text{ and } i \neq u$
4. $\sum_{j=0}^n \lambda_j \partial_{\lambda_j} - \beta_0.$

Moreover, for $\tilde{\beta} = (\beta_0, \beta) \in \mathfrak{A}_{\tilde{A}}$ and $\beta_0 \in (-1, 0]$ we have

$$F_{p+(n-d)}^H \mathcal{N}_u \simeq F_p^{\text{ord}} \mathcal{D}_{W_u \times V} / \mathcal{K}_{A_u}^{\tilde{\beta}^u}.$$

Proof. Recall that $\mathcal{N}_u = \check{\mathcal{M}}_{A_u}^{\beta^u} \boxtimes \mathcal{D}_V / \left((\partial_{\tilde{\lambda}_i})_{i \neq u}, \tilde{\lambda}_u \partial_{\tilde{\lambda}_u} - \beta_0 \right) = \mathcal{D}_{W_u \times V} / \tilde{\mathcal{K}}_{A_u}^{\tilde{\beta}^u}$, where

$$\tilde{\mathcal{K}}_{A_u}^{\tilde{\beta}^u} = \left((\check{E}_k^u + \beta_k^u)_{k=1, \dots, d}, (\check{\square}_m)_{m \in \mathbb{L}_{A_u}}, (\partial_{\tilde{\lambda}_j})_{j \neq u}, (\tilde{\lambda}_u \partial_{\tilde{\lambda}_u} - \beta_0) \right).$$

Using the coordinate transformation (44) we see that $\tilde{\mathcal{K}}_{A_u}^{\tilde{\beta}^u}$ is transformed into the ideal $\mathcal{K}_{A_u}^{\tilde{\beta}}$ generated by the operators

$$\begin{aligned} \sum_{\substack{i=0 \\ i \neq u}}^n a_{ki}^u (\partial_{w_{iu}} - \lambda_i \partial_{\lambda_u}) w_{iu} + \beta_k^u & \quad k = 1, \dots, d \\ \check{\square}_m = \prod_{\substack{m_i > 0 \\ i \neq u}} w_{iu}^{m_i} - \prod_{\substack{m_i < 0 \\ i \neq u}} w_{iu}^{-m_i} & \quad m \in \mathbb{L}_{A_u} \\ \partial_{\lambda_i} - w_{iu} \partial_{\lambda_u} & \quad \text{for } i = 0, \dots, n \text{ and } i \neq u \\ (\lambda_u + \sum_{\substack{j=0 \\ j \neq u}}^n \lambda_j w_{ju}) \partial_{\lambda_u} - \beta_0. & \end{aligned}$$

The last operator can be rewritten (using the relations $\partial_{\lambda_i} - w_{iu} \partial_{\lambda_u}$, i.e. the third class of operators) as

$$\sum_{j=0}^n \lambda_j \partial_{\lambda_j} - \beta_0 \equiv (\lambda_u + \sum_{\substack{j=0 \\ j \neq u}}^n \lambda_j w_{ju}) \partial_{\lambda_u} - \beta_0.$$

The Euler-type operators $\sum_{\substack{i=0 \\ i \neq u}}^n a_{ki}^u (\partial_{w_{iu}} - \lambda_i \partial_{\lambda_u}) w_{iu}$ can be further simplified by writing

$$\begin{aligned} \sum_{\substack{i=0 \\ i \neq u}}^n a_{ki}^u (\partial_{w_{iu}} - \lambda_i \partial_{\lambda_u}) w_{iu} + \beta_k^u & \equiv \sum_{\substack{i=0 \\ i \neq u}}^n a_{ki}^u (\partial_{w_{iu}} w_{iu} - \lambda_i \partial_{\lambda_i}) + \beta_k^u \\ & = \sum_{\substack{i=0 \\ i \neq u}}^n a_{ki}^u \partial_{w_{iu}} w_{iu} - \sum_{i=1}^n a_{ki} \lambda_i \partial_{\lambda_i} + a_{ku} \sum_{i=0}^n \lambda_i \partial_{\lambda_i} + \beta_k^u \\ & \equiv \sum_{\substack{i=0 \\ i \neq u}}^n a_{ki}^u \partial_{w_{iu}} w_{iu} - \sum_{i=1}^n a_{ki} \lambda_i \partial_{\lambda_i} + a_{ku} \beta_0 + \beta_k^u, \\ & = \sum_{\substack{i=0 \\ i \neq u}}^n a_{ki}^u \partial_{w_{iu}} w_{iu} - \sum_{i=1}^n a_{ki} \lambda_i \partial_{\lambda_i} + \beta_k, \end{aligned}$$

where the first equivalence follows by using the relation $\sum_{j=0}^n \lambda_j \partial_{\lambda_j} \equiv (\lambda_u + \sum_{\substack{j=0 \\ j \neq u}}^n \lambda_j w_{ju}) \partial_{\lambda_u}$ from above.

Hence we obtain the presentation $\mathcal{N}_u \simeq \mathcal{D}_{W_u \times V} / \mathcal{K}_{A_u}^{\tilde{\beta}}$, and the statement on the Hodge filtration follows directly from Proposition 5.9. \square

5.4 A Koszul complex

In this subsection, we will construct a strict resolution of the filtered module (\mathcal{N}_u, F^H) . For this purpose, we first describe an alternative presentation of the ideal $\mathcal{K}_{A_u}^{\tilde{\beta}} \subset \mathcal{D}_{W_u \times V}$. Let A_u^s be the $(d+1) \times (2n+1)$ -

matrix with columns $(0, \underline{a}_0 - \underline{a}_u), \dots, (0, \widehat{\underline{a}_u - \underline{a}_u}), \dots, (0, \underline{a}_n - \underline{a}_u), (1, \underline{a}_0), \dots, (1, \underline{a}_n)$ (here the symbol $\widehat{}$ means that the zero column $(0, \underline{a}_u - \underline{a}_u)$ is omitted). In other words, we have

$$A_u^s = \left(\begin{array}{ccc|c|ccc} 0 & \dots & 0 & 1 & 1 & \dots & 1 \\ \hline & & & 0 & & & \\ & & & \vdots & & & \\ & & & 0 & & & \end{array} \right).$$

We prove in Lemma 5.13 that the \mathcal{D} -module underlying \mathcal{N}_u is isomorphic to a partial Fourier-Laplace transformed GKZ-system with respect to the matrix A_u^s and parameter $\tilde{\beta}_u$. With the help of the results in section 3.1 we construct a \mathcal{D} -free strictly filtered resolution of the filtered module $(\mathcal{N}_u, F_\bullet^H)$ in Proposition 5.15.

As a first step we prove some properties of the matrices A_u^s .

Lemma 5.12. *If, as before, $\mathbb{Z}\tilde{A} = \mathbb{Z}^{d+1}$ and $\mathbb{N}\tilde{A} = \mathbb{Z}^{d+1} \cap \mathbb{R}_{\geq 0}\tilde{A}$ hold, then we have $\mathbb{Z}A_u^s = \mathbb{Z}^{d+1}$ and $\mathbb{N}A_u^s = \mathbb{Z}^{d+1} \cap \mathbb{R}_{\geq 0}A_u^s$.*

Proof. From $\mathbb{Z}\tilde{A} = \mathbb{Z}^{d+1}$ we conclude $\mathbb{Z}A_u^s = \mathbb{Z}^{d+1}$ since evidently $\mathbb{Z}\tilde{A} \subset \mathbb{Z}A_u^s$. Hence it remains to show that the semi-group $\mathbb{N}A_u^s$ is normal. We have

$$C_u \cdot A_u^s = \left(\begin{array}{ccc|c|ccc} 0 & \dots & 0 & 1 & 1 & \dots & 1 \\ \hline & & & 0 & & & \\ & & & \vdots & & & \\ & & & 0 & & & \end{array} \right) =: (\underline{\tilde{a}}_1^u, \dots, \underline{\tilde{a}}_n^u, \underline{\tilde{a}}_0^u, \underline{\tilde{a}}_1^u, \dots, \underline{\tilde{a}}_n^u) \in M((d+1) \times (2n+1), \mathbb{Z}),$$

where $C_u \in GL(d+1, \mathbb{Z})$ is the matrix already used in Lemma 5.8. It suffices to show the normality property for the semi-group $\mathbb{N}(C_u \cdot A_u^s)$ since C_u is an invertible linear mapping, hence a homeomorphism. Suppose that we are given a linear combination

$$\underline{v} = \sum_{i=1}^n \lambda_i \underline{\tilde{a}}_i^u + \sum_{j=0}^n \mu_j \underline{\tilde{a}}_j^u \in \mathbb{Z}^{d+1},$$

where $\lambda_i, \mu_j \in \mathbb{R}_{\geq 0}$. Then $\underline{v} = \sum_{i=1}^n (\lambda_i + \mu_i) \underline{\tilde{a}}_i^u + \underline{\tilde{a}}_0^u \cdot \left(\sum_{j=0}^n \mu_j \right)$. Clearly, $\sum_{j=0}^n \mu_j \in \mathbb{N}$, and moreover, the vector $\sum_{i=1}^n (\lambda_i + \mu_i) \underline{\tilde{a}}_i^u$ lies in $\mathbb{R}_{\geq 0}A_u$, but the latter semi-group is normal according to Lemma 5.8. Hence we have $\sum_{i=1}^n (\lambda_i + \mu_i) \underline{\tilde{a}}_i^u \in \mathbb{N}A_u$, and therefore $\underline{v} \in \mathbb{N}\tilde{A}_u \subset \mathbb{N}(C_u \cdot A_u^s)$, as required. \square

We now show that \mathcal{N}_u can be interpreted as a partial Fourier-Laplace transformed GKZ-system. For this we consider the GKZ system $\mathcal{M}_{A_u^s}$ on $\tilde{W}_u \times V$ with coordinates $(\hat{w}_{iu})_{i \neq u}, \lambda_0, \dots, \lambda_n$. Let $\text{FL}_{\tilde{W}_u}$ be the partial Fourier-Laplace transformation which interchanges $\partial_{\hat{w}_{iu}}$ with $(w_{iu})_{i \neq u}$ and \hat{w}_{iu} with $-\partial_{w_{iu}}$.

Lemma 5.13. *Let $\tilde{\mathcal{I}}_{A_u^s}^{(\vee)}$ be the left $\mathcal{D}_{W_u \times V}$ ideal generated by the operators*

$$\square_{(\underline{m}, \underline{l})}^{(\vee)} := \prod_{\substack{m_i > 0 \\ i \neq u}} w_{iu}^{m_i} \prod_{\substack{l_i > 0 \\ 0 \leq i \leq n}} \partial_{\lambda_i}^{l_i} - \prod_{\substack{m_i < 0 \\ i \neq u}} w_{iu}^{-m_i} \prod_{\substack{l_i > 0 \\ 0 \leq i \leq n}} \partial_{\lambda_i}^{-l_i},$$

where $(\underline{m}, \underline{l}) = ((m_i)_{i \neq u}, l_0, \dots, l_n) \in \mathbb{L}_{A_u^s}$,

$$\tilde{E}_k^u - \beta_k := - \sum_{\substack{i=0 \\ i \neq u}}^n a_{ki}^u \partial_{w_{iu}} w_{iu} + \sum_{i=1}^n a_{ki} \lambda_i \partial_{\lambda_i} - \beta_k \quad \text{for } k = 1, \dots, d$$

(notice that the operators \tilde{E}_k^u are the same operators as in Proposition 5.11 1. above, but multiplied with -1 , which is useful for a Fourier-Laplace transformation that will be performed below) and

$$\tilde{E}_0^u - \beta_0 := \sum_{i=0}^n \lambda_i \partial_{\lambda_i} - \beta_0.$$

Then we have $\widetilde{\mathcal{I}}_{A_u^s}^{(\vee)\widetilde{\beta}_u} = \widetilde{\mathcal{K}}_{A_u^s}^{\widetilde{\beta}_u}$, and hence the $\mathcal{D}_{W_u \times V}$ -module \mathcal{N}_u is isomorphic to $\mathcal{D}_{W_u \times V} / \widetilde{\mathcal{I}}_{A_u^s}^{(\vee)\widetilde{\beta}_u}$. In other words, we have an isomorphism

$$\mathcal{N}_u \simeq \text{FL}_{\widetilde{W}_u} \mathcal{M}_{A_u^s}^{\widetilde{\beta}_u}.$$

Proof. For the first statement, notice that $\square_{(\underline{m},0)}^{(\vee)}$ equals the operator $\check{\square}_{\underline{m}}$ from the definition of the ideal \mathcal{K}_u . On the other hand, one can obtain all operators $\square_{(\underline{m},l)}^{(\vee)}$ from the operators $\check{\square}_{(\underline{m}',0)}$ using the relations $\partial_{\lambda_i} - w_{iu} \partial_{\lambda_u}$. The last statement follows by interchanging $\partial_{w_{iu}}$ with $-\hat{w}_{iu}$ and w_{iu} with $\partial_{\hat{w}_{iu}}$ in the classes of operators of type 1., 2., 3. and 4. in the definition of the ideal $\mathcal{K}_{A_u^s}^{\widetilde{\beta}_u}$. \square

In order to construct a strictly filtered resolution of \mathcal{N}_u , we use the theory of Euler-Koszul complexes, which we explained in section 3.1. It will be applied to the $\mathcal{D}_{\widetilde{W}_u \times V}$ -module $\mathcal{M}_{A_u^s}^{\widetilde{\beta}_u}$. As before, we work at the level of global sections.

Let $F_{\bullet}^{\hat{\omega}} D_{\widetilde{W}_u \times V}$ the filtration on $D_{\widetilde{W}_u \times V}$ corresponding to the weight vector

$$\begin{aligned} \hat{\omega} &= ((\text{weight}(\hat{w}_{iu}))_{i \neq u}, (\text{weight}(\partial_{\hat{w}_{iu}}))_{i \neq u}, \text{weight}(\lambda_0), \dots, \text{weight}(\lambda_n), \text{weight}(\partial_{\lambda_0}), \dots, \text{weight}(\partial_{\lambda_n})) \\ &:= \left(\underbrace{(1, \dots, 1)}_{n\text{-times}}, \underbrace{(0, \dots, 0)}_{n\text{-times}}, \underbrace{(0, \dots, 0)}_{(n+1)\text{-times}}, \underbrace{(1, \dots, 1)}_{(n+1)\text{-times}} \right). \end{aligned}$$

Notice that this filtration corresponds to the order filtration $F_{\bullet}^{\text{ord}} D_{W_u \times V}$ under the Fourier-Laplace transformation functor $\text{FL}_{\widetilde{W}_u}$. We obtain a filtered resolution $((K_u^{\bullet}, d), F^{\hat{\omega}})$ of $M_{A_u^s}^{\widetilde{\beta}_u}$. Using Remark 3.3 we show that resolution is strict.

Lemma 5.14. *The Euler-Koszul complex $(K_u^{\bullet}, F^{\hat{\omega}})$ is a resolution of $(M_{A_u^s}^{\widetilde{\beta}_u}, F^{\hat{\omega}})$ in the category of filtered $D_{\widetilde{W}_u \times V}$ -modules (with respect to the filtration $F_{\bullet}^{\hat{\omega}} D_{\widetilde{W}_u \times V}$), i.e., we have a quasi-isomorphism $K_u^{\bullet} \rightarrow M_{A_u^s}$ and the complex K_u^{\bullet} is strictly filtered.*

Proof. By Remark 3.3 above it is enough to show that $H^{-i}(\text{Gr}_{\bullet}^{F^{\hat{\omega}}} K_u^{\bullet}) = 0$ for $i \geq 1$ and $H^0(\text{Gr}_{\bullet}^{F^{\hat{\omega}}} K_u^{\bullet}) \simeq \text{Gr}_{\bullet}^{F^{\hat{\omega}}} M_{A_u^s}$. Denote by $GD_{\widetilde{W}_u \times V} = \text{Gr}_{\bullet}^{\hat{\omega}} D_{\widetilde{W}_u \times V}$ the associated graded object of $D_{\widetilde{W}_u \times V}$, by $(\hat{v}_{iu})_{i \neq u}$ the symbol of $(\partial_{\hat{w}_{iu}})_{i \neq u}$ and by μ_j the symbol of ∂_{λ_j} in $GD_{\widetilde{W}_u \times V}$. Since $\square_{(\underline{m},l)}$ is homogeneous in (∂_{λ_j}) and $\text{ord}_{\hat{\omega}}(\partial_{\hat{w}_{iu}}) = 0$ for all $i \neq u$ we have

$$\text{Gr}^{\hat{\omega}}(D_{\widetilde{W}_u \times V} / D_{\widetilde{W}_u \times V} J_{A_u^s}^g) = GD_{\widetilde{W}_u \times V} / J_{A_u^s}^g$$

where $J_{A_u^s}^g$ is generated by

$${}^g \square_{(\underline{m},l)} := \prod_{\substack{m_i > 0 \\ i \neq u}} \hat{v}_{iu}^{m_i} \prod_{\substack{l_i > 0 \\ 0 \leq i \leq n}} \mu_i^{l_i} - \prod_{\substack{m_i < 0 \\ i \neq u}} \hat{v}_{iu}^{-k_i} \prod_{\substack{l_i > 0 \\ 0 \leq i \leq n}} \mu_i^{-l_i}$$

Notice that

$$GD_{\widetilde{W}_u \times V} / J_{A_u^s}^g \simeq \mathbb{C}[(\hat{w}_{iu})_{i \neq u}, \lambda_0, \dots, \lambda_n] \otimes_{\mathbb{C}} \mathbb{C}[\mathbb{N}A_u^s]$$

The associated graded complex $\text{Gr}^{\hat{\omega}} K_u^{\bullet}$ is isomorphic to a Koszul complex

$$\text{Gr}^{\hat{\omega}} K_u^{\bullet} \simeq \text{Kos}(GD_{\widetilde{W}_u \times V} / \hat{J}_{A_u^s}^g, ({}^g E_k^u)_{k=0, \dots, d})$$

where ${}^g E_k^u$ is defined by

$${}^g \hat{E}_k^u := \sum_{i \neq u} a_{ki}^u \hat{w}_{iu} \hat{v}_{iu} + \sum_{i=1}^n a_{ki} \lambda_i \mu_i \quad \text{for } k = 1, \dots, d$$

and

$${}^g E_0^u := \sum_{i=0}^n \lambda_i \mu_i$$

It is shown in [BZGM15, Theorem 1.2] that the ${}^g E_k^u$ are part of a system of parameters. Since $\mathbb{N}A_u^s$ is a normal semi-group (see Lemma 5.12 above), the ring $GD_{\tilde{W}_u \times V} / J_{A_u^s}^g$ is Cohen-Macaulay. Hence $({}^g E_k^u)_{k=0, \dots, d}$ is a regular sequence in $GD_{\tilde{W}_u \times V} / J_{A_u^s}^g$. This shows that $H^{-i}(\mathrm{Gr}_{\bullet}^{\hat{\omega}} K_u^{\bullet}) = 0$ for $i \geq 1$. On the other hand, it follows from [SST00, Theorem 4.3.5] that $H^0(\mathrm{Gr}_{\bullet}^{\hat{\omega}} K_u^{\bullet}) = GD_{\tilde{W}_u \times V} / (J_{A_u^s}^g + ({}^g E_k^u)_{k=0, \dots, d}) \simeq \mathrm{Gr}_{\bullet}^{\hat{\omega}} M_{A_u^s}^{\tilde{\beta}}$, as required. \square

As a consequence, we obtain the filtered resolution of \mathcal{N}_u we are looking for. Let $J_{A_u^s}^{(\vee)}$ be the ideal in $D_{W_u \times V}$ generated by the box operators $\square_{(\underline{m}, \underline{l})}^{(\vee)}$ for $(\underline{m}, \underline{l}) \in \mathbb{L}_{A_u^s}$. Put

$$K_u^{(\vee)-l} = \bigoplus_{0 \leq i_1 < \dots < i_l \leq d} D_{W_u \times V} / J_{A_u^s}^{(\vee)} e_{i_1, \dots, i_l}$$

and define

$$K_u^{\bullet(\vee)} := \mathrm{Kos}(D_{W_u \times V} / J_{A_u^s}^{(\vee)}, (E_k^u - \beta_k)_{k=0, \dots, d}),$$

where the E_k^u denote the (pairwise commuting) endomorphisms of $D_{W_u \times V} / J_{A_u^s}^{(\vee)}$ induced from right multiplication by E_k^u on $D_{W_u \times V}$. Define a filtration $\{F_{\bullet}^{(\vee)} K_u^{\bullet(\vee)}\}$ on $K_u^{\bullet(\vee)}$ by

$$F_p^{(\vee)} K_u^{(\vee)l} := \bigoplus_{0 \leq i_1 < \dots < i_p \leq d} F_{p-l+(n-d)}^{\mathrm{ord}} D_{W_u \times V} / J_{A_u^s}^{(\vee)}.$$

Then we have

Proposition 5.15. *We have a filtered quasi-isomorphism $(K_u^{\bullet(\vee)}, F_{\bullet}^{(\vee)}) \simeq (N_u, F_{\bullet}^{\mathrm{ord}}) \simeq (N_u, F_{\bullet+(n-d)}^H)$, i.e. the complex $(K_u^{\bullet(\vee)}, F_{\bullet}^{(\vee)})$ is a resolution of $(N_u, F_{\bullet+(n-d)}^H)$ in the category of filtered $D_{W_u \times V}$ -modules.*

Proof. The filtered quasi-isomorphism $(K_u^{\bullet(\vee)}, F_{\bullet}^{(\vee)}) \simeq (N_u, F_{\bullet}^{\mathrm{ord}})$ is obtained by applying the Fourier-Laplace functor $\mathrm{FL}_{\tilde{W}_u}$ to the (filtered) Euler-Koszul complex $(K_u^{\bullet(\vee)}, F_{\bullet}^{\hat{\omega}})$ from above (using Lemma 5.14). The second filtered (quasi-)isomorphism $(N_u, F_{\bullet}^{\mathrm{ord}}) \simeq (N_u, F_{\bullet+(n-d)}^H)$ is just the content of Proposition 5.11. \square

5.5 \mathcal{R} -modules

In Subsection 5.4 we explicitly computed the filtered \mathcal{D} -module $(\mathcal{N}, F_{\bullet}^H)$ in the charts $W_u \times V$. Since the direct image of \mathcal{N} under $\pi_2 : \mathbb{P}(W) \times V \rightarrow V$ is the GKZ-system we are looking for we would just need to compute a filtered version of $R\pi_{2*} DR_{\mathbb{P}(W) \times V/V}(\mathcal{N})$ using a Čech argument. It turns out that the theory of \mathcal{R} -modules is most suitable for this task. Hence here we lift the results of the Subsection 5.4 to the level of \mathcal{R} -modules. Following [Sab05], we first recall very briefly the basic notion of \mathcal{R} -modules and the Rees construction which provides a functor from the category of filtered \mathcal{D} -modules to that of \mathcal{R} -modules. In Proposition 5.17 we compute the corresponding Rees object of \mathcal{N}_u and of its resolution 5.18. We glue these resolutions in order to obtain a global resolution of the Rees object of \mathcal{N} in Proposition 5.19.

Let X be a smooth variety of dimension n . The order filtration of \mathcal{D}_X gives rise to the Rees ring $R_F \mathcal{D}_X$. Given a filtered \mathcal{D}_X -module $(\mathcal{M}, F_{\bullet} \mathcal{M})$ we construct the corresponding graded $R_F \mathcal{D}_X$ -module $R_F \mathcal{M} := \bigoplus_{k \in \mathbb{Z}} F_k \mathcal{M} z^k$. In local coordinates the sheaf of rings $R_F \mathcal{D}_X$ is given by

$$R_F \mathcal{D}_X = \mathcal{O}_X[z] \langle z \partial_{x_1}, \dots, z \partial_{x_n} \rangle$$

Denote by \mathcal{X} the product $X \times \mathbb{C}$. We will consider the sheaf

$$\mathcal{R}_{\mathcal{X}} = \mathcal{O}_{\mathcal{X}} \otimes_{\mathcal{O}_X[z]} R_F \mathcal{D}_X$$

and its ring of global sections

$$R_{\mathcal{X}} := \Gamma(\mathcal{X}, \mathcal{R}_{\mathcal{X}}) = \mathcal{O}_X(X)[z] \langle z \partial_{x_1}, \dots, z \partial_{x_n} \rangle$$

Given an $R_F \mathcal{D}_X$ -module $R_F \mathcal{M}$ the corresponding $\mathcal{R}_{\mathcal{X}}$ -module is

$$\mathcal{M} := \mathcal{O}_{\mathcal{X}} \otimes_{\mathcal{O}_{X[z]}} R_F \mathcal{M}$$

This gives an exact functor \mathcal{T} from the category of filtered \mathcal{D}_X -modules $MF(\mathcal{D}_X)$ to the category of $\mathcal{R}_{\mathcal{X}}$ -modules $Mod(\mathcal{R}_{\mathcal{X}})$

$$\begin{aligned} \mathcal{T} : MF(\mathcal{D}_X) &\longrightarrow Mod(\mathcal{R}_{\mathcal{X}}) \\ (\mathcal{M}, F_{\bullet} \mathcal{M}) &\mapsto \mathcal{M} \end{aligned}$$

We denote by $Mod_{qc}(\mathcal{R}_{\mathcal{X}})$ the category of $\mathcal{R}_{\mathcal{X}}$ -modules which are quasi-coherent $\mathcal{O}_{\mathcal{X}}$ -modules. We denote by $\Omega_{\mathcal{X}}^1 = z^{-1} \Omega_{X \times \mathbb{C}/\mathbb{C}}^1$ the sheaf of algebraic 1-forms on \mathcal{X} relative to the projection $\mathcal{X} \rightarrow \mathbb{C}$ having at most a pole of order one along $z = 0$. If we put $\Omega_{\mathcal{X}}^k = \wedge^k \Omega_{\mathcal{X}}^1$, we get a deRham complex

$$0 \longrightarrow \mathcal{O}_{\mathcal{X}} \xrightarrow{d} \Omega_{\mathcal{X}}^1 \xrightarrow{d} \dots \xrightarrow{d} \Omega_{\mathcal{X}}^n \longrightarrow 0$$

where the differential d is induced by the relative differential $d_{X \times \mathbb{C}/\mathbb{C}}$. If X is a smooth affine variety we get the following equivalence of categories.

Lemma 5.16. *Let X be a smooth affine variety. The functor*

$$\Gamma(\mathcal{X}, \bullet) : Mod_{qc}(\mathcal{R}_{\mathcal{X}}) \longrightarrow Mod(R_{\mathcal{X}})$$

is exact and gives an equivalence of categories.

Proof. The proof is completely parallel to the \mathcal{D} -module case (see e.g. [HTT08, Proposition 1.4.4]). \square

One can also define a notion of direct image in the category of \mathcal{R} -modules. Since we only need the case of a projection, we will restrict ourselves to this special situation. Let X, Y smooth algebraic varieties and $f : X \times Y \rightarrow Y$ be the projection to the second factor. Similarly as above we have a relative de Rham complex $\Omega_{\mathcal{X} \times \mathcal{Y}/\mathcal{Y}}^{\bullet} = z^{-1} \Omega_{X \times Y \times \mathbb{C}/Y \times \mathbb{C}}^{\bullet}$. If \mathcal{M} is an $\mathcal{R}_{\mathcal{X} \times \mathcal{Y}}$ -module the relative de Rham complex $DR_{\mathcal{X} \times \mathcal{Y}/\mathcal{Y}}(\mathcal{M})$ is locally given by

$$d(\omega \otimes m) = d\omega \otimes m + \sum_{i=1}^n \left(\frac{dx_i}{z} \wedge \omega \right) \otimes z \partial_{x_i} m$$

where $(x_i)_{1 \leq i \leq n}$ is a local coordinate of X . The direct image with respect to f is then defined as

$$f_+ \mathcal{M} := Rf_* DR_{\mathcal{X} \times \mathcal{Y}/\mathcal{Y}}(\mathcal{M})[n]$$

Recall that for a filtered \mathcal{D} -module $(\mathcal{M}, F_{\bullet} \mathcal{M})$ the direct image under f is given by

$$f_+ \mathcal{M} = Rf_* \left(0 \rightarrow \mathcal{M} \rightarrow \Omega_{X \times Y/Y}^1 \otimes \mathcal{M} \rightarrow \dots \rightarrow \Omega_{X \times Y/Y}^n \otimes \mathcal{M} \rightarrow 0 \right) [n]$$

together with its filtration

$$F_p f_+ \mathcal{M} = Rf_* \left(0 \rightarrow F_p \mathcal{M} \rightarrow \Omega_{X \times Y/Y}^1 \otimes F_{p+1} \mathcal{M} \rightarrow \dots \rightarrow \Omega_{X \times Y/Y}^n \otimes F_{p+n} \mathcal{M} \rightarrow 0 \right) [n]$$

Notice, however, that if $(\mathcal{M}, F_{\bullet} \mathcal{M})$ underlies a mixed Hodge module on $X \times Y$, then the Hodge filtration on the cohomology modules of the direct image complex is not, in general, given by this definition, unless X is projective. It is a straightforward exercise to check that the functor \mathcal{T} commutes with the direct image functor f_+ .

We will apply this to the filtered \mathcal{D} -module $(\mathcal{N}, F_{\bullet}^H)$ as defined in equation (40) in order to compute $\mathcal{H}^0 \pi_{2+} \mathcal{N} \simeq \mathcal{H}^{2n+d+1}(p_+(q^\dagger \mathcal{O}_T^\beta \otimes F^\dagger j_{\dagger} \mathcal{O}_{\mathbb{C}^*}^{-\beta_0-1}))$ together with its corresponding Hodge filtration. We will denote by $\mathcal{P} \times \mathcal{V}$ the space $\mathbb{P}(W) \times V \times \mathbb{C}$. The corresponding \mathcal{R} -module is

$$\mathcal{N} := \mathcal{T}(\mathcal{N}) = \mathcal{O}_{\mathcal{P} \times \mathcal{V}} \otimes_{\mathcal{O}_{\mathbb{P}(W) \times V}[z]}} R_{FH} \mathcal{N}$$

The direct image with respect to π_2 is then given by

$$\pi_{2+}\mathcal{N} \simeq R\pi_{2*} \left(0 \rightarrow \mathcal{N} \rightarrow \Omega^1_{\mathcal{D} \times \mathcal{V} / \mathcal{V}} \otimes \mathcal{N} \rightarrow \dots \rightarrow \Omega^n_{\mathcal{D} \times \mathcal{V} / \mathcal{V}} \otimes \mathcal{N} \rightarrow 0 \right) [n] \quad (45)$$

Since this is rather hard to compute, we will replace the complex

$$0 \rightarrow \mathcal{N} \rightarrow \Omega^1_{\mathcal{D} \times \mathcal{V} / \mathcal{V}} \otimes \mathcal{N} \rightarrow \dots \rightarrow \Omega^n_{\mathcal{D} \times \mathcal{V} / \mathcal{V}} \otimes \mathcal{N} \rightarrow 0$$

by a quasi-isomorphic one. For this we will construct a resolution of \mathcal{N} . Let $\mathcal{W}_u \times \mathcal{V} := W_u \times V \times \mathbb{C}$ and denote by \mathcal{N}_u the restriction of \mathcal{N} to $\mathcal{W}_u \times \mathcal{V}$. We write $R\mathcal{W}_u \times \mathcal{V} = \Gamma(\mathcal{W}_u \times \mathcal{V}, \mathcal{R}_{\mathcal{W}_u \times \mathcal{V}})$, then the module of global sections of \mathcal{N}_u is the $R\mathcal{W}_u \times \mathcal{V}$ -module

$$N_u := \Gamma(\mathcal{W}_u \times \mathcal{V}, \mathcal{N}_u).$$

Proposition 5.17. *The $R\mathcal{W}_u \times \mathcal{V}$ -module N_u is isomorphic to*

$$z^{n-d} \cdot R\mathcal{W}_u \times \mathcal{V} / \mathbb{I}_{A_u^s}$$

where $\mathbb{I}_{A_u^s}$ is generated by

$$\bar{\square}_{(m,l)} := \prod_{\substack{m_i > 0 \\ i \neq u}} w_{iu}^{m_i} \prod_{\substack{l_i > 0 \\ 0 \leq i \leq n}} (z\partial_{\lambda_i})^{l_i} - \prod_{\substack{m_i < 0 \\ i \neq u}} w_{iu}^{-m_i} \prod_{\substack{l_i < 0 \\ 0 \leq i \leq n}} (z\partial_{\lambda_i})^{-l_i},$$

where $(m, l) = ((m_i)_{i \neq u}, l_0, \dots, l_n) \in \mathbb{L}_{A_u^s}$,

$$\bar{E}_k^u - \beta_k := - \sum_{\substack{i=0 \\ i \neq u}}^n a_{ki}^u z \partial_{w_{iu}} w_{iu} + \sum_{i=1}^n a_{ki} \lambda_i z \partial_{\lambda_i} - \beta_k \quad \text{for } k = 1, \dots, d$$

and

$$\bar{E}_0^u - \beta_0 := \sum_{i=0}^n \lambda_i z \partial_{\lambda_i} - \beta_0$$

Proof. This follows easily from Lemma 5.13 and Lemma 5.16. \square

We will now define a Koszul complex K_u^\bullet in the category of R_u -modules which corresponds to the Koszul complex $K_u^{(\vee)}$ alluded to above. Write $\mathbb{J}_{A_u^s}$ for the left ideal in $R\mathcal{W}_u \times \mathcal{V}$ generated by all operators $\bar{\square}_{(m,l)}$ for $(m, l) \in \mathbb{L}_{A_u^s}$, then a computation similar to formula (7) shows that the maps

$$\begin{aligned} R\mathcal{W}_u \times \mathcal{V} / \mathbb{J}_{A_u^s} &\longrightarrow R\mathcal{W}_u \times \mathcal{V} / \mathbb{J}_{A_u^s} \\ P &\mapsto P \cdot (\bar{E}_k^u - \beta_k) \quad \text{for } k = 0, \dots, d \end{aligned} \quad (46)$$

are well-defined. Since $[\bar{E}_{k_1}^u - \beta_{k_1}, \bar{E}_{k_2}^u - \beta_{k_2}] = 0$ for $k_1, k_2 \in \{0, \dots, d\}$ we can build a Koszul complex

$$K_u^\bullet := \text{Kos} \left(z^{n-d} \cdot R\mathcal{W}_u \times \mathcal{V} / \mathbb{J}_{A_u^s}, (\bar{E}_k^u - \beta_k)_{k=0, \dots, d} \right)$$

whose terms are given by

$$z^{n-2d-1} \cdot R\mathcal{W}_u \times \mathcal{V} / \mathbb{J}_{A_u^s} \rightarrow \dots \rightarrow \bigoplus_{i=1}^n z^{n-d-1} \cdot R\mathcal{W}_u \times \mathcal{V} / \mathbb{J}_{A_u^s} e_1 \wedge \hat{e}_i \wedge \dots \wedge e_d \rightarrow z^{n-d} \cdot R\mathcal{W}_u \times \mathcal{V} / \mathbb{J}_{A_u^s} e_1 \wedge \dots \wedge e_d$$

Lemma 5.18. *The Koszul complex K_u^\bullet is a resolution of N_u .*

Proof. In order to prove the lemma it is enough to apply the exact Rees functor \mathcal{T} to the Koszul complex $K_u^{(\vee)}$ which is a strict resolution of N_u in the category of filtered $D_{W_u \times V}$ -modules by Proposition 5.15. \square

We denote by \mathcal{K}_u^\bullet the corresponding resolution of $\mathcal{N}_u = \mathcal{N}|_{W_u \times V}$. We are now able to construct a resolution of \mathcal{N} .

Proposition 5.19. *There exists a resolution \mathcal{K}^\bullet of \mathcal{N} in the category of $\mathcal{R}_{\mathcal{P} \times \mathcal{V}}$ -modules which is locally given by*

$$\Gamma(\mathcal{W}_u \times \mathcal{V}, \mathcal{K}^\bullet) = \mathbf{K}_u^\bullet$$

Proof. The resolution \mathcal{K}^\bullet is constructed by providing glueing maps between the $\mathbf{R}_{\mathcal{W}_{u_1 u_2} \times \mathcal{V}}$ -modules

$$\Gamma(\mathcal{W}_{u_1 u_2} \times \mathcal{V}, \mathcal{K}_{u_1}^\bullet) \simeq \mathbf{K}_{u_1}^\bullet[w_{u_2 u_1}^{-1}] \longrightarrow \Gamma(\mathcal{W}_{u_1 u_2} \times \mathcal{V}, \mathcal{K}_{u_2}^\bullet) \simeq \mathbf{K}_{u_2}^\bullet[w_{u_1 u_2}^{-1}]$$

which are compatible with the glueing maps on

$$\Gamma(\mathcal{W}_{u_1 u_2} \times \mathcal{V}, \mathcal{N}_{u_1}) \simeq \mathbf{N}_{u_1}[w_{u_2 u_1}^{-1}] \longrightarrow \Gamma(\mathcal{W}_{u_1 u_2} \times \mathcal{V}, \mathcal{N}_{u_2}) \simeq \mathbf{N}_{u_2}[w_{u_1 u_2}^{-1}]$$

Notice that the latter maps are given by

$$\begin{aligned} \mathbf{N}_{u_1}[w_{u_2 u_1}^{-1}] &\longrightarrow \mathbf{N}_{u_2}[w_{u_1 u_2}^{-1}] \\ P &\mapsto \iota_{u_1 u_2}(P)w_{u_1 u_2}^{n+1} \end{aligned}$$

which follows from Lemma 5.10 and by tracing back the functors applied to $g_{u+}\mathcal{O}_T$. Using the same argument as in Lemma 5.10 shows that the maps

$$\begin{aligned} \mathbf{K}_{u_1}^\bullet[w_{u_2 u_1}^{-1}] &\longrightarrow \mathbf{K}_{u_2}^\bullet[w_{u_1 u_2}^{-1}] \\ P &\mapsto \iota_{u_1 u_2}(P)w_{u_1 u_2}^{n+1} \end{aligned}$$

are well defined. We have to check that they give rise to a morphism of complexes. But this follows from the commutativity of the diagram

$$\begin{array}{ccc} P & \longrightarrow & \iota_{u_1 u_2}(P)w_{u_1 u_2}^{n+1} \\ \\ \mathbf{R}_{\mathcal{W}_{u_1} \times \mathcal{V}}/\mathbf{J}_{u_1} & \longrightarrow & \mathbf{R}_{\mathcal{W}_{u_2} \times \mathcal{V}}/\mathbf{J}_{u_2} \\ \uparrow \cdot \bar{E}_k^{u_1} - \beta_k & & \uparrow \cdot \bar{E}_k^{u_2} - \beta_k \\ \mathbf{R}_{\mathcal{W}_{u_1} \times \mathcal{V}}/\mathbf{J}_{u_1} & \longrightarrow & \mathbf{R}_{\mathcal{W}_{u_2} \times \mathcal{V}}/\mathbf{J}_{u_2} \\ \\ P & \longrightarrow & \iota_{u_1 u_2}(P)w_{u_1 u_2}^{n+1} \end{array}$$

□

5.6 A quasi-isomorphism

In order to compute the direct image of the \mathcal{R} -module \mathcal{N} under π_2 we have to deal with with the relative de Rham complex $DR_{\mathcal{P} \times \mathcal{V}|\mathcal{V}}(\mathcal{N})$ (cf. formula (45)). In this subsection we show in Proposition 5.20 that this complex is quasi-isomorphic to a complex \mathcal{L}^\bullet which is the top cohomology (with respect to the deRham differential) of the double complex $DR_{\mathcal{P} \times \mathcal{V}|\mathcal{V}}(\mathcal{K}^\bullet)$. In Proposition 5.21 we give a local description of this complex \mathcal{L}^\bullet on the charts $\mathcal{W}_u \times \mathcal{V}$.

As announced above we now apply the relative de Rham functor $DR_{\mathcal{P} \times \mathcal{V} | \mathcal{V}}$ to the resolution \mathcal{K}^\bullet and get a double complex $\Omega_{\mathcal{P} \times \mathcal{V} | \mathcal{V}}^{\bullet+n} \otimes \mathcal{K}^\bullet$:

$$\begin{array}{ccc}
\dots & \longrightarrow & \Omega_{\mathcal{P} \times \mathcal{V} | \mathcal{V}}^{n-1} \otimes \mathcal{K}^0 \xrightarrow{II d^{n,0}} \Omega_{\mathcal{P} \times \mathcal{V} | \mathcal{V}}^n \otimes \mathcal{K}^0 \\
& & \uparrow \scriptstyle{Id^{n-1,0}} \qquad \qquad \qquad \uparrow \scriptstyle{Id^{n,0}} \\
\dots & \longrightarrow & \Omega_{\mathcal{P} \times \mathcal{V} | \mathcal{V}}^{n-1} \otimes \mathcal{K}^{-1} \xrightarrow{II d^{n,-1}} \Omega_{\mathcal{P} \times \mathcal{V} | \mathcal{V}}^n \otimes \mathcal{K}^{-1} \\
& & \uparrow \qquad \qquad \qquad \uparrow \\
& & \vdots \qquad \qquad \qquad \vdots
\end{array}$$

The corresponding total complex is denoted by $Tot\left(\Omega_{\mathcal{P} \times \mathcal{V} | \mathcal{V}}^{\bullet+n} \otimes \mathcal{K}^\bullet\right)$.

Proposition 5.20. *The following natural morphisms of complexes*

$$\Omega_{\mathcal{P} \times \mathcal{V} | \mathcal{V}}^{\bullet+n} \otimes \mathcal{N} \longleftarrow Tot\left(\Omega_{\mathcal{P} \times \mathcal{V} | \mathcal{V}}^{\bullet+n} \otimes \mathcal{K}^\bullet\right) \longrightarrow \Omega_{\mathcal{P} \times \mathcal{V} | \mathcal{V}}^n \otimes \mathcal{K}^\bullet / II d\left(\Omega_{\mathcal{P} \times \mathcal{V} | \mathcal{V}}^{n-1} \otimes \mathcal{K}^\bullet\right) =: \mathcal{L}^\bullet$$

are quasi-isomorphisms.

Proof. Since the double complex $\Omega_{\mathcal{P} \times \mathcal{V} | \mathcal{V}}^{\bullet+n} \otimes \mathcal{K}^\bullet$ is bounded we can associate with it two spectral sequences which both converge. The first one is given by taking cohomology in the vertical direction which gives the IE_1 -page of the spectral sequence. Since \mathcal{K}^\bullet is a resolution of \mathcal{N} and $\Omega_{\mathcal{P} \times \mathcal{V} | \mathcal{V}}^l$ is a locally free (i.e. flat) $\mathcal{O}_{\mathcal{P} \times \mathcal{V}}$ -module for every $l = 1, \dots, n$, the only terms which are non-zero are the $IE_1^{0,q}$ -terms which are isomorphic to $\Omega_{\mathcal{P} \times \mathcal{V} | \mathcal{V}}^{q+n} \otimes \mathcal{N}$. Hence the first spectral sequence degenerates at the second page which shows that $\Omega_{\mathcal{P} \times \mathcal{V} | \mathcal{V}}^{\bullet+n} \otimes \mathcal{N} \leftarrow Tot\left(\Omega_{\mathcal{P} \times \mathcal{V} | \mathcal{V}}^{\bullet+n} \otimes \mathcal{K}^\bullet\right)$ is a quasi-isomorphism.

We now look at the second spectral sequence which is given by taking cohomology in the horizontal direction. We claim that $IE_1^{p,q} = 0$ for $q \neq 0$. It is enough to check this locally on the charts $\mathcal{W}_u \times \mathcal{V}$ using, moreover, Lemma 5.16 at the level of global sections. Notice that the complex

$$\Gamma(\mathcal{W}_u \times \mathcal{V}, \Omega_{\mathcal{P} \times \mathcal{V} | \mathcal{V}}^{\bullet+n} \otimes \mathcal{K}^l)$$

is isomorphic to a direct sum of Koszul complexes $Kos^\bullet(z^{-d-l}R_{\mathcal{W}_u \times \mathcal{V}}/J_{A_u^s}, \frac{1}{z}(z\partial_{w_{iu}} \cdot)_{i \neq u})$ where each summand is given by

$$z^{n-d-l}R_{\mathcal{W}_u \times \mathcal{V}}/J_{A_u^s} \longrightarrow \dots \longrightarrow z^{-d-l}R_{\mathcal{W}_u \times \mathcal{V}}/J_{A_u^s} e_1 \wedge \dots \wedge e_n.$$

The quotient $R_{\mathcal{W}_u \times \mathcal{V}}/J_{A_u^s}$ can be written as

$$\mathbb{C}[z, (z\partial_{w_{iu}})_{i \neq u}] \otimes_{\mathbb{C}[z]} \left(\mathbb{C}[z, \lambda_0, \dots, \lambda_n, (w_{iu})_{i \neq u}] \langle z\partial_{\lambda_0}, \dots, z\partial_{\lambda_n} \rangle / \left((\overline{\square}_{(m,l)})_{(m,l) \in \mathbb{L}_{A_u^s}} \right) \right)$$

Since the operators $z\partial_{w_{iu}} \cdot$ act only on the first term in the tensor product, we immediately see that $IE_1^{p,q} = 0$ for $q \neq 0$.

The fact that $Tot\left(\Omega_{\mathcal{P} \times \mathcal{V} | \mathcal{V}}^{\bullet+n} \otimes \mathcal{K}^\bullet\right) \rightarrow \mathcal{L}^\bullet$ is a quasi-isomorphism follows from the fact that $IE^{p,q} = 0$ for $q \neq 0$, i.e. the second spectral sequence degenerates at the second page. \square

The next result is an explicit local description of the complex \mathcal{L}^\bullet .

Proposition 5.21. *For any $u \in \{0, \dots, n\}$ define the ring*

$$S_{\mathcal{W}_u \times \mathcal{V}} := \mathbb{C}[z, \lambda_0, \dots, \lambda_n, (w_{iu})_{i \neq u}] \langle z\partial_{\lambda_0}, \dots, z\partial_{\lambda_n} \rangle$$

and denote by \mathcal{S} the sheaf of rings on $\mathcal{P} \times \mathcal{V}$ which is locally given by

$$\Gamma(\mathcal{W}_u \times \mathcal{V}, \mathcal{S}) = S_{\mathcal{W}_u \times \mathcal{V}}$$

with glueing maps

$$\begin{aligned} \mathbb{S}_{\mathcal{W}_{u_1} \times \mathcal{V}}[w_{u_2 u_1}^{-1}] &\longrightarrow \mathbb{S}_{\mathcal{W}_{u_2} \times \mathcal{V}}[w_{u_1 u_2}^{-1}] \\ P &\mapsto \iota_{u_1 u_2}(P) \end{aligned}$$

Denote by $J_{A_u^s}$ the left $\mathbb{S}_{\mathcal{W}_u \times \mathcal{V}}$ -ideal generated by the Box operators $\bar{\square}_{(m,l)}$ for $(m,l) \in \mathbb{L}_{A_u^s}$. Note that this is a slight abuse of notation, as the ideal generated by the same set of operators in the ring $\mathbb{R}_{\mathcal{W}_u \times \mathcal{V}}$ was also denoted by $J_{A_u^s}$, but which is justified by the fact that these generators do not contain the variables $z\partial_{w_{iu}}$. Then the complex \mathcal{L}^\bullet is given locally by

$$\Gamma(\mathcal{W}_u \times \mathcal{V}, \mathcal{L}^\bullet) \simeq \text{Kos}^\bullet(z^{-d} \mathbb{S}_{\mathcal{W}_u \times \mathcal{V}} / J_{A_u^s}, (\tilde{E}_k - \beta_k)_{k=0, \dots, d}) \quad (47)$$

whose terms are given by

$$z^{-2d-1} \mathbb{S}_{\mathcal{W}_u \times \mathcal{V}} / J_{A_u^s} \longrightarrow \dots \longrightarrow z^{-d} \mathbb{S}_{\mathcal{W}_u \times \mathcal{V}} / J_{A_u^s} e_1 \wedge \dots \wedge e_d$$

where

$$\begin{aligned} \tilde{E}_k - \beta_k &:= \sum_{i=1}^n a_{ki} \lambda_i z \partial_{\lambda_i} - \beta_k \quad \text{for } k = 1, \dots, d \\ \tilde{E}_0 - \beta_0 &:= \sum_{i=0}^n \lambda_i z \partial_{\lambda_i} - \beta_0 \end{aligned}$$

Proof. It follows from Proposition 5.20 that the 0-th cohomology of the complex $(\Omega_{\mathcal{P} \times \mathcal{V} / \mathcal{V}}^{\bullet+n} \otimes \mathcal{H}^p, \Pi d^{\bullet,p})$ is a direct sum of terms of the form $H^0(z^{-d} \text{Kos}^\bullet(\mathbb{R}_{\mathcal{W}_u \times \mathcal{V}} / J_{A_u^s}, \frac{1}{z}(z\partial_{w_{iu}} \cdot)_{i \neq u}))$. Taking the cokernel of left multiplication on $\mathbb{R}_{\mathcal{W}_u \times \mathcal{V}} / J_{A_u^s}$ by $z\partial_{w_{iu}}$ shows that we have an isomorphism of $\mathbb{S}_{\mathcal{W}_u \times \mathcal{V}}$ -modules

$$H^0(z^{-d} \text{Kos}^\bullet(\mathbb{R}_{\mathcal{W}_u \times \mathcal{V}} / J_{A_u^s}, (z\partial_{w_{iu}} \cdot)_{i \neq u})) \simeq z^{-d} \mathbb{S}_{\mathcal{W}_u \times \mathcal{V}} / J_{A_u^s}.$$

Hence equation (47) follows. \square

The ideals $J_{A_u^s}$ glue to an ideal $\mathcal{J} \subset \mathcal{S}$. Notice that the Euler vector fields $(\tilde{E}_k - \beta_k)_{k=0, \dots, d}$ are global sections of \mathcal{S} . We recall from Proposition 5.19 that the glueing maps for $\Gamma(\mathcal{W}_u \times \mathcal{V}, \Omega_{\mathcal{P} \times \mathcal{V}}^n \otimes \mathcal{H}^p)$ are given by:

$$\bigwedge_{\substack{i=0 \\ i \neq u_1}}^n dw_{iu_1} \otimes P \longmapsto \bigwedge_{\substack{i=0 \\ i \neq u_2}}^n dw_{iu_2} \cdot (w_{u_1 u_2})^{-n-1} \otimes \iota_{u_1 u_2}(P) w_{u_1 u_2}^{n+1}$$

Since both powers of $w_{u_1 u_2}$ on the right hand side cancel when considering the quotient \mathcal{L}^p , we see that

$$\mathcal{L}^\bullet \simeq \text{Kos}^\bullet(z^{-d} \mathcal{S} / \mathcal{J}, (\tilde{E}_k - \beta_k)_{k=0, \dots, d})$$

Summarizing, Proposition 5.20 and Proposition 5.21 show that instead of computing the direct image (45) we can compute

$$R\pi_{2*}(\mathcal{L}^\bullet) \simeq R\pi_{2*}(z^{-d} \text{Kos}^\bullet(\mathcal{S} / \mathcal{J}, (\tilde{E}_k - \beta_k)_{k=0, \dots, d}))$$

5.7 Computation of the direct image

In this subsection we continue our computation of $\pi_{2+} \mathcal{N} \simeq R\pi_{2*}(\mathcal{L}^\bullet)$. Because \mathcal{V} is affine it is enough to work at the level of global sections (cf. Lemma 5.16). We get:

$$\Gamma R\pi_{2*}(\mathcal{L}^\bullet) \simeq R\Gamma R\pi_{2*}(\mathcal{L}^\bullet) \simeq R\Gamma(\mathcal{L}^\bullet) \simeq R\Gamma(\text{Kos}^\bullet(z^{-d} \mathcal{S} / \mathcal{J}, (\tilde{E}_k - \beta_k)_{k=0, \dots, d})) \quad (48)$$

where the first isomorphism follows from the exactness of $\Gamma(\mathcal{V}, \bullet)$. Hence we have to consider the hypercohomology of a Koszul complex on $\mathcal{S} / \mathcal{J}$.

We will show that each term of this Koszul complex is Γ -acyclic. For this it is enough to show that \mathcal{S}/\mathcal{J} is Γ -acyclic. In Proposition 5.22 we show that \mathcal{S}/\mathcal{J} is acyclic and its corresponding associated graded module has an easy description in terms of the "semi" R-module S/J_{A^s} on $\mathbb{C}_z \times W \times V$ if all local cohomologies of S/J_{A^s} with respect to the ideal (w_0, \dots, w_n) vanish. Finally we show in Lemma 5.24 that the computation of these local cohomologies can be reduced to the computation of local cohomology groups of semigroup rings, which is a problem in commutative algebra and which we tackle in section 5.8.

Recall that $\mathcal{P} \times \mathcal{V} = \mathbb{C}_z \times \mathbb{P}(W) \times V$. We denote by $\mathcal{W} \times \mathcal{V}$ the space $\mathbb{C}_z \times W \times V$. Let

$$S := \mathbb{C}[z, w_0, \dots, w_n, \lambda_0, \dots, \lambda_n] \langle z\partial_{\lambda_0}, \dots, z\partial_{\lambda_n} \rangle$$

and consider the S-module

$$S/J_{A^s}$$

where the left ideal J_{A^s} is generated by

$$\square_{(m,l)} = \prod_{m_i > 0} w_i^{m_i} \prod_{l_i > 0} (z\partial_{\lambda_i})^{l_i} - \prod_{m_i < 0} w_i^{-m_i} \prod_{l_i < 0} (z\partial_{\lambda_i})^{-l_i} \quad \text{for } (m, l) \in \mathbb{L}_{A^s},$$

the matrix A^s is given by

$$A^s := (\underline{a}_0^s, \dots, \underline{a}_n^s, \underline{b}_0^s, \dots, \underline{b}_n^s) := \begin{pmatrix} 1 & 1 & \dots & 1 & 0 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 & 1 & 1 & \dots & 1 \\ 0 & a_{11} & \dots & a_{1n} & 0 & a_{11} & \dots & a_{1n} \\ \vdots & \vdots & & \vdots & \vdots & \vdots & & \vdots \\ 0 & a_{d1} & \dots & a_{dn} & 0 & a_{d1} & \dots & a_{dn} \end{pmatrix}$$

and \mathbb{L}_{A^s} is the \mathbb{Z} -module of relations among the columns of A^s . Notice that S/J_{A^s} is \mathbb{Z} -graded by the degree of the w_i . Denote by S_{w_u} the localization of S with respect to w_u , then one easily sees that the degree zero part $[S_{w_u}/J_{A^s}]_0$ of S_{w_u}/J_{A^s} is equal to $\Gamma(\mathcal{W}_u \times \mathcal{V}, \mathcal{S}/\mathcal{J}) \simeq S_{\mathcal{W}_u \times \mathcal{V}}/J_{A^s}$ if we identify w_i/w_u with w_{iu} . Let $\widetilde{S/J_{A^s}}$ be the associated sheaf on $\mathcal{P} \times \mathcal{V}$ having global sections $[S/J_{A^s}]_0$, then we obviously have

$$\widetilde{S/J_{A^s}} \simeq \mathcal{S}/\mathcal{J}$$

Define

$$\Gamma_*(\mathcal{S}/\mathcal{J}) := \bigoplus_{a \in \mathbb{Z}} \Gamma(\mathcal{P} \times \mathcal{V}, (\mathcal{S}/\mathcal{J})(a))$$

We want to use the following result applied to the graded module S/J_{A^s}

Proposition 5.22. [Gro61, Proposition 2.1.3] *There is the following exact sequence of \mathbb{Z} -graded S-modules*

$$0 \longrightarrow H_{(\underline{w})}^0(S/J_{A^s}) \longrightarrow S/J_{A^s} \longrightarrow \Gamma_*(\mathcal{S}/\mathcal{J}) \longrightarrow H_{(\underline{w})}^1(S/J_{A^s}) \longrightarrow 0$$

and for each $i \geq 1$, there is the following isomorphism

$$\bigoplus_{a \in \mathbb{Z}} H^i(\mathcal{P} \times \mathcal{V}, (\mathcal{S}/\mathcal{J})(a)) \simeq H_{(\underline{w})}^{i+1}(S/J_{A^s}) \quad (49)$$

where (\underline{w}) is the ideal in $\mathbb{C}[z, \lambda_0, \dots, \lambda_n, w_0, \dots, w_n]$ generated by w_0, \dots, w_n .

Proof. In the category of $\mathbb{C}[z, \lambda_0, \dots, \lambda_n, w_0, \dots, w_n]$ -modules, the statement follows from [Gro61, Proposition 2.1.3]. The statement in the category of S-modules follows from the proof given there. \square

In order to compute the local cohomology of S/J_{A^s} we introduce a variant of the Ishida complex (see e.g. [BH93, Theorem 6.2.5]). Let $T := \mathbb{C}[w_0, \dots, w_n, z\partial_{\lambda_0}, \dots, z\partial_{\lambda_n}] \subset S$ be a commutative subring and let $\mathbb{C}[\mathbb{N}A^s]$ be the affine semigroup algebra of A^s , i.e.

$$\mathbb{C}[\mathbb{N}A^s] = \{y^{\underline{c}} \in \mathbb{C}[y_0^\pm, \dots, y_{n+1}^\pm] \mid \underline{c} \in \mathbb{N}A^s \subset \mathbb{Z}^{d+2}\}$$

We have a map

$$\begin{aligned}\Phi_{A^s} : T &\longrightarrow \mathbb{C}[\mathbb{N}A^s] \\ w_i &\mapsto y_i^{a_i^s} \\ z\partial_{\lambda_i} &\mapsto y_i^{b_i^s}\end{aligned}$$

Notice that the kernel K_{A^s} of Φ_{A^s} is equal to the ideal in T generated by the elements $\square_{(k,l)}$, hence

$$T/K_{A^s} \simeq \mathbb{C}[\mathbb{N}A^s]$$

Remark 5.23. *The \mathbb{Z} -grading of T by the degree of the w_i induces a \mathbb{Z} -grading on $\mathbb{C}[\mathbb{N}A^s]$ since the operators $\square_{(k,l)}$ are homogeneous. The semi-group ring $\mathbb{C}[\mathbb{N}A^s] \subset \mathbb{C}[\mathbb{Z}^{d+2}]$ carries also a natural \mathbb{Z}^{d+2} -grading. Looking at the matrix A^s one sees that the \mathbb{Z} -grading coming from T is the first component of this \mathbb{Z}^{d+2} -grading.*

We regard $\mathbb{C}[\mathbb{N}A^s]$ as a T -module using the map Φ_{A^s} , which gives the isomorphisms

$$S/J_{A^s} \simeq S \otimes_T T/K_{A^s} \simeq S \otimes_T \mathbb{C}[\mathbb{N}A^s]$$

We want to express the local cohomology of S/J_{A^s} by the local cohomology of the commutative ring $\mathbb{C}[\mathbb{N}A^s]$. For this, let I be the ideal in $\mathbb{C}[\mathbb{N}A^s]$ generated by $y_0^{a_0^s}, \dots, y_n^{a_n^s}$, then we have the following change of rings formula:

Lemma 5.24. *There is the following isomorphism of \mathbb{Z} -graded S -modules:*

$$H_{(w)}^k(S/J_{A^s}) \simeq S \otimes_T H_I^k(\mathbb{C}[\mathbb{N}A^s])$$

Proof. Notice that if S was commutative this would be a standard property of the local cohomology groups. Here we have to adapt the proof slightly. First notice that it is enough to compute $H_{(w)}^k(S/J_{A^s})$ with an injective resolution of T -modules. To see why, let I^\bullet be an injective resolution (in the category of S -modules) of S/J_{A^s} . Since S is a free, hence flat, T -module, it follows from $\text{Hom}_S(S \otimes_T M, I) \simeq S \otimes_T \text{Hom}_T(M, I)$, that an injective S -module is also an injective T -module. Therefore we have

$$H_{(w)}^k(S/J_{A^s}) \simeq H^k \Gamma_{(w)}(I^\bullet) = H^k \Gamma_{I'}(I^\bullet) \simeq H_{I'}^k(S/J_{A^s})$$

where I' is the ideal in T generated by w_0, \dots, w_n and the second isomorphism follows from the equality

$$\Gamma_{(w)}(I^k) = \{x \in I^k \mid \forall i \exists k_i \text{ such that } w_i^{k_i} x = 0\} = \Gamma_{I'}(I^k)$$

Let J^\bullet be an injective resolution of T/K_{A^s} . In order to show the claim consider the following isomorphisms

$$\begin{aligned}S \otimes_T H_I^k(\mathbb{C}[\mathbb{N}A^s]) &\simeq S \otimes_T H_{I'}^k(T/K_{A^s}) \\ &\simeq S \otimes_T H^k \Gamma_{I'}(J^\bullet) \\ &\simeq H^k(S \otimes_T \Gamma_{I'}(J^\bullet)) \\ &\simeq H^k \Gamma_{I'}(S \otimes_T J^\bullet) \\ &\simeq H_{I'}^k(S/J_{A^s})\end{aligned}$$

where the third isomorphism follows from the fact that S is a flat T -module and the fifth isomorphism follows from the fact that $S \otimes_T J^\bullet$ is a T -injective resolution of $S/J_{A^s} \simeq S \otimes_T T/K_{A^s}$. \square

5.8 Local cohomology of semi-group rings

In this subsection we compute local cohomology groups of some special semigroup rings associated with the semigroups $\mathbb{N}A^s$. These local cohomology groups turned up in Lemma 5.24. We will use their vanishing in Subsection 5.9 (specifically in Corollary 5.32) to prove Γ -acyclicity of a Koszul complex.

We show that these local cohomology groups can be expressed as the cohomology of the Ishida complex of $\mathbb{C}[\mathbb{N}A^s]$ (cf. Proposition 5.25). Since A^s is a $(d+2) \times (2n+2)$ integer matrix the Ishida complex

carries a natural \mathbb{Z}^{d+2} -grading. The vanishing of certain graded pieces of the Ishida complex resp. of its cohomology depends on the position of the degree as seen as an element in \mathbb{R}^{d+2} in relation with the cone spanned by the columns of A^s (cf. Lemma 5.26 and Proposition 5.30).

Let \mathcal{F} be the face lattice of $\mathbb{R}_{\geq 0}A^s$ and denote by \mathcal{F}_σ the sublattice of faces which lie in the face σ spanned by $\underline{a}_0^s, \dots, \underline{a}_n^s$. For a face σ of $\mathbb{R}_{\geq 0}A^s$ consider the multiplicatively closed set

$$U_\sigma := \{y^\underline{c} \mid \underline{c} \in \mathbb{N}(A^s \cap \sigma)\}$$

and denote by $\mathbb{C}[\mathbb{N}A^s]_\sigma = \mathbb{C}[\mathbb{N}A^s + \mathbb{Z}(A^s \cap \sigma)]$ the localization with respect to U_σ . We put

$$L_\sigma^k = \bigoplus_{\substack{\tau \in \mathcal{F}_\sigma \\ \dim \tau = k}} \mathbb{C}[\mathbb{N}A^s]_\tau$$

and define maps $f^k : L_\sigma^k \rightarrow L_\sigma^{k+1}$ by specifying their components

$$f_{\tau', \tau}^k : \mathbb{C}[\mathbb{N}A^s]_{\tau'} \rightarrow \mathbb{C}[\mathbb{N}A^s]_\tau \quad \text{to be} \quad \begin{cases} 0 & \text{if } \tau' \not\subset \tau \\ \epsilon(\tau', \tau) \text{nat} & \text{if } \tau' \subset \tau \end{cases}$$

where ϵ is a suitable incidence function on \mathcal{F}_σ and nat is the natural localization morphism. The Ishida complex with respect to the face σ is

$$L_\sigma^\bullet : 0 \rightarrow L_\sigma^0 \rightarrow L_\sigma^1 \rightarrow \dots \rightarrow L_\sigma^{d+1} \rightarrow 0$$

The Ishida complex with respect to the face σ can be used to calculate local cohomology groups of $\mathbb{C}[\mathbb{N}A^s]$.

Proposition 5.25. *As above, denote by $I \subset \mathbb{C}[\mathbb{N}A^s]$ the ideal generated by the elements $\Phi_{A^s}(w_i) = y^{\underline{a}_i^s}$. Then for all k we have the isomorphism*

$$H_I^k(\mathbb{C}[\mathbb{N}A^s]) \simeq H^k(L_\sigma^\bullet)$$

Proof. The proof can be easily adapted from [BH93, Theorem 6.2.5]. For the convenience of the reader we sketch it here together with the necessary modifications. In order to show the claim we have to prove that the functors $N \mapsto H^k(L_\sigma^\bullet \otimes N)$ form a universal δ -functor (see e.g. [Har77]). If we can additionally show that

$$H_I^0(\mathbb{C}[\mathbb{N}A^s]) \simeq H^0(L_\sigma^\bullet) \tag{50}$$

the claim follows by [Har77, Corollary III.1.4]. Let $\mathcal{F}_\sigma(1)$ be the set of one-dimensional faces in \mathcal{F}_σ and notice that

$$H_{I'}^0(\mathbb{C}[\mathbb{N}A^s]) \simeq \ker \left(\mathbb{C}[\mathbb{N}A^s] \longrightarrow \bigoplus_{\tau \in \mathcal{F}_\sigma(1)} \mathbb{C}[\mathbb{N}A^s]_\tau \right) \simeq H^0(L_\sigma^\bullet \otimes_T M)$$

where $I' \subset \mathbb{C}[\mathbb{N}A^s]$ is the ideal generated by $\{y^{\underline{a}_i^s} \mid \mathbb{R}_{\geq 0}\underline{a}_i^s \in \mathcal{F}_\sigma(1)\}$. In order to show (50) we have to show that $\text{rad } I' = I$ since obviously $H_{I'}^0(\mathbb{C}[\mathbb{N}A^s]) = H_{\text{rad } I'}^0(\mathbb{C}[\mathbb{N}A^s])$. Since $I' \subset I$ and $I = \text{rad } I$ (I is a prime ideal corresponding to the face spanned by $\underline{a}_0^s, \dots, \underline{a}_n^s$), it is enough to check that a multiple of every $y^\underline{c} \in I$ lies in I' . But this follows easily from the fact that the elements $\{\underline{a}_i^s \mid \mathbb{R}_{\geq 0}\underline{a}_i^s \in \mathcal{F}_\sigma(1)\}$ span the same cone over \mathbb{Q} as the elements $\{\underline{a}_0^s, \dots, \underline{a}_n^s\}$.

The proof that $N \mapsto H^k(L_\sigma^\bullet \otimes_T N)$ is a δ -functor is completely parallel to the proof in [BH93]. \square

Notice that the complex L_σ^\bullet is \mathbb{Z}^{d+2} -graded since $\mathbb{C}[\mathbb{N}A^s]$ is \mathbb{Z}^{d+2} -graded. In order to analyze the cohomology of L_σ^\bullet we look at its \mathbb{Z}^{d+2} -graded parts. For this we have to determine when $(\mathbb{C}[\mathbb{N}A^s]_\tau)_x \neq 0$ (and therefore $(\mathbb{C}[\mathbb{N}A^s]_\tau)_x \simeq \mathbb{C}$) for $x \in \mathbb{Z}^{d+2}$.

We are following [BH93, Chapter 6.3]. Denote by C_{A^s} the cone $\mathbb{R}_{\geq 0}A^s \subset \mathbb{R}^{d+2}$. Let $x, y \in \mathbb{R}^{d+2}$. We say that y is visible from x if $y \neq x$ and the line segment $[x, y]$ does not contain a point $y' \in C_{A^s}$ with $y' \neq y$. A subset S is visible from X if each $v \in S$ is visible from x .

Recall that the cone C_{A^s} is given by the intersection of finitely many half-spaces

$$H_\tau^+ := \{x \in \mathbb{R}^{d+2} \mid \langle n_\tau, x \rangle \geq 0\} \quad \tau \in \mathcal{F}(d+1)$$

where $\mathcal{F}(d+1)$ is the set of $d+1$ -dimensional faces (facets) of C_{A^s} . We set

$$x^0 = \{\tau \mid \langle n_\tau, x \rangle = 0\}, \quad x^+ = \{\tau \mid \langle n_\tau, x \rangle > 0\}, \quad x^- = \{\tau \mid \langle n_\tau, x \rangle < 0\}$$

Lemma 5.26. [BH93, Lemma 6.3.2, 6.3.3]

1. A point $y \in C_{A^s}$ is visible from $x \in \mathbb{R}^{d+2} \setminus C_{A^s}$ if and only if $y^0 \cap x^- \neq \emptyset$.
2. Let $x \in \mathbb{Z}^{d+2}$ and let τ be a face of C_{A^s} . The \mathbb{C} -vector space $(\mathbb{C}[\text{INA}^s]_\tau)_x$ is non-zero if and only if τ is not visible from x .

Recall that the facet $\sigma \in \mathcal{F}(d+1)$ is spanned by $\underline{a}_0^s, \dots, \underline{a}_n^s$. It is the unique maximal element in the face lattice $\mathcal{F}_\sigma \subset \mathcal{F}$. Denote by H_σ its supporting hyperplane (i.e. $\sigma = C_{A^s} \cap H_\sigma$) which is given by


$$H_\sigma = \{x \in \mathbb{R}^{d+2} \mid \langle n_\sigma, x \rangle = 0\}$$

where $n_\sigma = (0, 1, 0, \dots, 0)$. Let $\tau \in \mathcal{F}_\sigma$ be a k -dimensional face contained in σ and set $I_\tau := \{i \mid \underline{a}_i^s \in \tau\}$. Notice that the vectors $\{\underline{a}_i^s \mid i \in I_\tau\}$ span the face τ . This face τ gives rise to two other faces, namely its "shadow" τ^s which is spanned by the vectors $\{\underline{b}_i^s \mid \underline{a}_i^s \in \tau\}$ and the unique $k+1$ -dimensional face τ^c which contains both τ and τ^s . Let $\{\tau_1, \dots, \tau_m\} = \mathcal{F}_\sigma(d)$ be the faces of dimension d contained in σ , which give rise to the facets $\tau_1^c, \dots, \tau_m^c$.

Example 5.27. Consider the matrix

$$A^s = \begin{pmatrix} 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 \\ 0 & 1 & -1 & 0 & 1 & -1 \end{pmatrix}$$

where the face σ is generated by $(1, 0, 0)$, $(1, 0, 1)$, $(1, 0, -1)$ and its shadow σ^s is generated by $(0, 1, 0)$, $(0, 1, 1)$, $(0, 1, -1)$. The facet τ^c is generated by τ and its shadow τ^s .



First notice that by Lemma 5.26.1 the facet σ is visible from a point $x \in \mathbb{R}^{d+2}$ if and only if $\langle n_\sigma, x \rangle < 0$. If $\langle n_\sigma, x \rangle \geq 0$ holds, it follows from Lemma 5.26.1. that a face $\tau_i \subset \sigma$ is visible from x if and only if the facet τ_i^c is visible from x , i.e. $\langle n_{\tau_i^c}, x \rangle < 0$.

We define

$$S := \mathbb{Z}^{d+2} \cap (\mathbb{R}(\underline{a}_0^s, \dots, \underline{a}_n^s) + \mathbb{R}_{\geq 0}(\underline{b}_0^s, \dots, \underline{b}_n^s)),$$

this is the set of \mathbb{Z}^{d+2} -degrees occurring in $\mathbb{C}[\text{INA}^s]_\sigma$. Notice also that we have

$$H_\sigma = \mathbb{R}(\underline{a}_0^s, \dots, \underline{a}_n^s) \quad \text{and} \quad H_\sigma^+ = \mathbb{R}(\underline{a}_0^s, \dots, \underline{a}_n^s) + \mathbb{R}_{\geq 0}(\underline{b}_0^s, \dots, \underline{b}_n^s).$$

Given a point $x \in S$ with $\langle n_\sigma, x \rangle \geq 0$ we will construct a point $y_x \in \mathbb{Z}^{d+2}$ which lies in H_σ such that τ_i is visible from x if and only if it is visible from y_x for all $i = 1, \dots, m$. Denote by z_x the projection

of x to the sub-vector space generated by $\underline{b}_0^s, \dots, \underline{b}_n^s$. Since the semi-group generated by these vectors is saturated, we can express z_x as a linear combination with positive integers

$$z_x = \sum_{i=0}^n r_i^x \underline{b}_i^s \quad \text{with } r_i^x \in \mathbb{N}$$

Since we have $0 = \langle n_{\tau_i^c}, \underline{a}_j^s - \underline{b}_j^s \rangle = \langle n_{\tau_i^c}, (1, -1, 0, \dots, 0) \rangle$ for any $\underline{a}_j^s, \underline{b}_j^s \in \tau_i^c$ the first two components of the vector $n_{\tau_i^c}$ are equal. Hence, if we set

$$y_x := x + \sum_{j=0}^n r_j^x \underline{a}_j^s - \sum_{i=0}^n r_i^x \underline{b}_i^s$$

we easily see that

$$\langle n_{\tau_i^c}, x \rangle = \langle n_{\tau_i^c}, y_x \rangle \quad \text{for } i = 1, \dots, m. \quad (51)$$

It follows that τ_i is visible from any point $x \in S$ if and only if it is visible from y_x , as required. Let us remark that the vectors x and y_x differ only in the first two components, because the same is true for the pair of vectors $(\underline{a}_i^s, \underline{b}_i^s)$ for all $i \in \{0, \dots, n\}$.

Lemma 5.28. *In the above situation, let $x \in S$. Then $y_x \in S \cap H_\sigma$ and we have*

$$(L_\sigma^\bullet)_x = (L_\sigma^\bullet)_{y_x}$$

Proof. For the first point, notice that the vector $x - z_x$ is precisely the projection of x to H_σ . On the other hand, we have $y_x = x - z_x + \sum_{i=0}^n n_i^x \underline{a}_i^s$, and $\sum_{i=0}^n n_i^x \underline{a}_i^s$ is an element of H_σ anyhow.

The second statement is an easy consequence of Lemma 5.26.2. More precisely, Equation (51) shows that the visibility of some facet τ_i^c is the same from x and from y_x . Moreover, σ is not visible from both x and y_x (i.e., $\langle n_\sigma, x \rangle \geq 0$, $\langle n_\sigma, y_x \rangle \geq 0$), hence, also the visibility of τ_i is the same from x and from y_x . We conclude that any localization $\mathbb{C}[\mathbb{N}A^s]_\tau$ (for any face $\tau \subset \sigma$) vanishes in degree x if and only if it vanishes in degree y_x . This yields the desired equality $(L_\sigma^\bullet)_x = (L_\sigma^\bullet)_{y_x}$. \square

We are now able to compute the cohomology of the Ishida complex with respect to the face σ . Set

$$H_{\tau_i^c}^- := \{x \in \mathbb{R}^{d+2} \mid \langle n_{\tau_i^c}, x \rangle < 0\} \quad \text{for } i = 1, \dots, m$$

and define

$$S^- := \mathbb{Z}^{d+2} \cap H_\sigma^+ \cap \bigcap_{i=1}^m H_{\tau_i^c}^-.$$


Notice that $S = \mathbb{Z}^{d+2} \cap H_\sigma^+$, hence we have a natural inclusion $S^- \subset S$.

Example 5.29. *We consider again the matrix*

$$A^s = \begin{pmatrix} 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 \\ 0 & 1 & -1 & 0 & 1 & -1 \end{pmatrix}$$

and take the point $x = (-1, 1, 1)$. Its projection to $\mathbb{R}(\underline{b}_0^s, \dots, \underline{b}_n^s)$ is $(0, 1, 1)$, hence we get

$$y_x = x + (1, 0, 1) - (0, 1, 1) = (0, 0, 1) \in H^\sigma.$$



Proposition 5.30. *Let A^s as above. Take any $x \in \mathbb{Z}^{d+2}$ and denote by L_σ^\bullet the Ishida complex with respect to the face σ generated by $\underline{a}_0^s, \underline{a}_1^s, \dots, \underline{a}_n^s$.*

1. *If $x \notin S$, then $(L_\sigma^\bullet)_x = 0$.*
2. *If $x \in S \setminus S^-$, then $H^i(L_\sigma^\bullet)_x = 0$ for all i*
3. *If $x \in S^-$, then $H^i(L_\sigma^\bullet)_x = 0$ for $i \neq d+1$ and $H^{d+1}(L_\sigma^\bullet)_x \simeq \mathbb{C}$.*

Proof. The first point follows from the fact that we have $(\mathbb{C}[\mathbb{N}A^s]_\sigma)_x = 0$ for $x \notin S$, hence $(L_\sigma^i)_x = 0$ for all i . For the proof of the second and third point, it is sufficient to consider the case where $x \in H_\sigma$: Namely, in both cases we have $x \in S$ so that Lemma 5.28 apply. We can thus replace x by y_x , i.e., $(L_\sigma^\bullet)_x = (L_\sigma^\bullet)_{y_x}$. Moreover, $x \in S^-$ if and only if $y_x \in S^- \cap H_\sigma$ by formula (51). Hence we will suppose in the remainder of this proof that $x \in S \cap H_\sigma$.

We will reduce statements 2. and 3. for $x \in S \cap H_\sigma$ to the computation of the local cohomology of a semi-group ring with respect to a maximal ideal via the Ishida complex as done in [BH93, Theorem 6.3.4]. For this, we will use the matrix $\tilde{A} = (\tilde{a}_0, \tilde{a}_1, \dots, \tilde{a}_n)$, which can be seen as the matrix of the first $n+1$ columns of A^s , with the second row deleted. The semigroup $\mathbb{N}\tilde{A}$ (resp. the cone $C_{\tilde{A}}$) embeds into $\mathbb{N}A^s$ (resp. into C_{A^s}) via the map $\tilde{a}_i \mapsto \underline{a}_i^s$, and these embeddings are compatible with the embeddings $\mathbb{R}^{r+1} \hookrightarrow \mathbb{R}^{r+2}$ (resp. $\mathbb{Z}^{r+1} \hookrightarrow \mathbb{Z}^{r+2}$) given by

$$(x_1, x_3, x_4, \dots, x_{d+2}) \mapsto (x_1, 0, x_3, x_4, \dots, x_{d+2}).$$

The following equality of semi-groups holds true:

$$S^- \cap H_\sigma = \mathbb{Z}^{d+1} \cap \text{Int}(-C_{\tilde{A}}), \quad (52)$$

where both intersections are taken in \mathbb{Z}^{d+2} . To show this, notice that $C_{\tilde{A}} = \mathbb{R}_{\geq 0}(\underline{a}_0^s, \dots, \underline{a}_n^s) \cap H_\sigma$, that $\tau_i^c \cap H_\sigma = \tau_i$ and hence

$$\bigcap_{i=1}^m (H_{\tau_i^c}^- \cap H_\sigma) = \text{Int}(-C_{\tilde{A}})$$

Consider the projection map

$$p: \mathbb{R}^{d+2} \longrightarrow \mathbb{R}^{d+1}$$

$$(x_1, x_2, x_3, \dots, x_{d+2}) \mapsto (x_1, x_3, \dots, x_{d+2})$$

which forgets the second component, then for all $\tau \subset \sigma$ and all elements $x \in S \cap H_\sigma$ we have that

$$(\mathbb{C}[\mathbb{N}A^s]_\tau)_x \simeq \left(\mathbb{C}[\mathbb{N}\tilde{A}]_{p(\tau)} \right)_{p(x)}$$

Under this isomorphism the \mathbb{Z}^{d+2} -graded part $(L_\sigma^\bullet)_x$ of the Ishida complex with respect to the face σ goes over to the \mathbb{Z}^{d+1} -graded part $(L^\bullet)_{p(x)}$ of the Ishida complex considered in [BH93] (i.e., the Ishida complex of the semi-group $\mathbb{C}[\mathbb{N}\tilde{A}]$ with respect to the maximal ideal generated by (w_0, \dots, w_n)). Using formula (52), the proposition follows now from Theorem 6.3.4 in loc.cit. \square

We finish this subsection by the following easy consequence, which will be crucial in the proof of the main result (Theorem 5.35 below).

Corollary 5.31. *In the above situation, we have $H^i(L_\sigma^\bullet) = 0$ for all $i \neq d+1$, $H^{d+1}(L_\sigma^\bullet)_x = 0$ for all $x \in \mathbb{Z}^{d+2} \setminus S^-$ and $\deg_{\mathbb{Z}}(H^{d+1}(L_\sigma^\bullet)_x) < 0$ for $x \in S^-$, where $\deg_{\mathbb{Z}}(-)$ refers to the \mathbb{Z} -grading of $H^i(L_\sigma^\bullet)$ corresponding to the first row of A^s .*

In other words, the cohomology groups of the Ishida complex (with respect to the face σ) are concentrated in negative degrees.

Proof. The first two statements are precisely those from Proposition 5.30, points 1. and 2. In order to show the third one, notice that for any $x \in S$, we have $\deg_{\mathbb{Z}}(x) \leq \deg_{\mathbb{Z}}(y_x)$ (this follows from the very definition of the vector y_x). Now let $x \in S^-$, and suppose that $H^{d+1}(L_\sigma^\bullet)_x \neq 0$. From Lemma 5.28 we deduce that

$$H^{d+1}(L_\sigma^\bullet)_x = H^{d+1}(L_\sigma^\bullet)_{y_x},$$

and as already remarked above, $y_x \in S^- \cap H_\sigma$ because $x \in S^-$. However, we deduce from formula (52) that $\deg_{\mathbb{Z}}(y_x) < 0$ if $y_x \in S^- \cap H_\sigma$, so that we obtain $\deg_{\mathbb{Z}}(H^{d+1}(L_\sigma^\bullet)_x) < 0$, as required. \square

5.9 Statement and proof of the main theorem

In this subsection we finally finish the computation of the direct image $\pi_{2+}\mathcal{N}$. The Γ -cyclicity of \mathcal{S}/\mathcal{J} (cf. 5.32) which follows directly from the results of the previous two subsections enables us to compute the global sections of $\pi_{2+}\mathcal{N}$ as the cohomology of a Koszul complex (cf. Proposition 5.33). Finally we are able to compute the Hodge filtration on the GKZ-system in Theorem 5.35.

Corollary 5.32. *The \mathcal{S} -modules \mathcal{S}/\mathcal{J} are Γ -acyclic.*

Proof. If we consider the degree zero part of formula (49), then it suffices to show that the \mathbb{Z} -graded local cohomology S -modules $H_{(w)}^*(S/J_{A_s})$ are concentrated in negative degrees. By Proposition 5.25 and Lemma 5.24, these local cohomology groups are calculate by the Ishida complex L_σ^\bullet , i.e., we have isomorphisms

$$H_{(w)}^k(S/J_{A_s}) \simeq S \otimes_T H_T^k(\mathbb{C}[\mathbb{N}A^s]) \simeq S \otimes_T H^k(L_\sigma^\bullet).$$

The cohomology groups $H^k(L_\sigma^\bullet)$ are concentrated in negative degrees by Corollary 5.31 (and tensoring with S does not change the \mathbb{Z} -degree which is counted with respect to the degree of the variables w_0, \dots, w_n). Hence the result follows. \square

Proposition 5.33. *There is the following isomorphism in $D^b(\mathcal{R}_V)$:*

$$\Gamma\pi_{2+}\mathcal{N} \simeq \Gamma(\text{Kos}^\bullet(z^{-d}\mathcal{S}/\mathcal{J}, (\tilde{E} - \beta_k)_{k=0, \dots, d}))$$

Proof. By formula (45), Proposition 5.20 and Proposition 5.21 we have the isomorphisms

$$\begin{aligned} \Gamma\pi_{2+}\mathcal{N} &\simeq \Gamma R\pi_{2*}(\Omega_{\mathcal{P} \times \mathcal{Y} / \mathcal{Y}}^{\bullet+n} \otimes \mathcal{N}) \\ &\simeq R\Gamma R\pi_{2*}(\Omega_{\mathcal{P} \times \mathcal{Y} / \mathcal{Y}}^{\bullet+n} \otimes \mathcal{N}) \\ &\simeq R\Gamma(\Omega_{\mathcal{P} \times \mathcal{Y} / \mathcal{Y}}^{\bullet+n} \otimes \mathcal{N}) \\ &\simeq R\Gamma(\mathcal{L}^\bullet) \end{aligned} \tag{53}$$

Using the last isomorphism in (48) and Corollary 5.32 we get

$$R\Gamma(\mathcal{L}^\bullet) \simeq R\Gamma(Kos^\bullet(z^{-d}\mathcal{S}/\mathcal{J}, (\tilde{E})_{k=0,\dots,d})) \simeq \Gamma(Kos^\bullet(z^{-d}\mathcal{S}/\mathcal{J}, (\tilde{E})_{k=0,\dots,d}))$$

□

Denote by $R_{\mathcal{Y}}$ the ring

$$R_{\mathcal{Y}} = \mathbb{C}[z, \lambda_0, \dots, \lambda_n] \langle z\partial_{\lambda_0}, \dots, z\partial_{\lambda_n} \rangle,$$

let $J_{\tilde{A}}^\lambda \subset R_{\mathcal{Y}}$ be the left ideal generated by

$$\square_{\underline{l}}^\lambda = \prod_{l_i > 0} (z\partial_{\lambda_i})^{l_i} - \prod_{l_i < 0} (z\partial_{\lambda_i})^{-l_i} \quad \text{for } \underline{l} \in \mathbb{L}_{\tilde{A}}$$

and let $I_{\tilde{A}}^\lambda \subset R_{\mathcal{Y}}$ be the left ideal generated by $J_{\tilde{A}}^\lambda$ and the operators

$$\begin{aligned} \tilde{E}_k - \beta_k &:= \sum_{i=1}^n a_{ki} \lambda_i z \partial_{\lambda_i} - \beta_k \quad \text{for } k = 1, \dots, d \\ \tilde{E}_0 - \beta_0 &:= \sum_{i=0}^n \lambda_i z \partial_{\lambda_i} - \beta_0 \end{aligned}$$

Lemma 5.34. *There is the following isomorphism of $R_{\mathcal{Y}}$ -modules*

$$\Gamma \mathcal{H}^0(\pi_{2+}\mathcal{N}) \simeq \mathcal{H}^0(\Gamma\pi_{2+}\mathcal{N}) \simeq z^{-d}R_{\mathcal{Y}}/I_{\tilde{A}}^\lambda$$

Proof. The first isomorphism follows from Lemma 5.16. The second isomorphism follows from Proposition 5.33, the isomorphism

$$\Gamma(\mathcal{S}/\mathcal{J}) \simeq R_{\mathcal{Y}}/J_{\tilde{A}}^\lambda$$

and the isomorphism

$$z^{-d}R_{\mathcal{Y}}/I_{\tilde{A}}^\lambda \simeq H^0\left(Kos^\bullet\left(z^{-d}R_{\mathcal{Y}}/J_{\tilde{A}}^\lambda, (\tilde{E}_k - \beta_k)_{k=0,\dots,d}\right)\right)$$

□

We are now able to prove the main theorem of this paper. Let \tilde{A} be the $(d+1) \times (n+1)$ integer matrix

$$\tilde{A} = \begin{pmatrix} 1 & 1 & \dots & 1 \\ 0 & a_{11} & \dots & a_{1n} \\ \vdots & \vdots & & \vdots \\ 0 & a_{d1} & \dots & a_{dn} \end{pmatrix}$$

given by a matrix $A = (a_{jk})$ such that $\mathbb{Z}\tilde{A} = \mathbb{Z}^{d+1}$ and such that $\mathbb{N}\tilde{A} = \mathbb{Z}^{d+1} \cap \mathbb{R}_{\geq 0}\tilde{A}$.

Theorem 5.35. *Let \tilde{A} be an integer matrix as above, $\tilde{\beta} \in \mathfrak{A}_{\tilde{A}}$ and $\beta_0 \in (-1, 0]$. The GKZ-system $\mathcal{M}_{\tilde{A}}^{\tilde{\beta}}$ carries the structure of a mixed Hodge module whose Hodge filtration is given by the shifted order filtration, i.e.*

$$(\mathcal{M}_{\tilde{A}}^{\tilde{\beta}}, F_\bullet^H) \simeq (\mathcal{M}_{\tilde{A}}^{\tilde{\beta}}, F_{\bullet+d}^{ord}).$$

Proof. Recall from Proposition 5.4 that we have the isomorphism

$${}^H\mathcal{M}_{\tilde{A}}^{\tilde{\beta}} \simeq \mathcal{H}^{2n+d+1}(p_*(q^{*p}\mathbb{C}_T^\beta \otimes F^*j_!^p\mathbb{C}_{\mathbb{C}^*}^{-\beta_0-1})) \in \text{MHM}(V).$$

The underlying \mathcal{D}_V -module of this mixed Hodge module is

$$\mathcal{H}^{2n+d+1}(p_+(q^\dagger\mathcal{O}_T^\beta \otimes_{\mathcal{O}} F^\dagger(j_!\mathcal{O}_{\mathbb{C}^*}^{-\beta_0-1}))) \simeq \mathcal{H}^0(\pi_{2+}\mathcal{N}).$$

We have already computed the Hodge filtration of \mathcal{N} (more precisely, we have computed it on the restrictions \mathcal{N}_u of \mathcal{N} to each chart $W_u \times V$ in Proposition 5.11). In order to compute the Hodge filtration under the direct image of π_2 , we will use the results obtained above and read off the Hodge filtration from the corresponding \mathcal{R}_V -module $\mathcal{F}(\mathcal{M}_{\tilde{A}}^{\tilde{\beta}}, F^H)$. We have the following isomorphisms

$$\Gamma \mathcal{F}(\mathcal{M}_{\tilde{A}}^{\tilde{\beta}}, F^H) \simeq \Gamma \mathcal{F}(\mathcal{H}^0(\pi_{2+}\mathcal{N}, F^H)) \simeq \Gamma \mathcal{H}^0(\pi_{2+}\mathcal{N}) \simeq z^{-d}R_{\mathcal{Y}}/I_{\tilde{A}}^\lambda$$

Using these isomorphisms the claim follows easily. □

5.10 Duality

For applications like the one presented in the next section, it will be useful to extend the computation of the Hodge filtration on $\mathcal{M}_{\tilde{A}}^{\beta}$ to the dual Hodge module $\mathbb{D}\mathcal{M}_{\tilde{A}}^{\beta}$. This is possible under the assumption made in the above main theorem (Theorem 5.35) plus the extra requirement that the semi-group ring $\mathbb{C}[\mathbb{N}\tilde{A}]$ is Gorenstein. More precisely, it follows from [Wal07], that under these assumptions, the \mathcal{D}_V -module $\mathbb{D}\mathcal{M}_{\tilde{A}}^{\beta}$ is still a GKZ-system. Hence it is reasonable to expect that its Hodge filtration will also be the order filtration up to a suitable shift.

The Gorenstein condition for normal semi-group rings has a well-known combinatorial expression (see [BH93, Corollary 6.3.8]), namely, $\mathbb{C}[\mathbb{N}\tilde{A}]$ is Gorenstein if and only if there is a vector \tilde{c} such that the set of interior points $\text{int}(\mathbb{N}\tilde{A})$ (i.e., the intersection of $\text{int}(\mathbb{R}_{\geq 0}\tilde{A}) \cap \mathbb{Z}^{d+1}$ is given by $\tilde{c} + \mathbb{N}\tilde{A}$.

Theorem 5.36. *Suppose that $\tilde{A} \in M(d+1 \times n+1, \mathbb{Z})$ is such that $\mathbb{Z}\tilde{A} = \mathbb{Z}^{d+1}$, $\mathbb{N}\tilde{A} = \mathbb{Z}^{d+1} \cap \mathbb{R}_{\geq 0}\tilde{A}$ and such that $\text{int}(\mathbb{N}\tilde{A}) = \tilde{c} + \mathbb{N}\tilde{A}$ for some $\tilde{c} = (c_0, c) \in \mathbb{Z}^{d+1}$ and $\beta \in \mathfrak{A}_A$. Then we have*

$$\mathbb{D}\mathcal{M}_{\tilde{A}}^{\tilde{\beta}} \simeq \mathcal{M}_{\tilde{A}}^{-\tilde{\beta}-\tilde{c}},$$

and the Hodge filtration on $\mathbb{D}\mathcal{M}_{\tilde{A}}^{\tilde{\beta}}$ is the order filtration, shifted by $n + c_0$, i.e., we have

$$F_p^H \mathbb{D}\mathcal{M}_{\tilde{A}}^{\tilde{\beta}} \simeq F_{p-n-c_0}^{\text{ord}} \mathcal{M}_{\tilde{A}}^{-\tilde{\beta}-\tilde{c}}.$$

Proof. The proof is very much parallel to [RS15, Proposition 2.19] resp. [RS17, Theorem 5.4], we will give the main ideas here once again for the convenience of the reader. We again work with the modules of global sections, and write $D_V := \mathbb{C}[\lambda_0, \dots, \lambda_n] \langle \partial_{\lambda_0}, \dots, \partial_{\lambda_n} \rangle$ and $S_{\tilde{A}}$ for the commutative ring $\mathbb{C}[\partial_{\lambda_0}, \dots, \partial_{\lambda_n}] / (\square_l)_{l \in \mathbb{L}_{\tilde{A}}}$. These rings are \mathbb{Z}^{d+1} -graded by $\deg(\lambda_i) = -\tilde{a}_i$, $\deg(\partial_{\lambda_i}) = \tilde{a}_i$.

In order to calculate $\mathbb{D}\mathcal{M}_{\tilde{A}}^{\tilde{\beta}}$ together with its Hodge filtration, we need to find a strictly filtered free resolution $(L_{\bullet}, F_{\bullet}) \xrightarrow{\sim} (M_{\tilde{A}}^{\tilde{\beta}}, F_{\bullet}^H) = (M_{\tilde{A}}^{\tilde{\beta}}, F_{\bullet+d}^{\text{ord}})$. We have already used in the previous sections of this paper resolutions of “Koszul”-type for various (filtered) \mathcal{D} -modules. Here we consider the Euler-Koszul complex

$$K^{\bullet} := \text{Kos}(D_V \otimes_{\mathbb{C}[\partial_{\lambda}]} S_{\tilde{A}}, (E_k - \beta_k)_{k=0, \dots, d}),$$

as defined in section 3.1 and a generalization to \mathbb{Z}^{d+1} -graded $\mathbb{C}[\partial_{\lambda}]$ -modules (for details see [MMW05]). A free resolution of $M_{\tilde{A}}^{\tilde{\beta}}$ is constructed as follows: Take a $\mathbb{C}[\partial]$ -free graded resolution of $T^{\bullet} \rightarrow S_{\tilde{A}}$, and define L^{\bullet} to be the total complex $\text{Tot}(K^{\bullet}(E - \beta, D_V \otimes_{\mathbb{C}[\partial_{\lambda}]} T^{\bullet}))$. Notice that the double complex $K^{\bullet}(E - \beta, D_V \otimes_{\mathbb{C}[\partial_{\lambda}]} T^{\bullet})$ exists since $K^{\bullet}(E - \beta, D_V \otimes_{\mathbb{C}[\partial]} -)$ is a functor from the category of \mathbb{Z}^{d+1} -graded $\mathbb{C}[\partial]$ -modules to the category of (bounded complexes of) \mathbb{Z}^{d+1} -graded D_V -modules. Then we have $L^{-k} = 0$ for all $k > n+1$ (notice that the length of the Euler-Koszul complexes is $d+1$, and the length of the resolution $T_{\bullet} \rightarrow P$ is $n-d+1$, hence the total complex has length $(d+1) + (n-d+1) - 1 = n+1$). Moreover, the last term L^{-n-1} of this complex is simply equal to D_V (and so is the first one L^0).

As we have $\text{int}(\mathbb{N}\tilde{A}) = \tilde{c} + \mathbb{N}\tilde{A}$, the ring $\mathbb{C}[\mathbb{N}\tilde{A}] \simeq S_{\tilde{A}}$ is Gorenstein, more precisely, we have $\omega_{S_{\tilde{A}}} \simeq S_{\tilde{A}}(\tilde{c})$, where $\omega_{S_{\tilde{A}}}$ is the canonical module of $S_{\tilde{A}}$. Then a spectral sequence argument (see also [Wal07, Proposition 4.1]), using

$$\text{Ext}_{\mathbb{C}[\partial]}^i(S_{\tilde{A}}, \omega_{\mathbb{C}[\partial]}) \simeq \begin{cases} 0 & \text{if } i < n-d \\ S_{\tilde{A}}(\tilde{c}) & \text{if } i = n-d \end{cases}$$

shows that

$$\mathbb{D}\mathcal{M}_{\tilde{A}}^{\tilde{\beta}} \cong M_{\tilde{A}}^{-\tilde{\beta}-\tilde{c}}.$$

In order to calculate the Hodge filtration on $M_{\tilde{A}}^{-\tilde{c}}$, we remark that the Euler-Koszul complex is naturally filtered by putting

$$F_p K^{-l} := \bigoplus_{0 \leq i_1 < \dots < i_l \leq l} F_{p+d-l}^{\text{ord}}(D_V \otimes_{\mathbb{C}[\partial]} S_{\tilde{A}}) e_{i_1 \dots i_l}.$$

Notice that $D_V \otimes_{\mathbb{C}[\partial]} S_{\tilde{A}} \simeq D_V / (\square_l)_{l \in \mathbb{L}_{\tilde{A}}}$, so that this D_V -module has an order filtration induced from $F_{\bullet}^{\text{ord}} D_V$. In order to show that $(K^{\bullet}, F_{\bullet}) \rightarrow (M_{\tilde{A}}^{\tilde{\beta}}, F^H)$ is a filtered quasi-isomorphism, it suffices (by

Lemma 3.2) to show that $\mathrm{Gr}_{\bullet}^F K^{\bullet} \rightarrow \mathrm{Gr}_{\bullet}^{F^H} M_{\tilde{A}}^{\tilde{\beta}}$ is a quasi-isomorphism. This follows from [SST00, Formula 4.32, Lemma 4.3.7], as $\mathbb{C}[\mathbb{N}\tilde{A}]$ is Cohen-Macaulay due to the normality assumption on \tilde{A} . The final step is to endow the free resolution $L^{\bullet} = \mathrm{Tot}(K^{\bullet}(E - \beta, D_V \otimes_{\mathbb{C}[\partial_{\lambda}]} T^{\bullet}))$ with a strict filtration F_{\bullet} and to show that $(L_{\bullet}, F_{\bullet}) \xrightarrow{\sim} (M_{\tilde{A}}^{\tilde{\beta}}, F_{\bullet}^H)$. As the resolution $T_{\bullet} \rightarrow S_{\tilde{A}}$ is taken in the category of \mathbb{Z}^{d+1} -graded $\mathbb{C}[\partial]$ -modules, the morphisms of this resolution are homogenous for the (\mathbb{Z}) -grading $\deg(\lambda_i) = -1$ and $\deg(\partial_{\lambda_i}) = 1$ (notice that this is the grading given by the first component of the \mathbb{Z}^{d+1} -grading of the ring $D_V \otimes_{\mathbb{C}[\partial]} S_{\tilde{A}}$). Hence these morphisms are naturally filtered for the order filtration $F_{\bullet}^{ord}(D_V \otimes_{\mathbb{C}[\partial]} S_{\tilde{A}})$ and they are even strict: for a map given by homogenous operators from $\mathbb{C}[\partial]$ taking the symbols has simply no effect, so that $\mathrm{Gr}_{\bullet}^F(D_V \otimes_{\mathbb{C}[\partial]} T_{\bullet}) \rightarrow \mathrm{Gr}_{\bullet}^{F^{ord}}(D_V \otimes_{\mathbb{C}[\partial]} S_{\tilde{A}})$ is a filtered quasi-isomorphism (and similarly for the sums occurring in the terms K^{-l}). However, we have to determine the \mathbb{Z} -degree (for the grading $\deg(\partial_{\lambda_i}) = 1$) of the highest (actually, the only nonzero) cohomology module $\mathrm{Ext}_{\mathbb{C}[\partial]}^{n-d}(S_{\tilde{A}}, \omega_{\mathbb{C}[\partial]})$: it is the first component of the difference of the degree of $\omega_{\mathbb{C}[\partial]}$ (i.e., the first component of the sum of the columns of \tilde{A}), which is $n + 1$, and the first component of the degree of $\omega_{S_{\tilde{A}}}$, which is c_0 . Now the shift of the filtration between $M_{\tilde{A}}^{\tilde{\beta}}$ and the dual module $M_{\tilde{A}}^{-\tilde{\beta}-(c_0, c)}$ is the sum of the length of the complex $K^{\bullet}(E - \beta, D_V \otimes_{\mathbb{C}[\partial]} S_{\tilde{A}})$, i.e., $d + 1$, and the above \mathbb{Z} -degree of $\mathrm{Ext}_{\mathbb{C}[\partial]}^{n-d}(S_{\tilde{A}}, \omega_{\mathbb{C}[\partial]})$, i.e. $n + 1 - c_0$. Hence the filtration $F_{\bullet} L^{-n-1}$ is again the shifted order filtration, more precisely, we have

$$F_p L^{-n-1} = F_{p+d-(d+1)-(n+1-c_0)}^{ord} D_V = F_{p-n-2+c_0}^{ord} D_V.$$

Now it follows from [Sai94, page 55] that

$$\mathbb{D}(M_{\tilde{A}}^{\tilde{\beta}}, F^H) \simeq \mathrm{Hom}_{D_V}((L^{\bullet}, F_{\bullet}), ((D_V \otimes \Omega_V^{n+1})^{\vee}, F_{\bullet, -2(n+1)} D_V \otimes (\Omega_V^{n+1})^{\vee}))$$

so that we finally obtain

$$F_p^H \mathbb{D} M_{\tilde{A}}^{\tilde{\beta}} = F_{p-n-c_0}^{ord} M_{\tilde{A}}^{-\tilde{\beta}-(c_0, c)}.$$

□

We now consider the special case $\beta = 0$. From Proposition 5.2, we know that up to multiplication by a non-zero constant, we have the morphism

$$\begin{aligned} \phi : F_{p+d-c_0}^{ord} M_{\tilde{A}}^{-(c_0, c)} &= F_{p+n+d}^H \mathbb{D}(M_{\tilde{A}}^0) = F_p^H \mathbb{D}(M_{\tilde{A}}^0)(-n-d) \longrightarrow F_p^H M_{\tilde{A}}^0 = F_{p+d}^{ord} M_{\tilde{A}}^0 \\ &P \longmapsto P \cdot \partial^{(c_0, c)} \end{aligned}$$

where $\partial^{(c_0, c)} := \prod_{i=0}^n \partial_{\lambda_i}^{k_i}$ for any $\underline{k} = (k_0, \dots, k_n)$ with $\tilde{A} \cdot \underline{k} = (c_0, c)$. Since \tilde{A} is homogeneous we have $\sum k_i = c_0$. As a consequence, we obtain the following result.

Corollary 5.37. *Under the above assumptions on \tilde{A} , the morphism*

$$\begin{aligned} \phi : (M_{\tilde{A}}^{-(c_0, c)}, F_{\bullet, -c_0}^{ord}) &\longrightarrow (M_{\tilde{A}}^0, F_{\bullet}^{ord}) \\ P &\longmapsto P \cdot \partial^{(c_0, c)} \end{aligned}$$

(where $\partial^{(c_0, c)}$ is as above) is strictly filtered.

Proof. Since both filtered modules $(M_{\tilde{A}}^{-(c_0, c)}, F_{\bullet, -c_0}^{ord})$ and $(M_{\tilde{A}}^0, F_{\bullet}^{ord})$ underly mixed Hodge modules on V by Theorem 5.35 and the morphism is induced from a morphism in $\mathrm{MHM}(V, \mathbb{C})$, we obtain the strictness statement we are looking for. □

Remark 5.38. *If $\mathbb{C}[\mathbb{N}\tilde{A}]$ is not Gorenstein but normal (and therefore Cohen-Macaulay) then the proof of Theorem 5.36 shows that*

$$\mathbb{D} M_{\tilde{A}}^{\tilde{\beta}} \simeq \mathcal{H}^0(E + \beta, D_V \otimes \mathrm{Ext}^{n-d}(S_{\tilde{A}}, \omega_{\mathbb{C}[\partial_{\lambda}]}) \simeq \mathcal{H}^0(E + \beta, D_V \otimes \omega_{S_{\tilde{A}}})$$

Recall that the canonical module $\omega_{S_{\tilde{A}}}$ of $S_{\tilde{A}}$ is isomorphic to $\mathbb{C}[\mathrm{int}(\mathbb{N}\tilde{A})]$ in the category of \mathbb{Z}^{d+1} -graded $\mathbb{C}[\partial_{\lambda}]$ -modules. The module $\omega_{S_{\tilde{A}}}$ carries a \mathbb{Z} -grading given by the first component of the \mathbb{Z}^{d+1} -grading.

Hence $D_V \otimes_{\mathbb{C}[\partial_\lambda]} \omega_{S_{\tilde{A}}}$ carries an order filtration which induces a filtration F_{\bullet}^{ord} on $\mathcal{H}^0(E + \beta, D_V \otimes \omega_{S_{\tilde{A}}})$. We therefore get

$$F_p^H \mathbb{D}\mathcal{M}^{\tilde{\beta}} \simeq F_{p-n-c_0}^{ord} \mathcal{H}^0(E + \beta, D_V \otimes \omega_{S_{\tilde{A}}}).$$

where $c_0 := \min\{\deg_{\mathbb{Z}}(P) \mid P \in \mathbb{C}[\text{int}(\mathbb{N}\tilde{A})]\}$. Let $\tilde{c} \in \text{deg}(\mathbb{C}[\text{int}(\mathbb{N}\tilde{A})])$ with $\tilde{c} = (c_0, c)$. Similar to [Wal07, Proposition 4.4] it can be shown that the inclusion $S_{\tilde{A}}[-\tilde{c}] \hookrightarrow \mathbb{C}[\text{int}(\mathbb{N}\tilde{A})]$ induces an isomorphism $M^{-\tilde{\beta}-\tilde{c}} \xrightarrow{\simeq} \mathcal{H}^0(E + \beta, D_V \otimes \omega_{S_{\tilde{A}}})$ however we do not expect $(M^{-\tilde{\beta}-\tilde{c}}, F_{\bullet}^{ord}) \rightarrow (\mathcal{H}^0(E + \beta, D_V \otimes \omega_{S_{\tilde{A}}}), F_{\bullet}^{ord})$ to be a filtered isomorphism.

5.11 Hodge structures on affine hypersurfaces of tori

In this subsection we explain how our main result implies in a rather direct way a classical theorem of Batyrev concerning the description of the Hodge filtration of the relative cohomology of smooth affine hypersurfaces in algebraic tori.

We first want to recall the sheaf theoretic definition of relative cohomology. Let X be a topological space and K be a closed subset. Denote by $j : X \setminus K \rightarrow X$ the open embedding of the complement. The relative cohomology of the pair (X, K) is defined as the following hypercohomology:

$$H^i(X, K; \mathbb{C}) := \mathbb{H}^i(X, j_* j^{-1} \mathbb{Q}_X)$$

If X and K are quasi-projective varieties the relative cohomology of the pair (X, K) carries a mixed Hodge structure, which is given by $\mathbb{H}^i(X, j_* j^{-1} \mathbb{Q}_X^H)$.

We want to compute this in the following situation: Consider as in section 5 the family of Laurent polynomials $\varphi_A : T \times \Lambda \rightarrow V = \mathbb{C}_{\lambda_0} \times \Lambda$, where $\mathbb{Z}\tilde{A} = \mathbb{Z}^{d+1}$ and $\mathbb{N}\tilde{A} = \mathbb{Z}^{d+1} \cap \mathbb{R}_{\geq 0}\tilde{A}$. Let $\Delta := \text{Conv}(\underline{a}_0, \underline{a}_1, \dots, \underline{a}_n)$ be the convex hull of the exponents of φ_A , where $\underline{a}_0 := 0$. Let $\tau \subset \Delta$ be a face of Δ , $x \in V$ and

$$F_{A,x}^\tau = \sum_{i: \underline{a}_i \in \tau} x_i t^{\underline{a}_i}$$

Definition 5.39. *The fiber $\varphi_A^{-1}(x)$ is non-degenerate if for every face τ of Δ the equations*

$$F_{A,x}^\tau = t_1 \frac{\partial F_{A,x}^\tau}{\partial t_1} = \dots = t_d \frac{\partial F_{A,x}^\tau}{\partial t_d} = 0$$

have no common solution in T .

Let $x \in V$ such that the fiber $\varphi_A^{-1}(x)$ is non-degenerate. We give a model of $H^i(T, \varphi_A^{-1}(x); \mathbb{C})$ as the quotient of a graded semi-group ring and compute explicitly its Hodge filtration. This recovers a result of Stienstra [Sti98, Theorem 7] using results of Batyrev [Bat93].

Lemma 5.40. *Let $x \in V$ and $i_x : \{x\} \rightarrow V$ be the inclusion. Suppose $\varphi_A^{-1}(x)$ is non-degenerate. Then*

1. *the fiber $\varphi_A^{-1}(x)$ is smooth.*
2. *the map i_x is noncharacteristic with respect to \mathcal{M}_A^0 .*

Proof. The first statement follows directly from the definition for $\tau = \Delta$. The second statement follows from [Ado94, Lemma 3.3]. \square

Consider the following diagram (cf. diagram (34))

$$\begin{array}{ccccc}
 Y & \longrightarrow & U & & \\
 \downarrow & & \downarrow j_U & \searrow \pi_2^U & \\
 T \times V & \xrightarrow{g \times id} & \mathbb{P}(W) \times V & \xrightarrow{\pi_2} & V \\
 \uparrow & & \uparrow i_Z & \nearrow \pi_2^Z & \\
 \Gamma & \longrightarrow & Z & &
 \end{array}$$

where Y resp. Γ are the pull-backs such that both squares on the left are cartesian. Notice that Γ is given as a subspace of $T \times V$ by the equation $\lambda_0 + \sum_{i=1}^n \lambda_i t^{a_i} = 0$; hence, Γ is the graph of φ_A and Y is its complement in $T \times V$. Restricting this diagram to some $x \in V$ we therefore get

$$\begin{array}{ccccc}
T \setminus \varphi^{-1}(x) & \xrightarrow{\bar{g}} & U_x & & \\
\bar{j} \downarrow & & \downarrow j & \searrow \pi_x^U & \\
T & \xrightarrow{g} & \mathbb{P}(W) & \xrightarrow{\pi_x} & \{x\} \\
\bar{i} \uparrow & & \uparrow i & \nearrow \pi_x^Z & \\
\varphi^{-1}(x) & \xrightarrow{\tilde{g}} & Z_x & &
\end{array} \tag{54}$$

We will need the following statement

Lemma 5.41. *Let $x \in V$ be such that $\varphi_A^{-1}(x)$ is smooth then we have an isomorphism in $D^b(\text{MHM}(\mathbb{P}(W)))$:*

$$g_* \bar{j}_! \bar{j}^{-1} \mathbb{Q}_T^H \simeq j_! \bar{g}_* \mathbb{Q}_{T \setminus \varphi^{-1}(x)}^H$$

Proof. The statement follows using the following chain of isomorphisms

$$g_* \bar{j}_! \bar{j}^{-1} \mathbb{Q}_T^H \stackrel{!}{\simeq} j_! j^{-1} g_* \mathbb{Q}_T^H \simeq j_! \bar{g}_* \bar{j}^{-1} \mathbb{Q}_T^H \simeq j_! \bar{g}_* \mathbb{Q}_{T \setminus \varphi^{-1}(x)}^H$$

where the second isomorphism follows from base change. It remains to show the first isomorphism. Notice that we have the following triangles

$$\begin{array}{ccc}
j_! j^{-1} g_* & \longrightarrow & g_* \longrightarrow i_! i^* g_* \xrightarrow{+1} \\
g_* \bar{j}_! \bar{j}^{-1} & \longrightarrow & g_* \longrightarrow g_* \bar{i}_! \bar{i}^* \xrightarrow{+1}
\end{array}$$

So it is enough to show $i_! i^* g_* \mathbb{Q}_T^H \simeq g_* \bar{i}_! \bar{i}^* \mathbb{Q}_T^H$. But this can be seen as follows:

$$i_! i^* g_* \mathbb{Q}_T^H \simeq i_! i^! g_* \mathbb{Q}_T^H[2] \simeq i_! \tilde{g}_* \tilde{i}^! \mathbb{Q}_T^H[2] \simeq i_! \tilde{g}_* \tilde{i}^* \mathbb{Q}_T^H \simeq g_* \bar{i}_! \bar{i}^* \mathbb{Q}_T^H$$

where we use the smoothness of $\varphi^{-1}(x)$ in the first and third isomorphism. \square

In order to proof the statement that the restriction of the GKZ-system is isomorphic to a relative cohomology group, we have to rewrite the GKZ-system as a Radon transform. For this, consider the following diagram

$$\begin{array}{ccccc}
T \setminus \varphi^{-1}(x) & \xrightarrow{\bar{g}} & U_x & \xrightarrow{\pi_x^U} & \{x\} \\
\bar{i}_x \downarrow & & \downarrow i_{U_x} & & \downarrow i_x \\
Y & \longrightarrow & U & \xrightarrow{\pi_2^U} & V \\
\pi_1^Y \downarrow & & \downarrow \pi_1^U & & \\
T & \xrightarrow{g} & \mathbb{P}(W) & &
\end{array}$$

where all squares are cartesian.

Proposition 5.42. *Let $x \in V$ be such that $\varphi_A^{-1}(x)$ is non-degenerate, then there is an isomorphism of mixed Hodge structures:*

$$i_x^* \mathcal{M}_A^0 \simeq H^d(T, \varphi_A^{-1}(x); \mathbb{C})$$

and $H^i(T, \varphi_A^{-1}(x); \mathbb{C}) = 0$ for $i \neq d$.

Proof. Consider the following isomorphisms

$$\begin{aligned}
i_x^* \pi_{2!}^U (\pi_1^U)^* g_*^p \mathbb{Q}_T^H &\simeq \pi_{x!}^U i_{U_x}^* (\pi_1^U)^* g_*^p \mathbb{Q}_T^H && \text{base change} \\
&\simeq \pi_{x!}^U (\pi_1^U \circ i_{U_x})^* g_*^p \mathbb{Q}_T^H \\
&\simeq \pi_{x!}^U (\pi_1^U \circ i_{U_x})^! g_*^p \mathbb{Q}_T^H && (\pi_1^U \circ i_{U_x}) \text{ open} \\
&\simeq \pi_{x!}^U \bar{g}_* (\pi_1^Y \circ \bar{i}_x)^! g_*^p \mathbb{Q}_T^H && \text{base change} \\
&\simeq \pi_{x!}^U \bar{g}_* (\pi_1^Y \circ \bar{i}_x)^* g_*^p \mathbb{Q}_T^H && (\pi_1^Y \circ \bar{i}_x) \text{ open} \\
&\simeq \pi_{x!}^U \bar{g}_*^p \mathbb{Q}_{T \setminus \varphi^{-1}(x)}^H
\end{aligned}$$

We can rewrite this further by looking at a part of diagram (54):

$$\begin{array}{ccccc}
T \setminus \varphi^{-1}(x) & \xrightarrow{\bar{g}} & U_x & & \\
\bar{j} \downarrow & & \downarrow j & \searrow \pi_x^U & \\
T & \xrightarrow{g} & \mathbb{P}(W) & \xrightarrow{\pi_x} & \{x\}
\end{array}$$

We have

$$\pi_{x*} g_* \bar{j}_! \bar{j}^{-1} {}^p \mathbb{Q}_T^H \simeq \pi_{x*} j_! \bar{g}_* {}^p \mathbb{Q}_{T \setminus \varphi^{-1}(x)}^H \simeq \pi_{x!} \bar{g}_* {}^p \mathbb{Q}_{T \setminus \varphi^{-1}(x)}^H \simeq i_x^* \pi_{2!}^U (\pi_1^U)^* g_* {}^p \mathbb{Q}_T^H$$

where the last isomorphism follows from the calculation above. If we take cohomology and keep in mind that i_x is non-characteristic we get

$$\begin{aligned}
H^i(T, \varphi_A^{-1}(x); \mathbb{C}) &\simeq H^{i-d}(\pi_{x*} g_* \bar{j}_! \bar{j}^{-1} {}^p \mathbb{Q}_T^H) \\
&\simeq \mathcal{H}^{i-d} i_x^* (\pi_{2!}^U (\pi_1^U)^* g_* {}^p \mathbb{Q}_T^H) \\
&\simeq i_x^* \mathcal{H}^{i-d+n+1} (\pi_{2!}^U (\pi_1^U)^* g_* {}^p \mathbb{Q}_T^H) \\
&\simeq i_x^* \mathcal{H}^{i-d+n+1} (*\mathcal{R}_c^\circ(g_* {}^p \mathbb{Q}_T^H))
\end{aligned}$$

Since $\mathcal{H}^k(*\mathcal{R}_c^\circ(g_* {}^p \mathbb{Q}_T^H)) = 0$ for $k \neq n+1$ and $\mathcal{H}^k(*\mathcal{R}_c^\circ(g_* {}^p \mathbb{Q}_T^H)) = \mathcal{M}_A^0$ for $k = n+1$ the claim follows. \square

Denote by $S_{\tilde{A}} := \mathbb{C}[\mathbb{N}\tilde{A}] \subset \mathbb{C}[u_0^\pm, \dots, u_d^\pm]$ the semigroup ring generated by $\underline{u}^{\tilde{a}_0}, \dots, \underline{u}^{\tilde{a}_n}$ where $\tilde{A} = (\tilde{a}_0, \dots, \tilde{a}_n)$ is the matrix from 33.

Define the following differential operators

$$D_k := \sum_{i=0}^n \left(\tilde{a}_{ki} x_i \underline{u}^{\tilde{a}_i} + u_i \partial_{u_i} \right) \quad \text{for } k = 0, \dots, d \text{ and fixed } x = (x_0, \dots, x_n) \in V$$

which act on $\mathbb{C}[u_0^\pm, \dots, u_d^\pm]$ and which preserve $S_{\tilde{A}}$. For $\underline{u}^{\tilde{l}} = \underline{u}^{l_0 \cdot \tilde{a}_0} \dots \underline{u}^{l_n \cdot \tilde{a}_n} \in S_{\tilde{A}}$ we define the degree $\deg(\underline{u}^{\tilde{l}}) = \sum_{i=0}^n l_i$. Define a descending filtration F^\bullet of \mathbb{C} -vector spaces on $S_{\tilde{A}}$ where $F^{d+1} S_{\tilde{A}} = 0$ and the filtration step $F^{d-k} S_{\tilde{A}}$ is spanned by monomials $\underline{u}^{\tilde{l}}$ with $\deg(\underline{u}^{\tilde{l}}) \leq k$.

Theorem 5.43. *Let $x \in V$ such that $\varphi_A^{-1}(x)$ is non-degenerate, then the following isomorphism of filtered vector spaces holds*

$$(i_x^* \mathcal{M}_{\tilde{A}}^0, F^\bullet) \simeq (S_{\tilde{A}} / (D_k S_{\tilde{A}})_{k=0, \dots, d}, F^{-\bullet})$$

Proof. Since we have assumed that i_x is non-characteristic with respect to \mathcal{M}_A^0 the only non-zero cohomology group of $i_x^* \mathcal{M}_{\tilde{A}}^0$ is

$$\mathcal{H}^0 i_x^* \mathcal{M}_{\tilde{A}}^0 \simeq (\lambda_i - x_i)_{i=0, \dots, n} \setminus \mathcal{M}_{\tilde{A}}^0$$

We define a \mathbb{C} -linear map

$$\begin{aligned}
\Psi' : S_{\tilde{A}} &\longrightarrow (\lambda_i - x_i)_{i=0, \dots, n} \setminus \mathcal{M}_{\tilde{A}}^0 \\
\underline{u}^{l_0 \cdot \tilde{a}_0} \dots \underline{u}^{l_n \cdot \tilde{a}_n} &\mapsto \partial_{\lambda_0}^{l_0} \dots \partial_{\lambda_n}^{l_n}
\end{aligned}$$

We want to show that this map factors over $S_{\tilde{A}} / (D_k S_{\tilde{A}})_{k=0, \dots, d}$ so that Ψ' descends to a map

$$\Psi : S_{\tilde{A}} / (D_k S_{\tilde{A}})_{k=0, \dots, d} \longrightarrow (\lambda_i - x_i)_{i=0, \dots, n} \setminus \mathcal{M}_{\tilde{A}}^0$$

Let $P = \underline{u}^{l_0 \cdot \tilde{a}_0} \dots \underline{u}^{l_n \cdot \tilde{a}_n}$, then $\Psi'(P) = \partial_{\lambda_0}^{l_0} \dots \partial_{\lambda_n}^{l_n}$ and

$$\begin{aligned}
\Psi'(D_k P) &= \Psi\left(\sum_{i=0}^n \tilde{a}_{ki} x_i \underline{u}^{a_i} P + \sum_{i=0}^n \tilde{a}_{ki} l_i P\right) \\
&= \left(\sum_{i=0}^n \tilde{a}_{ki} x_i \partial_{\lambda_i} + \sum_{i=0}^n \tilde{a}_{ki} l_i\right) \Psi(P) \\
&= \left(\sum_{i=0}^n \tilde{a}_{ki} \lambda_i \partial_{\lambda_i} + \sum_{i=0}^n \tilde{a}_{ki} l_i\right) \Psi(P) \\
&= \Psi(P) \cdot \left(\sum_{i=0}^n \tilde{a}_{ki} \lambda_i \partial_{\lambda_i}\right) \\
&= 0
\end{aligned}$$

We will now construct an inverse Θ to Ψ . If $P \in D_V$ is a normally ordered element, we denote by $\bar{P} \in S_{\tilde{A}}$ the element which is obtained from P by replacing λ_i with x_i and ∂_{λ_i} with $\underline{u}^{l_i a_i}$, i.e. if $P = \lambda_0^{k_0} \dots \lambda_n^{k_n} \partial_{\lambda_0}^{l_0} \dots \partial_{\lambda_n}^{l_n}$ the element \bar{P} is given by $x_0^{k_0} \dots x_n^{k_n} \underline{u}^{l_0 \cdot a_0} \dots \underline{u}^{l_n \cdot a_n}$. This gives the map

$$\begin{aligned}
\Theta' : (\lambda_i - x_i)_{i=0, \dots, n} \setminus D_V &\longrightarrow S_{\tilde{A}} / (D_k S_{\tilde{A}})_{k=0, \dots, d} \\
P &\mapsto \bar{P}
\end{aligned}$$

We want to show that Θ' factors over $(\lambda_i - x_i)_{i=0, \dots, n} \setminus M_A^0$ so that Θ' descends to a map

$$\begin{aligned}
\Theta : (\lambda_i - x_i)_{i=0, \dots, n} \setminus M_A^0 &\longrightarrow S_{\tilde{A}} / (D_k S_{\tilde{A}})_{k=0, \dots, d} \\
P &\mapsto \bar{P}
\end{aligned}$$

We have to show that $\Theta'(P \cdot E_k) = 0$ and $\Theta'(P \cdot \square_l) = 0$ for $k = 0, \dots, d$ and $l \in \mathbb{L}$. We can assume that P is a monomial $\lambda_0^{j_0} \dots \lambda_n^{j_n} \partial_{\lambda_0}^{l_0} \dots \partial_{\lambda_n}^{l_n}$:

$$\begin{aligned}
\Theta'(\lambda_0^{j_0} \dots \lambda_n^{j_n} \partial_{\lambda_0}^{l_0} \dots \partial_{\lambda_n}^{l_n} E_k) &= \Theta'(\lambda_0^{j_0} \dots \lambda_n^{j_n} \partial_{\lambda_0}^{l_0} \dots \partial_{\lambda_n}^{l_n} \left(\sum_{i=0}^n \tilde{a}_{ki} \lambda_i \partial_{\lambda_i}\right)) \\
&= \Theta'(\lambda_0^{j_0} \dots \lambda_n^{j_n} \left(\sum_{i=0}^n (\tilde{a}_{ki} \lambda_i \partial_{\lambda_i} + \tilde{a}_{ki} l_i)\right) \partial_{\lambda_0}^{l_0} \dots \partial_{\lambda_n}^{l_n}) \\
&= x_0^{j_0} \dots x_n^{j_n} \left(\sum_{i=0}^n (\tilde{a}_{ki} x_i \underline{u}^{a_i} + \tilde{a}_{ki} l_i)\right) \underline{u}^{l_0 \cdot a_0} \dots \underline{u}^{l_n \cdot a_n} \\
&= \left(\sum_{i=0}^n (\tilde{a}_{ki} x_i \underline{u}^{a_i} + \tilde{a}_{ki} l_i)\right) \cdot x_0^{j_0} \dots x_n^{j_n} \cdot \underline{u}^{l_0 \cdot a_0} \dots \underline{u}^{l_n \cdot a_n} \\
&= \left(\sum_{i=0}^n (\tilde{a}_{ki} x_i \underline{u}^{a_i} + u_i \partial_{u_i})\right) \cdot x_0^{j_0} \dots x_n^{j_n} \cdot \underline{u}^{l_0 \cdot a_0} \dots \underline{u}^{l_n \cdot a_n} \\
&= 0
\end{aligned}$$

and

$$\begin{aligned}
\Theta'(\lambda_0^{j_0} \dots \lambda_n^{j_n} \partial_{\lambda_0}^{l_0} \dots \partial_{\lambda_n}^{l_n} \square_l) &= \Theta'(\lambda_0^{j_0} \dots \lambda_n^{j_n} \partial_{\lambda_0}^{l_0} \dots \partial_{\lambda_n}^{l_n} \left(\prod_{l_i > 0} \partial_{\lambda_i}^{l_i} - \prod_{l_i < 0} \partial_{\lambda_i}^{-l_i}\right)) \\
&= x_0^{j_0} \dots x_n^{j_n} \underline{u}^{l_0 \cdot a_0} \dots \underline{u}^{l_n \cdot a_n} \left(\prod_{l_i > 0} \underline{u}^{l_i \cdot a_i} - \prod_{l_i < 0} \underline{u}^{-l_i \cdot a_i}\right) \\
&= x_0^{j_0} \dots x_n^{j_n} \underline{u}^{l_0 \cdot a_0} \dots \underline{u}^{l_n \cdot a_n} \left(\underline{u}^{\sum_{l_i > 0} l_i \cdot a_i} - \underline{u}^{-\sum_{l_i < 0} l_i \cdot a_i}\right) \\
&= 0
\end{aligned}$$

where the last equality follows from $\sum_{i=0}^n l_i \mathbf{q}_i = 0$. This shows that Ψ is an isomorphism. The statement about the Hodge filtration follows from Theorem 5.35, the fact that x is a smooth point of $\mathcal{M}_{\tilde{A}}^0$ and the definition of the inverse image of filtered \mathcal{D} -modules (cf. [Sai88] chapter 3.5, notice that no shift in the Hodge filtration occurs since we are dealing with left \mathcal{D} -modules instead of right \mathcal{D} -modules as in loc. cit.). \square

6 Landau-Ginzburg models and non-commutative Hodge structures

In this final section we will give a first application of our main result. It is concerned with Hodge-theoretic properties of differential systems occurring in toric mirror symmetry. More precisely, we will prove [RS17, Conjecture 6.15] showing that the so-called *reduced quantum \mathcal{D} -module* of a nef (also called weak Fano) complete intersection inside a smooth projective toric variety underlies a (variation of) non-commutative Hodge structure(s). We will recall the necessary notation and results of loc.cit. and then deduce this conjecture from our main Theorem 5.35. The basic strategy to obtain the proof of the conjecture is to identify the reduced quantum \mathcal{D} -module with an object which is the Fourier-Laplace transform of a filtered \mathcal{D} -module underlying a pure polarized Hodge module. The latter is nothing but the image of the duality morphism from Corollary 5.37. Notice that this corollary depends in an essential way on our main Theorem 5.35, since the strictness of the duality morphism with respect to the order filtrations holds only because the latter are (up to a shift) the Hodge filtrations of mixed Hodge modules. The identification with the reduced quantum \mathcal{D} -module relies on the explicit description of the latter from [MM17] (already used extensively in [RS17]).

Let X_Σ be a smooth, projective and toric variety with $\dim_{\mathbb{C}}(X_\Sigma) = k$. Put $m := k + b_2(X_\Sigma)$. Let $\mathcal{L}_1, \dots, \mathcal{L}_l$ be globally generated line bundles on X_Σ (in particular, they are nef according to [Ful93, Section 3.4]) and assume that $-K_{X_\Sigma} - \sum_{i=1}^l c_1(\mathcal{L}_i)$ is nef. Put $\mathcal{E} := \oplus_{i=1}^l \mathcal{L}_i$, and let \mathcal{E}^\vee be the dual vector bundle. Its total space $\mathbb{V}(\mathcal{E}^\vee) := \mathbf{Spec}_{\mathcal{O}_{X_\Sigma}}(\mathrm{Sym}_{\mathcal{O}_{X_\Sigma}}(\mathcal{E}))$ is a quasi-projective toric variety with defining fan Σ' . The matrix $A \in M((k+l) \times (m+l), \mathbb{Z})$ whose columns are the primitive integral generators of the rays of Σ' then satisfies the conditions in Theorem 5.36. More precisely, we have $\mathbb{Z}\tilde{A} = \mathbb{Z}^{d+1}$ and it follows from [RS17, Proposition 5.1] that the semi-group $\mathbb{N}\tilde{A}$ is normal and that we have $\mathrm{int}(\mathbb{N}\tilde{A}) = \tilde{c} + \mathbb{N}\tilde{A}$, where $\tilde{c} = \sum_{i=m+1}^{m+l} e_i = (l+1, \mathbf{0}, \mathbf{1})$, e_i being the i 'th standard vector in \mathbb{Z}^{1+m+l} .

The strictly filtered duality morphism ϕ from Corollary 5.37 is more concretely given as

$$\begin{aligned} \phi : (\mathcal{M}_{\tilde{A}}^{-(l+1, \mathbf{0}, \mathbf{1})}, F_{\bullet-l-1}^{\mathrm{ord}}) &\longrightarrow (\mathcal{M}_{\tilde{A}}^0, F_{\bullet}^{\mathrm{ord}}) \\ P &\longmapsto P \cdot \partial_{\lambda_0} \cdot \partial_{\lambda_{m+1}} \cdots \partial_{\lambda_{m+l}}. \end{aligned}$$

Proposition 6.1. *The image of ϕ underlies a pure Hodge module of weight $m+k+2l$, where the Hodge filtration is given by*

$$F_{\bullet}^H \mathrm{im}(\phi) = \mathrm{im}(\phi) \cap F_{\bullet+k+l}^{\mathrm{ord}} \mathcal{M}_{\tilde{A}}^0.$$

Proof. This is a consequence of [RS17, Theorem 2.16] and of Proposition 5.2. \square

A main point in the paper [RS17] is to consider the partial localized Fourier transformations of the GKZ-systems $\mathcal{M}_{\tilde{A}}^\beta$. We recall the main construction and refer to [RS17, Section 3.1] for details (in particular concerning the definition and properties of the Fourier-Laplace functor FL and its ‘‘localized’’ version $\mathrm{FL}^{\mathrm{loc}}$). Let (as done already in section in 5.1) Λ be the affine space \mathbb{C}^{m+l} with coordinates $\lambda_1, \dots, \lambda_{m+l}$ (so that $V = \mathbb{C}_{\lambda_0} \times \Lambda$) and put $\widehat{V} := \mathbb{C}_z \times \Lambda$. Let $\widehat{\mathcal{M}}_A^{(\beta_0, \beta)}$ be the $\mathcal{D}_{\widehat{V}}$ -module $\mathcal{D}_{\widehat{V}}[z^{-1}]/\mathcal{I}$, where \mathcal{I} is the left ideal generated by the operators $\widehat{\square}_l$ (for all $l \in \mathbb{L}_A$), $\widehat{E}_j - \beta_j z$ (for $j = 1, \dots, k+l$) and $\widehat{E} - \beta_0 z$, which are defined by

$$\begin{aligned} \widehat{\square}_l &:= \prod_{i: l_i < 0} (z \cdot \partial_{\lambda_i})^{-l_i} - \prod_{i: l_i > 0} (z \cdot \partial_{\lambda_i})^{l_i}, \\ \widehat{E} &:= z^2 \partial_z + \sum_{i=1}^{m+l} z \lambda_i \partial_{\lambda_i}, \\ \widehat{E}_j &:= \sum_{i=1}^{m+l} a_{ji} z \lambda_i \partial_{\lambda_i}. \end{aligned}$$

We denote the corresponding $\mathcal{D}_{\widehat{\nu}}$ -module by $\widehat{\mathcal{M}}_A^{(\beta_0, \beta)}$. Then we have (see [RS17, Lemma 3.2])

$$\mathrm{FL}_\Lambda^{\mathrm{loc}} \left(\mathcal{M}_A^{(\beta_0, \beta)} \right) = \widehat{\mathcal{M}}_A^{(\beta_0+1, \beta)}.$$

Consider the filtration on $\mathcal{D}_{\widehat{\nu}}$ for which z has degree -1 , ∂_z has degree 2 and $\deg(\lambda_i) = 0$, $\deg(\partial_{\lambda_i}) = 1$. Write $\mathrm{MF}^z(\mathcal{D}_{\widehat{\nu}})$ for the category of well-filtered $\mathcal{D}_{\widehat{\nu}}$ -modules (that is, $\mathcal{D}_{\widehat{\nu}}$ -modules equipped with a filtration compatible with the filtration on $\mathcal{D}_{\widehat{\nu}}$ just described and such that the corresponding Rees module is coherent over the corresponding Rees ring). Denote by G_\bullet the induced filtrations on the module $\widehat{\mathcal{M}}_A^{(\beta_0, \beta)}$, which are $\mathcal{R}_{\mathbb{C}_z \times \Lambda}$ -modules. We have

$$G_0 \widehat{\mathcal{M}}_A^{(\beta_0, \beta)} = \mathcal{R}_{\mathbb{C}_z \times \Lambda} / \mathcal{R}_{\mathbb{C}_z \times \Lambda}(\widehat{\square}_l)_{l \in \mathbb{L}_A} + \mathcal{R}_{\mathbb{C}_z \times \Lambda} \widehat{E} + \mathcal{R}_{\mathbb{C}_z \times \Lambda}(\widehat{E}_j)_{k=1, \dots, k+l}$$

and $G_k \widehat{\mathcal{M}}_A^{(\beta_0, \beta)} = z^k \cdot G_0 \widehat{\mathcal{M}}_A^{(\beta_0, \beta)}$. In general, the modules $\widehat{\mathcal{M}}_A^{(\beta_0, \beta)}$ and their filtration steps may be quite complicated. However, we have considered in [RS17] their restriction to a specific Zariski open subset $\Lambda^\circ \subset \left(\Lambda \setminus \bigcup_{i=1}^{m+l} \{w_i = 0\} \right) \subset \Lambda$ (called W° in [RS17, Remark 3.8]), which contains the critical locus of the family of Laurent polynomials associated with the matrix A (but excludes certain singularities at infinity of this family). Denote by ${}^\circ \widehat{\mathcal{M}}_A^{(\beta_0, \beta)}$ the restriction $(\widehat{\mathcal{M}}_A^{(\beta_0, \beta)})|_{\mathbb{C}_z \times \Lambda^\circ}$ together with the induced filtration $G_\bullet {}^\circ \widehat{\mathcal{M}}_A^{(\beta_0, \beta)}$. Then $G_k {}^\circ \widehat{\mathcal{M}}_A^{(\beta_0, \beta)}$ is $\mathcal{O}_{\mathbb{C}_z \times \Lambda^\circ}$ -locally free for all k . Moreover, the multiplication by z is invertible on ${}^\circ \widehat{\mathcal{M}}_A^{(\beta_0, \beta)}$, filtered with respect to G_\bullet (shifting the filtration by one) and so is its inverse. Hence, we have a strict morphism

$$\cdot z : ({}^\circ \widehat{\mathcal{M}}_A^{(\beta_0, \beta)}, G_\bullet) \longrightarrow ({}^\circ \widehat{\mathcal{M}}_A^{(\beta_0-1, \beta)}, G_{\bullet+1}).$$

We also need a slightly modified version of the Fourier-Laplace transformed GKZ-systems. More precisely, define the modules ${}^\circ \widehat{\mathcal{N}}_A^\beta$ as the cyclic quotients of $\mathcal{D}_{\mathbb{C}_z \times \Lambda^\circ}[z^{-1}]$ by the left ideal generated by $\widehat{\square}_l$ for $l \in \mathbb{L}_A$ and $\widehat{E}_j - z\beta_j$ for $j = 0, \dots, k+c$, where

$$\begin{aligned} \widehat{\square}_l &:= \prod_{i \in \{1, \dots, m\}: l_i > 0} \lambda_i^{l_i} (z \cdot \partial_i)^{l_i} \prod_{i \in \{m+1, \dots, m+l\}: l_i > 0} \prod_{\nu=1}^{l_i} (\lambda_i (z \cdot \partial_i) - z \cdot \nu) \\ &\quad - \prod_{i=1}^{m+l} \lambda_i^{l_i} \cdot \prod_{i \in \{1, \dots, m\}: l_i < 0} \lambda_i^{-l_i} (z \cdot \partial_i)^{-l_i} \prod_{i \in \{m+1, \dots, m+l\}: l_i < 0} \prod_{\nu=1}^{-l_i} (\lambda_i (z \cdot \partial_i) - z \cdot \nu). \end{aligned}$$

Consider the *invertible* morphism

$$\Psi : {}^\circ \widehat{\mathcal{N}}_A^{(0, \mathbf{0}, \mathbf{0})} \longrightarrow {}^\circ \widehat{\mathcal{M}}_A^{-(2l, \mathbf{0}, \mathbf{1})} \quad (55)$$

given by right multiplication with $z^l \cdot \prod_{i=m+1}^{m+l} \lambda_i$ (recall that $\lambda_i \neq 0$ on Λ°). We define $\widetilde{\phi}$ to be the composition $\widetilde{\phi} := \widehat{\phi} \circ \Psi$, where $\widehat{\phi}$ is the morphism

$$\widehat{\phi} : {}^\circ \widehat{\mathcal{M}}_A^{-(2l, \mathbf{0}, \mathbf{1})} \longrightarrow {}^\circ \widehat{\mathcal{M}}_A^{(-l, \mathbf{0}, \mathbf{0})},$$

given by right multiplication with $\partial_{\lambda_{m+1}} \cdots \partial_{\lambda_{m+l}}$. In concrete terms, we have:

$$\begin{aligned} \widetilde{\phi} : {}^\circ \widehat{\mathcal{N}}_A^{(0, \mathbf{0}, \mathbf{0})} &\longrightarrow {}^\circ \widehat{\mathcal{M}}_A^{(-l, \mathbf{0}, \mathbf{0})}, \\ x &\longmapsto \widehat{\phi}(x \cdot z^l \cdot \lambda_{m+1} \cdots \lambda_{m+l}) = x \cdot (z\lambda_{m+1}\partial_{m+1}) \cdots (z\lambda_{m+l}\partial_{m+l}). \end{aligned}$$

We have an induced filtration $G_\bullet {}^\circ \widehat{\mathcal{N}}_A^{(0, \mathbf{0}, \mathbf{0})}$ which satisfies

$$G_0 {}^\circ \widehat{\mathcal{N}}_A^{(0, \mathbf{0}, \mathbf{0})} = \mathcal{R}_{\mathbb{C}_z \times \Lambda^\circ} / \mathcal{R}_{\mathbb{C}_z \times \Lambda^\circ}(\widetilde{\square}_l)_{l \in \mathbb{L}_A} + \mathcal{R}_{\mathbb{C}_z \times \Lambda^\circ}(\widehat{E}_j - z\beta_j)_{j=0, \dots, m+l}$$

and $G_k {}^\circ \widehat{\mathcal{N}}_A^{(0, \mathbf{0}, \mathbf{0})} = z^k \cdot G_0 {}^\circ \widehat{\mathcal{N}}_A^{(0, \mathbf{0}, \mathbf{0})}$

In order to obtain the lattices G_\bullet we need to extend the functor $\mathrm{FL}_\Lambda^{\mathrm{loc}}$ to the category of filtered \mathcal{D} -modules.

Definition 6.2. Let $(\mathcal{M}, F_\bullet) \in \text{MF}(\mathcal{D}_V) = \text{MF}(\mathcal{D}_{\mathbb{C}_{\lambda_0} \times \Lambda})$. Define $\mathcal{M}[\partial_{\lambda_0}^{-1}] := \mathcal{D}_V[\partial_{\lambda_0}^{-1}] \otimes_{\mathcal{D}_V} \mathcal{M}$ and consider the natural localization morphism $\widehat{\text{loc}} : \mathcal{M} \rightarrow \mathcal{M}[\partial_{\lambda_0}^{-1}]$. We define the saturation of F_\bullet to be

$$F_k \mathcal{M}[\partial_{\lambda_0}^{-1}] := \sum_{j \geq 0} \partial_{\lambda_0}^{-j} \widehat{\text{loc}}(F_{k+j} \mathcal{M}). \quad (56)$$

and we denote by $G_\bullet \widehat{\mathcal{M}}$ the filtration induced from $F_k \mathcal{M}[\partial_{\lambda_0}^{-1}]$ on $\widehat{\mathcal{M}} := \text{FL}_\Lambda^{\text{loc}}(M) \in M_h(\mathcal{D}_{\widehat{V}}) = M_h(\mathcal{D}_{\mathbb{C}_z \times \Lambda})$. Notice that for $(\mathcal{M}, F_\bullet) = (\widehat{\mathcal{M}}_A^{(\beta_0, \beta)}, F_\bullet^{\text{ord}})$, the two definitions of G_\bullet coincide: As we have

$$F_k^{\text{ord}} \mathcal{M}_A^{(\beta_0-1, \beta)}[\partial_{\lambda_0}^{-1}] = \text{im}(\partial_{\lambda_0}^k \mathbb{C}[\lambda_0, \lambda_1, \dots, \lambda_{m+l}] \langle \partial_{\lambda_0}^{-1}, \partial_{\lambda_0}^{-1} \partial_{\lambda_1}, \dots, \partial_{\lambda_0}^{-1} \partial_{\lambda_{m+l}} \rangle) \text{ in } \mathcal{M}_A^{(\beta_0-1, \beta)}[\partial_{\lambda_0}^{-1}],$$

the filtration induced by $F_k^{\text{ord}} \mathcal{M}_A^{(\beta_0-1, \beta)}[\partial_{\lambda_0}^{-1}]$ on $\widehat{\mathcal{M}}_A^{(\beta_0, \beta)}$ is precisely $G_k \widehat{\mathcal{M}}_A^{(\beta_0, \beta)}$.

We denote by $(\text{FL}_\Lambda^{\text{loc}}, \text{Sat})$ the induced functor from the category $\text{MF}(\mathcal{D}_V)$ to the category $\text{MF}^z(\mathcal{D}_{\widehat{\Lambda}})$ which sends (\mathcal{M}, F_\bullet) to $(\widehat{\mathcal{M}}, G_\bullet)$.

From the above duality considerations, we deduce the following result.

Proposition 6.3. *The morphism*

$$\widetilde{\phi} : {}^\circ \widehat{\mathcal{N}}_A^{(0,0,0)} \longrightarrow {}^\circ \widehat{\mathcal{M}}_A^{(-l,0,0)}$$

is strict with respect to the filtration G_\bullet , in particular, we have

$$\widetilde{\phi} \left(G_0 {}^\circ \widehat{\mathcal{N}}_A^{(0,0,0)} \right) = G_0 {}^\circ \widehat{\mathcal{M}}_A^{(-l,0,0)} \cap \text{im}(\widetilde{\phi})$$

Moreover, the object $(\text{im}(\widetilde{\phi}), G_\bullet)$ is obtained via the functor $(\text{FL}_\Lambda^{\text{loc}}, \text{Sat})$ from $(\text{im}(\phi), F_\bullet^H = F_{\bullet+k+l}^{\text{ord}})$, which underlies a pure Hodge module of weight $m+k+2l$ by Proposition 6.1.

Proof. The morphism Ψ is invertible, filtered (shifting the filtration by $-l$) and its inverse is also filtered. Hence it is strict. Therefore the strictness of $\widetilde{\phi}$ follows from the strictness of $z\widehat{\phi}$. We will deduce it from the strictness property of the morphism ϕ in Corollary 5.37.

Notice that the morphism $\widehat{\phi}$ is obtained from ϕ by linear extension in $\partial_{\lambda_0}^{-1}$. Recall that the morphism

$$\phi : (\mathcal{M}_{\widehat{A}}^{-(l+1,0,1)}, F_\bullet^{\text{ord}}) \longrightarrow (\mathcal{M}_{\widehat{A}}^0, F_{\bullet+l+1}^{\text{ord}})$$

was strict, hence equation (56) yields the strictness of

$$\widehat{\phi} : (\widehat{\mathcal{M}}_A^{-(2l,0,1)}, G_\bullet) \longrightarrow (\widehat{\mathcal{M}}_A^{(-l,0,0)}, G_{\bullet+l})$$

Finally, as already noticed above, this yields the strictness of

$$\widetilde{\phi} = \widehat{\phi} \circ \Psi : (\widehat{\mathcal{N}}_A^{(0,0,0)}, G_\bullet) \longrightarrow (\widehat{\mathcal{M}}_A^{(-l,0,0)}, G_\bullet).$$

□

The next corollary is now a direct consequence of [Sab08, Corollary 3.15].

Corollary 6.4. *The free $\mathcal{O}_{\mathbb{C}_z \times \Lambda^\circ}$ -module $G_0 {}^\circ \widehat{\mathcal{M}}_A^{(-l,0,0)} \cap \text{im}(\widetilde{\phi})$ underlies a variation of pure polarized non-commutative Hodge structures on Λ° (see [Sab11] for a detailed discussion of this notion).*

The main result in [RS17] concerns a mirror statement for several quantum \mathcal{D} -modules which are associated with the toric variety X_Σ and the split vector bundle \mathcal{E} . In particular, one can consider the reduced quantum \mathcal{D} -module $\overline{\text{QDM}}(X_\Sigma, \mathcal{E})$ which is a vector bundle on $\mathbb{C}_z \times H^0(X_\Sigma, \mathbb{C}) \times B_\varepsilon^*$, where $B_\varepsilon^* := \{q \in (\mathbb{C}^*)^{b_2(X_\Sigma)}, |0 < |q| < \varepsilon\}$ together with a flat connection

$$\nabla : \overline{\text{QDM}}(X_\Sigma, \mathcal{E}) \rightarrow \overline{\text{QDM}}(X_\Sigma, \mathcal{E}) \otimes_{\mathcal{O}_{\mathbb{C}_z \times H^0(X_\Sigma, \mathbb{C}) \times B_\varepsilon^*}} z^{-1} \Omega_{\mathbb{C}_z \times H^0(X_\Sigma, \mathbb{C}) \times B_\varepsilon^*}^1 (\log(\{0\} \times H^0(X_\Sigma, \mathbb{C}) \times B_\varepsilon^*)).$$

We refer to [MM17] for a detailed discussion of the definition of $\overline{\text{QDM}}(X_\Sigma, \mathcal{E})$, a short version can be found in [RS17, Section 4.1]. Notice that in loc.cit., $\overline{\text{QDM}}(X_\Sigma, \mathcal{E})$ is defined on some larger set, but in mirror type statements only its restriction to $H^0(X_\Sigma, \mathbb{C}) \times \mathbb{C}_z \times B_\varepsilon^*$ is considered. In the sequel, we will need to consider a Zariski open subset of $\mathcal{KM}^\circ \subset (\mathbb{C}^*)^{b_2(X_\Sigma)}$ which contains B_ε^* . We recall the main result from [MM17], which gives a GKZ-type description of $\overline{\text{QDM}}(X_\Sigma, \mathcal{E})$. We present it in a slightly different form, taking into account [RS17, Proposition 6.9]. Let $\mathcal{R}_{\mathbb{C}_z \times \mathcal{KM}^\circ}$ be the sheaf of Rees rings on $\mathbb{C}_z \times \mathcal{KM}^\circ$, and $R_{\mathbb{C}_z \times \mathcal{KM}^\circ}$ its module of global sections. If we write q_1, \dots, q_r for the coordinates on $(\mathbb{C}^*)^r$ (with $r := b_2(X_\Sigma)$), then $R_{\mathbb{C}_z \times \mathcal{KM}^\circ}$ is generated by $zq_i \partial_{q_i}$ and $z^2 \partial_z$ over $\mathcal{O}_{\mathbb{C}_z \times \mathcal{KM}^\circ}$.

Theorem 6.5. *For any $\mathcal{L} \in \text{Pic}(X_\Sigma)$, write $\widehat{\mathcal{L}} \in R_{\mathbb{C}_z \times \mathcal{KM}^\circ}$ for the associated “quantized operator” as defined in [MM17, Notation 4.2.] or [RS17, Theorem 6.7]. Define the left ideal J of $R_{\mathbb{C}_z \times \mathcal{KM}^\circ}$ by*

$$J := R_{\mathbb{C}_z \times \mathcal{KM}^\circ}(Q_L)_{l \in \mathbb{L}_{A'}} + R_{\mathbb{C}_z \times \mathcal{KM}^\circ} \cdot \widehat{E},$$

where

$$\begin{aligned} Q_L &:= \prod_{i \in \{1, \dots, m\}: l_i > 0} \prod_{\nu=0}^{l_i-1} (\widehat{\mathcal{D}}_i - \nu z) \prod_{j \in \{1, \dots, c\}: l_{m+j} > 0} \prod_{\nu=1}^{l_{m+j}} (\widehat{\mathcal{L}}_j + \nu z) \\ &- \frac{q^L}{z} \cdot \prod_{i \in \{1, \dots, m\}: l_i < 0} \prod_{\nu=0}^{-l_i-1} (\widehat{\mathcal{D}}_i - \nu z) \prod_{j \in \{1, \dots, c\}: l_{m+j} < 0} \prod_{\nu=1}^{-l_{m+j}} (\widehat{\mathcal{L}}_j + \nu z), \\ \widehat{E} &:= z^2 \partial_z - \widehat{K}_{\mathbb{V}(\mathcal{E}^\vee)}. \end{aligned}$$

Here we write $\mathcal{D}_i \in \text{Pic}(X_\Sigma)$ for a line bundle associated with the torus invariant divisor D_i , where $i = 1, \dots, m$. Let $K \subset R_{\mathbb{C}_z \times \mathcal{KM}^\circ}$ be the ideal

$$K := \left\{ P \in R_{\mathbb{C}_z \times \mathcal{KM}^\circ} \mid \exists p \in \mathbb{Z}, k \in \mathbb{N} : \prod_{i=0}^k \prod_{j=1}^c (\widehat{\mathcal{L}} + p + i) P \in J \right\}$$

and \mathcal{K} the associated sheaf of ideals in $\mathcal{R}_{\mathbb{C}_z \times \mathcal{KM}^\circ}$.

Suppose as above that the bundle $-K_{X_\Sigma} - \sum_{j=1}^l \mathcal{L}_j$ is nef, and moreover that each individual bundle \mathcal{L}_j is ample. Then there is a map $\text{Mir} : B_\varepsilon^* \rightarrow H^0(X_\Sigma, \mathbb{C}) \times B_\varepsilon^*$ such that we have an isomorphism of $\mathcal{R}_{\mathbb{C}_z \times B_\varepsilon^*}$ -modules

$$(\mathcal{R}_{\mathbb{C}_z \times \mathcal{KM}^\circ} / \mathcal{K})|_{\mathbb{C}_z \times B_\varepsilon^*} \xrightarrow{\cong} (\text{id}_{\mathbb{C}_z} \times \text{Mir})^* \overline{\text{QDM}}(X_\Sigma, \mathcal{E}).$$

In order to relate the quantum \mathcal{D} -module $\overline{\text{QDM}}(X_\Sigma, \mathcal{E})$ with our results on GKZ-systems, we will use the restriction map $\bar{\rho} : \mathcal{KM}^\circ \hookrightarrow \Lambda$ as constructed in [RS17] (discussion before Definition 6.3. in loc.cit.). Then it follows from the results of loc.cit., Proposition 6.10, that we have an isomorphism of $\mathcal{R}_{\mathbb{C}_z \times \mathcal{KM}^\circ}$ -modules

$$\mathcal{R}_{\mathbb{C}_z \times \mathcal{KM}^\circ} / \mathcal{K} \cong (\text{id}_{\mathbb{C}_z} \times \bar{\rho})^* \left(\widetilde{\phi} \left(G_0 \widehat{\mathcal{N}}_A^{(0,0,0)} \right) \right)$$

Now we can deduce from Corollary 6.4 the main result of this section.

Theorem 6.6. *Consider the above situation of a k -dimensional toric variety X_Σ , globally generated line bundles $\mathcal{L}_1, \dots, \mathcal{L}_l$ such that $-K_{X_\Sigma} - \mathcal{E}$ is nef, where $\mathcal{E} = \bigoplus_{j=1}^l \mathcal{L}_j$, with \mathcal{L}_j ample for $j = 1, \dots, l$. Then the smooth $\mathcal{R}_{\mathbb{C}_z \times \mathcal{KM}^\circ}$ -module $(\text{id}_{\mathbb{C}_z} \times \text{Mir})^* \overline{\text{QDM}}(X_\Sigma, \mathcal{E})$ (i.e., the vector bundle over $\mathbb{C}_z \times \mathcal{KM}^\circ$ together with its connection operator ∇) underlies a variation of pure polarized non-commutative Hodge structures.*

Proof. The strictness of $\widetilde{\phi}$ as shown in Proposition 6.3 shows that $G_0 \widehat{\mathcal{M}}_A^{(-l,0,0)} \cap \text{im}(\widetilde{\phi}) = \widetilde{\phi}(G_0 \mathcal{N}^{(0,0,0)})$, hence, by Corollary 6.4, the module $\widetilde{\phi}(G_0 \mathcal{N}^{(0,0,0)})$ underlies a variation of pure polarized non-commutative Hodge structures on Λ^0 . Hence the assertion follows from the mirror statement of Theorem 6.5. \square

References

- [Ado94] Alan Adolphson, *Hypergeometric functions and rings generated by monomials*, Duke Math. J. **73** (1994), no. 2, 269–290.
- [Bat93] Victor V. Batyrev, *Variations of the mixed Hodge structure of affine hypersurfaces in algebraic tori*, Duke Math. J. **69** (1993), no. 2, 349–409.
- [BH93] Winfried Bruns and Jürgen Herzog, *Cohen-Macaulay rings*, Cambridge Studies in Advanced Mathematics, vol. 39, Cambridge University Press, Cambridge, 1993.
- [Bry86] Jean-Luc Brylinski, *Transformations canoniques, dualité projective, théorie de Lefschetz, transformations de Fourier et sommes trigonométriques*, Astérisque (1986), no. 140-141, 3–134, 251, Géométrie et analyse microlocales.
- [BZGM15] Christine Berkesch Zamaere, Stephen Griffeth, and Ezra Miller, *Systems of parameters and holonomicity of A -hypergeometric systems*, Pacific J. Math. **276** (2015), no. 2, 281–286.
- [CnDRS19] Alberto Castaño Domínguez, Thomas Reichelt, and Christian Sevenheck, *Examples of hypergeometric twistor \mathcal{D} -modules*, Algebra Number Theory **13** (2019), no. 6, 1415–1442.
- [CnDS19] Alberto Castaño Domínguez, Alberto and Christian Sevenheck, *Irregular Hodge filtration of some confluent hypergeometric systems*, J. Inst. Math. Jussieu (2019), 142.
- [DE03] Andrea D’Agnolo and Michael Eastwood, *Radon and Fourier transforms for \mathcal{D} -modules*, Adv. Math. **180** (2003), no. 2, 452–485.
- [DS13] Michael Dettweiler and Claude Sabbah, *Hodge theory of the middle convolution*, Publ. Res. Inst. Math. Sci. **49** (2013), no. 4, 761–800.
- [ESY17] Hélène Esnault, Claude Sabbah, and Jeng-Daw Yu, *E_1 -degeneration of the irregular Hodge filtration*, J. Reine Angew. Math. **729** (2017), 171–227.
- [Ful93] William Fulton, *Introduction to toric varieties*, Annals of Mathematics Studies, vol. 131, Princeton University Press, Princeton, NJ, 1993, The William H. Roever Lectures in Geometry.
- [GfGZ87] I. M. Gel’fand, M. I. Graev, and A. V. Zelevinskiĭ, *Holonomic systems of equations and series of hypergeometric type*, Dokl. Akad. Nauk SSSR **295** (1987), no. 1, 14–19.
- [GfZK89] I. M. Gel’fand, A. V. Zelevinskiĭ, and M. M. Kapranov, *Hypergeometric functions and toric varieties*, Funktsional. Anal. i Prilozhen. **23** (1989), no. 2, 12–26.
- [GKZ90] Israel M. Gel’fand, Mikhail M. Kapranov, and Andrei V. Zelevinsky, *Generalized Euler integrals and A -hypergeometric functions*, Adv. Math. **84** (1990), no. 2, 255–271.
- [Gro61] A. Grothendieck, *Éléments de géométrie algébrique. III. Étude cohomologique des faisceaux cohérents. I*, Inst. Hautes Études Sci. Publ. Math. (1961), no. 11, 167.
- [Har77] Robin Hartshorne, *Algebraic geometry*, Springer-Verlag, New York-Heidelberg, 1977, Graduate Texts in Mathematics, No. 52.
- [Hoc72] Melvin Hochster, *Rings of invariants of tori, Cohen-Macaulay rings generated by monomials, and polytopes*, Ann. of Math. (2) **96** (1972), 318–337.
- [HTT08] Ryoshi Hotta, Kiyoshi Takeuchi, and Toshiyuki Tanisaki, *\mathcal{D} -modules, perverse sheaves, and representation theory*, Progress in Mathematics, vol. 236, Birkhäuser Boston Inc., Boston, MA, 2008, Translated from the 1995 Japanese edition by Takeuchi.
- [Kat90] Nicholas M. Katz, *Exponential sums and differential equations*, Annals of Mathematics Studies, vol. 124, Princeton University Press, Princeton, NJ, 1990.

- [KKP17] Ludmil Katzarkov, Maxim Kontsevich, and Tony Pantev, *Bogomolov-Tian-Todorov theorems for Landau-Ginzburg models*, J. Differential Geom. **105** (2017), no. 1, 55–117.
- [MM04] Philippe Maisonobe and Zoghman Mebkhout, *Le théorème de comparaison pour les cycles évanescents*, Éléments de la théorie des systèmes différentiels géométriques (Philippe Maisonobe and Luis Narváez Macarro, eds.), Sémin. Congr., vol. 8, Soc. Math. France, Paris, 2004, Papers from the CIMPA Summer School held in Séville, September 2–13, 1996, pp. 311–389.
- [MM17] Etienne Mann and Thierry Mignon, *Quantum \mathcal{D} -modules for toric nef complete intersections*, Internat. J. Math. **28** (2017), no. 6, 1750047, 49.
- [MMW05] Laura Felicia Matusevich, Ezra Miller, and Uli Walther, *Homological methods for hypergeometric families*, J. Amer. Math. Soc. **18** (2005), no. 4, 919–941.
- [Moc15a] Takuro Mochizuki, *Mixed twistor \mathcal{D} -modules*, Lecture Notes in Mathematics, vol. 2125, Springer, Cham, 2015.
- [Moc15b] ———, *Twistor property of GKZ-hypergeometric systems*, preprint arXiv:1501.04146, 2015.
- [MS05] Ezra Miller and Bernd Sturmfels, *Combinatorial commutative algebra*, Graduate Texts in Mathematics, vol. 227, Springer-Verlag, New York, 2005.
- [Rei14] Thomas Reichelt, *Laurent Polynomials, GKZ-hypergeometric Systems and Mixed Hodge Modules*, Compositio Mathematica (**150**) (2014), 911–941.
- [RS15] Thomas Reichelt and Christian Sevenheck, *Logarithmic Frobenius manifolds, hypergeometric systems and quantum \mathcal{D} -modules*, Journal of Algebraic Geometry **24** (2015), no. 2, 201–281.
- [RS17] ———, *Non-affine Landau-Ginzburg models and intersection cohomology*, Ann. Sci. Éc. Norm. Supér. (4) **50** (2017), no. 3, 665–753.
- [RSW18] Thomas Reichelt, Christian Sevenheck, and Uli Walther, *On the b -functions of hypergeometric systems*, Int. Math. Res. Not. IMRN (2018), no. 21, 6535–6555.
- [Sab05] Claude Sabbah, *Polarizable twistor \mathcal{D} -modules*, Astérisque (2005), no. 300, vi+208.
- [Sab08] ———, *Fourier-Laplace transform of a variation of polarized complex Hodge structure.*, J. Reine Angew. Math. **621** (2008), 123–158.
- [Sab11] ———, *Non-commutative Hodge structures*, Ann. Inst. Fourier (Grenoble) **61** (2011), no. 7, 2681–2717.
- [Sab18] C. Sabbah, *Irregular Hodge theory*, Mm. Soc. Mat. Fr. (N.S.) **156** (2018), available at arXiv:1511.00176 [math.AG] (with the collaboration of Jeng-Daw Yu).
- [Sai88] Morihiko Saito, *Modules de Hodge polarisables*, Publ. Res. Inst. Math. Sci. **24** (1988), no. 6, 849–995 (1989).
- [Sai90] ———, *Mixed Hodge modules*, Publ. Res. Inst. Math. Sci. **26** (1990), no. 2, 221–333.
- [Sai93] ———, *On b -function, spectrum and rational singularity*, Math. Ann. **295** (1993), no. 1, 51–74.
- [Sai94] ———, *On the theory of mixed Hodge modules*, Selected papers on number theory, algebraic geometry, and differential geometry, Amer. Math. Soc. Transl. Ser. 2, vol. 160, Amer. Math. Soc., Providence, RI, 1994, Translated from Sūgaku, Translation edited by Katsumi Nomizu, pp. 47–61.
- [SST00] Mutsumi Saito, Bernd Sturmfels, and Nobuki Takayama, *Gröbner deformations of hypergeometric differential equations*, Algorithms and Computation in Mathematics, vol. 6, Springer-Verlag, Berlin, 2000.

- [Sti98] Jan Stienstra, *Resonant hypergeometric systems and mirror symmetry*, Integrable systems and algebraic geometry (Kobe/Kyoto, 1997), World Sci. Publ., River Edge, NJ, 1998, pp. 412–452.
- [SW08] Mathias Schulze and Uli Walther, *Irregularity of hypergeometric systems via slopes along coordinate subspaces*, Duke Math. J. **142** (2008), no. 3, 465–509.
- [SW09] ———, *Hypergeometric \mathcal{D} -modules and twisted Gauß-Manin systems*, J. Algebra **322** (2009), no. 9, 3392–3409.
- [Wal07] Uli Walther, *Duality and monodromy reducibility of A -hypergeometric systems*, Math. Ann. **338** (2007), no. 1, 55–74.

Thomas Reichelt
Mathematisches Institut
Universität Heidelberg
69120 Heidelberg
Germany
treichelt@mathi.uni-heidelberg.de

Christian Sevenheck
Fakultät für Mathematik
Technische Universität Chemnitz
09107 Chemnitz
Germany
christian.sevenheck@mathematik.tu-chemnitz.de