FOURIER FRAMES FOR THE CANTOR-4 SET

GABRIEL PICIOROAGA AND ERIC S. WEBER

ABSTRACT. The measure supported on the Cantor-4 set constructed by Jorgensen-Pedersen is known to have a Fourier basis, i.e. that it possess a sequence of exponentials which form an orthonormal basis. We construct Fourier frames for this measure via a dilation theory type construction. We expand the Cantor-4 set to a 2 dimensional fractal which admits a representation of a Cuntz algebra. Using the action of this algebra, an orthonormal set is generated on the larger fractal, which is then projected onto the Cantor-4 set to produce a Fourier frame.

Jorgensen and Pedersen [10] demonstrated that there exist singular measures ν which are spectral—that is, they possess a sequence of exponential functions which form an orthonormal basis in $L^2(\nu)$. The canonical example of such a singular and spectral measure is the uniform measure on the Cantor 4-set defined as follows:

$$C_4 = \{x \in [0,1] : x = \sum_{k=1}^{\infty} \frac{a_k}{4^k}, \ a_k \in \{0,2\}\}.$$

This is analogous to the standard middle third Cantor set where 4^k replaces 3^k . The set C_4 can also be described as the attractor set of the following iterated function system on \mathbb{R} :

$$\tau_0(x) = \frac{x}{4}, \qquad \tau_2(x) = \frac{x+2}{4}.$$

The uniform measure on the set C_4 then is the unique probability measure μ_4 which is invariant under this iterated function system:

$$\int f(x)d\mu_4(x) = \frac{1}{2} \left(\int f(\tau_0(x)) d\mu_4(x) + \int f(\tau_2(x)) d\mu_4(x) \right)$$

for all $f \in C(\mathbb{R})$, see [9] for details. The standard spectrum for μ_4 is $\Gamma_4 = \{\sum_{n=0}^N l_n 4^n : l_n \in \{0,1\}\}$, though there are many spectra [4, 2].

Remarkably, Jorgensen and Pedersen prove that the uniform measure μ_3 on the standard middle third Cantor set is not spectral. Indeed, there are no three mutually orthogonal exponentials in $L^2(\mu_3)$. Thus, there has been much attention on whether there exists a Fourier frame for $L^2(\mu_3)$ —the problem is still unresolved, but see [5, 6] for progress in this regard. In this paper, we will construct Fourier frames for $L^2(\mu_4)$ using a dilation theory type argument. The motivation is whether the construction we demonstrate here for μ_4 will be applicable to μ_3 . Fourier frames for μ_4 were constructed in [6] using a duality type construction.

Date: July 13, 2018.

²⁰¹⁰ Mathematics Subject Classification. Primary: 42C15, 28A80; Secondary 42B05, 46L89.

Key words and phrases. Fourier series, frames, fractals, iterated function system, Cuntz algebra.

A frame for a Hilbert space H is a sequence $\{x_n\}_{n\in I}\subset H$ such that there exists constants A, B>0 such that for all $v\in H$,

$$A||v||^2 \le \sum_{n \in I} |\langle v, x_n \rangle|^2 \le B||v||^2.$$

The largest A and smallest B which satisfy these inequalities are called the frame bounds. The frame is called a Parseval frame if both frame bounds are 1. The sequence $\{x_n\}_{n\in I}$ is a Bessel sequence if there exists a constant B which satisfies the second inequality, whether or not the first inequality holds; B is called the Bessel bound. A Fourier frame for $L^2(\mu_4)$ is a sequence of frequencies $\{\lambda_n\}_{n\in I}\subset\mathbb{R}$ together with a sequence of "weights" $\{d_n\}_{n\in I}\subset\mathbb{C}$ such that $x_n=d_ne^{2\pi i\lambda_n x}$ is a frame. Fourier frames (unweighted) for Lebesgue measure were introduced by Duffin and Schaffer [3], see also Ortega-Cerda and Seip [13].

It was proven in [8] that a frame for a Hilbert space can be dilated to a Riesz basis for a bigger space, that is to say, that any frame is the image under a projection of a Riesz basis. Moreover, a Parseval frame is the image of an orthonormal basis under a projection. This result is now known to be a consequence of the Naimark dilation theory. This will be our recipe for constructing a Fourier frame: constructing a basis in a bigger space and then projecting onto a subspace. We require the following result along these lines [1]:

Lemma 1. Let H be a Hilbert space, V, K closed subspaces, and let P_V be the projection onto V. If $\{x_n\}_{n\in I}$ is a frame in K with frame bounds A, B, then:

- 1. $\{P_V x_n\}_{n \in I}$ is a Bessel sequence in V with Bessel bound no greater than B;
- 2. if the projection $P_V: K \to V$ is onto, then $\{P_V x_n\}_{n \in I}$ is a frame in V;
- 3. if $V \subset K$, then then $\{P_V x_n\}_{n \in I}$ is a frame in V with frame bounds between A and B.

Note that if $V \subset K$ and $\{x_n\}_{n\in I}$ is a Parseval frame for K, then $\{P_Vx_n\}_{n\in I}$ is a Parseval frame for V. In the second item above, it is possible that the lower frame bound for $\{P_Vx_n\}$ is smaller than A, but the upper frame bound is still no greater than B.

The foundation of our construction is a dilation theory type argument. Our first step, described in Section 1, is to consider the fractal like set $C_4 \times [0,1]$, which we will view in terms of an iterated function system. This IFS will give rise to a representation of the Cuntz algebra \mathcal{O}_4 on $L^2(\mu_4 \times \lambda)$ since $\mu_4 \times \lambda$ is the invariant measure under the IFS. Then in Section 2, we will generate via the action of \mathcal{O}_4 an orthonormal set in $L^2(\mu_4 \times \lambda)$ whose vectors have a particular structure. In Section 3, we consider a subspace V of $L^2(\mu_4 \times \lambda)$ which can be naturally identified with $L^2(\mu_4)$, and then project the orthonormal set onto V to, ultimately, obtain a frame. Of paramount importance will be whether the orthonormal set generated by \mathcal{O}_4 spans the subspace V so that the projection yields a Parseval frame. Section 4 demonstrates concrete constructions in which this occurs, and identifies all possible Fourier frames that can be constructed using this method.

We note here that there may be Fourier frames for $L^2(\mu_4)$ which cannot be constructed in this manner, but we are unaware of such an example.

1. Dilation of the Cantor-4 Set

We wish to construct a Hilbert space H which contains $L^2(\mu_4)$ as a subspace in a natural way. We will do this by making the fractal C_4 bigger as follows. We begin with an iterated

function system on \mathbb{R}^2 given by:

$$\Upsilon_0(x,y) = (\frac{x}{4}, \frac{y}{2}), \ \Upsilon_1(x,y) = (\frac{x+2}{4}, \frac{y}{2}), \ \Upsilon_2(x,y) = (\frac{x}{4}, \frac{y+1}{2}), \ \Upsilon_3(x,y) = (\frac{x+2}{4}, \frac{y+1}{2}).$$

As these are contractions on \mathbb{R}^2 , there exists a compact attractor set, which is readily verified to be $C_4 \times [0,1]$. Likewise, by Hutchinson [9], there exists an invariant probability measure supported on $C_4 \times [0,1]$; it is readily verified that this invariant measure is $\mu_4 \times \lambda$, where λ denotes the Lebesgue measure restricted to [0,1]. Thus, for every continuous function $f: \mathbb{R}^2 \to \mathbb{C}$,

(1)
$$\int f(x,y) \ d(\mu_4 \times \lambda) = \frac{1}{4} \left(\int f(\frac{x}{4}, \frac{y}{2}) \ d(\mu_4 \times \lambda) + \int f(\frac{x+2}{4}, \frac{y}{2}) \ d(\mu_4 \times \lambda) + \int f(\frac{x+2}{4}, \frac{y+1}{2}) \ d(\mu_4 \times \lambda) + \int f(\frac{x+2}{4}, \frac{y+1}{2}) \ d(\mu_4 \times \lambda) \right).$$

The iterated function system Υ_j has a left inverse on $C_4 \times [0,1]$, given by

$$R: C_4 \times [0,1] \to C_4 \times [0,1]: (x,y) \mapsto (4x,2y) \mod 1,$$

so that $R \circ \Upsilon_j(x, y) = (x, y)$ for j = 0, 1, 2, 3.

We will use the iterated function system to define an action of the Cuntz algebra \mathcal{O}_4 on $L^2(\mu_4 \times \lambda)$. To do so, we choose filters

$$m_0(x,y) = H_0(x,y)$$

$$m_1(x,y) = e^{2\pi i x} H_1(x,y)$$

$$m_2(x,y) = e^{4\pi i x} H_2(x,y)$$

$$m_3(x,y) = e^{6\pi i x} H_3(x,y)$$

where

$$H_j(x,y) = \sum_{k=0}^{3} a_{jk} \chi_{\Upsilon_k(C_4 \times [0,1])}(x,y)$$

for some choice of scalar coefficients a_{jk} . In order to obtain a representation of \mathcal{O}_4 on $L^2(\mu_4 \times \lambda)$, we require that the above filters satisfy the matrix equation $\mathcal{M}^*(x,y)\mathcal{M}(x,y) = I$ for $\mu_4 \times \lambda$ almost every (x,y), where

$$\mathcal{M}(x,y) = \begin{pmatrix} m_0(\Upsilon_0(x,y)) & m_0(\Upsilon_1(x,y)) & m_0(\Upsilon_2(x,y)) & m_0(\Upsilon_3(x,y)) \\ m_1(\Upsilon_0(x,y)) & m_1(\Upsilon_1(x,y)) & m_1(\Upsilon_2(x,y)) & m_1(\Upsilon_3(x,y)) \\ m_2(\Upsilon_0(x,y)) & m_2(\Upsilon_1(x,y)) & m_2(\Upsilon_2(x,y)) & m_2(\Upsilon_3(x,y)) \\ m_3(\Upsilon_0(x,y)) & m_3(\Upsilon_1(x,y)) & m_3(\Upsilon_2(x,y)) & m_3(\Upsilon_3(x,y)) \end{pmatrix}$$

For our choice of filters, the matrix \mathcal{M} becomes

$$\mathcal{M}(x,y) = \begin{pmatrix} a_{00} & a_{01} & a_{02} & a_{03} \\ e^{\pi i x/2} a_{10} & -e^{\pi i x/2} a_{11} & e^{\pi i x/2} a_{12} & -e^{\pi i x/2} a_{13} \\ e^{\pi i x} a_{20} & e^{\pi i x} a_{21} & e^{\pi i x} a_{22} & e^{\pi i x} a_{23} \\ e^{3\pi i x/2} a_{30} & -e^{3\pi i x/2} a_{31} & e^{3\pi i x/2} a_{32} & -e^{3\pi i x/2} a_{33} \end{pmatrix},$$

which is unitary if and only if the matrix

$$H = \begin{pmatrix} a_{00} & a_{01} & a_{02} & a_{03} \\ a_{10} & -a_{11} & a_{12} & -a_{13} \\ a_{20} & a_{21} & a_{22} & a_{23} \\ a_{30} & -a_{31} & a_{32} & -a_{33} \end{pmatrix}$$

is unitary. For the remainder of this section, we assume that H is unitary.

Lemma 2. The operator $S_i: L^2(\mu_4 \times \lambda) \to L^2(\mu_4 \times \lambda)$ given by

$$[S_j f](x, y) = \sqrt{4}m_j(x, y)f(R(x, y))$$

is an isometry.

Proof. We calculate:

$$||S_j f||^2 = \int |\sqrt{4}m_j(x, y)f(R(x, y))|^2 d(\mu_4 \times \lambda)$$

$$= \frac{1}{4} \sum_{k=0}^3 \int 4|m_j(\Upsilon_k(x, y))f(R(\Upsilon_k(x, y)))|^2 d(\mu_4 \times \lambda)$$

$$= \int \left(\sum_{k=0}^3 |m_j(\Upsilon_k(x, y))|^2\right) |f(x, y)|^2 d(\mu_4 \times \lambda).$$

We used Equation (1) in the second line. The sum in the integral is the square of the Euclidean norm of the j-th row of the matrix \mathcal{M} , which is unitary. Hence, the sum is 1, so the integral is $||f||^2$, as required.

Lemma 3. The adjoint is given by

$$[S_j^* f](x,y) = \frac{1}{2} \sum_{k=0}^3 \overline{m_j(\Upsilon_k(x,y))} f(\Upsilon_j(x,y)).$$

Proof. Let $f, g \in L^2(\mu_4 \times \lambda)$. We calculate

$$\langle S_j f, g \rangle = \int \sqrt{4} m_j(x, y) f(R(x, y)) \overline{g(x, y)} \ d(\mu_4 \times \lambda)$$

$$= \frac{1}{4} \sum_{k=0}^3 \int \sqrt{4} m_j(\Upsilon_k(x, y)) f(R(\Upsilon_k(x, y))) \overline{g(\Upsilon_k(x, y))} \ d(\mu_4 \times \lambda)$$

$$= \int f(x, y) \overline{\left(\frac{1}{2} \sum_{k=0}^3 \overline{m_j(\Upsilon_k(x, y))} g(\Upsilon_k(x, y))\right)} \ d(\mu_4 \times \lambda)$$

where we use Equation (1) and the fact that R is a left inverse of Υ_k .

Lemma 4. The isometries S_j satisfy the Cuntz relations:

$$S_j^* S_k = \delta_{jk} I, \qquad \sum_{k=0}^3 S_k S_k^* = I.$$

Proof. We consider the orthogonality relation first. Let $f \in L^2(\mu_4 \times \lambda)$. We calculate:

$$[S_j^* S_k f](x,y) = \frac{1}{2} \sum_{\ell=0}^3 \overline{m_j(\Upsilon_\ell(x,y))} [S_k f](\Upsilon_\ell(x,y))$$

$$= \frac{1}{2} \sum_{\ell=0}^3 \overline{m_j(\Upsilon_\ell(x,y))} \sqrt{4} m_k (\Upsilon_\ell(x,y)) f(R(\Upsilon_\ell(x,y)))$$

$$= \left(\sum_{\ell=0}^3 \overline{m_j(\Upsilon_\ell(x,y))} m_k (\Upsilon_\ell(x,y))\right) f(x,y).$$

Note that the sum is the scalar product of the k-th row with the j-th row of the matrix \mathcal{M} , which is unitary. Hence, the sum is δ_{jk} as required.

Now for the identity relation, let $f, g \in L^2(\mu_4 \times \lambda)$. We calculate:

$$\begin{split} &\langle \sum_{k=0}^{3} S_{k} S_{k}^{*} f, g \rangle = \sum_{k=0}^{3} \langle S_{k}^{*} f, S_{k}^{*} g \rangle \\ &= \sum_{k=0}^{3} \int \left(\frac{1}{2} \sum_{\ell=0}^{3} \overline{m_{k}(\Upsilon_{\ell}(x,y))} f(\Upsilon_{\ell}(x,y)) \right) \left(\frac{1}{2} \sum_{n=0}^{3} \overline{m_{k}(\Upsilon_{n}(x,y))} g(\Upsilon_{n}(x,y)) \right) d(\mu_{4} \times \lambda) \\ &= \sum_{\ell=0}^{3} \sum_{n=0}^{3} \frac{1}{4} \int \left(\sum_{k=0}^{3} \overline{m_{k}(\Upsilon_{\ell}(x,y))} \overline{m_{k}(\Upsilon_{n}(x,y))} \right) f(\Upsilon_{\ell}(x,y)) \overline{g(\Upsilon_{n}(x,y))} d(\mu_{4} \times \lambda) \\ &= \frac{1}{4} \sum_{n=0}^{3} \int f(\Upsilon_{n}(x,y)) \overline{g(\Upsilon_{n}(x,y))} d(\mu_{4} \times \lambda) \\ &= \int f(x,y) \overline{g(x,y)} d(\mu_{4} \times \lambda) \\ &= \langle f,g \rangle. \end{split}$$

Note that the sum over k in the third line is the scalar product of the ℓ -th column with the n-th column of \mathcal{M} , so the sum collapses to $\delta_{\ell n}$. The sum on n in the fourth line collapses by Equation (1).

2. Orthonormal Sets in $L^2(\mu_4 \times \lambda)$

Since the isometries S_j satisfy the Cuntz relations, we can use them to generate orthonormal sets in the space $L^2(\mu_4 \times \lambda)$. We do so by having the isometries act on a generating vector. We consider words in the alphabet $\{0,1,2,3\}$; let W_4 denote the set of all such words. For a word $\omega = j_K j_{K-1} \dots j_1$, we denote by $|\omega| = K$ the length of the word, and define

$$S_{\omega}f = S_{j_K}S_{j_{K-1}}\dots S_{j_1}f.$$

Definition 1. Let

$$X_4 = \{\omega \in W_4 : |\omega| = 1\} \cup \{\omega \in W_4 : |\omega| \ge 2, \ j_1 \ne 0\}.$$

For convenience, we allow the empty word ω_{\emptyset} with length 0, and define $S_{\omega_{\emptyset}} = I$, the identity.

Lemma 5. Suppose $f \in L^2(\mu_4 \times \lambda)$ with ||f|| = 1, and that $S_0 f = f$. Then, $\{S_{\omega} f : \omega \in X_4\}$

is an orthonormal set.

Proof. Suppose $\omega, \omega' \in X_4$ with $\omega \neq \omega'$. First consider $|\omega| = |\omega'|$, with $\omega = j_K \dots j_1$ and $\omega' = i_K \dots i_1$. Suppose that ℓ is the largest index such that $j_\ell \neq i_\ell$. Then we have

$$\langle S_{\omega}f, S_{\omega'}f \rangle = \langle S_{j_{\ell}} \dots S_{j_{1}}f, S_{i_{\ell}} \dots S_{i_{1}}f \rangle = \langle S_{i_{\ell}}^{*}S_{j_{\ell}} \dots S_{j_{1}}f, S_{i_{\ell-1}} \dots S_{i_{1}}f \rangle = 0$$

by the orthogonality condition of the Cuntz relations.

Now, if $K = |\omega| > |\omega'| = M$, with $\omega' = i_M \dots i_1$, we define the word $\rho = i_M \dots i_1 0 \dots 0$ so that $|\rho| = K$. Note that $\rho \notin X_4$ so $\omega \neq \rho$. Note further that $S_{\omega'} f = S_{\rho} f$. Thus, by a similar argument to that above, we have

$$\langle S_{\omega}f, S_{\omega'}f\rangle = 0.$$

Remark 1. The set $\{S_{\omega}f : \omega \in X_4\}$ need not be complete. We will provide an example of this in Example 1 in Section 4.

Our goal is to project the set $\{S_{\omega}f : \omega \in X_4\}$ onto some subspace V of $L^2(\mu_4 \times \lambda)$ to obtain a frame. To that end, we need to know when the projection $\{P_VS_{\omega}f : \omega \in X_4\}$ is a frame, which by Lemma 1 requires the projection $P_V : K \to V$ to be onto, where K is the subspace spanned by $\{S_{\omega}f : \omega \in X_4\}$. The tool we will use is the following result, which is a minor adaptation of a result from [7]; we will not use this result directly, but will use all of the critical components.

Theorem 1. Let \mathcal{H} be a Hilbert space, $\mathcal{K} \subset \mathcal{H}$ a closed subspace, and $(S_i)_{i=0}^{N-1}$ be a representation of the Cuntz algebra \mathcal{O}_N . Let \mathcal{E} be an orthonormal set in \mathcal{H} and $f: X \to \mathcal{K}$ a norm continuous function on a topological space X with the following properties:

- i) $\mathcal{E} = \bigcup_{i=0}^{N-1} S_i \mathcal{E}$ where the union is disjoint.
- ii) $\overline{span}\{f(t): t \in X\} = \mathcal{K} \text{ and } ||f(t)|| = 1, \text{ for all } t \in X.$
- iii) There exist functions $\mathfrak{m}_i: X \to \mathbb{C}$, $g_i: X \to X$, $i = 0, \ldots, N-1$ such that

(2)
$$S_i^* f(t) = \mathfrak{m}_i(t) f(g_i(t)), \quad t \in X.$$

- iv) There exist $c_0 \in X$ such that $f(c_0) \in \overline{span}\mathcal{E}$.
- v) The only function $h \in \mathcal{C}(X)$ with $h \ge 0$, h(c) = 1, $\forall c \in \{x \in X : f(x) \in \overline{span}\mathcal{E}\}$, and

(3)
$$h(t) = \sum_{i=0}^{N-1} |\mathfrak{m}_i(t)|^2 h(g_i(t)), \quad t \in X$$

are the constant functions.

Then $\mathcal{K} \subset \overline{span}\mathcal{E}$.

3. The Projection

Recall the definition of the filters $m_j(x,y) = e^{2\pi i j x} H_j(x,y)$ from Section 1. We choose the filter coefficients a_{jk} so that the matrix H is unitary. We place the additional constraint that

$$a_{00} = a_{01} = a_{02} = a_{03} = \frac{1}{2},$$

so that $S_0\mathbb{1} = \mathbb{1}$, where $\mathbb{1}$ the function in $L^2(\mu_4 \times \lambda)$ which is identically 1. As $S_0\mathbb{1} = \mathbb{1}$, by Lemma 5, the set $\{S_\omega\mathbb{1} : \omega \in X_4\}$ is orthonormal. Moreover, we place the additional constraint that for every j, $a_{j0} + a_{j2} = a_{j1} + a_{j3}$, which will be required for our calculation of the projection.

Definition 2. We define the subspace $V = \{f \in L^2(\mu_4 \times \lambda) : f(x,y) = g(x)\chi_{[0,1]}(y), g \in L^2(\mu_4)\}$. Note that the subspace V can be identified with $L^2(\mu_4)$ via the isometric isomorphism $g \mapsto g(x)\chi_{[0,1]}(y)$. We will suppress the y variable in the future.

Definition 3. We define a function $c: X_4 \to \mathbb{N}_0$ as follows: for a word $\omega = j_K j_{K-1} \dots j_1$,

$$c(\omega) = \sum_{k=1}^{K} j_k 4^{K-k}.$$

Here $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. It is readily verified that c is a bijection.

Lemma 6. For a word $\omega = j_K j_{K-1} \dots j_1$,

$$S_{\omega} \mathbb{1} = e^{2\pi i c(\omega)x} \left(\prod_{k=1}^{K} 2H_{j_k}(R^{K-k}(x,y)) \right).$$

Proof. We proceed by induction on the length of the word ω . The equality is readily verified for $|\omega| = 1$. Let $\omega_0 = j_{K-1} j_{n-2} \dots j_1$. We have

$$\begin{split} S_{\omega} \mathbb{1} &= S_{j_K} S_{\omega_0} \mathbb{1} \\ &= S_{j_K} \left[e^{2\pi i c(\omega_0) x} \left(\prod_{k=1}^{K-1} 2H_{j_k} (R^{K-1-k}(x,y)) \right) \right] \\ &= 2 e^{2\pi i \lambda_{j_K} x} H_{j_K}(x,y) e^{2\pi i c(\omega_0) \cdot 4x} \left(\prod_{k=1}^{K-1} H_{j_k} (R^{K-k}(x,y)) \right) \\ &= 2 e^{2\pi i (\lambda_{j_K} + 4c(\omega_0)) x} H_{j_K}(R^{K-K}(x,y)) \left(\prod_{k=1}^{K-1} 2H_{j_k} (R^{K-k}(x,y)) \right) \\ &= e^{2\pi i c(\omega) x} \left(\prod_{k=1}^{K} 2H_{j_k} (R^{K-k}(x,y)) \right). \end{split}$$

The last line above is justified by the following calculation:

$$\lambda_{j_K} + 4c(\omega_0) = \lambda_{j_K} + 4\left(\sum_{k=1}^{K-1} \lambda_{j_k} 4^{K-1-k}\right)$$

$$= \lambda_{j_K} 4^{K-K} + \sum_{k=1}^{K-1} \lambda_{j_k} 4^{K-k}$$

$$= \sum_{k=1}^{K} \lambda_{j_k} 4^{K-k}$$

$$= c(\omega).$$

We wish to project the vectors $S_{\omega} \mathbb{1}$ onto the subspace V. The following lemma calculates that projection, where P_V denotes the projection onto the subspace V.

Lemma 7. If
$$f(x,y) = g(x)h(x,y)$$
 with $g \in L^2(\mu_4)$ and $h \in L^\infty(\mu_4 \times \lambda)$, then $[P_V f](x,y) = g(x)G(x)$

where $G(x) = \int_{[0,1]} h(x,y) d\lambda(y)$.

Proof. We verify that for every $F(x) \in L^2(\mu_4)$, f(x,y) - g(x)G(x) is orthogonal to F(x). We calculate utilizing Fubini's theorem:

$$\langle f - gG, F \rangle = \int \int g(x)h(x,y)\overline{F(x)} \ d(\mu_4 \times \lambda) - \int \int g(x)G(x)\overline{F(x)} \ d(\mu_4 \times \lambda)$$

$$= \int_{C_4} g(x)\overline{F(x)} \left(\int_{[0,1]} h(x,y) - G(x) \ d\lambda(y) \right) \ d\mu_4(x)$$

$$= \int_{C_4} g(x)\overline{F(x)} \left(G(x) - G(x) \right) \ d\mu_4(x)$$

$$= 0.$$

For the purposes of the following lemma, αx and βy are understood to be modulo 1.

Lemma 8. For any word $\omega = j_K j_{K-1} \dots j_1$,

$$\int \prod_{k=1}^{K} 2H_{j_k}(R^{k-1}(x,y)) \ d\lambda(y) = \prod_{k=1}^{K} 2\int H_{j_k}(4^{k-1}x,y) \ d\lambda(y).$$

Proof. Let $F_m(x,y) = \prod_{k=m}^{K} 2H_{j_k}(4^{k-1}x, 2^{k-m}y)$. Note that

$$F_m(x, \frac{y}{2}) = 2H_{j_m}(4^{m-1}x, \frac{y}{2}) \left(\prod_{k=m+1}^K 2H_{j_k}(4^{k-1}x, 2^{k-(m+1)}y) \right) = 2H_{j_m}(4^{m-1}x, \frac{y}{2}) F_{m+1}(x, y).$$

Likewise for $F_m(x, \frac{y+1}{2})$.

Since λ is the invariant measure for the iterated function system $y \mapsto \frac{y}{2}$, $y \mapsto \frac{y+1}{2}$, we calculate:

$$\int_{0}^{1} F_{m}(x,y) \ d\lambda(y) = \frac{1}{2} \left[\int_{0}^{1} F_{m}(x,\frac{y}{2}) \ d\lambda(y) + \int_{0}^{1} F_{m}(x,\frac{y+1}{2}) \ d\lambda(y) \right]$$

$$= \frac{1}{2} \left[\int_{0}^{1} 2H_{j_{m}}(4^{m-1}x,\frac{y}{2})F_{m+1}(x,y) + 2H_{j_{m}}(4^{m-1}x,\frac{y+1}{2})F_{m+1}(x,y) \ d\lambda(y) \right]$$

$$= \frac{1}{2} \left[\int_{0}^{1} 2a_{j_{m},q}F_{m+1}(x,y) + 2a_{j_{m},q+2}F_{m+1}(x,y) \ d\lambda(y) \right]$$

$$= \frac{1}{2} \left[2a_{j_{m},q} + 2a_{j_{m},q+2} \right] \cdot \left[\int_{0}^{1} F_{m+1}(x,y) \ d\lambda(y) \right]$$

$$= \left[\int_{0}^{1} 2H_{j_{m}}(4^{m-1}x,y) \ d\lambda(y) \right] \cdot \left[\int_{0}^{1} F_{m+1}(x,y) \ d\lambda(y) \right]$$

where q = 0 if $0 \le 4^{m-1}x < \frac{1}{2}$, and q = 1 if $\frac{1}{2} \le 4^{m-1}x < 1$.

The result now follows by a standard induction argument.

Proposition 1. Suppose the filters $m_j(x,y)$ are chosen so that

- i) the matrix H is unitary,
- ii) $a_{00} = a_{01} = a_{02} = a_{03} = \frac{1}{2}$, and
- iii) for j = 0, 1, 2, 3, $a_{j0} + a_{j2} = a_{j1} + a_{j3}$.

Then for any word $\omega = j_K \dots j_1$,

$$P_V S_{\omega} \mathbb{1} = d_{\omega} e^{2\pi i c(\omega)x},$$

where

(4)
$$d_{\omega} = \prod_{k=1}^{K} \left(a_{j_k 0} + a_{j_k 2} \right).$$

Proof. We apply the previous three Lemmas to obtain

$$[P_V S_{\omega} \mathbb{1}](x,y) = e^{2\pi i c(\omega)x} \int \prod_{k=1}^K 2H_{j_k}(4^{k-1}x,y)d\lambda(y)$$
$$= e^{2\pi i c(\omega)x} \prod_{k=1}^K 2\int H_{j_k}(4^{k-1}x,y)d\lambda(y)$$

By assumption iii), the integral $\int H_{j_k}(4^{k-1}x,y)d\lambda(y)$ is independent of x, and the value of the integral is $\frac{a_{j0}}{2} + \frac{a_{j2}}{2}$. Equation 4 now follows.

4. Concrete Constructions

We now turn to concrete constructions of Fourier frames for μ_4 . The hypotheses of Lemma 5 and Proposition 1 require H to be unitary and requires the matrix

$$A = \begin{pmatrix} a_{00} & a_{01} & a_{02} & a_{03} \\ a_{10} & a_{11} & a_{12} & a_{13} \\ a_{20} & a_{21} & a_{22} & a_{23} \\ a_{30} & a_{31} & a_{32} & a_{33} \end{pmatrix}$$

to have the first row be identically $\frac{1}{2}$ and to have the vector $\begin{pmatrix} 1 & -1 & 1 & -1 \end{pmatrix}^T$ in the kernel. We can use Hadamard matrices to construct examples of such a matrix A. Every 4×4 Hadamard matrix is a permutation of the following matrix:

$$U_{\rho} = \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & \rho & -\rho \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -\rho & \rho \end{pmatrix}$$

where ρ is any complex number of modulus 1.

If we set $H = U_{\rho}$, we obtain

(5)
$$A = \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & \rho & \rho \\ 1 & 1 & -1 & -1 \\ 1 & 1 & -\rho & -\rho \end{pmatrix}$$

which has the requisite properties to apply Lemma 5 and Proposition 1.

We define for k = 1, 2, 3, $l_k : \mathbb{N}_0 \to \mathbb{N}_0$ by $l_k(n)$ is the number of digits equal to k in the base 4 expansion of n. Note that $l_k(0) = 0$, and we follow the convention that $0^0 = 1$.

Theorem 2. For the choice A as in Equation (5) with $\rho \neq -1$, the sequence

(6)
$$\left\{ \left(\frac{1+\rho}{2} \right)^{l_1(n)} 0^{l_2(n)} \left(\frac{1-\rho}{2} \right)^{l_3(n)} e^{2\pi i n x} : n \in \mathbb{N}_0 \right\}$$

is a Parseval frame in $L^2(\mu_4)$.

Proof. By Lemma 5, we have that $\{S_{\omega}\mathbb{1} : \omega \in X_4\}$ is an orthonormal set. For a word $\omega = j_K j_{K-1} \dots j_1$, Proposition 1 yields that

$$P_V S_{\omega} \mathbb{1} = e^{2\pi i c(\omega)x} \prod_{k=1}^{K} (a_{j_k 0} + a_{j_k 2}).$$

Then, setting $n = c(\omega)$, we obtain

$$P_V S_{\omega} \mathbb{1} = e^{2\pi i n x} \left(a_{00} + a_{02} \right)^{K - l_1(n) - l_2(n) - l_3(n)} \prod_{i=1}^{3} \left(a_{i0} + a_{i2} \right)^{l_i(n)}.$$

Since

$$a_{00} + a_{02} = 1$$
, $a_{10} + a_{12} = \frac{1+\rho}{2}$, $a_{20} + a_{22} = 0$, $a_{30} + a_{32} = \frac{1-\rho}{2}$,

it follows that

$$P_V S_{\omega} \mathbb{1} = \left(\frac{1+\rho}{2}\right)^{l_1(n)} 0^{l_2(n)} \left(\frac{1-\rho}{2}\right)^{l_3(n)} e^{2\pi i n x}.$$

Since c is a bijection, the set $\{P_V S_{\omega} \mathbb{1} : \omega \in X_4\}$ coincides with the set in (6).

In order to establish that the set (6) is a Parseval frame, we wish to apply Lemma 1, which requires that the subspace V is contained in the closed span of $\{S_{\omega}1: \omega \in X_4\}$. Denote the closed span by \mathcal{K} . We will proceed in a manner similar to Theorem 1. Define the function $f: \mathbb{R} \to V$ by $f(t) = e_t$ where $e_t(x, y) = e^{2\pi i x t}$. Note that $f(0) = 1 \in \mathcal{K}$. Likewise, define a function $h_X: \mathbb{R} \to \mathbb{R}$ by

$$h_X(t) = \sum_{\omega \in X_4} |\langle f(t), S_\omega \mathbb{1} \rangle|^2 = ||P_{\mathcal{K}} f(t)||^2.$$

Claim 1. We have $h_X \equiv 1$.

Assuming for the moment that the claim holds, we deduce that $f(t) \in \mathcal{K}$ for every $t \in \mathbb{R}$. Since $\{f(\gamma) : \gamma \in \Gamma_4\}$ is an orthonormal basis for V, it follows that the closed span of $\{f(t) : t \in \mathbb{R}\}$ is all of V. We conclude that $V \subset \mathcal{K}$, and so Lemma 1 implies that $\{P_V S_\omega \mathbb{1} : \omega \in X_4\}$ is a Parseval frame for V, from which the Theorem follows.

Thus, we turn to the proof of Claim 1. First, we require $\{S_{\omega}1 : \omega \in X_4\} = \bigcup_{j=0}^3 \{S_j S_{\omega}1 : \omega \in X_4\}$, where the union is disjoint. Clearly, the RHS is a subset of the LHS, and the union is disjoint. Consider an element of the LHS: $S_{\omega}1$. If $|\omega| \geq 2$, we write $S_{\omega}1 = S_j S_{\omega_0}1$ for some j and some $\omega_0 \in X_4$, whence $S_{\omega}1$ is in the RHS. If $|\omega| = 1$, then we write $S_{\omega}1 = S_j1 = S_j S_01$, which is again an element of the RHS. Equality now follows.

As a consequence,

$$\begin{split} h_X(t) &= \sum_{\omega \in X_4} |\langle f(t), S_\omega \mathbb{1} \rangle|^2 \\ &= \sum_{j=0}^3 \sum_{\omega \in X_4} |\langle f(t), S_j S_\omega \mathbb{1} \rangle|^2 \\ &= \sum_{j=0}^3 \sum_{\omega \in X_4} |\langle S_j^* f(t), S_\omega \mathbb{1} \rangle|^2. \end{split}$$

We calculate:

$$\begin{split} [S_j^*f(t)](x,y) &= \frac{1}{2} \sum_{k=0}^3 \overline{m_j(\Upsilon_k(x,y))} e_t(\Upsilon_k(x,y)) \\ &= \frac{1}{2} \left[\overline{a_{j0}} e^{-2\pi i j x/4} e_t(\frac{x}{4},\frac{y}{2}) + e^{-\pi i j} \overline{a_{j1}} e^{-2\pi i j x/4} e_t(\frac{x+2}{4},\frac{y}{2}) \right. \\ &\qquad \qquad + \overline{a_{j2}} e^{-2\pi i j x/4} e_t(\frac{x}{4},\frac{y+1}{2}) + e^{-\pi i j} \overline{a_{j3}} e^{-2\pi i j x/4} e_t(\frac{x+2}{4},\frac{y+1}{2}) \right] \\ &= \frac{1}{2} \left[\overline{a_{j0}} e^{-2\pi i j x/4} e_t(\frac{x}{4},\frac{y}{2}) + e^{-\pi i j} \overline{a_{j1}} e^{-2\pi i j x/4} e^{\pi i t} e_t(\frac{x}{4},\frac{y}{2}) \right. \\ &\qquad \qquad + \overline{a_{j2}} e^{-2\pi i j x/4} e_t(\frac{x}{4},\frac{y}{2}) + e^{-\pi i j} \overline{a_{j3}} e^{-2\pi i j x/4} e^{\pi i t} e_t(\frac{x}{4},\frac{y}{2}) \right] \\ &= \frac{1}{2} \left[\overline{a_{j0}} + e^{-\pi i j} \overline{a_{j1}} e^{\pi i t} + \overline{a_{j2}} + e^{-\pi i j} \overline{a_{j3}} e^{\pi i t} \right] e^{-2\pi i j x/4} e_t(\frac{x}{4},\frac{y}{2}) \\ &= \frac{1}{2} \left[\overline{a_{j0}} + e^{-\pi i j} \overline{a_{j1}} e^{\pi i t} + \overline{a_{j2}} + e^{-\pi i j} \overline{a_{j3}} e^{\pi i t} \right] e^{2\pi i (\frac{x}{4} - j\frac{x}{4})} \\ &= \frac{1}{2} \left[\overline{a_{j0}} + e^{-\pi i j} \overline{a_{j1}} e^{\pi i t} + \overline{a_{j2}} + e^{-\pi i j} \overline{a_{j3}} e^{\pi i t} \right] e^{2\pi i (\frac{t-j}{4}x)} \\ &= \frac{1}{2} \left[\overline{a_{j0}} + e^{-\pi i j} \overline{a_{j1}} e^{\pi i t} + \overline{a_{j2}} + e^{-\pi i j} \overline{a_{j3}} e^{\pi i t} \right] e^{2\pi i (\frac{t-j}{4}x)} \\ &= \frac{1}{2} \left[\overline{a_{j0}} + e^{-\pi i j} \overline{a_{j1}} e^{\pi i t} + \overline{a_{j2}} + e^{-\pi i j} \overline{a_{j3}} e^{\pi i t} \right] e^{2\pi i (\frac{t-j}{4}x)} \\ &= \frac{1}{2} \left[\overline{a_{j0}} + e^{-\pi i j} \overline{a_{j1}} e^{\pi i t} + \overline{a_{j2}} + e^{-\pi i j} \overline{a_{j3}} e^{\pi i t} \right] e^{2\pi i (\frac{t-j}{4}x)} \\ &= \frac{1}{2} \left[\overline{a_{j0}} + e^{-\pi i j} \overline{a_{j1}} e^{\pi i t} + \overline{a_{j2}} + e^{-\pi i j} \overline{a_{j3}} e^{\pi i t} \right] e^{2\pi i (\frac{t-j}{4}x)} \\ &= \frac{1}{2} \left[\overline{a_{j0}} + e^{-\pi i j} \overline{a_{j1}} e^{\pi i t} + \overline{a_{j2}} + e^{-\pi i j} \overline{a_{j3}} e^{\pi i t} \right] e^{2\pi i (\frac{t-j}{4}x)} \\ &= \frac{1}{2} \left[\overline{a_{j0}} + e^{-\pi i j} \overline{a_{j1}} e^{\pi i t} + \overline{a_{j2}} + e^{-\pi i j} \overline{a_{j3}} e^{\pi i t} \right] e^{2\pi i (\frac{t-j}{4}x)} \\ &= \frac{1}{2} \left[\overline{a_{j0}} + e^{-\pi i j} \overline{a_{j1}} e^{\pi i t} + \overline{a_{j2}} + e^{-\pi i j} \overline{a_{j3}} e^{\pi i t} \right] e^{2\pi i (\frac{t-j}{4}x)} \\ &= \frac{1}{2} \left[\overline{a_{j0}} + e^{-\pi i j} \overline{a_{j1}} e^{\pi i t} + \overline{a_{j2}} + e^{-\pi i j} \overline{a_{j3}} e^{\pi i t} \right] e^{2\pi i (\frac{t-j}{4}x)}$$

Thus, we define

$$\mathfrak{m}_{j}(t) = \frac{1}{2} \left(\overline{a_{j0}} + \overline{a_{j2}} \right) + \frac{e^{-\pi i j}}{2} \left(\overline{a_{j1}} + \overline{a_{j3}} \right) e^{\pi i t},$$

and

$$g_j(t) = \frac{t-j}{4}.$$

As a consequence, we obtain

$$h_X(t) = \sum_{j=0}^{3} \sum_{\omega \in X_4} |\langle S_j^* f(t), S_\omega \mathbb{1} \rangle|^2$$

$$= \sum_{j=0}^{3} \sum_{\omega \in X_4} |\langle \mathfrak{m}_j(t) f(g_j(t)), S_\omega \mathbb{1} \rangle|^2$$

$$= \sum_{j=0}^{3} |\mathfrak{m}_j(t)|^2 h_X(g_j(t)).$$
(7)

Because of our choice of coefficients in the matrix A, which has the vector $\begin{pmatrix} 1 & -1 & 1 & -1 \end{pmatrix}^T$ in the kernel, we have for every j: $a_{j0} + a_{j2} = a_{j1} + a_{j3}$. Thus, if we let $b_j = \overline{a_{j0}} + \overline{a_{j2}}$, the functions \mathfrak{m}_j simplify to

$$\mathfrak{m}_j(t) = b_j e^{\pi i \frac{t}{2}} \cos(\pi \frac{t}{2})$$

for j = 0, 2, and

$$\mathfrak{m}_j(t) = -ib_j e^{\pi i \frac{t}{2}} \sin(\pi \frac{t}{2})$$

for j = 1, 3. Substituting these into Equation (7), (8)

$$h_X(t) = \cos^2\left(\frac{\pi t}{2}\right) h_X\left(\frac{t}{4}\right) + \sin^2\left(\frac{\pi t}{2}\right) \frac{|1+\overline{\rho}|^2}{2} h_X\left(\frac{t-1}{4}\right) + \sin^2\left(\frac{\pi t}{2}\right) \frac{|1-\overline{\rho}|^2}{2} h_X\left(\frac{t-3}{4}\right).$$

Claim 2. The function h_X can be extended to an entire function.

Assume for the moment that Claim 2 holds, we finish the proof of Claim 1. If $h_X(t) = 1$ for $t \in [-1, 0]$, then $h_X(z) = 1$ for all $z \in \mathbb{C}$, and Claim 1 holds.

Now, assume to the contrary that $h_X(t)$ is not identically 1 on [-1,0]. Since $0 \le h_X(t) \le 1$ for t real, then $\beta = \min\{h_X(t) : t \in [-1,0]\} < 1$. Because constant functions satisfy (8), $h_1 := h_X - \beta$ also satisfies Equation (8). There exists t_0 such that $h_1(t_0) = 0$ and $t_0 \ne 0$ as $h_X(0) = 1$. Since $h_1 \ge 0$ each of the terms in (8) must vanish:

(9)
$$\cos^2\left(\frac{\pi t_0}{2}\right) h_1\left(\frac{t_0}{4}\right) = 0$$

(10)
$$\sin^2\left(\frac{\pi t_0}{2}\right) \frac{|1+\overline{\rho}|^2}{2} h_1\left(\frac{t_0-1}{4}\right) = 0$$

(11)
$$\sin^2\left(\frac{\pi t_0}{2}\right) \frac{|1-\overline{\rho}|^2}{2} h_1\left(\frac{t_0-3}{4}\right) = 0$$

Our hypothesis is that $\rho \neq -1$, so in Equation (10), the coefficient $\frac{|1+\overline{\rho}|}{2} \neq 0$.

Case 1: If $t_0 \neq -1$ then Equation (9) implies $h_1(t_0/4) = 0 = h_1(g_0(t_0))$. Let $t_1 := g_0(t_0) \in (-1,0)$; iterating the previous argument implies that $h_1(g_0(t_1)) = 0$. Thus, we obtain an infinite sequence of zeroes of h_1 .

Case 2: If $t_0 = -1$, then the previous argument does not hold. However, we can construct another zero of $h_1, t'_0 \in (-1,0)$ to which the previous argument will hold. Indeed, if $t_0 = -1$,

Equation (10) implies $h_1((t_0-1)/4) = h_1(-1/2) = 0$. Let $t'_0 = -1/2$ and continue as in Case 1.

In either case, h_1 vanishes on a (countable) set with an accumulation point, and since h_1 is analytic it follows that $h_1 \equiv 0$, a contradiction, and Claim 1 holds.

Now, to prove Claim 2, we follow the proof of Lemma 4.2 of [10]. For a fixed $\omega \in X_4$, define $f_{\omega} : \mathbb{C} \to \mathbb{C}$ by

$$f_{\omega}(z) = \langle e_z, S_{\omega} \mathbb{1} \rangle = \int e^{2\pi i z x} \overline{[S_{\omega} \mathbb{1}](x, y)} \ d(\mu_4 \times \lambda).$$

Since the distribution $\overline{[S_{\omega}1](x,y)}$ $d(\mu_4 \times \lambda)$ is compactly supported, a standard convergence argument demonstrates that f_{ω} is entire. Likewise, $f_{\omega}^*(z) = \overline{f_{\omega}(\overline{z})}$ is entire, and for t real,

$$f_{\omega}(t)f_{\omega}^{*}(t) = (\langle e_{t}, S_{\omega} \mathbb{1} \rangle) \left(\overline{\langle e_{t}, S_{\omega} \mathbb{1} \rangle} \right) = |\langle e_{t}, S_{\omega} \mathbb{1} \rangle|^{2}.$$

Thus,

$$h_X(t) = \sum_{\omega \in X_4} f_{\omega}(t) f_{\omega}^*(t).$$

For $n \in \mathbb{N}$, let $h_n(z) = \sum_{|\omega| \le n} f_{\omega}(z) f_{\omega}^*(z)$, which is entire. By Hölder's inequality,

$$\sum_{\omega \in X_4} |f_{\omega}(z) f_{\omega}^*(z)| \le \left(\sum_{\omega \in X_4} |\langle e_z, S_{\omega} \mathbb{1} \rangle|^2 \right)^{1/2} \left(\sum_{\omega \in X_4} |\langle e_{\overline{z}}, S_{\omega} \mathbb{1} \rangle|^2 \right)^{1/2}$$

$$\le ||e_z|| ||e_{\overline{z}}||$$

$$\le e^{KIm(z)}$$

for some constant K. Thus, the sequence $h_n(z)$ converges pointwise to a function h(z), and are uniformly bounded on strips $Im(z) \leq C$. By the theorems of Montel and Vitali, the limit function h is entire, which coincides with h_X for real t, and Claim 2 is proved.

Example 1. As mentioned in Section 2, in general, $\{S_{\omega}\mathbb{1}\}$ need not be complete, and the exceptional point $\rho = -1$ in Theorem 2 provides the example. In the case $\rho = -1$, the set (6) becomes

$$\{d_n e^{2\pi i n x} : n \in \mathbb{N}_0\}$$

where the coefficients $d_n = 1$ if $n \in \Gamma_3$ and 0 otherwise. Here,

$$\Gamma_3 = \{ \sum_{n=0}^{N} l_n 4^n : l_n \in \{0, 3\} \}$$

and it is known [4] that the sequence $\{e^{2\pi inx}: n \in \Gamma_3\}$ is incomplete in $L^2(\mu_4)$. Thus, $\{P_V S_\omega \mathbb{1}\}$ is incomplete in V, so $\{S_\omega \mathbb{1}\}$ is incomplete in $L^2(\mu_4 \times \lambda)$.

We can generalize the construction of Theorem 2 as follows. We want to choose a matrix

$$A = \begin{pmatrix} \frac{1}{2} & \frac{1}{2} & \frac{1}{2} & \frac{1}{2} \\ h_{10} & h_{11} & h_{12} & h_{13} \\ h_{20} & h_{21} & h_{22} & h_{23} \\ h_{30} & h_{31} & h_{32} & h_{33} \end{pmatrix}$$

such that $\begin{pmatrix} 1 & -1 & 1 & -1 \end{pmatrix}^T$ is in the kernel of H and the matrix

$$H = \begin{pmatrix} \frac{1}{2} & \frac{1}{2} & \frac{1}{2} & \frac{1}{2} \\ h_{10} & -h_{11} & h_{12} & -h_{13} \\ h_{20} & h_{21} & h_{22} & h_{23} \\ h_{30} & -h_{31} & h_{32} & -h_{33} \end{pmatrix}$$

is unitary. We obtain a system of nonlinear equations in the 12 unknowns. To parametrize all solutions, we consider the following row vectors:

(12)
$$\vec{v}_0 = \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & 1 \end{pmatrix}$$
 $\vec{w}_0 = \frac{1}{2} \begin{pmatrix} 1 & -1 & 1 & -1 \end{pmatrix}$

(13)
$$\vec{v}_1 = \frac{1}{2} \begin{pmatrix} 1 & -1 & -1 & 1 \end{pmatrix}$$
 $\vec{w}_1 = \frac{1}{2} \begin{pmatrix} 1 & 1 & -1 & -1 \end{pmatrix}$ (14) $\vec{v}_2 = \frac{1}{2} \begin{pmatrix} 1 & 1 & -1 & -1 \end{pmatrix}$ $\vec{w}_2 = \frac{1}{2} \begin{pmatrix} 1 & -1 & -1 & 1 \end{pmatrix}$

(14)
$$\vec{v}_2 = \frac{1}{2} \begin{pmatrix} 1 & 1 & -1 & -1 \end{pmatrix}$$
 $\vec{w}_2 = \frac{1}{2} \begin{pmatrix} 1 & -1 & -1 & 1 \end{pmatrix}$

If we construct the matrix A so that the rows are linear combinations of $\{\vec{v}_0, \vec{v}_1, \vec{v}_2\}$, then A will satisfy the desired condition on the kernel. Note that if the j-th row of A is $\alpha_{i0}\vec{v}_0$ + $\alpha_{j1}\vec{v}_1 + \alpha_{j2}\vec{v}_2$ for j = 1, 3, then the j-th row of H is $\alpha_{j0}\vec{w}_0 + \alpha_{j1}\vec{w}_1 + \alpha_{j2}\vec{w}_2$, whereas if j = 0, 2, then the j-th row of H is equal to the j-th row of A.

Thus, we want to choose coefficients α_{jk} , j=0,1,2,3, k=1,2,3 so that the matrix

(15)
$$H = \begin{pmatrix} \alpha_{00}\vec{v}_0 + \alpha_{01}\vec{v}_1 + \alpha_{02}\vec{v}_2 \\ \alpha_{10}\vec{w}_0 + \alpha_{11}\vec{w}_1 + \alpha_{12}\vec{w}_2 \\ \alpha_{20}\vec{v}_0 + \alpha_{21}\vec{v}_1 + \alpha_{22}\vec{v}_2 \\ \alpha_{30}\vec{w}_0 + \alpha_{31}\vec{w}_1 + \alpha_{32}\vec{w}_2 \end{pmatrix}$$

is unitary. To satisfy the requirement on the first row, we choose $\alpha_{00} = 1$ and $\alpha_{01} = \alpha_{02} = 0$. Calculating the inner products of the rows of H, we obtain the following necessary and sufficient conditions:

(16)
$$|\alpha_{j0}|^2 + |\alpha_{j1}|^2 + |\alpha_{j2}|^2 = 1$$

$$\alpha_{00}\overline{\alpha_{20}} = 0$$

$$\alpha_{11}\overline{\alpha_{22}} + \alpha_{12}\overline{\alpha_{21}} = 0$$

(19)
$$\alpha_{10}\overline{\alpha_{30}} + \alpha_{11}\overline{\alpha_{31}} + \alpha_{12}\overline{\alpha_{32}} = 0$$

$$\alpha_{21}\overline{\alpha_{32}} + \alpha_{22}\overline{\alpha_{31}} = 0$$

Proposition 2. Fix $\alpha_{00} = 1$. There exists a solution to the Equations (16) - (20) if and only if $\alpha_{10}, \alpha_{30} \in \mathbb{C}$ with

(21)
$$|\alpha_{10}|^2 + |\alpha_{30}|^2 = 1.$$

Proof. (\Leftarrow) If $|\alpha_{10}|^2 = 1$, then we choose $\alpha_{21} = \alpha_{31} = 1$ and all other coefficients to be 0 to obtain a solution to Equations (16) - (20). Likewise, if $|\alpha_{10}|^2 = 0$, then choose $\alpha_{11} = \alpha_{21} = 1$ and all other coefficients to be 0.

Now suppose that $0 < |\alpha_{10}| < 1$, and we choose $\lambda = \frac{-\overline{\alpha_{10}}\alpha_{30}}{1 - |\alpha_{10}|^2}$. Then choose α_{11} and α_{12} such that $|\alpha_{11}|^2 + |\alpha_{12}|^2 = 1 - |\alpha_{10}|^2$. Now let $\alpha_{31} = \lambda \alpha_{11}$ and $\alpha_{32} = \lambda \alpha_{12}$. We have

(22)
$$\alpha_{10}\overline{\alpha_{30}} + \alpha_{11}\overline{\alpha_{31}} + \alpha_{12}\overline{\alpha_{32}} = \alpha_{10}\overline{\alpha_{30}} + \overline{\lambda}|\alpha_{11}|^2 + \overline{\lambda}|\alpha_{12}|^2$$
$$= \alpha_{10}\overline{\alpha_{30}} + \overline{\lambda}(1 - |\alpha_{10}|^2)$$
$$= 0,$$

so Equation (19) is satisfied.

Equation (17) forces $\alpha_{20} = 0$; choose α_{21} and α_{22} such that $|\alpha_{21}|^2 + |\alpha_{22}|^2 = 1$ and $\alpha_{11}\overline{\alpha_{21}} + \alpha_{12}\overline{\alpha_{22}} = 0$. Thus, Equations (18) and (20) are satisfied. Finally, regarding Equation (16), it is satisfied for j = 0, 1, 2 by construction. For j = 3, we calculate:

$$|\alpha_{30}|^{2} + |\alpha_{31}|^{2} + |\alpha_{32}|^{2} = |\alpha_{30}|^{2} + |\lambda|^{2} (|\alpha_{11}|^{2} + |\alpha_{12}|^{2})$$

$$= |\alpha_{30}|^{2} + \frac{|\alpha_{10}|^{2} |\alpha_{30}|^{2}}{(1 - |\alpha_{10}|^{2})^{2}} (1 - |\alpha_{10}|^{2})$$

$$= |\alpha_{30}|^{2} \left(1 + \frac{|\alpha_{10}|^{2}}{1 - |\alpha_{10}|^{2}}\right)$$

$$= \frac{|\alpha_{30}|^{2}}{1 - |\alpha_{10}|^{2}}$$

$$= 1$$

as required.

(\Rightarrow) Suppose that we have a solution to Equations (16) - (20). If $|\alpha_{10}| = 1$, then we must have $\alpha_{11} = \alpha_{12} = 0$, and thus Equation (19) requires $\alpha_{30} = 0$, so Equation (21) holds.

Now suppose $|\alpha_{10}| < 1$. Since $\alpha_{20} = 0$, we must have that $|\alpha_{21}|^2 + |\alpha_{22}|^2 = 1$. Combining this with Equations (18) and (20) imply that the matrix

$$\begin{pmatrix} \alpha_{11} & \alpha_{12} \\ \alpha_{31} & \alpha_{32} \end{pmatrix}$$

is singular. Thus, there exists a λ such that $\alpha_{31} = \lambda \alpha_{11}$ and $\alpha_{32} = \lambda \alpha_{12}$. Using the same computation as in Equation (22), we conclude that $\lambda = \frac{-\overline{\alpha_{10}}\alpha_{30}}{1 - |\alpha_{10}|^2}$; then Equation (23) implies (21).

The coefficient matrix we obtain from this construction is

$$H = \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & 1 \\ \alpha_{10} + \alpha_{11} + \alpha_{12} & \alpha_{10} - \alpha_{11} + \alpha_{12} & \alpha_{10} - \alpha_{11} - \alpha_{12} & \alpha_{10} + \alpha_{11} - \alpha_{12} \\ \alpha_{21} + \alpha_{22} & -\alpha_{21} + \alpha_{22} & -\alpha_{21} - \alpha_{22} & \alpha_{21} - \alpha_{22} \\ \alpha_{30} + \lambda \alpha_{11} + \lambda \alpha_{12} & \alpha_{30} - \lambda \alpha_{11} + \lambda \alpha_{12} & \alpha_{30} - \lambda \alpha_{11} - \lambda \alpha_{12} & \alpha_{30} + \lambda \alpha_{11} - \lambda \alpha_{12} \end{pmatrix}$$

where we are allowed to choose α_{11} , α_{12} , α_{21} and α_{22} subject to the normalization condition in Equation (16). However, those choices do not affect the construction, since if we apply Proposition 1 and the calculation from Theorem 2, we obtain

(24)
$$P_V S_{\omega} \mathbb{1} = (\alpha_{10})^{\ell_1(n)} \cdot (0)^{\ell_2(n)} \cdot (\alpha_{30})^{\ell_3(n)} e^{2\pi i n x}.$$

This will in fact be a Parseval frame for $L^2(\mu_4)$, provided $V \subset \mathcal{K}$, as in the proof of Theorem 2.

Theorem 3. Suppose $p, q \in \mathbb{C}$ with $|p|^2 + |q|^2 = 1$. Then

$$\{p^{\ell_1(n)} \cdot 0^{\ell_2(n)} \cdot q^{\ell_3(n)} e^{2\pi i n x} : n \in \mathbb{N}_0\}$$

is a Parseval frame for $L^2(\mu_4)$, provided $p \neq 0$.

Proof. Substitute $\alpha_{10} = p$ and $\alpha_{30} = q$ in Proposition 2 and Equation (24). As noted, we only need to verify $V \subset \mathcal{K}$. We proceed as in the proof of Theorem 2; indeed, define f, h_X , \mathfrak{m}_j and g_j as previously. We obtain $b_0 = 1$, $b_1 = \overline{p}$, $b_2 = 0$, and $b_3 = \overline{q}$, so Equation (8) becomes

$$h_X(t) = \cos^2\left(\frac{\pi t}{2}\right) h_X\left(\frac{t}{4}\right) + |\overline{p}|^2 \sin^2\left(\frac{\pi t}{2}\right) h_X\left(\frac{t-1}{4}\right) + |\overline{q}|^2 \sin^2\left(\frac{\pi t}{2}\right) h_X\left(\frac{t-3}{4}\right).$$

From here, the same argument shows that $h_X \equiv 1$, and $V \subset \mathcal{K}$.

5. Concluding Remarks

We remark here that the constructions given above for μ_4 does not work for μ_3 . Indeed, we have the following no-go result. To obtain the measure $\mu_3 \times \lambda$, we consider the iterated function system:

$$\Upsilon_0(x,y) = (\frac{x}{3}, \frac{y}{2}), \ \Upsilon_1(x,y) = (\frac{x+2}{3}, \frac{y}{2}), \ \Upsilon_2(x,y) = (\frac{x}{3}, \frac{y+1}{2}), \ \Upsilon_3(x,y) = (\frac{x+2}{3}, \frac{y+1}{2}).$$

Using the same choice of filters, the matrix $\mathcal{M}(x,y)$ reduces to

$$H = \begin{pmatrix} a_{00} & a_{01} & a_{02} & a_{03} \\ a_{10} & e^{4\pi i/3} a_{11} & a_{12} & e^{4\pi i/3} a_{13} \\ a_{20} & e^{2\pi i/3} a_{21} & a_{22} & e^{2\pi i/3} a_{23} \\ a_{30} & a_{31} & a_{32} & a_{33} \end{pmatrix}$$

which we require to be unitary. Additionally, we require the same conditions as for μ_4 , namely, the first row of H must have all entries $\frac{1}{2}$, and $a_{j0} + a_{j2} = a_{j1} + a_{j3}$. The inner product of the first two rows must be 0. Hence,

$$\frac{1}{2} \left(a_{10} + e^{4\pi i/3} a_{11} + a_{12} + e^{4\pi i/3} a_{13} \right) = \frac{1}{2} \left(a_{10} + a_{12} \right) \left(1 + e^{4\pi i/3} \right) = 0.$$

Consequently, $a_{10} + a_{12} = 0$. Likewise, $a_{20} + a_{22} = a_{30} + a_{32} = 0$. As a result,

$$H\begin{pmatrix} 1\\0\\1\\0 \end{pmatrix} = \begin{pmatrix} a_{00} + a_{02}\\a_{10} + a_{12}\\a_{20} + a_{22}\\a_{30} + a_{32} \end{pmatrix} = \begin{pmatrix} 1\\0\\0\\0 \end{pmatrix}$$

and so H cannot be unitary.

It may be possible to extend the construction for μ_4 to μ_3 by considering a representation of \mathcal{O}_n for some sufficiently large n, or by considering $\mu_3 \times \rho$ for some other fractal measure ρ rather than λ .

Acknowledgement. We thank Dorin Dutkay for assisting with the proof of Claim 1 in Theorem 2.

REFERENCES

- Akram Aldroubi, Portraits of frames, Proc. Amer. Math. Soc. 123 (1995), no. 6, 1661–1668. MR 1242070 (95g:46037)
- Xin-Rong Dai, Xing-Gang He, and Chun-Kit Lai, Spectral property of Cantor measures with consecutive digits, Adv. Math. 242 (2013), 187–208. MR 3055992
- 3. R. J. Duffin and A. C. Schaeffer, A class of nonharmonic Fourier series, Trans. Amer. Math. Soc. 72 (1952), 341–366. MR 0047179 (13,839a)
- 4. Dorin Ervin Dutkay, Deguang Han, and Qiyu Sun, On the spectra of a Cantor measure, Adv. Math. **221** (2009), no. 1, 251–276. MR 2509326 (2010f:28013)
- 5. Dorin Ervin Dutkay, Deguang Han, and Eric Weber, Bessel sequences of exponentials on fractal measures, J. Funct. Anal. **261** (2011), no. 9, 2529–2539. MR 2826404 (2012h:42059)
- 6. ______, Continuous and discrete Fourier frames for fractal measures, Trans. Amer. Math. Soc. **366** (2014), no. 3, 1213–1235. MR 3145729
- 7. Dorin Ervin Dutkay, Gabriel Picioroaga, and Myung-Sin Song, Orthonormal bases generated by Cuntz algebras, J. Math. Anal. Appl. 409 (2014), no. 2, 1128–1139. MR 3103223
- 8. Deguang Han and David R. Larson, Frames, bases and group representations, Mem. Amer. Math. Soc. 147 (2000), no. 697, x+94. MR 1686653 (2001a:47013)
- 9. John E. Hutchinson, Fractals and self-similarity, Indiana Univ. Math. J. **30** (1981), no. 5, 713–747. MR MR625600 (82h:49026)
- 10. Palle E. T. Jorgensen and Steen Pedersen, Dense analytic subspaces in fractal L^2 -spaces, J. Anal. Math. **75** (1998), 185–228. MR MR1655831 (2000a:46045)
- 11. Izabella Laba and Yang Wang, On spectral Cantor measures, J. Funct. Anal. 193 (2002), no. 2, 409–420. MR 1929508 (2003g:28017)
- 12. Jian-Lin Li, $\mu_{M,D}$ -orthogonality and compatible pair, J. Funct. Anal. **244** (2007), no. 2, 628–638. MR 2297038 (2008h:42046)
- 13. Joaquim Ortega-Cerdà and Kristian Seip, *Fourier frames*, Ann. of Math. (2) **155** (2002), no. 3, 789–806. MR MR1923965 (2003k:42055)
- 14. Robert S. Strichartz, Mock Fourier series and transforms associated with certain Cantor measures, J. Anal. Math. 81 (2000), 209–238. MR 1785282 (2001i:42009)

Department of Mathematical Sciences, 414 E. Clark St., University of South Dakota, Vermillion, SD 57069

E-mail address: gabriel.picioroaga@usd.edu

DEPARTMENT OF MATHEMATICS, IOWA STATE UNIVERSITY, 396 CARVER HALL, AMES, IA 50011 E-mail address: esweber@iastate.edu