

# ON SOME METABELIAN 2-GROUP WHOSE ABELIANIZATION IS OF TYPE $(2, 2, 2)$ AND APPLICATIONS

A. AZIZI, A. ZEKHNINI, AND M. TAOUS

ABSTRACT. Let  $G$  be some metabelian 2-group satisfying the condition  $G/G' \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ . In this paper, we construct all the subgroups of  $G$  of index 2 or 4, we give the abelianization types of these subgroups and we compute the kernel of the transfer map. Then we apply these results to study the capitulation problem of the 2-ideal classes of some fields  $\mathbf{k}$  satisfying the condition  $\text{Gal}(\mathbf{k}_2^{(2)}/\mathbf{k}) \simeq G$ , where  $\mathbf{k}_2^{(2)}$  is the second Hilbert 2-class field of  $\mathbf{k}$ .

## 1. INTRODUCTION

Let  $G$  be a group and let  $G' = [G, G]$  be its *derived group*, that is the subgroup of  $G$  generated by the commutators  $[x, y] = x^{-1}y^{-1}xy$ , where  $x$  and  $y$  are in  $G$ . Let  $\gamma_i(G)$  be the  $i$ -th term of the lower central series of  $G$  defined inductively by  $\gamma_1(G) = G$  and  $\gamma_{i+1}(G) = [\gamma_i(G), G]$ . The group  $G$  is said to be *nilpotent* if there exists a positive integer  $c$  such that  $\gamma_{c+1}(G) = 1$ ; the smallest integer  $c$  satisfying this equality is called *the nilpotency class* of  $G$ ; if the order of  $G$  is  $p^h$ , where  $p$  is a prime, then the number  $cc(G) = h - c$  is called the *coclass* of  $G$ . It is well known that a  $p$ -group  $G$  is nilpotent. Recall that a group  $G$  is said to be *metabelian* if its derived group  $G'$  is abelian. Finally, a subgroup  $H$  of a group  $G$ , not reduced to an element, is called *maximal* if it is the unique subgroup of  $G$  distinct from  $G$  containing  $H$ .

Consider the group family defined, for any integer  $n \geq 1$ , as follows

$$G_n = \langle \sigma, \tau, \rho : \rho^4 = \sigma^8 = \tau^{2^{n+2}} = 1, \sigma^4 = \tau^{2^{n+1}}, \rho^2 = \tau^{2^n} \sigma^2, \quad (1) \\ [\tau, \sigma] = 1, [\rho, \sigma] = \sigma^{-2}, [\rho, \tau] = \tau^2 \rangle.$$

In this paper, we construct all the subgroups of  $G_n$  of index 2 or 4, we give the abelianization types of these subgroups and we compute the kernel of the transfer map  $V_{G \rightarrow H} : G_n/G'_n \rightarrow H/H'$ , for any subgroup  $H$  of

---

2010 *Mathematics Subject Classification.* 11R11, 11R29, 11R32, 11R37.

*Key words and phrases.* 2-group, metabelian 2-group, capitulation, Hilbert class field.

$G_n$ , defined by the Artin map. Then we apply these results to study the capitulation problem of the 2-ideal classes of some fields  $\mathbf{k}$  satisfying the condition  $\text{Gal}(\mathbf{k}_2^{(2)}/\mathbf{k}) \simeq G_n$ , where  $\mathbf{k}_2^{(2)}$  is the second Hilbert 2-class field of  $\mathbf{k}$ . Finally, we illustrate our results by an example about an imaginary bicyclic biquadratic field.

## 2. RESULTS

Recall first that if  $x, y$  and  $z$  are elements of some group  $G$ , then we easily show that:

$$\begin{aligned} [xy, z] &= [x, z]^y [y, z] \text{ and} \\ [x, yz] &= [x, z][x, y]^z, \\ \text{where } x^y &= y^{-1}xy. \end{aligned}$$

Let  $G_n$  be the group family defined by the Equality (1). Since  $[\tau, \sigma] = 1$ ,  $[\rho, \sigma] = \sigma^{-2}$  and  $[\rho, \tau] = \tau^2$ , so  $G'_n = \langle \sigma^2, \tau^2 \rangle$ , which is abelian. Then  $G_n$  is metabelian and  $G_n/G'_n \simeq (2, 2, 2)$ , since  $\rho^2 = \tau^{2n} \sigma^2$ . Hence  $G_n$  admits seven subgroups of index 2, denote them by  $H_{i,2}$ , and seven subgroups of index 4, denote them by  $H_{i,4}$ , where  $1 \leq i \leq 7$ . These subgroups, their derived groups and their abelianizations are given in the Tables 1 and 2.

TABLE 1. Subgroups of  $G_n$  of index 2.

$i$	conditions	$H_{i,2}$	$H'_{i,2}$	$H_{i,2}/H'_{i,2}$
1		$\langle \sigma, \tau \rangle$	$\langle 1 \rangle$	$(2^2, 2^{n+2})$
2	$n = 1$	$\langle \sigma, \rho \rangle$	$\langle \sigma^2 \rangle$	$(2, 4)$
	$n \geq 2$	$\langle \sigma, \tau^2, \rho \rangle$	$\langle \sigma^2, \tau^4 \rangle$	$(2, 2, 2)$
3		$\langle \rho, \tau \rangle$	$\langle \tau^2 \rangle$	$(2, 4)$
4	$n = 1$	$\langle \sigma\tau, \rho, \sigma^2 \rangle$	$\langle (\sigma\tau)^2, \sigma^4 \rangle$	$(2, 2, 2)$
	$n \geq 2$	$\langle \sigma\tau, \rho \rangle$	$\langle (\sigma\tau)^2 \rangle$	$(2, 4)$
5		$\langle \sigma\rho, \tau \rangle$	$\langle \tau^2 \rangle$	$(2, 4)$
6	$n = 1$	$\langle \tau\rho, \sigma \rangle$	$\langle \sigma^2 \rangle$	$(2, 4)$
	$n \geq 2$	$\langle \tau\rho, \sigma, \tau^2 \rangle$	$\langle \sigma^2, \tau^4 \rangle$	$(2, 2, 2)$
7	$n = 1$	$\langle \sigma\tau, \tau\rho, \sigma^2 \rangle$	$\langle \sigma^4, (\sigma\tau)^4 \rangle$	$(2, 2, 2)$
	$n \geq 2$	$\langle \sigma\tau, \tau\rho \rangle$	$\langle (\sigma\tau)^2 \rangle$	$(2, 4)$

TABLE 2. Subgroups of  $G_n$  of index 4.

$i$	$H_{i,4}$	$H'_{i,4}$	$H_{i,4}/H'_{i,4}$
1	$\langle \sigma, \tau^2 \rangle$	$\langle 1 \rangle$	$(2^{\min(n,2)}, 2^{\max(n+1,3)})$
2	$\langle \tau, \sigma^2 \rangle$	$\langle 1 \rangle$	$(2, 2^{n+2})$
3	$\langle \rho, \tau^2 \rangle$	$\langle \tau^4 \rangle$	$(2, 4)$
4	$\langle \sigma\tau, \sigma^2 \rangle$	$\langle 1 \rangle$	$(4, 4)$ if $n = 1$ $(2, 2^{n+2})$ if $n \geq 2$
5	$\langle \sigma\rho, \tau^2 \rangle$	$\langle \tau^4 \rangle$	$(2, 4)$
6	$\langle \tau\rho, \tau^2 \rangle$	$\langle \tau^4 \rangle$	$(2, 4)$
7	$\langle \sigma\tau\rho, \tau^2 \rangle$	$\langle \tau^4 \rangle$	$(2, 4)$

To check the Tables entries we need the following lemma.

**Lemma 1.** *Let  $G_n = \langle \sigma, \tau, \rho \rangle$  denote the group defined above, then*

1.  $\rho^{-1}\sigma\rho = \sigma^3$ .
2.  $\rho^{-1}\tau\rho = \tau^{-1}$ .
3.  $\rho^2$  commutes with  $\sigma$  and  $\tau$ .
4.  $(\tau\rho)^2 = \rho^2$ .
5.  $(\sigma\tau\rho)^2 = (\sigma\rho)^2 = \rho^2\sigma^4$ .
6. For all  $r \in \mathbb{N}$ ,  $[\rho, \tau^{2^r}] = \tau^{2^{r+1}}$  and  $[\rho, \sigma^{2^r}] = \sigma^{-2^{r+1}}$ .

*Proof.* 1. and 2. are obvious, since  $[\rho, \sigma] = \sigma^{-2}$  and  $[\rho, \tau] = \tau^2$ .

3. As  $\rho^2 = \tau^{2^n}\sigma^2$ , so  $\rho^2 \in \langle \tau, \sigma \rangle$ , which is an abelian group, because  $[\tau, \sigma] = 1$ . Hence the result.

4.  $(\tau\rho)^2 = \tau\rho\tau\rho = \tau\rho^2\rho^{-1}\tau\rho = \tau\rho^2\tau^{-1} = \rho^2$ .

5. We proceed as in 4. to prove these results.

6. Since  $[\rho, \tau] = \tau^2$ , so  $[\rho, \tau^2] = \tau^4$ . By induction, we show that for all  $r \in \mathbb{N}^*$ ,  $[\rho, \tau^{2^r}] = \tau^{2^{r+1}}$ . Similarly, we prove that  $[\rho, \sigma^{2^r}] = \sigma^{-2^{r+1}}$ .  $\square$

Let us now prove some entries of the Tables 1 and 2, using the Lemma 1.

- For  $H_{1,2} = \langle \sigma, \tau, G'_n \rangle = \langle \sigma, \tau \rangle$ , we have  $H'_{1,2} = \langle 1 \rangle$ , since  $[\sigma, \tau] = 1$ . As  $\sigma^8 = \tau^{2^{n+2}} = 1$  and  $\sigma^4 = \tau^{2^{n+1}}$ , so  $H_{1,2}/H'_{1,2} \simeq (2^{\min(2, n+1)}, 2^{\max(3, n+2)}) = (2^2, 2^{n+2})$ .
- For  $H_{2,2} = \langle \sigma, \rho, G'_n \rangle = \langle \sigma, \rho, \tau^2, \sigma^2 \rangle = \langle \sigma, \rho, \tau^2 \rangle$ . If  $n = 1$ , then  $\rho^2 = \tau^2\sigma^2$ , hence  $H_{2,2} = \langle \sigma, \rho \rangle$ . Therefore  $H'_{2,2} = \langle \sigma^2 \rangle$  and  $H_{2,2}/H'_{2,2} \simeq (2, 4)$ . If  $n \geq 2$ , then  $H_{2,2} = \langle \sigma, \rho, \tau^2 \rangle$ . Therefore  $H'_{2,2} = \langle \sigma^2, \tau^4 \rangle$  and  $H_{2,2}/H'_{2,2} \simeq (2, 2, 2)$ .
- For  $H_{1,4} = \langle \sigma, G'_n \rangle = \langle \sigma, \sigma^2, \tau^2 \rangle = \langle \sigma, \tau^2 \rangle$ , we have  $H'_{1,4} = \langle 1 \rangle$ , since  $[\sigma, \tau] = 1$ . As  $\sigma^8 = \tau^{2^{n+2}} = 1$  and  $\sigma^4 = \tau^{2^{n+1}}$ , so  $H_{1,4}/H'_{1,4} \simeq (2^{\min(2, n)}, 2^{\max(3, n+1)})$ .
- For  $H_{2,4} = \langle \tau, G'_n \rangle = \langle \tau, \tau^2, \sigma^2 \rangle = \langle \tau, \sigma^2 \rangle$ , we have  $H'_{2,4} = \langle 1 \rangle$ , hence

$$H_{2,4}/H'_{2,4} \simeq (2^{\min(1,n+1)}, 2^{\max(2n+2)}) = (2, 2^{n+2}).$$

The other entries are checked similarly.

**Proposition 2.** *Let  $G_n$  be the group family defined by the Equality (1), then*

1. *The order of  $G_n$  is  $2^{n+5}$  and that of  $G'_n$  is  $2^{n+2}$ .*
2. *The coclass of  $G_n$  is 3 and its nilpotency class is  $n + 2$ .*

*Proof.* 1. Since  $\sigma^8 = \tau^{2^{n+2}} = 1$  and  $\sigma^4 = \tau^{2^{n+1}}$ , then

$$\langle \sigma, \tau \rangle \simeq (2^{\min(2,n+1)}, 2^{\max(3,n+2)}) = (2^2, 2^{n+2}).$$

Moreover, as  $\rho^2 = \tau^{2^n} \sigma^2$  and  $\rho^4 = \sigma^8 = \tau^{2^{n+2}} = 1$ , so

$$\langle \sigma, \tau, \rho \rangle \simeq (2^2, 2^{n+2}, 2).$$

Thus  $|G_n| = 2^{n+5}$ . Similarly, we prove that  $|G'_n| = 2^{n+2}$ , since  $G'_n = \langle \sigma^2, \tau^2 \rangle$ .

2. The lower central series of  $G_n$  is defined inductively by  $\gamma_1(G_n) = G_n$  and  $\gamma_{i+1}(G_n) = [\gamma_i(G_n), G_n]$ , that is the subgroup of  $G_n$  generated by the set  $\{[a, b] = a^{-1}b^{-1}ab/a \in \gamma_i(G_n), b \in G_n\}$ , so the coclass of  $G_n$  is defined to be  $cc(G_n) = h - c$ , where  $|G_n| = 2^h$  and  $c = c(G_n)$  is the nilpotency class of  $G_n$ . We easily get

$$\gamma_1(G) = G.$$

$$\gamma_2(G) = G' = \langle \sigma^2, \tau^2 \rangle.$$

$$\gamma_3(G) = [G', G] = \langle \sigma^4, \tau^4 \rangle.$$

Then Lemma 1(6) yields that  $\gamma_{j+1}(G) = [\gamma_j(G), G] = \langle \sigma^{2^j}, \tau^{2^j} \rangle$ . Hence, if we put  $v = \max(n + 1, 2) = n + 1$ , then we get  $\gamma_{v+2}(G) = \langle \sigma^{2^{v+1}}, \tau^{2^{v+1}} \rangle = \langle 1 \rangle$  and  $\gamma_{v+1}(G) = \langle \sigma^{2^v}, \tau^{2^v} \rangle \neq \langle 1 \rangle$ . As  $|G| = 2^{n+5}$ , so

$$c(G) = n + 2 \text{ and } cc(G) = n + 5 - (n + 2) = 3.$$

□

We continue with the following results.

**Proposition 3** ([8]). *Let  $H$  be a normal subgroup of  $G_n$ . For  $g \in G_n$ , put  $f = [\langle g \rangle, H : H]$  and let  $\{x_1, x_2, \dots, x_t\}$  be a set of representatives of  $G/\langle g \rangle H$ . The transfer map  $V_{G \rightarrow H} : G/G' \rightarrow H/H'$  is given by the following formula*

$$V_{G \rightarrow H}(gG') = \prod_{i=1}^t x_i^{-1} g^f x_i.H'. \quad (2)$$

Easily, we prove the following corollaries.

**Corollary 4.** *Let  $H$  be a subgroup of  $G_n$  of index 2. If  $G_n/H = \{1, zH\}$ , then  $V_{G \rightarrow H}(gG'_n) = \begin{cases} gz^{-1}gz.H' = g^2[g, z].H' & \text{if } g \in H, \\ g^2.H' & \text{if } g \notin H. \end{cases}$*

**Corollary 5.** *Let  $H$  be a normal subgroup of  $G_n$  of index 4. If  $G_n/H = \{1, zH, z^2H, z^3H\}$ , then*

$$V_{G \rightarrow H}(gG'_n) = \begin{cases} gz^{-1}gz^{-1}gz^{-1}gz^3.H' & \text{if } g \in H, \\ g^4.H' & \text{if } gH = zH, \\ g^2z^{-1}g^2z.H' & \text{if } g \notin H \text{ and } gH \neq zH. \end{cases}$$

**Corollary 6.** *Let  $H$  be a normal subgroup of  $G_n$  of index 4. If  $G_n/H = \{1, z_1H, z_2H, z_3H\}$  with  $z_3 = z_1z_2$ , then*

$$V_{G \rightarrow H}(gG'_n) = \begin{cases} gz_1^{-1}gz_1z_2^{-1}gz_1^{-1}gz_1z_2.H' & \text{if } g \in H, \\ g^2z_i^{-1}g^2z_i.H' & \text{if } gH = z_jH \text{ with } i \neq j. \end{cases}$$

**Theorem 7.** *Keep the previous notations. Then we have:*

1.  $\ker V_{G_n \rightarrow H_{1,2}} = \langle \tau G'_n \rangle$ .
2.  $\ker V_{G_n \rightarrow H_{2,2}} = \begin{cases} \langle \sigma G'_n, \rho G'_n \rangle & \text{if } n = 1. \\ \langle \sigma G'_n, \tau \rho G'_n \rangle & \text{if } n \geq 2. \end{cases}$
3.  $\ker V_{G_n \rightarrow H_{3,2}} = \langle \tau G'_n, \rho G'_n \rangle$ .
4.  $\ker V_{G_n \rightarrow H_{4,2}} = \begin{cases} \langle \tau \rho G'_n, \sigma \rho G'_n \rangle & \text{if } n = 1. \\ \langle \sigma \tau G'_n, \rho G'_n \rangle & \text{if } n \geq 2. \end{cases}$
5.  $\ker V_{G_n \rightarrow H_{5,2}} = \langle \tau G'_n, \sigma \rho G'_n \rangle$ .
6.  $\ker V_{G_n \rightarrow H_{6,2}} = \begin{cases} \langle \sigma G'_n, \tau \rho G'_n \rangle & \text{if } n = 1. \\ \langle \rho G'_n, \sigma G'_n \rangle & \text{if } n \geq 2. \end{cases}$
7.  $\ker V_{G_n \rightarrow H_{7,2}} = \begin{cases} \langle \sigma \tau G'_n, \rho G'_n \rangle & \text{if } n = 1. \\ \langle \sigma \tau G'_n, \tau \rho G'_n \rangle & \text{if } n \geq 2. \end{cases}$
8. *For all  $1 \leq i \leq 7$ ,  $\ker V_{G_n \rightarrow H_{i,4}} = G_n/G'_n$ .*

*Proof.* We prove only some assertions of the theorem.

1. We know, from the Table 1, that  $H_{1,2} = \langle \sigma, \tau \rangle$ , then  $G_n/H_{1,2} = \{1, \rho H_{1,2}\}$  and  $H'_{1,2} = \langle 1 \rangle$ . Hence, by Corollary 4 and Lemma 1, we get

- \*  $V_{G_n \rightarrow H_{1,2}}(\sigma G'_n) = \sigma^2[\sigma, \rho]H'_{1,2} = \sigma^2\sigma^2H'_{1,2} = \sigma^4H'_{1,2}$ .
- \*  $V_{G_n \rightarrow H_{1,2}}(\tau G'_n) = \tau^2[\tau, \rho]H'_{1,2} = \tau^2\tau^{-2}H'_{1,2} = H'_{1,2}$ .
- \*  $V_{G_n \rightarrow H_{1,2}}(\rho G'_n) = \rho^2H'_{1,2} \neq H'_{1,2}$ .
- \*  $V_{G_n \rightarrow H_{1,2}}(\sigma \tau G'_n) = (\sigma \tau)^2[\sigma \tau, \rho]H'_{1,2} = \tau^4H'_{1,2}$ .
- \*  $V_{G_n \rightarrow H_{1,2}}(\tau \rho G'_n) = (\tau \rho)^2H'_{1,2} = \rho^2H'_{1,2} \neq H'_{1,2}$ .
- \*  $V_{G_n \rightarrow H_{1,2}}(\sigma \rho G'_n) = (\sigma \rho)^2H'_{1,2} = \rho^2\sigma^4H'_{1,2} \neq H'_{1,2}$ .
- \*  $V_{G_n \rightarrow H_{1,2}}(\sigma \tau \rho G'_n) = (\sigma \tau \rho)^2H'_{1,2} = \rho^2\sigma^4H'_{1,2} \neq H'_{1,2}$ .

Therefore  $\ker V_{G_n \rightarrow H_{1,2}} = \langle \tau G'_n \rangle$ .

8. We know, from the Table 2, that  $H_{1,4} = \langle \sigma, \tau^2 \rangle$ , then  $G_n/H_{1,4} = \{1, \tau H_{1,4}, \rho H_{1,4}, \tau \rho H_{1,4}\}$  and  $H'_{1,4} = \langle 1 \rangle$ . Hence Corollary 6 and Lemma 1 yield that

$$\begin{aligned}
* \quad & V_{G_n \rightarrow H_{1,4}}(\sigma G'_n) = \sigma^8 H'_{1,4} = H'_{1,4}. \\
* \quad & V_{G_n \rightarrow H_{1,4}}(\tau G'_n) = \tau^2 \rho^{-1} \tau^2 \rho H'_{1,4} = H'_{1,4}. \\
* \quad & V_{G_n \rightarrow H_{1,4}}(\rho G'_n) = \rho^4 H'_{1,4} = H'_{1,4}.
\end{aligned}$$

Therefore  $\ker V_{G_n \rightarrow H_{1,4}} = \langle \sigma G'_n, \tau G'_n, \rho G'_n \rangle$ .

The other assertions are proved similarly.  $\square$

### 3. APPLICATIONS

Let  $\mathbf{k}$  be a number field and  $C_{\mathbf{k},2}$  be its 2-class group, that is the 2-Sylow subgroup of the ideal class group  $C_{\mathbf{k}}$  of  $\mathbf{k}$ , in the wide sens. Let  $\mathbf{k}_2^{(1)}$  be the Hilbert 2-class field of  $\mathbf{k}$  in the wide sens. Then the Hilbert 2-class field tower of  $\mathbf{k}$  is defined inductively by:  $\mathbf{k}_2^{(0)} = \mathbf{k}$  and  $\mathbf{k}_2^{(n+1)} = (\mathbf{k}_2^{(n)})^{(1)}$ , where  $n$  is a positive integer. Let  $\mathbb{M}$  be an unramified extension of  $\mathbf{k}$  and  $C_{\mathbb{M}}$  be the subgroup of  $C_{\mathbf{k}}$  associated to  $\mathbb{M}$  by the class field theory. Denote by  $j_{\mathbf{k} \rightarrow \mathbb{M}} : C_{\mathbf{k}} \rightarrow C_{\mathbb{M}}$  the homomorphism that associate to the class of an ideal  $\mathcal{A}$  of  $\mathbf{k}$  the class of the ideal generated by  $\mathcal{A}$  in  $\mathbb{M}$ , and by  $\mathcal{N}_{\mathbb{M}/\mathbf{k}}$  the norm of the extension  $\mathbb{M}/\mathbf{k}$ .

Throughout all this section, assume that  $\text{Gal}(\mathbf{k}_2^{(2)}/\mathbf{k}) \simeq G_n$ . Hence, according to the class field theory,  $C_{\mathbf{k},2} \simeq G_n/G'_n \simeq (2,2,2)$ , thus  $C_{\mathbf{k},2} = \langle \mathbf{a}, \mathbf{b}, \mathbf{c} \rangle \simeq \langle \sigma G'_n, \tau G'_n, \rho G'_n \rangle$ , where  $(\mathbf{a}, \mathbf{k}_2^{(2)}/\mathbf{k}) = \sigma G'_n$ ,  $(\mathbf{b}, \mathbf{k}_2^{(2)}/\mathbf{k}) = \tau G'_n$  and  $(\mathbf{c}, \mathbf{k}_2^{(2)}/\mathbf{k}) = \rho G'_n$ , with  $(\cdot, \mathbf{k}_2^{(2)}/\mathbf{k})$  denotes the Artin symbol in  $\mathbf{k}_2^{(2)}/\mathbf{k}$ .

It is well known that each subgroup  $H_{i,j}$ , where  $1 \leq i \leq 7$  and  $j = 2$  or  $4$ , of  $C_{\mathbf{k},2}$  is associated, by class field theory, to a unique unramified extension  $\mathbf{K}_{i,j}$  of  $\mathbf{k}_2^{(1)}$  such that  $H_{i,j}/H'_{i,j} \simeq C_{\mathbf{K}_{i,j},2}$ .

Our goal is to study the capitulation problem of the 2-ideal classes of  $\mathbf{k}$  in its unramified quadratic and biquadratic extensions  $\mathbf{K}_{i,2}$  and  $\mathbf{K}_{i,4}$ . By the class field theory, the kernel of  $j_{\mathbf{k} \rightarrow \mathbb{M}}$ ,  $\ker j_{\mathbf{k} \rightarrow \mathbb{M}}$ , is determined by the kernel of the transfer map  $V_{G \rightarrow H} : G/G' \rightarrow H/H'$ , where  $G = \text{Gal}(\mathbf{k}_2^{(2)}/\mathbf{k})$  and  $H = \text{Gal}(\mathbb{M}_2^{(2)}/\mathbb{M})$ .

**Theorem 8.** *Keep the previous notations. Then we have:*

1.  $\ker j_{\mathbf{k} \rightarrow \mathbf{K}_{1,2}} = \langle \mathbf{b} \rangle$ .
2.  $\ker j_{\mathbf{k} \rightarrow \mathbf{K}_{2,2}} = \begin{cases} \langle \mathbf{a}, \mathbf{c} \rangle & \text{if } n = 1. \\ \langle \mathbf{a}, \mathbf{bc} \rangle & \text{if } n \geq 2. \end{cases}$
3.  $\ker j_{\mathbf{k} \rightarrow \mathbf{K}_{3,2}} = \langle \mathbf{b}, \mathbf{c} \rangle$ .
4.  $\ker j_{\mathbf{k} \rightarrow \mathbf{K}_{4,2}} = \begin{cases} \langle \mathbf{ac}, \mathbf{bc} \rangle & \text{if } n = 1. \\ \langle \mathbf{ab}, \mathbf{c} \rangle & \text{if } n \geq 2. \end{cases}$
5.  $\ker j_{\mathbf{k} \rightarrow \mathbf{K}_{5,2}} = \langle \mathbf{b}, \mathbf{ac} \rangle$ .
6.  $\ker j_{\mathbf{k} \rightarrow \mathbf{K}_{6,2}} = \begin{cases} \langle \mathbf{a}, \mathbf{bc} \rangle & \text{if } n = 1. \\ \langle \mathbf{a}, \mathbf{c} \rangle & \text{if } n \geq 2. \end{cases}$

7.  $\ker j_{\mathbf{k} \rightarrow \mathbf{K}_{7,2}} = \begin{cases} \langle \mathbf{ab}, \mathbf{c} \rangle & \text{if } n = 1. \\ \langle \mathbf{ab}, \mathbf{bc} \rangle & \text{if } n \geq 2. \end{cases}$
8. For all  $1 \leq i \leq 7$ ,  $\ker j_{\mathbf{k} \rightarrow \mathbf{K}_{i,4}} = \mathbf{C}_{\mathbf{k},2}$ .
9. The 2-class group of  $\mathbf{k}_2^{(1)}$  is of type  $(2, 2^{n+1})$ .
10. The Hilbert 2-class field tower of  $\mathbf{k}$  stops at  $\mathbf{k}_2^{(2)}$ .

*Proof.* According to the Theorem 7, we have

(1) Since  $\ker V_{G_n \rightarrow H_{1,2}} = \langle \tau G'_n \rangle$ , so  $\ker j_{\mathbf{k} \rightarrow \mathbf{K}_{1,2}} = \langle \mathbf{b} \rangle$ .

(2) Similarly, as  $\ker V_{G_n \rightarrow H_{2,2}} = \begin{cases} \langle \sigma G'_n, \rho G'_n \rangle & \text{if } n = 1, \\ \langle \sigma G'_n, \tau \rho G'_n \rangle & \text{if } n \geq 2, \end{cases}$

then  $\ker j_{\mathbf{k} \rightarrow \mathbf{K}_{2,2}} = \begin{cases} \langle \mathbf{a}, \mathbf{c} \rangle & \text{if } n = 1. \\ \langle \mathbf{a}, \mathbf{bc} \rangle & \text{if } n \geq 2. \end{cases}$

The other assertions are proved similarly.

(9) It is well known that  $\mathbf{C}_{\mathbf{k}_2^{(1)},2} \simeq G'_n$ , where  $\mathbf{C}_{\mathbf{k}_2^{(1)},2}$  is the 2-class group of  $\mathbf{k}_2^{(1)}$ . As  $G'_n \simeq \langle \sigma^2, \tau^2 \rangle \simeq (2^{\min(1,n)}, 2^{\max(2,n+1)}) \simeq (2, 2^{n+1})$ , since  $\sigma^4 = \tau^{2^{n+1}}$ ; so the result derived.

(10)  $H_{1,4}$ ,  $H_{2,4}$  and  $H_{4,4}$  are the three subgroups of index 2 of the group  $H_{1,2}$ , then  $\mathbf{K}_{1,4}$ ,  $\mathbf{K}_{2,4}$  and  $\mathbf{K}_{4,4}$  are the three unramified quadratic extensions of  $\mathbf{K}_{1,2}$ . On the other hand, the 2-class groups of these fields are of rank 2, since, by the class field theory,  $\mathbf{C}_{\mathbf{K}_{i,j},2} \simeq H_{i,j}/H'_{i,j}$  with  $i = 1, 2$  or  $4$  and  $j = 2$  or  $4$ . Thus the Tables 1 and 2 imply that  $\mathbf{C}_{\mathbf{K}_{1,2},2} \simeq (2^2, 2^{n+2})$  and  $\mathbf{C}_{\mathbf{K}_{2,4},2} \simeq (2, 2^{n+2})$ . Hence  $h_2(\mathbf{K}_{2,4}) = \frac{h_2(\mathbf{K}_{1,2})}{2}$ , where  $h_2(K)$  denotes the 2-class number of the field  $K$ . Therefore we can apply Proposition 7 of [5], which says that  $\mathbf{K}_{1,2}$  has an abelian 2-class field tower if and only if it has a quadratic unramified extension  $\mathbf{K}_{2,4}/\mathbf{K}_{1,2}$  such that  $h_2(\mathbf{K}_{2,4}) = \frac{h_2(\mathbf{K}_{1,2})}{2}$ . Thus  $\mathbf{K}_{1,2}$  has abelian 2-class field tower which terminates at the first stage; this implies that the 2-class field tower of  $\mathbf{k}$  terminates at  $\mathbf{k}_2^{(2)}$ , since  $\mathbf{k} \subset \mathbf{K}_{1,2}$ . Moreover, we know, from Proposition 2, that  $|G_n| = 2^{n+5}$  and  $|G'_n| = 2^{n+2}$ , hence  $\mathbf{k}_2^{(1)} \neq \mathbf{k}_2^{(2)}$ .  $\square$

#### 4. EXAMPLE

Let  $p_1 \equiv p_2 \equiv 5 \pmod{8}$  be different primes. Denote by  $\mathbf{k}$  the imaginary bicyclic biquadratic field  $\mathbb{Q}(\sqrt{d}, i)$ , where  $d = 2p_1p_2$ . Let  $\mathbf{k}_2^{(1)}$  be the Hilbert 2-class field of  $\mathbf{k}$ ,  $\mathbf{k}_2^{(2)}$  its second Hilbert 2-class field and  $G$  be the Galois group of  $\mathbf{k}_2^{(2)}/\mathbf{k}$ . According to [4],  $\mathbf{k}$  has an elementary abelian 2-class group  $\mathbf{C}_{\mathbf{k},2}$  of rank 3, that is of type  $(2, 2, 2)$ . Put  $\mathbf{K} = \mathbf{k}(\sqrt{2}) = \mathbb{Q}(\sqrt{2}, \sqrt{p_1p_2}, \sqrt{-1})$ , and let  $q$  denote the unit index of  $\mathbf{K}^+ = \mathbb{Q}(\sqrt{2}, \sqrt{p_1p_2})$ . Denote by  $h_2(-p_1p_2)$  (resp.  $h_2(p_1p_2)$ ) the 2-class number of  $\mathbb{Q}(\sqrt{-p_1p_2})$

(resp.  $\mathbb{Q}(\sqrt{p_1 p_2})$ ), then, from [10],  $h_2(-p_1 p_2) = 2^{m+1}$  with  $m \geq 2$ , and  $h_2(p_1 p_2) = 2^n$  with  $n \geq 1$ . Assume that  $q = 2$ , then, by [3, Lemma 6],  $m = 2$  and  $n \geq 1$ , and by [3, Theorem 2],  $G \simeq G_n$ . The following result is proved in [3], and we give it here to illustrate the results shown above. For more details, the reader can see [3].

**Theorem 9.** *Let  $p_1 \equiv p_2 \equiv 5 \pmod{8}$  be two different primes. Put  $\mathbf{k} = \mathbb{Q}(\sqrt{2p_1 p_2}, i)$ ,  $\mathbf{k}$  has fourteen unramified extensions within his first Hilbert 2-class field,  $\mathbf{k}_2^{(1)}$  (see [1]). Denote by  $\mathbf{C}_{\mathbf{k},2}$  the 2-class group of  $\mathbf{k}$ . Then the following assertions hold.*

1. *Exactly four elements of  $\mathbf{C}_{\mathbf{k},2}$  capitulate in each unramified quadratic extension of  $\mathbf{k}$ , except one where there are only two.*
2. *All the 2-classes of  $\mathbf{k}$  capitulate in each unramified biquadratic extension of  $\mathbf{k}$ .*
3. *The Hilbert 2-class field tower of  $\mathbf{k}$  stops at  $\mathbf{k}_2^{(2)}$  (see [2]).*
4.  *$\mathbf{C}_{\mathbf{k}_2^{(1)},2} \simeq (2, 2^{n+1})$ .*
5. *The coclass of  $G$  is 3 and its nilpotency class is  $n + 2$ .*
6. *The 2-class groups of the unramified quadratic extensions of  $\mathbf{k}$  are of type  $(2, 4)$ ,  $(2, 2, 2)$  or  $(4, 2^{n+2})$ .*
7. *The types of the 2-class groups of the unramified biquadratic extensions of  $\mathbf{k}$  are exactly those given by the Table 2.*

## REFERENCES

- [1] A. Azizi, A. Zekhnini and M. Taous, *On the unramified quadratic and biquadratic extensions of the field  $\mathbb{Q}(\sqrt{d}, i)$* , IJA, Volume **6**, No. 24 (2012), 1169-1173.
- [2] A. Azizi, A. Zekhnini and M. Taous, *On the 2-class field tower of  $\mathbb{Q}(\sqrt{2p_1 p_2}, i)$  and the Galois group of its second Hilbert 2-class field*, Collect. Math. Volume 65, **1** (2014), 131-141.
- [3] A. Azizi, A. Zekhnini and M. Taous, *Structure of  $\text{Gal}(\mathbf{k}_2^{(2)}/\mathbf{k})$  for some fields  $\mathbf{k} = \mathbb{Q}(\sqrt{2p_1 p_2}, i)$  with  $\text{Cl}_2(\mathbf{k}) \simeq (2, 2, 2)$* , Abh. Math. Sem. Univ. Hamburg, Volume 84, **2** (2014), 203-231.
- [4] A. Azizi and M. Taous, *Détermination des corps  $\mathbf{k} = \mathbb{Q}(\sqrt{d}, i)$  dont les 2-groupes de classes sont de type  $(2, 4)$  ou  $(2, 2, 2)$* , Rend. Istit. Mat. Univ. Trieste. **40** (2008), 93-116.
- [5] E. Benjamin, F. Lemmermeyer and C. Snyder, *Real Quadratic Fields with Abelian 2-Class Field tower*, J. of Number Theory, Volume **73**, Number 2, December (1998), pp. 182-194 (13).
- [6] Gerald J. Janusz, *Algebraic Number Fields*, Academic Press, New York, 1973.
- [7] H. Kisilevsky, *Number fields with class number congruent to 4 mod 8 and Hilbert's theorem 94*, J. Number Theory **8** (1976), 271-279.
- [8] K. Miyake, *Algebraic Investigations of Hilbert's Theorem 94, the Principal Ideal theorem and Capitulation Problem*, Expos. Math. **7** (1989), 289-346.



- [9] O. Taussky, *A remark concerning Hilbert's Theorem 94*, J. Reine Angew. Math. **239/240** (1970), 435-438.
- [10] P. Kaplan, *Sur le 2-groupe de classes d'idéaux des corps quadratiques*. J. Reine angew. Math. **283/284** (1976), 313-363.

ABDELMALEK AZIZI AND ABDELKADER ZEKHNINI, DÉPARTEMENT DE MATHÉMATIQUES,  
FACULTÉ DES SCIENCES, UNIVERSITÉ MOHAMMED 1, OUJDA, MOROCCO

*E-mail address:* `abdelmalekazizi@yahoo.fr`

*E-mail address:* `zekha1@yahoo.fr`

MOHAMMED TAOUS, DÉPARTEMENT DE MATHÉMATIQUES, FACULTÉ DES SCIENCES  
ET TECHNIQUES, UNIVERSITÉ MOULAY ISMAIL, ERRACHIDIA, MOROCCO.

*E-mail address:* `taousm@hotmail.com`