

ON SOLVABLE SUBGROUPS OF THE CREMONA GROUP

JULIE DÉSERTE

ABSTRACT. The Cremona group $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ is the group of birational self-maps of $\mathbb{P}_{\mathbb{C}}^2$. Using the action of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ on the Picard-Manin space of $\mathbb{P}_{\mathbb{C}}^2$ we characterize its solvable subgroups. If $G \subset \text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ is solvable, non virtually abelian, and infinite, then up to finite index: either any element of G is of finite order or conjugate to an automorphism of $\mathbb{P}_{\mathbb{C}}^2$, or G preserves a unique fibration that is rational or elliptic, or G is, up to conjugacy, a subgroup of the group generated by one hyperbolic monomial map and the diagonal automorphisms.

We also give some corollaries.

1. INTRODUCTION

We know properties on finite subgroups ([16]), finitely generated subgroups ([6]), uncountable maximal abelian subgroups ([13]), nilpotent subgroups ([14]) of the Cremona group. In this article we focus on solvable subgroups of the Cremona group.

Let G be a group. Recall that $[g, h] = ghg^{-1}h^{-1}$ denotes the commutator of g and h . If Γ_1 and Γ_2 are two subgroups of G , then $[\Gamma_1, \Gamma_2]$ is the subgroup of G generated by the elements of the form $[g, h]$ with $g \in \Gamma_1$ and $h \in \Gamma_2$. We define the *derived series* of G by setting

$$G^{(0)} = G \quad G^{(n+1)} = [G^{(n)}, G^{(n)}] \quad \forall n \geq 0.$$

The *soluble length* $\ell(G)$ of G is defined by

$$\ell(G) = \min \{k \in \mathbb{N} \cup \{0\} \mid G^{(k)} = \{\text{id}\}\}$$

with the convention: $\min \emptyset = \infty$. We say that G is *solvable* if $\ell(G) < \infty$. The study of solvable groups started a long time ago, and any linear solvable subgroup is up to finite index triangularizable (Lie-Kolchin theorem, [23, Theorem 21.1.5]). The assumption "up to finite index" is essential: for instance the subgroup of $\text{PGL}(2, \mathbb{C})$ generated by $\begin{bmatrix} 1 & 0 \\ 1 & -1 \end{bmatrix}$ and $\begin{bmatrix} -1 & 1 \\ 0 & 1 \end{bmatrix}$ is isomorphic to \mathfrak{S}_3 so is solvable but is not triangularizable.

1991 *Mathematics Subject Classification.* 14E07, 14E05.

Author partially supported by ANR Grant "BirPol" ANR-11-JS01-004-01.

Theorem A. *Let G be an infinite, solvable, non virtually abelian subgroup of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$. Then, up to finite index, one of the following holds:*

- (1) *any element of G is either of finite order, or conjugate to an automorphism of $\mathbb{P}_{\mathbb{C}}^2$;*
- (2) *G preserves a unique fibration that is rational, in particular G is, up to conjugacy, a subgroup of $\text{PGL}(2, \mathbb{C}(y)) \rtimes \text{PGL}(2, \mathbb{C})$;*
- (3) *G preserves a unique fibration that is elliptic;*
- (4) *G is, up to birational conjugacy, contained in the group generated by*

$$\{(x^p y^q, x^r y^s), (\alpha x, \beta y) \mid \alpha, \beta \in \mathbb{C}^*\}$$

where $M = \begin{bmatrix} p & q \\ r & s \end{bmatrix}$ denotes an element of $\text{GL}(2, \mathbb{Z})$ with spectral radius > 1 . The group G preserves the two holomorphic foliations defined by the 1-forms $\alpha_1 x dy + \beta_1 y dx$ and $\alpha_2 x dy + \beta_2 y dx$ where (α_1, β_1) and (α_2, β_2) denote the eigenvectors of ${}^t M$.

Furthermore if G is uncountable, case 3. does not hold.

Examples. • Denote by S_3 the group generated by $\begin{bmatrix} 1 & 0 \\ 1 & -1 \end{bmatrix}$ and $\begin{bmatrix} -1 & 1 \\ 0 & 1 \end{bmatrix}$.

As we recall before $S_3 \simeq \mathfrak{S}_3$. Consider now the subgroup G of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ whose elements are the monomial maps $(x^p y^q, x^r y^s)$ with $\begin{bmatrix} p & q \\ r & s \end{bmatrix} \in S_3$. Then any element of G has finite order, and G is solvable; it gives an example of case 1.

- Other examples that illustrate case 1. are the following groups

$$\{(\alpha x + \beta y + \gamma, \delta y + \varepsilon) \mid \alpha, \delta \in \mathbb{C}^*, \beta, \gamma, \varepsilon \in \mathbb{C}\} \subset \text{Aut}(\mathbb{P}_{\mathbb{C}}^2),$$

and

$$E = \{(\alpha x + P(y), \beta y + \gamma) \mid \alpha, \beta \in \mathbb{C}^*, \gamma \in \mathbb{C}, P \in \mathbb{C}[y]\} \subset \text{Aut}(\mathbb{C}^2).$$

- The centralizer of a birational map of $\mathbb{P}_{\mathbb{C}}^2$ that preserves a unique fibration that is rational is virtually solvable ([9, Corollary C]); this example falls in case 2 (see §3.2).
- In [10, Proposition 2.2] Cornulier proved that the group

$$\langle (x+1, y), (x, y+1), (x, xy) \rangle$$

is solvable of length 3, and is not linear over any field; this example falls in case 2. The invariant fibration is given by $x = \text{cst}$.

Remark. In case 1. if there exists an integer d such that $\deg \phi \leq d$ for any ϕ in G , then there exist a smooth projective variety M and a birational map $\psi: M \dashrightarrow \mathbb{P}_{\mathbb{C}}^2$ such that $\psi^{-1}G\psi$ is a solvable subgroup of $\text{Aut}(M)$ (see §3.3). But there is some solvable subgroups G with only elliptic elements that do not satisfy this property:

the group E introduced in Examples 1. Let us mention an other example: Wright constructs abelian subgroups H of $\text{Aut}(\mathbb{C}^2)$ such that any element of H is of finite order, H is unbounded and does not preserve any fibration ([28]).

In §3 we prove Theorem A: we first assume that our solvable, infinite and non virtually abelian subgroup G contains a hyperbolic map, then that it contains a twist and no hyperbolic map, and finally that all elements of G are elliptic. In the last section (§4) we also

- recover the following fact: if G is an infinite nilpotent subgroup of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$, then G does not contain a hyperbolic map;
- remark that we can bound the soluble length of a nilpotent subgroup of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ by the dimension of $\mathbb{P}_{\mathbb{C}}^2$ as Epstein and Thurston did in the context of Lie algebras of rational vector fields on a connected complex manifold;
- give a negative answer to the following question of Favre: does any solvable and finitely generated subgroup G of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ contain a subgroup of finite index whose commutator subgroup is nilpotent? if we assume that $[G, G]$ is not a torsion group;
- give a description of the embeddings of the solvable Baumslag-Solitar groups into the Cremona group.

Acknowledgments. I would like to thank D. Cerveau for always having time to listen to me, for his precious advices, and for pointing out [24]. Thanks to S. Cantat for helpful remarks and comments, and also to J. Ribon for the example of §4.3. I would like also thank P. Sad for his permanent invitation at IMPA where I found the quietness to put my ideas in order. Thanks to the referee whose comments and suggestions help me to improve the presentation.

2. SOME PROPERTIES OF THE BIRATIONAL MAPS

First definitions. Let \mathcal{S} be a projective surface. We will denote by $\text{Bir}(\mathcal{S})$ the group of birational self-maps of \mathcal{S} ; in the particular case of the complex projective plane the group $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ is called **Cremona group**. Take ϕ in $\text{Bir}(\mathcal{S})$, we will denote by $\text{Ind}\phi$ the set of points of indeterminacy of ϕ ; the codimension of $\text{Ind}\phi$ is ≥ 2 .

A birational map from $\mathbb{P}_{\mathbb{C}}^2$ into itself can be written

$$(x : y : z) \dashrightarrow (\phi_0(x, y, z) : \phi_1(x, y, z) : \phi_2(x, y, z))$$

where the ϕ_i 's denote some homogeneous polynomials of the same degree and without common factors of positive degree. The **degree** of ϕ is equal to the degree of the ϕ_i 's. Let ϕ be a birational map of $\mathbb{P}_{\mathbb{C}}^2$. One can define the **dynamical degree** of ϕ as

$$\lambda(\phi) = \lim_{n \rightarrow +\infty} (\deg \phi^n)^{1/n}.$$

More generally let \mathcal{S} be a projective surface, and $\phi: \mathcal{S} \dashrightarrow \mathcal{S}$ be a birational map. Take any norm $\|\cdot\|$ on the Néron-Severi real vector space $N^1(\mathcal{S})$. If ϕ^* is the induced

action by ϕ on $N^1(\mathcal{S})$, we can define

$$\lambda(\phi) = \lim_{n \rightarrow +\infty} \|(\phi^n)^*\|^{1/n}.$$

Remark that this quantity is a birational invariant: if $\psi: \mathcal{S} \dashrightarrow \mathcal{S}'$ is a birational map, then $\lambda(\psi\phi\psi^{-1}) = \lambda(\phi)$.

Classification of birational maps. The algebraic degree is not a birational invariant, but the first dynamical degree is; more precisely one has a classification of birational maps based on the degree growth. Before stating it, let us first introduce the following definitions. Let ϕ be an element of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$. If

- $(\deg \phi^n)_{n \in \mathbb{N}}$ is bounded, we say that ϕ is an *elliptic map*,
- $(\deg \phi^n)_{n \in \mathbb{N}}$ grows linearly, we say that ϕ is a *Jonquières twist*,
- $(\deg \phi^n)_{n \in \mathbb{N}}$ grows quadratically, we say that ϕ is a *Halphen twist*,
- $(\deg \phi^n)_{n \in \mathbb{N}}$ grows exponentially, we say that ϕ is a *hyperbolic map*.

Theorem 2.1 ([15, 20, 3]). *Let ϕ be an element of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$. Then one and only one of the following cases holds*

- ϕ is elliptic, furthermore if ϕ is of infinite order, then ϕ is up to birational conjugacy an automorphism of $\mathbb{P}_{\mathbb{C}}^2$,
- ϕ is a Jonquières twist, ϕ preserves a unique fibration that is rational and every conjugate of ϕ is not an automorphism of a projective surface,
- ϕ is a Halphen twist, ϕ preserves a unique fibration that is elliptic and ϕ is conjugate to an automorphism of a projective surface,
- ϕ is a hyperbolic map.

In the three first cases, $\lambda(\phi) = 1$, in the last one $\lambda(\phi) > 1$.

The Picard-Manin and bubble spaces. Let \mathcal{S} , and \mathcal{S}_i be complex projective surfaces. If $\pi: \mathcal{S}_1 \rightarrow \mathcal{S}$ is a birational morphism, one gets $\pi^*: N^1(\mathcal{S}) \rightarrow N^1(\mathcal{S}_1)$ an embedding of Néron-Severi groups. Take two birational morphisms $\pi_1: \mathcal{S}_1 \rightarrow \mathcal{S}$ and $\pi_2: \mathcal{S}_2 \rightarrow \mathcal{S}$; the morphism π_2 is *above* π_1 if $\pi_1^{-1}\pi_2$ is regular. Starting with two birational morphisms one can always find a third one that covers the two first. Therefore the inductive limit of all groups $N^1(\mathcal{S}_i)$ for all surfaces \mathcal{S}_i above \mathcal{S} is well-defined; it is the *Picard-Manin space* $Z_{\mathcal{S}}$ of \mathcal{S} . For any birational map π , π^* preserves the intersection form and maps nef classes to nef classes hence the limit space $Z_{\mathcal{S}}$ is endowed with an intersection form of signature $(1, \infty)$ and a nef cone.

Let \mathcal{S} be a complex projective surface. Consider all complex and projective surfaces \mathcal{S}_i above \mathcal{S} , that is all birational morphisms $\pi_i: \mathcal{S}_i \rightarrow \mathcal{S}$. If p (resp. q) is a point of a complex projective surface \mathcal{S}_1 (resp. \mathcal{S}_2), and if $\pi_1: \mathcal{S}_1 \rightarrow \mathcal{S}$ (resp. $\pi_2: \mathcal{S}_2 \rightarrow \mathcal{S}$) is a birational morphism, then p is identified with q if $\pi_1^{-1}\pi_2$ is a local isomorphism in a neighborhood of q that maps q onto p . The *bubble space* $\mathcal{B}(\mathcal{S})$ is the union of all points of all surfaces above \mathcal{S} modulo the equivalence relation induced by this identification. If p belongs to $\mathcal{B}(\mathcal{S})$ represented by a point p on a surface $\mathcal{S}_i \rightarrow \mathcal{S}$,

denote by E_p the exceptional divisor of the blow-up of p and by e_p its divisor class viewed as a point in \mathcal{Z}_S . The following properties are satisfied

$$\begin{cases} e_p \cdot e_q = 0 & \text{if } p \neq q \\ e_p \cdot e_p = -1 \end{cases}$$

Hyperbolic space \mathbb{H}_S . Embed $N^1(S)$ as a subgroup of \mathcal{Z}_S ; this finite dimensional lattice is orthogonal to e_p for any $p \in \mathcal{B}(S)$, and

$$\mathcal{Z}_S = \left\{ D + \sum_{p \in \mathcal{B}(S)} a_p e_p \mid D \in N^1(S), a_p \in \mathbb{R} \right\}.$$

The **completed Picard-Manin space** $\overline{\mathcal{Z}}_S$ of S is the L^2 -completion of \mathcal{Z}_S ; in other words

$$\overline{\mathcal{Z}}_S = \left\{ D + \sum_{p \in \mathcal{B}(S)} a_p e_p \mid D \in N^1(S), a_p \in \mathbb{R}, \sum a_p^2 < +\infty \right\}.$$

The intersection form extends as an intersection form with signature $(1, \infty)$ on $\overline{\mathcal{Z}}_S$. Let

$$\overline{\mathcal{Z}}_S^+ = \{ d \in \overline{\mathcal{Z}}_S \mid d \cdot c \geq 0 \quad \forall c \in \overline{\mathcal{Z}}_S \}$$

be the nef cone of $\overline{\mathcal{Z}}_S$ and

$$\mathcal{L}\overline{\mathcal{Z}}_S = \{ d \in \overline{\mathcal{Z}}_S \mid d \cdot d = 0 \}$$

be the light cone of $\overline{\mathcal{Z}}_S$.

The **hyperbolic space** \mathbb{H}_S of S is then defined by

$$\mathbb{H}_S = \{ d \in \overline{\mathcal{Z}}_S^+ \mid d \cdot d = 1 \}.$$

Let us remark that \mathbb{H}_S is an infinite dimensional analogue of the classical hyperbolic space \mathbb{H}^n . The **distance** on \mathbb{H}_S is defined by

$$\cosh(\text{dist}(d, d')) = d \cdot d' \quad \forall d, d' \in \mathbb{H}_S.$$

The **geodesics** are intersections of \mathbb{H}_S with planes. The projection of \mathbb{H}_S onto $\mathbb{P}(\overline{\mathcal{Z}}_S)$ is one-to-one, and the boundary of its image is the projection of the cone of isotropic vectors of $\overline{\mathcal{Z}}_S$. Hence

$$\partial\mathbb{H}_S = \{ \mathbb{R}_+ d \mid d \in \overline{\mathcal{Z}}_S^+, d \cdot d = 0 \}.$$

Isometries of \mathbb{H}_S . If $\pi: S' \rightarrow S$ is a birational morphism, we get a canonical isometry π^* (and not only an embedding) between \mathbb{H}_S and $\mathbb{H}_{S'}$. This allows to define an action of $\text{Bir}(S)$ on \mathbb{H}_S . Consider a birational map ϕ on a complex projective surface S . There exists a surface S' , and $\pi_1: S' \rightarrow S$, $\pi_2: S' \rightarrow S$ two morphisms such that $\phi = \pi_2 \pi_1^{-1}$. One can define ϕ_\bullet by

$$\phi_\bullet = (\pi_2^*)^{-1} \pi_1^*;$$

in fact, one gets a faithful representation of $\text{Bir}(S)$ into the group of isometries of \mathbb{H}_S (see [6]).

The isometries of \mathbb{H}_S are classified in three types ([4, 19]). The **translation length** of an isometry ϕ_\bullet of \mathbb{H}_S is defined by

$$L(\phi_\bullet) = \inf \{ \text{dist}(p, \phi_\bullet(p)) \mid p \in \mathbb{H}_S \}$$

If the infimum is a minimum, then

- either it is equal to 0 and ϕ_\bullet has a fixed point in \mathbb{H}_S , ϕ_\bullet is thus **elliptic**,
- or it is positive and ϕ_\bullet is **hyperbolic**. Hence the set of points $p \in \mathbb{H}_S$ such that $\text{dist}(p, \phi_\bullet(p))$ is equal to $L(\phi_\bullet)$ is a geodesic line $\text{Ax}(\phi_\bullet) \subset \mathbb{H}_S$. Its boundary points are represented by isotropic vectors $\omega(\phi_\bullet)$ and $\alpha(\phi_\bullet)$ in $\overline{\mathcal{Z}}_S$ such that

$$\phi_\bullet(\omega(\phi_\bullet)) = \lambda(\phi) \omega(\phi_\bullet) \quad \phi_\bullet(\alpha(\phi_\bullet)) = \frac{1}{\lambda(\phi)} \alpha(\phi_\bullet)$$

The axis $\text{Ax}(\phi_\bullet)$ of ϕ_\bullet is the intersection of \mathbb{H}_S with the plane containing $\omega(\phi_\bullet)$ and $\alpha(\phi_\bullet)$; furthermore ϕ_\bullet acts as a translation of length $L(\phi_\bullet) = \log \lambda(\phi)$ along $\text{Ax}(\phi_\bullet)$ (see [8, Remark 4.5]). For all p in \mathbb{H}_S one has

$$\lim_{k \rightarrow +\infty} \frac{\phi_\bullet^{-k}(p)}{\lambda(\phi)} = \alpha(\phi_\bullet) \quad \lim_{k \rightarrow +\infty} \frac{\phi_\bullet^k(p)}{\lambda(\phi)} = \omega(\phi_\bullet).$$

When the infimum is not realized, $L(\phi_\bullet) = 0$ and ϕ_\bullet is **parabolic**: ϕ_\bullet fixes a unique line in $\overline{\mathcal{Z}}_S$; this line is fixed pointwise, and all orbits $\phi_\bullet^n(p)$ in \mathbb{H}_S accumulate to the corresponding boundary point when n goes to $\pm\infty$.

There is a strong relationship between this classification and the classification of birational maps of the complex projective plane ([6, Theorem 3.6]): if ϕ is an element of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$, then

- ϕ_\bullet is an elliptic isometry if and only if ϕ is an elliptic map;
- ϕ_\bullet is a parabolic isometry if and only if ϕ is a twist;
- ϕ_\bullet is a hyperbolic isometry if and only if ϕ is a hyperbolic map.

Tits alternative. Cantat proved the Tits alternative for the Cremona group ([6, Theorem C]): let G be a finitely generated subgroup of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$, then

- either G contains a free non abelian subgroup,
- or G contains a subgroup of finite index that is solvable.

As a consequence he studied finitely generated and solvable subgroups of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ without torsion ([6, Theorem 7.3]): let G be such a group, there exists a subgroup G_0 of G of finite index such that

- either G_0 is abelian,
- or G_0 preserves a foliation.

3. PROOF OF THEOREM A

3.1. Solvable groups of birational maps containing a hyperbolic map. Let us recall the following criterion (for its proof see for example [11]) used on many occasions by Klein, and also by Tits ([26]) known as Ping-Pong Lemma: *let H be a group acting on a set X , let Γ_1, Γ_2 be two subgroups of H , and let Γ be the subgroup generated by Γ_1 and Γ_2 . Assume that Γ_1 contains at least three elements, and Γ_2 at least two elements. Suppose that there exist two non-empty subsets X_1, X_2 of X such that*

$$X_2 \not\subset X_1, \quad \gamma(X_2) \subset X_1 \quad \forall \gamma \in \Gamma_1 \setminus \{\text{id}\}, \quad \gamma'(X_1) \subset X_2 \quad \forall \gamma' \in \Gamma_2 \setminus \{\text{id}\}.$$

*Then Γ is isomorphic to the free product $\Gamma_1 * \Gamma_2$.* The Ping-Pong argument allows us to prove the following:

Lemma 3.1. *A solvable, non abelian subgroup of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ cannot contain two hyperbolic maps ϕ and ψ such that $\{\omega(\phi_{\bullet}), \alpha(\phi_{\bullet})\} \neq \{\omega(\psi_{\bullet}), \alpha(\psi_{\bullet})\}$.*

Proof. Assume by contradiction that $\{\omega(\phi_{\bullet}), \alpha(\phi_{\bullet})\} \neq \{\omega(\psi_{\bullet}), \alpha(\psi_{\bullet})\}$. Then the Ping-Pong argument implies that there exist two integers n and m such that ψ^n and ϕ^m generate a subgroup of G isomorphic to the free group F_2 (see [6]). But $\langle \phi, \psi \rangle$ is a solvable group: contradiction. \square

Let G be an infinite solvable, non virtually abelian, subgroup of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$. Assume that G contains a hyperbolic map ϕ . Let $\alpha(\phi_{\bullet})$ and $\omega(\phi_{\bullet})$ be the two fixed points of ϕ_{\bullet} on $\partial\mathbb{H}_{\mathbb{P}_{\mathbb{C}}^2}$, and $Ax(\phi_{\bullet})$ be the geodesic passing through these two points. As G is solvable there exists a subgroup of G of index ≤ 2 that preserves $\alpha(\phi_{\bullet})$, $\omega(\phi_{\bullet})$, and $Ax(\phi_{\bullet})$ (see [6, Theorem 6.4]); let us still denote by G this subgroup. Note that there is no twist in G since a parabolic isometry has a unique fixed point on $\partial\mathbb{H}_{\mathbb{P}_{\mathbb{C}}^2}$. One has a morphism $\kappa: G \rightarrow \mathbb{R}_{>0}$ such that

$$\psi_{\bullet}(\ell) = \kappa(\psi)\ell$$

for any ℓ in $\overline{\mathbb{Z}_{\mathbb{P}_{\mathbb{C}}^2}}$ lying on $Ax(\phi_{\bullet})$.

The kernel of κ is an infinite subgroup that contains only elliptic maps. Indeed the set of elliptic elements of G coincides with $\ker \kappa$; and $[G, G] \subset \ker \kappa$ so if $\ker \kappa$ is finite, G is abelian up to finite index which is by assumption impossible.

Gap property. If ψ is an hyperbolic birational map of G , then $\kappa(\psi) = L(\psi_{\bullet}) = \log \lambda(\psi)$. Recall that $\lambda(\phi)$ is an algebraic integer with all Galois conjugates in the unit disk, that is a Salem number, or a Pisot number. The smallest known number is the Lehmer number $\lambda_L \simeq 1,176$ which is a root of $X^{10} + X^9 - X^7 - X^6 - X^5 - X^4 - X^3 + X + 1$. Blanc and Cantat prove in [2, Corollary 2.7] that there is a gap in the dynamical spectrum $\Lambda = \{\lambda(\phi) \mid \phi \in \text{Bir}(\mathbb{P}_{\mathbb{C}}^2)\}$: there is no dynamical degree in $]1, \lambda_L[$.

The gap property implies that in fact κ is a morphism from G to a subgroup of $\mathbb{R}_{>0}$ isomorphic to \mathbb{Z} .

Elliptic subgroups of the Cremona group with a large normalizer. Consider in $\mathbb{P}_{\mathbb{C}}^2$ the complement of the union of the three lines $\{x = 0\}$, $\{y = 0\}$ and $\{z = 0\}$. Denote by \mathcal{U} this open set isomorphic to $\mathbb{C}^* \times \mathbb{C}^*$. One has an action of $\mathbb{C}^* \times \mathbb{C}^*$ on \mathcal{U} by translation. Furthermore $\mathrm{GL}(2, \mathbb{Z})$ acts on \mathcal{U} by monomial maps

$$\begin{bmatrix} p & q \\ r & s \end{bmatrix} \mapsto ((x, y) \mapsto (x^p y^q, x^r y^s))$$

One thus has an injective morphism from $(\mathbb{C}^* \times \mathbb{C}^*) \rtimes \mathrm{GL}(2, \mathbb{Z})$ into $\mathrm{Bir}(\mathbb{P}_{\mathbb{C}}^2)$. Let G_{toric} be its image.

One can now apply [12, Theorem 4] that says that if there exists a short exact sequence

$$1 \longrightarrow A \longrightarrow N \longrightarrow B \longrightarrow 1$$

where $N \subset \mathrm{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ contains at least one hyperbolic element, and $A \subset \mathrm{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ is an infinite and elliptic¹ group, then N is up to conjugacy a subgroup of G_{toric} . Hence up to birational conjugacy $G \subset G_{\mathrm{toric}}$. Recall now that if ψ is a hyperbolic map of the form $(x^a y^b, x^c y^d)$, then to preserve $\alpha(\psi_{\bullet})$ and $\omega(\psi_{\bullet})$ is equivalent to preserve the eigenvectors of the matrix $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$. We can now thus state:

Proposition 3.2. *Let G be an infinite solvable, non virtually abelian, subgroup of $\mathrm{Bir}(\mathbb{P}_{\mathbb{C}}^2)$. If G contains a hyperbolic birational map, then G is, up to conjugacy and finite index, a subgroup of the group generated by*

$$\{(x^p y^q, x^r y^s), (\alpha x, \beta y) \mid \alpha, \beta \in \mathbb{C}^*\}$$

where $\begin{bmatrix} p & q \\ r & s \end{bmatrix}$ denotes an element of $\mathrm{GL}(2, \mathbb{Z})$ with spectral radius > 1 .

3.2. Solvable groups with a twist. Consider a solvable, non abelian subgroup G of $\mathrm{Bir}(\mathbb{P}_{\mathbb{C}}^2)$. Let us assume that G contains a twist ϕ ; the map ϕ preserves a unique fibration \mathcal{F} that is rational or elliptic. Let us prove that any element of G preserves \mathcal{F} . Denote by $\alpha(\phi_{\bullet}) \in \partial \mathbb{H}_{\mathbb{P}_{\mathbb{C}}^2}$ the fixed point of ϕ_{\bullet} . Take one element in $\mathcal{L}\overline{\mathcal{Z}}_{\mathbb{P}_{\mathbb{C}}^2}$ still denoted $\alpha(\phi_{\bullet})$ that represents $\alpha(\phi_{\bullet})$. Take $\varphi \in G$ such that $\varphi(\alpha(\phi_{\bullet})) \neq \alpha(\phi_{\bullet})$. Then $\psi = \varphi\phi\varphi^{-1}$ is parabolic and fixes the unique element $\alpha(\psi_{\bullet})$ of $\mathcal{L}\overline{\mathcal{Z}}_{\mathbb{P}_{\mathbb{C}}^2}$ proportional to $\varphi(\alpha(\phi_{\bullet}))$. Take $\varepsilon > 0$ such that $\mathcal{U}(\alpha(\phi_{\bullet}), \varepsilon) \cap \mathcal{U}(\alpha(\psi_{\bullet}), \varepsilon) = \emptyset$ where

$$\mathcal{U}(\alpha, \varepsilon) = \{\ell \in \mathcal{L}\overline{\mathcal{Z}}_{\mathbb{P}_{\mathbb{C}}^2} \mid \alpha \cdot \ell < \varepsilon\}.$$

Since ψ_{\bullet} is parabolic, then for n large enough $\psi_{\bullet}^n(\mathcal{U}(\alpha(\phi_{\bullet}), \varepsilon))$ is included in a $\mathcal{U}(\alpha(\psi_{\bullet}), \varepsilon)$. For m sufficiently large $\phi_{\bullet}^m \psi_{\bullet}^n(\mathcal{U}(\alpha(\phi_{\bullet}), \varepsilon)) \subset (\mathcal{U}(\alpha(\phi_{\bullet}), \varepsilon/2)) \subsetneq (\mathcal{U}(\alpha(\phi_{\bullet}), \varepsilon))$; hence $\phi_{\bullet}^m \psi_{\bullet}^n$ is hyperbolic. You can by this way build two hyperbolic maps whose sets of fixed points are distinct: this gives a contradiction with Lemma 3.1. So for any $\varphi \in G$ one has : $\alpha(\phi_{\bullet}) = \alpha(\varphi_{\bullet})$; one can thus state the following result.

1. A subgroup of $\mathrm{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ is elliptic if it fixes a point in $\mathbb{H}_{\mathbb{P}_{\mathbb{C}}^2}$.

Proposition 3.3. *Let G be a solvable, non abelian subgroup of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ that contains a twist ϕ . Then*

- *if ϕ is a Jonquières twist, then G preserves a rational fibration, that is up to birational conjugacy G is a subgroup of $\text{PGL}(2, \mathbb{C}(y)) \times \text{PGL}(2, \mathbb{C})$,*
- *if ϕ is a Halphen twist, then G preserves an elliptic fibration.*

Remark 3.4. Both cases are mutually exclusive.

Note that if G is a subgroup of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ that preserves an elliptic fibration, then G is countable ([5]). Let us explain briefly why. A smooth rational projective surface \mathcal{S} is a **Halphen surface** if there exists an integer $m > 0$ such that the linear system $| -mK_{\mathcal{S}} |$ is of dimension 1, has no fixed component, and has no base point. The smallest positive integer for which \mathcal{S} satisfies such a property is the **index** of \mathcal{S} . If \mathcal{S} is a Halphen surface of index m , then $K_{\mathcal{S}}^2 = 0$ and, by the genus formula, the linear system $| -mK_{\mathcal{S}} |$ defines a genus 1 fibration $\mathcal{S} \rightarrow \mathbb{P}_{\mathbb{C}}^1$. This fibration is **relatively minimal** in the sense that there is no (-1) -curve contained in a fiber. The following properties are equivalent:

- \mathcal{S} is a Halphen surface of index m ,
- there exists an irreducible pencil of curves of degree $3m$ with 9 base points of multiplicity m in $\mathbb{P}_{\mathbb{C}}^2$ such that \mathcal{S} is the blow-up of the 9 base points and $| -mK_{\mathcal{S}} |$ is the proper transform of this pencil (the base points set may contain infinitely near points).

As a corollary of the classification of relatively minimal elliptic surfaces the relative minimal model of a rational elliptic surface is a Halphen surface of index m ([22, Chapter 2, §10]). Up to conjugacy G is a subgroup of $\text{Aut}(\mathcal{S})$ where \mathcal{S} denotes a Halphen surface of index m . The action of G on $\text{NS}(\mathcal{S})$ is almost faithful, and G is a discrete (it preserves the integral structure of $\text{NS}(\mathcal{S})$) and virtually abelian (it preserves the intersection form and the class of the elliptic fibration) subgroup of $\text{Aut}(\mathcal{S})$. So one has:

Corollary 3.5. *If G is an uncountable, solvable, non abelian subgroup of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$, then G doesn't contain a Halphen twist.*

Example 3.6. Let us come back to the example given in §1. If $\phi \in \text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ preserves a unique fibration that is rational then one can assume that up to birational conjugacy this fibration is given, in the affine chart $z = 1$, by $y = \text{cst}$. If ϕ preserves $y = \text{cst}$ fiberwise, then

- ϕ is contained in a maximal abelian subgroup denoted $\text{Ab}(\phi)$ that preserves $y = \text{cst}$ fiberwise ([13]),
- the centralizer of ϕ is a finite extension of $\text{Ab}(\phi)$ (see [9, Theorem B]).

This allows us to establish that if ϕ preserves a fibration not fiberwise, then the centralizer of ϕ is virtually solvable. For instance if $\phi = (x + a(y), y + 1)$ (resp.

$(b(y)x, \beta y)$ or $(x + a(y), \beta y)$ with $\beta \in \mathbb{C}^*$ of infinite order) preserves a unique fibration, then the centralizer of ϕ is solvable and metabelian ([9, Propositions 5.1 and 5.2]).

3.3. Solvable groups with no hyperbolic map, and no twist. Let M be a smooth, irreducible, complex, projective variety of dimension n . Fix a Kähler form κ on M . If ℓ is a positive integer, denote by $\pi_i: M^\ell \rightarrow M$ the projection onto the i -th factor. The manifold M^ℓ is then endowed with the Kähler form $\sum_{i=1}^{\ell} \pi_i^* \kappa$ which induces a Kähler metric. To any $\phi \in \text{Bir}(M)$ one can associate its graph $\Gamma_\phi \subset M \times M$ defined as the Zariski closure of

$$\{(z, \phi(z)) \in M \times M \mid z \in M \setminus \text{Ind}\phi\}.$$

By construction Γ_ϕ is an irreducible subvariety of $M \times M$ of dimension n . Both projections $\pi_1, \pi_2: M \times M \rightarrow M$ restrict to a birational morphism $\pi_1, \pi_2: \Gamma_\phi \rightarrow M$.

The **total degree** $\text{tdeg}\phi$ of $\phi \in \text{Bir}(M)$ is defined as the volume of Γ_ϕ with respect to the fixed metric on $M \times M$:

$$\text{tdeg}\phi = \int_{\Gamma_\phi} (\pi_1^* \kappa + \pi_2^* \kappa)^n = \int_{M \setminus \text{Ind}\phi} (\kappa + \phi^* \kappa)^n.$$

Let $d \geq 1$ be a natural integer, and set

$$\text{Bir}_d(M) = \{\phi \in \text{Bir}(M) \mid \text{tdeg}\phi \leq d\}.$$

A subgroup G of $\text{Bir}(M)$ has **bounded degree** if it is contained in $\text{Bir}_d(M)$ for some $d \in \mathbb{N}^*$.

Any subgroup G of $\text{Bir}(M)$ that has bounded degree can be regularized, that is up to birational conjugacy all indeterminacy points of all elements of G disappear simultaneously:

Theorem 3.7 ([27]). *Let M be a complex projective variety, and let G be a subgroup of $\text{Bir}(M)$. If G has bounded degree, there exists a smooth, complex, projective variety M' , and a birational map $\psi: M' \dashrightarrow M$ such that $\psi^{-1}G\psi$ is a subgroup of $\text{Aut}(M')$.*

The proof of this result can be found in [21, 29]; an heuristic idea appears in [7].

4. APPLICATIONS

4.1. Nilpotent subgroups of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$. Let us recall that if G is a group, the **descending central series** of G is defined by

$$C^0G = G \quad C^{n+1}G = [G, C^nG] \quad \forall n \geq 0.$$

We say that G is **nilpotent** if there exists $j \geq 0$ such that $C^jG = \{\text{id}\}$. If j is the minimum non-negative number with such a property we say that G is of **nilpotent class** j . Nilpotent subgroups of the Cremona group have been described:

Theorem 4.1 ([14]). *Let G be a nilpotent subgroup of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$. Then*

- *either G is up to finite index metabelian,*
- *or G is a torsion group.*

We find an alternative proof of [14, Lemma 4.2] for G infinite:

Lemma 4.2. *Let G be an infinite, nilpotent, non virtually abelian subgroup of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$. Then G does not contain a hyperbolic map.*

Proof. The group G is also solvable. Assume by contradiction that G contains a hyperbolic map; then according to Theorem A up to birational conjugacy and finite index there exists $Y \subset \mathbb{C}^* \times \mathbb{C}^*$ infinite such that G is generated by $\phi = (x^p y^q, x^r y^s)$ and

$$\{(\alpha x, \beta y) \mid (\alpha, \beta) \in Y\}.$$

The group $C^1 G$ contains

$$\{[\phi, (\alpha x, \beta y)] \mid (\alpha, \beta) \in Y\} = \{(\alpha^{p-1} \beta^q x, \alpha^r \beta^{s-1} y) \mid (\alpha, \beta) \in Y\}$$

that is infinite since Y is infinite. Suppose that $C^i G$ contains the infinite set

$$\{(\alpha^{\ell_i} \beta^{n_i} x, \alpha^{k_i} \beta^{m_i} y) \mid (\alpha, \beta) \in Y\}$$

(ℓ_i, n_i, k_i and m_i are some functions in p, q, r and s); then $C^{i+1} G$ contains

$$\{(\alpha^{(p-1)\ell_i + qm_i} \beta^{(p-1)k_i + qn_i} x, \alpha^{r\ell_i + qm_i} \beta^{rk_i + (s-1)m_i} y) \mid (\alpha, \beta) \in Y\}$$

that is still infinite. \square

So any nilpotent and infinite subgroup of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ falls in case 1., 2., 3. of Theorem A. If it falls in case 2. or 3. then G is virtually metabelian ([14, Proof of Theorem 1.1.]). Finally if G falls in case 1., we can prove as in [14] that either G is a torsion group, or G is virtually metabelian.

4.2. Soluble length of a nilpotent subgroup of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$. Let us recall the following statement due to Epstein and Thurston ([17]): let M be a connected complex manifold. Let \mathfrak{h} be a nilpotent Lie subalgebra of the complex vector space of rational vector fields on M . Then $\mathfrak{h}^{(n)} = \{0\}$ if $n \geq \dim M$; hence the soluble length of \mathfrak{h} is bounded by the dimension of M . We have a similar statement in the context of birational maps; indeed a direct consequence of Theorem 4.1 is the following property: let $G \subset \text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ be a nilpotent subgroup of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ that is not a torsion group, then the soluble length of G is bounded by the dimension of $\mathbb{P}_{\mathbb{C}}^2$.

4.3. Favre's question. In [18] Favre asked few questions; among them there is the following: does any solvable, finitely generated subgroup G of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ contain a subgroup H of finite index such that $[H, H]$ is nilpotent? We will prove that the answer is no if $[G, G]$ is not a torsion group.

Take G a solvable and finitely generated subgroup of the Cremona group; besides suppose that $[G, G]$ is not a torsion group. Assume that the answer of Favre's question is yes. Up to finite index one can assume that $[G, G]$ is nilpotent. According to

Theorem 4.1 the group $G^{(1)} = [G, G]$ is up to finite index metabelian; in other words up to finite index $G^{(2)} = [G^{(1)}, G^{(1)}]$ is abelian and so $G^{(3)} = [G^{(2)}, G^{(2)}] = \{\text{id}\}$, *i.e.* the soluble length of G is bounded by 3 up to finite index. Consider the subgroup

$$\langle (x+y^2, y), (x(1+y), y), \left(x, \frac{y}{1+y}\right), (x, 2y) \rangle$$

of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$. It is solvable of length 4 (*see* [24]): contradiction.

4.4. Baumslag-Solitar groups. For any integers m, n such that $mn \neq 0$, the Baumslag-Solitar group $\text{BS}(m; n)$ is defined by the following presentation

$$\text{BS}(m; n) = \langle r, s \mid rs^m r^{-1} = s^n \rangle$$

In [3] we prove that there is no embedding of $\text{BS}(m; n)$ into $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ as soon as $|n|, |m|$, and 1 are distinct; it corresponds exactly to the case $\text{BS}(m; n)$ is not solvable. Indeed $\text{BS}(m; n)$ is solvable if and only if $|m| = 1$ or $|n| = 1$ (*see* [25, Proposition A.6]).

Proposition 4.3. *Let ρ be an embedding of $\text{BS}(1; n) = \langle r, s \mid rsr^{-1} = s^n \rangle$, with $n \neq 1$, into the Cremona group. Then*

- *the image of ρ doesn't contain a hyperbolic map,*
- *and*

$$\rho(s) = (x, y+1) \quad \rho(r) = (v(x), n(y+a(x)))$$

with $v \in \text{PGL}(2, \mathbb{C})$ and $a \in \mathbb{C}(x)$.

Proof. According to [3, Proposition 6.2, Lemma 6.3] one gets that $\rho(s) = (x, y+1)$ and $\rho(r) = (v(x), n(y+a(x)))$ for some $v \in \text{PGL}(2, \mathbb{C})$ and $a \in \mathbb{C}(x)$.

Furthermore $\rho(s)$ can neither be conjugate to an automorphism of the form $(\alpha x, \beta y)$ (*see* [1]), nor to a hyperbolic birational map of the form $(\gamma x^p y^q, \delta x^r y^s)$ with $\begin{bmatrix} p & q \\ r & s \end{bmatrix} \in \text{GL}(2, \mathbb{Z})$ of spectral radius > 1 . As a consequence Proposition 3.2 implies that $\rho(\text{BS}(1; n))$ does not contain a hyperbolic birational map. \square

REFERENCES

- [1] J. Blanc. Conjugacy classes of affine automorphisms of \mathbb{K}^n and linear automorphisms of \mathbb{P}^n in the Cremona groups. *Manuscripta Math.*, 119(2):225–241, 2006.
- [2] J. Blanc and S. Cantat. Dynamical degrees of birational transformations of projective surfaces, [arXiv:1307.0361](https://arxiv.org/abs/1307.0361). *J. Amer. Math. Soc.*, to appear.
- [3] J. Blanc and J. Déserti. Degree growth of birational maps of the plane. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, To appear.
- [4] M. R. Bridson and A. Haefliger. *Metric spaces of non-positive curvature*, volume 319 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin, 1999.

- [5] S. Cantat. *Dynamique des automorphismes des surfaces complexes compactes*. PhD thesis, École Normale Supérieure de Lyon, 1999.
- [6] S. Cantat. Sur les groupes de transformations birationnelles des surfaces. *Ann. of Math. (2)*, 174(1):299–340, 2011.
- [7] S. Cantat. Morphisms between Cremona groups, and characterization of rational varieties. *Compos. Math.*, 150(7):1107–1124, 2014.
- [8] S. Cantat and S. Lamy. Normal subgroups in the Cremona group. *Acta Math.*, 210(1):31–94, 2013. With an appendix by Yves de Cornulier.
- [9] D. Cerveau and J. Déserti. Centralisateurs dans le groupe de Jonquières. *Michigan Math. J.*, 61(4):763–783, 2012.
- [10] Y. Cornulier. Nonlinearity of some subgroups of the planar Cremona group. *Unpublished manuscript, 2013*, <http://www.normalesup.org/~cornulier/crelin.pdf>.
- [11] P. de la Harpe. *Topics in geometric group theory*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 2000.
- [12] T. Delzant and P. Py. Kähler groups, real hyperbolic spaces and the Cremona group. *Compos. Math.*, 148(1):153–184, 2012. With an appendix by S. Cantat.
- [13] J. Déserti. Sur les automorphismes du groupe de Cremona. *Compos. Math.*, 142(6):1459–1478, 2006.
- [14] J. Déserti. Sur les sous-groupes nilpotents du groupe de Cremona. *Bull. Braz. Math. Soc. (N.S.)*, 38(3):377–388, 2007.
- [15] J. Diller and C. Favre. Dynamics of bimeromorphic maps of surfaces. *Amer. J. Math.*, 123(6):1135–1169, 2001.
- [16] I. V. Dolgachev and V. A. Iskovskikh. Finite subgroups of the plane Cremona group. In *Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. I*, volume 269 of *Progr. Math.*, pages 443–548. Birkhäuser Boston, Inc., Boston, MA, 2009.
- [17] D. B. A. Epstein and W. P. Thurston. Transformation groups and natural bundles. *Proc. London Math. Soc. (3)*, 38(2):219–236, 1979.
- [18] C. Favre. Le groupe de Cremona et ses sous-groupes de type fini. *Astérisque*, (332):Exp. No. 998, vii, 11–43, 2010. Séminaire Bourbaki. Volume 2008/2009. Exposés 997–1011.
- [19] É. Ghys and P. de la Harpe, editors. *Sur les groupes hyperboliques d’après Mikhael Gromov*, volume 83 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, 1990. Papers from the Swiss Seminar on Hyperbolic Groups held in Bern, 1988.
- [20] M. H. Gizatullin. Rational G -surfaces. *Izv. Akad. Nauk SSSR Ser. Mat.*, 44(1):110–144, 239, 1980.
- [21] A. Huckleberry and D. Zaitsev. Actions of groups of birationally extendible automorphisms. In *Geometric complex analysis (Hayama, 1995)*, pages 261–285. World Sci. Publ., River Edge, NJ, 1996.
- [22] V. A. Iskovskikh and I. R. Shafarevich. Algebraic surfaces. In *Algebraic geometry, II*, volume 35 of *Encyclopaedia Math. Sci.*, pages 127–262. Springer, Berlin, 1996.
- [23] M. I. Kargapolov and Ju. I. Merzljakov. *Fundamentals of the theory of groups*, volume 62 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Berlin, 1979. Translated from the second Russian edition by Robert G. Burns.
- [24] M. Martelo and J. Ribón. Derived length of solvable groups of local diffeomorphisms. *Math. Ann.*, 358(3-4):701–728, 2014.
- [25] E. Souche. *Quasi-isométrie et quasi-plans dans l’étude des groupes discrets*. PhD thesis, Université de Provence, 2001.
- [26] J. Tits. Free subgroups in linear groups. *J. Algebra*, 20:250–270, 1972.
- [27] A. Weil. On algebraic groups of transformations. *Amer. J. Math.*, 77:355–391, 1955.

- [28] D. Wright. Abelian subgroups of $\text{Aut}_k(k[X, Y])$ and applications to actions on the affine plane. *Illinois J. Math.*, 23(4):579–634, 1979.
- [29] D. Zaitsev. Regularization of birational group operations in the sense of Weil. *J. Lie Theory*, 5(2):207–224, 1995.

INSTITUT DE MATHÉMATIQUES DE JUSSIEU-PARIS RIVE GAUCHE, UMR 7586, UNIVERSITÉ
PARIS 7, BÂTIMENT SOPHIE GERMAIN, CASE 7012, 75205 PARIS CEDEX 13, FRANCE.
E-mail address: `deserti@math.univ-paris-diderot.fr`