

A counterexample to the Hopf-Oleinik lemma (elliptic case) *

D. E. Apushkinskaya, A. I. Nazarov

December 25, 2015

Abstract

We construct a new counterexample to the Hopf-Oleinik boundary point lemma. It shows that for convex domains $C^{1, \text{Dini}}$ assumption on $\partial\Omega$ is the necessary and sufficient condition providing the Hopf-Oleinik type estimates.

Dedicated to Professor M.V. Safonov

1 Introduction

The influence of the properties of a domain to the behavior of a solution is one of the most important topic in the qualitative analysis of partial differential equations.

The significant result in this field is the Hopf-Oleinik lemma, known also as the "Boundary Point Principle". This celebrated lemma states:

Let u be a nonconstant solution to a second-order homogeneous uniformly elliptic nondivergence equation with bounded measurable coefficients, and let u attain its extremum at a point x^0 located on the boundary of a domain $\Omega \subset \mathbb{R}^n$. Then $\frac{\partial u}{\partial \mathbf{n}}(x^0)$ is necessarily nonzero provided that $\partial\Omega$ satisfies the proper assumptions at x^0 .

This result was established in a pioneering paper of S. Zaremba [Zar10] for the Laplace equation in a 3-dimensional domain Ω having interior touching ball at x^0 and generalized by G. Giraud [Gir32]-[Gir33] to equations with

*AMS Subject Classification: 35J15, 35B45

Key words: elliptic equations, Hopf-Oleinik lemma, Dini continuity, counterexample

Hölder continuous leading coefficients and continuous lower order coefficients in domains Ω belonging to the class $C^{1,\alpha}$ with $\alpha \in (0, 1)$.

Notice that a related assertion about the negativity on $\partial\Omega$ of the normal derivative of the Green's function corresponding to the Dirichlet problem for the Laplace operator was proved much earlier for 2-dimensional smooth domains by C. Neumann in [Neu88] (see also [Kor01]). The result of [Neu88] was extended for operators with the lower order coefficients by L. Lichtenstein [Lic24]. The same version of the Boundary Point Principle for the Laplacian and 3-dimensional domains satisfying a more flexible interior paraboloid condition was obtained by M.V. Keldysch and M.A. Lavrentiev in [KL37].

A crucial step in studying the Boundary Point Principle was made by E. Hopf [Hop52] and O.A. Oleinik [Ole52], who simultaneously and independently proved the statement for the general elliptic equations with bounded coefficients and domains satisfying an interior ball condition at x^0 .

Later the efforts of many mathematicians were focused on generalization of the Boundary Point Principle in several directions (for the details we refer the reader to [ABM⁺11] and [Alv11] and references therein). Among these directions are the extension of the class of operators and the class of solutions, as well as the weakening of assumptions on the boundary.

The widening of the class of operators to singular/degenerate ones was made in the papers [KH75], [KH77] and [ABM⁺11], while the uniform elliptic operators with unbounded lower order coefficients were studied in [Saf10] and [Naz12] (see also [NU09]). We mention also the publications [Tol83] and [MS15] where the Boundary Point Principle was established for a class of degenerate quasilinear operators including the p -Laplacian.

We note that before 2010 all the results were formulated for classical solutions, i.e. $u \in C^2(\Omega)$. The class of solutions was expanded in [Saf10] to strong generalized solutions with Sobolev's second order derivatives. The latter requirement seems to be natural in studying of nondivergence elliptic equations.

The reduction of the assumptions on the boundary of Ω up to $C^{1,\text{Dini}}$ -regularity was realized for various elliptic operators in the papers [Wid67], [Him70] and [Lie85] (see also [Saf08]). A weakened form of the Hopf-Oleinik lemma (the existence of a boundary point x^1 in any neighborhood of x^0 and a direction ℓ such that $\frac{\partial u}{\partial \ell}(x^1) \neq 0$) was proved in [Nad83] for a much wider class of domains including all Lipschitz ones. We mention also the paper [Swe97] where the behavior of superharmonic functions near the boundary of 2-dimensional domains with corners is described in terms of the main eigenfunction of the Dirichlet Laplacian.

The sharpness of some requirements was confirmed by corresponding counterexamples constructed in [Wid67], [Him70], [KH75], [Saf08], [ABM⁺11]

and [Naz12]. In particular, the counterexamples from [Wid67], [Him70] and [Saf08] show that the Hopf-Oleinik result fails for domains lying entirely in non-Dini paraboloids.

The main result of our paper is a new counterexample showing the sharpness of the Dini condition for the boundary of Ω . The simplest version of this counterexample can be formulated as follows:

Let Ω be a convex domain in \mathbb{R}^n , let $\partial\Omega$ in a neighborhood of the origin be described by the equation $x_n = F(x')$ with $F \geq 0$ and $F(0) = 0$, and let $u \in W_{n,loc}^2(\Omega) \cap C(\bar{\Omega})$ be a solution of the uniformly elliptic equation

$$-a^{ij}(x)D_iD_ju = 0 \quad \text{in } \Omega.$$

Suppose also that $u|_{\partial\Omega}$ vanishes at a neighborhood of the origin. If, in addition, the function $\delta(r) = \sup_{|x'| \leq r} \frac{F(x')}{|x'|}$ is not Dini continuous at zero, then $\frac{\partial u}{\partial \mathbf{n}}(0) = 0$.

Thus, it turns out that for convex domains the Dini continuity assumption on $\delta(r)$ is necessary and sufficient for the validity of the Boundary Point Principle. We emphasize that in our counterexample the Dini condition fails for supremum of $F(x')/|x'|$, while in all the previous results of this kind it fails for infimum of $F(x')/|x'|$. In other words, we show that the violating of the Dini condition just in one direction causes the lack of the Hopf-Oleinik lemma.

1.1 Notation and Conventions

Throughout the paper we use the following notation:

$x = (x', x_n) = (x_1, \dots, x_{n-1}, x_n)$ is a point in \mathbb{R}^n ;

$\mathbb{R}_+^n = \{x \in \mathbb{R}^n : x_n > 0\}$;

$|x|, |x'|$ are the Euclidean norms in the corresponding spaces;

$\chi_{\mathcal{E}}$ denotes the characteristic function of the set $\mathcal{E} \subset \mathbb{R}^n$;

Ω is a bounded domain in \mathbb{R}^n with boundary $\partial\Omega$;

$\mathcal{P}_{r,h}(\bar{x}') = \{x \in \mathbb{R}^n : |x' - \bar{x}'| < r, 0 < x_n < h\}$; $\mathcal{P}_r(\bar{x}') = \mathcal{P}_{r,r}(\bar{x}')$;

$\mathcal{P}_{r,h} = \mathcal{P}_{r,h}(0)$; $\mathcal{P}_r = \mathcal{P}_r(0)$;

$B_r(x^0)$ is the open ball in \mathbb{R}^n with center x^0 and radius r ; $B_r = B_r(0)$;

For $r_1 < r_2$ we define the annulus $\mathcal{B}(x^0, r_1, r_2) = B_{r_2}(x^0) \setminus \overline{B_{r_1}(x^0)}$.

$v_+ = \max\{v, 0\}$, $v_- = \max\{-v, 0\}$.

$\|\cdot\|_{\infty, \Omega}$ denotes the norm in $L_{\infty}(\Omega)$.

We adopt the convention that the indices i and j run from 1 to n . We also adopt the convention regarding summation with respect to repeated indices.

D_i denotes the operator of (weak) differentiation with respect to x_i ;
 $D = (D', D_n) = (D_1, \dots, D_{n-1}, D_n)$.

\mathcal{L} is a linear uniformly elliptic operator with measurable coefficients:

$$\mathcal{L}u \equiv -a^{ij}(x)D_iD_ju + b^i(x)D_iu, \quad \nu\mathcal{I}_n \leq (a^{ij}(x)) \leq \nu^{-1}\mathcal{I}_n, \quad (1)$$

where \mathcal{I}_n is identity $(n \times n)$ -matrix. We denote $\mathbf{b}(x) = (b^1(x), \dots, b^n(x))$.

We use letters C and N (with or without indices) to denote various constants. To indicate that, say, C depends on some parameters, we list them in the parentheses: $C(\dots)$.

Definition 1. We say that a function $\sigma : [0, 1] \rightarrow \mathbb{R}_+$ belongs to the class \mathcal{D}_1 if

- σ is increasing, and $\sigma(0) = 0$, and $\sigma(1) = 1$;
- $\sigma(t)/t$ is summable and decreasing.

Remark 1. Our assumption about decay of $\sigma(t)/t$ is not restrictive. Indeed, for any increasing function $\sigma : [0, 1] \rightarrow \mathbb{R}_+$ satisfying $\sigma(0) = 0$ and $\sigma(1) = 1$ and having summable $\sigma(t)/t$, we can define

$$\tilde{\sigma}(t) = t \sup_{\tau \in [t, 1]} \frac{\sigma(\tau)}{\tau}, \quad t \in (0, 1).$$

It is easy to see that $\tilde{\sigma} \in \mathcal{D}_1$, $\tilde{\sigma}(t)/t$ decreases and $\sigma(t) \leq \tilde{\sigma}(t)$, $\forall t \in (0, 1]$.

Definition 2. Let a function σ belong to the class \mathcal{D}_1 . We define the function \mathcal{J}_σ as follows

$$\mathcal{J}_\sigma(s) := \int_0^s \frac{\sigma(\tau)}{\tau} d\tau. \quad (2)$$

Remark 2. Decreasing of $\sigma(t)/t$ implies

$$\sigma(t) \leq \mathcal{J}_\sigma(t) \quad \forall t \in [0, 1]. \quad (3)$$

In addition, for $t \leq t_0 \leq 1$ we have

$$\sigma(t/t_0) = \frac{\sigma(t/t_0)}{t/t_0} \cdot t/t_0 \leq \frac{\sigma(t)}{t} \cdot t/t_0 = \frac{\sigma(t)}{t_0}, \quad (4)$$

and, similarly,

$$\mathcal{J}_\sigma(t/t_0) \leq \frac{\mathcal{J}_\sigma(t)}{t_0}. \quad (5)$$

Definition 3. We say that a function ζ satisfies the Dini condition at zero if

$$|\zeta(r)| \leq C\sigma(r),$$

and σ belongs to the class \mathcal{D}_1 .

2 Preliminaries

2.1 Properties of Ω

Let Ω be a bounded domain in \mathbb{R}^n . Without loss of generality we may assume $0 \in \partial\Omega$.

Suppose that Ω is locally convex in a neighborhood of the origin. Without restriction the latter means that for some $0 < \mathcal{R}_0 \leq 1$ we have

$$\mathcal{P}_{\mathcal{R}_0} \cap \Omega = \{(x', x_n) \in \mathbb{R}^n : |x'| \leq \mathcal{R}_0, F(x') < x_n < \mathcal{R}_0\},$$

where F is a convex nonnegative function satisfying $F(0) = 0$.

For $r \in (0, \mathcal{R}_0)$ we define the functions $\delta = \delta(r)$ and $\delta_1 = \delta_1(r)$ by the formulas

$$\delta(r) := \max_{|x'| \leq r} \frac{F(x')}{|x'|}, \quad \delta_1(r) := \max_{|x'| \leq r} |\nabla F(x')|. \quad (6)$$

Lemma 2.1. *The following statements hold:*

- (a) $\delta_1(r) \rightarrow 0$ as $r \rightarrow 0$ iff $\delta(r) \rightarrow 0$ as $r \rightarrow 0$.
- (b) $\delta_1(r)$ satisfies the Dini condition at zero iff $\delta(r)$ satisfies the Dini condition at zero.

Proof. By convexity of F , we have for any x' and z' the estimate

$$F(z') \geq F(x') + \nabla F(x') \cdot (z' - x'). \quad (7)$$

Therefore,

$$|\nabla F(x')| \geq \nabla F(x') \cdot \frac{x'}{|x'|} \geq \frac{F(x')}{|x'|},$$

and, consequently,

$$\delta_1(r) \geq \delta(r). \quad (8)$$

On the other hand, for any $r < \mathcal{R}_0/2$ we can find a point x'_* such that

$$|\nabla F(x'_*)| = \delta_1(r).$$

Choosing $z' = x'_* + r \frac{\nabla F(x'_*)}{|\nabla F(x'_*)|}$, we easily deduce from (7) the inequalities

$$|z'| \leq 2r \quad \text{and} \quad F(z') \geq r\delta_1(r),$$

which provide

$$\delta(2r) \geq \delta(|z'|) \geq \frac{\delta_1(r)}{2}. \quad (9)$$

Combining (8) and (9) we conclude that statement (a) is obvious and the integrals

$$\int_0^{\mathcal{R}_0} \frac{\delta(r)}{r} dr \quad \text{and} \quad \int_0^{\mathcal{R}_0} \frac{\delta_1(r)}{r} dr$$

converge simultaneously. \square

If $\delta(r)$ does not converge to zero as $r \rightarrow 0$, we can easily see that the domain Ω is contained in a dihedral wedge with the angle less than π and the edge going through the origin. For this case the statement of **Main Theorem** is proved already in [AN00, Theorem 4.3]. By this reason we will assume throughout this paper that

$$\delta(r) \rightarrow 0 \quad \text{as} \quad r \rightarrow 0. \quad (10)$$

In view of (10), it is evident that δ and δ_1 are moduli of continuity at the origin of the functions $F(x')/|x'|$ and $|\nabla F(x')|$, respectively.

2.2 Properties of $\mathcal{X}(\Omega)$

Let $\mathcal{X}(\Omega)$ be a function space with the norm $\|\cdot\|_{\mathcal{X},\Omega}$. For $\Omega_1 \subset \Omega$ we will assume that

$$\|f\|_{\mathcal{X},\Omega_1} = \|f \cdot \chi_{\Omega_1}\|_{\mathcal{X},\Omega}.$$

We suppose that $\mathcal{X}(\Omega)$ has the following properties:

- (i) For arbitrary measurable function g defined in Ω and any function $f \in \mathcal{X}(\Omega)$ the inequality $|g(x)| \leq |f(x)|$ implies $g \in \mathcal{X}(\Omega)$ and $\|g\|_{\mathcal{X},\Omega} \leq \|f\|_{\mathcal{X},\Omega}$;
- (ii) For $f_k \in \mathcal{X}(\Omega)$ the convergence $f_k \searrow 0$ a.e. in Ω implies $\|f_k\|_{\mathcal{X},\Omega} \rightarrow 0$.

Using the terminology of classic monograph of Kantorovich and Akilov [KA82] we may say that $\mathcal{X}(\Omega)$ is the ideal functional space with order continuous monotone norm (see [KA82, §3, Chapter IV, Part I] for more details).

We will also assume that

(iii) $\mathcal{X}_{loc}(\Omega)$ contains the Orlicz space $L_{\Phi,loc}(\Omega)$ with $\Phi(\varsigma) = e^\varsigma - \varsigma - 1$.

Finally, the basic assumption about $\mathcal{X}(\Omega)$ is the Aleksandrov-type maximum principle. Namely, we denote by $\mathcal{W}_{\mathcal{X},loc}^2(\Omega)$ the set of the functions u satisfying $D(Du) \in \mathcal{X}_{loc}(\Omega)$, and suppose that if $u \in \mathcal{W}_{\mathcal{X},loc}^2(\Omega) \cap \mathcal{C}(\overline{\Omega})$, $u|_{\partial\Omega} \leq 0$, and $|\mathbf{b}| \in \mathcal{X}(\Omega)$ then

$$u \leq N_0(n, \nu, \|\mathbf{b}\|_{\mathcal{X},\Omega}) \cdot \text{diam}(\Omega) \cdot \|(\mathcal{L}u)_+\|_{\mathcal{X},\{u>0\}}. \quad (11)$$

Remark 3. It is well known from [Ale60], [Bak61] and [Ale63] (see also survey [Naz05] for further references) that $L_n(\Omega)$ has property (11). It is also evident that properties (i)-(iii) are satisfied in $L_n(\Omega)$. Therefore, $L_n(\Omega)$ can be treated as a "basic" example of $\mathcal{X}(\Omega)$. As other examples of the space $\mathcal{X}(\Omega)$ we mention some Lebesgue weighted spaces with power weights (see [Naz01]).

Remark 4. Unlike the natural properties (i)-(ii), assumption (iii) is rather "technical" one. Without (iii), our arguments from the proof of Step 3 in Theorem 4.1 are not applicable to the approximating operator \mathcal{L}_ε . So, we can not withdraw (iii) in abstract setting. However, in all known examples of $\mathcal{X}(\Omega)$ the property (iii) is satisfied.

Remark 5. Some of the statements, that will be referred to in the sequel, were proved earlier just for the case $\mathcal{X}(\Omega) = L_n(\Omega)$. However, if all the arguments are based only on the Aleksandrov-type maximum principle, these statements remain valid for an arbitrary considered space $\mathcal{X}(\Omega)$. In such cases, we will refer without any further explanation.

We also need the following convergence lemmas.

Lemma 2.2. *Let $\{f_j\}$ be a sequence of measurable functions on Ω , and let $f \in \mathcal{X}(\Omega)$. Suppose also that $f_j \rightarrow 0$ in measure on Ω , and $|f_j(x)| \leq |f(x)|$. Then*

$$\|f_j\|_{\mathcal{X},\Omega} \rightarrow 0 \quad \text{as } j \rightarrow \infty. \quad (12)$$

Proof. We argue by a contradiction. Suppose (12) fails. Then there exists a subsequence $\{f_{j_k}\}$ satisfying

$$\|f_{j_k}\|_{\mathcal{X},\Omega} \geq \varepsilon > 0, \quad \forall k \in \mathbb{N}. \quad (13)$$

Due to the Riesz theorem, there exists also a sub-subsequence $\{f_{j_{k_l}}\}$ such that

$$f_{j_{k_l}} \rightarrow 0 \quad \text{a.e. in } \Omega.$$

For simplicity of notation we renumber the latter subsequence $\{f_{j_{k_l}}\}$ and denote its elements again by f_j .

Setting $\tilde{f}_k := \sup_{j \geq k} |f_j|$ we can easily see that $\tilde{f}_k \searrow 0$ a.e. in Ω . Now, taking into account properties (i) and (ii) of the space $\mathcal{X}(\Omega)$ we immediately get a contradiction with inequalities (13). The proof is complete. \square

Lemma 2.3. *Let $f \in \mathcal{X}(\Omega)$, and let $\mu(\rho) := \sup_{x \in \Omega} \|f\|_{\mathcal{X}, B_\rho(x) \cap \Omega}$.*

Then

$$\mu(\rho) \rightarrow 0 \quad \text{as } \rho \rightarrow 0.$$

Proof. For every $\rho > 0$ there exists a point $x^* = x^*(\rho) \in \Omega$ such that

$$\|f\|_{\mathcal{X}, B_\rho(x^*) \cap \Omega} \geq \frac{1}{2} \mu(\rho).$$

Next, for the sequence $f_\rho := f \cdot \chi_{B_\rho(x^*)}$ it is evident that $|f_\rho| \rightarrow 0$ in measure on Ω . Application of Lemma 2.2 finishes the proof. \square

Remark 6. We call $\mu(\rho) := \sup_{x \in \Omega} \|f\|_{\mathcal{X}, B_\rho(x) \cap \Omega}$ the modulus of continuity of function f in $\mathcal{X}(\Omega)$.

Lemma 2.4. *Let $D(Du) \in \mathcal{X}(\Omega)$, let \mathcal{L} be defined by (1), and let $\mathcal{L}u \in \mathcal{X}(\Omega)$. There exist the family of operators*

$$\mathcal{L}_\varepsilon = -a_\varepsilon^{ij}(x) D_i D_j + b_\varepsilon^i(x) D_i$$

with smooth coefficients a_ε^{ij} and bounded coefficients b_ε^i satisfying

$$\nu \mathcal{I}_n \leq (a_\varepsilon^{ij}(x)) \leq \nu^{-1} \mathcal{I}_n, \quad x \in \Omega, \quad (14)$$

$$|b_\varepsilon^i(x)| \leq |b^i(x)|, \quad x \in \Omega, \quad (15)$$

$$\|(\mathcal{L} - \mathcal{L}_\varepsilon)u\|_{\mathcal{X}, \Omega} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0, \quad (16)$$

respectively.

Proof. We start with extension of a^{ij} on the whole \mathbb{R}^n by the identity matrix and denote by a_ε^{ij} the standard mollification of extended functions a^{ij} . By construction, the coefficients a_ε^{ij} are smooth functions converging as $\varepsilon \rightarrow 0$ to a^{ij} a.e. in Ω . Moreover, it is clear that inequalities (14) are true.

Further, we set

$$\tilde{b}_\varepsilon^i(x) := \min \{|b^i(x)|, \varepsilon^{-1}\} \cdot \text{sign } b^i(x). \quad (17)$$

In view of (17), it is evident that $\tilde{b}_\varepsilon^i D_i u$ converges as $\varepsilon \rightarrow 0$ to $b^i D_i u$ almost everywhere in Ω . We claim that it is possible to change \tilde{b}_ε^i such that the "corrected coefficients" b_ε^i satisfy

$$|b_\varepsilon^i D_i u| \leq |b^i D_i u| \quad \text{in } \Omega. \quad (18)$$

Indeed, if $|\tilde{b}_\varepsilon^i D_i u| \leq |b^i D_i u|$ in Ω then (18) holds with $b_\varepsilon^i \equiv \tilde{b}_\varepsilon^i$. Otherwise, consider a point $x^0 \in \Omega$ where $|\tilde{b}_\varepsilon^i(x^0) D_i u(x^0)| > |b^i(x^0) D_i u(x^0)|$.

- a) Let $\tilde{b}_\varepsilon^i(x^0) D_i u(x^0) > b^i(x^0) D_i u(x^0) \geq 0$. In this case we decrease all the coefficients $\tilde{b}_\varepsilon^i(x^0)$ corresponding to the positive summands such that the both sums $b_\varepsilon^i D_i u$ and $b^i D_i u$ becomes equal.
- b) Let $\tilde{b}_\varepsilon^i(x^0) D_i u(x^0) < b^i(x^0) D_i u(x^0) \leq 0$. In this case we decrease all the coefficients $\tilde{b}_\varepsilon^i(x^0)$ corresponding to the negative summands such that the both sums $b_\varepsilon^i D_i u$ and $b^i D_i u$ becomes equal.
- c) Finally, let $\tilde{b}_\varepsilon^i(x^0) D_i u(x^0)$ and $b^i(x^0) D_i u(x^0)$ have different signs. In this case we apply to $-\tilde{b}_\varepsilon^i(x^0)$ the arguments from case a) or from case b), respectively.

Due to construction, the "corrected sum" $b_\varepsilon^i D_i u$ also converges as $\varepsilon \rightarrow 0$ to $b^i D_i u$ a.e. in Ω , and pointwise inequalities (15) hold true.

Finally, taking into account (18) and applying Lemma 2.2 we get (16). \square

3 Gradient estimates near the boundary

Lemma 3.1. *Let $\mathcal{N} \subset \mathbb{R}_+^n$ be an open set, let $\gamma = \frac{\nu}{\sqrt{n-1}}$, let $\rho > 0$, and let*

$$\Pi_\rho = \{y \in \mathbb{R}^n : |y_i| < \rho \text{ for } i = 1, \dots, n-1; \quad 0 < y_n < \gamma\rho\}.$$

We assume that $|\mathbf{b}| \in \mathcal{X}(\mathcal{N})$ and a function v satisfies the conditions

$$v \in \mathcal{W}_{\mathcal{X},loc}^2(\mathcal{N}), \quad v \geq 0 \quad \text{in } \Pi_\rho, \quad v \geq k = \text{const} > 0 \quad \text{on } \partial\mathcal{N} \cap \bar{\Pi}_\rho.$$

Then

$$v \geq C_1 k - C_2 k \|\mathbf{b}\|_{\mathcal{X}, \mathcal{N} \cap \Pi_\rho} - C_3 \rho \|(\mathcal{L}v)_-\|_{\mathcal{X}, \mathcal{N} \cap \Pi_\rho} \quad \text{in } \mathcal{N} \cap B_{\frac{\gamma\rho}{4}}(z),$$

where $z = (0, \dots, 0, \frac{1}{2}\gamma\rho)$, while $C_1 = \frac{1}{16}(1 - \gamma^2)$, $C_2 = C_2(n, \nu, \|\mathbf{b}\|_{\mathcal{X}, \mathcal{N}})$, and $C_3 = C_3(n, \nu, \|\mathbf{b}\|_{\mathcal{X}, \mathcal{N}})$.

Proof. The proof is similar in spirit to [AU95, Lemma 1].

Consider the barrier function

$$\psi(y) = k \left[\left(1 - \frac{y_n}{\gamma\rho}\right)^2 - \frac{|y'|^2}{\rho^2} \right].$$

An elementary computation gives

$$\mathcal{L}\psi \leq k \left(\frac{2(n-1)}{\rho^2} \nu^{-1} - \frac{2}{\gamma^2 \rho^2} \nu \right) + |\mathbf{b}| |D\psi| \leq N_1(n, \nu) |\mathbf{b}| \frac{k}{\rho} \quad \text{in } \Pi_\rho.$$

Moreover, setting

$$\begin{aligned} \mathcal{S}_1 &= \{y \in \partial(\mathcal{N} \cap \Pi_\rho) : |y_i| = \rho \text{ for some } i = 1, \dots, n-1\}, \\ \mathcal{S}_2 &= \{y \in \partial(\mathcal{N} \cap \Pi_\rho) : y_n = \gamma\rho\} \end{aligned}$$

we have

$$\begin{aligned} \psi|_{\mathcal{S}_1 \cup \mathcal{S}_2} &\leq 0 \leq v, \\ \psi|_{\partial\mathcal{N} \cap \bar{\Pi}_\rho} &\leq k \leq v|_{\partial\mathcal{N} \cap \bar{\Pi}_\rho}. \end{aligned}$$

Applying inequality (11) in $\mathcal{N} \cap \Pi_\rho$ to the difference $\psi - v$ we obtain

$$\psi - v \leq N_0 \cdot \text{diam}(\Pi_\rho) \cdot \|(\mathcal{L}\psi - \mathcal{L}v)_+\|_{\mathcal{X}, \mathcal{N} \cap \Pi_\rho} \quad \text{in } \mathcal{N} \cap \Pi_\rho,$$

and, consequently,

$$\begin{aligned} v &\geq k \left[\left(1 - \frac{\frac{3}{4}\gamma\rho}{\gamma\rho}\right)^2 - \frac{\gamma^2 \rho^2}{16\rho^2} \right] - C_2 k \|\mathbf{b}\|_{\mathcal{X}, \mathcal{N} \cap \Pi_\rho} - C_3 \rho \|(\mathcal{L}v)_-\|_{\mathcal{X}, \mathcal{N} \cap \Pi_\rho} \\ &= \frac{(1 - \gamma^2)}{16} k - C_2 k \|\mathbf{b}\|_{\mathcal{X}, \mathcal{N} \cap \Pi_\rho} - C_3 \rho \|(\mathcal{L}v)_-\|_{\mathcal{X}, \mathcal{N} \cap \Pi_\rho} \quad \text{in } \mathcal{N} \cap B_{\frac{\gamma\rho}{4}}(z). \end{aligned}$$

□

Our next statement is a version of Theorem 2.3 [Naz12].

Lemma 3.2. *Let $v \in \mathcal{W}_{\mathcal{X},loc}^2(\Omega) \cap \mathcal{C}(\overline{\Omega})$, let $v|_{\partial\Omega} = 0$, and let $|\mathbf{b}| \in \mathcal{X}(\Omega)$. Suppose also that for all $\rho \leq \rho_* \leq 1$ the inequalities*

$$\|b^n\|_{\mathcal{X},\mathcal{P}_\rho\cap\Omega} \leq \mathfrak{B}\sigma(\rho/\rho_*), \quad \|(\mathcal{L}v)_+\|_{\mathcal{X},\mathcal{P}_\rho\cap\Omega} \leq \mathfrak{F}\sigma(\rho/\rho_*)$$

hold true. Here \mathfrak{B} and \mathfrak{F} are some positive constants, while a function σ belongs to \mathcal{D}_1 .

Then

$$\sup_{0 < x_n < \rho} \frac{v(0, x_n)}{x_n} \leq C_4 \left(\rho^{-1} \sup_{\mathcal{P}_\rho \cap \Omega} v + \mathfrak{F}\mathcal{J}_\sigma(\rho/\rho_*) \right), \quad \forall \rho \leq \rho_*. \quad (19)$$

Here the constant C_4 depends on $n, \nu, \mathfrak{B}, \sigma$, and on the moduli of continuity of $|\mathbf{b}'|$ in $\mathcal{X}(\mathcal{P}_{\rho_} \cap \Omega)$, whereas \mathcal{J}_σ is a function defined by formula (2).*

Remark 7. We recall that $0 \in \partial\Omega$.

Proof. First, we assume that $\rho \leq \bar{\rho}$, where $\bar{\rho} \leq \rho_*$ will be fixed later. Following [Naz12] we introduce the sequence of cylinders $\mathcal{P}_{\rho_k, h_k}$, $k \geq 0$, where $\rho_k = 2^{-k}\rho$, $h_k = \zeta_k \rho_k$, while the sequence $\zeta_k \downarrow 0$ will be chosen later.

We set $w_k = v - M_k x_n$, where the quantities M_k , $k \geq 1$ are defined as

$$M_k = \sup_{\mathcal{P}_{\rho_k, h_{k-1}} \cap \Omega} \frac{v(x)}{\max\{x_n, h_k\}} \geq \sup_{\{\mathcal{P}_{\rho_k, h_{k-1}} \setminus \mathcal{P}_{\rho_k, h_k}\} \cap \Omega} \frac{v(x)}{x_n}.$$

It is easy to see that $w_k \leq 0$ on $\partial\Omega \cap \overline{\mathcal{P}_{\rho_k, h_k}}$, while the definition of M_k gives $w_k \leq 0$ on the top of the cylinder $\mathcal{P}_{\rho_k, h_k}$.

Let $x^0 \in \mathcal{P}_{\rho_k - h_k, h_k} \cap \Omega$. Taking into account Remark 5 we apply the so-called "boundary growth lemma" (see, for instance, [LU85, Lemma 2.5] or [Saf10, Lemma 2.6] or [Naz12, Lemma 2.2]) to the (positive) function $M_k h_k - w_k$ in $\mathcal{P}_{h_k}(x^0) \cap \Omega$. It gives for $x \in \mathcal{P}_{h_k/2, h_k}(x^0) \cap \Omega$

$$\begin{aligned} M_k h_k - w_k(x) &\geq M_k h_k [\vartheta - N_2 \| |\mathbf{b}| \|_{\mathcal{X}, \mathcal{P}_{\rho_k} \cap \Omega}] \\ &\quad - N_3 h_k \| (\mathcal{L}w_k)_+ \|_{\mathcal{X}, \mathcal{P}_{h_k}(x^0) \cap \Omega}, \end{aligned} \quad (20)$$

where $\vartheta = \vartheta(n, \nu, \sigma, \mathfrak{B}) \in (0, 1)$, the positive constant N_2 depends on the same parameters as ϑ whereas the positive constant N_3 is completely defined by the values of n, ν and \mathfrak{B} . We suppose that $\bar{\rho}$ is so small that the quantity in the square brackets is greater than $\vartheta/2$. Further, direct calculation shows that the assumptions of our lemma imply

$$\begin{aligned} \| (\mathcal{L}w_k)_+ \|_{\mathcal{X}, \mathcal{P}_{h_k}(x^0) \cap \Omega} &\leq \| (\mathcal{L}v)_+ \|_{\mathcal{X}, \mathcal{P}_{h_k}(x^0) \cap \Omega} + M_k \| b^n \|_{\mathcal{X}, \mathcal{P}_{h_k}(x^0) \cap \Omega} \\ &\leq (\mathfrak{F} + M_k \mathfrak{B}) \sigma(\rho_k/\rho_*). \end{aligned}$$

Substituting the last inequality into (20) and taking supremum w.r.t. x^0 we obtain

$$\sup_{\mathcal{P}_{\rho_k - h_k, h_k} \cap \Omega} w_k \leq M_k h_k [1 - \vartheta/2 + N_2 \mathfrak{B} \sigma(\rho_k/\rho_*)] + N_3 h_k \mathfrak{F} \sigma(\rho_k/\rho_*).$$

Repeating previous arguments provides for integer $m \leq \frac{\rho_k}{h_k}$ the inequality

$$\sup_{\mathcal{P}_{\rho_k - m h_k, h_k} \cap \Omega} w_k \leq M_k h_k \left[(1 - \vartheta/2)^m + N_2 \mathfrak{B} \frac{\sigma(\rho_k/\rho_*)}{\vartheta/2} \right] + N_3 h_k \mathfrak{F} \frac{\sigma(\rho_k/\rho_*)}{\vartheta/2}.$$

Setting $m = \lfloor \frac{\rho_{k+1}}{h_k} \rfloor$, we arrive at

$$\begin{aligned} \sup_{\mathcal{P}_{\rho_{k+1}, h_k} \cap \Omega} w_k &\leq \frac{M_k h_k}{1 - \vartheta/2} \left(\exp\left(-\lambda \frac{\rho_{k+1}}{h_k}\right) + N_2 \mathfrak{B} \frac{\sigma(\rho_k/\rho_*)}{\vartheta/2} \right) \\ &\quad + N_3 h_k \mathfrak{F} \frac{\sigma(\rho_k/\rho_*)}{(1 - \vartheta/2)\vartheta/2}, \end{aligned}$$

where $\lambda = -\ln(1 - \vartheta/2) > 0$.

Therefore, for $x \in \mathcal{P}_{\rho_{k+1}, h_k} \cap \Omega$

$$\frac{w_k(x)}{\max\{x_n, h_{k+1}\}} \leq M_k \gamma_k + N_3 \mathfrak{F} \frac{\sigma(\rho_k/\rho_*)}{(1 - \vartheta/2)\vartheta/2} \cdot \frac{2\zeta_k}{\zeta_{k+1}}, \quad (21)$$

where $\gamma_k = \frac{1}{1 - \vartheta/2} \frac{2\zeta_k}{\zeta_{k+1}} \cdot \left(\exp\left(-\frac{\lambda}{2\zeta_k}\right) + N_2 \mathfrak{B} \frac{\sigma(\rho_k/\rho_*)}{\vartheta/2} \right)$.

Estimate (21) implies

$$\begin{aligned} M_{k+1} &\leq M_k (1 + \gamma_k) + N_3 \mathfrak{F} \frac{\sigma(\rho_k/\rho_*)}{(1 - \vartheta/2)\vartheta/2} \cdot \frac{2\zeta_k}{\zeta_{k+1}} \\ &\leq M_1 \cdot \prod_{j=1}^k (1 + \gamma_j) + 2N_3 \mathfrak{F} \cdot \sum_{j=1}^k \sigma(\rho_j/\rho_*) \frac{\zeta_j}{\zeta_{j+1}} \cdot \prod_{l=j}^{k-1} (1 + \gamma_l). \end{aligned}$$

We set $\zeta_k = \frac{1}{k+k_0}$ and choose k_0 so large and $\bar{\rho}/\rho_*$ so small that $\gamma_1 \leq \frac{1}{2}$. Note that $k_0 = k_0(n, \nu, \sigma, \mathfrak{B})$ while $\bar{\rho}/\rho_*$ depends on the same parameters as k_0 and, in addition, on the moduli of continuity of $|\mathbf{b}'|$ in $\mathcal{X}(\mathcal{P}_{\rho_*} \cap \Omega)$.

Now we observe that the first term in γ_k forms a convergent series. The same is true for the second term, since

$$\sum_{k=1}^{\infty} \sigma(2^{-k} \rho/\rho_*) \asymp \int_0^{\infty} \sigma(2^{-s} \rho/\rho_*) ds \asymp \mathcal{J}_{\sigma}(\rho/\rho_*).$$

Therefore, the infinite product $\Pi = \prod_k (1 + \gamma_k)$ also converges, and we obtain for $k > 1$ the inequality

$$\begin{aligned} M_k &\leq \Pi \cdot \left(M_1 + 2N_3 \mathfrak{F} \cdot \sum_{j=1}^k \sigma(\rho_j/\rho_*) \frac{\zeta_j}{\zeta_{j+1}} \right) \\ &\leq \Pi \cdot (M_1 + N_4(n, \nu, \sigma, \mathfrak{B}) \mathfrak{F} \mathcal{J}_\sigma(\rho/\rho_*)). \end{aligned} \quad (22)$$

Thus, all M_k are bounded. It remains only to note that

$$M_1 \leq \frac{1}{h_1} \sup_{\mathcal{P}_{\rho/2} \cap \Omega} v. \quad (23)$$

Combining (22) and (23), we arrive at

$$\sup_{0 < x_n < \rho/2} \frac{v(0, x_n)}{x_n} \leq N_5(n, \nu, \sigma, \mathfrak{B}) \left(\rho^{-1} \sup_{\mathcal{P}_{\rho/2} \cap \Omega} v + \mathfrak{F} \mathcal{J}_\sigma(\rho/\rho_*) \right). \quad (24)$$

Further, it is easy to find a majorant for $\frac{v(0, x_n)}{x_n}$ for any $x_n \in [\rho/2, \rho]$ since

$$\sup_{\rho/2 \leq x_n < \rho} \frac{v(0, x_n)}{x_n} \leq 2\rho^{-1} \sup_{\rho/2 \leq x_n < \rho} v(0, x_n) \leq 2\rho^{-1} \sup_{\mathcal{P}_\rho \cap \Omega} v. \quad (25)$$

Combination of (24) and (25) implies (19) with $C_4 = \max\{N_5, 2\}$ for $\rho \leq \bar{\rho}$.

Now, we consider $\rho > \bar{\rho}$. If $x_n < \bar{\rho}$ then the estimate

$$\frac{v(0, x_n)}{x_n} \leq 2N_5 \left(\bar{\rho}^{-1} \sup_{\mathcal{P}_{\bar{\rho}} \cap \Omega} v + \mathfrak{F} \mathcal{J}_\sigma(\rho/\rho_*) \right) \quad (26)$$

follows from the above arguments. Otherwise, i.e. for $x_n \geq \bar{\rho}$, inequality (26) is especially true. Thus, for $\rho > \bar{\rho}$ we again arrive at (19) with $C_4 = \max\{N_5, 2\} \bar{\rho}^{-1}$. The proof is complete. \square

4 Main results

Recall that Ω satisfies the assumptions from Subsection 2.1. Throughout this section we shall suppose that \mathcal{L} is defined by (1), $|\mathbf{b}| \in \mathcal{X}(\Omega)$, and a function u satisfies the following assumptions:

$$u \in \mathcal{W}_{\mathcal{X},loc}^2(\Omega) \cap \mathcal{C}(\bar{\Omega}), \quad \mathcal{L}u = 0 \quad \text{in } \Omega, \quad u|_{\partial\Omega \cap \bar{\mathcal{P}}_{\mathcal{R}_0}} = 0. \quad (27)$$

Theorem 4.1. *Let the inequality*

$$\sup_{x \in \mathcal{P}_{\mathcal{R}_0/2}} \|b^n\|_{\mathcal{X}, \mathcal{P}_\rho(x') \cap \Omega} \leq \mathfrak{B} \sigma(\rho/\mathcal{R}_0)$$

hold true for all $\rho \leq \mathcal{R}_0/2$. Here \mathfrak{B} is a positive constant, and a function $\sigma \in \mathcal{D}_1$ satisfies

$$\mathcal{J}_\sigma(t) = o(\delta(t)) \quad \text{as } t \rightarrow 0. \quad (28)$$

Then, there exists a sufficiently small positive number R_0 completely defined by $n, \nu, \mathcal{R}_0, \mathfrak{B}$, by the functions σ, δ , and by the moduli of continuity of $|b'|$ in $\mathcal{X}(\Omega)$ such that for any $r \in (0, R_0/2)$ we have

$$\operatorname{osc}_{\Omega \cap \mathcal{P}_{r/4}} \frac{u(x)}{x_n} \leq (1 - \varkappa \delta(r)) \operatorname{osc}_{\Omega \cap \mathcal{P}_{2r}} \frac{u(x)}{x_n}. \quad (29)$$

Here the constant $\varkappa \in (0; 1)$ is completely determined by n, ν .

Proof. The proof will be divided into 3 steps.

1. Our arguments are adapted from [AU95, Lemma 2] and [Ura96, Lemma 3]. Let us denote

$$m^\pm = \sup_{\Omega \cap \mathcal{P}_{2r}} \pm \frac{u(x)}{x_n}, \quad \omega = m^+ + m^- = \operatorname{osc}_{\Omega \cap \mathcal{P}_{2r}} \frac{u(x)}{x_n}.$$

Since $u|_{\partial\Omega} = 0$ we have $m^\pm \geq 0$. Therefore, at least one of the numbers m^\pm is not less than $\frac{\omega}{2}$, and both of the numbers m^\pm are less than ω .

Let $m^+ \geq \frac{\omega}{2}$ for definiteness. Then we consider the nonnegative function $v(x) = m^+ x_n - u(x)$ in $\Omega \cap \mathcal{P}_{2r}$; (if $m^- > \frac{\omega}{2}$ then we consider the function $v(x) = m^- x_n + u(x)$).

Due to definition of δ , for any sufficiently small $r > 0$ we can find a point $x^* \in \partial\mathcal{P}_r \cap \partial\Omega$ such that $x_n^* = r\delta(r)$. Without loss of generality we may assume that $x_1^* = r$ and $x_\tau^* = 0$ for $\tau = 2, \dots, n-1$.

Next we assign to x^* a local orthogonal coordinate system y_1, \dots, y_n such that

- (a) y_1 -axis is directed along the projection of the vector $(x_1^*, \dots, x_{n-1}^*)$ onto tangential hyperplane to $\partial\Omega$ at x^* ;
- (b) y_2, \dots, y_{n-1} -axes are parallel to x_2, \dots, x_{n-1} -axes, respectively;
- (c) y_n -axis is directed inside Ω .

Due to the extremal property of x^* the axes y_1, \dots, y_{n-1} lie in the supporting hyperplane to $\partial\Omega$ at x^* . Moreover, if x^* is a smooth point of $\partial\Omega$ then y_n is directed along the inward normal to $\partial\Omega$.

Setting $\gamma = \frac{\nu}{\sqrt{n-1}}$ we consider in y -coordinates the cylinder

$$\Pi := \left\{ y \in \mathbb{R}^n : \left| y_1 - \frac{r}{2} \right| < \frac{r}{2}, |y_\tau| < \frac{r}{2}, 0 < y_n < \frac{1}{2}\gamma r \right\},$$

and the ball $B_{\rho_0}(z^0)$ with $\rho_0 = \frac{1}{8}\gamma r$ and $z^0 = (\frac{r}{2}, 0, \dots, 0, \frac{1}{4}\gamma r)$.

It should be emphasized that from now on, all considerations will be carried out in x -coordinates.




Figure 1: Schematic view of Π and $B_{\rho_0}(z^0)$.

We claim that

$$B_{\rho_0}(z^0) \subset \Omega. \quad (30)$$

Indeed, assume that (30) fails. Then there is a point $\hat{x} \in B_{\rho_0}(z^0)$ satisfying (in x -coordinates) the inequalities

$$F(\hat{x}') \geq \hat{x}_n \geq z_n^0 - \rho_0. \quad (31)$$

Since $\hat{x} \in B_{\rho_0}(z^0)$ it is clear that $|\hat{x}'| \leq 2r$ and

$$F(\hat{x}') \leq 2r\delta(2r).$$

On the other hand, denoting by φ the angle between x_n - and y_n -axis (see Fig. 1) we conclude that

$$z_n^0 - \rho_0 = r\delta(r) + \frac{r}{2} \sin \varphi + \frac{\gamma r}{4} \cos \varphi - \frac{\gamma r}{8} \geq \frac{\gamma r}{8} (2 \cos \varphi - 1).$$

Thus (31) is transformed into

$$\gamma (2 \cos \varphi - 1) \leq 16\delta(2r). \quad (32)$$

In view of (10) and Lemma 2.1, one can choose R_0 so small that $\delta_1(R_0) \leq 3/4$. It guarantees for all $r \leq R_0/2$ the inequalities

$$\cos \varphi = \frac{1}{\sqrt{1 + \tan^2 \varphi}} \geq \frac{1}{\sqrt{1 + \delta_1^2(r)}} \geq \frac{1}{\sqrt{1 + \delta_1^2(R_0)}} \geq \frac{4}{5}. \quad (33)$$

Now, combining (33) and (32) we get a contradiction with relation (10) provided $\delta(R_0)$ being small enough. The proof of (30) is complete.

2. With (30) at hands, we observe that

$$\inf\{x_n : x \in \Omega \cap \Pi\} \geq r\delta(r).$$

On the other hand, the condition $u = 0$ for $x \in \partial\Omega \cap \Pi$ gives the estimate

$$v = m^+ x_n \geq \frac{\omega}{2} x_n \quad \text{on} \quad \partial\Omega \cap \Pi.$$

Hence,

$$v \geq \frac{\omega}{2} r\delta(r) =: k_0 \quad \text{on} \quad \partial\Omega \cap \Pi. \quad (34)$$

So, we can apply Lemma 3.1 to the function v in cylinder Π . This gives the estimate

$$\inf_{B_{\rho_0}(z^0)} v \geq \left(k_0 [C_1 - C_2 \|\mathbf{b}\|_{\mathcal{X}, \Omega \cap \mathcal{P}_{2r}}] - C_3 \omega r \|b^n\|_{\mathcal{X}, \Omega \cap \mathcal{P}_{2r}} \right)_+,$$

where C_1 , C_2 and C_3 are the constants from Lemma 3.1. Decreasing R_0 , if necessary, we may assume that $\|\mathbf{b}\|_{\mathcal{X}, \Omega \cap \mathcal{P}_{R_0}} \leq C_1 / (2C_2)$. Thus, we arrive at

$$\inf_{B_{\rho_0}(z^0)} v \geq \left(k_0 \frac{C_1}{2} - C_3 \omega r \|b^n\|_{\mathcal{X}, \Omega \cap \mathcal{P}_{2r}} \right)_+ =: k_1. \quad (35)$$

Consider now an arbitrary point $\tilde{z} = (\tilde{z}', r/4 + \rho_0/8)$ such that $|\tilde{z}'| \leq \frac{r}{4}$. Observe also that $B_{\rho_0}(\tilde{z}) \subset \Omega$, otherwise we get a contradiction with definition of $\delta(r)$.

We claim that

$$\inf_{B_{\rho_0/8}(\tilde{z})} v \geq (k_0 \tilde{C}_1 - \tilde{C}_2 \omega r \|b^n\|_{\mathcal{X}, \Omega \cap \mathcal{P}_{2r}})_+, \quad (36)$$

where $\tilde{C}_1 = \tilde{C}_1(n, \nu)$, whereas \tilde{C}_2 is determined completely by n , ν , and $\|\mathbf{b}\|_{\mathcal{X}, \Omega}$. Indeed, due to convexity of Ω , for l running from 1 to a finite number $\mathfrak{N} = \mathfrak{N}(n, \nu)$ chosen so that

$$\frac{4}{3\rho_0} |z^0 - \tilde{z}| \leq \mathfrak{N} \leq \frac{2}{\rho_0} |z^0 - \tilde{z}|, \quad (37)$$

and for points $z^{[l]} := z^0 - \frac{l}{\mathfrak{N}}(z^0 - \tilde{z})$ we have $B_{\rho_0}(z^{[l]}) \subset \Omega$. It should be emphasized that the lower and the upper bounds in (37) do not depend on r .

In view of (35) we can compare in $\mathcal{B}(z^{[1]}, \rho_0/8, \rho_0)$ the function v with the standard barrier function

$$w(x) = k_1 \frac{|x - z^{[1]}|^{-s} - \rho_0^{-s}}{(\rho_0/8)^{-s} - \rho_0^{-s}}.$$

If $s = n\nu^{-2}$ then elementary calculation guarantees the estimates

$$\begin{aligned} \mathcal{L}w &\leq |\mathbf{b}||Dw| \leq c(n, \nu)k_1|\mathbf{b}|\rho_0^{-1} \quad \text{in } \mathcal{B}(z^{[1]}, \rho_0/8, \rho_0), \\ w(x) = k_1 &\leq v(x) \quad \text{on the sphere } |x - z^{[1]}| = \frac{\rho_0}{8} \\ w(x) = 0 &\leq v(x) \quad \text{on the sphere } |x - z^{[1]}| = \rho_0. \end{aligned}$$

Application of the maximum principle (11) in $\mathcal{B}(z^{[1]}, \rho_0/8, \rho_0)$ to the difference $w - v$ gives us the inequality

$$v(x) \geq (k_1 [w(x) - 2cN_0\|\mathbf{b}\|_{\mathcal{X}, \Omega \cap \mathcal{P}_{2r}}] - N_0 \frac{\gamma^r}{4} \omega \|b^n\|_{\mathcal{X}, \Omega \cap \mathcal{P}_{2r}})_+.$$

Since $B_{\rho_0/8}(z^{[2]}) \subset \mathcal{B}(z^{[1]}, \rho_0/8, 7\rho_0/8)$, the evident bound $w \geq \theta(n, \nu)$ holds true in $B_{\rho_0/8}(z^{[2]})$.

Decreasing R_0 , if necessary, we ensure that $\|\mathbf{b}\|_{\mathcal{X}, \Omega \cap \mathcal{P}_{R_0}} \leq (4cN_0)^{-1} \theta$. This implies

$$\inf_{B_{\rho_0/8}(z^{[2]})} v(x) \geq \left(\frac{k_1 \theta}{2} - N_0 \frac{\gamma^r}{4} \omega \|b^n\|_{\mathcal{X}, \Omega \cap \mathcal{P}_{2r}} \right)_+ =: k_2.$$

Repeating this procedure for $\mathcal{B}(z^{[l]}, \rho_0/8, \rho_0)$ and $l = 2, \dots, \mathfrak{N}$ we arrive at (36) with $\tilde{C}_1 = (\theta/2)^\mathfrak{N}$ and $\tilde{C}_2 = N_0 \frac{\gamma}{4} \cdot \frac{1 - (\theta/2)^\mathfrak{N}}{1 - (\theta/2)}$.

Furthermore, it is clear that

$$\left(k_0\tilde{C}_1 - \tilde{C}_2r\omega\|b^n\|_{\mathcal{X},\Omega\cap\mathcal{P}_{2r}}\right)_+ \geq \omega r \left(\frac{1}{2}\tilde{C}_1\delta(r) - \tilde{C}_2\mathfrak{B}\sigma(r/\mathcal{R}_0)\right)_+,$$

while inequalities (3) and (4) guarantee that

$$\sigma(r/\mathcal{R}_0) \leq \frac{\mathcal{J}_\sigma(r)}{\mathcal{R}_0}.$$

Decreasing again R_0 and taking into account the assumption (28) and the above inequalities, we can transform (36) into the form

$$\inf_{B_{\rho_0/8}(\tilde{z})} v \geq \frac{1}{4}\tilde{C}_1\omega r\delta(r) =: \tilde{k}. \quad (38)$$

3. Now, we take a small $\eta > 0$, define the set

$$\mathcal{A}_\eta := \mathcal{B}(\tilde{z}, \rho_0/8, \tilde{z}_n) \cap \Omega \cap \{x \in \mathcal{P}_{\mathcal{R}_0} : F(x') + \eta < x_n < \mathcal{R}_0\}$$

and introduce in \mathcal{A}_η the barrier function

$$W(x) = \mu\tilde{k} \frac{|x - \tilde{z}|^{-s} - (\tilde{z}_n)^{-s}}{(\rho_0/8)^{-s} - (\tilde{z}_n)^{-s}},$$

where $s = n\nu^{-2}$ and $0 < \mu \leq 1$.

Notice that $D(Du) \in \mathcal{X}(\mathcal{A}_\eta)$. Using Lemma 2.4 we construct the family of operators \mathcal{L}_ε satisfying $\|\mathcal{L}_\varepsilon u\|_{\mathcal{X},\mathcal{A}_\eta} \rightarrow 0$ as $\varepsilon \rightarrow 0$.

Arguing in the spirit of the proof of Lemma 4.2 [LU88], we define $v_1(x)$ and $v_2(x)$ as solutions of the following problems:

$$\begin{cases} \mathcal{L}_\varepsilon v_1 = b_\varepsilon^i D_i W & \text{in } \mathcal{A}_\eta \\ v_1 = v & \text{on } \partial\mathcal{A}_\eta \end{cases}, \quad \begin{cases} \mathcal{L}_\varepsilon v_2 = b_\varepsilon^i D_i W - b_\varepsilon^n m^+ & \text{in } \mathcal{A}_\eta \\ v_2 = 0 & \text{on } \partial\mathcal{A}_\eta \end{cases}.$$

It is well known (see, for instance, [Kry08, Chapter 6]) that $D(Dv_1)$ and $D(Dv_2)$ belong to the space $BMO_{loc}(\mathcal{A}_\eta)$. Moreover, the John-Nirenberg theorem [JN61] (see also [Duo01, §4, Chapter 6]) implies that $D(Dv_i)$, $i = 1, 2$, belong to the Orlicz space $L_{\Phi,loc}(\mathcal{A}_\eta)$ with $\Phi(\varsigma) = e^\varsigma - \varsigma - 1$. So, taking into account the property (iii) we may conclude that $v_i \in \mathcal{W}_{\mathcal{X},loc}^2(\mathcal{A}_\eta)$, $i = 1, 2$.

Furthermore, in view of (38) and the direct calculation, we have the inequalities

$$\begin{aligned}\mathcal{L}_\varepsilon W &\leq b_\varepsilon^i D_i W \quad \text{in } \mathcal{A}_\eta, \\ W(x) = \mu \tilde{k} &\leq v(x) = v_1(x) \quad \text{on the sphere } |x - \tilde{z}| = \frac{\rho_0}{8}, \\ W(x) = 0 &\leq v(x) = v_1(x) \quad \text{on } \partial\mathcal{A}_\eta \cap \{x \in \mathbb{R}^n : |x - \tilde{z}| = \tilde{z}_n\}.\end{aligned}$$

On the rest of $\partial\mathcal{A}_\eta$ we have $x_n = F(x') + \eta$ and, consequently, $\text{dist}\{x, \partial\Omega\} \leq \eta$. Since $u \in \mathcal{C}(\overline{\Omega})$, the latter inequality implies the estimate $u \leq H(\eta)$ there, and therefore,

$$v_1(x) = v(x) = m^+ x_n - u \geq \frac{\omega}{2} x_n - H(\eta),$$

where H is a nonnegative function tending to zero as $\eta \rightarrow 0$.

In addition, it is easy to verify that

$$W(x) \leq \mu N_6(n, \nu) \tilde{C}_1 \omega \delta(r) x_n \quad \text{in } \overline{\mathcal{B}}(\tilde{z}, \rho_0/8, \tilde{z}_n).$$

Choosing $\mu = \min \left\{ 1; \left(2N_6 \tilde{C}_1 \right)^{-1} \right\}$, we get

$$v_1(x) \geq W(x) - H(\eta) \quad \text{on } \partial\mathcal{A}_\eta.$$

The maximum principle (11) applied to the difference $W - H(\eta) - v_1$ in \mathcal{A}_η provides the inequality

$$v_1(x) \geq W(x) - H(\eta) \geq \mu N_7(n, \nu) \tilde{C}_1 \omega \delta(r) (\tilde{z}_n - |x - \tilde{z}|) - H(\eta).$$

It follows from the last inequality with $x = (\tilde{z}', x_n) \in \Omega$ and $0 < x_n \leq \tilde{z}_n - \rho_0/8 = r/4$ that

$$v_1(\tilde{z}', x_n) \geq N_8(n, \nu) \omega \delta(r) x_n - H(\eta). \quad (39)$$

Next, we look for a majorant for v_2 . With this aim in view, we extend the coefficients a_ε^{ij} continuously and the coefficients b_ε^i by zero to the whole annulus $\mathcal{B}(\tilde{z}, \rho_0/8, \tilde{z}_n)$, and denote by $\tilde{v}_2(x)$ the solution of the problem

$$\begin{aligned}\mathcal{L}_\varepsilon \tilde{v}_2 &= \begin{cases} (\mathcal{L}_\varepsilon v_2)_+ & \text{in } \mathcal{A}_\eta, \\ 0 & \text{in } \mathcal{B}(\tilde{z}, \rho_0/8, \tilde{z}_n) \setminus \mathcal{A}_\eta; \end{cases} \\ \tilde{v}_2 &= 0 \quad \text{on } \partial\mathcal{B}(\tilde{z}, \rho_0/8, \tilde{z}_n).\end{aligned}$$

The maximum principle guarantees

$$v_2 \leq \tilde{v}_2 \quad \text{in } \mathcal{A}_\eta. \quad (40)$$

Direct computations show that for $\rho \leq r/4$ the barrier function W satisfies in the set $\mathcal{E}_\rho := \mathcal{P}_\rho(\tilde{z}', 0) \cap \mathcal{B}(\tilde{z}, \rho_0/8, \tilde{z}_n)$ the following inequalities

$$\begin{aligned} |D_n W| &\leq |DW| \leq N_9(n, \nu) \mu \frac{\tilde{k}}{r} \leq N_9 \omega \delta(r), \\ |D'W| &\leq N_9 \mu \frac{\tilde{k} \rho}{r^2} \leq N_9 \omega \frac{\delta(r) \rho}{r}. \end{aligned}$$

So, in view of (15) and (10), we have for all $\rho \leq r/4$ the bounds

$$\begin{aligned} \|(\mathcal{L}_\varepsilon \tilde{v}_2)_+\|_{\mathcal{X}, \mathcal{E}_\rho} &\leq \|b^n\|_{\mathcal{X}, \mathcal{E}_\rho} \left(m^+ + \|D_n W\|_{\infty, \mathcal{E}_\rho} \right) + \|\mathbf{b}'\|_{\mathcal{X}, \mathcal{E}_\rho} \|D'W\|_{\infty, \mathcal{E}_\rho} \\ &\leq N_{10}(n, \nu) \omega \left[\mathfrak{B}\sigma(\rho/\mathcal{R}_0) + \frac{\delta(r)}{r} \rho \|\mathbf{b}'\|_{\mathcal{X}, \mathcal{A}_\eta} \right]. \end{aligned}$$

Since the function $\rho \mapsto \left[\mathfrak{B}\sigma(\rho/\mathcal{R}_0) + \frac{\delta(r)}{r} \rho \|\mathbf{b}'\|_{\mathcal{X}, \mathcal{A}_\eta} \right]$ satisfies the Dini condition at zero, there exist the uniquely defined function $\sigma_1 \in \mathcal{D}_1$ and a constant \mathfrak{B}_1 such that

$$\mathfrak{B}\sigma(\rho/\mathcal{R}_0) + \frac{\delta(r)}{r} \rho \|\mathbf{b}'\|_{\mathcal{X}, \mathcal{A}_\eta} = \mathfrak{B}_1 \sigma_1(4\rho/r).$$

Thus, we may apply Lemma 3.2 to the function \tilde{v}_2 . It gives for $\rho = r/4$ the estimate

$$\sup_{0 < x_n < r/4} \frac{\tilde{v}_2(\tilde{z}', x_n)}{x_n} \leq C_4 \left((r/4)^{-1} \sup_{\mathcal{E}_{r/4}} \tilde{v}_2 + N_{10} \omega \mathfrak{B}_1 \mathcal{J}_{\sigma_1}(1) \right). \quad (41)$$

It is easy to see that

$$\mathfrak{B}_1 \mathcal{J}_{\sigma_1}(1) = \mathfrak{B} \mathcal{J}_\sigma \left(\frac{r}{4\mathcal{R}_0} \right) + \frac{\delta(r)}{4} \|\mathbf{b}'\|_{\mathcal{X}, \mathcal{A}_\eta}.$$

Furthermore, applying (11) to \tilde{v}_2 and to the operator \mathcal{L}_ε in $\mathcal{B}(\tilde{z}, \rho_0/8, \tilde{z}_n)$, we obtain

$$\sup_{\mathcal{E}_{r/4}} \tilde{v}_2 \leq \sup_{\mathcal{B}(\tilde{z}, \rho_0/8, \tilde{z}_n)} \tilde{v}_2 \leq N_{11}(n, \nu, \|\mathbf{b}\|_{\mathcal{X}, \Omega}) \omega r \left[\mathfrak{B}\sigma \left(\frac{r}{\mathcal{R}_0} \right) + \delta(r) \|\mathbf{b}'\|_{\mathcal{X}, \mathcal{A}_\eta} \right].$$

Substitution of the above estimates in (41) and having regard to (3) provide

$$\sup_{0 < x_n < r/4} \frac{\tilde{v}_2(\tilde{z}', x_n)}{x_n} \leq N_{12} \omega \left[\mathfrak{B} \mathcal{J}_\sigma \left(\frac{r}{\mathcal{R}_0} \right) + \delta(r) \|\mathbf{b}'\|_{\mathcal{X}, \mathcal{A}_\eta} \right], \quad (42)$$

where the constant N_{12} depends only on n , ν and $\|\mathbf{b}\|_{\mathcal{X}, \Omega}$.

Taking into account the inequality (5), the assumption (28), and the evident relation $\|\mathbf{b}'\|_{\mathcal{X}, \mathcal{A}} = o(1)$ as $r \rightarrow 0$, we decrease R_0 such that the property

$$\left[\mathfrak{B} \mathcal{J}_\sigma \left(\frac{r}{\mathcal{R}_0} \right) + \delta(r) \|\mathbf{b}'\|_{\mathcal{X}, \mathcal{A}_\eta} \right] \leq \frac{N_8}{2N_{12}} \delta(r) \quad (43)$$

holds true for all $r \leq R_0$.

Finally, combining (39)-(40) with (42)-(43) we arrive at the estimate

$$v_1(\tilde{z}', x_n) - v_2(\tilde{z}', x_n) \geq \frac{N_8}{2} \omega \delta(r) x_n - H(\eta) \quad (44)$$

for $r \leq R_0$ and $x = (\tilde{z}', x_n) \in \Omega$ with $x_n \in [F(\tilde{z}') + \eta, r/4]$.

Considering in \mathcal{A}_η the function $v_3(x) = v(x) - v_1(x) + v_2(x)$ one can easily see that

$$\mathcal{L}_\varepsilon v_3 = -\mathcal{L}_\varepsilon u \rightarrow 0 \quad \text{in } \mathcal{X}(\mathcal{A}_\eta) \quad \text{as } \varepsilon \rightarrow 0.$$

In addition, $v_3 = 0$ on $\partial \mathcal{A}_\eta$. Applying the maximum principle (11) to $\pm v_3$ and to the operator \mathcal{L}_ε we obtain that the difference $v_1(x) - v_2(x)$ converges to $v(x)$ uniformly in \mathcal{A}_η . Therefore, passing in (44) first to the limit as $\varepsilon \rightarrow 0$ and then as $\eta \rightarrow 0$, we get

$$\frac{v(x)}{x_n} \geq \frac{N_8}{2} \omega \delta(r). \quad (45)$$

for $r \leq R_0$ and $x = (\tilde{z}', x_n) \in \Omega$ with $x_n \in [F(\tilde{z}'), r/4]$.

Since \tilde{z}' can be chosen arbitrarily with only $|\tilde{z}'| \leq \frac{r}{4}$, the estimate (45) gives (29) with $\varkappa = N_8/2$. \square

Theorem 4.2 (Main Theorem). *Let the assumptions of Theorem 4.1 hold, and let $\delta(r) = \max_{|x'| \leq r} \frac{F(x')}{|x'|}$ do not satisfy the Dini condition at zero.*

Then for any function u satisfying (27) the equality

$$\frac{\partial u}{\partial \mathbf{n}}(0) = 0$$

holds true.

Proof. Consider the sequence $r_k = 8^{-k}R_0$, $k \geq 0$, where R_0 is the constant from Theorem 4.1.

Application of Theorem 4.1 to u guarantees for $k \geq 0$ the following inequalities

$$\operatorname{osc}_{\Omega \cap \mathcal{P}_{r_{k+1}}} \frac{u(x)}{x_n} \leq (1 - \varkappa \delta(r_k/2)) \operatorname{osc}_{\Omega \cap \mathcal{P}_{r_k}} \frac{u(x)}{x_n} \leq \operatorname{osc}_{\Omega \cap \mathcal{P}_{R_0}} \frac{u(x)}{x_n} \cdot \prod_{j=0}^k (1 - \varkappa \delta(r_j/2)).$$

Since

$$\sum_{j=0}^{\infty} \ln(1 - \varkappa \delta(r_j/2)) \asymp - \sum_{j=0}^{\infty} \delta(r_j/2) \asymp - \int_0^{r_0} \frac{\delta(r)}{r} dr = -\infty,$$

we have

$$\prod_{j=0}^k (1 - \varkappa \delta(r_j/2)) \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

We recall also that Lemma 3.2 implies the finiteness of the quantity $\operatorname{osc}_{\Omega \cap \mathcal{P}_{R_0}} \frac{u(x)}{x_n}$.

Thus, taking into account that $u|_{\partial \Omega \cap \mathcal{P}_{R_0}} = 0$ we get

$$\left| \frac{\partial u}{\partial \mathbf{n}}(0) \right| = \left| \lim_{x_n \rightarrow 0} \frac{u(0, x_n)}{x_n} \right| \leq \lim_{k \rightarrow \infty} \left| \operatorname{osc}_{\Omega \cap \mathcal{P}_{r_k}} \frac{u(x)}{x_n} \right| = 0,$$

and complete the proof. \square

Acknowledgement

The authors would like to thank the anonymous referee for essential comments.

This work was supported by the Russian Foundation of Basic Research (RFBR) through the grant 15-01-07650 and by the St. Petersburg State University grant 6.38.670.2013.

References

- [ABM⁺11] R. Alvarado, D. Brigham, V. Maz'ya, M. Mitrea, and E. Ziadé, *On the regularity of domains satisfying a uniform hour-glass condition and a sharp version of the Hopf-Oleinik boundary point principle*, Probl. Mat. Anal. **57** (2011), 3–68 [Russian], English transl. in J. Math. Sci. (N.Y.) **176**, no. 3 (2011), 281–360.

- [Ale60] A. D. Aleksandrov, *Certain estimates concerning the Dirichlet problem*, Dokl. Akad. Nauk SSSR **134** (1960), no. 5, 1000–1004 [Russian], English transl. in Soviet Math. Dokl. **1** (1961), 1151–1154. MR 0147776 (26 #5290)
- [Ale63] ———, *Uniqueness conditions and estimates for a solution of the Dirichlet problem*, Vest. Leningr. Univ. Ser. Mat. Mekh. Astron. **18** (1963), no. 3, 5–29 [Russian]. MR 0164135 (29 #1434)
- [Alv11] R. Alvarado, *Topics in harmonic analysis and partial differential equations: extension theorems and geometric maximum principles*, Masters Thesis, University of Missouri, 2011.
- [AN00] D. E. Apushkinskaya and A. I. Nazarov, *The Dirichlet problem for quasilinear elliptic equations in domains with smooth closed edges*, Probl. Mat. Anal. **21** (2000), 3–29 [Russian], English transl. in J. Math. Sci. (N.Y.) **105**, no. 5 (2001), 2299–2318. MR 1855435 (2002m:35061)
- [AU95] D. E. Apushkinskaya and N. N. Ural'tseva, *On the behavior of the free boundary near the boundary of the domain*, Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **221** (1995), 5–19 [Russian], English transl. in J. Math. Sci. (N.Y.) **87**, no. 2 (1997), 3267–3276. MR 1359745 (96m:35340)
- [Bak61] I. Ya. Bakel'man, *On the theory of quasilinear elliptic equations*, Sib. Mat. Zh. **2** (1961), no. 179–186 [Russian]. MR 0126604 (23 A3900)
- [Duo01] J. Duoandikoetxea, *Fourier analysis*, Graduate Studies in Mathematics, vol. 29, American Mathematical Society, Providence, RI, 2001.
- [Gir32] G. Giraud, *Generalisation des problèmes sur les opérations du type elliptique*, Bull. des Sciences Math. **56** (1932), 316–352.
- [Gir33] ———, *Problèmes de valeurs à la frontière relatifs à certaines données discontinues*, Bull. Soc. Math. France **61** (1933), 1–54. MR 1504997
- [Him70] B. N. Himčenko, *On the behavior of solutions of elliptic equations near the boundary of a domain of type $A^{(1)}$* , Dokl. Akad. Nauk SSSR **193** (1970), 304–305 [Russian], English transl. in Soviet Math. Dokl. **11** (1970), 943–944. MR 0273190 (42 #8071)

- [Hop52] E. Hopf, *A remark on linear elliptic differential equations of second order*, Proc. Amer. Math. Soc. **3** (1952), 791–793. MR 0050126 (14,280b)
- [JN61] F. John and L. Nirenberg, *On functions of bounded mean oscillation*, Comm. Pure Appl. Math. **14** (1961), no. 415–426.
- [KA82] L. V. Kantorovich and G. P. Akilov, *Functional analysis*, second ed., Translated from the Russian by H.L. Silcock, Pergamon Press, Oxford-Elmsford, N.Y., 1982.
- [KH75] L. I. Kamynin and B. N. Himčenko, *Theorems of Giraud type for a second order elliptic operator that is weakly degenerate near the boundary*, Dokl. Akad. Nauk SSSR **224** (1975), no. 4, 752–755 [Russian], English transl. in Soviet Math. Dokl. **16**, No. 5 (1975), 1287-1291.
- [KH77] ———, *Theorems of Giraud type for second order equations with a weakly degenerate non-negative characteristic part*, Sibirsk. Mat. Ž. **18** (1977), no. 1, 103–121 [Russian], English transl. in Sib. Math. J. **18** (1977), 76-91.
- [KL37] M. V. Keldysh and M. A. Lavrent’ev, *On the uniqueness of the Neumann problem*, Dokl. Akad. Nauk SSSR **16** (1937), no. 3, 151–152 [Russian].
- [Kor01] A. Korn, *Lehrbuch der potentialtheorie. II. allgemeine theorie des logarithmischen potentials und der potentialfunctionen in der ebene*, Berlin: F. Dümmler, 1901.
- [Kry08] N. V. Krylov, *Lectures on elliptic and parabolic equations in Sobolev spaces*, Graduate Studies in Mathematics, vol. 96, American Mathematical Society, Providence, RI, 2008.
- [Lic24] L. Lichtenstein, *Neue Beiträge zur Theorie der linearen partiellen Differentialgleichungen zweiter Ordnung vom elliptischen Typus*, Math. Zeitschr. **20** (1924), 194–212.
- [Lie85] G. M. Lieberman, *Regularized distance and its applications*, Pacific J. Math. **117** (1985), no. 2, 329–352. MR 779924 (87j:35101)
- [LU85] O. A. Ladyzhenskaya and N. N. Ural’tseva, *Estimates of the Hölder constant for functions satisfying a uniformly elliptic or*

- uniformly parabolic quasilinear inequality with unbounded coefficients*, Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) **147** (1985), 72–94, 204, Boundary value problems of mathematical physics and related problems in the theory of functions, No. 17. MR MR821476 (87e:35090)
- [LU88] O. A. Ladyzhenskaya and N. N. Ural'tseva, *Estimates on the boundary of a domain for the first derivatives of functions satisfying an elliptic and parabolic inequality*, Trudy MIAN SSSR **179** (1988), 102–125 [Russian], English transl. in Proc. Steklov Inst. Math., Issue 2 (1989), 109–135. MR 0964915 (89k:35083)
- [MS15] H. Mikayelyan and H. Shahgholian, *Hopf's lemma for a class of singular/degenerate PDE's*, Ann. Acad. Sci. Fenn. **40** (2015), 475–484.
- [Nad83] N. S. Nadirashvili, *On the question of the uniqueness of the solution of the second boundary value problem for second-order elliptic equations*, Mat. Sb. (N.S.) **122 (164)** (1983), no. 3, 341–359 [Russian], English transl. in Math. USSR - Sbornik, **50**, No. 2 (1985), 25–341. MR 721393 (85f:35070)
- [Naz01] A. I. Nazarov, *Estimates for the maximum of solutions of elliptic and parabolic equations in terms of weighted norms of the right-hand side*, Algebra i Analiz **13** (2001), no. 2, 151–164 [Russian], English transl. in St. Petersburg Math. J., **13**, No. 2 (2002), 269–279. MR 1834864 (2002j:35049)
- [Naz05] ———, *The maximum principle of A.D. Aleksandrov*, Sovrem. Mat. Prilozh. (2005), no. 29, 129–145 [Russian], English transl. in J. Math. Sci. (N.Y.) **142**, no. 3 (2007), 2154–2171. MR 2465040 (2011b:35051)
- [Naz12] ———, *A centennial of the Zaremba-Hopf-Oleinik lemma*, SIAM J. Math. Anal. **44** (2012), no. 1, 437–453. MR 2888295
- [Neu88] C. Neumann, *Über die Methode des arithmetischen Mittels*, Abhand. der Königl. Sächsischen Ges. der Wissenschaften. Leipzig **10** (1888), 662–702.
- [NU09] A. I. Nazarov and N. N. Ural'tseva, *Qualitative properties of solutions to elliptic and parabolic equations with unbounded lower-order coefficients*, preprint 2009-05, St. Petersburg Math. Soc. El. Prepr. Archive, 2009.

- [Ole52] O. A. Oleĭnik, *On properties of solutions of certain boundary problems for equations of elliptic type*, Mat. Sb. (N.S.) **30 (72)** (1952), 695–702. MR 0050125 (14,280a)
- [Saf08] M. V. Safonov, *Boundary estimates for positive solutions to second order elliptic equations*, preprint, <http://arxiv.org/abs/0810.0522>, 2008.
- [Saf10] ———, *Non-divergence elliptic equations of second order with unbounded drift*, Nonlinear partial differential equations and related topics, vol. 229, Amer. Math. Soc. Transl. Ser. 2, no. 211–232, Amer. Math. Soc., Providence, RI, 2010, pp. 211–232.
- [Swe97] G. Sweers, *Hopf’s lemma and two dimensional domains with corners*, Rend. Istit. Mat. Univ. Trieste **XXVIII** (1997), 383–419.
- [Tol83] P. Tolksdorf, *On the Dirichlet problem for quasilinear equations in domains with conical boundary points*, Comm. Partial Differential Equations **8** (1983), no. 7, 773–817.
- [Ura96] N. N. Ural’tseva, *C^1 regularity of the boundary of a noncoincident set in a problem with an obstacle*, Algebra i Analiz **8** (1996), no. 2, 205–221, English transl. in *J. Math. Sci. (N.Y.)*, **87** (2) (1997), 3267–3276. MR 1392033 (97m:35105)
- [Wid67] K.-O. Widman, *Inequalities for the Green function and boundary continuity of the gradient of solutions of elliptic differential equations*, Math. Scand. **21** (1967), 17–37. MR 0239264 (39 #621)
- [Zar10] S. Zaremba, *Sur un problème mixte relatif à l’ équation de Laplace*, Bull. Acad. Sci. Cracovie. Cl. Sci. Math. Nat. Ser. A (1910), 313–344.

DEPARTMENT OF MATHEMATICS, SAARLAND UNIVERSITY, P.O. BOX 151150, SAARBRÜCKEN 66041, GERMANY and FACULTY OF MATHEMATICS AND MECHANICS, ST. PETERSBURG STATE UNIVERSITY, UNIVERSITetskii PR. 28, ST. PETERSBURG 198504, RUSSIA

E-mail address: darya@math.uni-sb.de

ST. PETERSBURG DEPARTMENT OF STEKLOV INSTITUTE, FONTANKA 27, ST. PETERSBURG 191023, RUSSIA and FACULTY OF MATHEMATICS AND MECHANICS, ST. PETERSBURG STATE UNIVERSITY, UNIVERSITetskii PR. 28, ST. PETERSBURG 198504, RUSSIA

E-mail address: al.il.nazarov@gmail.com