

How to deal with number of decision making units and variables in data envelopment analysis

Dariush Khezrimotlagh*

Department of Applied Statistics, Faculty of Economics and Administration, University of Malaya, Kuala Lumpur, Malaysia, dariush@um.edu.my, January 01, 2014.

Abstract

Enough number of Decision Making Units (DMUs) in comparison with number of input and output variables has been a concern of using Data Envelopment Analysis (DEA) in the last three decades. There are several studies in the literature of DEA which have been tried to handle this issue by providing additional procedures to increase the number of DMUs, decreasing the number of variables or finding a relation between the number of DMUs and variables. However, this paper illustrates a simple methodology to rectify this concern without any extra conditions and additional methodologies. A geometric reason is illustrated to depict the validity of method which is measured by Kourosh and Arash Model (KAM). The technique is quite simple with no computational complexities of current methodologies even if the number of DMUs is less than the number of variables. A real-life numerical example of 32 DMUs with 45 variables demonstrates the advantages of proposed technique.

Keywords: DEA, KAM, Efficiency, Technical efficiency, Ranking.

1. Introduction

Charnes *et al.* (1978) proposed Data Envelopment Analysis (DEA) to assess the relative efficiency of a set of homogenous Decision Making Units (DMUs) with multiple inputs and multiple outputs. DEA has dramatically been developed in the last three decades, and known as a popular non-parametric technique which only requires a simple set of inputs and outputs values. A description on DEA literature can be seen in Ray (2005).

Although, DEA provides a good number of new insights and additional information not available from conventional econometric methods (Seiford and Thrall, 1990), there is a concern to use DEA due to have enough number of DMUs in comparison with number of variables. Indeed, as the number of DMUs decreases or the number of variables increases, the technically efficient DMUs are increased which decreases discriminative power of DEA.

There are several studies in the literature of DEA which have provided methodologies to reduce the number of variables, or find a relation between the number of DMUs and variables such as the studies of Golany and Roll (1989), Banker *et al.* (1989), Pedraja-Chaparro *et al.* (1999), Dyson *et al.* (2001), Jenkins and Anderson (2003), Cooper *et al.* (2007), Morita and Avkiran (2009) and Osman *et al.* (2011). However, this study unlike the previous studies illustrates that there is no concern about the number of DMUs in comparison with the number of variables while the new proposed technique is applied. The new technique can be measured by Kourosh and Arash Model (KAM) which was recently proposed by Khezrimotlagh *et al.* (2013) to improve the DEA capabilities to distinguish between efficient and technically efficient DMUs.

The rest of this paper is organized in five sections. Section 2 is a short background on KAM. Section 3 represents a methodology to distinguish between technically efficient DMUs without concerning about the number of DMUs and variables. A real numerical example proposed by Osman *et al.* (2011) is represented in Section 4 to depict the advantages of KAM to deal with this issue and the paper is concluded in the last section. Simulations are also performed using Microsoft Excel Solver as it required simple linear programming.

2. Background on KAM

Suppose that there are n DMUs (DMU $_i$, $i = 1, 2, \dots, n$) with m non-negative inputs (x_{ij} , $j = 1, 2, \dots, m$) and p non-negative outputs (y_{ik} , $k = 1, 2, \dots, p$), such that, at least one of the inputs and one of the outputs

* e-mail address: khezrimotlagh@gmail.com

of each DMU are not zero, and for every i there is a j such that $x_{ij} \neq 0$ and also for every i there is a k such that $y_{ik} \neq 0$. The linear KAM is as follows for an appropriate $\epsilon = (\epsilon^-, \epsilon^+) \in \mathbb{R}_+^{m+p}$, where ϵ^- is $(\epsilon_1^-, \epsilon_2^-, \dots, \epsilon_m^-)$ and ϵ^+ is $(\epsilon_1^+, \epsilon_2^+, \dots, \epsilon_p^+)$ (Khezrimotlagh *et al.* 2013):

$$\begin{aligned} & \max \sum_{j=1}^m w_j^- s_{lj}^- + \sum_{k=1}^p w_k^+ s_{lk}^+, \\ & \text{Such that} \\ & \sum_{i=1}^n \lambda_i x_{ij} + s_{lj}^- = x_{lj} + \epsilon_j^-, \quad \forall j, & \sum_{i=1}^n \lambda_i y_{ik} - s_{lk}^+ = y_{lk} - \epsilon_k^+, \quad \forall k, \\ & x_{lj} - s_{lj}^- \geq 0, \quad \forall j, & y_{lk} + s_{lk}^+ - 2\epsilon_k^+ \geq 0, \quad \forall k, \\ & \sum_{i=1}^n \lambda_i = 1; & \lambda_i \geq 0, \quad \forall i, \\ & s_{lj}^- \geq 0, \quad \forall j, & s_{lk}^+ \geq 0, \quad \forall k. \end{aligned}$$

The best technical target and score with ϵ -Degree of Freedom (DF) are respectively depicted as:

$$\begin{cases} x_{lj}^* = x_{lj} - s_{lj}^- + \epsilon_j^-, \quad \forall j, \\ y_{lk}^* = y_{lk} + s_{lk}^+ - \epsilon_k^+, \quad \forall k, \end{cases} \quad KA_{\epsilon}^* = \frac{\sum_{k=1}^p w_k^+ y_{lk} / \sum_{j=1}^m w_j^- x_{lj}}{\sum_{k=1}^p w_k^+ y_{lk}^* / \sum_{j=1}^m w_j^- x_{lj}^*}.$$

After the optimization the following definition is able to identify the efficient DMUs regarding the goals of evolution (Khezrimotlagh *et al.* 2013).

Definition: A technical efficient DMU is KAM efficient with ϵ -DF in inputs and outputs if $KA_0^* - KA_{\epsilon}^* \leq \delta$. Otherwise, it is inefficient with ϵ -DF in inputs and outputs. The proposed amount for δ is ' $10^{-1}\epsilon$ ' or ' $\epsilon/(m+p)$ '.

3. Deal with number of DMUs and variables

It has almost always been suggested in the literature of DEA that there should be an enough number of observations in comparison with a selected number of factors in order to discriminate between DMUs suitably. For example, Pedraja-Chaparro *et al.* (1999) said DEA loses its discrimination power in terms of number of efficient and inefficient units when the value of $n/(m+p)$ is too small. Golany and Roll (1989) suggested that n should be greater than $2 \times (m+p)$, whereas Banker *et al.* (1989), Fried-man and Sinuany-Stern (1998) and Cooper *et al.* (2007) proposed that it should be greater than $3 \times (m+p)$. Dyson (2001) recommended that n should be greater than $2 \times m \times p$. However, there should not be a concern about number of DMUs and selecting factors, although, a good number of DMUs is more appropriate. In this section is illustrated how to deal with the number of DMUs even if it is less than the number of variables.

Suppose that there are two non-dominated DMUs A and B with single constant input and two output values which are commensurate with unity scale. Two different situations for A and B can be considered in output spaces as depicted in Figures 1 and 2.




Figure 1: Two technically efficient DMUs.




Figure 2: Different frontiers in output space.

Since the output values are considered commensurate with unity scale, it is clear to know that the situation of B is better than the situation of A, because the efficiency scores of A and B are $(4+10)/\text{Input}$ and $(10+9)/\text{Input}$, respectively. Now let us change the direct view to measure the efficiency of A, and look at the neighbors of A in Production Possibility Set (PPS). In other words, assume that, for instance A', is under evaluation. As the arrows in Figures 1 and 2 depict, A' does not suggest to A while optimum slacks are calculated, and it is suggested toward B. However, as the place of B is changed to B₁ and B₂, that is, the situation of B becomes weaker in comparison with the situation of A, the optimum slacks for A' become smaller and smaller, respectively. When the situation of B is changed to B₂, A' is suggested to A which shows that the situation of A is good enough, that is, the situation of A should be the same as the situation of B₂ or better than it.

For every two set of DMUs the above simple method can be exemplified. If in practice the frontier is the same as AB₁ it is obvious that there is no differences between the efficiency score of A and B₁, but if there is a very small difference between their efficiency, measuring the efficiency of their neighbors is enough to discriminate between two DMUs.

Now suppose that the number of DMUs is increased as shown in Figures 3 and 4. The above illustrated methodology is easily able to distinguish between DMUs A-H. However, if DMUs lie on the segment line AH (Figure 3), there is no difference between their efficiency scores, however, the differences will be increased as Figure 4 illustrated. In Figure 3, if neighbors of DMUs with subscript 1 are considered there is no differences in their technical efficiency scores. But it has a huge difference when DMUs with subscript 1 in Figure 4 is considered.




Figure 3: Eight technically efficient DMUs.




Figure 4: Two different frontiers in outputs space.

Therefore, number of DMUs does not have any effects on measuring the efficiency score of DMUs while the used technique is to measure the mix efficiency scores of DMUs' neighbors. It is enough to check one or two neighbors of a DMU to see whether its efficiency is the same as the evaluated DMU. If the differences between the efficiency scores of an evaluated DMU and its neighbors are great, the DMU should not be known as an efficient DMU, although, it may be a technically efficient DMU. As Khezrimotlagh *et al.* (2013) quoted the technical efficiency is a necessary condition for being efficiency and it is not enough to call a DMU as fully efficient. Efficiency also depends to internal factors while productivity depends to both internal and external factors, and their definitions should not wrongly be interpreted.

Moreover, radial models are rarely suitable for measuring the relative efficiency scores of DMUs and benchmark them. They only consider one of the input and output orientations. Although, Khezrimotlagh *et al.* (2012) demonstrated the problem of radial models to benchmark DMUs, a simple counter example is to compare the fractions $3/4$ and $4/5$ (or $4/3$ or $5/4$) which show the ratio of output/input for two different targets. If a DMU can select one of the input (output) 4 and 5, an input (output) orientation model suggests input 4 (output 5), however, this means selecting a target with smaller efficiency score. In other words, radial models suppose that all parts of the Farrell frontier have the same efficiency scores, whereas Khezrimotlagh *et al.* (2013) argued this shortcoming in the construction and literature of DEA, and recommended to avoid using radial models.

The above valuable technique can easily be measured by KAM and the proposed definition in Section 2 is also good enough to suggest efficient DMUs from technically efficient ones. If a DMU is in a good place

in comparison with other available DMUs, its neighbors should be suggested to the DMU and the differences of their efficiency scores should approximately be the same. In other words, KAM by considering a commensurate error in data, measures the efficiency scores of DMUs' neighbors, and identify the efficient DMUs with epsilon degree of freedom. Next section depicts the advantages of the above technique while the number of DMUs is 32 and the number of variables is 45.

4. A real-life numerical example

Let us consider the example of Osman *et al.* (2011) with 32 nurses and 45 factors. They classified 20 of factors into 6 input groups and the remained factors into 9 output groups by calculating the average of factors and rounding data with two decimal digits. It is clear this assessing misses some of the information, however using the technology of KAM is simply able to discriminate between these nurses without missing any of information. Applying 0-KAM in Variable Returns to Scale (VRS) technology shows that all nurses were technically efficient except the second nurse who was inefficient. Indeed, VRS is considered due to increase the number of technically efficient DMUs.

In order to apply KAM, suppose that $\varepsilon = 0.001$, that is, only thousandth errors in each factor which is quite negligible according to data and what were missed in the practice of Osman *et al.* (2011). Select the components of epsilon vector as $\varepsilon_j^- = \varepsilon \times x_{1j}$, and $\varepsilon_k^+ = \varepsilon \times x_{1k}$, for $j = 1, 2, \dots, 20$ and $k = 1, 2, \dots, 25$, which means KAM measures the efficiency score of an evaluated DMU's neighbor which has only one tenth percentages errors in its data in comparison with the evaluated DMU. For instance, the distance of the selected neighbor from the first nurse in the PPS is 0.022 only which is quite negligible. Assume that the weights are considered as $w_j^- = 1/x_{1j}$, and $w_k^+ = 1/y_{1k}$, for $j = 1, 2, \dots, 20$ and $k = 1, 2, \dots, 25$.

Now let us apply 0.001-KAM in VRS for these 32 nurses with 20 input and 25 output variables. The results are depicted in the following figure which was sorted from most efficient nurses to least efficient nurses.




Figure 5: 0.001-KAM efficiency scores for 32 nurses with 45 variables.

Nurses 32, 30, 25, 9, 8 and 22 are the first six efficient nurses and nurses 5, 11, 6, 12, 29 and 2 are the last six ones. The average of 32 efficiency scores is 0.996514 with the standard division of 0.016887, and the interquartile mean is 0.999657 with the standard division of 0.000105. According to Definition of Section 2, 0.001-KAM only knows the first four nurses as efficient with 0.001-DF where $\delta = 0.0001$. It means, if only tenth percentages errors are introduced in data, only nurses 32, 30, 25 and 9 have good combinations of their factors and other nurses, although, they are technically efficient, they should improve their efficiency.

For another experience, suppose that $\varepsilon = 0.1$ with the previous assumptions. Figure 6 depicts the results of 0.1-KAM which was sorted from the most efficient nurses to least efficient ones. The average of efficiency scores is 0.956893 with standard division of 0.03565 and the interquartile mean is 0.967210 with the standard division of 0.009959. The first four nurses 32, 30, 25 and 9 have still the best rank, but nurse 1 gets the rank better than nurses 8 and 22 while 10 percentages errors are introduced in the data. Moreover, the technically efficient nurses 9 and 12 even get the rank worse than inefficient nurse 2 with 0.1-DF.

The above valuable outcome is only one of the advantages of KAM to improve discrimination power of DEA. KAM can consider variety of weights for factors when the weights are unknown. If the weights of

factors are available the results of KAM is the same as the results of allocation models (Khezrimotlagh, 2014a). A discussion on how to select an epsilon for applying KAM can also be seen in (Khezrimotlagh 2014b).




Figure 6: 0.1-KAM efficiency scores for 32 nurses with 45 variables.

5. Conclusion

This paper clearly illustrates how DEA is easily able to deal with number of DMUs in comparison with number of variables without any additional procedures, hybrid methodologies and computational complexities. The paper presents an obvious geometrical reason to prove that there is no concerns about number of DMUs with arbitrary number of variables in DEA while KAM is applied. A real life application of 32 DMUs with 45 factors represents the advantages of KAM in comparison with current parametric and non-parametric methodologies to measure the performance evaluations of DMUs.

References

- Banker, R.D., Charnes, A., Cooper, W.W., Swarts, W., Thomas, D., (1989). An introduction to data envelopment analysis with some of its models and their uses. *Research in governmental and nonprofit accounting* 5 (1) 125-163.
- Charnes A., Cooper, W.W., Rhodes, E. (1978). Measuring the inefficiency of decision making units. *European Journal of Operational Research* 2 (6) 429-444.
- Cooper, W.W., Seiford, L.M., Tone, K., (2007). *Data envelopment analysis: a comprehensive text with models, applications, references and DEA-solver software*, 2nd edition. Springer, New York.
- Dyson, R., Allen, R., Camanho, A., Podinovski, V., Sarrico, C., Shale, E., (2001). Pitfalls and protocols in DEA. *European Journal of Operational Research*. 132 (2) 245-259.
- Friedman, L., Sinuany-Stern, Z., (1998). Combining ranking scales and selecting variables in the DEA context: The case of industrial branches. *Computers and Operations Research* 25 (9) 781-791.
- Golany, B.A., Roll, Y., (1989). An application procedure for data envelopment analysis. *Omega* 17 (3) 237-250.
- Jenkins, L., Anderson, M., 2003. A multivariate statistical approach to reducing the number of variables in data envelopment analysis. *European Journal of Operational Research*, 147 (1) 51-61.
- Khezrimotlagh, D., Salleh, S., Mohsenpour, Z., (2012). Benchmarking inefficient Decision Making Units in DEA. *Journal of Basic and Applied Scientific Research* 2 (12) 12056-12065.
- Khezrimotlagh, D., Salleh S., Mohsenpour, Z., (2013). A New Method for Evaluating Decision Making Units in DEA, *Journal of the Operational Research Society*, 65 (1) 694–707.
- Khezrimotlagh, D., (2014a). Profit Efficiency by Kourosh and Arash Model, *Applied Mathematical Sciences*, 8(24): 1165-1170.
- Khezrimotlagh, D., (2014b). How to Select an Epsilon in Kourosh and Arash Model, *The 12th International Conference on Data Envelopment Analysis*, Kuala Lumpur, Malaysia, April 14-17.
- Morita, H., Avkiran, N.K., (2009). Selecting inputs and Outputs in Data Envelopment Analysis by Designing Statistical Experiments. *Journal of the Operations Research Society of Japan*, 52 (2) 163-173.

- Osman, I. H., Berbary, L. N., Sidani, Y., Al-Ayoubi, B., Emrouznejad, A., (2011). Data envelopment analysis model for the appraisal and relative performance evaluation of nurses at an intensive care unit. *Journal of medical systems*, 35 (5) 1039-1062.
- Pedraja-Chaparro, F., Salinas-Jimenez, J., and Smith, P., (1999). On the quality of the data envelopment analysis model. *The Journal of Operational Research Society*, 50 (6) 636-644.
- Ray, S.C., (2004). *Data Envelopment Analysis: Theory and Techniques for Economics and Operations Research*. Cambridge University Press, New York.
- Seiford, L.M., Thrall, R.M., (1990). Recent developments in DEA: the mathematical programming approach to frontier analysis. *Journal of econometrics* 46 (1) 7-38.