

ON THE SELMER GROUP ATTACHED TO A MODULAR FORM AND AN ALGEBRAIC HECKE CHARACTER

Yara Elias¹

Department of Mathematics, McGill University
yara.elias@mail.mcgill.ca

Abstract

We construct an Euler system of generalized Heegner cycles to bound the Selmer group associated to a modular form and an algebraic Hecke character. The main argument is based on Kolyvagin's method adapted by Bertolini and Darmon [2] and by Nekovář [17] while the key object of the Euler system, the generalized Heegner cycles, were first considered by Bertolini, Darmon and Prasanna in [4].

Contents

1	Introduction	2
2	Motive associated to a modular form and a Hecke character	5
3	p-adic Abel-Jacobi map	7
4	Generalized Heegner cycles	8
5	Euler system properties	10
6	Kolyvagin cohomology classes	13
7	Global extensions by Kolyvagin classes	16
8	Complex conjugation	19
9	Local to Global study	21
10	Reciprocity law and local triviality	25

Acknowledgments. It is a pleasure to thank my advisor Henri Darmon for numerous discussions of the subject of this article as well as for his suggestions, corrections and valuable feedback on the writing of this monograph. I am grateful to Olivier Fouquet, Eyal Goren, Ariel Shnidman and the anonymous referees for many corrections and suggestions. This work was finalized at the Max Planck Institute for Mathematics, I am thankful for its hospitality.

¹Supported by a doctoral scholarship of the Fonds Québécois de la Recherche sur la Nature et les Technologies.

1 Introduction

Kolyvagin [14, 11] constructs an Euler system based on Heegner points and uses it to bound the size of the Selmer group of certain (modular) elliptic curves E defined over \mathbb{Q} , over imaginary quadratic fields K assuming the non-vanishing of a suitable Heegner point. In particular, this implies that

$$\text{rank}(E(K)) = 1,$$

and the Tate-Shafarevich group $\text{III}(E/K)$ is finite. Bertolini and Darmon adapt Kolyvagin's descent to Mordell-Weil groups over ring class fields [2]. More precisely, they show that for a given character χ of $\text{Gal}(K_c/K)$ where K_c is the ring class field of K of conductor c ,

$$\text{rank}(E(K_c)^\chi) = 1$$

assuming that the projection of a suitable Heegner point is non-zero. More generally, one can associate to a modular form f of even weight $2r$ and level $\Gamma_0(N)$ a p -adic Galois representation $T_p(f)$ [13, 22]. For a given number field K , there is a p -adic Abel-Jacobi map

$$\Phi_{f,K} : \text{CH}^r(X/K)_0 \longrightarrow H^1(K, T_p(f)),$$

where

- X is the Kuga-Sato variety of dimension $2r - 1$, that is, a compact desingularization of the $2r - 2$ -fold fibre product of the universal generalized elliptic curve over the modular curve $X_1(N)$,
- $\text{CH}^r(X/K)_0$ is the r -th Chow group of X over K , that is the group of homologically trivial cycles on X defined over K of codimension r modulo rational equivalence,
- $H^1(K, T_p(f))$ stands for the first Galois cohomology group of $\text{Gal}(\overline{K}/K)$ acting on $T_p(f)$.

Nekovář [17] adapts the method of Euler systems to modular forms of higher even weight to describe the image by the Abel-Jacobi map $\Phi_{f,K}$ of Heegner cycles on the associated Kuga-Sato varieties, hence showing that

$$\dim_{\mathbb{Q}_p}(\Phi_{f,K}(e_f \text{CH}^r(X/K)_0) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p) = 1$$

for a suitable projector e_f , assuming the non-vanishing of a suitable Heegner cycle. In Article [8], we combined these two approaches to study modular forms of higher even weight twisted by ring class characters χ of imaginary quadratic fields and showed that

$$\dim_{\mathbb{Q}_p}(\Phi_{f,\chi,K}(e_{f,\chi} \text{CH}^r(X/K)_0) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p) = 1$$

for a suitable projector $e_{f,\chi}$, assuming the non-vanishing of a suitable generalized Heegner cycle. In this article, we study the Selmer group associated to a modular form of even weight $r + 2$ and

an unramified algebraic Hecke character ψ of K of infinity type $(r, 0)$. Our setting involves the *generalized Heegner cycles* introduced by Bertolini, Darmon and Prasanna in [4].

Our motivation stems from the Beilinson-Bloch-Kato conjecture that predicts that

$$\dim_{\mathbb{Q}_p}(\Phi_{f,\psi,K}(e_{f,\psi} \text{CH}^r(W/K)_0) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p) = \text{ord}_{s=r+1} L(f \otimes \theta_\psi, s),$$

where

$$\theta_\psi = \sum_{\mathfrak{a} \subset \mathcal{O}_K} \psi(\mathfrak{a}) q^{N(\mathfrak{a})}$$

is the theta series associated to ψ [20, 13], W is a Kuga-Sato like variety, and $e_{f,\psi}$ is a suitable projector.

Let f be a normalized newform of level $\Gamma_0(N)$ where $N \geq 5$ and even weight $r+2 > 2$. Denote by $K = \mathbb{Q}(\sqrt{-D})$ an imaginary quadratic field with odd discriminant satisfying the Heegner hypothesis, that is primes dividing N split in K . For simplicity, we assume that

$$|\mathcal{O}_K^\times| = 2.$$

Let

$$\psi : \mathbb{A}_K^\times \longrightarrow \mathbb{C}^\times$$

be an unramified algebraic Hecke character of K of infinity type $(r, 0)$. There is an elliptic curve A defined over the Hilbert class field K_1 of K with complex multiplication by \mathcal{O}_K . We further assume that A is a \mathbb{Q} -curve, that is A is K_1 -isogenous to its conjugates in $\text{Aut}(K_1)$. This is possible by the assumption on the parity of D , (see [9, Section 11]). Consider a prime p not dividing $ND\phi(N)N_A N_\psi$, where N_A is the conductor of A and N_ψ is the conductor of ψ . We denote by V_f the f -isotypic part of the p -adic étale realization of the motive associated to f by Scholl [22] and Deligne [7] twisted by $\frac{r+2}{2}$ and by V_ψ the p -adic étale realization of the motive associated to ψ twisted by $\frac{r}{2}$. More precisely, V_ψ is the ψ -isotypic component of the first Galois cohomology group of

$$\text{res}_{K_1/\mathbb{Q}}(A) = \prod_{\sigma \in \text{Gal}(K_1/\mathbb{Q})} A^\sigma$$

where A^σ is the σ -conjugate of A , (see Section 2 for more details). Let \mathcal{O}_F be the ring of integers of

$$F = \mathbb{Q}(a_1, a_2, \dots, b_1, b_2, \dots),$$

where the a_i 's are the coefficients of f and the b_i 's are the coefficients of the theta series

$$\theta_\psi = \sum_{\mathfrak{a} \subset \mathcal{O}_K} \psi(\mathfrak{a}) q^{N(\mathfrak{a})}$$

associated to ψ . Then V_f and V_ψ will be viewed (by extending scalars appropriately) as free $\mathcal{O}_F \otimes \mathbb{Z}_p$ -modules of rank 2. We denote by

$$V = V_f \otimes_{\mathcal{O}_F \otimes \mathbb{Z}_p} V_\psi$$

the p -adic étale realization of the tensor product of V_f and V_ψ and let V_\wp be its localization at a prime \wp in F dividing p . Let $\mathcal{O}_{F,\wp}$ be the localization of \mathcal{O}_F at \wp . Then V_\wp is a four dimensional representation of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ over $\mathcal{O}_{F,\wp}$ with coefficients in

$$\text{End}(\text{Res}_{K_1, \mathbb{Q}}(A)/\mathbb{Q}) = \bigoplus_{\sigma \in \text{Gal}(K_1/\mathbb{Q})} \text{Hom}(A, A^\sigma),$$

(see Section 2).

By the Heegner hypothesis, there is an ideal \mathcal{N} of \mathcal{O}_K satisfying

$$\mathcal{O}_K/\mathcal{N} = \mathbb{Z}/N\mathbb{Z}.$$

We can therefore fix $\Gamma_1(N)$ level structure on A , that is a point of exact order N defined over the ray class field L_1 of K of conductor \mathcal{N} . Consider a pair (φ_1, A_1) where A_1 is an elliptic curve defined over K_1 with level N structure and

$$\varphi_1 : A \longrightarrow A_1$$

is an isogeny over \overline{K} . We associate to it a codimension $r+1$ cycle on V

$$\Upsilon_{\varphi_1} = \text{Graph}(\varphi_1)^r \subset (A \times A_1)^r \simeq (A_1)^r \times A^r$$

and define a *generalized Heegner cycle* of conductor 1

$$\Delta_{\varphi_1} = e_r \Upsilon_{\varphi_1},$$

where e_r is an appropriate projector (1). Then Δ_{φ_1} is defined over L_1 . We consider the corestriction

$$P(\varphi_1) = \text{cor}_{L_1, K} \Phi_{f, \psi, L_1}(\Delta_{\varphi_1}) \in H^1(K, V_\wp/p)$$

where Φ_{f, ψ, L_1} is the p -adic étale Abel-Jacobi map. The Selmer group

$$S \subseteq H^1(K, V_\wp/p)$$

consists of the cohomology classes which localizations at a prime v of K lie in

$$\begin{cases} H^1(K_v^{ur}/K_v, V_\wp/p) & \text{for } v \text{ not dividing } NN_A N_\psi p \\ H_f^1(K_v, V_\wp/p) & \text{for } v \text{ dividing } p \end{cases}$$

where K_v is the completion of K at v , and $H_f^1(K_v, V_\wp/p) = H_{\text{cris}}^1(K_v, V_\wp/p)$ is the *finite part* of $H^1(K_v, V_\wp/p)$ [5]. Note that the assumptions we make will ensure that $H^1(K_v^{ur}/K_v, V_\wp/p) = 0$ for v dividing $NN_A N_\psi$. We denote by $Fr(v)$ the arithmetic Frobenius element generating $\text{Gal}(K_v^{ur}/K_v)$, and by $I_v = \text{Gal}(\overline{K}_v/K_v^{ur})$.

Let h be the class number of K , and let $\mathfrak{a}_1, \dots, \mathfrak{a}_h$ be a set of representatives of the ideal classes of \mathcal{O}_K . For a cohomology class c in $H^1(K, V_\wp/p)$, we define

$$c^\Psi = \frac{1}{h} \sum_{i=1}^h \psi^{-1}(\mathfrak{a}_i) \mathfrak{a}_i \cdot c$$

where $\mathfrak{a}_i \cdot c$ is the image of c by the map $H^1(K, V_{\wp}/p) \rightarrow H^1(K, V_{\wp}/p/V_{\wp}/p[\mathfrak{a}_i])$ induced by the projection map

$$V_{\wp}/p \rightarrow V_{\wp}/p/V_{\wp}/p[\mathfrak{a}_i].$$

Then $(c^\psi)^\psi = c^\psi$ and it is independent of the choice of representatives $\mathfrak{a}_1, \dots, \mathfrak{a}_h$.

Theorem 1.1. *Let p be such that*

$$(p, ND\phi(N)N_A N_\psi) = 1.$$

Suppose that V_{\wp}/p is a simple $\text{Aut}_{L_1}(V_{\wp}/p)$ -module and suppose that $\text{Gal}(\overline{L_1}/L_1)$ does not act trivially on V_{\wp}/p . Suppose further that the eigenvalues of $\text{Fr}(v)$ acting on V_{\wp}^v are not equal to 1 modulo p for v dividing $NN_A N_\psi$. Assume that $P(\varphi_1)^\psi \neq 0$ in $H^1(K, V_{\wp}/p)^\psi$. Then the $\overline{\psi}$ -eigenspace of the Selmer group $S^{\overline{\psi}}$ has rank 1 over $\mathcal{O}_{F, \wp}/p$.

To prove Theorem 1.1, we first consider the p -adic étale realization of the twisted motive V associated to f and ψ in the middle étale cohomology of the associated Kuga-Sato variety. This provides us with a p -adic Abel-Jacobi map that lands in the Selmer group S . Next, we construct an Euler system of generalized Heegner cycles which were first considered by Bertolini, Darmon and Prasanna in [4]. These algebraic cycles lie in the domain of the p -adic Abel Jacobi map. In order to bound the rank of the $\overline{\psi}$ -eigenspace of the Selmer group $S^{\overline{\psi}}$, we apply Kolyvagin's descent using local Tate duality, the local reciprocity law, an appropriate global pairing of S and Cebotarev's density theorem.

Our development is an adaptation of Nekovář's techniques [17] and Bertolini and Darmon's approach [2]. The main novelty is that the algebraic Hecke character ψ is of infinite type. In particular, the Galois representation associated to V is four-dimensional over $\mathcal{O}_{F, \wp}/p$.

2 Motive associated to a modular form and a Hecke character

In this section, we describe the construction of the four dimensional $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ -representation

$$V_{\wp} = (V_f \otimes_{\mathcal{O}_F \otimes \mathbb{Z}_p} V_{\psi})_{\wp},$$

where \wp is a prime of F dividing p . Denote by $Y_1(N)$ the affine modular curve over \mathbb{Q} parametrising elliptic curves with level $\Gamma_1(N)$. Let $j : Y_1(N) \hookrightarrow X_1(N)$ be its proper compact desingularization classifying generalized elliptic curves of level $\Gamma_1(N)$. The assumption $N \geq 5$ allows for the definition of the generalized universal elliptic curve $\pi : \mathcal{E} \rightarrow X_1(N)$. Denote by W_r the Kuga-Sato variety of dimension $r+1$, that is a compact desingularization of the r -fold fiber product of \mathcal{E} over $X_1(N)$. We let W be the $2r+1$ -dimensional variety defined by

$$W = W_r \times A^r,$$

where A is as in Section 1. We denote by $[\alpha]$ the element of $\text{End}_{K_1}(A) \otimes_{\mathbb{Z}} \mathbb{Q}$ corresponding to an element α of K . Given two non-singular varieties X and Y over a number field, the group of correspondences $\text{Corr}^m(X, Y) = \text{CH}^{\dim(X)+m}(X \times Y)$ is the group of algebraic cycles of codimension

$\dim(X) + m$ on $X \times Y$ modulo rational equivalence, (see [3, Section 2] for more details). Consider the projectors

$$e_A^{(1)} = \left(\frac{\sqrt{-D} + [\sqrt{-D}]}{2\sqrt{-D}} \right)^{\otimes r} + \left(\frac{\sqrt{-D} - [\sqrt{-D}]}{2\sqrt{-D}} \right)^{\otimes r}, \quad e_A^{(2)} = \left(\frac{1 - [-1]}{2} \right)^{\otimes r}$$

and

$$e_A = e_A^{(1)} \circ e_A^{(2)}$$

in $\mathbb{Q}[\text{End}(A)]^r$. These projectors $e_A^{(1)}$, $e_A^{(2)}$ and e_A belong to the group of correspondences

$$\text{Corr}^0(A^r, A^r)_{\mathbb{Q}} = \text{Corr}^0(A^r, A^r) \otimes_{\mathbb{Z}} \mathbb{Q}$$

from A^r to itself. Let

$$\Gamma_r = (\mathbb{Z}/N \rtimes \mu_2)^r \rtimes \Sigma_r$$

where $\mu_2 = \{\pm 1\}$ and Σ_r is the symmetric group on r elements. Then Γ_r acts on W_r , (see [22, Sections 1.1.0 and 1.1.1] for more details.) The projector e_W in $\mathbb{Z} \left[\frac{1}{2Nr!} \right] [\Gamma_r]$ associated to Γ_r , called Scholl's projector, belongs to the group of zero correspondences $\text{Corr}^0(W_r, W_r)_{\mathbb{Q}}$ from W_r to itself over \mathbb{Q} , (see [3, Section 2.1]). Let

$$e_r = e_W e_A, \tag{1}$$

be the projector in the group of zero correspondences $\text{Corr}^0(W, W)_{\mathbb{Q}}$ from W to itself over \mathbb{Q} . We consider the sheafs

$$\mathcal{F} = j_* \text{Sym}^r(R^1 \pi_* \mathbb{Z}_p) \quad \text{and} \quad \mathcal{F}_A = j_* \text{Sym}^r(R^1 \pi_* \mathbb{Z}_p) \otimes e_A H_{\text{et}}^r(\overline{A^r}, \mathbb{Z}_p).$$

Proposition 2.1. *The étale cohomology group*

$$H_{\text{et}}^1(X_1(N) \otimes \overline{\mathbb{Q}}, \mathcal{F}_A)$$

is isomorphic to

$$e_r H_{\text{et}}^{2r+1}(\overline{W} \otimes \overline{\mathbb{Q}}, \mathbb{Z}_p)$$

and

$$H_{\text{et}}^1(X_1(N) \otimes \overline{\mathbb{Q}}, \mathcal{F}) \otimes e_A H_{\text{et}}^r(\overline{A^r}, \mathbb{Z}_p).$$

Also, we have

$$e_r H_{\text{et}}^i(\overline{W} \otimes \overline{\mathbb{Q}}, \mathbb{Z}_p) = 0$$

for all $i \neq 2r + 1$.

Proof. The proof is a combination of [22, theorem 1.2.1] and [4, proposition 2.4]. Note that the proof in [22, theorem 1.2.1] involves \mathbb{Q}_p coefficients but it is still valid in our setting, (see the Remark following [17, Proposition 2.1]). \square

Let $B = \Gamma_0(N)/\Gamma_1(N)$. We define

$$\tilde{V} = e_B H_{et}^1(X_1(N) \otimes \overline{\mathbb{Q}}, \mathcal{F}_A)(r+1)$$

where $e_B = \frac{1}{|B|} \sum_{b \in B} b$. Given rational primes ℓ coprime to $NN_A N_\psi$, the Hecke operators T_ℓ provide correspondences on $X_1(N)$ [22], inducing endomorphisms of \tilde{V} . Letting

$$I = \text{Ker}\{\mathbb{T} \longrightarrow \mathcal{O}_F : T_\ell \mapsto a_\ell b_\ell, \forall \ell \nmid NN_A N_\psi\},$$

we can define the (f, ψ) -isotypic component of \tilde{V} by

$$V = \tilde{V}/I\tilde{V}.$$

Hence, there is a map $m : \tilde{V} \longrightarrow V$ that is equivariant under the action of Hecke operators T_ℓ , for ℓ not dividing $NN_A N_\psi$ and under the action of the Galois group $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$. The f -isotypic component of $e_B H_{et}^1(X_1(N) \otimes \overline{\mathbb{Q}}, \mathcal{F})\left(\frac{r}{2} + 1\right)$ gives rise (by extending scalars appropriately) to V_f and $e_A H'_{et}(\overline{A^r}, \mathbb{Z}_p)\left(\frac{r}{2}\right)$ gives rise to V_ψ . They are free $\mathcal{O}_F \otimes \mathbb{Z}_p$ -modules of rank 2. Hence,

$$V_{\wp} = (V_f \otimes_{\mathcal{O}_F \otimes \mathbb{Z}_p} V_\psi)_{\wp}$$

is a four dimensional representation of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ over $\mathcal{O}_{F, \wp}$ with coefficients in

$$\text{End}(A/\mathbb{Q}) = \bigoplus_{\sigma \in \text{Gal}(H/\mathbb{Q})} \text{Hom}(A, A^\sigma).$$

3 p-adic Abel-Jacobi map

We use Proposition 2.1 to view the p -adic étale realization of the twisted motive V associated to f and ψ in the middle étale cohomology of the associated Kuga-Sato varieties. For an integer n with $(n, pNN_A N_\psi) = 1$, let

$$L_n = L_1 \cdot K_n$$

be the compositum of the ring class field K_n of K of conductor n with the ray class field L_1 of K of conductor \mathcal{N} . Consider the p -adic étale Abel-Jacobi map

$$\text{CH}^{r+1}(W/L_n)_0 \longrightarrow H^1(L_n, H_{et}^{2r+1}(W \otimes \overline{\mathbb{Q}}, \mathbb{Z}_p(r+1))),$$

where $\text{CH}^{r+1}(W/L_n)_0$ is the group of homologically trivial cycles of codimension $r+1$ on W defined over L_n , modulo rational equivalence. This map factors through $e_r(\text{CH}^{r+1}(W/L_n)_0 \otimes \mathbb{Z}_p)$ as the Abel-Jacobi map commutes with correspondences on W . Composing the Abel-Jacobi map with the projectors e_r and e_B and with $m : \tilde{V} \longrightarrow V$, we obtain a map

$$\Phi_{f, \psi, L_n} : e_r(\text{CH}^{r+1}(W/L_n)_0 \otimes \mathbb{Z}_p) \longrightarrow H^1(L_n, V),$$

which is $\mathbb{T}[\text{Gal}(L_n/\mathbb{Q})]$ -equivariant.

Conjectures and motivation. Beilinson [1, Conjecture 5.9] predicts that

$$\dim_{\mathbb{Q}} e_r \mathrm{CH}^{r+1}(W/K)_0 = \mathrm{ord}_{s=r+1} L(f \otimes \theta_{\psi}, s).$$

Bloch and Kato [5] conjecture that

$$\Phi_{f, \psi, K} : e_r \mathrm{CH}^{r+1}(W/K)_0 \otimes_{\mathbb{Q}} \mathbb{Q}_p \longrightarrow S \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$$

is an isomorphism. As a consequence, one expects that

$$\dim_{\mathbb{Q}_p}(S \otimes_{\mathbb{Z}_p} \mathbb{Q}_p) = \mathrm{ord}_{s=r+1} L(f \otimes \theta_{\psi}, s).$$

In certain settings to be relaxed in his forthcoming work, Shnidman [23, Remark X.2] relates $L'(f \otimes \theta_{\psi}, r+1)$ to the complex valued height $\langle \cdot, \cdot \rangle_B$ of Δ_{φ_1} defined by Beilinson [1], that is

$$L'(f \otimes \theta_{\psi}, r+1) = \langle \Delta_{\varphi_1}, \Delta_{\varphi_1} \rangle_B$$

up to an explicit constant. However, it is worth noting that the pairing $\langle \cdot, \cdot \rangle_B$ is not known to be non-degenerate.

Kolyvagin's results [14] combined with those of Gross and Zagier [12] prove the Birch and Swinnerton-Dyer conjecture for analytic rank less than or equal to 1. This is the particular case of the Beilinson-Bloch-Kato conjectures where the modular form f is associated to an elliptic curve and ψ is the trivial character. Nekovář's results [17, 18] that correspond to the setting where ψ is trivial provide further evidence towards a p -adic analog of the Beilinson-Bloch-Kato conjecture of the form

$$\dim_{\mathbb{Q}_p}(\mathrm{Im}(\Phi_{f, K}) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p) = \mathrm{ord}_{s=r+1} L_p(f, s)$$

due to Perrin-Riou [6, section 2.8], [19]. Shnidman [23] relates the order of vanishing

$$\mathrm{ord}_{s=r+1} L_p(f \otimes \theta_{\psi}, s)$$

of the p -adic L -function at $s = r + 1$ to the height of the image by the p -adic Abel-Jacobi map of an appropriate generalized Heegner cycle of conductor 1. In this article, we prove that

$$\dim_{\mathbb{Q}_p}(\mathrm{Im}(\Phi_{f, \psi, K})^{\overline{\psi}} \otimes_{\mathbb{Z}_p} \mathbb{Q}_p) = 1.$$

4 Generalized Heegner cycles

We describe the construction of generalized Heegner cycles following Bertolini, Darmon and Prasanna [4]. Consider pairs (φ_i, A_i) where A_i is an elliptic curve with level N structure and

$$\varphi_i : A \longrightarrow A_i$$

is an isogeny over \overline{K} . Two pairs

$$(\varphi_i, A_i), (\varphi_j, A_j)$$

are said to be *isomorphic* if there is a \bar{K} -isomorphism $\alpha : A_i \rightarrow A_j$ satisfying $\alpha \circ \varphi_i = \varphi_j$. Recall that \mathcal{N} is an ideal of \mathcal{O}_K such that $\mathcal{O}_K/\mathcal{N} = \mathbb{Z}/N\mathbb{Z}$. Let $Isog^{\mathcal{N}}(A)$ denote the isomorphism classes of pairs (φ_i, A_i) with $\ker(\varphi_i) \cap A[\mathcal{N}]$ trivial. For (φ_i, A_i) in $Isog^{\mathcal{N}}(A)$, we associate a codimension $r+1$ cycle on W

$$\Upsilon_{\varphi_i} = \text{graph}(\varphi_i)^r \subset (A \times A_i)^r \simeq (A_i)^r \times A^r \subset W_r \times A^r$$

and define a *generalized Heegner cycle*

$$\Delta_{\varphi_i} = e_r \Upsilon_{\varphi_i}.$$

We view $\text{graph}(\varphi_i)$ as a subset of $A_i \times A$. Denote by D_{A_i} the element

$$(\text{graph}(\varphi_i) - 0 \times A - \deg(\varphi_i)(A_i \times 0)) \text{ in } NS(A_i \times A),$$

where $NS(A_i \times A)$ is the Néron-Severi group of $A_i \times A$. Then by an adaptation of [18, Paragraph II.3.2], we have

$$\Delta_{\varphi_i} = D_{A_i}^r.$$

Let us assume that the index i of A_i indicates that $\text{End}(A_i)$, which is an order in \mathcal{O}_K , has conductor i . Then Δ_{φ_i} is defined over the compositum of the abelian extension \tilde{K} of K over which the isomorphism class of A is defined, with the ring class field K_i of conductor i . Since \tilde{K} is the smallest extension of K_1 over which $\text{Gal}(\bar{K}/\tilde{K})$ acts trivially on $A[\mathcal{N}]$, it is equal to the ray class field L_1 of K of conductor \mathcal{N} . Therefore, Δ_{φ_i} is defined over

$$L_i = L_1 K_i.$$

Hence,

$$\Delta_{\varphi_i} \text{ belongs to } \text{CH}^{r+1}(W/L_i).$$

In fact, Δ_{φ_i} is homologically trivial on W as shown in [4, proposition 2.7]. In the rest of this section, we consider elements (φ_i, A_i) and (φ_j, A_j) in $Isog^{\mathcal{N}}(A)$.

Lemma 4.1. *Consider the map*

$$g \times I : A_i \times A \rightarrow A_j \times A,$$

where $g : A_i \rightarrow A_j$ is an isogeny of elliptic curves and $I : A \rightarrow A$ is the identity map. Then

$$(g \times I)_* D_{A_i} = \deg(g) \frac{\deg(\varphi_i)}{\deg(\varphi_j)} D_{A_j}.$$

Proof. We denote the intersection pairing of two divisors by a dot. We have

$$(g \times I)_* D_{A_i} \cdot (g \times I)_* D_{A_i} = \deg(g)^2 D_{A_i} \cdot D_{A_i},$$

where

$$\begin{aligned}
& D_{A_i} \cdot D_{A_i} \\
&= (\text{graph}(\varphi_i) - 0 \times A - \deg(\varphi_i)A_i \times 0) \cdot (\text{graph}(\varphi_i) - 0 \times A - \deg(\varphi_i)A_i \times 0) \\
&= \text{graph}(\varphi_i) \cdot \text{graph}(\varphi_i) + 0 \times A \cdot 0 \times A + \deg(\varphi_i)A_i \times 0 \cdot \deg(\varphi_i)A_i \times 0 \\
&\quad - 2\text{graph}(\varphi_i) \cdot 0 \times A - 2\text{graph}(\varphi_i) \cdot \deg(\varphi_i)A_i \times 0 + 2\deg(\varphi_i)A_i \times 0 \cdot 0 \times A \\
&= 0 + 0 + 0 - 2\deg^2(\varphi_i) - 2\deg(\varphi_i) + 2\deg(\varphi_i) \\
&= -2\deg^2(\varphi_i).
\end{aligned}$$

In the previous computation, the equality $\text{graph}(\varphi_i) \cdot \text{graph}(\varphi_i) = 0$ follows from the implication

$$(x, \varphi_i(x)) = (x, \varphi_i(x) + P) \implies P = 0$$

for a translation of $\varphi_i(x)$ by some quantity P . Hence,

$$(g \times I)_* D_{A_i} \cdot (g \times I)_* D_{A_i} = -2\deg(g)^2 \deg^2(\varphi_i).$$

The induced map

$$(g \times I)_* : NS(A_i \times A) \longrightarrow NS(A_j \times A)$$

respects the subgroups generated by complex multiplication cycles. Then, since

$$(g \times I)_* D_{A_i} = kD_{A_j},$$

where $A_j = g(A_i)$ and $k > 0$, we have

$$(g \times I)_* D_{A_i} \cdot (g \times I)_* D_{A_i} = k^2 D_{A_j} \cdot D_{A_j} = -2k^2 \deg^2(\varphi_j).$$

The equality $-2\deg(g)^2 \deg^2(\varphi_i) = -2k^2 \deg^2(\varphi_j)$ then implies that

$$k = \deg(g) \frac{\deg(\varphi_i)}{\deg(\varphi_j)},$$

and

$$(g \times I)_* D_{A_i} = \deg(g) \frac{\deg(\varphi_i)}{\deg(\varphi_j)} D_{A_j}.$$

□

5 Euler system properties

We study certain global and local norm compatibilities of generalized Heegner cycles satisfying the properties of Euler systems. We have $\mathcal{O}_F \otimes \mathbb{Z}_p = \bigoplus_{\varphi_i|p} \mathcal{O}_{F, \varphi_i}$ where $\mathcal{O}_{F, \varphi_i}$ is the completion of

\mathcal{O}_F at the prime \wp_i dividing p . Recall that $V_{\wp} = (V_f \otimes_{\mathcal{O}_F} V_{\psi})_{\wp}$ where \wp is a prime of F dividing p . For a Galois representation V ,

$$F(V)$$

will designate the smallest extension of F such that $\text{Gal}(\overline{F}/F(V))$ acts trivially on V . We denote by $\text{Frob}_v(F_1/F_2)$ the conjugacy class of the Frobenius substitution of the prime $v \in F_2$ in $\text{Gal}(F_1/F_2)$ and by $\text{Frob}_{\infty}(F_1/\mathbb{Q})$ the conjugacy class of the complex conjugation τ in $\text{Gal}(F_1/\mathbb{Q})$. A rational prime ℓ is called a *Kolyvagin prime* if

$$(\ell, NDN_A N_{\psi} p) = 1 \text{ and } a_{\ell} b_{\ell} \equiv \ell + 1 \equiv a_{\ell}^2 - b_{\ell}^2 + 4 \equiv 0 \pmod{p}. \quad (2)$$

Let

$$L = K(\mu_p)(V_{\wp}/p),$$

where μ_p is the group of p -th roots of unity. Condition (2) is equivalent to

$$\text{Frob}_{\ell}(L/\mathbb{Q}) = \text{Frob}_{\infty}(L/\mathbb{Q}). \quad (3)$$

Indeed, it is enough to compare the characteristic polynomial of the complex conjugation $(x^2 - 1)^2 = x^4 - 2x^2 + 1$ acting on V_{\wp}/p with roots -1 and 1 , each with multiplicity 2, with the twist by $r+1$ of the characteristic polynomial of the Frobenius substitution at ℓ acting on V_{\wp}/p with roots

$$\alpha_1 \alpha_3, \alpha_1 \alpha_4, \alpha_2 \alpha_3, \text{ and } \alpha_2 \alpha_4$$

satisfying

$$\alpha_1 \alpha_2 = \ell^r, \alpha_1 + \alpha_2 = b_{\ell}, \alpha_3 \alpha_4 = \ell^{r+1}, \alpha_3 + \alpha_4 = a_{\ell}.$$

The characteristic polynomial of $\text{Frob}(\ell)$ acting on V_{\wp}/p is

$$\begin{aligned} & (x - \alpha_1 \alpha_3)(x - \alpha_1 \alpha_4)(x - \alpha_2 \alpha_3)(x - \alpha_2 \alpha_4) \\ &= (x^2 - (\alpha_1 \alpha_3 + \alpha_1 \alpha_4)x + \alpha_1^2 \alpha_3 \alpha_4)(x^2 - (\alpha_2 \alpha_3 + \alpha_2 \alpha_4)x + \alpha_2^2 \alpha_3 \alpha_4) \\ &= (x^2 - \alpha_1 a_{\ell} x + \ell^{r+1} \alpha_1^2)(x^2 - \alpha_2 a_{\ell} x + \ell^{r+1} \alpha_2^2) \end{aligned}$$

We use the equality $(\alpha_1 + \alpha_2)^2 = \alpha_1^2 + \alpha_2^2 + 2\alpha_1 \alpha_2$ that is $b_{\ell}^2 - 2\ell^r = \alpha_1^2 + \alpha_2^2$ to conclude that the latter equals

$$\begin{aligned} & x^4 - (\alpha_2 a_{\ell} + \alpha_1 a_{\ell}) x^3 + (\ell^{r+1} \alpha_1^2 + \ell^{r+1} \alpha_2^2 + \alpha_1 \alpha_2 a_{\ell}^2) x^2 \\ & \quad - \ell^{r+1} (\alpha_1 a_{\ell} \alpha_2^2 + \alpha_2 a_{\ell} \alpha_1^2) x + \ell^{2r+2} \alpha_1^2 \alpha_2^2 \\ &= x^4 - a_{\ell} b_{\ell} x^3 + (\ell^{r+1} b_{\ell}^2 - 2\ell^{2r+1} + a_{\ell}^2 \ell^r) x^2 - \ell^{2r+1} a_{\ell} (\alpha_1 + \alpha_2) x + \ell^{4r+2} \\ &= x^4 - a_{\ell} b_{\ell} x^3 + (\ell^{r+1} b_{\ell}^2 - 2\ell^{2r+1} + a_{\ell}^2 \ell^r) x^2 - \ell^{2r+1} b_{\ell} a_{\ell} x + \ell^{4r+2}. \end{aligned}$$

To twist this characteristic polynomial by ℓ^{r+1} , it is enough to map $x \mapsto \ell^{r+1} x$. We obtain

$$\begin{aligned} & \ell^{4r+4} x^4 - a_{\ell} b_{\ell} \ell^{3r+3} x^3 + \ell^{2r+2} (\ell^{r+1} b_{\ell}^2 - 2\ell^{2r+1} + a_{\ell}^2 \ell^r) x^2 - \ell^{3r+2} b_{\ell} a_{\ell} x + \ell^{4r+2} \\ &= \ell^{4r+4} \left(x^4 - \frac{a_{\ell} b_{\ell}}{\ell^{r+1}} x^3 + \frac{\ell^{r+1} b_{\ell}^2 - 2\ell^{2r+1} + a_{\ell}^2 \ell^r}{\ell^{2r+2}} x^2 - \frac{b_{\ell} a_{\ell}}{\ell^{r+2}} x + \frac{1}{\ell^2} \right). \end{aligned}$$

On the one hand, using the congruences

$$a_\ell b_\ell \equiv \ell + 1 \equiv a_\ell^2 - b_\ell^2 + 4 \equiv 0 \pmod{p},$$

we find that the characteristic polynomial

$$x^4 - 2x^2 + 1$$

of the complex conjugation τ acting on V_{\wp}/p is congruent to the characteristic polynomial of $\text{Frob}(\ell)$ acting on V_{\wp}/p . On the other hand, comparing the action of the Frobenius element Frob_ℓ and the complex conjugation τ on ζ_p , where ζ_p is a p -th root of unity, we obtain

$$\zeta_p^\ell = \text{Frob}_\ell(\zeta_p) = \text{Frob}_\infty(\zeta_p) = \zeta_p^{-1}.$$

This implies that $\ell \equiv -1 \pmod{p}$. As a consequence, Condition (2) is necessary to satisfy Equality (3).

Let $n = \ell_1 \cdots \ell_k$ be a squarefree integer where ℓ_i is a Kolyvagin prime for $i = 1, \dots, k$. The Galois group $\text{Gal}(K_n/K_1)$ is the product over the primes ℓ dividing n of the cyclic groups

$$G_\ell = \text{Gal}(K_\ell/K_1)$$

of order $\ell + 1$. We denote by σ_ℓ a generator of G_ℓ . The extensions L_1 and K_n are disjoint over K_1 . Indeed, if $I = L_1 \cap K_n$ is a non-trivial extension of K_1 , then $\text{Gal}(I/K_1) = \prod_{\ell_j} \text{Gal}(K_{\ell_j}/K)$ for some ℓ_j dividing n . This implies that the primes ℓ_j ramify in I and hence also in L_1 , a contradiction since $(n, N) = 1$. Hence

$$G_n = \text{Gal}(L_n/L_1) \simeq \text{Gal}(K_n/K_1).$$

The Frobenius condition on ℓ implies that it is inert in K . Denote by λ the unique prime in K above ℓ . Writing n as $n = \ell m$, we have that λ splits completely in L_m since it is unramified in L_m and has the same image as $\text{Frob}_\infty(L/K) = \tau^2 = \text{Id}$ by the Artin map. A prime λ_m of L_m above λ ramifies completely in L_n . We denote by λ_n the unique prime in L_n above λ_m . Consider the image of Δ_{φ_n} by the Abel-Jacobi map

$$\Phi_{f, \psi, L_n} : e_r(\text{CH}^{r+1}(W/L_n)_0 \otimes \mathbb{Z}_p) \longrightarrow H^1(L_n, V).$$

Proposition 5.1. *Consider $(A_n, \varphi_n) \sim (A_m, \varphi_m) \in \text{Isog}^{\mathcal{N}}(A)$ where $n = \ell m$ for an odd prime ℓ . Then*

$$T_\ell \Phi_{f, \psi, L_m}(\Delta_{\varphi_m}) = \text{cor}_{L_n, L_m} \Phi_{f, \psi, L_n}(\Delta_{\varphi_n}) = a_\ell b_\ell \Phi_{f, \psi, L_m}(\Delta_{\varphi_m}).$$

Proof. By [21, corollary 11.4],

$$T_\ell(\Delta_{\varphi_m}) = \sum_{n_i} \Delta_{\varphi_{n_i}},$$

where the generalized Heegner cycles $\Delta_{\varphi_{n_i}}$ correspond to elements

$$(A_{n_i}, \varphi_{n_i}) \sim (A_m, \varphi_m)$$

in $\text{Isog}^{\mathcal{N}}(A)$ for elliptic curves A_{n_i} that are ℓ -isogenous to A_m respecting level N structure. These elliptic curves A_{n_i} correspond to gA_m where

$$g \in \text{Gal}(L_n/L_m) \simeq \text{Gal}(K_n/K_m) \simeq \text{Gal}(K_\ell/K_1).$$

Hence

$$\sum_{n_i} \Delta_{\varphi_{n_i}} = \sum_{g \in \text{Gal}(L_n/L_m)} g \Delta_{\varphi_n} = \text{cor}_{L_n, L_m}(\Delta_{\varphi_n}) = a_\ell b_\ell \Delta_{\varphi_m},$$

where the last equality follows from the action of T_ℓ on V . Finally, we apply the p-adic Abel-Jacobi map which commutes with T_ℓ to obtain $T_\ell \Phi_{f, \psi, L_m}(\Delta_{\varphi_m}) = \text{cor}_{L_n, L_m} \Phi_{f, \psi, L_n}(\Delta_{\varphi_n})$. \square

For an element $c \in H^1(F, M)$, we denote by $\text{res}_v(c) \in H^1(F_v, M)$ the image of c by the restriction map $H^1(F, M) \rightarrow H^1(F_v, M)$ induced from the inclusion

$$\text{Gal}(\overline{F}_v/F_v) \hookrightarrow \text{Gal}(\overline{F}/F).$$

Proposition 5.2. *Consider $(A_n, \varphi_n) \sim (A_m, \varphi_m) \in \text{Isog}^{\mathcal{N}}(A)$ where $n = \ell m$. Then*

$$\text{res}_{\lambda_n} \Phi_{f, \psi, L_n}(\Delta_{\varphi_n}) = k \text{Frob}_\ell(L_n/L_m) \text{res}_{\lambda_m} \Phi_{f, \psi, L_m}(\Delta_{\varphi_m})$$

for $k = \ell \frac{\deg(\varphi_i)}{\deg(\varphi_j)}$.

Proof. Since λ_m completely ramifies in L_n , we have

$$\mathcal{O}_{L_m}/\lambda_m \simeq \mathcal{O}_{L_n}/\lambda_n,$$

which is isomorphic to the finite field with ℓ^2 elements as ℓ is inert in K . As a consequence, the reductions of the elliptic curves A_n and A_m at λ_n and λ_m are supersingular. Hence the ℓ -isogeny from A_n to A_m reduces to the Frobenius morphism Frob_ℓ . Therefore, we have $\text{Frob}_\ell(A_m) \equiv A_n \pmod{\lambda_n}$. By Proposition 4.1, this implies

$$(\text{Frob}_\ell \times I)_* D_{A_m} \equiv k D_{A_n} \pmod{\lambda_n}$$

where $k = \ell \frac{\deg(\varphi_i)}{\deg(\varphi_j)}$ from which the result follows. \square

6 Kolyvagin cohomology classes

We denote by

$$\Phi_{f, \psi, L_n}(\Delta_{\varphi_n})_{\wp} \in H^1(L_n, V_{\wp})$$

the image of $\Phi_{f, \psi, L_n}(\Delta_{\varphi_n}) \in H^1(L_n, V)$ by the map $H^1(L_n, V) \rightarrow H^1(L_n, V_{\wp})$ induced by the projection $V \rightarrow V_{\wp}$. Let

$$y_{\varphi_n} = \text{red}(\Phi_{f, \psi, L_n}(\Delta_{\varphi_n})_{\wp}) \in H^1(L_n, V_{\wp}/p)$$

be the image of $\Phi_{f,\psi,L_n}(\Delta_{\varphi_n})_{\wp} \in H^1(L_n, V_{\wp})$ by the map $H^1(L_n, V_{\wp}) \longrightarrow H^1(L_n, V_{\wp}/p)$ induced by the projection $V_{\wp} \rightarrow V_{\wp}/p$. We use certain operators (4) defined by Kolyvagin in order to lift the cohomology classes $y_{\varphi_n} \in H^1(L_n, V_{\wp}/p)$ to Kolyvagin cohomology classes $P(\varphi_n) \in H^1(K, V_{\wp}/p)$, for appropriate n .

Lemma 6.1. *For all n ,*

$$H^0(L_n, V_{\wp}/p) = H^0(L_1, V_{\wp}/p) = 0$$

and $\text{Gal}(L_n(V_{\wp}/p)/L_n) \simeq \text{Gal}(L_1(V_{\wp}/p)/L_1)$

Proof. The extensions L_n/L_1 and $L_1(V_{\wp}/p)/L_1$ are unramified outside primes dividing n and $N_A N_{\psi} N_p$. Therefore, $L_n \cap L_1(V_{\wp}/p)$ is unramified over L_1 and is hence contained in L_1 , the maximal unramified extension of K of conductor \mathcal{N} . Hence,

$$H^0(L_n, V_{\wp}/p) = H^0(L_1, V_{\wp}/p).$$

The result follows from the assumption that V_{\wp}/p is a simple $\text{Aut}_{L_1}(V_{\wp}/p)$ -module and $\text{Gal}(\overline{L_1}/L_1)$ does not act trivially on V_{\wp}/p . \square

Proposition 6.2. *The restriction map*

$$\text{res}_{L_1, L_n} : H^1(L_1, V_{\wp}/p) \longrightarrow H^1(L_n, V_{\wp}/p)^{G_n}$$

is an isomorphism.

Proof. Recall that $G_n = \text{Gal}(L_n/L_1)$ and let $G = \text{Gal}(\overline{\mathbb{Q}}/L_n)$. The result follows from the inflation-restriction sequence

$$0 \rightarrow H^1(L_n/L_1, (V_{\wp}/p)^G) \xrightarrow{\text{inf}} H^1(L_1, V_{\wp}/p) \xrightarrow{\text{res}} H^1(L_n, V_{\wp}/p)^{G_n} \rightarrow H^2(L_n/L_1, (V_{\wp}/p)^G),$$

since $H^0(L_n, V_{\wp}/p) = 0$ by Lemma 6.1. \square

Let

$$\text{Tr}_{\ell} = \sum_{i=0}^{\ell} \sigma_{\ell}^i, \quad D_{\ell} = \sum_{i=1}^{\ell} i \sigma_{\ell}^i \quad \text{in } \mathbb{Z}[G_{\ell}]. \quad (4)$$

They are related by

$$(\sigma_{\ell} - 1)D_{\ell} = \ell + 1 - \text{Tr}_{\ell}. \quad (5)$$

Define

$$D_n = \prod_{\ell|n} D_{\ell} \in \mathbb{Z}[G_n].$$

Proposition 6.3.

$$D_n y_{\varphi_n} \in H^1(L_n, V_{\wp}/p)^{G_n}.$$

Proof. It is enough to show that for all ℓ dividing n ,

$$(\sigma_\ell - 1)D_n y_{\varphi_n} = 0.$$

We have

$$(\sigma_\ell - 1)D_n = (\sigma_\ell - 1)D_\ell D_m = (\ell + 1 - \text{Tr}_\ell)D_m,$$

where the last equality follows by Relation (5). Since $\text{res}_{L_m, L_n} \circ \text{cor}_{L_n, L_m} = \text{Tr}_\ell$,

$$\begin{aligned} & (\ell + 1 - \text{Tr}_\ell)D_m \text{red}(\Phi_{f, \psi, L_n}(\Delta_{\varphi_n})_{\wp}) \\ &= (\ell + 1)D_m \text{red}(\Phi_{f, \psi, L_n}(\Delta_{\varphi_n})_{\wp}) - D_m a_\ell b_\ell \text{red}(\Phi_{f, \psi, L_m}(\Delta_{\varphi_m})_{\wp}) \\ &\equiv 0 \pmod{p}. \end{aligned}$$

by Proposition 5.1 and Condition 2. □

As a consequence, the cohomology classes $D_n y_{\varphi_n} \in H^1(L_n, V_{\wp}/p)^{G_n}$ can be lifted to cohomology classes $c(\varphi_n) \in H^1(L_1, V_{\wp}/p)$ such that

$$\text{res}_{L_1, L_n} c(\varphi_n) = D_n y_{\varphi_n}.$$

We define

$$P(\varphi_n) = \text{cor}_{L_1, K} c(\varphi_n) \text{ in } H^1(K, V_{\wp}/p).$$

For a place v of K and a cohomology class c in $H^1(K, V_{\wp}/p)$, we denote by $\text{res}_v(c)$ the image of c by the map $H^1(K, V_{\wp}/p) \rightarrow H^1(K_v, V_{\wp}/p)$ induced by the inclusion $G_{K_v} \hookrightarrow G_K$.

Proposition 6.4. *Let v be a prime of L_1 .*

1. *If $v | N_A N_\psi N$, then $\text{res}_v(P(\varphi_n))$ is trivial.*
2. *If $v \nmid N_A N_\psi N n p$, then $\text{res}_v(P(\varphi_n))$ lies in $H^1(K_v^{ur}/K_v, V_{\wp}/p)$.*

Proof. 1. We denote by

$$V_{\wp}/p^{\text{dual}} = \text{Hom}(V_{\wp}/p, \mathbb{Z}/p\mathbb{Z}(1))$$

the local Tate dual of V_{\wp}/p . The local Euler characteristic formula [16, Section 1.2] yields

$$|H^1(K_v, V_{\wp}/p)| = |H^0(K_v, V_{\wp}/p)| \times |H^2(K_v, V_{\wp}/p)|.$$

Local Tate duality then implies

$$|H^1(K_v, V_{\wp}/p)| = |H^0(K_v, V_{\wp}/p)| \times |H^0(K_v, V_{\wp}/p^{\text{dual}})| = |H^0(K_v, V_{\wp}/p)|^2$$

as V_{\wp}/p is self-dual. The Weil conjectures and the assumption on $Fr(v)$ imply that

$$((V_{\wp}/p)^{I_v})^{Fr(v)} = 0$$

where $\langle Fr(v) \rangle = \text{Gal}(K_v^{ur}/K_v)$ and $I = \text{Gal}(\overline{K}_v/K_v^{ur})$ is the inertia group. Therefore,

$$H^0(K_v, V_{\wp}/p) = ((V_{\wp}/p)^{I_v})^{Fr(v)} = 0.$$

2. If v does not divide $N_A N_{\psi} N n p$, then

$$\text{res}_{L_{1,v}, L_{n,v'}} \text{res}_v c(\varphi_n) = \text{res}_{v'} D_n y_n \in H^1(\overline{L}_{n,v'}/L_{n,v'}, V_{\wp}/p)$$

for v' above v in L_n . The exact sequence

$$\cdots \rightarrow H^1(L_{n,v'}^{ur}/L_{n,v'}, (V_{\wp}/p)^{I_v}) \rightarrow H^1(L_{n,v'}, V_{\wp}/p) \xrightarrow{\text{res}} H^1(\overline{L}_{n,v'}/L_{n,v'}^{ur}, V_{\wp}/p) \rightarrow \cdots$$

allows us to view the cohomology class $\text{res}_{v'} D_n y_n$ that belongs to $\text{Ker}(\text{res})$ as an element in

$$H^1(L_{n,v'}^{ur}/L_{n,v'}, V_{\wp}/p).$$

Since v is unramified in L_n/L_1 , then $\text{res}_v c(\varphi_n)$ belongs to $H^1(L_{1,v}^{ur}/L_{1,v}, V_{\wp}/p)$. □

7 Global extensions by Kolyvagin classes

We consider the restriction d of an element c of $H^1(K, V_{\wp}/p)$ to $H^1(L, V_{\wp}/p)$. Then d factors through some finite extension \tilde{L} of L . We denote by

$$L(c) = \tilde{L}^{\ker(d)}$$

the subfield of \tilde{L} fixed by $\ker(d)$. Note that $L(c)$ is an extension of L . In this section, we study extensions of L by Kolyvagin cohomology classes c and $P(\varphi_q)$, where $P(\varphi_q)$ will play a crucial role in the proof of Theorem 1.1. We recall the statement of the theorem.

Theorem 1.1. Let p be such that

$$(p, ND\phi(N)N_A) = 1.$$

Suppose that V_{\wp}/p is a simple $\text{Aut}_{L_1}(V_{\wp}/p)$ -module and suppose that $\text{Gal}(\overline{L}_1/L_1)$ does not act trivially on V_{\wp}/p . Suppose further that the eigenvalues of $Fr(v)$ acting on $V_{\wp}^{I_v}$ are not equal to 1 modulo p for v dividing $NN_A N_{\psi}$. If $P(\varphi_1)^{\psi}$ is non-zero in $H^1(K, V_{\wp}/p)^{\psi}$, then the $\overline{\psi}$ -eigenspace of the Selmer group $S^{\overline{\psi}}$ has rank 1 over $\mathcal{O}_{F, \wp}/p$.

Recall that $L = K(\mu_p)(V_{\wp}/p)$.

Lemma 7.1. *We have*

$$H^1(\mathrm{Aut}_K(V_{\wp}/p), V_{\wp}/p) = H^1(L/K, V_{\wp}/p) = 0.$$

Proof. First note that if $p \nmid |\mathrm{Aut}_K(V_{\wp}/p)|$, then

$$H^1(\mathrm{Aut}_K(V_{\wp}/p), V_{\wp}/p) = 0.$$

If p divides $|\mathrm{Aut}_K(V_{\wp}/p)|$, then since V_{\wp}/p is irreducible as an $\mathrm{Aut}_K(V_{\wp}/p)$ -module, Dickson's lemma [24, Theorem 6.21] implies that $\mathrm{Aut}_K(V_{\wp}/p)$ contains $\mathrm{SL}_2(F_q)$ for some q . In particular, it contains $2I$ where I is the identity map. Sah's lemma [15, 8.8.1] states that if G is a group, M a G -representation, and g an element of $\mathrm{Center}(G)$, then the map $x \rightarrow (g - 1)x$ is the zero map on $H^1(G, M)$. Therefore, by Sah's lemma, the map $x \mapsto (2I - I)x = Ix$ is the zero map on

$$H^1(\mathrm{Aut}_K(V_{\wp}/p), V_{\wp}/p)$$

implying that $H^1(\mathrm{Aut}_K(V_{\wp}/p), V_{\wp}/p) = 0$. As a consequence, $H^1(K(\mu_p)(V_{\wp}/p)/K, V_{\wp}/p) = 0$ since p does not divide $|\mathrm{Gal}(K(\mu_p)/K)| = p - 1$. \square

We denote the Galois group $\mathrm{Gal}(L/K)$ by G . The restriction map

$$r : H^1(K, V_{\wp}/p) \rightarrow H^1(L, V_{\wp}/p)^G = \mathrm{Hom}_G(\mathrm{Gal}(\overline{\mathbb{Q}}/L), V_{\wp}/p)$$

has kernel

$$\mathrm{Ker}(r) = H^1(L/K, V_{\wp}/p) = 0$$

by Lemma 7.1. Consider the evaluation pairing

$$[\cdot, \cdot] : r(S) \times \mathrm{Gal}(\overline{\mathbb{Q}}/L) \rightarrow V_{\wp}/p. \quad (6)$$

We denote by $\mathrm{Gal}_S(\overline{\mathbb{Q}}/L)$ the annihilator of $r(S)$. Let L^S be the extension of L fixed by $\mathrm{Gal}_S(\overline{\mathbb{Q}}/L)$ and G_S the Galois group $\mathrm{Gal}(L^S/L)$.

Remark 7.2. The element $P(\varphi_1)$ belongs to S by Proposition 6.4. Also, $L(P(\varphi_1))$ is a subfield of L^S . Indeed, assume $\rho \in \mathrm{Gal}_S(\overline{\mathbb{Q}}/L)$, then $[\rho, \rho] = 0$ for all $\rho \in S$. Hence, $P(\varphi_1)$ defines a cocycle of S by

$$\rho \rightarrow \rho(P(\varphi_1)) - P(\varphi_1) = 0.$$

This implies that ρ fixes $L(P(\varphi_1))$, a subfield of L^S .

Proposition 7.3. *There is a Kolyvagin prime q such that*

$$\mathrm{res}_{\beta} P(\varphi_1)^{\psi} \neq 0,$$

where β is a prime dividing q in K .

Proof. By Cebratev's density theorem, there is a Kolyvagin prime q such that

$$\text{Frob}_q(L^S/\mathbb{Q}) = \tau h, \quad h^{1+\tau} \notin \text{Gal}(L^S/L(P(\varphi_1)^\Psi))$$

for some h in $\text{Gal}(L^S/L)$. The restriction of τh to L is indeed τ . Let β be a prime of L above q . Since $\text{Frob}_\beta(L(P(\varphi_1)^\Psi)/L) = (\tau h)^2$ does not fix $P(\varphi_1)^\Psi$, β is not in the kernel of the Artin map. Hence β does not split completely in $L(P(\varphi_1)^\Psi)$. Therefore, $\text{res}_\beta P(\varphi_1)^\Psi \neq 0$. \square

Consider the following extensions

$$\begin{array}{ccccc}
 & & H_0 = H_1 H_2 & & \\
 & \nearrow & \uparrow & \nwarrow & \\
 H_1 = L(P(\varphi_1)) & & v_0 & & H_2 = L(P(\varphi_q)) \\
 & \nwarrow & \vdots & \nearrow & \\
 & & L = K(\mu_p)(V_{\wp}/p) & & \\
 & \swarrow & \downarrow & \searrow & \\
 & & v_1 & & v_2
 \end{array}$$

where q is a Kolyvagin prime satisfying

$$\text{res}_\beta P(\varphi_1)^\Psi \neq 0$$

for some β in L above q (see Proposition (7.3)). We define

$$V_i = \text{Gal}(H_i/F) \quad \text{for } i = 0, 1, 2.$$

An automorphism of $L(P(\varphi_1))/L$ corresponds to the multiplication of the generators of $L(P(\varphi_1))$ over L by an element of $(V_{\wp})_p \simeq V_{\wp}/p$. Hence

$$V_1 = \text{Gal}(L(P(\varphi_1))/L) \simeq V_{\wp}/p.$$

Similarly, we have $V_2 \simeq V_{\wp}/p$. We recall that h is the class number of K , and $\mathfrak{a}_1, \dots, \mathfrak{a}_h$ are a set of representatives of the ideal classes of \mathcal{O}_K . For a cohomology class c in $H^1(K, V_{\wp}/p)$, we defined

$$c^\Psi = \frac{1}{h} \sum_{i=1}^h \psi^{-1}(\mathfrak{a}_i) \mathfrak{a}_i \cdot c,$$

where $\mathfrak{a}_i \cdot c$ is the image of c by the map $H^1(K, V_{\wp}/p) \longrightarrow H^1(K, V_{\wp}/p/V_{\wp}/p [\mathfrak{a}_i])$ induced by the projection map

$$V_{\wp}/p \longrightarrow V_{\wp}/p/V_{\wp}/p [\mathfrak{a}_i].$$

Note that c^Ψ is independent of the choice of representatives $\mathfrak{a}_1, \dots, \mathfrak{a}_h$. Furthermore, $(c^\Psi)^\Psi = c^\Psi$ lies in $H^1(K, V_{\wp}/p)^\Psi$. We denote by

$$H_1^\Psi = L(P(\varphi_1)^\Psi), \quad H_1^{\overline{\Psi}} = L(P(\varphi_1)^{\overline{\Psi}}), \quad H_2^\Psi = L(P(\varphi_q)^\Psi), \quad H_2^{\overline{\Psi}} = L(P(\varphi_q)^{\overline{\Psi}}),$$

and we let

$$V_i^\Psi = \text{Gal}(H_i^\Psi/H_i), \quad V_i^{\overline{\Psi}} = \text{Gal}(H_i^{\overline{\Psi}}/H_i), \quad \text{for } i = 1, 2.$$

We will show that

$$H_1^\Psi \cap H_2^\Psi = L.$$

Lemma 7.4. *There is an isomorphism of $\mathcal{O}_{F,\wp}/p$ -modules*

$$H^1(K_\lambda^{ur}/K_\lambda, V_{\wp}/p) \longrightarrow H^1(K_\lambda^{ur}, V_{\wp}/p)$$

mapping $\text{res}_\lambda P(\varphi_{m\ell})$ to $\text{res}_\lambda P(\varphi_m)$. Also, $\text{res}_\lambda P(\varphi_\ell)$ is ramified.

Proof. This is an adaptation of [8, Section 5] that uses the properties of the Euler system considered in Proposition 5.1 and Proposition 5.2. \square

Recall that τ denotes complex conjugation.

Proposition 7.5. *The extensions H_1^Ψ and H_2^Ψ are linearly disjoint over L .*

Proof. Linearly independent cocycles c_1 and c_2 of $H^1(K, V_{\wp}/p)$ over $\mathcal{O}_{F,\wp}/p$ can be viewed by Lemma (7.1) as linearly independent homomorphisms h_1 and h_2 in $\text{Hom}_{\text{Gal}(L/K)}(L, V_{\wp}/p)$ over $\mathcal{O}_{F,\wp}/p$. Linearly independent homomorphisms h_1 and h_2 induce linearly disjoint extensions $\overline{L}^{\text{Ker}(h_1)}$ and $\overline{L}^{\text{Ker}(h_2)}$ of L . Hence, if H_1 and H_2 were not linearly disjoint over L , we would have that

$$u_1 c_1 + u_2 c_2 = 0, \quad \text{for some } u_1, u_2 \text{ in } (\mathcal{O}_{F,\wp}/p)^*,$$

where

$$c_1 = P(\varphi_1)^\Psi \quad \text{and} \quad c_2 = P(\varphi_q)^\Psi.$$

Lemma (7.4) implies that $\text{res}_\beta P(\varphi_q)^\Psi$ is ramified for $\beta \in K$ dividing q and provides an isomorphism sending $\text{res}_\beta P(\varphi_q)^\Psi$ to $\text{res}_\beta P(\varphi_1)^\Psi$. The latter implies by Proposition (7.3) that $\text{res}_\beta P(\varphi_q)^\Psi$ is non-zero. This is inconsistent with the equality $u_1 c_1 + u_2 c_2 = 0$ since the nontrivial cocycles c_1, c_2 have different ramification properties. \square

8 Complex conjugation

We study the action of complex conjugation on the image by the p -adic Abel-Jacobi map of generalized Heegner cycles. Recall that D_{A_j} is the element

$$(\text{graph}(\varphi_j) - 0 \times A - \text{deg}(\varphi_j)(A_j \times 0)) \text{ in } NS(A_j \times A).$$

In this section, we write D_{A_j, φ_j} instead of D_{A_j} alone in order to keep track of the underlying map φ_j .

Lemma 8.1. *There is an element σ in $\text{Gal}(K_j/K)$ such that*

$$\tau\Phi(\Delta_{\varphi_j})_{\wp} = (-1)^{\frac{r+2}{2}} \varepsilon_L N^{r/2} \deg^{-r}(\sigma) \sigma\Phi(\Delta_{\varphi_j})_{\wp},$$

where ε_L is the sign of the functional equation of $L(f, s)$.

Proof. We have that $(\tau \times I)_*(D_{A_j, \varphi_j}) = D_{\tau(A_j), \overline{\varphi_j}}$. Article [10] shows that $\tau A_j = W_N(\sigma A_j)$ for some σ in $G(K_j/K)$, hence

$$(\tau \times I)_*(D_{A_j, \varphi_j}^r) = D_{\tau(A_j), \overline{\varphi_j}}^r = D_{W_N(\sigma A_j), W_N \circ \sigma \circ \varphi_j}^r.$$

Consider the map

$$W \times I : W_N \times A \longrightarrow W_N \times A : ((E, P), A) \longrightarrow ((E/\langle P \rangle, P'), A),$$

where P' is such that the Weil pairing $\langle P, P' \rangle$ of P with P' satisfies $\langle P, P' \rangle = \zeta_N$ for some choice ζ_N of an N -th root of unity ζ_N . Note that W has degree N . Also,

$$W_* f(\tau) d_\tau d_z = (-1)^{\frac{r+2}{2}} \varepsilon_L N^{r/2} f(\tau) d_\tau d_z.$$

This implies as in [17, Proposition 6.2] that

$$(W \times I)_* D_{W_N(\sigma A_j), W_N \circ \sigma \circ \varphi_j}^r = (-1)^{\frac{r+2}{2}} \varepsilon_L N^{r/2} D_{W_N(\sigma A_j), W_N \circ \sigma \circ \varphi_j}^r,$$

while Proposition 4.1 implies that the former equals $N^r \frac{\deg^r(\varphi_i)}{\deg^r(\varphi_j)} D_{\sigma A_j, \sigma \circ \varphi_j}^r$. Hence,

$$D_{W_N(\sigma A_j), W_N \circ \sigma \circ \varphi_j}^r = (-1)^{\frac{r+2}{2}} \varepsilon_L k_1 N^{r/2} D_{\sigma A_j, \sigma \circ \varphi_j}^r,$$

where $k_1 = \frac{\deg^r(\varphi_i)}{\deg^r(\varphi_j)}$. Applying Proposition 4.1 to the map $(\sigma \times I)$, we obtain

$$(\sigma \times I)_*(D_{A_j, \varphi_j}^r) = k_2 D_{\sigma A_j, \sigma \circ \varphi_j}^r,$$

where $k_2 = \deg^r(\sigma) \frac{\deg^r(\varphi_i)}{\deg^r(\varphi_j)} = \deg^r(\sigma) k_1$. Hence,

$$(\tau \times I)_*(D_{A_j, \varphi_j}^r) = D_{W_N(\sigma A_j), W_N \circ \sigma \circ \varphi_j}^r = (-1)^{\frac{r+2}{2}} \varepsilon_L N^{r/2} \deg^{-r}(\sigma) (\sigma \times I)_*(D_{A_j, \varphi_j}^r).$$

Therefore

$$\tau\Phi(\Delta_{\varphi_j})_{\wp} = (-1)^{\frac{r+2}{2}} \varepsilon_L N^{r/2} \deg^{-r}(\sigma) \sigma\Phi(\Delta_{\varphi_j})_{\wp}.$$

□

Lemma 8.2. Let $\varepsilon = (-1)^{\frac{r+2}{2}} \varepsilon_L$ and $k = \deg^{-r}(\sigma) N^{r/2}$. Then

$$\tau P(\varphi_1)^\Psi = \varepsilon k \sigma P(\varphi_1)^\Psi \text{ and } \tau P(\varphi_q)^\Psi = -\varepsilon k \sigma P(\varphi_q)^\Psi.$$

Proof. For an element c in $H^1(K, V_{\wp}/p)$, we have that

$$\tau \cdot c^\Psi = \frac{1}{h} \sum_{i=1}^h \overline{\psi}^{-1}(\mathfrak{a}_i) \tau \cdot (\mathfrak{a}_i \cdot c) = \frac{1}{h} \sum_{i=1}^h \overline{\psi}^{-1}(\mathfrak{a}_i) \tau(\mathfrak{a}_i \cdot c) \tau^{-1} = \frac{1}{h} \sum_{i=1}^h \overline{\psi}^{-1}(\mathfrak{a}_i) \mathfrak{a}_i \cdot \tau c \tau^{-1} = (\tau c)^\Psi.$$

For $c = P(\varphi_j)$ and σ in $\text{Gal}(K_j/K)$, we have that

$$\sigma \cdot c^\Psi = \frac{1}{h} \sum_{i=1}^h \psi^{-1}(\mathfrak{a}_i) \sigma \cdot (\mathfrak{a}_i \cdot c) = \frac{1}{h} \sum_{i=1}^h \psi^{-1}(\mathfrak{a}_i) \sigma(\mathfrak{a}_i \cdot c) \sigma^{-1} = \frac{1}{h} \sum_{i=1}^h \psi^{-1}(\mathfrak{a}_i) \mathfrak{a}_i \cdot \sigma c \sigma^{-1} = (\sigma c)^\Psi.$$

Hence

$$\tau P(\varphi_1)^\Psi = (\tau P(\varphi_1))^\Psi = \varepsilon k \sigma P(\varphi_1)^\Psi.$$

The identity $\tau D_q = -D_q \tau$ indicates that

$$\tau P(\varphi_q)^\Psi = (\tau P(\varphi_q))^\Psi = -\varepsilon k \sigma P(\varphi_q)^\Psi.$$

□

For $(v_{11}, v_{12}, v_{21}, v_{22})$ in $V_1^\Psi V_1^\Psi V_2^\Psi V_2^\Psi$,

$$\tau(v_{11}, v_{12}, v_{21}, v_{22}) \tau = (\varepsilon k \sigma \tau v_{12}, \varepsilon k \sigma \tau v_{11}, -\varepsilon k \sigma \tau v_{22}, -\varepsilon k \sigma \tau v_{21}).$$

We define

$$U_0 = \{(v_{11}, v_{12}, v_{21}, v_{22}) \mid \varepsilon k \sigma \tau v_{1i} + v_{1i}, -\varepsilon k \sigma \tau v_{2j} + v_{2j}, i, j = 1, 2 \text{ generate } V_{\wp}/p\}.$$

Then U_0^+ generates V_0^+ . We let

$$L(U_0) = \{\ell \text{ rational prime} \mid \text{Frob}_\ell(H_0/\mathbb{Q}) = [\tau u], u \in U_0\}.$$

9 Local to Global study

Local Tate duality. Let K_λ be a local field with residue field F_q and let A be a finite group with an unramified action of $\text{Gal}(\overline{K}_\lambda/K_\lambda)$ killed by a prime p . Assume p divides $q - 1$ so that $\mu_p \subset K_\lambda$ and let $A' = \text{Hom}(A, \mu_p)$. We denote by K_λ' , the maximal tamely ramified extension of K_λ , and by $H_{ur}^1(K_\lambda, *)$, the group $H^1(K_\lambda^{ur}/K_\lambda, *)$. The natural pairing $A \times A' \longrightarrow \mu_p$ yields the cup product pairing

$$H^1(K_\lambda, A) \times H^1(K_\lambda, A') \longrightarrow H^2(K_\lambda, \mu_p) = \mathbb{Z}/p\mathbb{Z}$$

which induces a perfect local Tate pairing

$$H^1(K_\lambda^{ur}/K_\lambda, A) \times H^1(K_\lambda, A')/H^1(K_\lambda^{ur}/K_\lambda, A) \longrightarrow \mathbb{Z}/p\mathbb{Z}.$$

Let

$$\alpha : H^1(K_\lambda^{ur}/K_\lambda, A) \xrightarrow{\sim} A/(\phi - 1)A$$

be the evaluation map at the Frobenius element ϕ where

$$\text{Gal}(K_\lambda^{ur}/K_\lambda) = \langle \phi \rangle.$$

Then α is an isomorphism. The exact sequence of Galois groups

$$0 \longrightarrow \text{Gal}(\overline{K}_\lambda/K_\lambda^t) \longrightarrow \text{Gal}(\overline{K}_\lambda/K_\lambda^{ur}) \longrightarrow \text{Gal}(K_\lambda^t/K_\lambda^{ur}) \longrightarrow 0$$

induces the exact sequence

$$H^1(K_\lambda^t/K_\lambda^{ur}, A') \longrightarrow H^1(K_\lambda^{ur}, A') \longrightarrow H^1(K_\lambda^t, A') \longrightarrow 0,$$

where $H^1(K_\lambda^t, A') = 0$ since $\text{Gal}(\overline{K}_\lambda/K_\lambda^t)$ is a pro- q group. Therefore,

$$H^1(K_\lambda^{ur}, A') \simeq H^1(K_\lambda^t/K_\lambda^{ur}, A') \simeq \text{Hom}(\mathbb{Z}/p\mathbb{Z}(1), A') \simeq \text{Hom}(\mu_p, A').$$

Hence we have an isomorphism

$$H^1(K_\lambda^{ur}, A') \xrightarrow{\sim} \text{Hom}(\mu_p, A').$$

The exact sequence of Galois cohomology groups

$$0 \longrightarrow H^1(K_\lambda^{ur}/K, A') \longrightarrow H^1(K_\lambda, A') \longrightarrow H^1(K_\lambda^{ur}, A')^\phi \longrightarrow 0$$

allows us to identify $H^1(K_\lambda, A')/H^1(K_\lambda^{ur}/K_\lambda, A')$ with

$$H^1(K_\lambda^{ur}, A')^\phi \simeq \text{Hom}(\mu_p, A')^\phi.$$

Hence, we obtain a perfect local pairing

$$\langle \cdot, \cdot \rangle_\lambda : H^1(K_\lambda^{ur}/K_\lambda, A) \times H^1(K_\lambda^{ur}, A')^\phi \longrightarrow \mathbb{Z}/p\mathbb{Z}.$$

Set up of the proof. Given a Kolyvagin prime ℓ , the Frobenius condition implies that it is inert in K . We denote by λ the prime of K lying above ℓ . We have a perfect local pairing

$$\langle \cdot, \cdot \rangle_\lambda : H^1(K_\lambda^{ur}/K_\lambda, (V_{\wp}/p)^{I_\lambda}) \times H^1(K_\lambda^{ur}, V_{\wp}/p) \longrightarrow \mathbb{Z}/p,$$

where $I_\lambda = \text{Gal}(\overline{K}_\lambda/K_\lambda^{ur})$ and $\mathcal{O}_{F, \wp}$ -linear isomorphisms

$$\{H^1(K_\lambda^{ur}, V_{\wp}/p)\}^{\text{dual}} \simeq H^1(K_\lambda^{ur}/K_\lambda, (V_{\wp}/p)^{I_\lambda}) \simeq (V_{\wp}/p)^{I_\lambda}/(\phi - 1), \quad (7)$$

where ϕ generates $\text{Gal}(K_\lambda^{ur}/K_\lambda)$. We denote by

$$res_\lambda : H^1(K, V_\phi/p) \longrightarrow H^1(K_\lambda, V_\phi/p)$$

the restriction map from $H^1(K, V_\phi/p)$ to $H^1(K_\lambda, V_\phi/p)$ induced by the embedding

$$\text{Gal}(\overline{K}_\lambda/K_\lambda) \hookrightarrow \text{Gal}(\overline{K}/K).$$

Restricting the domain of res_λ to the Selmer group S , we obtain

$$res_\lambda : S \longrightarrow H^1(K_\lambda^{ur}/K_\lambda, (V_\phi/p)^{I_\lambda}).$$

Taking the $\mathbb{Z}/p\mathbb{Z}$ -dual of this map, we obtain a homomorphism

$$\omega_\ell : H^1(K_\lambda^{ur}, V_\phi/p) \longrightarrow S^{dual}.$$

We denote

$$X_\ell = \omega_\ell(H^1(K_\lambda^{ur}, V_\phi/p))$$

in S^{dual} . We aim to bound the size of $S^{dual, \psi}$ by studying the set

$$\{X_\ell^\psi\}_{\ell \in L(U_0)}.$$

Proposition 9.1. *The modules $\{X_\ell\}_{\ell \in L(U_0)}$ generate S^{dual} .*

Proof. Let $G = \text{Gal}(L/K)$. Consider an element s of S . The restriction map

$$H^1(K, V_\phi/p) \xrightarrow{\text{res}} H^1(L, V_\phi/p)^G$$

is injective by Lemma (7.1) as

$$\text{Ker}(\text{res}) = H^1(L/K, V_\phi/p) = 0.$$

We identify s with its image by restriction in

$$H^1(L, V_\phi/p)^G \subset \text{Hom}_G(L, V_\phi/p).$$

We will show that

$$\{res_\lambda\}_{\ell \in L(U_0)} : S \longrightarrow \{H^1(K_\lambda^{ur}/K_\lambda, (V_\phi/p)^{I_\lambda})\}_{\ell \in L(U_0)}$$

is injective. As a consequence, the induced map between the duals

$$\{H^1(K_\lambda^{ur}, V_\phi/p)\}_{\ell \in L(U_0)} \longrightarrow S^{dual}$$

is surjective. Hence, it is enough to show that $res_\lambda(s) = 0$ for all $\ell \in L(U_0)$ implies $s = 0$. Consider \widetilde{H}_0 , the minimal Galois extension of \mathbb{Q} containing H_0 such that s factors through $\text{Gal}(\widetilde{H}_0/L)$. Let

x be an element of $\text{Gal}(\widetilde{H}_0/L)$ such that $x|_{H_0}$ belongs to U_0 . By Chebotarev's density theorem, there exists ℓ in $L(U_0)$ such that $\text{Frob}_\ell(\widetilde{H}_0/\mathbb{Q}) = [\tau x]$. The hypothesis $\text{res}_\lambda(s) = 0$ implies that $s(\text{Frob}_{\lambda_L}(\widetilde{H}_0/L)) = 0$ for λ_L above ℓ in L since $\text{Frob}_{\lambda_L}(\widetilde{H}_0/L)$ is a generator of $\text{Gal}(\widetilde{H}_0, \lambda_{\widetilde{H}_0}^-/L_{\lambda_L})$, where $\lambda_{\widetilde{H}_0}^-$ is above λ_L in \widetilde{H}_0 . In fact,

$$\text{Frob}_{\lambda_L}(\widetilde{H}_0/L) = (\tau x)^{|D(L/\mathbb{Q})|} = (\tau x)^2 = x^\tau x = (x^+)^2,$$

where $|D(L/\mathbb{Q})|$ is the order of the decomposition group $D(L/\mathbb{Q})$, also the order of the residue extension. Therefore, $s(x^+) = 0$ for all $x \in \text{Gal}(\widetilde{H}_0/L)$ such that $x|_{H_0}$ belongs to U_0 . Since U_0^+ generates V_0^+ , we have that s vanishes on $\text{Gal}(\widetilde{H}_0/L)^+$. Hence, $\text{Im}(s)$ lies in V_{\wp}/p^- , the minus eigenspace of V_{\wp}/p for the action of τ . In particular, it cannot be a proper G -submodule of V_{\wp}/p . \square

Proposition 9.2. *The elements*

$$\text{res}_\lambda P(\varphi_\ell)^\Psi \text{ and } \text{res}_\lambda P(\varphi_{\ell q})^\Psi$$

generate $H^1(K_\lambda^{ur}, V_{\wp}/p)^\Psi$.

Proof. We have

$$H^1(K_\lambda^{ur}, V_{\wp}/p) \simeq V_{\wp}/p(K_\lambda)$$

by (7). The module $V_{\wp}/p(K_\lambda)^\Psi$ is of rank 2 over $\mathcal{O}_{F,\wp}/p$, hence, so is $H^1(K_\lambda^{ur}, V_{\wp}/p)^\Psi$. The Frobenius condition on ℓ implies that

$$g_1 = \text{res}_\lambda P(\varphi_1)^\Psi \text{ and } g_2 = \text{res}_\lambda P(\varphi_q)^\Psi$$

are linearly independent in $H^1(K_\lambda^{ur}, V_{\wp}/p)^\Psi$ over $\mathcal{O}_{F,\wp}/p$. Indeed, if they were linearly dependent, then

$$g_1^{(\tau x_{11})^2} - g_1, \text{ and } g_2^{(\tau x_{21})^2} - g_2$$

where $\text{Frob}_\ell(H_0/\mathbb{Q}) = \tau u = (\tau x_{11}, \tau x_{12}, \tau x_{21}, \tau x_{22})$ would also be linearly dependent. The Frobenius condition implies that

$$\{\text{Frob}_\ell(H_i/L) = x_{i1}^\tau x_{i1} = (\tau x_{i1})^2, \quad i = 1, 2\}$$

generate (V_{\wp}/p) , which yields a contradiction since $(\tau x_{11})^2$ acts on $\text{res}_\lambda P(\varphi_1)^\Psi$ which generates H_1^Ψ by $g_1^{(\tau x_{11})^2} - g_1$ and $(\tau x_{21})^2$ acts on $\text{res}_\lambda P(\varphi_q)^\Psi$ which generates H_1^Ψ by $g_2^{(\tau x_{21})^2} - g_2$. Therefore, by Lemma (7.4), $\text{res}_\lambda P(\varphi_\ell)^\Psi$ and $\text{res}_\lambda P(\varphi_{\ell q})^\Psi$ are linearly independent in $H^1(K_\lambda^{ur}, V_{\wp}/p)^\Psi$ over $\mathcal{O}_{F,\wp}/p$. \square

10 Reciprocity law and local triviality

In this section, we use the local reciprocity law as well as the local properties of the Kolyvagin cohomology classes $P(\varphi_n)$ to study the modules X_ℓ for ℓ in $L(U_0)$.

Proposition 10.1. *We have*

$$\sum_{\lambda|\ell|n} \langle s_\lambda^\Psi, \text{res}_\lambda P(\varphi_n)^\Psi \rangle_\lambda = 0.$$

Proof. The proof follows [17, proposition 11.2(2)] where both the reciprocity law and the ramification properties of $P(\varphi_n)$ in proposition 6.4 are used. \square

Proposition 10.2. 1. *The element $\omega_\ell(\text{res}_\lambda P(\varphi_\ell)^\Psi)$ vanishes in X_ℓ^Ψ for ℓ in $L(U_0)$.*

2. *The modules $\{X_\ell^\Psi\}_{\ell \in L(U_0)}$ are generated over $O_{F,\wp}/p$ by $\omega_q(\text{res}_\beta P(\varphi_{\ell_0q})^\Psi)$.*

3. *The module S^Ψ is of rank 1 over $O_{F,\wp}/p$.*

Proof. 1. Recall that λ is the prime above ℓ in K . The image of $\text{res}_\lambda P(\varphi_\ell)^\Psi$ by the map

$$\omega_\ell : H^1(K_\lambda^{ur}, V_{\wp}/p)^\Psi \longrightarrow X_\ell^\Psi$$

is the homomorphism

$$S \longrightarrow \mathbb{Z}/p\mathbb{Z} : s^\Psi \mapsto \langle s_\lambda^\Psi, P(\varphi_\ell)^\Psi \rangle_\lambda.$$

Proposition 10.1 implies that

$$\langle s_\lambda^\Psi, P(\varphi_\ell)^\Psi \rangle_\lambda = 0.$$

Hence, the image by ω_ℓ of $\text{res}_\lambda P(\varphi_\ell)^\Psi$, one of the two generators of $H^1(K_\lambda^{ur}, V_{\wp}/p)^\Psi$ by proposition 9.2 vanishes.

2. Let β be the prime above q in K . Proposition (10.1) implies that

$$\langle s_\lambda^\Psi, P(\varphi_{\ell q})^\Psi \rangle_\lambda + \langle s_\beta^\Psi, P(\varphi_{\ell q})^\Psi \rangle_\beta = 0.$$

Hence,

$$\omega_\ell(\text{res}_\lambda P(\varphi_{\ell q})^\Psi) + \omega_q(\text{res}_\beta P(\varphi_{\ell q})^\Psi) = 0.$$

Therefore, X_ℓ^Ψ is generated by $\omega_q(\text{res}_\beta P(\varphi_{\ell q})^\Psi)$ for all $\ell \in L(U_0)$. As a consequence,

$$\{X_\ell^\Psi\}_{\ell \in L(U_0)} \subseteq X_q^\Psi,$$

where the rank one module X_q^Ψ is generated over $O_{F,\wp}/p$ by $\omega_q(\text{res}_\beta P(\varphi_{\ell_0q})^\Psi)$ for some ℓ_0 in $L(U_0)$.

3. By proposition (9.1), the set $\{X_\ell^\Psi\}_{\ell \in L(U_0)}$ generates $S^{\text{dual},\Psi}$. Furthermore, $P(\varphi_1)^\Psi$ belongs to S^Ψ by Proposition (6.4) and is non-zero by the hypothesis on $P(\varphi_1)^\Psi$ and Proposition (8.2). Therefore,

$$\text{rank}(S^\Psi) = \text{rank}(S^{\text{dual},\Psi}) = 1$$

over $O_{F,\wp}/p$. \square

References

- [1] A. A. Beilinson. Height pairing between algebraic cycles. In *K-theory, arithmetic and geometry (Moscow, 1984–1986)*, volume 1289 of *Lecture Notes in Math.*, pages 1–25. Springer, Berlin, 1987.
- [2] Massimo Bertolini and Henri Darmon. Kolyvagin’s descent and Mordell-Weil groups over ring class fields. *J. Reine Angew. Math.*, 412:63–74, 1990.
- [3] Massimo Bertolini, Henri Darmon, and Kartik Prasanna. Chow-Heegner points on CM elliptic curves and values of p -adic L -functions. *International Mathematics Research Notices*, 2012.
- [4] Massimo Bertolini, Henri Darmon, and Kartik Prasanna. Generalized Heegner cycles and p -adic Rankin L -series. *Duke Math. J.*, 162(6):1033–1148, 2013.
- [5] Spencer Bloch and Kazuya Kato. L -functions and Tamagawa numbers of motives. In *The Grothendieck Festschrift, Vol. I*, volume 86 of *Progr. Math.*, pages 333–400. Birkhäuser Boston, Boston, MA, 1990.
- [6] Pierre Colmez. Fonctions L p -adiques. *Séminaire Bourbaki*, 41:21–58, 1998-1999.
- [7] Pierre Deligne. Formes Modulaires et Représentations de $GL(2)$. In *Modular Functions of One Variable II*, volume 349 of *Lecture Notes in Math.*, pages 55–105. Springer Berlin Heidelberg, 1973.
- [8] Yara Elias. Kolyvagin’s method for Chow groups of Kuga-Sato varieties over ring class fields. *Ann. Math. Qué.*, pages 1–21, 2015.
- [9] Benedict H. Gross. *Arithmetic on Elliptic Curves with Complex Multiplication*. PhD thesis, Harvard University, 1980.
- [10] Benedict H. Gross. Heegner points on $X_0(N)$. In *Modular forms (Durham, 1983)*, Ellis Horwood Ser. Math. Appl.: Statist. Oper. Res., pages 87–105. Horwood, Chichester, 1984.
- [11] Benedict H. Gross. Kolyvagin’s work on modular elliptic curves. In *L -functions and arithmetic (Durham, 1989)*, volume 153 of *London Math. Soc. Lecture Note Ser.*, pages 235–256. Cambridge Univ. Press, Cambridge, 1991.
- [12] Benedict H. Gross and Don B. Zagier. Heegner points and derivatives of L -series. *Invent. Math.*, 84(2):225–320, 1986.
- [13] Uwe Jannsen. Algebraic cycles, K -theory, and extension classes. In *Mixed Motives and Algebraic K -Theory*, volume 1400 of *Lecture Notes in Math.*, pages 57–188. Springer Berlin Heidelberg, 1990.

- [14] Victor A. Kolyvagin. *Euler Systems*, volume 2 of *The Grothendieck Festschrift*. Birkhäuser Boston, Boston, MA, 1990.
- [15] Serge Lang. *Fundamentals of Diophantine Geometry*. Springer, 1983.
- [16] James S. Milne. *Arithmetic Duality Theorems*. Perspectives in mathematics. Academic Press, 1986.
- [17] Jan Nekovář. Kolyvagin’s method for Chow groups of Kuga-Sato varieties. *Invent. Math.*, 107(1):99–125, 1992.
- [18] Jan Nekovář. On the p -adic height of Heegner cycles. *Math. Ann.*, 302(1):609–686, 1995.
- [19] Bernadette Perrin-Riou. Points de Heegner et dérivées de fonctions L p -adiques. *Invent. Math.*, 89(3):455–510, 1987.
- [20] Peter Schneider. Introduction to the Beilinson conjectures. *Perspectives in Mathematics*, 4:1–36, 1988.
- [21] Chad Schoen. On the computation of the cycle class map for nullhomologous cycles over the algebraic closure of a finite field. *Ann. Sci.éc. Norm. Supér.*, 28(1):1–50, 1995.
- [22] Anthony J. Scholl. Motives for modular forms. *Invent. Math.*, 100(1):419–430, 1990.
- [23] Ariel Shnidman. p -adic heights of generalized Heegner cycles. *arXiv: 1407.0785v2*, pages 1–34, 2014.
- [24] Michio Suzuki. *Group theory*. Number vol. 1 in Grundlehren der mathematischen Wissenschaften. Springer-Verlag, 1982.