

THE WEIGHTED SINGER CONJECTURE FOR COXETER GROUPS IN DIMENSIONS THREE AND FOUR

WIKTOR J. MOGILSKI

ABSTRACT. Given a Coxeter system (W, S) there is a contractible simplicial complex Σ called the Davis complex on which W acts properly and cocompactly. In [7], the weighted L^2 -(co)homology groups of Σ were defined, and in [5], the Singer conjecture for Coxeter groups was appropriately formulated for weighted L^2 -(co)homology theory. In this article, we prove the weighted version of the Singer conjecture in dimension three under the assumption that the nerve of the Coxeter group is not dual to a hyperbolic simplex, and in dimension four under additional restrictions. We then prove a general version of the conjecture where the nerve of the Coxeter group is assumed to be a flag triangulation of a 3-manifold.

Given a Coxeter system (W, S) with corresponding nerve L , Davis constructs a contractible simplicial complex Σ_L on which W acts properly and cocompactly (for details see [3, 4]). Given an S -tuple $\mathbf{q} = (q_s)_{s \in S}$ of positive real numbers satisfying $q_s = q_{s'}$ if s and s' are conjugate in W , one defines the weighted L^2 -(co)homology spaces $L_{\mathbf{q}}^2 H_k(\Sigma_L)$ (see [5]). They are special in the sense that they admit a notion of dimension: one can attach a nonnegative real number to each of the Hilbert spaces $L_{\mathbf{q}}^2 H_k(\Sigma_L)$ called the von Neumann dimension. Hence one can define weighted L^2 -Betti numbers, denoted by $L_{\mathbf{q}}^2 b_k(\Sigma_L)$. More details on this theory can be found in [3, 5, 7]. When $\mathbf{q} = \mathbf{1}$, the groups $L_{\mathbf{1}}^2 H_k(\Sigma_L)$ are the L^2 -(co)homology groups of Σ_L considered in [6]. There, the classical Singer conjecture for manifolds is reformulated for Coxeter groups, using the fact that if L is a triangulation of the $(n-1)$ -sphere, then Σ_L is an n -manifold.

Conjecture 1 (Singer Conjecture). *Suppose that L is a triangulation of S^{n-1} . Then*

$$L_{\mathbf{1}}^2 H_k(\Sigma_L) = 0 \text{ for } k \neq \frac{n}{2}.$$

Conjecture 1 is known for elementary reasons for $n \leq 2$ and holds by a result of Lott and Lück [9], in conjunction with the validity of the Geometrization Conjecture for 3-manifolds [12], for $n = 3$. It was proved by Davis–Okun [6] for the case where W is right-angled and $n \leq 4$, and furthermore it was shown that Conjecture 1 for n odd implies Conjecture 1 for n even, under the assumption that W is right-angled. It was later proved for the case where W is an even Coxeter group and $n = 4$ by Schroeder [13], under the restriction that L is a flag complex. Recently Okun–Schreve [11, Theorem 4.9] gave a proof of Conjecture 1 for the case $n = 4$, so now Conjecture 1 is known in full generality for dimensions $n \leq 4$.

In [5], the Singer conjecture for Coxeter groups was formulated for weighted L^2 -(co)homology:

Conjecture 2 (Weighted Singer Conjecture). *Suppose that L is a triangulation of S^{n-1} . Then*

$$L_{\mathbf{q}}^2 H_k(\Sigma_L) = 0 \text{ for } k > \frac{n}{2} \text{ and } \mathbf{q} \leq \mathbf{1}.$$

By weighted Poincaré duality, this is equivalent to the conjecture that if $\mathbf{q} \geq \mathbf{1}$ and $k < \frac{n}{2}$, then $L_{\mathbf{q}}^2 H_k(\Sigma_L)$ vanishes. Conjecture 2 is known for elementary reasons for $n \leq 2$, and in [5], it was proved for the case where W is right-angled and $n \leq 4$, and furthermore it was shown that Conjecture 2 for n odd implies Conjecture 2 for n even (also under the assumption that W is right-angled). In this article we prove Conjecture 2 for $n = 3, 4$, under some additional restrictions on the nerve L .

Theorem 3. *Suppose that the nerve L of a Coxeter group is a triangulation of S^2 not dual to a hyperbolic 3-simplex. Then*

$$L_{\mathbf{q}}^2 H_k(\Sigma_L) = 0 \text{ for } k > 1 \text{ and } \mathbf{q} \leq \mathbf{1}.$$

Note that, in conjunction with weighted Poincaré duality and [7, Theorem 7.1], Theorem 3 explicitly describes the behavior of the $L_{\mathbf{q}}^2$ -(co)homology groups for all \mathbf{q} : they are always concentrated in a single dimension. Let \mathcal{R} denote the region of convergence of the growth series of the corresponding Coxeter group.

Corollary 4. *Suppose that the nerve L of a Coxeter group is a triangulation of S^2 not dual to a hyperbolic 3-simplex.*

- *If $\mathbf{q} \in \bar{\mathcal{R}}$, then $L_{\mathbf{q}}^2 H_*(\Sigma_L)$ is concentrated in dimension 0.*
- *If $\mathbf{q} \notin \mathcal{R}$ and $\mathbf{q} \leq \mathbf{1}$, then $L_{\mathbf{q}}^2 H_*(\Sigma_L)$ is concentrated in dimension 1.*
- *If $\mathbf{q} \notin \mathcal{R}^{-1}$ and $\mathbf{q} \geq \mathbf{1}$, then $L_{\mathbf{q}}^2 H_*(\Sigma_L)$ is concentrated in dimension 2.*
- *If $\mathbf{q} \in \bar{\mathcal{R}}^{-1}$, then $L_{\mathbf{q}}^2 H_*(\Sigma_L)$ is concentrated in dimension 3.*

In either case one can use [7, Corollary 3.4], along with a standard computation for growth series [3, Theorem 17.1.9], to explicitly compute each $L_{\mathbf{q}}^2$ -Betti number.

The Coxeter groups whose nerves are dual to hyperbolic 3-simplices are sometimes in literature called Lánner groups, and there is only nine Lánner groups in dimension three (see [8, Table 6.9]). So there are only nine groups standing in the way of proving Conjecture 2 in full generality for dimension three.

In dimension four we prove the following theorem. First, recall that a subcomplex A of a simplicial complex L is *full* if the vertices of a simplex of L lie in A , then the simplex must lie in A .

Theorem 5. *Suppose that the nerve L of a Coxeter group is a triangulation of S^3 . Furthermore, suppose that there exists a vertex of L such that its link is a full subcomplex of L and not dual to a hyperbolic 3-simplex. Then*

$$L_{\mathbf{q}}^2 H_k(\Sigma_L) = 0 \text{ for } k > 2 \text{ and } \mathbf{q} \leq \mathbf{1}.$$

We obtain the following corollary.

Corollary 6. *Suppose that the nerve L of a Coxeter group is a flag triangulation of S^3 . Then*

$$L_{\mathbf{q}}^2 H_k(\Sigma_L) = 0 \text{ for } k > 2 \text{ and } \mathbf{q} \leq \mathbf{1}.$$

Proof. Since L is flag, it follows that the link of every vertex is a full subcomplex of L . Furthermore, the link of every vertex is not the boundary of a 3-simplex (and in particular, not dual to a 3-simplex). Theorem 5 now completes the proof. ■

We then conclude the article by proving the following generalization of the above corollary.

Theorem 7. *Suppose that the nerve L of a Coxeter group is a flag triangulation of a 3-manifold. Then*

$$L_{\mathbf{q}}^2 H_k(\Sigma_L) = 0 \text{ for } k > 2 \text{ and } \mathbf{q} \leq \mathbf{1}.$$

Consider the case where the nerve L is a triangulation of the n -disk (in this case Σ_L is a manifold with boundary). The following theorem will serve as one of the main ingredients in our proofs.

Theorem 8 ([10, Theorem 5.16]). *Suppose that the nerve L of a Coxeter group is a triangulation of the n -disk, where $n = 2, 3$. If $\mathbf{q} \leq \mathbf{1}$ and $k > \frac{n}{2}$, then $L_{\mathbf{q}}^2 b_k(\Sigma_L) = 0$.*

A special case of Andreev's theorem. For the purpose of this article, we use the notation $\Sigma_L = X$ whenever Σ_L admits a W_L -invariant metric making it isometric to X . The following theorem is now a special case of Andreev's theorem [1, Theorem 2].

Theorem 9. *Suppose that the nerve L is a triangulation of S^2 , but not the boundary of a 3-simplex, and let (W, S) be the corresponding Coxeter system. Furthermore, suppose that*

- *For every $T \subset S$, W_T is not a Euclidean reflection group.*
- *$W \neq W_T \times D_\infty$, where $T \subset S$ spans empty triangle in L and D_∞ is the infinite dihedral group.*

Then $\Sigma_L = \mathbb{H}^3$.

Proof. The nerve L of a Coxeter system (W, S) has a natural piecewise spherical structure, and under this structure, if $s, t \in S$ are connected by an edge in L , then the edge has length $\pi - \pi/m_{st}$, where $(st)^{m_{st}} = 1$. Hence L inherits the structure of a *metric flag complex* [3, Lemma 12.3.1], meaning that any collection of pairwise connected edges of L spans a simplex if and only if there exists a spherical simplex with the corresponding edge lengths.

Suppose now that L is a triangulation of S^2 . Let C be an empty circuit in L and suppose that C is not the boundary of two adjacent triangles. We say that C is a *Euclidean circuit* if the corresponding Coxeter group W_C is a Euclidean reflection group. It follows from L being a metric flag complex that a Euclidean circuit is always a full subcomplex of L , and in particular, W_C is a special subgroup of W . Conditions (i) and (ii) can now be translated from the conditions given in Andreev's Theorem (for details on how to derive them, see for example [14, Section 3]). ■

Equidistant hypersurfaces. Suppose that the Coxeter group W has nerve L that is a triangulation of S^2 and that $\Sigma_L = \mathbb{H}^3$. Let D denote the Davis chamber (in \mathbb{H}^3) and let W_M be a special subgroup of W . We now consider the (possibly infinite) convex polytope $W_M D$ in \mathbb{H}^3 .

For $t > 0$, let S_t denote the t -distant surface from a component S of $\partial W_M D$. Then S_t is a smooth surface (see [2, Proposition II.2.2.1]). In fact, S_t is a union of pieces of which there are three types: hyperbolic, Euclidean, and spherical, each of which are the equidistant pieces from faces, edges, and vertices of S , respectively. The Euclidean pieces look like rectangles that are each adjacent to two hyperbolic pieces and two spherical pieces, and the spherical pieces are adjacent to Euclidean pieces.

As $W_M D$ is convex, the nearest point projection $p : \mathbb{H}^3 \cup \partial\mathbb{H}^3 \rightarrow W_M D$ is defined. If we fix $t > r > 0$, then p induces a map $p_{tr} : S_t \rightarrow S_r$.

Lemma 10. *The map $p_{tr} : S_t \rightarrow S_r$ induced by nearest point projection is $\frac{\tanh(t)}{\tanh(r)}$ -quasiconformal.*

Proof. It suffices to check what p_{tr} does on each of the three types of pieces. First, note that a face of S is simply the intersection of $\partial W_M D$ with a hyperbolic plane in \mathbb{H}^3 . Thus p_{tr} simply scales the corresponding hyperbolic pieces on S_t and S_r by a constant factor. Hence p_{tr} is conformal there. Similarly, the map p_{tr} is conformal on the spherical pieces.

Second, we consider the Euclidean piece in S_t equidistant from an edge of S . A Euclidean piece looks like a rectangle adjacent to two hyperbolic pieces at two parallel edges (parallel in the intrinsic Euclidean geometry), and the the map induced by nearest point projection $S_t \rightarrow S$ scales by a factor of $1/\cosh(t)$ in the direction of those edges. The other two edges of the Euclidean piece are each adjacent to a spherical piece. An edge like this is the arc of a circle with radius t centered at a vertex in S . Thus the edge has length $\theta \sinh(t)$, where θ is the dihedral angle at the corresponding edge of S . Hence the map p_{tr} scales by a factor of $\cosh(r)/\cosh(t)$ in the direction of the edges adjacent to the hyperbolic pieces, and scales the edges adjacent to the spherical pieces by a factor of $\sinh(r)/\sinh(t)$. Therefore p_{tr} is $\frac{\tanh(t)}{\tanh(r)}$ -quasiconformal on the Euclidean pieces. ■

Proof of Theorem 3. Suppose that M is a complete smooth Riemannian manifold. Given a nonnegative measurable function $f : M \rightarrow [0, \infty)$, we define a new norm on the C^∞ k -forms called the L_f^2 norm by

$$\|\omega\|_f^2 = \int_M \|\omega\|_p^2 f(p) dV,$$

where $\|\omega\|_p^2$ is the pointwise norm and dV is the volume form of M . Let $L_f^2 \mathcal{C}^*(M)$ denote the weighted L^2 de Rham complex defined using the L_f^2 norm.

Lemma 11. *Let M and N be smooth surfaces and suppose that $\phi : M \rightarrow N$ is a K -quasiconformal diffeomorphism. Let $g : N \rightarrow [0, \infty)$ be the function defined by $g(p) = f(\phi^{-1}(p))$. Then for every $\omega \in L_g^2 \mathcal{C}^1(N)$, we have that*

$$\frac{1}{K} \|\omega\|_g^2 \leq \|\phi^*(\omega)\|_f^2 \leq K \|\omega\|_g^2.$$

Proof. The pointwise norm of a 1-form is $\|\omega\|_p = \sup_{\substack{x \in T_p M \\ \|x\|=1}} \omega(x)$, where $T_p M$ is the tangent space of M at p . Since ϕ is K -quasiconformal, its differential $d\phi$ maps the circle $\{x \in T_p M \mid \|x\| = 1\}$ to an ellipse with semi-axis $b(p) \leq a(p)$ satisfying $\frac{a(p)}{b(p)} \leq K$. Thus for any $\omega \in L_g^2 \mathcal{C}^1(N)$,

$$b(p) \|\omega\|_{\phi(p)} \leq \|\phi^*(\omega)\|_p \leq a(p) \|\omega\|_{\phi(p)}.$$

Now, let dV_M and dV_N be the respective volume forms of M and N . We have that

$$(f dV_M)_p = \frac{(g(\phi) \phi^*(dV_N))_p}{a(p)b(p)},$$

so for L_f^2 norms we have

$$\begin{aligned}
 \|\phi^*(\omega)\|_f^2 &= \int_M \|\phi^*(\omega)\|_p^2 f(p) dV_M \\
 &\leq \int_M \frac{a(p)}{b(p)} \|\omega\|_{\phi(p)}^2 g(\phi(p)) \phi^*(dV_N) \\
 &\leq K \int_M \|\omega\|_{\phi(p)}^2 g(\phi(p)) \phi^*(dV_N) \\
 &= K \int_N \|\omega\|_x^2 g(x) dV_N = K \|\omega\|_g^2
 \end{aligned}$$

The remaining inequality follows similarly. \blacksquare

Suppose that the nerve L of W is a triangulation of S^2 and that $\Sigma_L = \mathbb{H}^3$. Define f to be the function $f(p) = q_w$, where $w \in W_L$ is a word of shortest length such that $p \in wD$ (here D is the Davis chamber). Let $L_{\mathbf{q}}^2 \mathcal{H}^*(\mathbb{H}^3)$ denote the weighted L^2 de Rham cohomology defined using this f .

Let W_M be an infinite special subgroup of W and let S be one of the components of $\partial W_M D$. Put coordinates (x, t) on \mathbb{H}^3 so that $t \in \mathbb{R}$ is the oriented distance from $p \in \mathbb{H}^3$ to the closest point $x \in S$. Fix $r > 0$, and for $t \geq r$ let S_t denote the hypersurface consisting of points of (oriented) distance t from S . Let $p_{tr} : S_t \rightarrow S_r$ be the map induced by nearest point projection, and let ϕ_{tr} denote the inverse of p_{tr} . By Lemma 10, p_{tr} is $K(t)$ -quasiconformal, with $K(t) = \frac{\tanh(t)}{\tanh(r)}$, and hence so is its inverse $\phi_{tr} : S_r \rightarrow S_t$. Let $i_r : S_r \rightarrow \mathbb{H}^3$ and $i_t : S_t \rightarrow \mathbb{H}^3$ be the inclusions. Then i_r and $i_t \circ \phi_{tr}$ are properly homotopic.

We now adapt the argument after [5, Theorem 16.10] to prove the following lemma.

Lemma 12. *If $\mathbf{q} \geq 1$, then the map $i_r^* : L_{\mathbf{q}}^2 \mathcal{H}^1(\mathbb{H}^3) \rightarrow L_{\mathbf{q}}^2 \mathcal{H}^1(S_r)$ induced by the inclusion i_r is the zero map.*

Proof. Set $g(x, y) = f(x, 0)$, so $f(x, y) \geq g(x, y)$, and let ω be a closed L_f^2 1-form on \mathbb{H}^3 . We now show that the restriction $i_r^*(\omega)$ to S_r represents the zero class in reduced L_f^2 -cohomology. For the remainder of the proof, we will use the notation $\|[\alpha]\|_g$ and $\|\alpha\|_x$ to denote the respective L_g^2 norm and pointwise norm of the harmonic representative of the cohomology class $[\alpha]$.

Suppose for a contradiction that $[i_r^*(\omega)] \neq 0$. Then $\|i_r^*(\omega)\|_g \geq \|[i_r^*(\omega)]\|_g > 0$. By Lemma 11, it follows that $\|\phi_{tr}^*(i_t^*(\omega))\|_g^2 \leq K(t) \|i_t^*(\omega)\|_g^2$, and since i_r and $i_t \circ \phi_{tr}$ are properly homotopic, $[i_r^*(\omega)] = [\phi_{tr}^*(i_t^*(\omega))]$. Therefore

$$K(t) \|i_t^*(\omega)\|_g^2 \geq \|[i_r^*(\omega)]\|_g^2 > 0.$$

Now, $i_t^*(\omega)$ is just a restriction of ω , so we have the pointwise inequality $\|\omega\|_x \geq \|i_t^*(\omega)\|_x$. Using Fubini's Theorem, we compute

$$\begin{aligned}
 \|\omega\|_g^2 &= \int_{\mathbb{H}^3} \|\omega\|_x^2 g(x, y) dV \geq \int_r^\infty \int_{S_t} \|\omega\|_x^2 g(x, y) dAdt \geq \int_r^\infty \int_{S_t} \|i_t^*(\omega)\|_x^2 g(x, y) dAdt \\
 &= \int_r^\infty \|i_t^*(\omega)\|_g^2 dt \geq \int_r^\infty \frac{\tanh(r)}{\tanh(t)} \|[i_r^*(\omega)]\|_g^2 dt = \infty.
 \end{aligned}$$

Since $\|\omega\|_f \geq \|\omega\|_g$, this contradicts the assumption that the L_f^2 norm of ω is finite. \blacksquare

Suppose that L is the nerve of a Coxeter group W_L and that A is a full subcomplex of L . For the proofs that follow, note that $\dim_{\mathbf{q}} L_{\mathbf{q}}^2 H_k(W_L \Sigma_A) = L_{\mathbf{q}}^2 b_k(\Sigma_A)$ (see [3, pg. 352 (vi)]).

Lemma 13. *Suppose that the nerve L is a triangulation of S^2 and that there exists a full subcomplex 1-sphere M of L that separates L into two full 2-disks L_1 and L_2 with boundary M . Furthermore, suppose that one of the following holds:*

- (i) $\Sigma_M = \mathbb{R}^2$.
- (ii) $\Sigma_L = \mathbb{H}^3$.

Then

$$L_{\mathbf{q}}^2 H_k(\Sigma_L) \text{ for } k \geq 2 \text{ and } \mathbf{q} \leq 1.$$

Proof. Since Σ_L is a 3-manifold, it follows that $L_{\mathbf{q}}^2 b_3(\Sigma_L) = 0$ [3, Proposition 20.4.1]. Hence we must show that $L_{\mathbf{q}}^2 b_2(\Sigma_L) = 0$. Consider the following Mayer–Vietoris sequence applied to $L = L_1 \cup_M L_2$:

$$\cdots \rightarrow L_{\mathbf{q}}^2 H_2(W_L \Sigma_{L_1}) \oplus L_{\mathbf{q}}^2 H_2(W_L \Sigma_{L_2}) \rightarrow L_{\mathbf{q}}^2 H_2(\Sigma_L) \rightarrow L_{\mathbf{q}}^2 H_1(W_L \Sigma_M) \rightarrow \cdots$$

By Theorem 8, we have that $L_{\mathbf{q}}^2 H_2(W_L \Sigma_{L_1}) = L_{\mathbf{q}}^2 H_2(W_L \Sigma_{L_2}) = 0$. If (i) holds, then [5, Corollary 14.5] implies that $L_{\mathbf{q}}^2 H_1(\Sigma_M) = 0$, and we are done. If (ii) holds, we argue that the connecting homomorphism $\partial_* : L_{\mathbf{q}}^2 H_2(\Sigma_L) \rightarrow L_{\mathbf{q}}^2 H_1(W_L \Sigma_M)$ is the zero map. By [5, Lemma 16.2], we reduce the proof to showing that the map induced by inclusion $i_* : L_{\mathbf{q}^{-1}}^2 H_1(W_L \Sigma_M) \rightarrow L_{\mathbf{q}^{-1}}^2 H_1(\Sigma_L)$ is the zero map, and since $W_L \Sigma_M$ is a disjoint union of copies of Σ_M , it is enough to show that the restriction of i_* to one summand $L_{\mathbf{q}^{-1}}^2 H_1(\Sigma_M)$ is zero.

Consider the infinite convex polytope $W_M D$, where D is the Davis chamber for W . We have that W_M acts properly and cocompactly on $W_M D$ by isometries. In particular, if S is one of the components of $\partial W_M D$, then W_M acts properly and cocompactly on S , and therefore $L_{\mathbf{q}^{-1}}^2 H^*(\Sigma_M) \cong L_{\mathbf{q}^{-1}}^2 \mathcal{H}^*(S)$. Hence we are done if we show that map $i^* : L_{\mathbf{q}^{-1}}^2 \mathcal{H}^1(\mathbb{H}^3) \rightarrow L_{\mathbf{q}^{-1}}^2 \mathcal{H}^1(S)$ induced by the inclusion $i : S \rightarrow \mathbb{H}^3$ is the zero map.

Fix $r > 0$, and let S_r be the r -distance surface from S . S_r and S are properly homotopy equivalent, and this equivalence induces a weak isomorphism between $L_{\mathbf{q}^{-1}}^2 \mathcal{H}^*(S)$ and $L_{\mathbf{q}^{-1}}^2 \mathcal{H}^*(S_r)$. Thus we have reduced the proof to showing that the map $i_r^* : L_{\mathbf{q}^{-1}}^2 \mathcal{H}^1(\mathbb{H}^3) \rightarrow L_{\mathbf{q}^{-1}}^2 \mathcal{H}^1(S_r)$ induced by the inclusion $i_r : S_r \rightarrow \mathbb{H}^3$ is the zero map, and therefore we are done by Lemma 12. \blacksquare

Remark 14. In [5, Section 16] W is strictly assumed to be right-angled, but the proof of [5, Lemma 16.2] does not use this, as it only uses properties of weighted L^2 -(co)homology.

Proof of Theorem 3. We first suppose that $\Sigma_L = \mathbb{H}^3$. We need to find a full subcomplex M of L satisfying the hypothesis of Lemma 13. First we suppose that L is a flag complex. Let v be a vertex of L and set $M = Lk(v)$. Since L is flag, M is a full subcomplex of L , and since L is a triangulation of the 2-sphere, it follows that M is a 1-sphere, and we are done. Now suppose that L is not flag. Since L is not the boundary of a 3-simplex, there exists an empty 2-simplex in L . Let M denote this empty 2-simplex. Then M separates L into two full 2-disks, both with

boundary M , and we are done. We now suppose that $\Sigma_L \neq \mathbb{H}^3$ and use Theorem 9 to perform a case-by-case analysis.

Case I: W contains a Euclidean special subgroup W_T . Let M be the full subcomplex of L corresponding to W_T . Then M separates L into two 2-disks both with boundary M and hence Lemma 13 (i) implies the assertion.

Case II: $W = W_T \times D_\infty$, where $T \subset S$ spans empty triangle in L . Either $\Sigma_L = \mathbb{R}^3$ or $\Sigma_L = \mathbb{H}^2 \times \mathbb{R}$. In both cases we are done by the weighted Künneth formula.

Case III: L is the boundary of a 3-simplex. By assumption, L is not dual to a hyperbolic simplex, so $\Sigma_L = \mathbb{R}^3$. Therefore we are done by [5, Corollary 14.5]. ■

Proof of Theorem 5. In this case, Σ_L is a 4-manifold, and hence $L_{\mathbf{q}}^2 b_4(\Sigma_L) = 0$ [3, Proposition 20.4.1]. It remains to show that $L_{\mathbf{q}}^2 b_3(\Sigma_L) = 0$. Suppose that the nerve L is a triangulation of S^3 and let $s \in L$ be a vertex. We make the following observations:

- The nerve L_{S-s} of the Coxeter system $(W_{S-s}, S-s)$ is a 3-disk.
- The nerve $St(s)$ of the Coxeter group $W_{St(s)}$ is a 3-disk.
- The nerve $Lk(s)$ of the Coxeter group $W_{Lk(s)}$ is a 2-sphere.

This is because the subcomplexes $St(s)$, $Lk(s)$, and L_{S-s} of L correspond to the closed star of the vertex s , link of the vertex s , and complement of the open star of s , respectively, which are all by assumption full subcomplexes of L .

Consider the following Mayer–Vietoris sequence:

$$\cdots \rightarrow L_{\mathbf{q}}^2 H_3(W_L \Sigma_{L_{S-s}}) \oplus L_{\mathbf{q}}^2 H_3(W_L \Sigma_{St(s)}) \rightarrow L_{\mathbf{q}}^2 H_3(\Sigma_L) \rightarrow L_{\mathbf{q}}^2 H_2(W_L \Sigma_{Lk(s)}) \rightarrow \cdots$$

By Theorem 8, $L_{\mathbf{q}}^2 b_3(\Sigma_{St(s)}) = 0$ and $L_{\mathbf{q}}^2 b_3(\Sigma_{L_{S-s}}) = 0$, and by Theorem 3, $L_{\mathbf{q}}^2 b_2(\Sigma_{Lk(s)}) = 0$. Therefore by the above sequence, $L_{\mathbf{q}}^2 b_3(\Sigma_L) = 0$.

Proof of Theorem 7. Given a Coxeter system (W, S) we let Δ denote the standard $(|S| - 1)$ -simplex with codimension-one faces indexed by S . Thus every $T \subset S$ corresponds to a codimension- $|T|$ face of Δ , denoted by Δ_T , with corresponding barycenter v_T . The Davis chamber D is the subcomplex of the barycentric subdivision of Δ spanned by the barycenters v_T with $T \in \mathcal{S}$, where \mathcal{S} is the poset of subsets $T \subset S$ such that W_T is finite. Set $D_T = D \cap \Delta_T$.

For each $T \in \mathcal{S}$, let c_T denote the union of simplices $c \subset \Sigma_L$ such that $c \cap D_T = v_T$. The boundary of c_T is then cellulated by wc_U , where $w \in W_T$ and $U \subset T$. With its simplicial structure, the boundary ∂c_T is the Coxeter complex corresponding to the Coxeter system (W_T, T) , which is a sphere since W_T is finite. It follows that c_T and its translates are disks, which are called *Coxeter cells of type T* . We denote Σ_L with this decomposition into Coxeter cells by Σ_{cc} .

Now, for for $U \subset S$, set $\mathcal{S}(U) := \{T \in \mathcal{S} \mid T \subset U\}$. Define $\Sigma(U)$ to be the subcomplex of Σ_{cc} consisting of all (closed) Coxeter cells of type T with $T \in \mathcal{S}(U)$. Given $T \in \mathcal{S}(U)$, we define the following subcomplexes of $\Sigma(U)$:

$$\begin{aligned} \Omega_{UT} &: \text{the union of closed cells of type } T', \text{ with } T' \in \mathcal{S}(U)_{\geq T}, \\ \partial\Omega_{UT} &: \text{the cells of } \Omega(U, T) \text{ of type } T'', \text{ with } T'' \notin \mathcal{S}(U)_{\geq T}. \end{aligned}$$

The pair $(\Omega(U, T), \partial\Omega(U, T))$ is the (U, T) -ruin. For brevity, we sometimes write $(\Omega(U, T), \partial)$ to denote the (U, T) -ruin. Note that if $T = \emptyset$, then $\Omega(U, T) = \Sigma(U)$ and $\partial\Omega(U, T) = \emptyset$.

For $s \in T$, set $U' = U - s$ and $T' = T - s$. As in [5, Proof of Theorem 8.3], we have the following weak exact sequence:

$$\cdots \rightarrow L_{\mathbf{q}}^2 H_*(\Omega(U', T'), \partial) \rightarrow L_{\mathbf{q}}^2 H_*(\Omega(U, T'), \partial) \rightarrow L_{\mathbf{q}}^2 H_*(\Omega(U, T), \partial) \rightarrow \cdots$$

For the special case when $U = S$ and $T = \{s\}$ the above sequence becomes:

$$\cdots \rightarrow L_{\mathbf{q}}^2 H_*(\Sigma(S - s)) \rightarrow L_{\mathbf{q}}^2 H_*(\Sigma(S)) \rightarrow L_{\mathbf{q}}^2 H_*(\Omega(S, T), \partial) \rightarrow \cdots$$

Lemma 15. *Suppose that L is a flag triangulation of a 3-manifold. Then for every $t \in L$, $L_{\mathbf{q}}^2 H_*(\Omega(S, t), \partial\Omega(S, t)) = 0$ for $* > 2$ and $\mathbf{q} \leq \mathbf{1}$.*

Proof. First, for $t \in L$, note that the (S, t) -ruin has the property that

$$\Omega(S, t) = \Omega(St(t), t),$$

where $St(t) = \{s \in S \mid m_{st} < \infty\}$. Set $Lk(t) = St(t) - t$, and so we have the following weak exact sequence:

$$\cdots \rightarrow L_{\mathbf{q}}^2 H_*(\Sigma(Lk(t))) \rightarrow L_{\mathbf{q}}^2 H_*(\Sigma(St(t))) \rightarrow L_{\mathbf{q}}^2 H_*(\Omega(S, t), \partial\Omega(S, t)) \rightarrow \cdots$$

Note that

$$L_{\mathbf{q}}^2 b_*(\Sigma(St(t))) = L_{\mathbf{q}}^2 b_*(\Sigma_{St(t)}) \text{ and } L_{\mathbf{q}}^2 b_*(\Sigma(Lk(t))) = L_{\mathbf{q}}^2 b_*(\Sigma_{Lk(t)}),$$

where $\Sigma_{St(t)}$ and $\Sigma_{Lk(t)}$ are the Davis complexes corresponding to the subgroups $W_{St(t)}$ and $W_{Lk(t)}$, respectively. Since L is flag, the respective nerves of the groups $W_{St(t)}$ and $W_{Lk(t)}$ are a 3-disk and a 2-sphere. Furthermore, the nerve of $W_{Lk(t)}$ is not the boundary of a 3-simplex (again, L is flag). By Theorem 8, $L_{\mathbf{q}}^2 b_k(\Sigma_{St(t)}) = 0$ for $k > 2$, and by Theorem 3, $L_{\mathbf{q}}^2 b_k(\Sigma_{Lk(t)}) = 0$ for $k > 1$. Therefore weak exactness of the sequence implies that $L_{\mathbf{q}}^2 H_*(\Omega(S, t), \partial) = 0$ for $* > 2$. ■

Lemma 16 (Compare [13, Lemma 4.1]). *For every $T \in \mathcal{S}^{(2)}$ and $U \subset S$ with $T \subset U$, we have $L_{\mathbf{q}}^2 H_4(\Omega(U, T), \partial\Omega(U, T)) = 0$ for $\mathbf{q} \leq \mathbf{1}$.*

Proof. The proof of [13, Lemma 4.1] goes through to show that $L_{\mathbf{1}}^2 H_4(\Omega(U, T), \partial) = 0$, the main point being that L is a flag triangulation of a 3-manifold, and so it follows that Σ_{cc} is a 4-pseudomanifold (i.e. every 3-cell of Σ_L is contained in precisely two 4-cells). The argument in [10, Lemma 4.8] now completes the proof. ■

Proof of Theorem 7. With the above lemmas, we now follow [13, Proof of The Main Theorem] line by line. For every $U \subset S$ and $t \in U$, we have the following weak exact sequence:

$$\cdots \rightarrow L_{\mathbf{q}}^2 H_*(\Sigma(U - t)) \rightarrow L_{\mathbf{q}}^2 H_*(\Sigma(U)) \rightarrow L_{\mathbf{q}}^2 H_*(\Omega(U, t), \partial) \rightarrow \cdots$$

By Lemma 16 and [13, Proposition 4.2], $L_{\mathbf{q}}^2 H_*(\Omega(U, t), \partial) = 0$ for $* > 2$, and hence by exactness,

$$L_{\mathbf{q}}^2 H_*(\Sigma(U - t)) \cong L_{\mathbf{q}}^2 H_*(\Sigma(U)) \text{ for } * > 2.$$

It follows that $L_{\mathbf{q}}^2 H_*(\Sigma(S)) \cong L_{\mathbf{q}}^2 H_*(\Sigma(\emptyset))$ for $* > 2$, and hence the theorem. ■

ACKNOWLEDGEMENTS

The author would like to thank his Ph.D. advisor Boris Okun for his invaluable guidance and discussions.

REFERENCES

- [1] E. M. Andreev, *On convex polyhedra of finite volume in Lobačcevskii space*, Math. USSR Sbornik, 12(2):255-259, 1970.
- [2] R. D. Canary, A. Marden, D. B. A. Epstein, *Fundamentals of Hyperbolic Manifolds: Selected Expositions*, London Mathematical Society Lecture Note Series 328, 2006.
- [3] M.W. Davis, *The Geometry and Topology of Coxeter Groups*, Princeton University Press, Princeton, 2007.
- [4] M.W. Davis, *Groups generated by reflections and aspherical manifolds not covered by Euclidean space*, Ann. of Math. 117 (1983) 293-324.
- [5] M.W. Davis, J. Dymara, T. Januszkiewicz, B. Okun, *Weighted L^2 -cohomology of Coxeter groups*, Geometry & Topology 11 (2007), 47-138.
- [6] M.W. Davis and B. Okun, *Vanishing theorems and conjectures for the ℓ^2 -homology of right-angled Coxeter groups*, Geometry & Topology, 5:7-74, 2001.
- [7] J. Dymara, *Thin buildings*, Geometry & Topology, 10:667-694, 2006.
- [8] J. Humphreys, *Reflection Groups and Coxeter Groups*, Cambridge University Press, Cambridge, 1990.
- [9] J. Lott and W. Lück, *L^2 -topological invariants of 3-manifolds*, Invent. Math. 120 (1995), 15-60.
- [10] W.J. Mogilski, *The fattened Davis complex and weighted L^2 -(co)homology of Coxeter groups*, arXiv:1502.07783 [math.AT] (2015).
- [11] B. Okun and K. Schreve, *The L^2 -(co)homology of groups with hierarchies*, arXiv:1407.1340 [math.GT] (2014).
- [12] G. Perelman, *Finite extinction time for the solutions to the Ricci flow on certain three-manifolds*, arXiv:math.DG/0307245 (2003).
- [13] T.A. Schroeder, *The ℓ^2 -homology of even Coxeter groups*, Algebraic & Geometric Topology, 9(2):1089-1104, 2009. DOI number: 10.2140/agt.2009.9.1089.
- [14] T.A. Schroeder, *On the three-dimensional Singer Conjecture for Coxeter groups*, arXiv:0909.0071 [math.GT] (2009).

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF WISCONSIN-MILWAUKEE, MILWAUKEE, WI 53211, USA

E-mail address: mogilski@uwm.edu