

Radiative rates for E1, E2, M1, and M2 transitions in the Br-like ions Sr IV, Y V, Zr VI, Nb VII, and Mo VIII

Kanti M. Aggarwal^{a,*}, Francis P. Keenan^a

^a*Astrophysics Research Centre, School of Mathematics and Physics, Queen's University Belfast, Belfast BT7 1NN, Northern Ireland, UK*

Abstract

Energies and lifetimes are reported for the lowest 375 levels of five Br-like ions, namely Sr IV, Y V, Zr VI, Nb VII, and Mo VIII, mostly belonging to the $4s^24p^5$, $4s^24p^44\ell$, $4s4p^6$, $4s^24p^45\ell$, $4s^24p^34d^2$, $4s4p^54\ell$, and $4s4p^55\ell$ configurations. Extensive configuration interaction has been included and the general-purpose relativistic atomic structure package (GRASP) has been adopted for the calculations. Additionally, radiative rates are listed among these levels for all E1, E2, M1, and M2 transitions. From a comparison with the measurements, the majority of our energy levels are assessed to be accurate to better than 2%, although discrepancies between theory and experiment for a few are up to 6%. An accuracy assessment of the calculated radiative rates (and lifetimes) is more difficult, because no prior results exist for these ions.

arXiv:1504.00335v1 [physics.atom-ph] 1 Apr 2015

*Corresponding author.

Email address: K.Aggarwal@qub.ac.uk (Kanti M. Aggarwal)

Contents

1. Introduction	3
2. Energy levels	3
3. Radiative rates	5
4. Lifetimes	6
5. Conclusions	6
Appendix A. Supplementary data	6
References	7
Explanation of Tables	15
Tables	
1. Energies (Ryd) for the lowest 375 levels of Sr IV and their lifetimes (τ , s). $a\pm b \equiv a \times 10^{\pm b}$. See page 15 for Explanation of Tables	18
2. Energies (Ryd) for the lowest 375 levels of Y V and their lifetimes (τ , s). $a\pm b \equiv a \times 10^{\pm b}$. See page 15 for Explanation of Tables	19
3. Energies (Ryd) for the lowest 375 levels of Zr VI and their lifetimes (τ , s). $a\pm b \equiv a \times 10^{\pm b}$. See page 15 for Explanation of Tables	20
4. Energies (Ryd) for the lowest 375 levels of Nb VII and their lifetimes (τ , s). $a\pm b \equiv a \times 10^{\pm b}$. See page 15 for Explanation of Tables	21
5. Energies (Ryd) for the lowest 375 levels of Mo VIII and their lifetimes (τ , s). $a\pm b \equiv a \times 10^{\pm b}$. See page 15 for Explanation of Tables	22
6. Transition wavelengths (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij} , dimensionless), and line strengths (S, in atomic units) for electric dipole (E1), and A_{ji} for E2, M1, and M2 transitions in Sr IV. The last column gives R(E1), the ratio of velocity and length forms of A- values for E1 transitions. $a\pm b \equiv a \times 10^{\pm b}$. See page 15 for Explanation of Tables	23
7. Transition wavelengths (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij} , dimensionless), and line strengths (S, in atomic units) for electric dipole (E1), and A_{ji} for E2, M1, and M2 transitions in Y V. The last column gives R(E1), the ratio of velocity and length forms of A- values for E1 transitions. $a\pm b \equiv a \times 10^{\pm b}$. See page 16 for Explanation of Tables	23
8. Transition wavelengths (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij} , dimensionless), and line strengths (S, in atomic units) for electric dipole (E1), and A_{ji} for E2, M1, and M2 transitions in Zr VI. The last column gives R(E1), the ratio of velocity and length forms of A- values for E1 transitions. $a\pm b \equiv a \times 10^{\pm b}$. See page 16 for Explanation of Tables	23
9. Transition wavelengths (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij} , dimensionless), and line strengths (S, in atomic units) for electric dipole (E1), and A_{ji} for E2, M1, and M2 transitions in Nb VII. The last column gives R(E1), the ratio of velocity and length forms of A- values for E1 transitions. $a\pm b \equiv a \times 10^{\pm b}$. See page 16 for Explanation of Tables	24
10. Transition wavelengths (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij} , dimensionless), and line strengths (S, in atomic units) for electric dipole (E1), and A_{ji} for E2, M1, and M2 transitions in Mo VIII. The last column gives R(E1), the ratio of velocity and length forms of A- values for E1 transitions. $a\pm b \equiv a \times 10^{\pm b}$. See page 17 for Explanation of Tables	24

1. Introduction

Atomic data for energy levels and radiative decay rates (A-values) are required for the modelling of plasmas, such as astrophysical and fusion. Generally, comparatively lighter ions ($Z \leq 30$) are of relevance to both astrophysical and fusion plasmas, but the heavier ones are mostly useful for the latter, mainly for assessing and controlling the radiation loss. With the ongoing ITER project, the need for atomic data for a wide range of ions has become greater. Therefore, in a recent paper, we [1] have reported atomic data for an important Br-like tungsten ion, W XL. In this paper we provide similar results for five additional Br-like ions with $38 \leq Z \leq 42$, i.e. Sr IV, Y V, Zr VI, Nb VII, and Mo VIII.

Laboratory measurements for energy levels are available for all five Br-like ions of interest, but are limited to only a few levels, except for Sr IV. These measurements have been compiled by the NIST (National Institute of Standards and Technology) team [2], and are available at their website http://physics.nist.gov/PhysRefData/ASD/levels_form.html. However, similar data for A-values [3] are limited to the magnetic dipole (M1) and electric quadrupole (E2) transitions between the ground state levels ($4s^24p^5\ ^2P_{3/2}^o - ^2P_{1/2}^o$) of several Br-like ions, including those considered here, insufficient for modelling applications. Therefore, in a recent paper Singh et al. [4] have calculated energies and lifetimes for the lowest 31 levels, belonging to the $4s^24p^5$, $4s^24p^44d$ and $4s4p^6$ configurations, of the five Br-like ions listed above. Additionally, they also listed A-values for electric dipole (E1) transitions, but only from the ground state $4s^24p^5\ ^2P_{3/2,1/2}^o$ to higher-lying levels. For the calculations, they adopted the GRASP code [5] but included only limited CI (configuration interaction). However, these 31 levels are not the lowest, because many of the $4s^24p^45s$ configuration intermix with these – see Tables 1–5. More importantly, their reported results cannot be reproduced. Discrepancies with our calculated energy levels, obtained with the same code, configurations and approximations, are only up to 0.15 Ryd, but are much larger (up to 5 orders of magnitude) for A-values and lifetimes. In our earlier paper [6] we demonstrated that these discrepancies are due to the results of [4] being unreliable. However, in [6] the atomic data are restricted to the 31 levels of the $4s^24p^5$, $4s^24p^44d$ and $4s4p^6$ configurations, although calculations were performed for up to 3990 levels. Therefore, in this paper we report complete results among the lowest 375 levels of these ions.

2. Energy levels

For our calculations we have adopted the fully relativistic multi-configuration Dirac-Fock (MCDF) code, originally developed by Grant et al. [5]. The code is based on the jj coupling scheme and includes higher-order relativistic corrections arising from the Breit interaction and QED (quantum electrodynamics) effects, which are important for the heavy ions considered here. However, this initial version of the MCDF code has undergone multiple revisions, and is now better known as GRASP [7]. The version adopted in our work has been revised by one of its authors (Dr. P. H. Norrington), is referred to as GRASP0, and is freely available at <http://web.am.qub.ac.uk/DARC/>. It provides comparable results with other revisions, such as GRASP2K [8, 9].

In our calculations, extensive configuration interaction (CI) has been included among 39 configurations, namely $4s^24p^5$, $4s^24p^44d/4f$, $4s4p^6$, $4p^64d/4f$, $4s4p^54d/4f$, $4s^24p^34f^2/4d^2/4d4f$, $4s^24p^24d^3$, $4s^24p4d^4$, $4s^24p^24d^24f$, $4s4p^34d^3$, $4p^54d^2$, $3d^94s^24p^54d/4f$, $3d^94s^24p^6$, $4s4p^55\ell$, $4p^65\ell$, $4s^24p^45\ell$, and $3d^94s^24p^55\ell$. These configurations generate 3990 levels in total and have been carefully chosen, mainly based on the interacting range of their energies. The specific number of levels each configuration generates and their energy ranges are listed in Table 1 of our earlier paper [6].

As noted earlier for some Ni ions [10] the interacting energy ranges of the configurations significantly influence the accuracy of the calculations. We also point out that these configurations are not the same as included for Br-like W XL [1], although many are common. Furthermore, additional configurations, such as $3p^5 3d^{10} 4s^2 4p^6$, $3p^5 3d^{10} 4s^2 4p^5 4d$ and $3p^5 3d^{10} 4s^2 4p^5 4f$, have also been tested but excluded from the final calculations, because their impact on the lower energy levels is insignificant, mainly because they generate levels lying at much higher energies. Nevertheless, we discuss their impact later in the section. Finally, as in most of our earlier work, we have adopted the option of ‘extended average level’, in which a weighted (proportional to $2j+1$) trace of the Hamiltonian matrix is minimised. This yields results comparable to other options, such as ‘average level’ – see for example, Aggarwal et al. for several ions of Kr [11] and Xe [12]. The ions considered here have nuclear charges between those of Kr and Xe.

In Tables 1–5 we list our lowest 375 energy levels for five Br-like ions with $38 \leq Z \leq 42$. Comparisons with the measurements compiled by NIST have already been discussed in our earlier work [6], but only for 31 levels of the $4s^2 4p^5$, $4s^2 4p^4 4d$ and $4s 4p^6$ configurations. However, NIST energies are also available for a few other higher lying levels, particularly those of $4p^4 5s$ for all ions, and of many more configurations of Sr IV. Therefore, in Table A we compare our calculated energies with the NIST listings for the common levels of Sr IV. For most levels the theoretical energies agree with the NIST values within 2%, but discrepancies for a few are up to 6%, particularly among the lowest 60 levels in Table A.

We would like to stress here that the LSJ labels assigned to the levels in Tables 1–5 are not always unique, because levels from different configurations mix strongly and sometimes the eigenvector of the same level/configuration dominates for two (or even three) levels. This is a common problem for all calculations, particularly when the CI is important as in the present case. However, the ambiguity in the configuration/level designation is only for a few levels in each ion. In Table B we list the mixing coefficients for the lowest 50 levels of Sr IV – see in particular, levels 14, 18, 39, and 50, which are highly mixed. Hence, the LSJ designations provided in Tables 1–5 may sometimes be interchangeable, although the associated J^π values are definitive for all levels and ions.

We now investigate if a larger CI expansion is helpful in improving the accuracy of our calculated energy levels. Adopting the *Flexible Atomic Code* (FAC) of Gu [13], available from the website <http://sprg.ssl.berkeley.edu/~mfgu/fac/>, we have performed a series of calculations with up to 12 137 levels. The additional levels belong to the $4p^6 6\ell$, $4s 4p^5 6\ell$, $4s^2 4p^4 6\ell$, $4p^6 7\ell$, $4s 4p^5 7\ell$, $4s^2 4p^4 7\ell$, $4s^2 4p^3 5\ell^2$, $4s 4p^4 5\ell^2$, $3p^5 3d^{10} 4s^2 4p^6$, $3p^5 3d^{10} 4s^2 4p^5 4d$, and $3p^5 3d^{10} 4s^2 4p^5 4f$ configurations. FAC is also a fully relativistic code and results obtained for energy levels and A-values are (generally) comparable to GRASP, as already shown for several other ions, see for example Aggarwal et al. for Kr [11] and Xe [12] ions. This larger calculation can be more easily performed with this code within a reasonable time frame of a few days, because of its higher efficiency. However, energies obtained from this larger FAC calculation agree with GRASP for the levels of the $4s^2 4p^5$, $4s^2 4p^4 4d$ and $4s 4p^6$ configurations, within 0.01 Ryd for all ions, as shown in our earlier work [6]. In Table C we make a similar comparison for all levels of the $4s^2 4p^5$, $4s 4p^6$, $4s^2 4p^4 4d/4f$, and $4s^2 4p^4 5\ell$ configurations of Sr IV. These levels are not listed in the ascending order of energy, but in the order of the configurations for easier understanding. Furthermore, the energy differences between the two independent calculations are listed in both magnitude and percentage for a ready reference. Although energy differences for most levels are below 1%, discrepancies are up to 3.5% for a few, particularly for those of the $n=4$ configurations for which energies from FAC are invariably higher. Correspondingly energy differences between the NIST compilation and FAC calculation has increased for most levels, and therefore a

larger expansion is not helpful. A closer agreement between theoretical and experimental energies for these levels may perhaps be achieved by the inclusion of an enormously large CI (involving over a million levels or configuration state functions, CSF), as shown by Froese Fischer [14] for W XL, where she performed extremely large calculations for a limited set of levels. Unfortunately, for the greater number of levels involved in the present work and for five ions, such calculations are not possible with our computational resources.

3. Radiative rates

The absorption oscillator strength (f_{ij}) for a transition $i \rightarrow j$ is a dimensionless quantity and is related to the radiative rate A_{ji} (in s^{-1}) by the following expression:

$$f_{ij} = \frac{mc}{8\pi^2 e^2} \lambda_{ji}^2 \frac{\omega_j}{\omega_i} A_{ji} = 1.49 \times 10^{-16} \lambda_{ji}^2 \frac{\omega_j}{\omega_i} A_{ji} \quad (1)$$

where m and e are the electron mass and charge, respectively, c the velocity of light, λ_{ji} the transition wavelength in Å, and ω_i and ω_j the statistical weights of the lower i and upper j levels, respectively. Similarly, the oscillator strength f_{ij} (dimensionless), A-values and the line strength S (in atomic units, 1 a.u. = $6.460 \times 10^{-36} \text{ cm}^2 \text{ esu}^2$) are related by the standard equations given in [1].

In our calculations with the GRASP code the S- (and subsequently A- and f-) values have been determined in both the length and velocity forms, equivalent to the Babushkin and Coulomb gauges in the relativistic nomenclature. In Tables 6–10 we present results for transitions from the lowest three to higher excited levels, with the full tables available online in the electronic version. Included in these tables are the transition (energies) wavelengths (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij} , dimensionless), and line strengths (S in a.u.) for all 16 568 E1 transitions among the lowest 375 levels. The listed wavelengths (and other parameters) are based on the Breit and QED-corrected theoretical energies, given in Tables 1–5, where the *indices* used to represent the lower and upper levels of a transition are also defined. However, for the 29 166 E2, 23 514 M1, and 20 476 M2 transitions, only the A-values are included in Tables 6–10, because the corresponding results for f- or S-values can be easily obtained using Eqs. (1-5) given in [1]. Since the length form is considered to be comparatively more accurate, results listed in Tables 6–10 are in this form alone. However, for accuracy assessment we have also listed the ratio R of the velocity and length forms for all E1 transitions.

For comparison purpose there are no available data in the literature for A-values. However, based on calculations with differing amount of CI, codes, and the ratio of the forms in length and velocity, the accuracy of our A-values has been assessed in our earlier paper [6]. It is estimated to be $\sim 20\%$ for transitions with $f \geq 0.01$. However, that comparison was limited to transitions among the 31 levels of the $4s^2 4p^5$, $4s^2 4p^4 4d$ and $4s 4p^6$ configurations, and hence is not applicable to a wider range of transitions listed in Tables 6–10. For example, there are 2431 transitions of Sr IV in Table 6 for which $f(\text{E1}) \geq 0.01$, and $\sim 59\%$ differ by over 20%. Their number reduces to 20% if R is taken to be within 50% and only 3% transitions differ by over an order of magnitude – see for example: 4–67, 5–66, and 6–65, all of which have $f < 0.1$. For weaker transitions the ratio R may be up to several orders of magnitude.

In Table D we compare our GRASP results for A- and f- values with the corresponding FAC calculations, for transitions from the lowest two to higher excited levels, up to 49, of Sr IV. Unfortunately, differences for a few transitions

are up to a factor of 5. Most such transitions are comparatively weak (such as 1–11, 1–33 and 2–18), and larger differences for these are not surprising, because of differences in the methodology and the CI. However, the 1–32 transition is strong ($f = 0.84$) but the corresponding result from FAC is lower by a factor of three. Inclusion of larger CI generally increases the accuracy of A-values. However, in the present case it may not be true as was discussed earlier in section 2 for energy levels. Finally, there are no discrepancies with the Biémont et al. [3] A-values for the M1 and E2 ($4s^24p^5$) $^2P_{3/2}^o - ^2P_{1/2}^o$ transitions.

4. Lifetimes

The lifetime τ of a level j is determined as follows:

$$\tau_j = \frac{1}{\sum_i A_{ji}}. \quad (2)$$

In Tables 1–5 we list lifetimes for the lowest 375 levels of Sr IV, Y V, Zr VI, Nb VII, and Mo VIII. The calculations of τ include A-values from all types of transitions, i.e. E1, E2, M1, and M2. Unfortunately, there are no measurements or other theoretical results available for comparison purposes.

5. Conclusions

Adopting the GRASP code, energy levels and radiative rates (for E1, E2, M1, and M2 transitions) for the lowest 375 levels of 5 Br-like ions (Sr IV, Y V, Zr VI, Nb VII, and Mo VIII) have been presented. Additionally, lifetimes for these levels are also listed although no other data are available in the literature with which to compare our results. For most energy levels and for all ions, the accuracy of our results is estimated to be better than 2%. This assessment is based on comparisons with the results listed by NIST. However, there is scope for improvement because for a few levels the discrepancies between theory and measurement are up to 6%. Although length and velocity forms of the A-values have been discussed and comparisons with the FAC code with larger CI have been made, it is not possible to assign any accuracy estimate for the radiative data. Future calculations and measurements (particularly for lifetimes) will be helpful in the accuracy assessment of our listed data.

Acknowledgments

KMA is thankful to AWE Aldermaston for financial support.

Appendix A. Supplementary data

Owing to space limitations, only parts of Tables 6–10 are presented here, the full tables being made available as supplemental material in conjunction with the electronic publication of this work. Supplementary data associated with this article can be found, in the online version, at [doi:10.1080/00107179.2015.1058888](https://doi.org/10.1080/00107179.2015.1058888).

References

- [1] K.M. Aggarwal, F.P. Keenan, *At. Data Nucl. Data Tables* 100 (2014) 1399.
- [2] A. Kramida, Y. Ralchenko, J. Reader, and NIST ASD Team, 2015 NIST Atomic Spectra Database (version 5.1)
Online at: <http://physics.nist.gov/asd>
- [3] E. Biémont, R.D. Cowan, J.E. Hansen, *Phys. Scr.* 37 (1988) 850.
- [4] S. Aggarwal, A.K. Singh, M. Mohan, *Phys. Scr.* 88 (2013) 035301.
- [5] I.P. Grant, B.J. McKenzie, P.H. Norrington, D.F. Mayers, N.C. Pyper, *Comput. Phys. Commun.* 21 (1980) 207.
- [6] K.M. Aggarwal, F.P. Keenan, *Phys. Scr.* 89 (2014) 125404.
- [7] K.G. Dyall, I.P. Grant, C.T. Johnson, F.A. Parpia, E.P. Plummer, *Comput. Phys. Commun.* 55 (1989) 425.
- [8] P. Jönsson, X. He, C.F. Fischer, I.P. Grant, *Comput. Phys. Commun.* 177 (2007) 597.
- [9] P. Jönsson P, G. Gaigalas, J. Bieroń, C.F. Fischer, I.P. Grant, *Comput. Phys. Commun.* 184 (2013) 2197.
- [10] K.M. Aggarwal, F.P. Keenan, *At. Data Nucl. Data Tables* 85 (2003) 453.
- [11] K.M. Aggarwal, F.P. Keenan, K.D. Lawson, *At. Data Nucl. Data Tables* 94 (2008) 323.
- [12] K.M. Aggarwal, F.P. Keenan, K.D. Lawson, *At. Data Nucl. Data Tables* 96 (2010) 123.
- [13] M.F. Gu, *Can. J. Phys.* 86 (2008) 675.
- [14] C. Froese Fischer, *Atoms* 2 (2014) 1.

Table A

Comparison of energies (Ryd) for the lowest 221 levels of Sr IV. See Table 1 for the full listing of the levels.

Index	Configuration	Level	NIST	Present	Index	Configuration	Level	NIST	Present
1	4s ² 4p ⁵	2P _{3/2} ^o	0.00000	0.00000	71	4s ² 4p ⁴ 5d	4D _{3/2}	2.98773	3.04428
2	4s ² 4p ⁵	2P _{1/2} ^o	0.08865	0.08515	72	4s ² 4p ⁴ (³ P)4f	4G _{11/2} ^o	3.05620	3.04851
3	4s4p ⁶	2S _{1/2}	1.37149	1.44244	73	4s ² 4p ⁴ (³ P)4f	2F _{7/2} ^o	3.05986	3.05012
4	4s ² 4p ⁴ (³ P)4d	4D _{7/2}	1.70553	1.75452	74	4s ² 4p ⁴ (³ P)4f	2F _{5/2} ^o	3.05919	3.05057
5	4s ² 4p ⁴ (³ P)4d	4D _{5/2}	1.70590	1.75612	75	4s ² 4p ⁴ (³ P)4f	2G _{9/2} ^o	3.05985	3.05418
6	4s ² 4p ⁴ (³ P)4d	4D _{3/2}	1.71340	1.76432	76	4s ² 4p ⁴ 5d	4D _{1/2}	3.02153	3.05480
7	4s ² 4p ⁴ (³ P)4d	4D _{1/2}	1.72338	1.77444	77	4s ² 4p ⁴ (³ P)4f	4D _{3/2} ^o	3.06845	3.05433
8	4s ² 4p ⁴ (³ P)4d	4F _{9/2}	1.79574	1.85784	79	4s ² 4p ⁴ 5d	2F _{7/2}	3.00694	3.07641
9	4s ² 4p ⁴ (³ P)4d	4F _{7/2}	1.82563	1.89004	80	4s ² 4p ⁴ (³ P)4f	4D _{5/2} ^o	3.10718	3.08883
10	4s ² 4p ⁴ (³ P)4d	4F _{5/2}	1.85301	1.91248	81	4s ² 4p ⁴ 5d	4P _{1/2}	3.06490	3.09218
11	4s ² 4p ⁴ (³ P)4d	4F _{3/2}	1.86062	1.92068	82	4s ² 4p ⁴ (³ P)4f	2D _{3/2} ^o	3.07140	3.09237
12	4s ² 4p ⁴ (¹ D)4d	2P _{1/2}	1.82736	1.92312	83	4s ² 4p ⁴ (³ P)4f	4D _{7/2} ^o	3.11125	3.09267
13	4s ² 4p ⁴ (³ P)4d	4P _{1/2}	1.86576	1.95489	84	4s ² 4p ⁴ (³ P)4f	4G _{9/2} ^o	3.10886	3.09969
14	4s ² 4p ⁴ (³ P)4d	4P _{3/2}	1.86636	1.95507	85	4s ² 4p ⁴ 5d	4P _{3/2}	3.07975	3.11020
15	4s ² 4p ⁴ (¹ D)4d	2D _{3/2}	1.88199	1.97045	86	4s ² 4p ⁴ 5d	4F _{5/2}	3.07235	3.11358
16	4s ² 4p ⁴ (³ P)4d	2F _{7/2}	1.89068	1.97942	87	4s ² 4p ⁴ (³ P)4f	4G _{7/2} ^o	3.12808	3.11450
17	4s ² 4p ⁴ (³ P)4d	4P _{5/2}	1.90398	1.98759	88	4s ² 4p ⁴ (³ P)4f	4G _{5/2} ^o	3.13039	3.11686
18	4s ² 4p ⁴ (¹ D)4d	2P _{3/2}	1.90647	1.99565	89	4s ² 4p ⁴ (³ P)4f	2G _{7/2} ^o	3.05986	3.12866
19	4s ² 4p ⁴ (¹ D)4d	2D _{5/2}	1.93164	2.01910	90	4s ² 4p ⁴ 5d	4F _{7/2}	3.07004	3.13151
20	4s ² 4p ⁴ (¹ D)4d	2G _{9/2}	1.96093	2.04505	91	4s ² 4p ⁴ 5d	4F _{3/2}	3.07975	3.13216
21	4s ² 4p ⁴ (¹ D)4d	2G _{7/2}	1.96094	2.04924	92	4s ² 4p ⁴ 5d	4P _{5/2}	3.10783	3.13371
22	4s ² 4p ⁴ (³ P)4d	2F _{5/2}	1.95873	2.05338	93	4s ² 4p ⁴ 5d	2P _{1/2}	3.06490	3.13400
23	4s ² 4p ⁴ 5s	4P _{5/2}	2.08365	2.17395	95	4s ² 4p ⁴ 5d	2F _{5/2}	3.08538	3.15224
24	4s ² 4p ⁴ (¹ D)4d	2F _{5/2}	2.05829	2.18489	96	4s ² 4p ⁴ 5d	2D _{3/2}	3.10940	3.15983
25	4s ² 4p ⁴ (¹ D)4d	2F _{7/2}	2.07858	2.20325	98	4s ² 4p ⁴ (¹ D)4f	2D _{5/2} ^o	3.20007	3.20054
26	4s ² 4p ⁴ 5s	4P _{3/2}	2.11606	2.21122	99	4s4p ⁵ (³ P)4d	4F _{9/2} ^o	3.19353	3.20534
27	4s ² 4p ⁴ 5s	4P _{1/2}	2.15902	2.24728	100	4s4p ⁵ (³ P)4d	4F _{7/2} ^o	3.19595	3.20625
28	4s ² 4p ⁴ 5s	2P _{3/2}	2.17080	2.26566	102	4s ² 4p ⁴ (¹ D)4f	2D _{3/2} ^o	3.21335	3.21271
29	4s ² 4p ⁴ (¹ S)4d	2D _{3/2}	2.20683	2.29294	104	4s4p ⁵ (³ P)4d	4F _{5/2} ^o	3.21161	3.22170
30	4s ² 4p ⁴ 5s	2P _{1/2}	2.20611	2.30524	105	4s ² 4p ⁴ (¹ D)4f	2P _{3/2} ^o	3.20835	3.22357
31	4s ² 4p ⁴ (¹ S)4d	2D _{5/2}	2.23930	2.31329	106	4s ² 4p ⁴ (¹ D)4f	2F _{7/2} ^o	3.22531	3.22933
32	4s ² 4p ⁴ 5s	2D _{5/2}	2.27859	2.39785	107	4s ² 4p ⁴ (¹ D)4f	2H _{9/2} ^o	3.21820	3.23064
33	4s ² 4p ⁴ 5s	2D _{3/2}	2.28276	2.40477	108	4s ² 4p ⁴ (¹ D)4f	2H _{11/2} ^o	3.21852	3.23116
34	4s ² 4p ⁴ (³ P)4d	2P _{3/2}	2.29991	2.41725	109	4s4p ⁵ (³ P)4d	4F _{3/2} ^o	3.22985	3.23979
35	4s ² 4p ⁴ (³ P)4d	2P _{1/2}	2.30762	2.44445	110	4s ² 4p ⁴ (¹ D)4f	2F _{5/2} ^o	3.23839	3.24733
36	4s ² 4p ⁴ (³ P)4d	2D _{5/2}	2.32210	2.45347	140	4s ² 4p ⁴ 5d	2D _{5/2}	3.40200	3.46151
37	4s ² 4p ⁴ 5p	4P _{5/2}	2.43798	2.49996	144	4s ² 4p ⁴ 5f	4F _{9/2} ^o	3.44051	3.47997
38	4s ² 4p ⁴ 5p	4P _{3/2}	2.43791	2.50245	145	4s ² 4p ⁴ 5f	4F _{5/2} ^o	3.44937	3.48051
39	4s ² 4p ⁴ (¹ D)4d	2S _{1/2}	2.34510	2.51590	146	4s ² 4p ⁴ 5f	4G _{11/2} ^o	3.44247	3.48097
40	4s ² 4p ⁴ (³ P)4d	2D _{3/2}	2.40740	2.52376	147	4s ² 4p ⁴ 5f	4F _{7/2} ^o	3.46247	3.48102
41	4s ² 4p ⁴ 5p	4P _{1/2} ^o	2.46361	2.52932	148	4s ² 4p ⁴ 5d	2D _{3/2}	3.40550	3.48141
42	4s ² 4p ⁴ 5p	4D _{7/2} ^o	2.47181	2.54782	150	4s ² 4p ⁴ 5f	2G _{9/2} ^o	3.44023	3.48431
43	4s ² 4p ⁴ 5p	2D _{5/2} ^o	2.47253	2.55249	152	4s ² 4p ⁴ 5f	2F _{5/2} ^o	3.43363	3.48880
44	4s ² 4p ⁴ 5p	2P _{1/2} ^o	2.51560	2.58741	153	4s ² 4p ⁴ 5f	2F _{7/2} ^o	3.44485	3.48903
45	4s ² 4p ⁴ 5p	4D _{3/2} ^o	2.53403	2.59655	155	4s ² 4p ⁴ 5g	4G _{11/2}	3.48916	3.52370
46	4s ² 4p ⁴ 5p	4D _{1/2} ^o	2.53254	2.60789	156	4s ² 4p ⁴ 5g	4G _{9/2}	3.48954	3.52378
47	4s ² 4p ⁴ 5p	2P _{3/2} ^o	2.56562	2.61162	157	4s ² 4p ⁴ 5g	4G _{7/2}	3.48938	3.52395
48	4s ² 4p ⁴ 5p	4D _{5/2} ^o	2.54395	2.61764	158	4s ² 4p ⁴ 5g	2G _{9/2}	3.48922	3.52408
49	4s ² 4p ⁴ 5s	2S _{1/2}	2.50314	2.63469	159	4s ² 4p ⁴ 5g	4G _{5/2}	3.49138	3.52626
50	4s ² 4p ⁴ 5p	2D _{3/2} ^o	2.51655	2.64532	160	4s ² 4p ⁴ 5g	4F _{7/2}	3.49153	3.52643
51	4s ² 4p ⁴ 5p	4S _{3/2} ^o	2.57292	2.65539	161	4s ² 4p ⁴ 5g	4H _{13/2}	3.49268	3.52782
52	4s ² 4p ⁴ 5p	2S _{1/2} ^o	2.57379	2.66029	162	4s ² 4p ⁴ 5g	2H _{11/2}	3.49276	3.52792
53	4s ² 4p ⁴ 5p	2F _{5/2} ^o	2.63042	2.73474	163	4s ² 4p ⁴ 5g	4F _{3/2}	3.49426	3.52971
54	4s ² 4p ⁴ 5p	2F _{7/2} ^o	2.64552	2.74818	164	4s ² 4p ⁴ 5g	2F _{5/2}	3.49432	3.52980
55	4s ² 4p ⁴ 5p	2D _{3/2} ^o	2.66504	2.78784	176	4s ² 4p ⁴ 5g	4G _{5/2}	3.56527	3.59757
56	4s ² 4p ⁴ 5p	2D _{5/2} ^o	2.68932	2.80035	177	4s ² 4p ⁴ 5g	4F _{7/2}	3.56540	3.59770
57	4s ² 4p ⁴ 5p	2P _{3/2} ^o	2.68703	2.82130	178	4s ² 4p ⁴ 5g	4H _{11/2}	3.56600	3.59835
58	4s ² 4p ⁴ 5p	2P _{1/2} ^o	2.71289	2.85686	179	4s ² 4p ⁴ 5g	4H _{9/2}	3.56612	3.59847

Table A (continued)

Index	Configuration	Level	NIST	Present	Index	Configuration	Level	NIST	Present
59	4s ² 4p ⁴ 5p	² P _{3/2} ^o	2.87771	2.94839	180	4s ² 4p ⁴ 5g	⁴ H _{7/2}	3.56953	3.60217
60	4s ² 4p ⁴ 5p	² P _{1/2} ^o	2.86746	2.94916	181	4s ² 4p ⁴ 5g	⁴ F _{9/2}	3.56952	3.60218
61	4s ² 4p ⁴ (³ P)4f	⁴ F _{9/2} ^o	2.99398	2.96793	182	4s ² 4p ⁴ 5g	⁴ F _{9/2}	3.57084	3.60346
62	4s ² 4p ⁴ (³ P)4f	⁴ F _{7/2} ^o	2.99724	2.97174	183	4s ² 4p ⁴ 5g	² F _{7/2}	3.57086	3.60348
63	4s ² 4p ⁴ (³ P)4f	⁴ F _{5/2} ^o	3.00727	2.97953	212	4s ² 4p ⁴ 5g	² I _{13/2}	3.67276	3.73417
64	4s ² 4p ⁴ (³ P)4f	⁴ F _{3/2} ^o	3.01458	2.99056	215	4s ² 4p ⁴ 5g	² F _{7/2}	3.67594	3.73733
65	4s4p ⁵ (³ P)4d	⁴ P _{1/2} ^o	3.04607	2.99545	216	4s ² 4p ⁴ 5g	² F _{5/2}	3.67602	3.73738
66	4s4p ⁵ (³ P)4d	⁴ P _{3/2} ^o	3.05629	3.00729	218	4s ² 4p ⁴ 5g	² G _{9/2}	3.67975	3.74211
67	4s4p ⁵ (³ P)4d	⁴ P _{5/2} ^o	3.08185	3.02837	219	4s ² 4p ⁴ 5g	² G _{7/2}	3.67984	3.74212
68	4s ² 4p ⁴ 5d	⁴ D _{7/2}	2.98136	3.03583	220	4s ² 4p ⁴ 5g	² H _{9/2}	3.68028	3.74256
69	4s ² 4p ⁴ 5d	⁴ D _{5/2}	3.03631	3.03752	221	4s ² 4p ⁴ 5g	² H _{11/2}	3.68029	3.74259
70	4s ² 4p ⁴ (³ P)4f	⁴ D _{1/2} ^o	3.05667	3.04175					

NIST: http://physics.nist.gov/PhysRefData/ASD/levels_form.html

GRASP: present calculations from the GRASP code with 3990 levels

Table B

Mixing coefficients (MC) for the lowest 50 levels of Sr IV. Numbers outside and inside a bracket correspond to MC and the level, respectively. See Table 1 for definition of all levels.

Index	Configuration	Level	Mixing coefficients
1	$4s^2 4p^5$	$2P_{3/2}^o$	0.97(1)
2	$4s^2 4p^5$	$2P_{1/2}^o$	0.97(2)
3	$4s 4p^6$	$2S_{1/2}$	0.79(3)-0.59(39)
4	$4s^2 4p^4(^3P)4d$	$4D_{7/2}$	0.97(4)
5	$4s^2 4p^4(^3P)4d$	$4D_{5/2}$	0.96(5)
6	$4s^2 4p^4(^3P)4d$	$4D_{3/2}$	-0.96(6)
7	$4s^2 4p^4(^3P)4d$	$4D_{1/2}$	-0.97(7)
8	$4s^2 4p^4(^3P)4d$	$4F_{9/2}$	0.96(8)+0.23(20)
9	$4s^2 4p^4(^3P)4d$	$4F_{7/2}$	0.93(9)+0.22(16)+0.20(21)
10	$4s^2 4p^4(^3P)4d$	$4F_{5/2}$	0.97(10)
11	$4s^2 4p^4(^3P)4d$	$4F_{3/2}$	0.95(11)+0.22(29)
12	$4s^2 4p^4(^1D)4d$	$2P_{1/2}$	-0.22(7)-0.26(13)-0.64(35)+0.67(12)
13	$4s^2 4p^4(^3P)4d$	$4P_{1/2}$	-0.94(13)+0.22(35)
14	$4s^2 4p^4(^3P)4d$	$4P_{3/2}$	0.77(14)+0.33(34)-0.43(18)
15	$4s^2 4p^4(^1D)4d$	$2D_{3/2}$	-0.53(40)+0.32(34)+0.60(15)-0.29(18)-0.29(29)
16	$4s^2 4p^4(^3P)4d$	$2F_{7/2}$	-0.26(9)+0.78(16)+0.45(21)+0.29(25)
17	$4s^2 4p^4(^3P)4d$	$4P_{5/2}$	0.89(17)+0.22(36)+0.25(31)
18	$4s^2 4p^4(^1D)4d$	$2P_{3/2}$	-0.59(14)+0.22(40)+0.49(34)-0.24(15)-0.50(18)
19	$4s^2 4p^4(^1D)4d$	$2D_{5/2}$	-0.37(17)-0.46(22)+0.46(36)-0.55(19)+0.27(31)
20	$4s^2 4p^4(^1D)4d$	$2G_{9/2}$	0.23(8)-0.96(20)
21	$4s^2 4p^4(^1D)4d$	$2G_{7/2}$	0.42(16)-0.86(21)+0.24(25)
22	$4s^2 4p^4(^3P)4d$	$2F_{5/2}$	-0.76(22)-0.28(36)-0.37(24)+0.40(19)
23	$4s^2 4p^4 5s$	$4P_{5/2}$	0.96(23)
24	$4s^2 4p^4(^1D)4d$	$2F_{5/2}$	0.31(22)-0.90(24)-0.21(19)
25	$4s^2 4p^4(^1D)4d$	$2F_{7/2}$	0.37(16)-0.91(25)
26	$4s^2 4p^4 5s$	$4P_{3/2}$	0.77(26)+0.60(28)+0.21(33)
27	$4s^2 4p^4 5s$	$4P_{1/2}$	0.97(27)+0.22(49)
28	$4s^2 4p^4 5s$	$2P_{3/2}$	0.63(26)-0.75(28)
29	$4s^2 4p^4(^1S)4d$	$2D_{3/2}$	0.51(15)+0.20(18)+0.78(29)
30	$4s^2 4p^4 5s$	$2P_{1/2}$	0.97(30)-0.21(49)
31	$4s^2 4p^4(^1S)4d$	$2D_{5/2}$	0.39(19)+0.85(31)
32	$4s^2 4p^4 5s$	$2D_{5/2}$	-0.28(36)-0.92(32)
33	$4s^2 4p^4 5s$	$2D_{3/2}$	0.25(28)-0.96(33)
34	$4s^2 4p^4(^3P)4d$	$2P_{3/2}$	0.65(34)-0.26(15)+0.58(18)-0.24(97)
35	$4s^2 4p^4(^3P)4d$	$2P_{1/2}$	-0.65(35)-0.64(12)+0.26(93)
36	$4s^2 4p^4(^3P)4d$	$2D_{5/2}$	0.66(36)+0.49(19)-0.24(31)-0.32(32)-0.32(101)
37	$4s^2 4p^4 5p$	$4P_{5/2}^o$	-0.34(48)+0.92(37)
38	$4s^2 4p^4 5p$	$4P_{3/2}^o$	-0.22(45)+0.86(38)-0.25(51)+0.23(47)-0.24(57)
39	$4s^2 4p^4(^1D)4d$	$2S_{1/2}$	-0.40(3)-0.57(39)+0.59(49)+0.24(122)
40	$4s^2 4p^4(^3P)4d$	$2D_{3/2}$	-0.68(40)-0.41(15)+0.36(29)+0.33(96)
41	$4s^2 4p^4 5p$	$4P_{1/2}^o$	-0.82(41)-0.36(44)+0.23(52)+0.32(58)
42	$4s^2 4p^4 5p$	$4D_{7/2}^o$	0.97(42)+0.23(54)
43	$4s^2 4p^4 5p$	$2D_{5/2}^o$	0.53(48)+0.21(37)+0.78(43)+0.24(53)
44	$4s^2 4p^4 5p$	$2P_{1/2}^o$	-0.28(46)+0.51(41)-0.56(44)+0.30(52)+0.46(58)
45	$4s^2 4p^4 5p$	$4D_{3/2}^o$	0.74(45)+0.28(38)+0.47(50)-0.28(47)
46	$4s^2 4p^4 5p$	$4D_{1/2}^o$	0.91(46)+0.22(41)+0.24(52)+0.21(60)
47	$4s^2 4p^4 5p$	$2P_{3/2}^o$	0.56(45)-0.31(50)+0.59(47)-0.38(57)
48	$4s^2 4p^4 5p$	$4D_{5/2}^o$	0.76(48)+0.29(37)-0.57(43)
49	$4s^2 4p^4 5s$	$2S_{1/2}$	0.35(3)+0.48(39)+0.73(49)
50	$4s^2 4p^4 5p$	$2D_{3/2}^o$	0.33(38)+0.55(51)-0.54(50)-0.30(47)+0.36(57)

Table C

Comparison of energies (Ryd) for some levels of Sr IV between the GRASP (3990 levels) and FAC (12 137 levels) calculations. See Table 1 for indices of all levels.

Index	Configuration	Level	GRASP	FAC	ΔE	Percentage Difference
1	4s ² 4p ⁵	² P _{3/2} ^o	0.00000	0.00000	0.000	0.0
2	4s ² 4p ⁵	² P _{1/2} ^o	0.08515	0.08812	-0.003	-3.5
3	4s4p ⁶	² S _{1/2}	1.44244	1.46391	-0.021	-1.5
4	4s ² 4p ⁴ (³ P)4d	⁴ D _{7/2}	1.75452	1.78496	-0.030	-1.7
5	4s ² 4p ⁴ (³ P)4d	⁴ D _{5/2}	1.75612	1.78502	-0.029	-1.6
6	4s ² 4p ⁴ (³ P)4d	⁴ D _{3/2}	1.76432	1.79324	-0.029	-1.6
7	4s ² 4p ⁴ (³ P)4d	⁴ D _{1/2}	1.77444	1.80563	-0.031	-1.8
8	4s ² 4p ⁴ (³ P)4d	⁴ F _{9/2}	1.85784	1.88277	-0.025	-1.3
9	4s ² 4p ⁴ (³ P)4d	⁴ F _{7/2}	1.89004	1.91435	-0.024	-1.3
10	4s ² 4p ⁴ (³ P)4d	⁴ F _{5/2}	1.91248	1.94092	-0.028	-1.5
11	4s ² 4p ⁴ (³ P)4d	⁴ F _{3/2}	1.92068	1.94816	-0.027	-1.4
12	4s ² 4p ⁴ (¹ D)4d	² P _{1/2}	1.92312	1.94315	-0.020	-1.0
13	4s ² 4p ⁴ (³ P)4d	⁴ P _{1/2}	1.95489	1.97497	-0.020	-1.0
14	4s ² 4p ⁴ (³ P)4d	⁴ P _{3/2}	1.95507	1.97562	-0.021	-1.1
15	4s ² 4p ⁴ (¹ D)4d	² D _{3/2}	1.97045	1.99005	-0.020	-1.0
16	4s ² 4p ⁴ (³ P)4d	² F _{7/2}	1.97942	1.99863	-0.019	-1.0
17	4s ² 4p ⁴ (³ P)4d	⁴ F _{5/2}	1.98759	2.00848	-0.021	-1.1
18	4s ² 4p ⁴ (¹ D)4d	² P _{3/2}	1.99565	2.01784	-0.022	-1.1
19	4s ² 4p ⁴ (¹ D)4d	² D _{5/2}	2.01910	2.03959	-0.020	-1.0
20	4s ² 4p ⁴ (¹ D)4d	² G _{9/2}	2.04505	2.06394	-0.019	-0.9
21	4s ² 4p ⁴ (¹ D)4d	² G _{7/2}	2.04924	2.06975	-0.021	-1.0
22	4s ² 4p ⁴ (³ P)4d	² F _{5/2}	2.05338	2.07603	-0.023	-1.1
24	4s ² 4p ⁴ (¹ D)4d	² F _{5/2}	2.18489	2.19528	-0.010	-0.5
25	4s ² 4p ⁴ (¹ D)4d	² F _{7/2}	2.20325	2.21470	-0.011	-0.5
29	4s ² 4p ⁴ (¹ S)4d	² D _{3/2}	2.29294	2.30686	-0.014	-0.6
31	4s ² 4p ⁴ (¹ S)4d	² D _{5/2}	2.31329	2.32847	-0.015	-0.7
34	4s ² 4p ⁴ (³ P)4d	² P _{3/2}	2.41725	2.45337	-0.036	-1.5
35	4s ² 4p ⁴ (³ P)4d	² P _{1/2}	2.44445	2.48054	-0.036	-1.5
36	4s ² 4p ⁴ (³ P)4d	² D _{5/2}	2.45347	2.48627	-0.033	-1.3
40	4s ² 4p ⁴ (³ P)4d	² D _{3/2}	2.52376	2.56654	-0.043	-1.7
61	4s ² 4p ⁴ (³ P)4f	⁴ F _{9/2} ^o	2.96793	3.06145	-0.094	-3.2
62	4s ² 4p ⁴ (³ P)4f	⁴ F _{7/2} ^o	2.97174	3.06398	-0.092	-3.1
63	4s ² 4p ⁴ (³ P)4f	⁴ F _{5/2} ^o	2.97953	3.07113	-0.092	-3.1
64	4s ² 4p ⁴ (³ P)4f	⁴ F _{3/2} ^o	2.99056	3.08276	-0.092	-3.1
70	4s ² 4p ⁴ (³ P)4f	⁴ D _{1/2} ^o	3.04175	3.12919	-0.087	-2.9
72	4s ² 4p ⁴ (³ P)4f	⁴ G _{11/2} ^o	3.04851	3.13369	-0.085	-2.8
73	4s ² 4p ⁴ (³ P)4f	² F _{7/2} ^o	3.05012	3.13581	-0.086	-2.8
74	4s ² 4p ⁴ (³ P)4f	² F _{5/2} ^o	3.05057	3.13449	-0.084	-2.8
75	4s ² 4p ⁴ (³ P)4f	² G _{9/2} ^o	3.05418	3.13770	-0.084	-2.7
77	4s ² 4p ⁴ (³ P)4f	⁴ D _{3/2} ^o	3.05433	3.14399	-0.090	-2.9
80	4s ² 4p ⁴ (³ P)4f	⁴ D _{5/2} ^o	3.08883	3.18068	-0.092	-3.0
82	4s ² 4p ⁴ (³ P)4f	² D _{3/2} ^o	3.09237	3.18028	-0.088	-2.8
83	4s ² 4p ⁴ (³ P)4f	⁴ D _{7/2} ^o	3.09267	3.18379	-0.091	-2.9
84	4s ² 4p ⁴ (³ P)4f	⁴ G _{9/2} ^o	3.09969	3.18757	-0.088	-2.8
87	4s ² 4p ⁴ (³ P)4f	⁴ G _{7/2} ^o	3.11450	3.20436	-0.090	-2.9
88	4s ² 4p ⁴ (³ P)4f	⁴ G _{5/2} ^o	3.11686	3.20497	-0.088	-2.8
89	4s ² 4p ⁴ (³ P)4f	² G _{7/2} ^o	3.12866	3.21737	-0.089	-2.8
94	4s ² 4p ⁴ (³ P)4f	² D _{5/2} ^o	3.14073	3.22766	-0.087	-2.8
98	4s ² 4p ⁴ (¹ D)4f	² D _{5/2} ^o	3.20054	3.28899	-0.088	-2.8
102	4s ² 4p ⁴ (¹ D)4f	² D _{3/2} ^o	3.21271	3.29760	-0.085	-2.6
103	4s ² 4p ⁴ (¹ D)4f	² P _{1/2} ^o	3.22083	3.29974	-0.079	-2.4
105	4s ² 4p ⁴ (¹ D)4f	² P _{3/2} ^o	3.22357	3.30477	-0.081	-2.5
106	4s ² 4p ⁴ (¹ D)4f	² F _{7/2} ^o	3.22933	3.31772	-0.088	-2.7
107	4s ² 4p ⁴ (¹ D)4f	² H _{9/2} ^o	3.23064	3.31041	-0.080	-2.5
108	4s ² 4p ⁴ (¹ D)4f	² H _{11/2} ^o	3.23116	3.30982	-0.079	-2.4
110	4s ² 4p ⁴ (¹ D)4f	² F _{5/2} ^o	3.24733	3.33465	-0.087	-2.7
113	4s ² 4p ⁴ (¹ D)4f	² G _{7/2} ^o	3.27844	3.35699	-0.079	-2.4
114	4s ² 4p ⁴ (¹ D)4f	² G _{9/2} ^o	3.39470	3.35967	0.035	1.0
137	4s ² 4p ⁴ (¹ S)4f	² F _{7/2} ^o	3.42216	3.50584	-0.084	-2.4

Table C (continued)

Index	Configuration	Level	GRASP	FAC	ΔE	Percentage Difference
23	4s ² 4p ⁴ 5s	⁴ P _{5/2}	2.17395	2.15116	0.023	1.0
26	4s ² 4p ⁴ 5s	⁴ P _{3/2}	2.21122	2.18695	0.024	1.1
27	4s ² 4p ⁴ 5s	⁴ P _{1/2}	2.24728	2.22491	0.022	1.0
28	4s ² 4p ⁴ 5s	² P _{3/2}	2.26566	2.24073	0.025	1.1
30	4s ² 4p ⁴ 5s	² P _{1/2}	2.30524	2.28028	0.025	1.1
32	4s ² 4p ⁴ 5s	² D _{5/2}	2.39785	2.37129	0.027	1.1
33	4s ² 4p ⁴ 5s	² D _{3/2}	2.40477	2.37447	0.030	1.3
49	4s ² 4p ⁴ 5s	² S _{1/2}	2.63469	2.61923	0.015	0.6
37	4s ² 4p ⁴ 5p	⁴ P _{5/2} ^o	2.49996	2.48873	0.011	0.4
38	4s ² 4p ⁴ 5p	⁴ P _{3/2} ^o	2.50245	2.48963	0.013	0.5
41	4s ² 4p ⁴ 5p	⁴ P _{1/2} ^o	2.52932	2.51683	0.012	0.5
42	4s ² 4p ⁴ 5p	⁴ D _{7/2} ^o	2.54782	2.53549	0.012	0.5
43	4s ² 4p ⁴ 5p	² D _{5/2} ^o	2.55249	2.53759	0.015	0.6
44	4s ² 4p ⁴ 5p	² P _{1/2} ^o	2.58741	2.57098	0.016	0.6
45	4s ² 4p ⁴ 5p	⁴ D _{3/2} ^o	2.59655	2.57548	0.021	0.8
46	4s ² 4p ⁴ 5p	⁴ D _{1/2} ^o	2.60789	2.59592	0.012	0.5
47	4s ² 4p ⁴ 5p	² P _{3/2} ^o	2.61162	2.59364	0.018	0.7
48	4s ² 4p ⁴ 5p	⁴ D _{5/2} ^o	2.61764	2.60448	0.013	0.5
50	4s ² 4p ⁴ 5p	² D _{3/2} ^o	2.64532	2.63046	0.015	0.6
51	4s ² 4p ⁴ 5p	⁴ S _{3/2} ^o	2.65539	2.63687	0.019	0.7
52	4s ² 4p ⁴ 5p	² S _{1/2} ^o	2.66029	2.64431	0.016	0.6
53	4s ² 4p ⁴ 5p	² F _{5/2} ^o	2.73474	2.71481	0.020	0.7
54	4s ² 4p ⁴ 5p	² F _{7/2} ^o	2.74818	2.72829	0.020	0.7
55	4s ² 4p ⁴ 5p	² D _{3/2} ^o	2.78784	2.72420	0.064	2.3
56	4s ² 4p ⁴ 5p	² D _{5/2} ^o	2.80035	2.77807	0.022	0.8
57	4s ² 4p ⁴ 5p	² P _{3/2} ^o	2.82130	2.77417	0.047	1.7
58	4s ² 4p ⁴ 5p	² P _{1/2} ^o	2.85686	2.76898	0.088	3.1
59	4s ² 4p ⁴ 5p	² P _{3/2} ^o	2.94839	2.93990	0.008	0.3
60	4s ² 4p ⁴ 5p	² P _{1/2} ^o	2.94916	2.92947	0.020	0.7
68	4s ² 4p ⁴ 5d	⁴ D _{7/2}	3.03583	3.02896	0.007	0.2
69	4s ² 4p ⁴ 5d	⁴ D _{5/2}	3.03752	3.03032	0.007	0.2
71	4s ² 4p ⁴ 5d	⁴ D _{3/2}	3.04428	3.03583	0.008	0.3
76	4s ² 4p ⁴ 5d	⁴ D _{1/2}	3.05480	3.04434	0.010	0.3
78	4s ² 4p ⁴ 5d	⁴ F _{9/2}	3.06161	3.04947	0.012	0.4
79	4s ² 4p ⁴ 5d	² F _{7/2}	3.07641	3.05836	0.018	0.6
81	4s ² 4p ⁴ 5d	⁴ P _{1/2}	3.09218	3.07179	0.020	0.7
85	4s ² 4p ⁴ 5d	⁴ P _{3/2}	3.11020	3.08704	0.023	0.7
86	4s ² 4p ⁴ 5d	⁴ F _{5/2}	3.11358	3.08903	0.025	0.8
90	4s ² 4p ⁴ 5d	⁴ F _{7/2}	3.13151	3.11869	0.013	0.4
91	4s ² 4p ⁴ 5d	⁴ F _{3/2}	3.13216	3.12164	0.011	0.3
92	4s ² 4p ⁴ 5d	⁴ P _{5/2}	3.13371	3.12056	0.013	0.4
93	4s ² 4p ⁴ 5d	² P _{1/2}	3.13400	3.11552	0.018	0.6
95	4s ² 4p ⁴ 5d	² F _{5/2}	3.15224	3.13245	0.020	0.6
96	4s ² 4p ⁴ 5d	² D _{3/2}	3.15983	3.13212	0.028	0.9
97	4s ² 4p ⁴ 5d	² P _{3/2}	3.19133	3.16505	0.026	0.8
101	4s ² 4p ⁴ 5d	² D _{5/2}	3.20706	3.16648	0.041	1.3
111	4s ² 4p ⁴ 5d	² G _{7/2}	3.26191	3.24433	0.018	0.5
112	4s ² 4p ⁴ 5d	² G _{9/2}	3.26332	3.24632	0.017	0.5
116	4s ² 4p ⁴ 5d	² F _{5/2}	3.29350	3.26558	0.028	0.8
120	4s ² 4p ⁴ 5d	² F _{7/2}	3.30039	3.27235	0.028	0.8
122	4s ² 4p ⁴ 5d	² S _{1/2}	3.32014	3.27611	0.044	1.3
124	4s ² 4p ⁴ 5d	² P _{3/2}	3.32718	3.27597	0.051	1.5
125	4s ² 4p ⁴ 5d	² D _{5/2}	3.33937	3.27877	0.061	1.8
126	4s ² 4p ⁴ 5d	² P _{1/2}	3.35834	3.30026	0.058	1.7
127	4s ² 4p ⁴ 5d	² D _{3/2}	3.36312	3.29584	0.067	2.0
140	4s ² 4p ⁴ 5d	² D _{5/2}	3.46151	3.45032	0.011	0.3
148	4s ² 4p ⁴ 5d	² D _{3/2}	3.48141	3.45692	0.024	0.7
144	4s ² 4p ⁴ 5f	⁴ F _{9/2} ^o	3.47997	3.47217	0.008	0.2
145	4s ² 4p ⁴ 5f	⁴ F _{5/2} ^o	3.48051	3.47523	0.005	0.2
146	4s ² 4p ⁴ 5f	⁴ G _{11/2} ^o	3.48097	3.47133	0.010	0.3
147	4s ² 4p ⁴ 5f	⁴ F _{7/2} ^o	3.48102	3.47195	0.009	0.3

Table C (continued)

Index	Configuration	Level	GRASP	FAC	ΔE	Percentage Difference
149	4s ² 4p ⁴ 5f	⁴ F _{3/2} ^o	3.48378	3.47994	0.004	0.1
150	4s ² 4p ⁴ 5f	² G _{9/2} ^o	3.48431	3.47429	0.010	0.3
151	4s ² 4p ⁴ 5f	⁴ D _{1/2} ^o	3.48779	3.49314	-0.005	-0.2
152	4s ² 4p ⁴ 5f	² F _{5/2} ^o	3.48880	3.48798	0.001	0.0
153	4s ² 4p ⁴ 5f	² F _{7/2} ^o	3.48903	3.47946	0.010	0.3
154	4s ² 4p ⁴ 5f	² D _{3/2} ^o	3.49258	3.49269	0.000	0.0
165	4s ² 4p ⁴ 5f	⁴ D _{5/2} ^o	3.53113	3.54555	-0.014	-0.4
166	4s ² 4p ⁴ 5f	⁴ D _{7/2} ^o	3.54266	3.54353	-0.001	0.0
168	4s ² 4p ⁴ 5f	⁴ F _{3/2} ^o	3.54904	3.54683	0.002	0.1
169	4s ² 4p ⁴ 5f	⁴ G _{9/2} ^o	3.55098	3.54323	0.008	0.2
170	4s ² 4p ⁴ 5f	⁴ G _{5/2} ^o	3.55152	3.54853	0.003	0.1
171	4s ² 4p ⁴ 5f	² D _{5/2} ^o	3.55627	3.55359	0.003	0.1
172	4s ² 4p ⁴ 5f	² G _{7/2} ^o	3.55878	3.55219	0.007	0.2
173	4s ² 4p ⁴ 5f	² F _{5/2} ^o	3.58254	3.56150	0.021	0.6
174	4s ² 4p ⁴ 5f	² F _{7/2} ^o	3.58681	3.55546	0.031	0.9
199	4s ² 4p ⁴ 5f	² H _{9/2} ^o	3.67855	3.66615	0.012	0.3
200	4s ² 4p ⁴ 5f	² F _{7/2} ^o	3.67863	3.65744	0.021	0.6
201	4s ² 4p ⁴ 5f	² H _{11/2} ^o	3.68163	3.66678	0.015	0.4
202	4s ² 4p ⁴ 5f	² P _{3/2} ^o	3.68273	3.66060	0.022	0.6
204	4s ² 4p ⁴ 5f	² P _{1/2} ^o	3.68414	3.66073	0.023	0.6
205	4s ² 4p ⁴ 5f	² F _{5/2} ^o	3.68956	3.67138	0.018	0.5
206	4s ² 4p ⁴ 5f	² G _{7/2} ^o	3.69939	3.68352	0.016	0.4
207	4s ² 4p ⁴ 5f	² G _{9/2} ^o	3.69954	3.68388	0.016	0.4
208	4s ² 4p ⁴ 5f	² D _{5/2} ^o	3.70006	3.69248	0.008	0.2
209	4s ² 4p ⁴ 5f	² D _{3/2} ^o	3.70421	3.67756	0.027	0.7
228	4s ² 4p ⁴ 5f	² F _{7/2} ^o	3.78894	3.70501	0.084	2.2
237	4s ² 4p ⁴ 5f	² F _{7/2} ^o	3.86843	3.86864	0.000	0.0
238	4s ² 4p ⁴ 5f	² F _{5/2} ^o	3.86872	3.87176	-0.003	-0.1
155	4s ² 4p ⁴ 5g	⁴ G _{11/2} ^o	3.52370	3.50394	0.020	0.6
156	4s ² 4p ⁴ 5g	⁴ G _{9/2} ^o	3.52378	3.50403	0.020	0.6
157	4s ² 4p ⁴ 5g	⁴ G _{7/2} ^o	3.52395	3.50424	0.020	0.6
158	4s ² 4p ⁴ 5g	² G _{9/2} ^o	3.52408	3.50441	0.020	0.6
159	4s ² 4p ⁴ 5g	⁴ G _{5/2} ^o	3.52626	3.50674	0.020	0.6
160	4s ² 4p ⁴ 5g	⁴ F _{7/2} ^o	3.52643	3.50694	0.019	0.6
161	4s ² 4p ⁴ 5g	⁴ H _{13/2} ^o	3.52782	3.50823	0.020	0.6
162	4s ² 4p ⁴ 5g	² H _{11/2} ^o	3.52792	3.50835	0.020	0.6
163	4s ² 4p ⁴ 5g	⁴ F _{3/2} ^o	3.52971	3.51049	0.019	0.5
164	4s ² 4p ⁴ 5g	² F _{5/2} ^o	3.52980	3.51057	0.019	0.5
176	4s ² 4p ⁴ 5g	⁴ G _{5/2} ^o	3.59757	3.57825	0.019	0.5
177	4s ² 4p ⁴ 5g	⁴ F _{7/2} ^o	3.59770	3.57847	0.019	0.5
178	4s ² 4p ⁴ 5g	⁴ H _{11/2} ^o	3.59835	3.57901	0.019	0.5
179	4s ² 4p ⁴ 5g	⁴ H _{9/2} ^o	3.59847	3.57918	0.019	0.5
180	4s ² 4p ⁴ 5g	⁴ H _{7/2} ^o	3.60217	3.58334	0.019	0.5
181	4s ² 4p ⁴ 5g	⁴ F _{9/2} ^o	3.60218	3.58329	0.019	0.5
182	4s ² 4p ⁴ 5g	⁴ F _{9/2} ^o	3.60346	3.58595	0.018	0.5
183	4s ² 4p ⁴ 5g	² F _{7/2} ^o	3.60348	3.58602	0.017	0.5
210	4s ² 4p ⁴ 5g	² D _{3/2} ^o	3.73107	3.70493	0.026	0.7
211	4s ² 4p ⁴ 5g	² D _{5/2} ^o	3.73109	3.70492	0.026	0.7
212	4s ² 4p ⁴ 5g	² I _{13/2} ^o	3.73417	3.70776	0.026	0.7
213	4s ² 4p ⁴ 5g	² I _{11/2} ^o	3.73417	3.70777	0.026	0.7
215	4s ² 4p ⁴ 5g	² F _{7/2} ^o	3.73733	3.71146	0.026	0.7
216	4s ² 4p ⁴ 5g	² F _{5/2} ^o	3.73738	3.71152	0.026	0.7
218	4s ² 4p ⁴ 5g	² G _{9/2} ^o	3.74211	3.71630	0.026	0.7
219	4s ² 4p ⁴ 5g	² G _{7/2} ^o	3.74212	3.71632	0.026	0.7
220	4s ² 4p ⁴ 5g	² H _{9/2} ^o	3.74256	3.71678	0.026	0.7
221	4s ² 4p ⁴ 5g	² H _{11/2} ^o	3.74259	3.71680	0.026	0.7
245	4s ² 4p ⁴ 5g	² G _{7/2} ^o	3.91647	3.90515	0.011	0.3
246	4s ² 4p ⁴ 5g	² G _{9/2} ^o	3.91647	3.90513	0.011	0.3

Table D

Comparison of radiative rates (A , s^{-1}) and oscillator strengths (f , dimensionless) for some transitions of Sr IV between the GRASP (3990 levels) and FAC (12 137 levels) calculations. The last column gives the ratio of $f(\text{GRASP})$ and $f(\text{FAC})$. $a \pm b \equiv a \times 10^{\pm b}$. See Table 1 for definition of levels.

I	J	A(GRASP)	f (GRASP)	A(FAC)	f (FAC)	R
1	3	3.546+08	1.061-02	4.074+08	1.183-02	0.90
1	5	1.024+07	6.197-04	1.181+07	6.920-04	0.90
1	6	2.556+06	1.022-04	4.138+06	1.602-04	0.64
1	7	4.257+05	8.415-06	1.383+06	2.640-05	0.32
1	10	8.106+07	4.138-03	1.047+08	5.188-03	0.80
1	11	7.990+06	2.696-04	3.035+06	9.957-05	2.71
1	12	2.844+05	4.787-06	6.461+05	1.065-05	0.45
1	13	2.573+08	4.192-03	3.068+08	4.896-03	0.86
1	14	1.768+08	5.757-03	2.049+08	6.534-03	0.88
1	15	2.167+08	6.948-03	1.644+08	5.167-03	1.34
1	17	7.565+07	3.576-03	5.897+07	2.730-03	1.31
1	18	3.643+04	1.139-06	8.054+05	2.463-05	0.05
1	19	2.304+08	1.055-02	1.552+08	6.967-03	1.51
1	22	1.617+08	7.161-03	1.117+08	4.838-03	1.48
1	23	3.618+08	1.430-02	2.404+08	9.700-03	1.47
1	24	2.220+08	8.685-03	2.939+08	1.139-02	0.76
1	26	6.895+09	1.756-01	5.090+09	1.325-01	1.32
1	27	2.116+08	2.608-03	1.087+08	1.367-03	1.91
1	28	1.199+10	2.908-01	7.208+09	1.787-01	1.63
1	29	1.901+09	4.502-02	2.203+09	5.153-02	0.87
1	30	1.192+10	1.396-01	7.625+09	9.128-02	1.53
1	31	1.568+09	5.473-02	1.729+09	5.955-02	0.92
1	32	2.596+10	8.432-01	8.509+09	2.826-01	2.98
1	33	1.017+07	2.189-04	2.236+06	4.938-05	4.43
1	34	5.777+10	1.231+00	6.554+10	1.356+00	0.91
1	35	3.057+10	3.184-01	4.402+10	4.454-01	0.71
1	36	5.781+10	1.793+00	7.786+10	2.352+00	0.76
1	39	3.467+10	3.410-01	2.042+10	2.002-01	1.70
1	40	2.517+09	4.919-02	2.378+09	4.494-02	1.09
1	49	1.530+10	1.372-01	2.093+10	1.899-01	0.72
2	3	1.726+08	1.166-02	1.977+08	1.300-02	0.90
2	6	2.955+05	2.609-05	4.498+05	3.852-05	0.68
2	7	3.608+05	1.574-05	6.745+05	2.847-05	0.55
2	11	3.165+07	2.339-03	3.536+07	2.545-03	0.92
2	12	4.018+06	1.481-04	8.324+06	3.011-04	0.49
2	13	3.089+06	1.100-04	7.375+06	2.579-04	0.43
2	14	8.485+05	6.042-05	6.851+05	4.788-05	1.26
2	15	2.526+08	1.770-02	2.282+08	1.570-02	1.13
2	18	2.973+07	2.028-03	1.228+07	8.211-04	2.47
2	26	2.644+08	1.457-02	1.919+08	1.085-02	1.34
2	27	3.231+08	8.606-03	2.624+08	7.154-03	1.20
2	28	9.353+08	4.898-02	5.716+08	3.071-02	1.59
2	29	3.473+08	1.774-02	4.531+08	2.291-02	0.77
2	30	9.899+09	2.500-01	6.323+09	1.638-01	1.53
2	33	1.240+10	5.740-01	6.956+09	3.313-01	1.73
2	34	6.604+08	3.024-02	7.921+08	3.525-02	0.86
2	35	2.783+10	6.224-01	2.360+10	5.133-01	1.21
2	39	3.792+10	7.990-01	3.969+10	8.354-01	0.96
2	40	7.014+10	2.937+00	7.831+10	3.174+00	0.93
2	49	7.946+09	1.522-01	1.475+10	2.867-01	0.53

Explanation of Tables

Table 1. Energies (Ryd) for the lowest 375 levels of Sr IV and their lifetimes (τ , s).

Index	Level Index
Configuration	The configuration to which the level belongs
Level	The <i>LSJ</i> designation of the level
Energy	Present energies from the GRASP code with 3990 level calculations
τ (s)	Lifetime of the level in s

Table 2. Energies (Ryd) for the lowest 375 levels of Y V and their lifetimes (τ , s).

Index	Level Index
Configuration	The configuration to which the level belongs
Level	The <i>LSJ</i> designation of the level
Energy	Present energies from the GRASP code with 3990 level calculations
τ (s)	Lifetime of the level in s

Table 3. Energies (Ryd) for the lowest 375 levels of Zr VI and their lifetimes (τ , s).

Index	Level Index
Configuration	The configuration to which the level belongs
Level	The <i>LSJ</i> designation of the level
Energy	Present energies from the GRASP code with 3990 level calculations
τ (s)	Lifetime of the level in s

Table 4. Energies (Ryd) for the lowest 375 levels of Nb VII and their lifetimes (τ , s).

Index	Level Index
Configuration	The configuration to which the level belongs
Level	The <i>LSJ</i> designation of the level
Energy	Present energies from the GRASP code with 3990 level calculations
τ (s)	Lifetime of the level in s

Table 5. Energies (Ryd) for the lowest 375 levels of Mo VIII and their lifetimes (τ , s).

Index	Level Index
Configuration	The configuration to which the level belongs
Level	The <i>LSJ</i> designation of the level
Energy	Present energies from the GRASP code with 3990 level calculations
τ (s)	Lifetime of the level in s

Table 6. Transition wavelengths (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij} , dimensionless), and line strengths (S , in atomic units) for electric dipole (E1), and A_{ji} for electric quadrupole (E2), magnetic dipole (M1), and magnetic quadrupole (M2) transitions of Sr IV. The ratio $R(E1)$ of velocity and length forms of A- values for E1 transitions is listed in the last column.

i and j	The lower (i) and upper (j) levels of a transition as defined in Table 1.
λ_{ij}	Transition wavelength (in Å)
A_{ji}^{E1}	Radiative transition probability (in s^{-1}) for the E1 transitions
f_{ij}^{E1}	Absorption oscillator strength (dimensionless) for the E1 transitions
S_{ji}^{E1}	Line strength in atomic unit (a.u.), 1 a.u. = 6.460×10^{-36} cm ² esu ² for the E1 transitions
A_{ji}^{E2}	Radiative transition probability (in s^{-1}) for the E2 transitions
A_{ji}^{M1}	Radiative transition probability (in s^{-1}) for the M1 transitions
A_{ji}^{M2}	Radiative transition probability (in s^{-1}) for the M2 transitions
$R(E1)$	Ratio of velocity and length forms of A- (or f- and S-) values for the E1 transitions
$a \pm b$	$\equiv a \times 10^{\pm b}$

Table 7. Transition wavelengths (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij} , dimensionless), and line strengths (S , in atomic units) for electric dipole (E1), and A_{ji} for electric quadrupole (E2), magnetic dipole (M1), and magnetic quadrupole (M2) transitions of Y V. The ratio $R(E1)$ of velocity and length forms of A - values for E1 transitions is listed in the last column.

i and j	The lower (i) and upper (j) levels of a transition as defined in Table 2.
λ_{ij}	Transition wavelength (in Å)
A_{ji}^{E1}	Radiative transition probability (in s^{-1}) for the E1 transitions
f_{ij}^{E1}	Absorption oscillator strength (dimensionless) for the E1 transitions
S^{E1}	Line strength in atomic unit (a.u.), 1 a.u. = 6.460×10^{-36} cm ² esu ² for the E1 transitions
A_{ji}^{E2}	Radiative transition probability (in s^{-1}) for the E2 transitions
A_{ji}^{M1}	Radiative transition probability (in s^{-1}) for the M1 transitions
A_{ji}^{M2}	Radiative transition probability (in s^{-1}) for the M2 transitions
$R(E1)$	Ratio of velocity and length forms of A - (or f - and S -) values for the E1 transitions
$a \pm b$	$\equiv a \times 10^{\pm b}$

Table 8. Transition wavelengths (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij} , dimensionless), and line strengths (S , in atomic units) for electric dipole (E1), and A_{ji} for electric quadrupole (E2), magnetic dipole (M1), and magnetic quadrupole (M2) transitions of Zr VI. The ratio $R(E1)$ of velocity and length forms of A - values for E1 transitions is listed in the last column.

i and j	The lower (i) and upper (j) levels of a transition as defined in Table 3.
λ_{ij}	Transition wavelength (in Å)
A_{ji}^{E1}	Radiative transition probability (in s^{-1}) for the E1 transitions
f_{ij}^{E1}	Absorption oscillator strength (dimensionless) for the E1 transitions
S^{E1}	Line strength in atomic unit (a.u.), 1 a.u. = 6.460×10^{-36} cm ² esu ² for the E1 transitions
A_{ji}^{E2}	Radiative transition probability (in s^{-1}) for the E2 transitions
A_{ji}^{M1}	Radiative transition probability (in s^{-1}) for the M1 transitions
A_{ji}^{M2}	Radiative transition probability (in s^{-1}) for the M2 transitions
$R(E1)$	Ratio of velocity and length forms of A - (or f - and S -) values for the E1 transitions
$a \pm b$	$\equiv a \times 10^{\pm b}$

Table 9. Transition wavelengths (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij} , dimensionless), and line strengths (S , in atomic units) for electric dipole (E1), and A_{ji} for electric quadrupole (E2), magnetic dipole (M1), and magnetic quadrupole (M2) transitions of Nb VII. The ratio $R(E1)$ of velocity and length forms of A - values for E1 transitions is listed in the last column.

i and j	The lower (i) and upper (j) levels of a transition as defined in Table 4.
λ_{ij}	Transition wavelength (in Å)
A_{ji}^{E1}	Radiative transition probability (in s^{-1}) for the E1 transitions
f_{ij}^{E1}	Absorption oscillator strength (dimensionless) for the E1 transitions
S^{E1}	Line strength in atomic unit (a.u.), 1 a.u. = 6.460×10^{-36} cm ² esu ² for the E1 transitions
A_{ji}^{E2}	Radiative transition probability (in s^{-1}) for the E2 transitions
A_{ji}^{M1}	Radiative transition probability (in s^{-1}) for the M1 transitions
A_{ji}^{M2}	Radiative transition probability (in s^{-1}) for the M2 transitions
$R(E1)$	Ratio of velocity and length forms of A - (or f - and S -) values for the E1 transitions
$a \pm b$	$\equiv a \times 10^{\pm b}$

Table 10. Transition wavelengths (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij} , dimensionless), and line strengths (S , in atomic units) for electric dipole (E1), and A_{ji} for electric quadrupole (E2), magnetic dipole (M1), and magnetic quadrupole (M2) transitions of Mo VIII. The ratio $R(E1)$ of velocity and length forms of A- values for E1 transitions is listed in the last column.

i and j	The lower (i) and upper (j) levels of a transition as defined in Table 5.
λ_{ij}	Transition wavelength (in Å)
A_{ji}^{E1}	Radiative transition probability (in s^{-1}) for the E1 transitions
f_{ij}^{E1}	Absorption oscillator strength (dimensionless) for the E1 transitions
S^{E1}	Line strength in atomic unit (a.u.), 1 a.u. = 6.460×10^{-36} cm ² esu ² for the E1 transitions
A_{ji}^{E2}	Radiative transition probability (in s^{-1}) for the E2 transitions
A_{ji}^{M1}	Radiative transition probability (in s^{-1}) for the M1 transitions
A_{ji}^{M2}	Radiative transition probability (in s^{-1}) for the M2 transitions
$R(E1)$	Ratio of velocity and length forms of A- (or f- and S-) values for the E1 transitions
$a \pm b$	$\equiv a \times 10^{\pm b}$

Table 1Energies (Ryd) for the lowest 375 levels of Sr IV and their lifetimes (τ , s). $a\pm b \equiv a \times 10^{\pm b}$. See page 15 for Explanation of Tables

Index	Configuration	Level	Energy	τ (s)
1	4s ² 4p ⁵	² P _{3/2} ^o	0.00000
2	4s ² 4p ⁵	² P _{1/2} ^o	0.08515	6.816-02
3	4s4p ⁶	² S _{1/2}	1.44244	1.897-09
4	4s ² 4p ⁴ (³ P)4d	⁴ D _{7/2}	1.75452	1.196-01
5	4s ² 4p ⁴ (³ P)4d	⁴ D _{5/2}	1.75612	9.770-08
6	4s ² 4p ⁴ (³ P)4d	⁴ D _{3/2}	1.76432	3.506-07
7	4s ² 4p ⁴ (³ P)4d	⁴ D _{1/2}	1.77444	1.272-06
8	4s ² 4p ⁴ (³ P)4d	⁴ F _{9/2}	1.85784	7.869-01
9	4s ² 4p ⁴ (³ P)4d	⁴ F _{7/2}	1.89004	2.948-01
10	4s ² 4p ⁴ (³ P)4d	⁴ F _{5/2}	1.91248	1.234-08
11	4s ² 4p ⁴ (³ P)4d	⁴ F _{3/2}	1.92068	2.523-08
12	4s ² 4p ⁴ (¹ D)4d	² P _{1/2}	1.92312	2.324-07
13	4s ² 4p ⁴ (³ P)4d	⁴ P _{1/2}	1.95489	3.840-09
14	4s ² 4p ⁴ (³ P)4d	⁴ P _{3/2}	1.95507	5.630-09
15	4s ² 4p ⁴ (¹ D)4d	² D _{3/2}	1.97045	2.131-09
16	4s ² 4p ⁴ (³ P)4d	² F _{7/2}	1.97942	7.706-02
17	4s ² 4p ⁴ (³ P)4d	⁴ P _{5/2}	1.98759	1.322-08
18	4s ² 4p ⁴ (¹ D)4d	² P _{3/2}	1.99565	3.359-08
19	4s ² 4p ⁴ (¹ D)4d	² D _{5/2}	2.01910	4.340-09
20	4s ² 4p ⁴ (¹ D)4d	² G _{9/2}	2.04505	5.624-02
21	4s ² 4p ⁴ (¹ D)4d	² G _{7/2}	2.04924	6.753-02
22	4s ² 4p ⁴ (³ P)4d	² F _{5/2}	2.05338	6.185-09
.....				
.....				
.....				

Table 2Energies (Ryd) for the lowest 375 levels of Y V and their lifetimes (τ , s). $a\pm b \equiv a \times 10^{\pm b}$. See page 15 for Explanation of Tables

Index	Configuration	Level	Energy	τ (s)
1	$4s^2 4p^5$	$^2P_{3/2}^o$	0.00000
2	$4s^2 4p^5$	$^2P_{1/2}^o$	0.10989	3.171-02
3	$4s 4p^6$	$^2S_{1/2}$	1.64719	1.011-09
4	$4s^2 4p^4(^3P)4d$	$^4D_{5/2}$	2.04945	6.765-08
5	$4s^2 4p^4(^3P)4d$	$^4D_{7/2}$	2.04923	6.441-02
6	$4s^2 4p^4(^3P)4d$	$^4D_{3/2}$	2.05923	1.654-07
7	$4s^2 4p^4(^3P)4d$	$^4D_{1/2}$	2.07282	2.471-07
8	$4s^2 4p^4(^3P)4d$	$^4F_{9/2}$	2.16295	4.710-01
9	$4s^2 4p^4(^3P)4d$	$^4F_{7/2}$	2.20263	1.559-01
10	$4s^2 4p^4(^3P)4d$	$^4F_{5/2}$	2.23193	7.627-09
11	$4s^2 4p^4(^1D)4d$	$^2P_{1/2}$	2.23511	2.682-08
12	$4s^2 4p^4(^3P)4d$	$^4F_{3/2}$	2.23938	2.229-08
13	$4s^2 4p^4(^3P)4d$	$^4P_{3/2}$	2.27611	2.341-09
14	$4s^2 4p^4(^3P)4d$	$^4P_{1/2}$	2.27701	2.526-09
15	$4s^2 4p^4(^1D)4d$	$^2D_{3/2}$	2.29308	1.809-09
16	$4s^2 4p^4(^3P)4d$	$^2F_{7/2}$	2.30266	4.083-02
17	$4s^2 4p^4(^3P)4d$	$^4P_{5/2}$	2.31963	8.375-09
18	$4s^2 4p^4(^1D)4d$	$^2P_{3/2}$	2.32879	4.434-08
19	$4s^2 4p^4(^1D)4d$	$^2D_{5/2}$	2.35618	4.202-09
20	$4s^2 4p^4(^1D)4d$	$^2G_{9/2}$	2.37673	3.031-02
21	$4s^2 4p^4(^1D)4d$	$^2G_{7/2}$	2.38088	3.458-02
22	$4s^2 4p^4(^3P)4d$	$^2F_{5/2}$	2.39509	4.518-09
23	$4s^2 4p^4(^1D)4d$	$^2F_{5/2}$	2.53383	3.111-09
24	$4s^2 4p^4(^1D)4d$	$^2F_{7/2}$	2.55947	1.294-02
25	$4s^2 4p^4(^1S)4d$	$^2D_{3/2}$	2.65401	2.818-10
26	$4s^2 4p^4(^1S)4d$	$^2D_{5/2}$	2.68370	6.120-10
.....				
.....				
.....				

Table 3Energies (Ryd) for the lowest 375 levels of Zr VI and their lifetimes (τ , s). $a\pm b \equiv a \times 10^{\pm b}$. See page 15 for Explanation of Tables

Index	Configuration	Level	Energy	τ (s)
1	4s ² 4p ⁵	² P _{3/2} ^o	0.00000
2	4s ² 4p ⁵	² P _{1/2} ^o	0.13831	1.591-02
3	4s4p ⁶	² S _{1/2}	1.85108	6.388-10
4	4s ² 4p ⁴ (³ P)4d	⁴ D _{5/2}	2.33330	4.910-08
5	4s ² 4p ⁴ (³ P)4d	⁴ D _{7/2}	2.33503	3.883-02
6	4s ² 4p ⁴ (³ P)4d	⁴ D _{3/2}	2.34467	8.945-08
7	4s ² 4p ⁴ (³ P)4d	⁴ D _{1/2}	2.36246	9.255-08
8	4s ² 4p ⁴ (³ P)4d	⁴ F _{9/2}	2.45699	3.080-01
9	4s ² 4p ⁴ (³ P)4d	⁴ F _{7/2}	2.50424	9.109-02
10	4s ² 4p ⁴ (¹ D)4d	² P _{1/2}	2.53670	1.274-08
11	4s ² 4p ⁴ (³ P)4d	⁴ F _{5/2}	2.54249	4.784-09
12	4s ² 4p ⁴ (³ P)4d	⁴ F _{3/2}	2.54737	1.877-08
13	4s ² 4p ⁴ (³ P)4d	⁴ P _{3/2}	2.58564	1.409-09
14	4s ² 4p ⁴ (³ P)4d	⁴ P _{1/2}	2.58623	1.618-09
15	4s ² 4p ⁴ (¹ D)4d	² D _{3/2}	2.60562	1.547-09
16	4s ² 4p ⁴ (³ P)4d	² F _{7/2}	2.61536	2.261-02
17	4s ² 4p ⁴ (³ P)4d	⁴ P _{5/2}	2.63869	6.561-09
18	4s ² 4p ⁴ (¹ D)4d	² P _{3/2}	2.65265	3.524-08
19	4s ² 4p ⁴ (¹ D)4d	² D _{5/2}	2.68382	3.878-09
20	4s ² 4p ⁴ (¹ D)4d	² G _{9/2}	2.70085	1.701-02
21	4s ² 4p ⁴ (¹ D)4d	² G _{7/2}	2.70386	1.885-02
22	4s ² 4p ⁴ (³ P)4d	² F _{5/2}	2.72617	3.338-09
23	4s ² 4p ⁴ (¹ D)4d	² F _{5/2}	2.87088	2.591-09
24	4s ² 4p ⁴ (¹ D)4d	² F _{7/2}	2.90457	7.495-03
25	4s ² 4p ⁴ (¹ S)4d	² D _{3/2}	3.00386	2.173-10
26	4s ² 4p ⁴ (¹ S)4d	² D _{5/2}	3.04607	4.521-10
27	4s ² 4p ⁴ (³ P)4d	² P _{3/2}	3.22804	7.724-12
28	4s ² 4p ⁴ (³ P)4d	² P _{1/2}	3.26317	7.264-12
29	4s ² 4p ⁴ (³ P)4d	² D _{5/2}	3.29411	6.139-12
30	4s ² 4p ⁴ (¹ D)4d	² S _{1/2}	3.35217	6.177-12
31	4s ² 4p ⁴ (³ P)4d	² D _{3/2}	3.41203	6.164-12
....				
....				
....				

Table 4Energies (Ryd) for the lowest 375 levels of Nb VII and their lifetimes (τ , s). $a\pm b \equiv a \times 10^{\pm b}$. See page 15 for Explanation of Tables

Index	Configuration	Level	Energy	τ (s)
1	4s ² 4p ⁵	² P _{3/2} ^o	0.00000
2	4s ² 4p ⁵	² P _{1/2} ^o	0.17076	8.452-03
3	4s4p ⁶	² S _{1/2}	2.05605	4.422-10
4	4s ² 4p ⁴ (³ P)4d	⁴ D _{5/2}	2.61194	3.708-08
5	4s ² 4p ⁴ (³ P)4d	⁴ D _{7/2}	2.61624	2.526-02
6	4s ² 4p ⁴ (³ P)4d	⁴ D _{3/2}	2.62476	5.270-08
7	4s ² 4p ⁴ (³ P)4d	⁴ D _{1/2}	2.64747	4.359-08
8	4s ² 4p ⁴ (³ P)4d	⁴ F _{9/2}	2.74502	2.149-01
9	4s ² 4p ⁴ (³ P)4d	⁴ F _{7/2}	2.79938	5.802-02
10	4s ² 4p ⁴ (¹ D)4d	² P _{1/2}	2.83382	8.504-09
11	4s ² 4p ⁴ (³ P)4d	⁴ F _{3/2}	2.84881	1.221-08
12	4s ² 4p ⁴ (³ P)4d	⁴ F _{5/2}	2.84929	3.065-09
13	4s ² 4p ⁴ (³ P)4d	⁴ P _{1/2}	2.88874	1.058-09
14	4s ² 4p ⁴ (³ P)4d	⁴ P _{3/2}	2.89050	9.676-10
15	4s ² 4p ⁴ (¹ D)4d	² D _{3/2}	2.91412	1.302-09
16	4s ² 4p ⁴ (³ P)4d	² F _{7/2}	2.92434	1.277-02
17	4s ² 4p ⁴ (³ P)4d	⁴ P _{5/2}	2.95152	5.236-09
18	4s ² 4p ⁴ (¹ D)4d	² P _{3/2}	2.97333	2.515-08
19	4s ² 4p ⁴ (¹ D)4d	² D _{5/2}	3.00809	3.319-09
20	4s ² 4p ⁴ (¹ D)4d	² G _{7/2}	3.02353	1.080-02
21	4s ² 4p ⁴ (¹ D)4d	² G _{9/2}	3.02294	9.866-03
22	4s ² 4p ⁴ (³ P)4d	² F _{5/2}	3.05268	2.428-09
23	4s ² 4p ⁴ (¹ D)4d	² F _{5/2}	3.20237	2.085-09
24	4s ² 4p ⁴ (¹ D)4d	² F _{7/2}	3.24509	4.512-03
25	4s ² 4p ⁴ (¹ S)4d	² D _{3/2}	3.34908	1.597-10
26	4s ² 4p ⁴ (¹ S)4d	² D _{5/2}	3.40722	3.339-10
27	4s ² 4p ⁴ (³ P)4d	² P _{3/2}	3.60985	6.411-12
28	4s ² 4p ⁴ (³ P)4d	² P _{1/2}	3.64668	5.835-12
29	4s ² 4p ⁴ (³ P)4d	² D _{5/2}	3.68826	4.983-12
30	4s ² 4p ⁴ (¹ D)4d	² S _{1/2}	3.73214	5.337-12
31	4s ² 4p ⁴ (³ P)4d	² D _{3/2}	3.83277	5.015-12
....				
....				
....				

Table 5Energies (Ryd) for the lowest 375 levels of Mo VIII and their lifetimes (τ , s). $a \pm b \equiv a \times 10^{\pm b}$. See page 15 for Explanation of Tables

Index	Configuration	Level	Energy	τ (s)
1	$4s^2 4p^5$	$2P_{3/2}^o$	0.00000
2	$4s^2 4p^5$	$2P_{1/2}^o$	0.20760	4.703-03
3	$4s 4p^6$	$2S_{1/2}$	2.26319	3.251-10
4	$4s^2 4p^4 ({}^3P) 4d$	$4D_{5/2}$	2.88728	2.896-08
5	$4s^2 4p^4 ({}^3P) 4d$	$4D_{7/2}$	2.89480	1.737-02
6	$4s^2 4p^4 ({}^3P) 4d$	$4D_{3/2}$	2.90130	3.320-08
7	$4s^2 4p^4 ({}^3P) 4d$	$4D_{1/2}$	2.92955	2.339-08
8	$4s^2 4p^4 ({}^3P) 4d$	$4F_{9/2}$	3.02938	1.576-01
9	$4s^2 4p^4 ({}^3P) 4d$	$4F_{7/2}$	3.08991	3.971-02
10	$4s^2 4p^4 ({}^1D) 4d$	$2P_{1/2}$	3.12966	6.658-09
11	$4s^2 4p^4 ({}^3P) 4d$	$4F_{3/2}$	3.14527	6.446-09
12	$4s^2 4p^4 ({}^3P) 4d$	$4F_{5/2}$	3.15470	2.007-09
13	$4s^2 4p^4 ({}^3P) 4d$	$4P_{1/2}$	3.18757	7.094-10
14	$4s^2 4p^4 ({}^3P) 4d$	$4P_{3/2}$	3.19427	7.159-10
15	$4s^2 4p^4 ({}^1D) 4d$	$2D_{3/2}$	3.22159	1.084-09
16	$4s^2 4p^4 ({}^3P) 4d$	$2F_{7/2}$	3.23328	7.330-03
17	$4s^2 4p^4 ({}^3P) 4d$	$4P_{5/2}$	3.26103	4.093-09
18	$4s^2 4p^4 ({}^1D) 4d$	$2P_{3/2}$	3.29402	1.771-08
19	$4s^2 4p^4 ({}^1D) 4d$	$2D_{5/2}$	3.33205	2.629-09
20	$4s^2 4p^4 ({}^1D) 4d$	$2G_{7/2}$	3.34260	6.421-03
21	$4s^2 4p^4 ({}^1D) 4d$	$2G_{9/2}$	3.34577	5.890-03
22	$4s^2 4p^4 ({}^3P) 4d$	$2F_{5/2}$	3.37774	1.771-09
23	$4s^2 4p^4 ({}^1D) 4d$	$2F_{5/2}$	3.53186	1.672-09
24	$4s^2 4p^4 ({}^1D) 4d$	$2F_{7/2}$	3.58450	2.803-03
25	$4s^2 4p^4 ({}^1S) 4d$	$2D_{3/2}$	3.69325	1.160-10
26	$4s^2 4p^4 ({}^1S) 4d$	$2D_{5/2}$	3.77098	2.444-10
27	$4s^2 4p^4 ({}^3P) 4d$	$2P_{3/2}$	3.98377	5.455-12
28	$4s^2 4p^4 ({}^3P) 4d$	$2P_{1/2}$	4.01921	4.794-12
29	$4s^2 4p^4 ({}^3P) 4d$	$2D_{5/2}$	4.07180	4.199-12
30	$4s^2 4p^4 ({}^1D) 4d$	$2S_{1/2}$	4.10826	4.764-12
31	$4s^2 4p^4 ({}^3P) 4d$	$2D_{3/2}$	4.24675	4.229-12
....				
.....				
.....				

Table 6

Transition wavelengths (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij} , dimensionless), and line strengths (S, in atomic units) for electric dipole (E1), and A_{ji} for E2, M1, and M2 transitions in Sr IV. The last column gives R(E1), the ratio of velocity and length forms of A- values for E1 transitions. $a\pm b \equiv a \times 10^{\pm b}$. See page 15 for Explanation of Tables

i	j	λ_{ij}	A_{ji}^{E1}	f_{ij}^{E1}	S^{E1}	A_{ji}^{E2}	A_{ji}^{M1}	A_{ji}^{M2}	R(E1)
1	2	1.070+04	0.000+00	0.000+00	0.000+00	1.735-02	1.465+01	0.000+00	0.0+00
1	3	6.318+02	3.546+08	1.061-02	8.826-02	0.000+00	0.000+00	6.059+00	2.3-01
1	4	5.194+02	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	8.362+00	0.0+00
1	5	5.189+02	1.024+07	6.198-04	4.235-03	0.000+00	0.000+00	1.837-01	9.5-01
1	6	5.165+02	2.556+06	1.022-04	6.954-04	0.000+00	0.000+00	1.370-01	7.8-01
1	7	5.135+02	4.257+05	8.415-06	5.691-05	0.000+00	0.000+00	1.152+00	4.8-01
1	9	4.821+02	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	7.617-01	0.0+00
1	10	4.765+02	8.106+07	4.138-03	2.597-02	0.000+00	0.000+00	7.902-02	9.2-01
.....									
.....									
.....									

Table 7

Transition wavelengths (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij} , dimensionless), and line strengths (S, in atomic units) for electric dipole (E1), and A_{ji} for E2, M1, and M2 transitions in Y V. The last column gives R(E1), the ratio of velocity and length forms of A- values for E1 transitions. $a\pm b \equiv a \times 10^{\pm b}$. See page 16 for Explanation of Tables

i	j	λ_{ij}	A_{ji}^{E1}	f_{ij}^{E1}	S^{E1}	A_{ji}^{E2}	A_{ji}^{M1}	A_{ji}^{M2}	R(E1)
1	2	8.292+03	0.000+00	0.000+00	0.000+00	4.269-02	3.149+01	0.000+00	0.0+00
1	3	5.532+02	6.761+08	1.551-02	1.130-01	0.000+00	0.000+00	1.012+01	3.6-01
1	4	4.446+02	1.478+07	6.572-04	3.848-03	0.000+00	0.000+00	3.417-01	9.3-01
1	5	4.447+02	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	1.553+01	0.0+00
1	6	4.425+02	5.943+06	1.745-04	1.017-03	0.000+00	0.000+00	2.254-01	9.0-01
1	7	4.396+02	2.088+06	3.025-05	1.751-04	0.000+00	0.000+00	2.553+00	8.4-01
1	9	4.137+02	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	1.855+00	0.0+00
1	10	4.083+02	1.311+08	4.915-03	2.643-02	0.000+00	0.000+00	2.078-01	9.3-01
.....									
.....									
.....									

Table 8

Transition wavelengths (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij} , dimensionless), and line strengths (S, in atomic units) for electric dipole (E1), and A_{ji} for E2, M1, and M2 transitions in Zr VI. The last column gives R(E1), the ratio of velocity and length forms of A- values for E1 transitions. $a\pm b \equiv a \times 10^{\pm b}$. See page 16 for Explanation of Tables

i	j	λ_{ij}	A_{ji}^{E1}	f_{ij}^{E1}	S^{E1}	A_{ji}^{E2}	A_{ji}^{M1}	A_{ji}^{M2}	R(E1)
1	2	6.589+03	0.000+00	0.000+00	0.000+00	9.836-02	6.277+01	0.000+00	0.0+00
1	3	4.923+02	1.081+09	1.964-02	1.273-01	0.000+00	0.000+00	1.579+01	4.2-01
1	4	3.905+02	2.037+07	6.986-04	3.593-03	0.000+00	0.000+00	5.571-01	9.2-01
1	5	3.903+02	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	2.575+01	0.0+00
1	6	3.887+02	1.118+07	2.532-04	1.296-03	0.000+00	0.000+00	3.260-01	9.6-01
1	7	3.857+02	5.771+06	6.436-05	3.269-04	0.000+00	0.000+00	5.114+00	9.6-01
1	9	3.639+02	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	3.938+00	0.0+00
1	10	3.592+02	1.219+07	1.179-04	5.579-04	0.000+00	0.000+00	3.468+01	1.4+00
.....									
.....									
.....									

Table 9

Transition wavelengths (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij} , dimensionless), and line strengths (S, in atomic units) for electric dipole (E1), and A_{ji} for E2, M1, and M2 transitions in Nb VII. The last column gives R(E1), the ratio of velocity and length forms of A- values for E1 transitions. $a\pm b \equiv a \times 10^{\pm b}$. See page 16 for Explanation of Tables

i	j	λ_{ij}	A_{ji}^{E1}	f_{ij}^{E1}	S^{E1}	A_{ji}^{E2}	A_{ji}^{M1}	A_{ji}^{M2}	R(E1)
1	2	5.337+03	0.000+00	0.000+00	0.000+00	2.150-01	1.181+02	0.000+00	0.0+00
1	3	4.432+02	1.574+09	2.318-02	1.353-01	0.000+00	0.000+00	2.345+01	4.4-01
1	4	3.489+02	2.697+07	7.383-04	3.392-03	0.000+00	0.000+00	8.343-01	9.1-01
1	5	3.483+02	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	3.959+01	0.0+00
1	6	3.472+02	1.886+07	3.409-04	1.558-03	0.000+00	0.000+00	4.284-01	9.8-01
1	7	3.442+02	1.293+07	1.148-04	5.203-04	0.000+00	0.000+00	9.512+00	9.8-01
1	9	3.255+02	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	7.463+00	0.0+00
1	10	3.216+02	2.169+07	1.681-04	7.120-04	0.000+00	0.000+00	4.979+01	1.4+00
.....									
.....									
.....									

Table 10

Transition wavelengths (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij} , dimensionless), and line strengths (S, in atomic units) for electric dipole (E1), and A_{ji} for E2, M1, and M2 transitions in Mo VIII. The last column gives R(E1), the ratio of velocity and length forms of A- values for E1 transitions. $a\pm b \equiv a \times 10^{\pm b}$. See page 17 for Explanation of Tables

i	j	λ_{ij}	A_{ji}^{E1}	f_{ij}^{E1}	S^{E1}	A_{ji}^{E2}	A_{ji}^{M1}	A_{ji}^{M2}	R(E1)
1	2	4.389+03	0.000+00	0.000+00	0.000+00	4.485-01	2.122+02	0.000+00	0.0+00
1	3	4.026+02	2.157+09	2.621-02	1.390-01	0.000+00	0.000+00	3.360+01	4.5-01
1	4	3.156+02	3.453+07	7.734-04	3.214-03	0.000+00	0.000+00	1.176+00	9.0-01
1	5	3.148+02	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	5.756+01	0.0+00
1	6	3.141+02	2.953+07	4.367-04	1.806-03	0.000+00	0.000+00	5.210-01	9.8-01
1	7	3.111+02	2.562+07	1.859-04	7.613-04	0.000+00	0.000+00	1.662+01	9.7-01
1	9	2.949+02	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	1.285+01	0.0+00
1	10	2.912+02	2.995+07	1.903-04	7.298-04	0.000+00	0.000+00	6.731+01	1.3+00
.....									
.....									
.....									