

# Metastable Fractal Aggregates as a Result of Competition Between Diffusion-Limited Aggregation and Dissociation

Yuriy G. Gordienko and Elena E. Zasimchuk

Institute of Metal Physics of the National Academy of Sciences  
36 Vernadsky St., Kiev 252142, Ukraine  
*email:* gord@imp.kiev.ua

## Abstract

The cellular automaton model is used to simulate diffusion and aggregation with dissociation of point particles in 2D. A continuous phase transition is found that separates creation of compact aggregates and fractal ones. The transition is the function of pair-interaction energy ( $E_b$ ), type of neighborhood and temperature  $T$ . Manifestations of the transition in real physical systems are discussed.

## 1 Introduction

In the modern theory of fractal growth the significant efforts were dedicated to defining models ([2], [3]) and developing the theoretical concepts necessary to understand these model and to compute their properties analytically [1]. The crucial elements of the simulation models and theoretical schemes are:

- irreversibility;
- screening and freezing;
- spontaneous evolution to a fractal structure.

Moreover, the fractal properties are presumed to be well defined only asymptotically with respect to the large time and scale limits. It means that the problem of fractal growth consists in considering regions of the system that are very far from growing interface and that will not continue to be modified by further growth. But there are many experimental evidents of surface aggregates of different morphologies: fractal and dendritic [4], compact and fractal [5], convex and concave ones [6], which were determined by *kinetics* of 2D growth processes. Recent simulations brought to light dependence of aggregate morphology from coverage [7] in irreversible diffusion-limited aggregation (DLA). These observations of transition from fractal to compact aggregates can be explained by deposition of atoms in internal regions of the aggregates. But the passage from *irreversible* to *reversible* aggregation was also achieved by tuning a value of pair bond energy [8]. It points out an influence of dissociation processes on morphology of aggregates.

Investigation of morphology as a function of deposition conditions during DLA with dissociation (DLAD) is of great interest in the spirit of the great importance of building low-dimensional nanostructures with desirable geometry. That is why in the following the cellular automaton (CA) model of metastable fractal aggregates created by DLAD is considered (Sec.2). In Sec.3 the dependence of fractal dimensions as a function of kinetics of DLAD is presented and the results are discussed in Sec.4.

## 2 Cellular automaton model of diffusion-limited aggregation with dissociation

The rules of our CA model are similar to the classic DLA model [2] with the following addition: particles are randomly generated beyond a growing aggregate with a constant rate  $F$  and diffuse with a diffusion rate  $D \sim \exp(-E_m/k_B T)$ , where  $E_m$  is a migration energy. And they can dissociate from an aggregate with the following rate  $R \sim D_e \exp(-\sum_{NNs} E_b/k_B T)$ , where  $E_b$  is a pair bond energy,  $D_e \sim \exp(-E_e/k_B T)$  is a diffusion rate of particles in aggregate (for example, in the simplest case  $D = D_e$  [8]). Local interaction with the nearest neighbors is taken into account by summation. Our simulations involved a square lattice with sizes ranging from  $(100 \times 100)$  to  $(600 \times 600)$  with periodic boundary conditions. Simulations were conducted for different values of the ratio  $\kappa = \exp((E_b + E_m - E_e)/k_B T)$  and the ratio  $D/F$  ranging from  $10^4$  to  $10^9$ . The CA evolves in discrete time steps and the state of each site at the next time step is determined from the state of the site itself and those of the nearest-neighbor sites (NNs). The von Neumann neighborhood (with four NNs) and Moore one (with eight NNs and diagonal coupling) were used.

## 3 Metastable fractal aggregates with quasi-stable fractal dimensions




Figure 1: Typical morphologies created by diffusion-limited aggregation with dissociation (DLAD) on initial stages for the von Neumann neighborhood at  $D/F = 10^6$  for  $(100 \times 100)$  sections: a)  $\kappa = 20$ ; b)  $\kappa = 100$ ; c)  $\kappa = 200$ ; d)  $\kappa = 1000$ .

Fig.1 illustrates typical morphologies for the von Neumann neighborhood on initial stages at  $D/F = 10^6$  for  $(100 \times 100)$  sections: a)  $\kappa = 20$ ; b)  $\kappa = 100$ ; c)  $\kappa = 200$ ; d)  $\kappa = 1000$ . Fractal dimensions were determined from fits to the linear region in log-log plots, which were obtained by density-density correlation function ( $d_c$ ) and box counting measurement ( $d_b$ ). For this purpose a central part of an aggregate was considered as an inscribed square of maximum dimension. On initial stages of aggregation (the number of particle in aggregate

Table 1: Fractal dimensions ( $d_c$ ) for the von Neumann neighborhood obtained by density-density correlation measurements. Mean values of the interpolation lines are denoted by word "mean", standard deviations are shown in row "sd" and correlation coefficients are given in row "corr".

$\kappa$	20	40	100	150	200	400	1000	2000
mean	1.956	1.869	1.857	1.794	1.755	1.669	1.653	1.652
sd	0.012	0.014	0.005	0.008	0.003	0.007	0.004	0.006
corr	0.9998	0.9997	0.9999	0.9999	0.9999	0.9996	0.9999	0.9996

Table 2: Fractal dimensions ( $d_c$ ) for the Moore neighborhood obtained by density-density correlation measurements. Mean values of the interpolation lines are denoted by word "mean", standard deviations are shown in row "sd" and correlation coefficients are given in row "corr".

$\kappa$	4	10	20	40	100	200	400
mean	1.945	1.862	1.776	1.71	1.708	1.652	1.662
sd	0.007	0.007	0.003	0.007	0.006	0.018	0.027
corr	0.9997	0.9998	0.9999	0.9997	0.9997	0.9992	0.9980

$n < 10^2$ ) length scale is very small for reliable calculation of fractal dimensions. In the range  $10^2 < n < 10^3$  the dimensions grow and achieve asymptotic values for  $n \approx 10^3$ . The values of  $d_c$  shown in Table 1 for the von Neumann neighborhood and Table 2 for the Moore neighborhood with the main statistical characteristics of interpolation lines. Box counting measurements demonstrate several linear regions in log-log plots: from initial region with  $d_b = 2$  to a staircase part with alternating regions of  $d_b \approx 1$  and  $d_b \approx 2$ . The regions have small length scales for reliable determination of  $d_b$ . Although a core of aggregates becomes compact with time, they continue to generate "DLA-like" branches. When deposition was ceased, aggregates underwent significant restructructions. The previous simulations shown that aggregate splitted in many small parts and coalesced with creation of one large compact aggregate with  $d_c = 2$ .

## 4 Simulated phase diagrams as a new tool for investigation of surfaces

In the well-known work of Witten and Sander [2] DLA with sticking probability, which depends on the number of bonds to filled neighbor sites  $B$ , was considered *without* dissociation. Particle can stick to an aggregate only during a fraction  $s$  of its encounters:  $s = t^{3-B}$ , where  $t < 1$ . In addition to their conclusion that "long-range properties seem to be unaltered" for  $0.1 < t < 0.5$  we should like to emphasize the crucial changes in DLA caused by dissociation. It should be noted that the whole aggregate can split in unlinked parts, but scale invariance defined by density-density correlations remains.

The aforementioned results allow to propose the new method for investigation of adatom-adatom and adatom-surface interactions by means of investigation of scale invariant aggregates. For this purpose careful simulations of DLAD for different deposition rate,  $\kappa$ , type of interaction are necessary. It will give model phase diagrams for any kind of real aggregating

adatoms and substrate. Testing the real aggregation processes for different deposition conditions and comparing theirs with the simulated model phase diagrams will allow to bring to light new features of local interaction at surfaces. We used the naive Arrhenius-type expressions for dissociation rates. It is obvious the more realistic scheme will need the more sophisticated algorithms and codes with more reliable results.

In addition to this we recently observed the very complicated band patterns in metals during plastic deformation of metals [9]. They can be explained by development of disturbances of concentration of point-type defects and their agglomerates [10]. The crucial point of this assumption is based on possibility of point defects to aggregate in scale invariant structures with different dimensions and consequently different abilities to capture point-type defects. Fractal dimensions of aggregates are dependent on pair interaction energy and they define the selection rate among heterogeneities with further band pattern formation. The extended version of aforementioned investigation in 3D will also allow to make clear mechanisms of pattern formation in plastically deformed metals.

**Acknowledgments.** The authors gratefully acknowledge Prof.A.A.Snarski for helpful comments on this work.

## References

- [1] Erzan, A, Pietronero, L.: The fixed-scale transformation approach to fractal growth. *Reviews of Modern Physics* **67** (1995) 545-604
- [2] Witten, T, A, Sander, L, M.: Diffusion-limited aggregation. *Physical Review B* **27** (1983) 5686-5697
- [3] Niemeyer, J, Pietronero, L, Wiesmann H, J.: Fractal dimension of dielectric breakdown. *Physical Review Letters* **52** (1984) 1033-1036
- [4] Brune, H, Romainczyk, C, Roder, H, Kern, K.: Mechanism of the transition from fractal to dendritic growth of surface aggregates. *Nature* **369** (1994) 469-471
- [5] Roder, H, Hahn, E, Brune, H, Bucher, J-P, Kern, K.: Building one- and two-dimensional nanostructures by diffusion-controlled aggregation at surfaces. *Nature* **366** (1993) 141-143
- [6] Michely, Th, Hohage, M, Bott, M, Comsa, G.: Inversion of growth speed anisotropy in two dimensions. *Physical Review Letters* **70** (1993) 3943-3946
- [7] Amar, J, G, Family, F, Lam, P-M.: Dynamic scaling of the island-size distribution and percolation in a model of submonolayer molecular-beam epitaxy. *Physical Review B* **50** (1994) 8781-8797
- [8] Ratsch, C, Zangwill, A, Smilauer, P, Vvedensky, D, D.: Saturation and scaling of epitaxial island densities. *Physical Review Letters* **72** (1994) 3194-3197
- [9] Gordienko, Yu, G, Zasimchuk, E, E.: Single-crystal indicators of fatigue and plastic deformation damage. *Proc. of Second Europ. Conf. on Smart Structures and Materials (Glasgow, UK)*, ed. by P.Gardiner, A.McDonach, R.S.McEwen, *Proc. SPIE* **2361** (1994) 312-315
- [10] Gordienko, Yu, G, Zasimchuk, E, E.: Synergetic model of structure formation during plastic deformation of crystals. *Philosophical Magazine* **A70** 1 (1994) 99-107