

# Symmetric energy-momentum tensor: the Abraham form and the explicitly covariant formula

V.V. Nesterenko\* and A.V. Nesterenko

*Bogoliubov Laboratory of Theoretical Physics,  
Joint Institute for Nuclear Research,  
141980 Dubna, Russia*

## Abstract

We compare the known in literature, explicitly covariant 4-dimensional formula for the symmetric energy-momentum tensor of electromagnetic field in a medium and the energy-momentum tensor derived by Abraham in the 3-dimensional vector form. It is shown that these two objects coincide only on the physical configuration space  $\bar{\Gamma}$ , formed by the field vectors and the velocity of the medium, which satisfy the Minkowski constitutive relations. It should be emphasized that the 3-dimensional vector formulae for the components of the energy-momentum tensor were obtained by Abraham only on  $\bar{\Gamma}$ , and the task of their extension to the whole unconditional configuration space  $\Gamma$  was not posed. In order to accomplish the comparison noted above we derive the covariant formula anew by another method, namely, by generalizing the Abraham reasoning. The comparison conducted enables one to treat the explicitly covariant formula as a unique consistent extension of the Abraham formulae to the whole configuration space  $\Gamma$ . Thus the question concerning the relativistic covariance of the original 3-dimensional Abraham formulae defined on  $\bar{\Gamma}$  is solved positively. We discuss in detail the relativistic covariance of the 3-dimensional vector formulae for individual components of the 4-dimensional tensors in electrodynamics which is manifested in the form-invariance of these formulae under Lorentz transformations.

PACS numbers: 03.30.+p, 03.50.De, 41.20. -q

Keywords: Symmetric energy-momentum tensor, the Abraham tensor, the Minkowski energy-momentum tensor, Minkowski's phenomenological electrodynamics, electrodynamics of continuous moving media, relativistic covariance and form-invariance.

---

\* E-mail: nestr@theor.jinr.ru

## CONTENTS

I. Introduction	3
II. The Minkowski energy-momentum tensor	5
III. The Abraham energy-momentum tensor	7
IV. The symmetric energy-momentum tensor	13
V. Form-invariance of the vector formulae for the energy-momentum tensor	18
A. The methods of defining the Minkowski tensor and the form-invariance of the vector formulae involved	18
1. Definition of the Minkowski tensor by equation (2.4)	18
2. Definition of the Minkowski tensor by the vector equations (2.5)–(2.8)	19
B. Form-invariance of the symmetric energy-momentum tensor and the Abraham tensor in the vector notation	20
VI. Conclusion	22
Acknowledgments	24
References	24

## I. INTRODUCTION

In the electrodynamics of continuous media, it is generally accepted that the symmetric energy-momentum tensor of electromagnetic field was derived by Abraham (see, for example, the comprehensive survey by Pauli [1]). However, it is worth noting that Abraham has derived the components of his energy-momentum tensor only assuming that the Minkowski constitutive relations are satisfied, i.e., only on the physical configuration space  $\bar{\Gamma}$  of changing the field vectors  $\mathbf{E}$ ,  $\mathbf{H}$ ,  $\mathbf{D}$ ,  $\mathbf{B}$  and the velocity of the medium  $\mathbf{v}$ ,  $v < c$ . The task to extend the Abraham formulae to the whole unconditional configuration space  $\Gamma$  was not posed. This peculiarity of the Abraham formulae completely concerns also the 4-dimensional presentation of these formulae proposed by Grammel [2]. This fact explains in particular the lack of uniqueness of the Grammel formulae (three versions) and their explicit dependence on  $\varepsilon$  and  $\mu$ . Obviously, the Abraham and Grammel formulae should be used only on  $\bar{\Gamma}$ , where they coincide and do not depend explicitly on  $\varepsilon$  and  $\mu$ , but not on the whole configuration space  $\Gamma$ . In the Pauli survey [1] and in the subsequent papers dealing with the Abraham and Grammel formulae these peculiarities were not noted and not taken into account [3–8].

Let us make more precise the used terminology. Following Pauli, we define the symmetric energy-momentum tensor by its known values in co-moving frame [1, Eqs. (298) and (300)]. By Abraham’s formulae, or simply by the Abraham tensor, we call, following again Pauli [1], the set of formulae for all the components of the energy-momentum tensor, which were obtained by Abraham on  $\bar{\Gamma}$  (see §§38, 39 in his book [9] and also in his articles [10, 11]). These formulae will also be brought about in the present paper in Section III. The formulae derived by Abraham are valid only on  $\bar{\Gamma}$ , therefore in advance it is not apparently that they define the components of a 4-dimensional relativistic vector. This point is also investigated in our paper.

The symmetric energy-momentum tensor, as it was specified by Pauli and defined on the whole configuration space  $\Gamma$ , has been constructed considerably later than the Abraham works [12–15]. These authors proposed the explicitly covariant 4-dimensional formula for this tensor, which is applicable both in the rest frame and for moving media. It is important that this formula does not contain explicitly the material characteristics  $\varepsilon$  and  $\mu$  of the medium. In the books [12, 13] the covariant formula is given without derivation. In Ref. [14] the symmetric energy momentum tensor was derived by “abrahamization” of the Minkowski

tensor. Further development and application of this method see in [16–18]. The technique used in [15] will be discussed in Section VB of the present paper. The authors of all these works call their tensor by the Abraham tensor. However, the comparison of these two objects was not conducted. In the present paper this gap will be filled up.

For this aim, we generalize the Abraham reasoning which he followed when deriving his energy-momentum tensor on  $\bar{\Gamma}$ . This enables us to construct the symmetric energy-momentum tensor on the whole configuration space  $\Gamma$  in another way different from the calculations<sup>1</sup> in papers [14, 15]. In our approach, the identity of the symmetric energy-momentum tensor and the Abraham tensor on  $\bar{\Gamma}$  is proved easily.<sup>2</sup> In addition, we derive at the same time the explicit 3-dimensional vector formulae for all the components of the symmetric energy-momentum tensor. The latter is important due to the following.

In practical calculations, one uses, as a rule, the individual components of the energy-momentum tensor in 3-dimensional vector notation. In particular, such a situation takes place in the electrodynamics of moving bodies. The point is that the Lorentz vector describing the velocity of the medium

$$u_\alpha = \gamma\{\mathbf{q}, i\}, \quad \mathbf{q} = \mathbf{v}/c, \quad \gamma^{-1} = \sqrt{1 - \mathbf{q}^2}, \quad u_\alpha u_\alpha = -1, \quad \alpha = 1, 2, 3, 4 \quad (1.1)$$

cannot be itself considered as a small parameter. Only its spatial part,  $\mathbf{q}$ , may be such a parameter. The 3-dimensional vector formulae for the energy-momentum tensor in the Minkowski form and in the Abraham form are known [1, 9]. In the present paper, the analogous formulae will be derived for the symmetric energy-momentum tensor. A special attention will be paid to the discussion of the relativistic covariance of these formulae.

The layout of the paper is as follows. In Section II, the formulae are brought which define the energy-momentum tensor in the Minkowski form and the Minkowski constitutive relations are presented. This material is substantially used in the following. In Section III, the derivation of the energy-momentum tensor by Abraham is traced in detail. In Section IV the generalization of the Abraham reasoning enables us to construct the symmetric energy-momentum tensor on the whole configuration space  $\Gamma$ . Here the identity of the symmetric energy-momentum tensor and the Abraham tensor on  $\bar{\Gamma}$  is proved. At the same time, the explicit 3-dimensional vector formulae are derived for all the components of the symmetric

---

<sup>1</sup> The method used in [15] is discussed in Section VB of the present paper.

<sup>2</sup> To do the same proceeding from the formula obtained in [14, 15] is a formidable task.

energy-momentum tensor. These formulae are analogous to those derived by Abraham. In Section V, we discuss different methods of defining the tensors of the second rank in electrodynamics and the form-invariance of the pertinent 3-dimensional vector formulae. It is shown in detail in what way this form-invariance becomes apparent in the case of the energy-momentum tensor in the Minkowski form, in the Abraham form, and in the case of the symmetric energy-momentum tensor. In Conclusion, Section VI, the obtained results are summarized briefly and some comments are made.

We do not touch the Abraham-Minkowski controversy and the problem of determining the “correct” energy-momentum tensor in macroscopic electrodynamics. There is a large body of the literature on this subject, see, for example, the reviews [3, 5–8, 14, 15] and references therein.

We use the following notations. The Greek indices take values 1, 2, 3, 4, whereas the Latin indices assume the values  $x, y, z$ . The unrationalized Gaussian units are used for the electromagnetic field and the notations generally accepted in macroscopic electrodynamics [19] are adopted.

## II. THE MINKOWSKI ENERGY-MOMENTUM TENSOR

We will frequently refer to the energy-momentum tensor in the Minkowski form. Therefore, we bring here the formulae needed. This tensor is a straightforward generalization of the energy-momentum tensor of electromagnetic field in vacuum [20],

$$T_{\alpha\beta} = T_{\beta\alpha} = \frac{1}{4\pi} \left( F_{\alpha\gamma} F_{\gamma\beta} - \frac{\delta_{\alpha\beta}}{4} F_{\gamma\delta} F_{\delta\gamma} \right), \quad (2.1)$$

where  $F_{\alpha\beta} = -F_{\beta\alpha}$  is electromagnetic field tensor in vacuum,  $F_{ij} = \varepsilon_{ijk} H_k$ ,  $F_{4j} = -F_{j4} = iE_j$ .

Minkowski [1, 19] has formulated relativistic macroscopic electrodynamics in terms of two tensors which are defined by the formulae

$$F_{ij} = \varepsilon_{ijk} B_k, \quad F_{4j} = -F_{j4} = iE_j; \quad (2.2)$$

$$H_{ij} = \varepsilon_{ijk} H_k, \quad H_{4j} = -H_{j4} = iD_j. \quad (2.3)$$

Generalizing (2.1), Minkowski proposed the following energy-momentum tensor of electro-

magnetic field in a material medium [1]:

$$T_{\alpha\beta}^{\text{M}} = \frac{1}{4\pi} \left( F_{\alpha\gamma} H_{\gamma\beta} - \frac{\delta_{\alpha\beta}}{4} F_{\gamma\nu} H_{\nu\gamma} \right). \quad (2.4)$$

This definition holds both for the medium at rest and for the moving medium. According to construction, the velocity of the medium does not enter into Eq. (2.4). The Minkowski energy-momentum tensor is not symmetric  $T_{\alpha\beta}^{\text{M}} \neq T_{\beta\alpha}^{\text{M}}$  and its trace vanishes  $T_{\alpha\alpha}^{\text{M}} = 0$ .

By making use of the 3-dimensional vector notation we can present the components of the tensor  $T_{\alpha\beta}^{\text{M}}$  (2.4), in the arbitrary inertial reference frame, in the form [1]

$$T_{\alpha\beta}^{\text{M}} = \begin{pmatrix} \sigma_{ij}^{\text{M}} & -ic \mathbf{g}^{\text{M}} \\ -\frac{i}{c} \mathbf{S}^{\text{M}} & w^{\text{M}} \end{pmatrix}, \quad (2.5)$$

where

$$\sigma_{ij}^{\text{M}} = \frac{1}{4\pi} \left\{ E_i D_j + H_i B_j - \frac{\delta_{ij}}{2} (\mathbf{E}\mathbf{D} + \mathbf{H}\mathbf{B}) \right\}, \quad (2.6)$$

$$\frac{1}{c} \mathbf{S}^{\text{M}} = \frac{1}{c} \mathbf{S} = \frac{1}{4\pi} [\mathbf{E}\mathbf{H}], \quad c\mathbf{g}^{\text{M}} = \frac{1}{4\pi} [\mathbf{D}\mathbf{B}], \quad (2.7)$$

$$w^{\text{M}} = \frac{1}{8\pi} (\mathbf{E}\mathbf{D} + \mathbf{H}\mathbf{B}). \quad (2.8)$$

Equation (2.7) involves the Poynting vector

$$\mathbf{S} = \frac{c}{4\pi} [\mathbf{E}\mathbf{H}]. \quad (2.9)$$

In macroscopic electrodynamics [1, 19], the important role is played by the Minkowski constitutive relations:

$$D_\alpha = \varepsilon E_\alpha, \quad (2.10)$$

$$B_{\alpha\beta\gamma} = \mu H_{\alpha\beta\gamma}. \quad (2.11)$$

We use the notation from [21]

$$E_\alpha \equiv F_{\alpha\beta} u_\beta = \gamma \{ \mathbf{E} + [\mathbf{q}\mathbf{B}], i(\mathbf{q}\mathbf{E}) \}, \quad D_\alpha \equiv H_{\alpha\beta} u_\beta = \gamma \{ \mathbf{D} + [\mathbf{q}\mathbf{H}], i(\mathbf{q}\mathbf{D}) \}, \quad (2.12)$$

$$B_{\alpha\beta\gamma} = F_{\alpha\beta} u_\gamma + F_{\beta\gamma} u_\alpha + F_{\gamma\alpha} u_\beta, \quad H_{\alpha\beta\gamma} = H_{\alpha\beta} u_\gamma + H_{\beta\gamma} u_\alpha + H_{\gamma\alpha} u_\beta, \quad (2.13)$$

where  $u_\alpha$  is the four-vector of the medium velocity (1.1). In terms of the 3-dimensional vector notation Eqs. (2.10), (2.11) acquire the form [1, 19]

$$\mathbf{D} + [\mathbf{q}\mathbf{H}] = \varepsilon (\mathbf{E} + [\mathbf{q}\mathbf{B}]), \quad (2.14)$$

$$\mathbf{B} - [\mathbf{q}\mathbf{E}] = \mu (\mathbf{H} - [\mathbf{q}\mathbf{D}]). \quad (2.15)$$

Only due to the constitutive relations (2.10), (2.11) or (2.14), (2.15) the formal scheme of the macroscopic electrodynamics acquires the physical content [1, § 33]. The values of the vectors  $\mathbf{E}$ ,  $\mathbf{H}$ ,  $\mathbf{D}$ ,  $\mathbf{B}$  and the velocity of the medium  $\mathbf{v}$ ,  $v < c$ , which satisfy the Minkowski constitutive relations (2.10), (2.11) will be referred to as *the physical configuration space*  $\bar{\Gamma}$ .

In the relativistic electrodynamics of moving media, since the Minkowski works, the 4-dimensional generalization of the Poynting vector (2.9) is used (the so-called Ruhstrahlvektor [1])

$$\Omega_\alpha = H_{\alpha\beta\gamma} u_\beta E_\gamma. \quad (2.16)$$

For simplicity, we shall call  $\Omega_\alpha$  the Minkowski vector. In the co-moving reference frame ( $K'$ ), Eq. (2.16) yields [1, 21]

$$\frac{c}{4\pi} (\Omega'_1, \Omega'_2, \Omega'_3) = \mathbf{S}', \quad \Omega'_4 = 0. \quad (2.17)$$

Following Pauli [1], we mark quantities in a co-moving frame by prime. From Eqs. (2.17) and (1.1) it follows

$$u_\alpha \Omega_\alpha = 0. \quad (2.18)$$

### III. THE ABRAHAM ENERGY-MOMENTUM TENSOR

Abraham imposes the following conditions on the energy-momentum tensor to be found. This tensor should be symmetric; in the co-moving reference frame ( $\mathbf{v} = 0$ ) and under fulfillment of the constitutive relations<sup>3</sup>

$$\mathbf{D} = \varepsilon \mathbf{E}, \quad \mathbf{B} = \mu \mathbf{H} \quad (3.1)$$

(isotropic medium) the components of this tensor should assume the values

$$\sigma'_{ij}{}^A = \frac{1}{2} (\sigma'_{ij}{}^M + \sigma'_{ji}{}^M), \quad (3.2)$$

$$\frac{1}{c} \mathbf{S}'^A = c \mathbf{g}'^A = \frac{1}{c} \mathbf{S}' = \frac{1}{4\pi} [\mathbf{E}' \mathbf{H}'], \quad (3.3)$$

$$w'^A = \frac{1}{8\pi} (\mathbf{E}' \mathbf{D}' + \mathbf{H}' \mathbf{B}'). \quad (3.4)$$

---

<sup>3</sup> The account of the constitutive relations in Abraham's derivation of the energy-momentum tensor is not noted in the Pauli review [1] and in the subsequent papers, see, for example [3, 5–8, 15].

We have to stress here that all these requirements are not sufficient to uniquely define the energy-momentum tensor on the whole configuration space  $\Gamma$ . Indeed, Abraham, having been fixed the reference frame ( $\mathbf{v} = 0$ ), assigns the components of his tensor only on the physical configuration space  $\bar{\Gamma}$  (i.e., under fulfillment of the constitutive relations (3.1)), but not on the whole configuration space  $\Gamma$ , as the theory of relativity demands.

However, Abraham did not use the theory of relativity in constructing his energy-momentum tensor. “He went along the way of extrapolations proceeding from the medium at rest, relying on rather arbitrary assumptions and very poor experimental data” [5, p. 289, Russian edition]). Following this way, Abraham has derived the energy-momentum tensor first in the framework of his approach to the electrodynamics of moving bodies [10, §§ 38, 39] and later on in application to the Minkowski electrodynamics [11]. In the paper [11] Abraham approached the most close to using the tensor formalism in the problem under consideration. The Abraham reasoning [11] was “translated” to the standard tensor language by Grammel [11], the pertinent detailed calculations were also conducted by Kafka [21]. We shall follow the papers [2, 11, 21].

By making use of the standard tensor notation the Abraham reasoning [11] can be reformulated like this. In the co-moving reference frame ( $\mathbf{v} = 0$ ) and when the constitutive relations (3.1) hold, the symmetric tensor

$$\frac{1}{2}(T_{\alpha\beta}^M + T_{\beta\alpha}^M) \quad (3.5)$$

gives the following value for the density of the energy flux:

$$\frac{c}{4\pi} \frac{1}{2}([\mathbf{DB}] + [\mathbf{EH}]|_{\bar{\Gamma}; \mathbf{v}=0}) = \frac{\varepsilon\mu + 1}{2} \frac{c}{4\pi} [\mathbf{EH}]. \quad (3.6)$$

In order to get here the Poynting vector (2.9), in accordance with the requirement (3.3), one has to subtract from (3.6) the following quantity

$$\frac{\varepsilon\mu - 1}{2} \frac{c}{4\pi} [\mathbf{EH}] = \frac{c}{4\pi} \frac{1}{2}([\mathbf{DB}] - [\mathbf{EH}]|_{\bar{\Gamma}; \mathbf{v}=0}). \quad (3.7)$$

In the co-moving frame, the spatial component of the 4-vector

$$\frac{\varepsilon\mu - 1}{2} \frac{c}{4\pi} \Omega_\alpha, \quad (3.8)$$

where  $\Omega_\alpha$  is the Minkowski vector (2.16), (2.17), reproduces the left-hand side of Eq. (3.7). Bearing this in mind, it is natural to add to the tensor (3.5) the symmetric tensor constructed



from two 4-vectors  $u_\alpha$  and  $(\varepsilon\mu - 1)\Omega_\beta$ ,

$$T_{\alpha\beta}^{(1)} = \frac{1}{2}(T_{\alpha\beta}^M + T_{\beta\alpha}^M) + \frac{1}{4\pi} \frac{\varepsilon\mu - 1}{2}(u_\alpha\Omega_\beta + u_\beta\Omega_\alpha). \quad (3.9)$$

Taking into account (1.1) and (2.17), one can easily verify that the tensor (3.9) in the rest frame ( $\mathbf{v} = 0$ ) and under the fulfillment the constitutive relations (3.1) satisfies the conditions (3.2)–(3.4).<sup>4</sup> The tensor (3.9) in view of (2.18) has vanishing trace.

Formally, the tensor (3.9) is defined on the whole configuration space  $\Gamma$ . However, in order to carry out the needed subtraction at  $\mathbf{v} = 0$  and fulfill condition (3.3), this tensor should be considered only on  $\bar{\Gamma}$ . Obviously, this restriction must also be kept in the case of moving medium ( $\mathbf{v} \neq 0$ ), in order to have a smooth dependence on  $\mathbf{v}$  of the tensor looking for.

Abraham proceeds exactly in this way. He defines the energy-momentum tensor on  $\bar{\Gamma}$  as the contraction of the tensor  $T_{\alpha\beta}^{(1)}$  on  $\bar{\Gamma}$  (see Ref. [11, p. 42]):

$$T_{\alpha\beta}^A \Big|_{\bar{\Gamma}} = T_{\alpha\beta}^{(1)} \Big|_{\bar{\Gamma}}. \quad (3.10)$$

The task to define the energy-momentum tensor on the whole configuration space  $\Gamma$  was not posed by Abraham.

By making use of Eqs. (2.10), (2.11), we can represent the 4-vector  $(\varepsilon\mu - 1)\Omega_\alpha$  on  $\bar{\Gamma}$  in the form [21, Eq. (106)]

$$(\varepsilon\mu - 1)\Omega_\alpha = u_\beta(D_\gamma B_{\alpha\beta\gamma} - E_\gamma H_{\alpha\beta\gamma}) \Big|_{\bar{\Gamma}}. \quad (3.11)$$

Further, it is worthwhile to use the property (2.18) and to express the 4-vector (3.11) in terms of one 3-dimensional vector  $\mathfrak{W}$  (Eq. (107) in Ref. [21]),

$$(\varepsilon\mu - 1)\boldsymbol{\Omega} = \frac{\mathfrak{W}}{\gamma}, \quad (\varepsilon\mu - 1)\Omega_4 = \frac{i}{\gamma}(\mathbf{q}\mathfrak{W}). \quad (3.12)$$

In the case of isotropic medium moving with the velocity  $\mathbf{v}$  all the components of the energy-momentum tensor (3.9), (3.10),

$$T_{\alpha\beta}^A = \begin{pmatrix} \sigma_{ij}^A & -ic\mathbf{g}^A \\ -\frac{i}{c}\mathbf{S}^A & w^A \end{pmatrix}, \quad (3.13)$$

---

<sup>4</sup> The subtraction needed is brought about due to the negative sign in front of  $\mathbf{S}^M$  and  $\mathbf{g}^M$  in (2.5).

were expressed by Abraham in terms of the field vectors  $\mathbf{E}$ ,  $\mathbf{H}$ ,  $\mathbf{D}$ ,  $\mathbf{B}$ , the velocity of the medium  $\mathbf{q}$ , and the vector  $\mathfrak{W}$  (Eqs. (24<sub>a,b,c</sub>) in Ref. [11])

$$\sigma_{ij}^A = \frac{1}{2}(\sigma_{ij}^M + \sigma_{ji}^M) + \frac{1}{8\pi}(q_i \mathfrak{W}_j + q_j \mathfrak{W}_i), \quad (3.14)$$

$$\frac{1}{c} \mathbf{S}^A = \frac{1}{8\pi} \{[\mathbf{EH}] + [\mathbf{DB}] - \mathfrak{W} - \mathbf{q}(\mathbf{q}\mathfrak{W})\} = c\mathbf{g}^A, \quad (3.15)$$

$$w^A = w^M - \frac{1}{4\pi}(\mathbf{q}\mathfrak{W}). \quad (3.16)$$

In turn, the vector  $\mathfrak{W}$ , defined in (3.12), is also expressed in terms of the field vectors and the velocity of the medium. Abraham (Ref. [11], p. 42) uses here the constitutive relations in the 3-dimensional vector form (2.14), (2.15). Kafka applies here the 4-dimensional notation (Ref. [21], pp. 48–51). As a preliminary, Kafka proves the relation (Eq. (119) in Ref. [21])

$$(\varepsilon\mu - 1)(u_\alpha \Omega_\beta - u_\beta \Omega_\alpha)|_{\bar{\Gamma}} = F_{\alpha\gamma} H_{\gamma\beta} - F_{\beta\gamma} H_{\gamma\alpha}|_{\bar{\Gamma}}, \quad (3.17)$$

which holds on  $\bar{\Gamma}$ .

We shall not reproduce here a rather cumbersome proof of this formula [21] and only note the following. On the left-hand side of Eq. (3.17), the dependence on  $\varepsilon\mu$  disappear as a result of using the constitutive relations (2.10), (2.11). When the pair of indices  $\alpha\beta$  in Eq. (3.17) assumes the values 14, 24, 34, the following equality arises here:

$$-i\{\mathfrak{W} - \mathbf{q}(\mathbf{q}\mathfrak{W})\} = -i\{[\mathbf{DB}] - [\mathbf{EH}]\}, \quad (3.18)$$

or in another form

$$\mathfrak{W} = [\mathbf{DB}] - [\mathbf{EH}] + \mathbf{q}(\mathbf{q}\mathfrak{W}). \quad (3.19)$$

From (3.19) it follows

$$\begin{aligned} (\mathbf{q}\mathfrak{W}) &= (\mathbf{q}, [\mathbf{DB}] - [\mathbf{EH}]) + q^2(\mathbf{q}\mathfrak{W}) = \\ &= \gamma^2(\mathbf{q}, [\mathbf{DB}] - [\mathbf{EH}]). \end{aligned} \quad (3.20)$$

Finally, we arrive at the result

$$\mathfrak{W} = [\mathbf{DB}] - [\mathbf{EH}] + \gamma^2 \mathbf{q}(\mathbf{q}, [\mathbf{DB}] - [\mathbf{EH}]). \quad (3.21)$$

The substitution of (3.21) into (3.15) and (3.16) yields

$$\frac{1}{c} \mathbf{S}^A = c\mathbf{g}^A = \frac{1}{4\pi}[\mathbf{EH}] - \frac{\mathbf{q}}{4\pi(1-q^2)}(\mathbf{q}, [\mathbf{DB}] - [\mathbf{EH}]), \quad (3.22)$$

$$w^A = \frac{1}{8\pi}(\mathbf{ED} + \mathbf{HB}) - \frac{1}{4\pi(1-q^2)}(\mathbf{q}, [\mathbf{DB}] - [\mathbf{EH}]). \quad (3.23)$$

Here we shall not write out the bulky formulae which are obtained as a result of substituting Eq. (3.21) into (3.14). Their explicit form is obvious. The energy-momentum tensor constructed in this way is symmetric.

Let us note once more that Eqs. (3.14)–(3.16) and (3.21)–(3.23) determine the components of the Abraham energy-momentum only on  $\bar{\Gamma}$  and do not include the material characteristics of the medium. The Minkowski constitutive relations (2.10), (2.11) or (2.14), (2.15) are used by Abraham specifically. Namely, they are not used directly to express one pair of field vectors from the set  $\mathbf{E}$ ,  $\mathbf{H}$ ,  $\mathbf{D}$ ,  $\mathbf{B}$  in terms of the rest pair, but special formulae are used which contain all the vectors  $\mathbf{E}$ ,  $\mathbf{H}$ ,  $\mathbf{D}$ ,  $\mathbf{B}$ ,  $\mathbf{v}$  and the material characteristics  $\varepsilon$ ,  $\mu$ , these formulae being valid only under fulfillment of the Minkowski constitutive relations, i.e., on  $\bar{\Gamma}$ . An important example of such a formula is the equality (3.17). When the pair of indices  $\alpha\beta$  in Eq. (3.17) takes the values 23, 31, and 12, one obtains the equality [21]

$$[\mathbf{q}\mathfrak{W}] = [\mathbf{ED}] + [\mathbf{HB}].$$

Taking into account (3.19), we deduce

$$[\mathbf{q}, [\mathbf{DB}] - [\mathbf{EH}]] = [\mathbf{ED}] + [\mathbf{HB}]. \quad (3.24)$$

It is obvious that this equality holds only under fulfillment of the Minkowski relations (2.14), (2.15).

Abraham deduced his formulae only on  $\bar{\Gamma}$ . Therefore, it is not clear whether 10 quantities defined in this way can be definitely considered as the components of a 4-dimensional tensor.<sup>5</sup> In order to verify this one has, first of all, to define the quantities under consideration on the whole configuration space  $\Gamma$ . This will be done in Sec. IV.

Taking advantage of formula (3.17) we can represent the tensor  $T_{\alpha\beta}^{(1)}$  (3.9), defined on  $\bar{\Gamma}$ , in another form. Let us write the definition (3.9) and formula (3.17), multiplied by 1/2, on the neighbouring lines,

$$4\pi T_{\alpha\beta}^{(1)} = \frac{1}{2} F_{\alpha\gamma} H_{\gamma\beta} + \frac{1}{2} F_{\beta\gamma} H_{\gamma\alpha} - \frac{1}{4} \delta_{\alpha\beta} F_{\gamma\delta} H_{\delta\gamma} + \frac{\varepsilon\mu - 1}{2} (u_\alpha \Omega_\beta + u_\beta \Omega_\alpha), \quad (3.25)$$

$$0 = \frac{1}{2} F_{\alpha\gamma} H_{\gamma\beta} - \frac{1}{2} F_{\beta\gamma} H_{\gamma\alpha} - \frac{\varepsilon\mu - 1}{2} (u_\alpha \Omega_\beta - u_\beta \Omega_\alpha). \quad (3.26)$$

---

<sup>5</sup> Appealing to the works [2, 11, 21], Abraham believed that the energy-momentum tensor, constructed by him, possesses correct properties under Lorentz transformations [9, p. 309].

Adding and subtracting the left-hand sides and the right-hand sides of these equations we obtain on  $\bar{\Gamma}$ ,

$$T_{\alpha\beta}^A = T_{\alpha\beta}^{(1)} = T_{\alpha\beta}^{(2)} = T_{\alpha\beta}^{(3)}, \quad (3.27)$$

where

$$T_{\alpha\beta}^{(2)} = T_{\alpha\beta}^M + \frac{\varepsilon\mu - 1}{4\pi} u_\beta \Omega_\alpha, \quad (3.28)$$

$$T_{\alpha\beta}^{(3)} = T_{\beta\alpha}^M + \frac{\varepsilon\mu - 1}{4\pi} u_\alpha \Omega_\beta. \quad (3.29)$$

If we use in (3.26) Eq. (3.17), multiplied by an arbitrary real number, then we obviously can continue to infinity the chain of equalities in (3.27), and consequently in the definition (3.10). However, the tensors  $T_{\alpha\beta}^{(i)}$ ,  $i = 2, 3, \dots$  constructed in this way give nothing new in comparison with  $T_{\alpha\beta}^{(1)}$ . On  $\bar{\Gamma}$ , they are equal to  $T_{\alpha\beta}^{(1)}$  and to the Abraham tensor according to (3.10), that is all these tensors are simply another form of representation for the Abraham tensor. In spite of the fact that these tensors represented in terms of vector  $\mathfrak{W}$  in accordance with (3.12), do not, in general case, coincide with Eqs. (3.14)–(3.16) and are not explicitly symmetric, but upon replacing  $\mathfrak{W}$  by (3.21), the components of these tensors are defined by (3.22)–(3.23) and Maxwell stress tensors also coincide.

Let us show this using the tensor  $T_{\alpha\beta}^{(2)}$ , which is, according to construction, the most close to the Minkowski energy-momentum tensor. The components of the tensor  $T_{\alpha\beta}^{(2)}$  are defined by Eqs. (3.12) and (3.28):

$$\sigma_{ij}^{(2)} = \sigma_{ij}^M + \frac{1}{4\pi} \mathfrak{W}_i q_j, \quad (3.30)$$

$$\frac{1}{c} \mathbf{S}^{(2)} = \frac{1}{4\pi} [\mathbf{EH}] + \frac{1}{4\pi} \mathbf{q}(\mathbf{q}\mathfrak{W}), \quad (3.31)$$

$$c\mathbf{g}^{(2)} = \frac{1}{4\pi} [\mathbf{DB}] - \frac{1}{4\pi} \mathfrak{W}, \quad (3.32)$$

$$w^{(2)} = w^M - \frac{1}{4\pi} (\mathbf{q}\mathfrak{W}). \quad (3.33)$$

It is this form that was used by Abraham for the representation of the components of his energy-momentum tensor in his book [9, §39, Eqs. (199e), (199d), (201), and (201a)]. Replacing in (3.31) and (3.32) the vector  $\mathfrak{W}$  by (3.21), we can easily obtain (3.22) and (3.23). The identity of the 3-dimensional Maxwell stress tensor represented in the forms (3.30) and (3.14) is proved by Abraham in his article [11, pp. 42, 43].

Pauli in his survey [1, Eq. (303)], does not note that the tensors  $T_{\alpha\beta}^{(i)}$ ,  $i = 1, 2, 3$  are equal to each other only on  $\bar{\Gamma}$  and that in the rest frame these tensors coincide only due to the

constitutive relations (3.1). In view of this the status of Eq. (303) in the Pauli treatment [1] was left unclear.<sup>6</sup>

The employment of the tensors  $T_{\alpha\beta}^{(i)}$ ,  $i = 1, 2, 3$  outside  $\bar{\Gamma}$ [4] is not physically justified because outside  $\bar{\Gamma}$  these tensors have no relation to the Abraham energy-momentum tensor. Indeed, outside  $\bar{\Gamma}$  these tensors are different and already due to this reason one cannot consider them as an extension of the Abraham formulae (3.14)–(3.16), (3.21)–(3.23) on the whole configuration space  $\Gamma$ . The consistent extension of the Abraham formulae on  $\Gamma$  will be implemented in Sec. IV of the present paper.

#### IV. THE SYMMETRIC ENERGY-MOMENTUM TENSOR

In the Introduction, it was explained that the term “symmetric energy-momentum tensor” implies the following. This tensor is defined by known values of its components in the rest frame [1, Eqs. (298) and (300)] and is determined on the whole configuration space  $\Gamma$ . In the present Section we construct this tensor generalizing the Abraham reasoning considered in the preceding Sec. III. This will enable us to prove, rather simple, the identity of the symmetric energy-momentum tensor and the Abraham tensor on  $\bar{\Gamma}$ . Let us recall that the term “Abraham tensor” we kept for Eqs. (3.13)–(3.16), (3.21)–(3.23) derived by Abraham himself only on  $\bar{\Gamma}$ .

It is obvious that conditions (3.2)–(3.4) enable us to define the symmetric energy-momentum tensor on  $\Gamma$  if we demand their fulfilment in the co-moving reference frame without using the constitutive relations (3.1). Below we demonstrate this.

In the generic inertial reference frame, tensor (3.5) yields the density of the energy flux,

$$\frac{c}{4\pi} \frac{1}{2} ([\mathbf{DB}] + [\mathbf{EH}]). \quad (4.1)$$

In order to get here Poynting vector (2.9), in agreement with condition (3.3), one must subtract from (4.1) the following quantity:

$$\frac{c}{4\pi} \frac{1}{2} ([\mathbf{DB}] - [\mathbf{EH}]). \quad (4.2)$$

Let us consider the 4-vector  $\tilde{\Omega}_\alpha$  which is obtained from the Minkowski vector  $\Omega_\alpha$  (2.16) by the substitution

$$\mathbf{E} \rightarrow \mathbf{D}, \quad \mathbf{H} \rightarrow \mathbf{B}, \quad \mathbf{D} \rightarrow \mathbf{E}, \quad \mathbf{B} \rightarrow \mathbf{H}. \quad (4.3)$$

---

<sup>6</sup> Pauli’s assertion [1] “The identity of three expressions (303) follows from their coincidence in the co-moving reference frame  $K'$ ” is simply wrong, being literally understood.

Using Eq. (2.16), one can construct for  $\tilde{\Omega}_\alpha$  the explicitly covariant formula,

$$\tilde{\Omega}_\alpha = B_{\alpha\beta\gamma} u_\beta D_\gamma. \quad (4.4)$$

In the co-moving reference frame, Eq. (4.4) gives

$$\frac{c}{4\pi} (\tilde{\Omega}'_1, \tilde{\Omega}'_2, \tilde{\Omega}'_3) = \frac{c}{4\pi} [\mathbf{D}'\mathbf{B}'], \quad \tilde{\Omega}'_4 = 0 \quad (4.5)$$

(compare with Eqs. (2.17), (2.9)).

Now it is clear that the 4-vector (3.11), which was used by Abraham in constructing the energy-momentum tensor only on  $\bar{\Gamma}$ , should be replaced by the difference

$$\tilde{\Omega}_\alpha - \Omega_\alpha = u_\beta (D_\gamma B_{\alpha\beta\gamma} - E_\gamma H_{\alpha\beta\gamma}) = \quad (4.6)$$

$$= F_{\alpha\nu} H_{\nu\lambda} u_\lambda - H_{\alpha\nu} F_{\nu\lambda} u_\lambda = \tilde{\omega}_\alpha - \omega_\alpha, \quad (4.7)$$

where

$$\omega_\alpha = H_{\alpha\nu} F_{\nu\lambda} u_\lambda, \quad \tilde{\omega}_\alpha = F_{\alpha\nu} H_{\nu\lambda} u_\lambda. \quad (4.8)$$

In place of  $T_{\alpha\beta}^{(1)}$  defined in (3.9), we have now, on the whole configuration space  $\Gamma$ , the tensor which we call, for definiteness, *the symmetric energy-momentum tensor*:

$$T_{\alpha\beta}^{\text{sym}} = \frac{1}{2} (T_{\alpha\beta}^{\text{M}} + T_{\beta\alpha}^{\text{M}}) + A_{\alpha\beta}, \quad (4.9)$$

where

$$A_{\alpha\beta} = A_{\beta\alpha} = \frac{1}{8\pi} \{u_\alpha (\tilde{\Omega}_\beta - \Omega_\beta) + u_\beta (\tilde{\Omega}_\alpha - \Omega_\alpha)\} = \quad (4.10)$$

$$= \frac{1}{8\pi} \{u_\alpha (F_{\beta\nu} H_{\nu\lambda} u_\lambda - H_{\beta\nu} F_{\nu\lambda} u_\lambda) + u_\beta (F_{\alpha\nu} H_{\nu\lambda} u_\lambda - H_{\alpha\nu} F_{\nu\lambda} u_\lambda)\}. \quad (4.11)$$

Taking into account Eqs. (2.17) and (4.5), one can easily verify that in the rest frame ( $\mathbf{v} = 0$ ) the tensor  $T_{\alpha\beta}^{\text{sym}}$  satisfies the conditions (3.2)–(3.4) without utilizing the constitutive relations (3.1).

As was noted in the Introduction, the symmetric energy-momentum tensor (4.9), (4.11) was constructed in recent papers [14, 15] (the approach used in [15] is discussed in Section V B of the present paper). The authors called it by the Abraham tensor but juxtaposition of Eqs. (4.9), (4.11) and Eqs. (3.14)–(3.16), (3.21)–(3.23) was not implemented in [14, 15]. In our approach, the coincidence of these tensors on  $\bar{\Gamma}$  follows immediately from the comparison of (4.6) and (3.11). In the analytical way, it is proved as follows. Taking advantage of

relation (3.17), valid on  $\bar{\Gamma}$  and being read from right to left, and using property (2.18) of the Minkowski vector  $\Omega_\alpha$ , Equation (4.11) can be transformed to the form

$$A_{\alpha\beta}|_{\bar{\Gamma}} = \frac{1}{4\pi} \frac{\varepsilon\mu - 1}{2} (u_\alpha \Omega_\beta + u_\beta \Omega_\alpha)|_{\bar{\Gamma}}. \quad (4.12)$$

Keeping in mind Eqs. (4.9), (3.9), and (3.10), we obtain

$$T_{\alpha\beta}^{\text{sym}}|_{\bar{\Gamma}} = T_{\alpha\beta}^{(1)}|_{\bar{\Gamma}} = T_{\alpha\beta}^{\text{A}}|_{\bar{\Gamma}}. \quad (4.13)$$

Below we prove equality (4.13) once more by making use of the 3-dimensional vector formula (3.24) in place of (3.17).

When the energy-momentum tensor applies to practical calculations, the explicit form of its components is, as a rule, needed in terms of the 3-dimensional vector notation. We turn now to the construction of such formulae for the symmetric energy-momentum tensor on  $\Gamma$ .

First we derive the 3-dimensional vector representation for the auxiliary 4-vectors  $\omega_\alpha$  and  $\tilde{\omega}_\alpha$ , introduced in Eqs. (4.7) and (4.8). From (1.1) and (2.12) it follows

$$\omega_\alpha = \gamma\{[\mathbf{EH}] - [\mathbf{H}[\mathbf{qB}]] + \mathbf{D}(\mathbf{qE}); \mathbf{i}(\mathbf{ED}) - \mathbf{i}(\mathbf{q}[\mathbf{DB}])\}. \quad (4.14)$$

Obviously, the 4-vector  $\tilde{\omega}_\alpha$  is derived from  $\omega_\alpha$  by substitution (4.3),

$$\tilde{\omega}_\alpha = \gamma\{[\mathbf{DB}] - [\mathbf{B}[\mathbf{qH}]] + \mathbf{E}(\mathbf{qD}); \mathbf{i}(\mathbf{ED}) - \mathbf{i}(\mathbf{q}[\mathbf{EH}])\}. \quad (4.15)$$

The difference  $\tilde{\omega}_\alpha - \omega_\alpha$  becomes

$$\begin{aligned} \tilde{\omega}_\alpha - \omega_\alpha = \gamma\{ & [\mathbf{DB}] + \mathbf{E}(\mathbf{qD}) + \mathbf{H}(\mathbf{qB}) - [\mathbf{EH}] - \mathbf{D}(\mathbf{qE}) - \mathbf{B}(\mathbf{qH}); \\ & \mathbf{i}(\mathbf{q}[\mathbf{DB}]) - \mathbf{i}(\mathbf{q}[\mathbf{EH}])\}. \end{aligned} \quad (4.16)$$

By analogy with (3.12) it is convenient to represent the difference (4.6), (4.7), and (4.16) in the following form:

$$\tilde{\Omega} - \Omega = \frac{\mathbf{W}}{\gamma}, \quad \tilde{\Omega}_4 - \Omega_4 = \frac{\mathbf{i}}{\gamma}(\mathbf{qW}), \quad (4.17)$$

where the vector  $\mathbf{W}$  is

$$\mathbf{W} = \gamma^2\{[\mathbf{DB}] - [\mathbf{EH}] + \mathbf{E}(\mathbf{qD}) + \mathbf{H}(\mathbf{qB}) - \mathbf{B}(\mathbf{qH}) - \mathbf{D}(\mathbf{qE})\}. \quad (4.18)$$

Equation (4.18) yields in particular

$$(\mathbf{qW}) = \gamma^2\{(\mathbf{q}[\mathbf{DB}]) - (\mathbf{q}[\mathbf{EH}])\}. \quad (4.19)$$

Let us introduce the standard notation for the components of the tensor  $T_{\alpha\beta}^{\text{sym}}$  in a generic inertial reference frame  $K$ , where the medium velocity is  $\mathbf{v}$ ,

$$T_{\alpha\beta}^{\text{sym}} = \begin{pmatrix} \sigma_{ij}^{\text{sym}} & -i c \mathbf{g}^{\text{sym}} \\ -\frac{i}{c} \mathbf{S}^{\text{sym}} & w^{\text{sym}} \end{pmatrix}. \quad (4.20)$$

Equations (4.9), (4.10), and (4.17) yield for the individual components of the symmetric tensor (4.20) the following 3-dimensional vector formulae:

$$\sigma_{ij}^{\text{sym}} = \frac{1}{2} (\sigma_{ij}^{\text{M}} + \sigma_{ji}^{\text{M}}) + \frac{1}{8\pi} (q_i W_j + q_j W_i), \quad (4.21)$$

$$\frac{1}{c} \mathbf{S}^{\text{sym}} = \frac{1}{8\pi} \{ [\mathbf{E}\mathbf{H}] + [\mathbf{D}\mathbf{B}] - \mathbf{W} - \mathbf{q}(\mathbf{q}\mathbf{W}) \} = c \mathbf{g}^{\text{sym}}, \quad (4.22)$$

$$w^{\text{sym}} = w^{\text{M}} - \frac{1}{4\pi} (\mathbf{q}\mathbf{W}). \quad (4.23)$$

These formulae are also obtained by the substitution  $\mathfrak{W} \rightarrow \mathbf{W}$  in analogous equations (3.14)–(3.16) for the Abraham tensor. Equations (4.21)–(4.23) and (4.18) determine, in an explicit form, the symmetric energy-momentum tensor (4.20) on the whole configuration space  $\Gamma$ .

Comparing Eqs. (3.20) and (4.19), we infer that the energy density given by the symmetric energy-momentum tensor (4.23) and by the Abraham tensor (3.16) are equal to each other on the whole configuration space  $\Gamma$  and are defined by (3.23),

$$w^{\text{sym}} = w^{\text{A}} = \frac{1}{8\pi} (\mathbf{E}\mathbf{D} + \mathbf{H}\mathbf{B}) - \frac{1}{4\pi(1-q^2)} (\mathbf{q}, [\mathbf{D}\mathbf{B}] - [\mathbf{E}\mathbf{H}]). \quad (4.24)$$

In the general case, the remaining components of these tensors, by virtue of (3.21) and (4.18), do not coincide outside  $\bar{\Gamma}$ . However, when the problem under consideration allows to direct one of the coordinate axes (for example, the  $x$ -axis) along the medium velocity  $\mathbf{v}$ ,

$$\mathbf{q} = (q_x, 0, 0), \quad (4.25)$$

then some other components of these tensors will also equal on  $\Gamma$ , namely, the diagonal components of the Maxwell stress tensors,

$$\sigma_{xx}^{\text{sym}} = \sigma_{xx}^{\text{A}}, \quad \sigma_{yy}^{\text{sym}} = \sigma_{yy}^{\text{A}}, \quad \sigma_{zz}^{\text{sym}} = \sigma_{zz}^{\text{A}} \quad (4.26)$$

and the components

$$\sigma_{yz}^{\text{sym}} = \sigma_{zy}^{\text{sym}} = \sigma_{yz}^{\text{A}} = \sigma_{zy}^{\text{A}}. \quad (4.27)$$



The first equality in (4.26) is the consequence of the relation

$$\mathbf{W}_x = \gamma^2 \{ [\mathbf{DB}]_x - [\mathbf{EH}]_x \} = \mathfrak{W}_x, \quad (4.28)$$

which holds under condition (4.25) (see Eqs. (4.18) and (3.21)). The remaining equalities in (4.26) and equalities (4.27) follow from (4.25).

Now we prove that the 3-dimensional vectors  $\mathbf{W}$  (4.18) and  $\mathfrak{W}$  (3.21) equal on  $\bar{\Gamma}$ . For this purpose, we construct the vector product of  $\mathbf{q}$  and the relation (3.24) valid on  $\bar{\Gamma}$ ,

$$\begin{aligned} [\mathbf{q}[\mathbf{q}([\mathbf{DB}] - [\mathbf{EH}])]] &= \mathbf{q}(\mathbf{q}, [\mathbf{DB}] - [\mathbf{EH}]) - q^2([\mathbf{DB}] - [\mathbf{EH}]) = \\ &= [\mathbf{q}, [\mathbf{ED}]] + [\mathbf{q}[\mathbf{HB}]] = \mathbf{E}(\mathbf{qD}) - \mathbf{D}(\mathbf{qE}) + \mathbf{H}(\mathbf{qB}) - \mathbf{B}(\mathbf{qH}). \end{aligned} \quad (4.29)$$

The substitution of (4.29) into (4.18) yields

$$\begin{aligned} \mathbf{W} &= \gamma^2 \{ [\mathbf{DB}] - [\mathbf{EH}] - q^2([\mathbf{DB}] - [\mathbf{EH}]) + \mathbf{q}(\mathbf{q}, [\mathbf{DB}] - [\mathbf{EH}]) \} = \\ &= [\mathbf{DB}] - [\mathbf{EH}] + \gamma^2 \mathbf{q}(\mathbf{q}, ([\mathbf{DB}] - [\mathbf{EH}])) = \mathfrak{W}. \end{aligned} \quad (4.30)$$

It implies that the symmetric energy-momentum tensor (4.20)–(4.23) and the Abraham tensor (3.13)–(3.16) are identical on  $\bar{\Gamma}$ . In other words, this inference may be formulated like this: the correct extension of the Abraham formulae (3.13)–(3.16) onto the whole configuration space  $\Gamma$  is accomplished by symmetric energy-momentum tensor (4.20)–(4.23). Obviously, this extension is unique as the tensor  $T_{\alpha\beta}^{\text{sym}}$  is defined by its value in  $K'$  (3.2)–(3.4) uniquely.

Closing this section, we make an important remark. In practical calculations, aimed at obtaining the final physical result, it is obligatory assumed to use Minkowski constitutive relations (2.10), (2.11) or (2.14), (2.15). Therefore, in such studies it makes sense to utilize, from the very beginning, the energy-momentum tensor in the Abraham form as the respective formulae are substantially more compact in comparison with those given by the symmetric tensor. This concerns the density of the energy current, the linear momentum density (3.22), and especially the components of the 3-dimensional Maxwell stress tensor in the form (3.30).

## V. FORM-INVARIANCE OF THE VECTOR FORMULAE FOR THE ENERGY-MOMENTUM TENSOR

### A. The methods of defining the Minkowski tensor and the form-invariance of the vector formulae involved

In this subsection, we compare two methods of defining the 4-dimensional tensor: in the explicitly covariant form and in the 3-dimensional vector form. As an example, we shall use the Minkowski energy-momentum tensor. In this case one can clearly show the covariance properties of the 3-dimensional vector formulae for the Minkowski tensor, which result in their form-invariance.

#### 1. Definition of the Minkowski tensor by equation (2.4)

First we consider the definition of the Minkowski tensor by 4-dimensional explicitly covariant expression (2.4), and, in addition, we postulate that the latter is valid both for the medium at rest and for the moving medium. It is worth emphasizing that the velocity of the medium does not enter Eq. (2.4). In terms of the 3-dimensional vector notation the components of the tensor  $T_{\alpha\beta}^{\text{M}}$  (2.4), in the arbitrary inertial reference frame  $K$ , are given by Eqs. (2.5)–(2.8).

It is obvious that all the components of the Minkowski tensor (2.5), expressed in terms of the field vectors  $\mathbf{E}, \mathbf{H}, \mathbf{D}, \mathbf{B}$  according to Eqs. (2.6)–(2.8), keep their functional dependence, specified by these formulae, in a generic inertial reference frame. Indeed, when passing from Eq. (2.4) to Eqs. (2.5)–(2.8) the reference frame was not fixed or specified. This assertion is true both for the medium at rest and for the moving medium since the velocity of medium enter neither explicitly covariant formula (2.4) nor the consequent vector formulae (2.5)–(2.8).

Thus the transformation of the tensor  $T_{\alpha\beta}^{\text{M}}$ , defined in a generic reference frame  $K$  originally by an explicitly covariant formula (2.4) and afterwards written in the vector form (2.5)–(2.8), to a new inertial reference frame,  $K''$ , is simply reduced to the substitution  $\mathbf{F} \rightarrow \mathbf{F}''$  in Eqs. (2.6)–(2.8),

$$T_{\alpha\beta}^{\prime\prime\text{M}}(\mathbf{F}'') = T_{\alpha\beta}^{\text{M}}(\mathbf{F}''), \quad (5.1)$$

where  $\mathbf{F}$  means the set of the field vectors,

$$\mathbf{F} = \{\mathbf{E}, \mathbf{H}, \mathbf{D}, \mathbf{B}\}. \quad (5.2)$$

As before, the field vectors without prime belong to an arbitrary inertial reference frame  $K$  and the vectors with two primes pertain to analogous frame  $K''$ . This form-invariance property of the 3-dimensional vector formulae (2.5)–(2.8) is a straightforward consequence of the relativistic covariance of definition (2.4).

Certainly, the numerical values of the individual components of the tensor, calculated by Eqs. (2.5)–(2.8) at respective points  $r_\alpha = (x, y, z, ict)$  and  $r'' = (x'', y'', z'', ict'')$ , vary in accordance with the transformation law for the tensor

$$T''^{\text{M}}_{\alpha\beta}(\mathbf{F}''(r''_\gamma)) = \Lambda_{\alpha\delta}\Lambda_{\beta\rho}T^{\text{M}}_{\delta\rho}(\mathbf{F}(r_\sigma)), \quad (5.3)$$

where  $\Lambda_{\alpha\beta}$  is the matrix of the Lorentz transformation connecting the points  $r''_\alpha$  and  $r_\beta$ ,

$$r''_\alpha = \Lambda_{\alpha\beta}r_\beta. \quad (5.4)$$

The assertion above, concerning the Lorentz transformation of the components of the tensor  $T^{\text{M}}_{\alpha\beta}$  in the vectorial form, holds obviously for an arbitrary tensor of any rank.

## 2. Definition of the Minkowski tensor by the vector equations (2.5)–(2.8)

The Minkowski energy-momentum tensor, as a generic tensor, can be uniquely defined by fixing its components in a specified reference frame. Let us assume that in the co-moving frame  $K'$ , where the medium is at rest, the components of the tensor  $T'^{\text{M}}_{\alpha\beta}$  are defined by vector formulae (2.5)–(2.8). All the vectors and their components in these formulae should be provided with the prime, in accord with our notation. In order to find the components of the tensor under consideration in an arbitrary inertial reference frame we can proceed in two ways: (i) to construct the 4-dimensional explicitly covariant formula, which reproduces Eqs. (2.5)–(2.8) in the reference frame  $K'$ , for example, generalizing the energy-momentum tensor in the vacuum (2.1) and (2.4); (ii) by making use of Eq. (5.3) to transform the 3-dimensional Eqs. (2.5)–(2.8) to the generic reference frame  $K$ . Here we cannot take advantage of the form-invariance (5.1) since Eqs. (2.5)–(2.8), according to the task posing, define the Minkowski tensor not in an arbitrary reference frame  $K$ , but in the co-moving frame  $K'$ .

The first way was, in fact, traced above (Section V A 1). In order to apply here the Lorentz transformations one has to resort to Eq. (5.2) which in the case under consideration becomes

$$T_{\alpha\beta}^{\text{M}}(\mathbf{F}(r_\gamma)) = \Lambda_{\alpha\delta}\Lambda_{\beta\varrho}T'_{\delta\varrho}{}^{\text{M}}(\mathbf{F}(r'_\sigma)). \quad (5.5)$$

According to the conditions of the task, the components of the tensor  $T'_{\delta\varrho}{}^{\text{M}}(\mathbf{F}')$  in (5.5) are given by the vectorial equations (2.5)–(2.8). Further one has to express the field vectors  $\mathbf{F}'$  on the right-hand side of Eq. (5.5) in terms of  $\mathbf{F}$ . Having in mind that for Eqs. (2.5)–(2.8) there exist 4-dimensional explicitly covariant representation (2.4), one can predict beforehand the result of these transformations<sup>7</sup>,

$$T_{\alpha\beta}^{\text{M}}(\mathbf{F}(r_\gamma)) = T'_{\alpha\beta}{}^{\text{M}}(\mathbf{F}(r_\gamma)). \quad (5.6)$$

Nevertheless, in a recent paper [22], these transformations were carried out for the component  $T_{xx}^{\text{M}}$  and its form-invariance was shown.<sup>8</sup> This property of the Minkowski tensor was treated in Ref. [22] not completely correct, as the manifestation of *the relativistic invariance* of this tensor (see critical note in [15]). As was explained above, this form-invariance is in fact the consequence of *the covariance property* of individual components of generic tensor written *in vector notation*.

In pertinent literature [20], the form-invariance of the vectorial formulae, determining the components of the 4-dimensional vectors and tensors in the theory of electromagnetic field is not considered, probably owing to clearness of this property. Nevertheless, we believe that detailed discussion of this point presented above will be useful.

## B. Form-invariance of the symmetric energy-momentum tensor and the Abraham tensor in the vector notation

For these tensors, we have explicitly covariant 4-dimensional representations (4.9), (4.10) and (3.9), (3.10). Therefore, 3-dimensional vectorial Eqs. (4.20)–(4.23), (4.18) and (3.13)–(3.16), (3.21) or (3.22), (3.23) derived from them preserve the form of functional dependence

<sup>7</sup> The employment of the Lorentz transformations in construction of the symmetric energy-momentum tensor is discussed in Section V B.

<sup>8</sup> In Ref. [22] the co-moving frame was denoted by  $K$  and generic inertial frame by  $K'$ . The component  $T_{xx}^{\text{M}}$ , defined in  $K$  by Eq. (2.6) (present paper), was transformed into the reference frame  $K'$ . It was shown that  $T'_{xx}{}^{\text{M}}(\mathbf{F}') = T_{xx}^{\text{M}}(\mathbf{F})$ .

on field vectors  $\mathbf{F}$  (5.2) and velocity of medium  $\mathbf{v}$  when passing from one inertial reference frame to another. The transformation rule (5.1) generalized to this case reads

$$T_{\alpha\beta}''^{\text{sym}}(\mathbf{F}'', \mathbf{v}'') = T_{\alpha\beta}^{\text{sym}}(\mathbf{F}'', \mathbf{v}''), \quad (5.7)$$

$$T_{\alpha\beta}''^{\text{A}}(\mathbf{F}'', \mathbf{v}'') = T_{\alpha\beta}^{\text{A}}(\mathbf{F}'', \mathbf{v}''). \quad (5.8)$$

This rule enables one to find the tensors  $T_{\alpha\beta}''^{\text{sym}}(\mathbf{F}'', \mathbf{v}'')$  and  $T_{\alpha\beta}''^{\text{A}}(\mathbf{F}'', \mathbf{v}'')$  in an generic inertial reference frame  $K''$ , provided the reference frame  $K$  with tensors  $T_{\alpha\beta}^{\text{sym}}(\mathbf{F}, \mathbf{v})$  and  $T_{\alpha\beta}^{\text{A}}(\mathbf{F}, \mathbf{v})$  is a reference frame of *a generic type*, in particular,  $\mathbf{v} \neq 0$ , i.e.,  $K$  is not the co-moving frame.<sup>9</sup> Hence, the transformation rules (5.7) and (5.8) do not allow one to find the symmetric energy-momentum tensor and the Abraham tensor in arbitrary reference frame  $K''$  proceeding from Eqs. (3.2)–(3.4) for these tensors in the rest frame  $K'$  with  $\mathbf{v} = 0$  (without allowance for the constitutive relations (3.1) and with this allowance, respectively). Here the method must be used, which was employed in Sections III and IV, or one has to resort to the Lorentz transformations (5.5). The latter became in the case of the tensor  $T_{\alpha\beta}^{\text{sym}}$

$$T_{\alpha\beta}^{\text{sym}}(\mathbf{F}(r_\gamma)) = \Lambda_{\alpha\delta}\Lambda_{\beta\rho}T_{\delta\rho}'^{\text{sym}}(\mathbf{F}'(r'_\sigma)), \quad (5.9)$$

where  $\Lambda_{\alpha\beta}$  is the matrix of the Lorentz transformation connecting the points  $r_\alpha$  and  $r'_\delta$ ,

$$r_\alpha = \Lambda_{\alpha\delta}r'_\delta. \quad (5.10)$$

As before, the variables in an arbitrary inertial reference frame  $K$  are not marked by prime, and primed variables are referred to the co-moving frame  $K'$ . Obviously, the velocity of generic inertial reference frame  $K$  with respect to the rest frame  $K'$  is  $-\mathbf{v}$ , where  $\mathbf{v}$  is the velocity of the medium in  $K'$ . Hence, the matrix of the Lorentz transformation in (5.10) is defined by the velocity  $-\mathbf{v}$ :  $\Lambda_{\alpha\beta} = \Lambda_{\alpha\beta}(-\mathbf{v})$ .

The tensor  $T_{\alpha\beta}'^{\text{sym}}(\mathbf{F}')$  in the rest frame  $K'$  is specified by Eqs. (3.2)–(3.4). Further the field vectors  $\mathbf{F}'$  on the right-hand side of Eq. (5.9) must be expressed in terms of the field vectors  $\mathbf{F}$  in the generic inertial reference frame  $K$ . The latter is implemented by the transformation  $\Lambda(+\mathbf{v})$ .

The Lorentz transformations, considered above, were carried out in [15] and, what is more, for all the components of the tensor  $T_{\alpha\beta}^{\text{sym}}$  at once and for arbitrary direction of the medium velocity with respect to the coordinate axes.

---

<sup>9</sup> One cannot agree with the assertion, [13] that Abraham declined the form-invariance principle when constructing his energy-momentum tensor.

For a single component of the symmetric energy-momentum tensor  $T_{xx}^{\text{sym}} = \sigma_{xx}^{\text{sym}}$  the analogous transformations were in fact performed in the work [22], the  $x$ -axis being directed along the velocity of the medium  $\mathbf{v}$ ,

$$\mathbf{v} = (v_x, 0, 0). \quad (5.11)$$

The authors of Ref. [22] erroneously interpreted the obtained dependence of the component  $T_{xx}^{\text{sym}}$  on  $\mathbf{v}$  as an indication that this energy-momentum tensor “is not relativistically invariant”. As a matter of fact, in that paper the exact expression was derived for the  $xx$ -component of the symmetric energy-momentum tensor in a generic inertial reference frame, where the medium has the velocity  $v_x/c = -\beta$ . One can be easily convinced of this by comparing Equation (23) in the work [22], preliminarily changing the sign of  $\beta$ , with the  $xx$ -component of the Abraham tensor in the form  $T_{\alpha\beta}^{(2)}$  (3.28), (3.30)–(3.33). This tensor, borrowed from Ref. [9], is presented in the Appendix 4 of the survey [5] under the condition (5.11). Here one has to bear in mind the following. The diagonal elements of the tensor  $T_{\alpha\beta}^{(2)}$ , Eq. (3.28), and of the tensor  $T_{\alpha\beta}^A$ , Eqs. (3.10), (3.14)–(3.16), obviously coincide on the whole configuration space  $\Gamma$ . Further, when condition (4.25) is valid, equalities (4.26) and (4.27) hold. Thus,  $\sigma_{xx}^{(2)} = \sigma_{xx}^{\text{sym}}$  on  $\Gamma$ . In Ref. [15, p. 1361, Russian edition] it was explained in what way the setting of the problem in the work [22] should be supplemented in order to get the correct inference given above. However, this inference was not formulated in Ref. [15].

## VI. CONCLUSION

Let us summarize briefly the results obtained in our paper.

1. The Abraham energy-momentum tensor (3.13), (3.30)–(3.33) and the covariant Grammel formulae (3.10), (3.9), (3.27)–(3.29) for this tensor are defined only on  $\bar{\Gamma}$ , where they coincide and do not depend on the material characteristics of the medium  $\varepsilon$  and  $\mu$ .
2. Employment of the Abraham and Grammel formulae outside  $\bar{\Gamma}$  is not justified physically.
3. The generalization of the Abraham reasoning enables us to construct the symmetric energy-momentum tensor on the whole configuration space  $\Gamma$ .

4. The symmetric energy-momentum tensor coincides with the Abraham tensor on  $\bar{\Gamma}$  and provides the unique relativistically covariant extension of the Abraham tensor on the whole configuration space  $\Gamma$ .
5. In practical calculations, the energy-momentum tensor can be used in the Abraham form which is more compact in comparison with the symmetric tensor.
6. The vectorial 3-dimensional formulae for the components of the symmetric energy-momentum tensor have the same form (functional dependence) in the arbitrary inertial reference frame, i.e., these formulae are form-invariant. The same statement is also true for the Abraham tensor.

The physical basis of the Abraham tensor and the symmetric energy-momentum tensor is the definition of the linear momentum density of electromagnetic field in a medium, which is postulated, in the rest frame, in the form (3.3). Just this definition is the physical reason which distinguishes the Abraham approach from the Minkowski tensor and other versions of the electromagnetic energy-momentum tensor in the medium. The symmetry property of the Abraham tensor (3.13)–(3.16) and of the tensor (4.9)–(4.11), as well as their explicit dependence on the velocity of the medium, are, really, the consequences of the condition (3.3).

On the face of it, one may infer from (4.9) that these properties, symmetry, and explicit dependence on the medium velocity, are implemented independently of each other. Indeed, the symmetric tensor is already produced by the first two terms in (4.9),<sup>10</sup> and to take into account the dependence on the velocity of the medium, one has to add one more term  $A_{\alpha\beta}(u) = A_{\beta\alpha}(u)$ . As a matter of fact, it is not the case. Namely, the symmetrization alone, without introducing an explicit dependence on  $u_\alpha$ , does not enable one to meet on  $\Gamma$  conditions (3.2)–(3.4) determining the tensor  $T_{\alpha\beta}^{\text{sym}}$ .

Not numerous references, concerning the Abraham tensor, are, practically all, enumerated in the preset paper [1, 9–11]. Unfortunately, in the textbooks the Abraham energy-momentum tensor is, as a rule, not considered.

Until now, the problem of defining the energy-momentum tensor of electromagnetic field in the medium has been considered substantially for the medium at rest. However the

---

<sup>10</sup> The symmetrized Minkowski energy-momentum tensor is used in practical calculations [23].

complete solution of the problem in question demands also determination of the dependence of this tensor on the medium velocity. One may hope that precise and comprehensive consideration of the mathematical aspects relating to the Abraham approach will be helpful for these investigations.

## ACKNOWLEDGMENTS

VVN thanks I. Brevik for useful advices on the subject under study, A.A. Rukhadze for providing the reference to the paper [15], and Yu.N. Obukhov for references [12, 14].

- 
- [1] W. Pauli, *Theory of Relativity* (New York: Pergamon Press, 1958) §§ 33, 35; [W. Pauli, *Relativitätstheorie* (Leipzig: Teubner, 1921) §§ 33, 35].
  - [2] R. Grammel, *Ann. Physik* **346**, 570 (1913).
  - [3] V.L. Ginzburg and V.A. Ugarov, *Sov. Phys. Usp.* **19**, 94 (1976) [*Usp. Fiz. Nauk* **118**, 175 (1976)].
  - [4] U. Leonhardt, *Phys. Rev. A* **73**, 032108 (2006).
  - [5] D.V. Skobel'tsyn, *Sov. Phys. Usp.* **16**, 381 (1973) [*Usp. Fiz. Nauk* **110**, 253 (1973)].
  - [6] I. Brevik, *Mat. Phys. Medd. Dan. Vid. Selsk.* **37**, No. 11, 1 (1970); **37**, No. 13, 1 (1970).
  - [7] I. Brevik, *Phys. Rep.* **52**, 133 (1979).
  - [8] V.L. Ginzburg, *Sov. Phys. Usp.* **16**, 434 (1973) [*Usp. Fiz. Nauk* **110**, 309 (1973)].
  - [9] M. Abraham, *Theorie der Elektrizität*, Bd. 2, 4. Aufl. (Leipzig: Teubner, 1920).
  - [10] M. Abraham, *Palermo Rend.* **28**, 1 (1909); [[http://en.wikisource.org/wiki/Author:Max\\_Abraham](http://en.wikisource.org/wiki/Author:Max_Abraham)]
  - [11] M. Abraham, *Palermo Rend.* **30**, 33 (1910); [[http://en.wikisource.org/wiki/Author:Max\\_Abraham](http://en.wikisource.org/wiki/Author:Max_Abraham)]
  - [12] E. Schmutzer, *Relativistische Physik* (Leipzig: Akademie Verlag, 1968) p. 408.
  - [13] S.R. de Groot and L.G. Suttorp, *Foundations of Electrodynamics* (Amstredam: North-Holland, 1972), Ch. V, §7.
  - [14] Yu.N. Obukhov, *Ann. Phys. (Berlin)*, **17**, No. 9/10, 830 (2008).
  - [15] V.P. Makarov and A.A. Rukhadze, *Phys. Usp.* **52**, 937 (2011) [*Usp. Fiz. Nauk* **181**, 1357 (2011)].
  - [16] J. Gratus, Yu.N. Obukhov, and R.N. Tucker, *Annals Phys.* **327**, 2560 (2012).



- [17] T. Ramos, G.F. Guillermo, Yu.N. Obukhov, Phys. Lett. A **375**, 1703 (2011).
- [18] T. Ramos, G.F. Guillermo, Yu.N. Obukhov, J. Opt. **17**, 025611 (2015).
- [19] L.D. Landau and E.M. Lifshitz, *Electrodynamics of Continuous Media* (Oxford: Pergamon Press, 1984), §76.
- [20] L.D. Landau and E.M. Lifshitz, *The Classical Theory of Fields* (Oxford: Pergamon Press, 1980).
- [21] H. Kafka, Ann. Physik **58**, 1 (1919).
- [22] V.G. Veselago and V.V. Shchavlev, Phys. Usp. **53**, 317 (2010) [Usp. Fiz. Nauk **180**, 331 (2010)].
- [23] S. Goto, R.W. Tucker, and T.J. Walton, Proc. R. Soc. A **467**, 59 (2011); **467**, 79 (2011).