Dynamics of ellipses inscribed in quadrilaterals

Alan Horwitz Professor Emeritus of Mathematics Penn State Brandywine 25 Yearsley Mill Rd. Media, PA 19063 alh4@psu.edu

5/5/15

Abstract

Let Q be a convex quadrilateral in the xy plane, let int(Q) denote the interior of Q, and let $\partial(Q)$ denote the boundary of Q. Let D_1 and D_2 denote the diagonals of Q and let P denote their point of intersection. Let $P_0 = (x_0, y_0)$ be a point in $\overline{Q} = int(Q) \cup \partial(Q)$. We prove the following:

(i) If $P_0 \in int(Q), P_0 \notin D_1 \cup D_2$, then there are exactly two ellipses inscribed in Q which pass through P_0 .

(ii) If $P_0 \in int(Q)$ and $P_0 \in D_1 \cup D_2$, but $P_0 \neq P$, then there is exactly one ellipse inscribed in Q which passes through P_0 .

(iii) There is no ellipse inscribed in Q which passes through P.

(iv) If $P_0 \in \partial(Q)$, but P_0 is not one of the vertices of Q, then there is exactly one ellipse inscribed in Q which passes through P_0 (and is thus tangent to Q at one of its sides).

Introduction

Suppose that we are given a point, P_0 , in the interior of a convex quadrilateral, Q, in the xy plane. Is there an ellipse, E, inscribed in Q which also passes through P_0 ? If yes, how many such ellipses? By inscribed in Q we mean that E lies in Q and is tangent to each side of Q. Looked at in a dynamic sense: Imagine a particle constrained to travel along the path of an ellipse inscribed in a convex quadrilateral, Q. Thus the particle bounces off each side of Q along its path. Of course there are infinitely many such paths. Can we also specify a point in Q that the particle must pass through ? If yes, is such a path then unique ? We show below(Theorem 1.1) that the path is unique when P_0 lies on one of the diagonals of Q(but does not equal their intersection point), while there are two such paths if P_0 does not lie on one of the diagonals of Q. Finally, if P_0 equals the intersection point of the diagonals of Q, then no ellipse inscribed in Q passes through P_0 . We also prove that there is a unique ellipse inscribed in Q which is tangent a given point on the boundary of Q, assuming, of course, that that point is not one of the vertices of Q. Using affine invariance, it suffices to prove Theorem 1.1 when Q is the convex quadrilateral with vertices (0,0), (0,1), (1,0), and (s,t), where $s > 0, t > 0, s + t > 1, s \neq 1 \neq t$. The proof depends heavily on the general equation of an ellipse inscribed in Qgiven in Proposition 3.1 below. It would be interesting to try to extend some of the results of Theorem 1.1 to other families of simple closed convex curves inscribed in a convex quadrilateral. It seems difficult, though, to come up with the general equation of such families of curves as we do with ellipses. For a paper somewhat similar to this one, but involving ellipses inscribed in triangles, see [3].

1 Main Result

Theorem 1.1 Let Q be a convex quadrilateral in the xy plane, let int(Q) denote the interior of Q, and let $\partial(Q)$ denote the boundary of Q. Let D_1 and D_2 denote the diagonals of Q and let P denote their point of intersection. Let $P_0 = (x_0, y_0)$ be a point in $\overline{Q} = int(Q) \cup \partial(Q)$.

(i) If $P_0 \in int(Q), P_0 \notin D_1 \cup D_2$, then there are exactly two ellipses inscribed in Q which pass through P_0 .

(ii) If $P_0 \in int(Q)$ and $P_0 \in D_1 \cup D_2$, but $P_0 \neq P$, then there is exactly one ellipse inscribed in Q which passes through P_0 .

(iii) There is no ellipse inscribed in Q which passes through P.

(iv) If $P_0 \in \partial(Q)$, but P_0 is not one of the vertices of Q, then there is exactly one ellipse inscribed in Q which passes through P_0 (and is thus tangent to Q at one of its sides).

Remark 1.1 Instead of just looking at the class of ellipses inscribed in convex quadrilaterals, Q, one might see whether the results of Theorem 1.1 still hold for other families of simple closed convex curves inscribed in Q. For example, one could start with $x^n + y^n = 1$ and apply all nonsingular affine transformations to generate such a family. It seems clear geometrically (we do not have a rigorous proof) that there is no simple closed convex curve inscribed in Q. So Theorem 1.1(iii) would still hold for any family of simple closed convex curves inscribed in Q. However, the other parts of Theorem 1.1 would not necessarily hold since they depend on that particular family of simple closed convex curves.

By Theorem 1.1 we have the following:

Corollary 1 If two ellipses inscribed in a convex quadrilateral intersect at a point, then that point of intersection cannot lie on either diagonal of the quadrilateral.

2 Preliminary Results

A problem, often referred to in the literature as Newton's problem, was to determine the locus of centers of ellipses inscribed in a convex quadrilateral, Q, in the xy plane. Chakerian([1]) gives a partial solution of Newton's problem using orthogonal projection, which is the solution actually given by Newton.

Theorem 2.1 (Newton)Let M_1 and M_2 be the midpoints of the diagonals of Q. If E is an ellipse inscribed in Q, then the center of E must lie on the open line segment, Z, connecting M_1 and M_2 .

In [2], [4], and [5] we proved several results about ellipses inscribed in quadrilaterals. In particular, in [2] we proved the following converse of Newton's Theorem.

Theorem 2.2 Let Q be a convex quadrilateral in the xy plane and let M_1 and M_2 be the midpoints of the diagonals of Q. Let Z be the open line segment connecting M_1 and M_2 . If $(h, k) \in Z$, then there is a unique ellipse with center (h, k) inscribed in Q.

Remark 2.1 Theorem 2.2 gives a one-one map from the interval I to ellipses inscribed in Q.

We use Theorem 2.2 in this paper to derive the general equation of an ellipse inscribed in Q(see Proposition 3.1). We state the following known result without proof. The first inequality insures that the conic is an ellipse, while the second insures that the ellipse is non-trivial.

Lemma 2.1 The equation $Ax^2 + By^2 + 2Cxy + Dx + Ey + F = 0$, with A, B > 0, is the equation of an ellipse if and only if $AB - C^2 > 0$ and $AE^2 + BD^2 + 4FC^2 - 2CDE - 4ABF > 0$.

We prove Theorem 1.1 first when no two sides of Q are parallel and then when Q is a trapezoid. We do not give all of the details of the proof when Q is a parallelogram.

3 No Two Sides Parallel

Assume first that Q does not have two parallel sides. It suffices, by affine invariance, to prove Theorem 1.1 when Q is the convex quadrilateral with vertices (0,0), (0,1), (1,0), and $(s,t) \in G$, where

$$G = \{(s,t) : s > 0, t > 0, s+t > 1, s \neq 1 \neq t\}.$$

Thus we assume throughout the rest of this section that Q has this form. It is useful to state some facts about Q and related notation.

• The sides of Q are given by $S_1 = \overline{(0,0)(1,0)}, S_2 = \overline{(0,0)(0,1)},$

 $S_3 = \overline{(s,t)(1,0)}$, and $S_4 = \overline{(0,1)(s,t)}$, and the corresponding lines which make up $\partial(Q)$ are given by L_1 : $y = 0, L_2$: $x = 0, L_3$: $y = \frac{t}{s-1}(x-1)$, and L_4 : $y = 1 + \frac{t-1}{s}x.$ • The diagonals of Q are $y = \frac{t}{s}x$ and y = 1 - x and they intersect at

 $\left(\frac{s}{s+t}, \frac{t}{s+t}\right).$ • The midpoints of the diagonals of Q are the points • The line through the midpoints of the diagonals of Q has equation $y = M_1 = \left(\frac{1}{2}, \frac{1}{2}\right)$ and $M_2 = \left(\frac{1}{2}s, \frac{1}{2}t\right)$. • The line through the midpoints of the diagonals of Q has equation y = 0

L(x), where

$$L(x) = \frac{1}{2} \frac{s - t + 2x(t - 1)}{s - 1}$$

/

If s > 1, then

•
$$\partial(Q) = \{(x,y) : 0 \le x \le 1, y = 0\} \cup \{(x,y) : 1 \le x \le s, y = \frac{t}{s-1}(x-1)\} \cup \{(x,y) : 0 \le x \le s, y = 1 + \frac{t-1}{s}x\} \cup \{(x,y) : x = 0, 0 \le y \le 1\}.$$

• $\operatorname{int}(Q) = \{(x,y) : 0 < x \le 1, 0 < y < 1 + \frac{t-1}{s}x\} \cup \{(x,y) : 1 \le x < s, \frac{t}{s-1}(x-1) < y < 1 + \frac{t-1}{s}x\}.$
If $s < 1$, then
• $\partial(Q) = \{(x,y) : 0 \le x \le s, y = 0\} \cup \{(x,y) : s \le x \le 1, y = \frac{t}{s-1}(x-1)\} \cup \{(x,y) : 0 \le x \le s, y = 1 + \frac{t-1}{s}x\} \cup \{(x,y) : x = 0, 0 \le y \le 1\}.$
• $\operatorname{int}(Q) = \{(x,y) : 0 < x \le s, 0 < y < 1 + \frac{t-1}{s}x\} \cup \{(x,y) : s \le x < 1, 0 < y < \frac{t}{s-1}(x-1)\} \cup \{(x,y) : s \le x < 1, 0 < y < \frac{t}{s-1}(x-1)\}$
• $\operatorname{Int}(Q) = \{(x,y) : 0 < x \le s, 0 < y < 1 + \frac{t-1}{s}x\} \cup \{(x,y) : s \le 1, 0 < y < \frac{t}{s-1}(x-1)\}$
• $\operatorname{Int}(Q) = \{(x,y) : 0 < x \le s, 0 < y < 1 + \frac{t-1}{s}x\} \cup \{(x,y) : s \le 1, 0 < y < \frac{t}{s-1}(x-1)\}$
• $\operatorname{Let} I = (\frac{1}{2}, \frac{1}{2}s) \text{ if } s > 1 \text{ and } I = (\frac{1}{2}s, \frac{1}{2}) \text{ if } s < 1$. Then any point on a open line segment connecting M_1 and M_2 has the form $(h, L(h)), h \in I$.

the Before giving our first main result, we need the following simple lemma from [2]:

Lemma 3.1 If $(s,t) \in G$, then s + 2h(t-1) > 0 for all $h \in I$.

Equation (3.1) below was derived using some formulas given in [2] for the foci, center, and semi-major and minor axes of an ellipse inscribed in Q. Once the coefficents of the ellipse equation were simplified, we ended up with a simplified equation, which is given in Proposition 3.1(i) below. It is, as expected, much easier to prove that the equation given below is correct than to give the details of the actual derivation, which we do not provide here.

Proposition 3.1 (i) E is an ellipse inscribed in Q if and only if the general equation of E is given by

$$4(s-1)^{2}(L(h))^{2}(x-h)^{2} + 4(s-1)^{2}h^{2}(y-L(h))^{2}$$

-4(s-1)(2(t-1)h^{2} + (s-t+2)h-s)(x-h)(y-L(h)) (3.1)
= (2h-1)(2(t-1)h+s)(s-2h), h \in I.

(ii) If E is an ellipse given in (i) for some $h \in I$, then E is tangent to the four sides of Q at the points $\zeta_1 = \left(\frac{s-2h}{2(t-1)h+s-t}, 0\right) \in S_1$,

$$\begin{aligned} \zeta_2 &= \left(0, \frac{1}{2} \frac{s - 2h}{(s - 1)h}\right) \in S_2, \\ \zeta_3 &= \left(\frac{s + 2h(t - 1)}{t + s - 2h}, \frac{(2h - 1)t^2}{(s - 1)(s + t - 2h)}\right) \in S_3, \text{ and} \\ \zeta_4 &= \left(\frac{(2h - 1)s^2}{s(s + t - 2) - 2(t - 1)h}, \frac{(s + 2h(t - 1))(s - 1)}{s(s + t - 2) - 2(t - 1)h}\right) \in S_4. \end{aligned}$$

Proof. First, suppose that *E* is given by (3.1). Then by simplifying one can show that *E* has the form $Ax^2 + By^2 + 2Cxy + Dx + Ey + F = 0$, where $A = 4(s-1)^2(L(h))^2$, $B = 4(s-1)^2h^2$, $C = -2(s-1)(2(t-1)h^2 + (s-t+2)h-s)$, D = -2(s-2h)(s-t+2h(t-1)), E = -4h(s-1)(s-2h), and $F = (s-2h)^2$. Then $AB - C^2 = 16(s-1)^4h^2(L(h))^2 - 4(s-1)^2(2(t-1)h^2 + (s-t+2)h-s)^2 = 4(2h-1)(s-1)^2(s-2h)(2(t-1)h+s) > 0$ and $AE^2 + BD^2 + 4FC^2 - 2CDE - 4ABF = 16(2h-1)^2(s-1)^2(s-2h)^2(s+2h(t-1))^2 > 0$ by Lemma 3.1 and the fact that $h \in I$. Thus, by Lemma 2.1, (3.1) defines the equation of an ellipse for any $h \in I$. Now let F(x, y) equal the left hand side of (3.1). Then $F\left(\frac{s-2h}{2(t-1)h+s-t}, 0\right) = F\left(0, \frac{1}{2}\frac{s-2h}{(s-1)h}\right) = F\left(\frac{(2h-1)s^2}{(s+t-2)-2(t-1)h}, \frac{(s+2h(t-1))(s-1)}{s(s+t-2)-2(t-1)h}\right) = (2h-1)(2(t-1)h+s)(s-2h)$, which implies that the four points ζ_1 thrue ζ_1 is an *F*. For fixed *h* differentiating both sides of the acustion in (3.1) with

(2h-1)(2(t-1)h+s)(s-2h), which implies that the four points ζ_1 thru ζ_4 lie on E. For fixed h, differentiating both sides of the equation in (3.1) with respect to x yields:

$$8(s-1)^{2}(L(h))^{2}(x-h) + 8(s-1)^{2}h^{2}(y-L(h))\frac{dy}{dx}$$

-4(s-1)(2(t-1)h^{2} + (s-t+2)h - s)(x-h)\frac{dy}{dx}
-4(s-1)(2(t-1)h^{2} + (s-t+2)h - s)(y-L(h)) = 0.

 \mathbf{SO}

$$\begin{aligned} \frac{dy}{dx} &= D(x,y) = (4(s-1)(2(t-1)h^2 + (s-t+2)h-s)(y-L(h)) - 8(s-1)^2(L(h))^2(x-h))/\\ &\quad (8(s-1)^2h^2(y-L(h)) - 4(s-1)(2(t-1)h^2 + (s-t+2)h-s)(x-h)), \end{aligned}$$

which represents the slope of the ellipse. $D\left(\frac{s-2h}{s-2h-t+2ht},0\right) = 0 = \text{slope}$ of L_1 , $D\left(\frac{s-2h+2ht}{t+s-2h}, \frac{2t^2h-t^2}{(s-1)(t+s-2h)}\right) = \frac{t}{s-1} = \text{slope}$ of L_2 , and $D\left(\frac{2s^2h-s^2}{s^2-2s+ts+2h-2ht}, \frac{(s-2h+2ht)(s-1)}{s^2-2s+ts+2h-2ht}\right) = \frac{t-1}{s} = \text{slope}$ of L_3 . When $x = 0, y = \frac{s-2h}{2(s-1)h}$, the denominator of D(x,y) equals 0, but the nu-

merator of D(x, y) equals $2(2h-1)(s-2h)\frac{s+2h(t-1)}{h} \neq 0$ by Lemma 3.1 and the fact that $h \in I$. Thus L_2 is the tangent line at ζ_2 . It follows easily that ζ_1 thru ζ_4 lie on the line segments S_1 thru S_4 , respectively. For any simple closed convex curve, such as an ellipse, tangent to each side of Q then implies that that curve lies in Q. That proves that E is inscribed in Q. Second, suppose that E is an ellipse inscribed in Q. By Theorem 2.2, E has center $(h_1, L(h_1))$ for some $h_1 \in I$. We have just shown that (3.1) represents a family of ellipses inscribed in Q as h varies over I, and each ellipse given by (3.1) clearly has center (h, L(h)) for some $h \in I$. Let \tilde{E} be the ellipse given by (3.1) with $h = h_1$. Hence \tilde{E} also has center $(h_1, L(h_1))$ and is inscribed in Q. By Theorem 2.2(see Remark 2.1), $\tilde{E} = E$ and the general equation of E must be given by (3.1). That proves (i). We have also just shown that if E is given by (3.1), then E is tangent to the four sides of Q at the four points ζ_1 thru ζ_4 , which proves (ii).

Lemma 3.2 Let

$$f(x,y) = ((t-1)x - (s-1)y)^2 + 2(t-1)x + 2(s-1)y + 1, (s,t) \in G.$$
(3.2)
Then $f(x,y) > 0$ for any $(x,y) \in \overline{Q} = int(Q) \cup \partial(Q).$

Proof. While it is obvious that f(x, y) > 0 in \overline{Q} if both s and t are > 1, we can do the following if one or both of those values is > 1. First, expressing f as a quadratic in s we have

$$\begin{aligned} f(x,y) &= y^2 s^2 + \left(-2y^2 + 2y - 2xty + 2xy\right) s + \\ x^2 - 2x^2 t + 2xty - 2xy + x^2 t^2 + y^2 - 2x + 2xt - 2y + 1, \end{aligned}$$

which has discriminant $(-2y^2 + 2y - 2xty + 2xy)^2 - 4y^2(x^2 - 2x^2t + 2xty - 2xy + x^2t^2 + y^2 - 2x + 2xt - 2y + 1) = -16xy^2(t-1)$. If t > 1, then the discriminant

is negative and thus f has no roots with x and y real. Since $f\left(\frac{s}{s+t}, \frac{t}{s+t}\right) = 4st\frac{s+t-1}{(s+t)^2} > 0$, f(x,y) > 0 in \overline{Q} . Similarly, expressing f as a quadratic in t we have

$$f(x,y) = x^{2}t^{2} + (2xy + 2x - 2xys - 2x^{2})t + x^{2} + 1 + 2xys - 2xy - 2y^{2}s + y^{2} - 2x + y^{2}s^{2} + 2ys - 2y,$$

which has discriminant $(2xy + 2x - 2xys - 2x^2)^2 - 4x^2(x^2 + 1 + 2xys - 2xy - 2y^2s + y^2 - 2x + y^2s^2 + 2ys - 2y) = -16x^2y(s-1)$. Arguing as above, if s > 1, then f(x, y) > 0 in \overline{Q} . So assume now that s < 1 and t < 1, and so $I = \left(\frac{1}{2}s, \frac{1}{2}\right)$. $\frac{\partial f(x, y)}{\partial x} = 0$ and $\frac{\partial f(x, y)}{\partial y} = 0$ implies that (t-1)x + 1 - (s-1)y = 0 and (t-1)x - 1 - (s-1)y = 0, which in turn implies that (t-1)x - (s-1)y = -1 and (t-1)x - (s-1)y = 1. Thus f has no critical points in int(Q). To check f on $\partial(Q)$:

$$f(x,0) = ((t-1)x+1)^2, 0 \le x \le s.$$

$$f\left(x, \frac{t}{s-1}(x-1)\right) = (x+t-1)^2, s \le x \le 1.$$

$$f\left(x, 1+\frac{t-1}{s}x\right) = \frac{(s^2+(t-1)x)^2}{s^2}, 0 \le x \le s.$$

$$f(0,y) = ((s-1)y+1)^2, 0 \le y \le 1.$$

(t-1)x+1=0 implies that $x=\frac{1}{1-t}>1>s$, and thus $(t-1)x+1\neq 0$ if $0\leq x\leq s$, which implies that f(x,0)>0.

If $s \le x \le 1$, then $x+t-1 \ge s+t-1 > 0$, which implies that $f\left(x, \frac{t}{s-1}(x-1)\right) > 0$.

If $0 \le x \le s$, then $s^2 + (t-1)x \ge s^2 + (t-1)s = s(s+t-1) > 0$, which implies that $f\left(x, 1 + \frac{t-1}{s}x\right) > 0$.

Finally, (s-1)y + 1 = 0 implies that $y = \frac{1}{1-s} > 1$, which implies that f(0, y) > 0. Since f has no critical points in int(Q) and f is positive on $\partial(Q)$, f must positive in \overline{Q} .

Proof. (of Theorem 1.1 when no two sides are parallel): For fixed x and y, one can rewrite (3.1) in the form $p_{x,y}(h) = 0$, where

$$p_{x,y}(h) = 4(s-1)^2 (L(h))^2 (x-h)^2 + 4(s-1)^2 h^2 (y-L(h))^2 -4(s-1)(2(t-1)h^2 + (s-t+2)h-s) (x-h) (y-L(h)) -(2h-1)(2(t-1)h+s) (s-2h).$$

 $p_{x,y}$ depends on s and t as well as x and y, but we suppress that dependence in our notation. Let f(x, y) be given by (3.2). Some simplification shows that $p_{x,y}$ is really the following quadratic polynomial in h:

$$p_{x,y}(h) = 4f(x,y)h^{2} +$$

$$4(((t-1)x+1)((s-t)x-s) - (s-1)y(s+(s-t+2)x))h +$$

$$((s-t)x-s)^{2} + 4s(s-1)xy.$$
(3.3)

By Proposition 3.1(i), the equation of any ellipse, E, inscribed in Q is given by $p_{x,y}(h) = 0, h \in I$. Hence, for given (x_0, y_0) , the number of times an ellipse E with equation given by (3.1) passes through (x_0, y_0) equals the number of distinct roots of $p_{x_0,y_0}(h) = 0$ in I. Evaluating $p_{x,y}$ at the endpoints of I yields

$$p_{x,y}\left(\frac{1}{2}\right) = (x+y-1)^2 (s-1)^2 \ge 0$$

$$p_{x,y}\left(\frac{s}{2}\right) = (sy-xt)^2 (s-1)^2 \ge 0.$$
(3.4)

By (3.3), $p'_{x,y}(h_0) = 0$, where

$$h_0 = -\frac{1}{2} \frac{((t-1)x+1)((s-t)x-s) - (s-1)y(s+(s-t+2)x)}{((t-1)x+1)^2 + (s-1)y((s-1)y+2 - 2(t-1)x)}.$$

 h_0 depends on s, t, x, and y, but we suppress that dependence in our notation. After some simplification, it follows easily that

$$p_{x,y}(h_0) = \frac{4(s-1)^2 xy((x-1)t - (s-1)y)((t-1)x - s(y-1))}{f(x,y)}.$$
 (3.5)

We now assume throughout that s > 1 and thus $I = \left(\frac{1}{2}, \frac{1}{2}s\right)$. The case s < 1 follows similarly and we omit the details. Suppose that $(x, y) \in \operatorname{int}(Q)$. We shall prove that $p_{x,y}(h_0) < 0$. First, $y < 1 + \frac{t-1}{s}x$, 0 < x < s implies that sy - s < (t-1)x, so (t-1)x - s(y-1) > 0. Second, if $1 \le x < s$, then $\frac{t}{s-1}(x-1) < y$ since $(x, y) \in \operatorname{int}(Q)$, while if 0 < x < 1, then $\frac{t}{s-1}(x-1) < 0 < y$.

Thus (x-1)t - (s-1)y < 0, 0 < x < s and it follows that the numerator of $p_{x,y}(h_0)$ is negative, while the denominator of $p_{x,y}(h_0)$ is positive by Lemma 3.2. Summarizing:

$$p'_{x,y}(h_0) = 0 \text{ and } (x, y) \in \operatorname{int}(Q) \Rightarrow p_{x,y}(h_0) < 0.$$
 (3.6)

We now prove that $\frac{1}{2} < h_0 < \frac{s}{2}$. One could try to prove this directly, but we found it easier to use the following approach. Let $S = \{(x, y) \in int(Q) : h_0 \in I\}$. We shall prove that S is nonempty and both open and closed in int(Q) in the

Euclidean norm. Since int(Q) is connected, that will prove that S = int(Q). To prove that S is open, suppose that $(x, y) \in S$, so that $p'_{x,y}(h_0) = 0, \frac{1}{2} < h_0 < \frac{s}{2}$. Suppose also that $\{(x_k, y_k)\} \in int(Q)$ converges to (x, y) in the Euclidean norm, and let $h_{0,k}$ be the root of p'_{x_k,y_k} . Clearly $\frac{1}{2} < h_{0,k} < \frac{s}{2}$ for sufficiently large k, which implies that $(x_k, y_k) \in S$ for sufficiently large k and thus S is open. To prove that S is closed, suppose that $(x_k, y_k) \in S$ for all k and that $\{(x_k, y_k)\}$ converges to $(x, y) \in int(Q)$ in the Euclidean norm. Again, let $h_{0,k}$ be the root of p_{x_k,y_k} .Let $\{h_{0,k_j}\}$ be any convergent subsequence of $\{h_{0,k}\}$ with $h_{0,k_j} \to h_0$. Since $p'_{x_k,y_k}(h_{0,k_j}) = 0$ and $\frac{1}{2} < h_{0,k_j} < \frac{s}{2}$, $p'_{x,y}(h_0) = 0$ and $\frac{1}{2} \le h_0 \le \frac{s}{2}$ If $h_0 = \frac{1}{2}$ or $h_0 = \frac{s}{2}$, then $p_{x,y}\left(\frac{1}{2}\right) < 0$ or $p_{x,y}\left(\frac{s}{2}\right) < 0$ by (3.6), which contradicts (3.4). Thus $\frac{1}{2} < h_0 < \frac{s}{2}$ and so S is closed. Finally, to show that S is nonempty, let $x = \frac{s^2}{s+t}$ and $y = \frac{t}{s+t}$. Since the diagonals of Q intersect at $\left(\frac{s}{s+t}, \frac{t}{s+t}\right), (x,y) \in \operatorname{int}(Q).$ The corresponding $p_{x,y}$ is given by $p_{x,y}(h) =$ $\frac{4st(s+t-1)}{(s+t)^2}(2h-1)(2h-s), \text{ and } p'_{x,y}(h) = \frac{8st(s+t-1)}{(s+t)^2}(4h-s-1), \text{ which}$ implies that $h_0 = \frac{1}{4}s + \frac{1}{4}$. Since $h_0 - \frac{1}{2} = \frac{1}{4}(s-1) > 0$ and $h_0 - \frac{s}{2} = -\frac{1}{4}(s-1) < 0$ $0, \frac{1}{2} < h_0 < \frac{s}{2}$. Hence $\left(\frac{s}{s+t}, \frac{t}{s+t}\right) \in S$ and thus S is nonempty. Hence we have shown that S = int(Q), which implies that $h_0 \in I$ for any $(x, y) \in int(Q)$. Since $p_{x,y}(h_0) < 0$ and $h_0 \in I$, by (3.4) we have

$$(x,y) \in \operatorname{int}(Q) \Rightarrow p_{x,y} \text{ has two distinct roots in } \bar{I} = \left[\frac{1}{2}, \frac{1}{2}s\right].$$
 (3.7)

To prove (i), let $P_0 \in \operatorname{int}(Q)$, $P_0 \notin D_1 \cup D_2$. Then $x_0 + y_0 - 1 \neq 0 \neq sy_0 - tx_0$, which implies, by (3.4), that $p_{x_0,y_0}\left(\frac{1}{2}\right) > 0$ and $p_{x_0,y_0}\left(\frac{s}{2}\right) > 0$. By (3.7), p_{x_0,y_0} has two distinct roots, h_1 and h_2 , in *I*. Let E_j be the ellipse with equation given by (3.1) with $h = h_j$, j = 1, 2. By Proposition 3.1(i), E_j is inscribed in *Q*. Since $p_{x_0,y_0}(h_j) = 0$, E_j passes through P_0 since (3.1) holds with $x = x_0, y = y_0$, and $h = h_j, j = 1, 2$. Note that E_1 and E_2 are distinct since they have different centers. Now suppose that an ellipse $\tilde{E} \neq E_j, j = 1, 2$ is inscribed in *Q* and passes through P_0 , and suppose that \tilde{E} has center $(h_3, L(h_3)), h_3 \in I$. By Theorem 2.2(see Remark 2.1), $h_3 \neq h_j$ for j = 1 or 2. By Proposition 3.1(i), the equation of \tilde{E} is given by (3.1) with $h = h_3$. Since \tilde{E} passes through P_0 , $p_{x_0,y_0}(h_3) = 0$, which would give the quadratic p_{x_0,y_0} at least three distinct roots. Hence the only ellipses inscribed in *Q* which pass through P_0 are the ones corresponding to h_1 and h_2 , and so there are precisely two distinct ellipses inscribed in *Q* which pass through P_0 . To prove (ii), let $P_0 \in D_1 \cup D_2$, but $P_0 \neq P = \left(\frac{s}{s+t}, \frac{t}{s+t}\right)$. Then either $x_0 + y_0 - 1 = 0$ or $sy_0 - tx_0 = 0$, but not both, which implies that $p_{x_0,y_0}\left(\frac{1}{2}\right) = 0$ and $p_{x_0,y_0}\left(\frac{s}{2}\right) > 0$, or $p_{x_0,y_0}\left(\frac{1}{2}\right) > 0$ and $p_{x_0,y_0}\left(\frac{s}{2}\right) = 0$. In either case, only one of the roots, say h_1 , of p_{x_0,y_0} lies in I, while $h_2 = \frac{1}{2}$ or $\frac{s}{2}$, where h_2 is the other root of p_{x_0,y_0} . Let E be the ellipse with equation given by (3.1) with $h = h_1$. Arguing as above, E is inscribed in Q and passes through P_0 . Now suppose that an ellipse $E \neq E$ is inscribed in Q and passes through P_0 , and suppose that \tilde{E} has center $(h_3, L(h_3)), h_3 \in I$. By Theorem 2.2(see Remark 2.1), $h_3 \neq h_1$ and $h_3 \neq h_2$ since $h_2 \notin I$. By Proposition 3.1(i), the equation of \tilde{E} is given by (3.1) with $h = h_3$. Since \tilde{E} passes through $P_0, p_{x_0,y_0}(h_3) = 0$, which again would give the quadratic p_{x_0,y_0} at least three distinct roots. Hence the only ellipse inscribed in Q which passes through P_0 is the one corresponding to h_1 , and so there is precisely one ellipse inscribed in Q which passes through P_0 . To prove (iii), let $P_0 = P = \left(\frac{s}{s+t}, \frac{t}{s+t}\right)$. By (3.4), p_{x_0,y_0} vanishes at both endpoints of I, which implies that p_{x_0,y_0} has no roots in I. If there were an ellipse, E, inscribed in Q and passing through P_0 , then the equation of E is given by (3.1) with $h = h_3$ for some $h_3 \in I$. But then $p_{x_0,y_0}(h_3) = 0$, which contradicts the fact that p_{x_0,y_0} has no roots in I. Hence there is no ellipse inscribed in Q which passes thru P. Finally, to prove (iv), suppose that $P_0 \in \partial(Q)$, but P_0 is not one of the vertices of Q. By taking limits of points $P_0 \in int(Q)$, it follows by (3.7) that p_{x_0,y_0} has two roots in \overline{I} , which are not necessarily distinct (indeed we shall prove that they are not distinct). Since $P_0 \in \partial(Q)$, either $x_0 = 0, y_0 = 0, t (x_0 - 1) - (s - 1)y_0 = 0$, or $(t-1)x_0 - s(y_0 - 1) = 0$. Then by (3.5), $p_{x_0,y_0}(h_0) = 0$, which implies that p_{x_0,y_0} has a double root at $h_0 \in \tilde{I}$ since $p'_{x_0,y_0}(h_0) = 0$ by definition. Since P_0 is not a vertex of Q, P_0 cannot lie on either diagonal of Q. Thus by (3.4), $p_{x_0,y_0}\left(\frac{1}{2}\right) > 0$ and $p_{x_0,y_0}\left(\frac{s}{2}\right) > 0$, which implies that $h_0 \in I$. Let E be the ellipse with equation given by (3.1) with $h = h_0$. Since $p_{x_0,y_0}(h_0) = 0$, E passes through P_0 . Also, since $p'_{x_0,y_0}(h_0) = 0$, it is not hard to show that E must be tangent at P_0 to a side of Q. One could also argue that E must be inscribed in Q by Proposition 3.1(i). Arguing as above using the fact that p_{x_0,y_0} cannot have three or more distinct roots, there is no ellipse $E \neq E$ inscribed in Q and which also passes through P_0 , which proves uniqueness.

Remark 3.1 We could also have used Proposition 3.1(ii) to prove Theorem 1.1(iv).

Examples: (1) $s = \frac{1}{2}$, $t = \frac{3}{4}$, $x_0 = \frac{1}{3}$, $y_0 = \frac{3}{4}$, and Q is the convex quadrilateral with vertices (0,0), (0,1), (1,0), and $\left(\frac{1}{2}, \frac{3}{4}\right)$. Then $I = \left(\frac{1}{4}, \frac{1}{2}\right)$ and $P_0 = \left(\frac{1}{3}, \frac{3}{4}\right) \in \operatorname{int}(Q), P_0 \notin D_1 \cup D_2$. By Theorem 1.1(i), there are exactly two ellipses, E_1 and E_2 , inscribed in Q and which pass through P_0 . $p_{x_0,y_0}(h) = \frac{97}{144}h^2 - \frac{37}{72}h + \frac{13}{144}$, which has roots $\frac{37}{97} \pm \frac{6}{97}\sqrt{3} \in I$. Letting $h = \frac{37}{97} - \frac{6}{97}\sqrt{3}$ in (3.1) yields the equation of E_1 : $(29\,673 - 4104\sqrt{3}) x^2 + (23\,632 - 7104\sqrt{3}) y^2 + (26\,808 - 25\,488\sqrt{3}) xy + (18\,864\sqrt{3} - 38\,340) x + (19\,104\sqrt{3} - 37\,104) y = 9792\sqrt{3} - 17\,316$, and letting $h = \frac{37}{97} + \frac{6}{97}\sqrt{3}$ in (3.1) yields the equation of E_2 : $(29\,673 + 4104\sqrt{3}) x^2 + (23\,632 + 7104\sqrt{3}) y^2 + (26\,808 + 25\,488\sqrt{3}) xy - (18\,864\sqrt{3} + 38\,340) x - (19\,104\sqrt{3} + 37\,104) y = -9792\sqrt{3} - 17\,316$. (2) $s = 4, t = 2, x_0 = \frac{1}{2}, y_0 = \frac{1}{4}$, and Q is the convex quadrilateral with

(2) $s = 4, t = 2, x_0 = \frac{1}{2}, y_0 = \frac{1}{4}$, and Q is the convex quantilateral with vertices (0,0), (0,1), (1,0), and (4,2). Then $I = \left(\frac{1}{2},2\right)$ and $P_0 = \left(\frac{1}{2},\frac{1}{4}\right) \in int(Q)$. Since $sy_0 - tx_0 = 0$, $P_0 \in D_1 \cup D_2$, and since $P_0 \neq P = \left(\frac{2}{3},\frac{1}{3}\right)$, by Theorem 1.1(ii), there is exactly one ellipse, E, inscribed in Q which passes through P_0 . $p_{x_0,y_0}(h) = \frac{57}{4}h^2 - 36h + 15$, which has roots $h_1 = \frac{10}{19} \in I$ and $h_2 = 2 \in \overline{I} - I$. Letting $h = \frac{10}{19}$ in (3.1) yields the equation of E: $63\,916x^2 + 68\,400y^2 + 110\,352xy - 123\,424x - 127\,680y + 64\,960 = 5376$.

4 Trapezoid

Assume now that Q is a trapezoid. It suffices, by affine invariance, to prove Theorem 1.1 where Q is the trapezoid with vertices (0,0), (1,0), (0,1), and $(1,t), 0 < t \neq 1$. It is useful to state some facts about Q and related notation.

• The sides of Q are given by $S_1 = \overline{(0,0)(1,0)}, S_2 = \overline{(0,0)(0,1)}, S_3 = \overline{(1,t)(1,0)}$, and $S_4 = \overline{(0,1)(1,t)}$, and the corresponding lines which make up $\partial(Q)$ are given by

 $L_1: y = 0, L_2: x = 0, L_3: x = 1, \text{ and } L_4: y = 1 + (t - 1)x.$

• The diagonals of Q are y = tx and y = 1 - x, and they intersect at $\left(\frac{1}{1+t}, \frac{t}{1+t}\right)$.

• The midpoints of the diagonals of Q are the points $M_1 = \left(\frac{1}{2}, \frac{1}{2}\right)$ and $\begin{pmatrix} 1 & 1 \\ 2 & 2 \end{pmatrix}$

$$M_2 = \left(\frac{1}{2}, \frac{1}{2}t\right).$$

• The open line segment joining M_1 and M_2 is $\left\{h = \frac{1}{2}, k \in I\right\}$, where $I = \left(\frac{1}{2}, \frac{1}{2}t\right)$ if t > 1, or $I = \left(\frac{1}{2}t, \frac{1}{2}\right)$ if t < 1. • $\partial(Q) = \{(x, y) : 0 \le x \le 1, y = 0\} \cup \{(x, y) : x = 1, 0 \le y \le t\}$ $\cup \{(x,y): 0 \le x \le 1, y = 1 + (t-1)x\} \cup \{(x,y): x = 0, 0 \le y \le 1\} \text{ and } int (Q) = \{(x,y): 0 < x < 1, 0 < y < 1 + (t-1)x\}.$ • Note that

$$(t-1)x + 1 > 0, 0 \le x \le 1.$$
(4.1)

First we need the following proposition, which is the analogy of Proposition 3.1 for trapezoids.

Proposition 4.1 Let *Q* be the trapezoid with vertices (0,0), (1,0), (0,1), and $(1,t), 0 < t \neq 1$.

(i) E is an ellipse inscribed in Q if and only if the general equation of E is given by

$$4k^{2}(t-1)^{2}x^{2} + (t-1)^{2}y^{2} + 4(1-t)(tk-t+k)xy +4k(t-1)(2k-t)x + 2(t-1)(2k-t)y = -(2k-t)^{2}, k \in I.$$
(4.2)

(ii) If E is the ellipse given in (i), then E is tangent to the four sides of Q at the points $\zeta_1 = \left(\frac{2k-t}{2k(1-t)}, 0\right) \in S_1, \ \zeta_2 = \left(0, \frac{2k-t}{1-t}\right) \in S_2, \ \zeta_3 = \left(1, t\frac{1-2k}{1-t}\right) \in S_3, \ and \ \zeta_4 = \left(\frac{1-2k}{(1-t)(t+1-2k)}, \frac{t}{t+1-2k}\right) \in S_4.$

Proof. First, suppose that *E* is given by (4.2). Then *E* has the form $Ax^2 + By^2 + 2Cxy + Dx + Ey + F = 0$, where $A = 4k^2(t-1)^2$, $B = (t-1)^2$, C = 2(1-t)(tk-t+k), D = 4k(t-1)(2k-t), E = 2(t-1)(2k-t), and $F = (2k-t)^2$. Then $AB - C^2 = 4t(2k-1)(t-2k)(t-1)^2 > 0$ since $k \in I$ implies that t < 2k < 1 if t < 1, and 1 < 2k < t if t > 1. Also, $AE^2 + BD^2 + 4FC^2 - 2CDE - 4ABF = 16t^2(t-1)^2(2k-1)^2(2k-t)^2 > 0$ since t < 2k < 1 if t < 1, and 1 < 2k < t if t > 1. By Lemma 2.1, (4.2) defines the equation of an ellipse for any $k \in I$. Now let $F(x, y) = 4k^2(t-1)^2x^2 + (t-1)^2y^2 + 4(1-t)(tk-t+k)xy+4k(t-1)(2k-t)x+2(t-1)(2k-t)y+(2k-t)^2$, the left hand side of (4.2). Then $F\left(\frac{2k-t}{2k(1-t)}, 0\right) = F\left(0, \frac{2k-t}{1-t}\right) = F\left(1, t\frac{1-2k}{1-t}\right) = F\left(\frac{1-2k}{(1-t)(t+1-2k)}, \frac{t}{t+1-2k}\right) = -(2k-t)^2$, which implies that the four points ζ_1 thru ζ_4 lie on *E*. Differentiating both sides of the equation in (4.2) with respect to *x* yields $8k^2(t-1)^2x+2(t-1)^2y\frac{dy}{dx}+4(1-t)(tk-t+k)y+4(1-t)(tk-t+k)x\frac{dy}{dx}+4k(t-1)(2k-t)+2(t-1)(2k-t)\frac{dy}{dx}=0$, which implies that $\frac{dy}{dx} = D(x, y) = \frac{4k^2(t-1)x-2(tk-t+k)y+2k(2k-t)}{2(tk-t+k)x-(t-1)y-(2k-t)}$. Now $D\left(\frac{2k-t}{2k(1-t)}, 0\right) = 0$ = slope of L_1 and $D\left(\frac{1-2k}{(1-t)(t+1-2k)}, \frac{t}{t+1-2k}\right) = t-1$ = slope of L_4 .

When x = 0 and $y = \frac{2k-t}{1-t}$, the denominator of D(x, y) equals 0, but the numerator of D(x, y) equals $\frac{2t(2k-t)(2k-1)}{t-1} \neq 0$. Thus L_2 is the tangent line at ζ_2 . When x = 1 and $y = t\frac{1-2k}{1-t}$, again the denominator of D(x, y) equals 0, but the numerator of D(x, y) equals $\frac{2t(1-2k)(2k-t)}{t-1} \neq 0$. Thus L_3 is the tangent line at ζ_3 . It follows easily that ζ_1 thru ζ_4 lie on the line segments S_1 thru S_4 , respectively. For any simple closed convex curve, such as an ellipse, tangent to each side of Q then implies that that curve lies in Q. That proves that E is inscribed in Q. Second, suppose that E is an ellipse inscribed in Q. By Theorem 2.2, E has center $\left(\frac{1}{2}, k\right)$ for some $k \in I$. We have just shown that (4.2) represents a family of ellipse given by (4.2) has center $\left(\frac{1}{2}, k\right)$ for some $k \in I$. Arguing exactly as in the case above when Q does not have two parallel sides, it follows that the general equation of E must be given by (4.2). That proves (i). We have also just shown that if E is given by (4.2), then E is tangent to the four sides of Q at the four points ζ_1 thru ζ_4 , which proves (ii).

Lemma 4.1 Let $g(x,y) = (x-1)^2 + (y-tx)(x-1) + txy$ and $h(x,y) = t(t-1)x^2 - (t+1)xy + tx + y$. Then g(x,y) > 0 and h(x,y) > 0 if $(x,y) \in int(Q)$.

Proof. Suppose that $(x, y) \in int(Q)$. Then 0 < y < 1 + (t - 1)x, which implies that $g(x, y) > (x - 1)^2 + (1 + (t - 1)x - tx)(x - 1) + tx = tx > 0$.

Now express h as a linear function of y, L(y), for fixed x and t: $L(y) = (1 - (t+1)x)y + t(t-1)x^2 + tx$. Then by (4.1), $L(0) = t(t-1)x^2 + tx = tx((t-1)x+1) > 0$ and L(1 + (t-1)x) = (1-x)((t-1)x+1) > 0, which implies that L(y) > 0 for all 0 < y < 1 + (t-1)x.

(of Theorem 1.1 for trapezoids)For fixed x and y, one can write (4.2) in the form $p_{x,y}(k) = 0$, where $p_{x,y}$ is the polynomial in k given by

$$p_{x,y}(k) = 4 (xt - x + 1)^2 k^2 +4(tx - yxt^2 - xt^2 - y + yt - t + yx)k +(t + y)^2 + ty (yt - 2t - 2y) + 4txy (t - 1).$$
(4.3)

 $p_{x,y}$ depends on t as well, but we suppress that dependence in our notation. By Proposition 4.1(i), the equation of any ellipse, E, inscribed in Q is given by $p_{x,y}(k) = 0, k \in I$. Hence, for given (x, y), the number of times an ellipse E with equation given by (4.2) passes through (x, y) equals the number of distinct roots of $p_{x,y}(k) = 0$ in *I*. Evaluating $p_{x,y}$ at the endpoints of *I* yields

$$p_{x,y}\left(\frac{1}{2}\right) = (y+x-1)^2 (t-1)^2, \qquad (4.4)$$
$$p_{x,y}\left(\frac{t}{2}\right) = (t-1)^2 (tx-y)^2.$$

Now $p'_{x,y}(k_0) = 0$, where

$$k_0 = \frac{(t^2 - 1)xy + t(t - 1)x - (t - 1)y + t}{2((t - 1)x + 1)^2}.$$
(4.5)

We now assume throughout that t > 1 and thus $I = \left(\frac{1}{2}, \frac{1}{2}t\right)$. The case t < 1 follows similarly and we omit the details. To prove (i), let $P_0 \in int(Q), P_0 \notin D_1 \cup D_2$. A simple computation yields

$$p_{x_0,y_0}(k_0) = \frac{4t \left(t-1\right)^2 x_0 \left(x_0-1\right) y_0 \left((t-1)x_0-y_0+1\right)}{\left((t-1)x_0+1\right)^2}.$$
(4.6)

Note that the denominator in (4.6) is nonzero by (4.1). Now $(x_0, y_0) \in int(Q)$ implies that $(t-1)x_0-y_0+1 > 0$ and $x_0-1 < 0$. Hence $p_{x_0,y_0}(k_0) < 0$. Unlike the case above when Q does not have two parallel sides, here we shall prove directly that $\frac{1}{2} < k_0 < \frac{t}{2}$. First, $k_0 - \frac{1}{2} = \frac{(t^2 - 1)x_0y_0 + t(t-1)x_0 - (t-1)y_0 + t}{2((t-1)x_0+1)^2} - \frac{1}{2} = (t-1)\frac{(x_0-1)^2 + (x_0-1)y_0 - tx_0(x_0-y_0-1)}{2((t-1)x_0+1)^2} = (t-1)g(x_0,y_0) > 0$ by

Lemma 4.1. Second, $\frac{t}{2} - k_0 = \frac{t}{2} - \frac{(t^2 - 1)x_0y_0 + t(t - 1)x_0 - (t - 1)y_0 + t}{2((t - 1)x_0 + 1)^2} = t(t - 1)x^2 - (t + 1)x_0y_0 + tx_0 + y_0$

$$(t-1)\frac{t(t-1)x_0^2 - (t+1)x_0y_0 + tx_0 + y_0}{2((t-1)x_0 + 1)^2} = (t-1)h(x_0, y_0) > 0, \text{ again by}$$

Lemma 4.1. That proves that $\frac{1}{2} < k_0 < \frac{t}{2}$ when $P_0 \in \operatorname{int}(Q)$. Also, $P_0 \notin D_1 \cup D_2$ implies that $y_0 + x_0 - 1 \neq 0 \neq tx_0 - y_0$, which implies, by (4.4), that $p_{x_0,y_0}\left(\frac{1}{2}\right) > 0$ and $p_{x_0,y_0}\left(\frac{t}{2}\right) > 0$. Thus p_{x_0,y_0} must have two distinct roots in *I*. Arguing as we did for the case when no two sides of *Q* are parallel, it follows that precisely two distinct ellipses inscribed in *Q* pass through P_0 . To prove (ii), let $P_0 \in D_1 \cup D_2$, but $P_0 \neq P = \left(\frac{1}{1+t}, \frac{t}{1+t}\right)$. Then either $x_0 + y_0 - 1 = 0$ or $tx_0 - y_0 = 0$, but not both, which implies, by (4.4), that $p_{x_0,y_0}\left(\frac{1}{2}\right) = 0$ and $p_{x_0,y_0}\left(\frac{t}{2}\right) > 0$, or $p_{x_0,y_0}\left(\frac{1}{2}\right) > 0$ and $p_{x_0,y_0}\left(\frac{t}{2}\right) = 0$. In either case p_{x_0,y_0} has exactly one real root in *I* since $p_{x_0,y_0}(k_0) < 0$. Again, arguing as we did for the case when no two sides of *Q* are parallel, it follows that precisely one distinct ellipse inscribed in *Q* passes through P_0 . To prove (ii),

let $P_0 = P = \left(\frac{1}{1+t}, \frac{t}{1+t}\right)$. Then $x_0 + y_0 - 1 = tx_0 - y_0 = 0$, which implies, by (4.4), that $p_{x_0,y_0}\left(\frac{1}{2}\right) = p_{x_0,y_0}\left(\frac{t}{2}\right) = 0$ and hence p_{x_0,y_0} has no roots in I. As above, there is no ellipse inscribed in Q which passes thru P_0 . Finally we prove (iv). Suppose that $P_0 \in \partial(Q)$, but P_0 is not one of the vertices of Q. Since $P_0 \in \partial(Q)$, either $x_0 = 0, x_0 = 1, y_0 = 0$, or $(t-1)x_0 - y_0 + 1 = 0$. Then by (4.6), $p_{x_0,y_0}(k_0) = 0$, which implies that p_{x_0,y_0} has a double root at $k_0 \in \overline{I}$ since $p'_{x_0,y_0}(k_0) = 0$ by definition. Since P_0 is not a vertex of Q, P_0 cannot lie on either diagonal of Q. Thus by (4.4), $p_{x_0,y_0}\left(\frac{1}{2}\right) > 0$ and $p_{x_0,y_0}\left(\frac{t}{2}\right) > 0$, which implies that $k_0 \in I$. Let E be the ellipse with equation given by (4.2) with $k = k_0$. Since $p_{x_0,y_0}(k_0) = 0$, E passes through P_0 . Also, since $p'_{x_0,y_0}(k_0) = 0$, it is not hard to show that E must be tangent at P_0 to a side of Q. One could also argue that E must be inscribed in Q by Proposition 4.1(i). Using the fact that p_{x_0,y_0} cannot have three or more distinct roots, it follows easily that there is no ellipse $\tilde{E} \neq E$ inscribed in Q and which also passes through P_0 , which proves uniqueness. That completes the proof of Theorem 1.1 when Q is a trapezoid.

Example: Consider the trapezoid, \tilde{Q} , with vertices (-1, 2), (3, 4), (3, -1), and (9, 2). We want to find an an ellipse inscribed in \tilde{Q} which passes through the point (4, 2) int (\tilde{Q}) . First define the affine transformation

5 Parallelogram

Finally, assume that Q is a parallelogram. Since any parallelogram is affine equivalent to the unit square, it suffices to assume that Q is the unit square. In [[5]] it was shown that if Z is the rectangle with vertices (0,0), (l,0), (0,k), and (l,k), where l, k > 0, then the general equation of an ellipse inscribed in Z is given by $k^2x^2 + l^2y^2 - 2l(k-2v)xy - 2lkvx - 2l^2vy + l^2v^2 = 0, 0 < v < k$. It then follows immediately(or one can easily prove this directly) that the general equation of an ellipse inscribed in the unit square is given by $x^2 + y^2 + 2(2v-1)xy - 2vx - 2vy + v^2 = 0, 0 < v < 1$. The rest of the proof of Theorem 1.1 is similar to the proof for the trapezoid case and we omit the details.

6 Algorithms

Throughout, A denotes a non-singular affine map of the xy plane to itself.

• Given a point, P_0 , in the interior of a quadrilateral, Q, find all ellipses inscribed in \tilde{Q} which pass through \tilde{P}_0 .

Case 1: \hat{Q} does not have two parallel sides.

Step 1: Find an A which maps \tilde{Q} to the convex quadrilateral, Q, with vertices (0,0), (0,1), (1,0), and (s,t), where $s > 0, t > 0, s + t > 1, s \neq 1 \neq t$. Let $I = \left(\frac{1}{2}, \frac{1}{2}s\right)$ if s > 1, $I = \left(\frac{1}{2}s, \frac{1}{2}\right)$ if s < 1. A sends \tilde{P}_0 to the point $P_0 = (x_0, y_0)$ lying in the interior of Q.

Step 2: Find the roots of the quadratic polynomial in h given in (3.3), with $x = x_0$ and $y = y_0$. If P_0 does not lie on either diagonal of Q, then p_{x_0,y_0} has two roots in I, h_1 and h_2 . There are then two distinct ellipses, E_1 and E_2 , inscribed in Q which pass through P_0 . The equations of E_1 and E_2 are given by (3.1) with $h = h_j, j = 1, 2$.

If P_0 lies on one of the diagonals of Q, but P_0 does not equal the intersection point of the diagonals, then p_{x_0,y_0} has one root in I, h_1 . There is then one ellipse, E, inscribed in Q which passes through P_0 . The equation of E is given by (3.1) with $h = h_1$.

In each case above, use the map A to obtain the corresponding equation of each ellipse inscribed in \tilde{Q} which passes through \tilde{P}_0 .

Finally, if \tilde{P}_0 equals the intersection point of the diagonals of \tilde{Q} , then there is no ellipse inscribed in \tilde{Q} which passes through \tilde{P}_0 .

Case 2: Q is a trapezoid.

Step 1: Find an A which maps \tilde{Q} to the trapezoid, Q, with vertices (0,0), (1,0), (0,1), and $(1,t), 0 < t \neq 1$. Let $I = \left(\frac{1}{2}, \frac{1}{2}t\right)$ if t > 1, or $I = \left(\frac{1}{2}t, \frac{1}{2}\right)$ if t < 1. A sends \tilde{P}_0 to the point $P_0 = (x_0, y_0)$ lying in the interior of Q.

Step 2: Find the roots of the quadratic polynomial in k given in (4.6), with $x = x_0$ and $y = y_0$.

As with case 1, if P_0 does not lie on either diagonal of Q, then p_{x_0,y_0} has two roots in I, k_1 and k_2 . There are then two distinct ellipses, E_1 and E_2 , inscribed in Q which pass through P_0 . The equations of E_1 and E_2 are given by (4.2) with $k = k_i, j = 1, 2$.

If P_0 lies on one of the diagonals of Q, but P_0 does not equal the intersection point of the diagonals, then p_{x_0,y_0} has one root in I, k_1 . There is then one ellipse, E, inscribed in Q which passes through P_0 . The equation of E is given by (4.2) with $k = k_1$.

In each case above, use the map A to obtain the corresponding equation of each ellipse inscribed in \tilde{Q} which passes through \tilde{P}_0 .

Finally, if \tilde{P}_0 equals the intersection point of the diagonals of \tilde{Q} , then there is no ellipse inscribed in \tilde{Q} which passes through \tilde{P}_0 .

Case 3: \tilde{Q} is a parallelogram.

The details here are similar to the two cases above. One can first use an affine map to send \tilde{Q} to the unit square.

References

- G. D. Chakerian, A Distorted View of Geometry, MAA, Mathematical Plums, Washington, DC, 1979, 130-150.
- [2] Alan Horwitz, Ellipses of maximal area and of minimal eccentricity inscribed in a convex quadrilateral, Australian Journal of Mathematical Analysis and Applications, 2(2005), Issue 1, Article 4, 1-12.
- [3] Alan Horwitz, Dynamics of ellipses inscribed in triangles, submitted to the Journal of Classical Analysis.
- [4] Alan Horwitz, An area inequality for ellipses inscribed in quadrilaterals, Journal of Mathematical Inequalities, 4(2010), Issue 3, 431–443.
- [5] Alan Horwitz, Ellipses Inscribed in Parallelograms, Australian Journal of Mathematical Analysis and Applications, Volume 9, Issue 1(2012), 1–12.