

SOME PROPERTIES OF CONVECTION IN HYBRID STARS

A.V. Yudin^{1,2*}, M. Hempel³, D.K. Nadyozhin^{1,4}, T.L. Razinkova¹

¹*Institute for Theoretical and Experimental Physics, Moscow*

²*Novosibirsk State University*

³*Department of Physics, University of Basel, Basel, Switzerland*

⁴*SRC “Kurchatov Institute”, Moscow*

Abstract. It is shown that the unusual thermodynamic properties of matter within the region of two-phase coexistence in hybrid stars result in a change of the standard condition for beginning of convection. In particular, the thermal flux transported by convection may be directed towards the stellar center. We discuss favorable circumstances leading to such an effect of “inverse convection” and its possible influence on the thermal evolution of hybrid stars.

Key words: phase transition, quark matter, convection, hybrid stars, supernovae.

* e-mail <yudin@itep.ru>

INTRODUCTION

Neutron stars, together with stellar mass black holes, develop from collapsing cores of massive stars at the final stages of their evolution. The birth of a neutron star most likely manifests itself as a supernova explosion. The major part of matter in neutron stars proves to be in an extreme state with a density exceeding the nuclear one $\rho_n \approx 2.6 \times 10^{14} \text{ g/cm}^3$.

The possibility of phase transitions in nuclear matter was first supposed by Gurevich (1939). Then Ivanenko and Kurdgelaidze (1965) and Itoh (1970) advanced hypotheses concerning stars composed of quark matter. Nowadays there exists an extensive literature on neutron stars, quark stars, and neutron stars containing quark cores (the so-called hybrid stars). The excellent monograph of Haensel et al. (2007) can be recommended for further reading.

In a hybrid star the quark core is separated from the outer nuclear matter envelope by an intermediate layer where the phase transition (PT) between nuclear and quark matter occurs. Within this region of coexistence of nuclear and quark phases there is the possibility that the pressure is decreasing with growing temperature at constant density. Such an effect was mentioned, for instance, by Yudin et al. (2013), Hempel et al. (2013), and Iosilevskiy (2014). In single phases of quark or hadronic matter usually the opposite is the case, i.e., the pressure is increasing with growing temperature at constant density. According to the Clapeyron-Clausius formula (Landau and Lifshitz 1980) the derivative of pressure over temperature $(\frac{\partial P}{\partial T})_{\text{pt}}$ along the phase transition line in the phase diagram is given by

$$\left(\frac{\partial P}{\partial T}\right)_{\text{pt}} = \frac{S_2 - S_1}{\frac{1}{\rho_2} - \frac{1}{\rho_1}} < 0. \quad (1)$$

Here $S_{1,2}$ and $\rho_{1,2}$ are the entropies per baryon and densities of phases 1 and 2 in the region of their coexistence. We consider here the simplest Maxwellian description of the PT. In this case, $(\frac{\partial P}{\partial T})_{\text{pt}}$ in Eq. (1) can be replaced by $(\frac{\partial P}{\partial T})_{\rho}$, as long as ρ corresponds to a value where one has phase coexistence. The negative sign of the derivative in Eq. (1) thus appears because the quark phase has a higher entropy per baryon, $S_2 > S_1$ (note that $\rho_2 > \rho_1$). This also implies that to go from the low-density phase 1 to the high-density phase 2 at fixed temperature the system absorbs a thermal energy per baryon of $\Delta q = T(S_2 - S_1) > 0$.

Let us consider a schematic view of the phase diagram in the density-temperature




Fig. 1. Schematic view of a phase diagram for the transition between hadronic and quark matter.

plane (Fig. 1). The boundaries of the phases are shown by bold dashed lines. The region between the lines corresponds to two-phase coexistence.¹ An isentropic curve ($S = \text{const}$) is shown by the solid line. One can observe that in the phase-coexistence region the temperature begins to decrease along the isentropic curve when the density increases. Such a behavior of the PT was indicated by a number of authors, and especially by Steiner et al. (2000). Note that the temperate-pressure diagram would look qualitatively similar as Fig. 1.

The main idea of our work is to investigate the properties of convection expected in the region of phase coexistence. Consider for example a convective element that starts its adiabatic motion in the interior of a hybrid star at point A and moves outwards reaching point A' . So it keeps the initial entropy but has along its way the same pressure and hence temperature (let us remind that we consider Maxwellian PT) as the environment. If on its way the convective element proves to be more dense than the environment (point C) it will begin to sink. Such a configuration is convectively stable. On the contrary, when the state of the environment corresponds to point B , the density of the convective element is lower than that of the environment and it continues to rise — the configuration is convectively

¹Often the region of phase coexistence is also called the “mixed phase”.

unstable. The entropy of environment in point B is higher than in point A' . Therefore the condition of appearance of convection reads

$$\frac{dS}{dr} > 0. \quad (2)$$

This condition has the inequality sign opposite to the common Schwarzschild criterion. Our paper is devoted to the consideration of such uncommon circumstance and following consequences.

THE CONDITION OF APPEARANCE OF CONVECTION

The criterion of appearance of convection in General Relativity can be deduced in the same way as for Newtonian gravity (Thorne 1966). One has only to take into account that the energy per unit volume ϵ (with the rest mass energy being included) plays the role of density. For the beginning of convection the internal energy ϵ_{ce} of an ascending convective element should remain less than that of the surrounding matter ϵ_{sm} and vice versa for an descending element: $(\epsilon_{ce} - \epsilon_{sm})\Delta r < 0$, where Δr is a radial displacement.

Taking into account that the convective element moves adiabatically we can write down the following equations for changes in internal energies ϵ_{ce} and ϵ_{sm} on the way Δr :

$$\begin{cases} \Delta\epsilon_{ce} = \left(\frac{\partial\epsilon}{\partial P}\right)_{S,Y} \Delta P, \\ \Delta\epsilon_{sm} = \left(\frac{\partial\epsilon}{\partial P}\right)_{S,Y} \Delta P + \left(\frac{\partial\epsilon}{\partial S}\right)_{P,Y} \Delta S + \left(\frac{\partial\epsilon}{\partial Y}\right)_{P,S} \Delta Y, \end{cases} \quad (3)$$

where P is the pressure, S is the entropy per baryon, and Y is an additional dimensionless independent variable describing the chemical composition of matter. For supernova matter the values of the electron fraction $Y = Y_e$ or lepton charge fraction $Y = Y_e + Y_\nu$ are usually specified. In the case of several independent variables Y_i describing the composition one has to use sum over i in the last term of the above expression for $\Delta\epsilon_{sm}$. In the equation above we have thus assumed that the convective element maintains its original value of Y .

Now we can write the criterion of appearance of convection in the well known Ledoux form

$$\left(\frac{\partial\epsilon}{\partial S}\right)_{P,Y} \frac{dS}{dr} + \left(\frac{\partial\epsilon}{\partial Y}\right)_{P,S} \frac{dY}{dr} > 0. \quad (4)$$

For simplicity we assume below $Y = \text{const}$. Then the onset of convection depends on the distribution of entropy in the star (term dS/dr in Eq. (4)) and the sign of the term

$\left(\frac{\partial \epsilon}{\partial S}\right)_{P,Y}$. In the nonrelativistic limit we have $\epsilon \approx \rho c^2$ with ρ being the baryon mass density. Therefore up to a factor c^2 the multiplier in front of dS/dr is given by

$$\left(\frac{\partial \rho}{\partial S}\right)_P = -\frac{\rho^2}{\left(\frac{\partial P}{\partial T}\right)_S} = -\frac{T\rho}{P\gamma c_V} \left(\frac{\partial P}{\partial T}\right)_\rho, \quad (5)$$

where we introduce the adiabatic index γ and the specific heat capacity c_V :

$$\gamma \equiv \left(\frac{\partial \ln P}{\partial \ln \rho}\right)_S, \quad c_V \equiv T \left(\frac{\partial S}{\partial T}\right)_\rho. \quad (6)$$

For matter under common conditions the right hand side of Eq. (5) is obviously strictly negative and we obtain the criterion for onset of convection in the Schwarzschild form:

$$\frac{dS}{dr} < 0. \quad (7)$$

Hence, a negative gradient of entropy causes the onset of convection. However, within the phase-coexistence region the derivatives $\left(\frac{\partial P}{\partial T}\right)_S$ and $\left(\frac{\partial P}{\partial T}\right)_\rho$ can be negative and as a result the Schwarzschild criterion changes its sign. The negative entropy gradient in this region ensures the convective stability while the the positive one stimulates the development of an convective instability.

Let us consider now the general case. One can easily show that the factor in front of dS/dr in Eq. (4) is equal to

$$\left(\frac{\partial \epsilon}{\partial S}\right)_{P,Y} = \rho T \left[1 - \frac{\epsilon + P}{T \left(\frac{\partial P}{\partial T}\right)_S} \right]. \quad (8)$$

In the nonrelativistic limit the second term in square brackets is much larger than 1 and equivalent to Eq. (5). If $\left(\frac{\partial P}{\partial T}\right)_S < 0$ the sign of this term as well as the criterion of convection change again.

A REMARK ABOUT ISOTHERMS

Let us make a slight digression to illustrate the formulae of the previous section with an example of convection in isothermal stars which are in the state of total thermal equilibrium. In the limit of Newtonian gravity the temperature is constant throughout the star, $T = \text{const.}$, and we have

$$\frac{dS}{dr} = \left(\frac{\partial S}{\partial P}\right)_T \frac{dP}{dr}. \quad (9)$$

Using Eq. (5) we obtain

$$\left(\frac{\partial \rho}{\partial S}\right)_P \frac{dS}{dr} = \frac{T}{\rho^2 c_P} \left(\frac{\partial \rho}{\partial T}\right)_P^2 \frac{dP}{dr} \leq 0, \quad (10)$$

where $c_P = T \left(\frac{\partial S}{\partial T}\right)_P$ is the heat capacity at constant pressure and the last inequality is valid because the pressure never increases toward the surface of a star in hydrostatic equilibrium. Thus we have the well known result that a nonrelativistic isotherm is always convectively stable.

For the relativistic isotherms the condition of thermal equilibrium is $Te^\nu = \text{const.}$ (see Tolman 1969). Here the function ν is connected with the time component of the metric tensor ($e^{2\nu} = g_{00}$) and can be found from Einstein equations:

$$\frac{d\nu}{dr} = -\frac{1}{P + \epsilon} \frac{dP}{dr}. \quad (11)$$

Therefore for a relativistic isotherm the temperature falls by going outwards in the star proportionally to $\frac{dP}{dr}$:

$$\frac{1}{T} \frac{dT}{dr} = \frac{1}{P + \epsilon} \frac{dP}{dr}. \quad (12)$$

In this case the convective stability is not obvious. Using Eq. (12) one can find the entropy gradient:

$$\frac{dS}{dr} = \left(\frac{\partial S}{\partial T}\right)_P \frac{dT}{dr} + \left(\frac{\partial S}{\partial P}\right)_T \frac{dP}{dr} = \frac{c_P}{P + \epsilon} \left[1 - \frac{\epsilon + P}{T \left(\frac{\partial P}{\partial T}\right)_S}\right] \frac{dP}{dr}. \quad (13)$$

Then finally using also Eq. (8) we obtain

$$\left(\frac{\partial \epsilon}{\partial S}\right)_{P,Y} \frac{dS}{dr} = \frac{\rho T c_P}{P + \epsilon} \left[1 - \frac{\epsilon + P}{T \left(\frac{\partial P}{\partial T}\right)_S}\right]^2 \frac{dP}{dr} \leq 0. \quad (14)$$

Now one can see that also the relativistic isotherm is always convectively stable.

RESULTS OF CALCULATIONS

As an example of our calculations with an realistic equation of state (EOS) we discuss the results shown in Figs. 2 and 3, where the dependencies of the temperature T and derivative $\left(\frac{\partial \epsilon}{\partial S}\right)_{P,Y}$ on baryon density n_b are shown for different values of the entropy per baryon S .² The dashed curves in Figs. 2 and 3 correspond to the EOS for nuclear (hadronic) matter from Shen et al. (1998), Shen et al. (2011) (abbreviated STOS) whereas the solid ones refer to the hybrid EOS from Sagert et al. (2009), Fischer et al. (2011) which includes the

²We use natural units where $\hbar = c = k_B = 1$.




Fig. 2. The temperature versus baryon density n_b for hadronic (Shen et al. 2011) and hybrid equations of state (Sagert et al. 2009) (dashed and solid curves, respectively) for various different values of the entropy per baryon S and a fixed lepton fraction of $Y_L = 0.4$.

phase transition to quark matter using the quark bag model described by bag constant B . The quark bag constant $B^{1/4}$ and the lepton number per baryon $Y = Y_L = Y_e + Y_\nu$ are assumed to be 165 MeV and 0.4, respectively. The neutrino component is presumed to be in equilibrium with matter. Global charge neutrality is assumed for phase coexistence, which, as was first pointed out by Glendenning (1992), results in a Gibbs or “non-congruent” (Hempel et al. 2013) phase transition.

The derivative shown in Fig. 3 corresponds to a secondary thermodynamic derivative. We have calculated it from thermodynamic consistent, tri-cubic interpolation of the free energy. The used scheme is not able to handle properly the onset and end of the phase transition, leading to the spikes visible in Fig. 3 for the hybrid EOS that have to be considered as numerical artifacts.

The region of phase coexistence is distinctly identified with the negative temperature




Fig. 3. Derivative $\left(\frac{\partial \epsilon}{\partial S}\right)_{P, Y}$ multiplied by $\frac{1}{n_b T}$ versus baryon density n_b for hadronic (Shen et al. 2011) and hybrid equations of state (Sagert et al. 2009) (dashed and solid curves, respectively) for various different values of the entropy per baryon S and a fixed lepton fraction of $Y_L = 0.4$.

gradient in Fig. 2 and positive sign of derivative $\left(\frac{\partial \epsilon}{\partial S}\right)_{P, Y} > 0$ in Fig. 3. Exactly here the “inverse” convection could be developed. One can observe such an effect also in fig. 6 from Fischer et al. (2011) where the results of calculations for different values of lepton number are presented.

However, the situation is rather complex. First, along the same isentrope there are regions both with positive and negative sign of $\left(\frac{\partial \epsilon}{\partial S}\right)_{P, Y}$. Second, we have observed that the region of inverse convection is maximal for symmetrical matter with $Y_e = 0.5$ and shrinks as Y_e decreases when the neutronization is increasing. Therefore, the existence of the inverse convection and its parameters depend on:

1. the equations of state that are assumed both for hadronic and quark matter;
2. the method of the PT description such as Maxwellian, Gibbs (total thermodynamic

equilibrium if Coulomb interactions are neglected), or as possible intermediate combinations (for a detailed discussion of various cases see Hempel et al. 2009). In reality, of course, this freedom of choosing the PT type is just because of our current lack of knowledge about important parameters of microphysical interaction among the phases such as surface tension (see e.g. Maruyama et al. 2007);

3. the state of matter in the particular physical scenario considered, i.e., the values of the thermodynamic state parameters that are reached; first of all the entropy and the lepton number, and second the degree of matter transparency for neutrinos.

In the near future we plan to investigate the above components that have an effect on the inverse convection and the possible influence of such convection on the subsequent evolution of hybrid stars.

DISCUSSION AND CONCLUSION

First, we need to emphasize that the possibility of “inverse” convection discussed here is the direct consequence of the unusual property of the deconfinement PT expressed by Eq. (1). For example, for the nuclear liquid-gas PT the condition for convection will have the ordinary form as for single phases of quark and hadronic matter. For a comparison of these two PT, see, e.g., Hempel et al. (2013).

The consequences of possible existence of the “inverse” convection zone in a hybrid star can be conceived by looking at Fig. 1. In the case of well developed convection an ascending matter element that travels from A to A' has a lower entropy than the environment and thus a lower heat content. Similarly, an descending element has a higher heat content than the environment. Therefore in contrast to ordinary convection, within the “inverse” convection zone the heat flux is directed inwards in a star.

Let us consider the situation when the new quark phase appears inside a hot protoneutron star, formed shortly after the bounce of the core of a collapsing massive star (see more details about the collapse and its connection with supernova explosions in Sagert et al. 2009 and Fischer et al. 2011). We consider that the conversion of hadronic into quark matter proceeds on a timescale much faster than the hydrodynamic evolution of the proto-neutron star (for other scenarios, see the recent review by Drago and Pagliara 2015 and references therein). In this case, every matter element, transformed from the

hadron phase to the quark one keeps its entropy per baryon S and consequently decreases its temperature (see our Fig. 1 and also fig. 4 in Steiner et al. 2000). Thus, the temperature of the hybrid star quark core turns out to be lower than that of the surrounding hadronic envelope. After the inverse convection has set in, the heat flux is directed inwards, being transported by “inverse” convective streams inside the phase coexistence domain. This could also be accompanied by an inward neutrino flux which will try to remove the temperature difference between the quark core and the envelope, but this process requires special investigation.

After thermal relaxation the temperatures of the quark and hadron phases become similar. However, the entropy of the quark phase S_q exceeds that of the hadron phase S_h . Hence, within the region of phase coexistence we have the negative entropy gradient that guarantees convective stability under the condition of the “inverse” convection. After such internal thermal relaxation the subsequent thermal evolution of the hybrid star probably follows the standard picture (see, e.g. Nakazato et al. 2013). The detailed study of the thermal processes in question is of considerable interest and may tell a lot about the birth and thermal evolution of hybrid stars.

Recently Roberts et al. (2012) showed that different symmetry energies of the nuclear EOS induce different strengths of (normal) convection inside the cooling proto-neutron star after a core-collapse supernova explosion. The corresponding neutrino signal would allow to discriminate different hadronic EOS. It would be very interesting to investigate the neutrino signal of the “inverse” convection proposed here, and whether or not it contains a particular fingerprint of quark matter.

Currently, the neutrino-driven scenario is the favored explosion mechanism of core-collapse supernovae. In this scenario turbulence and convection are crucial to achieve sufficient neutrino heating of matter to trigger an explosion. If the “inverse” convection appeared already in the early post-bounce phase of the supernovae, therefore in principle it could also impact the explosion dynamics. These possibilities require further investigation by detailed numerical simulations.

Possible direct effects of the pressure decrease in the region of phase-coexistence on the stability of proto-neutron stars will be the subject of a forthcoming study by some of the authors.

Acknowledgements

The authors are thankful to the Swiss National Science Foundation (SNSF) for support (grant SCOPES No. IZ73Z0 152485). A.Y. thanks also the RF Government grant No. 11.G34.31.0047. M.H. is supported by the SNSF and partial support comes from NewCompStar, COST Action MP1304. We also would like to thank the anonymous referee for his/her constructive suggestions.

REFERENCES

1. A. Drago, G. Pagliara, arXiv:1509.02134, (2015)
2. T. Fischer, I. Sagert, G. Pagliara, M. Hempel, J. Schaffner-Bielich, T. Rauscher, F.-K. Thielemann, R. Kappeli, G. Martinez-Pinedo, M. Liebendörfer, *Astrophys. J. Suppl. Ser.*, **194**, 39, 28 pp., (2011)
3. N. K. Glendenning, *Phys. Rev. D*, 46, 4, pp. 1274-1287, (1992)
4. I. Gurevich, *Nature*, **144** 326-327, (1939)
5. P. Haensel, A. Y. Potekhin, D. G. Yakovlev, *Neutron Stars 1 (Equation of State and Structure)*, Springer, 619pp, (2007)
6. M. Hempel, V. Dexheimer, S. Schramm, I. Iosilevskiy, *Phys. Rev. C*, **88**, 1, (2013)
7. M. Hempel, G. Pagliara, and J. Schaffner-Bielich, *Phys. Rev. D*, **80**, 12, (2009)
8. I. Iosilevskiy, arXiv:1403.8053 [physics.plasm-ph], (2014)
9. N. Itoh, *Prog. Theor. Phys.*, **44**, 291, (1970)
10. D. D. Ivanenko, D. F. Kurdgelaidze, *Astrofizika*, **1**, No. 4, 479–482, (1965)
11. Landau L.D. and Lifshitz E.M., *Statistical Physics*, Part 1, 3d Edition, Oxford: Pergamon Press, (1980)
12. T. Maruyama, S. Chiba, H-J. Schulze, and T. Tatsumi, *Phys. Rev. D* 76, 12, (2007)
13. K. Nakazato, K. Sumiyoshi, H. Suzuki, T. Totani, H. Umeda, S. Yamada, *Astrophys. J. Suppl. Ser.* **205**, 1, (2013)
14. L. F. Roberts, G. Shen, V. Cirigliano, J. A. Pons, S. Reddy, S. E. Woosley, *Phys. Rev. Lett.*, 108, 061103, (2012)
15. I. Sagert, T. Fischer, M. Hempel, G. Pagliara, J. Schaffner-Bielich, A. Mezzacappa, F.-K. Thielemann, M. Liebendörfer, *Phys. Rev. Lett.*, **102**, 081101, (2009)
16. H. Shen, H. Toki, K. Oyamatsu, K. Sumiyoshi, *Prog. Theor. Phys.*, **100**, 1013-1031, (1998)
17. H. Shen, H. Toki, K. Oyamatsu, K. Sumiyoshi, *Astrophys. J. Suppl.*, **197**, 20, (2011)
18. A. Steiner, M. Prakash, J. M. Lattimer, *Phys. Rev. B*, **486**, 239-248, (2000)
19. K. Thorne, *Astrophys. J.*, **144**, 201, (1966)

20. R.C. Tolman “Relativity Thermodynamics and Cosmology”, Oxford at the Clarendon Press, (1969)
21. A.V. Yudin, T.L. Razinkova, D.K. Nadyozhin Astronomy Letters, **39**, 161, (2013)