Forecasting Electricity Spot Prices using Lasso: On Capturing the Autoregressive Intraday Structure

Florian Ziel

January 26, 2016

Abstract

In this paper we present a regression based model for day-ahead electricity spot prices. We estimate the considered linear regression model by the lasso estimation method. The lasso approach allows for many possible parameters in the model, but also shrinks and sparsifies the parameters automatically to avoid overfitting. Thus, it is able to capture the autoregressive intraday dependency structure of the electricity price well. We discuss in detail the estimation results which provide insights to the intraday behavior of electricity prices. We perform an out-of-sample forecasting study for several European electricity markets. The results illustrate well that the efficient lasso based estimation technique can exhibit advantages from two popular model approaches.

Introduction and motivation

In day-ahead electricity spot price forecasting there exists a vast range of differently structured models. A recent overview with a classification is given in Weron (2014).

Most of the day-ahead electricity spot prices are traded on an hourly (or half-hourly) basis. This leads to the obvious fact, that we have 24 (or 48) observations a day if we ignore the clock change issue. Subsequently we only consider hourly traded prices, but for half-hourly data the adjusted facts hold as well. All electricity price models rely heavily on this fact of having 24 observations a day. These models can be divided into two classes independent of whether the model is designed e.g. for point or for probabilistic forecasts.

Modeling approaches which belong to the first class construct 24 separate models, one model for each hour of the day. These models can be e.g. a standard time series method (e.g. Paraschiv et al. (2014), Nowotarski and Weron (2014), Maciejowska and Weron (2015)) or a regime switching model (e.g. Karakatsani and Bunn (2008b)). In general, the model approach is usually motivated by the fact that there is one auction each day (mostly in the morning or at lunch time) for the next 24 hours¹. Hence, we observe the 24 traded prices at once. Thus, modeling a 24-dimensional process by 24 small models is an obvious way to proceed.

The second class contains electricity spot price models that propose a single model for the full observed data. Here a 24 hours ahead forecast is computed. This approach is motivated by seeing the price process as one time series that evolves over time as related processes like electricity load or renewable energy generation. As in the first model class it can be of various model types, like time series based approaches (e.g. Haldrup et al. (2010), Cruz et al. (2011), Keles et al. (2012), Paraschiv et al. (2015) or Ziel et al. (2015a)) or machine learning models (e.g. Conejo et al. (2005), Singhal and Swarup (2011) or Chaâbane (2014)).

From the applied point of view both model approaches are valid as they can have a good forecasting performance in particular situations. Nevertheless, the first one seems to be more plausible, as it matches more directly the data generating process. However, in the first class with 24 models for each hour we have usually less observations and consequently 24 simple and low parametrized models (see e.g. Weron and Misiorek (2008) or Maciejowska et al. (2015)). More crucially, most methods assume the same model structure for all 24 hours. The model structure is often assumed in such a way, that the price at hour *i* depends on the price of the previous day at hour *i*, but not on a different hour *j*. Many models do not allow for such a dependency to simplify the model structure, even though there might be strong dependency present. Here models of the second class can have an advantage. As they use more observations for the estimation the resulting model tends to have more involved parameters that can theoretically capture the dependence, if the model design allows for it. Interestingly the vast majority of machine learning models (esp. artificial neural networks) fall into this second model class that estimates one big model, see e.g. Conejo et al. (2005), Singhal and Swarup (2011), Catalao et al. (2011), Shafie-Khah et al. (2011), Hu and Taylor (2014) or Abedinia et al. (2015)). It seems that these estimation techniques rely much more on the fact of having a lot of observations than time series based approaches.

To better motivate and understand this dependency problem we introduce $P_{d,h}$ as electricity price at day d and hour h. Throughout the paper we ignore we clock change issue, and interpolate the missing 2am hour in March and average the double occurring 2am hour in October so that there are 24 observable prices each day.

Now we have a closer look at the linear dependency structure between the price $P_{d,h}$ at day d and hour h and the price $P_{d-1,l}$ at the previous day d-1 and hour l. The corresponding sample correlations of the correlations

¹Once a year for the last Sunday March only 23 and for last Sunday in October 25 hours due to daylight saving time

 $\mathbb{C}or(P_{d,h}, P_{d-1,l})$ are exemplary visualized in Figure (1) for two selected markets, namely the EPEX spot price for Germany and Austria, and the APX spot price for Netherlands.



Figure 1: Sample correlations of $\mathbb{C}or(P_{d,h}, P_{d-1,l})$ for $h, l = 0, \dots, 23$ and two markets from 17.12.2009 to 12.08.2014.

The highlighted diagonal elements measure the linear dependency between the electricity price at hour h to its previous day value at the same hour h. We observe relatively high correlations on the diagonal. But, only for some hours in the night, the late afternoon and the evening they are the largest of each row. For most of the hours from midnight to the afternoon this is not the case. They exhibit their largest correlation with hours in the late evening. The most distinct lagged cross-hour dependency concerns $Y_{d,0}$. It depends much more on the last hour price $Y_{d-1,23}$ of the previous day than on the same hour price $Y_{d-1,0}$.

It is clear that this complex dependency structure should be covered by an appropriate spot price model. As mentioned before, most of the models of the first class do not capture this dependency structure, whereas the second one in principle does. But, the second class that uses one big model has the problem, that e.g. the forecast $\hat{Y}_{d+1,23}$ will depend on the previous hour predictions $\hat{Y}_{d+1,0}, \ldots, \hat{Y}_{d+1,22}$. Thus the prediction error $\hat{\varepsilon}_{d+1,h} = Y_{d+1,h} - \hat{Y}_{d+1,h}$ tend to increase in h as we have to compute 24 hours ahead forecast. And even though models of the second class can capture the cross-hour dependency, we observe that the full dependency pattern of Figure 1 is time-varying. Thus, it is required for models of the second class to utilize time-varying parameters for the autoregressive effect. This is usually not included in many models of this model class, see e.g. Conejo et al. (2005), Singhal and Swarup (2011), Catalao et al. (2011), Shafie-Khah et al. (2011), Hu and Taylor (2014) or Abedinia et al. (2015).

Interestingly, when we compare forecasting performance of relatively simple models of both model classes, models of the second class perform better for the first half of the day, whereas models of the first class are better in the second half of the day. This observation is also part of the results of the empirical section. Nevertheless, it is not clear which modeling method is superior in terms of forecasting performance.

In this paper we introduce a simple linear model approach that falls into the first model class. It contains 24 simple structured models and captures the mentioned dependency pattern directly. Thus, it joins the advantages of both model classes which leads to an improved overall forecasting performance. The model captures the full linear autoregressive dependency structure as seen in Figure 1, especially time-varying cross-hour dependencies. Furthermore, it also allows for a time-varying day-of-the-week effect that will be discussed more detailed in the next section. In electricity price forecasting literature such flexibility is a clear novelty even though there are models available that cover the same feature. For example, an autoregressive principal component analysis based approach as used in Maciejowska and Weron (2013) and Maciejowska and Weron (2015) can cover this behavior. However, there the impact is not modeled directly but indirectly through the underlying dimension reduction approach.

We restrict the model to these two modeling aspects (time-varying cross-hour dependencies and time-varying dayof-the-week effects) to highlight the advantage of the modeling and estimation approach. Especially, we do not focus on well known features like volatility clustering, non-linearity effects (esp. price spikes), long-term trends, structural breaks, public holidays effects, effects of other external regressors like renewable energy feed-in, or the day-light saving time effect. However extensions to all these directions are possible. We discuss this issue briefly in the model section and final discussion.

The crucial part of the estimation relies on high-dimensional statistics. We utilize the lasso (least absolute shrinkage and selection operator) method introduced by Tibshirani (1996), as a regularized estimation technique. Due to the shrinkage properties of the estimators we can handle models with many potential parameters. As the estimation procedure has the sparsity property, many of the potential parameters are not included in the finally estimated model. This property is great to avoid the overfitting problem.

The considered estimation technique is very fast. The estimation and forecasting of the 24 models is feasible in a few seconds without big effort. Thus, the model is also suitable as being an element of a forecast combination approach, see e.g. Bordignon et al. (2013) or Maciejowska et al. (2015). For such applications fast estimation methods are clearly preferable.

The lasso based approach is not completely new in electricity price forecasting literature. It is used by Ziel et al. (2015a) and Ludwig et al. (2015) for electricity price forecasting. Both models use the variable selection property, but consider a model of the second class and share the mentioned disadvantages. The approach of Ziel et al. (2015a) uses time-varying coefficients to capture the mentioned time-varying changes in the dependency structure. However, the time-varying coefficients are modeled using B-splines, that only allow for smooth changes over the day. In contrast, our model belongs to the first model class and allows for more flexibility and abrupt changes in the cross-hour dependency. Additionally, Ziel et al. (2015a) incorporate the impact of load and wind and solar net feed-in such as volatility clustering, public holiday and daylight saving time effects. In Ludwig et al. (2015) the parameters are assumed to be constant over time, so the lasso estimation works just as features selection and ignores time-varying dependencies. They use wind speed and temperature data to model exogenous impacts. In recent research of Dudek (2016) the lasso approach is applied to short-term electricity load forecasting.

The paper is structured as follows: In the second section we introduce the time series model and estimation method. In the third one we discuss several benchmarks for the empirical study which is performed in the fourth section. For the out-of-sample sample forecasting study we consider 10 different hourly European day-ahead electricity spot prices. We close with a summary and some conclusions.

$\mathbf{2}$ Time series model and estimation technique

The time series model that we consider is an autoregressive regression approach, that is similar to the one used in Weron and Misiorek (2008) or Maciejowska et al. (2015). In Weron and Misiorek (2008) the electricity price at hour hdepends linearly on the price at the same hour lagged by 1, 2 and 7 days, such as dummies on the weekdays Sunday, Monday and Saturday. However, the choice of the lags 1, 2 and 7 such as the selection of the weekday dummies is the same for all 24 hours and it is very restrictive. Concerning the weekday structure it is advisable to take a closer look at the weekly mean price of the electricity price. The weekly sample mean of the German/Austrian EPEX spot price and the Dutch APX spot price is given in Figure 2.



(a) EPEX Germany and Austria

Figure 2: Weekly sample mean of $P_{d,h}$ for two markets.

There we see the $7 \times 24 = 168$ sample mean values grouped by the 7 weekdays and 24 hours of the day. We can observe that the sample mean of the Sunday and Saturday values is clearly different from the mean on working days. On the working days from Monday to Friday the mean behavior is relatively similar. But for the morning hours on Monday and the afternoon and evening hours on Friday we observe clear deviations. This transition towards and from the weekend is known for the German/Austrian EPEX and modeled by Ziel et al. (2015a) by an approach of the second model class using time-varying parameters. Most importantly it suggests that the impact of the weekday

dummies in an autoregressive regression model might be different for different hours of the day. Hence, we require flexibility in the model that is able tackle these possible changes in the day-of-the-week dependency structure.

For the time series model that we introduce for the electricity price $P_{d,h}$ we assume that the processes $(P_{d,h})_{d\in\mathbb{Z}}$ are stationary for each hour h. Furthermore, we do not directly model $P_{d,h}$ itself, but its zero mean version $Y_{d,h} = P_{d,h} - \mu_h$ with $\mu_h = \mathbb{E}(P_{d,h})$. We can estimate μ_h easily by the sample mean.

To model the mentioned weekday effects we define the weekday indicators

$$W_k(d) = \begin{cases} 1 & , \mathcal{W}(d) = k \\ 0 & , \mathcal{W}(d) \neq k \end{cases}$$

Here $\mathcal{W}(d)$ is a function that gives a number which represent each day of the week. We use without loss of generality the enumeration k = 0 for a Sunday, k = 1 for a Monday, ..., k = 6 for a Saturday. As we want to point out the effect of the weekdays, we propose two models, one without the weekday effects, the other one with possible weekday effects.

For each hour h the two considered time series models for $Y_{d,h}$ are given by

$$Y_{d,h} = \sum_{l=1}^{24} \sum_{k \in \mathcal{I}(h,l)} \phi_{h,k,l} Y_{d-k,l} + \varepsilon_{d,h} \quad \text{and}$$
(1)

$$Y_{d,h} = \sum_{l=1}^{24} \sum_{k \in \mathcal{I}(h,l)} \phi_{h,k,l} Y_{d-k,l} + \sum_{k=0}^{6} \psi_{h,k} W_k(d) + \varepsilon_{d,h}$$
(2)

with parameters $\phi_{h,k,l}$ and $\psi_{h,k}$, $\mathcal{I}(h,l)$ as an index set of possible lags and $\varepsilon_{d,h}$ as error term. We assume that the error process $(\varepsilon_{d,h})_{d\in\mathbb{Z}}$ is uncorrelated with constant variance σ_h^2 . The parameters $\phi_{h,k,l}$ model the linear autoregressive impact of the past prices and $\psi_{h,k}$ a weekday effect.

The index sets $\mathcal{I}(h, l)$ are very important for the model. They specify the possible model structure. Of course larger sets $\mathcal{I}(h, l)$ increase the probability of overfitting, even though the used regularization technique reduces this problem a lot. If we choose $\mathcal{I}(h, l)$ too small we might miss important information in the dependency structure. Therefore we should choose $\mathcal{I}(h, l)$ reasonably, in the sense that we should only include those lags where we, as a modeler, think that a dependence is plausible. In our situation we choose

$$\mathcal{I}(h,l) = \begin{cases} \{1,\dots,36\} & , h = l \\ \{1,\dots,8\} & , h \neq l \end{cases}.$$
(3)

Thus, we include a possible relationship between $Y_{d,h}$ and the previous 36 days (five weeks plus a day) at the same hour h, but also a dependence to the past 8 days of the electricity prices of other hours.

Furthermore, model equation (2) (also (1)) is a regression model, so it is very modular. It is extremely easy to add new regressors, like lagged electricity load or renewable energy feed-in, seasonal cycles, trends, or event dummies. These additional regressors can be used to cover some of the ignored stylized facts, like long-term trends, structural breaks, public holidays and day-light saving time effects. Even non-linear impacts in $Y_{d-k,h}$ like in a threshold-AR process are possible, as long as the relationship is linear in the parameters. Another direction of extensions concerns the volatility of the residuals. Ziel et al. (2015a) and Ziel (2015) show that an extension to AR-ARCH type processes is possible as well, using an iterative lasso estimation technique.

For the estimation of the parameters in (1) or (2) we will use the corresponding ordinary least square (OLS) representation. We denote it by

$$Y_{d,h} = \mathbf{X}'_{d,h} \boldsymbol{\beta}_h + \varepsilon_{d,h}.$$
(4)

with the vector of regressors $\mathbf{X}_{d,h} = (\mathbf{X}_{d,h,1}, \dots, \mathbf{X}_{d,h,m_h})'$ of length m_h and parameter vector $\boldsymbol{\beta}_h = (\beta_{h,1}, \dots, \beta_{h,m_h})'$ of length m_h . Furthermore, we require for the lasso estimation technique the standardized version of (4). Thus, we introduce it with

$$\widetilde{Y}_{d,h} = \widetilde{X}'_{d,h}\widetilde{\beta}_h + \widetilde{\varepsilon}_{d,h}.$$
(5)

Here $Y_{d,h}$ and the elements of $X_{d,h}$ are scaled in such a way that they have a variance of 1. In the empirical part we scale using the corresponding sample standard deviations. If we have $\tilde{\beta}_h$ we can calculate the parameter vector β_h of (4) easily by rescaling.

The mentioned lasso estimation technique is a penalized regression approach that penalizes the parameters on the sum of the absolute values. Let D be the number of observable days. Then the lasso estimator $\hat{\beta}_h$ for $\hat{\beta}_h$ is given by

$$\widehat{\widetilde{\boldsymbol{\beta}}}_{h} = \operatorname*{arg\,min}_{\boldsymbol{\beta} \in \mathbb{R}^{m_{h}}} \sum_{d=1}^{D} \left(\widetilde{Y}_{d,h} - \widetilde{\boldsymbol{X}}_{d,h}^{\prime} \boldsymbol{\beta} \right)^{2} + \lambda_{h} \sum_{i=1}^{m_{h}} |\beta_{i}|$$
(6)

where $\lambda_h \geq 0$ is a tuning parameter. Note that for $\lambda_h = 0$ we receive the standard OLS estimator. For extremely huge tuning parameters λ_h the estimator will return the estimate $\hat{\beta}_h = \mathbf{0} = (0, \dots, 0)'$ where all parameters are shrinked to zero. In general the lasso estimator is a biased estimator. However, it can be tuned in such a way that the out-of-sample performance is better than the OLS estimator performance. The lasso estimator has an automatic sparsity property, so many of the potential parameters are not included in the finally estimated model. The larger λ_h the less parameters are included in the estimated model. But lasso does not only sparsify, it also regularizes the model. So the estimated parameter that are not estimated to be exactly zero are usually smaller (in absolute values) than the corresponding OLS solution with the same parameters. To be precise: the in-sample residual sum of square (RSS) is larger for the lasso solution than for the corresponding OLS solution with the same non-zero parameters. From a Baysian point of view lasso acts as a prior on the parameters that is centered at zero. For more information on the lasso algorithm from the theoretical and applied point of view we recommend the recently published lasso overview book Hastie et al. (2015).

As estimation algorithm the very fast coordinate descent algorithm of Friedman et al. (2007) is available. It solves the lasso problem on a given grid Λ of λ -values that is commonly chosen as a exponential grid. For this exponential grid Λ it is important that the corresponding solutions cover the full range from the $\hat{\beta} = 0$ solution to a solution that contains all possible parameters as the OLS solution. In detail we define the grid as $\Lambda = \{2^k | k \in \mathcal{G}\} \cup \{0\}$ where \mathcal{G} is an equidistant grid from 4 to -15 of length 50. We use the R package glmnet for implementation, see e.g. Friedman et al. (2010) for more details.

We choose the tuning parameters $\lambda_h \in \mathbf{\Lambda}$ for each hour h by minimizing the Bayesian information criterion (BIC) for each of the 24 single models. The BIC is a very conservative information criteria that avoids overestimation. However other information criteria, like the Akaike information criterion (AIC) or cross-validation based criteria, are possible as well.

3 Benchmarks models

As benchmarks we allow only those processes that model linear dependency structure in the data. These models are usually those ones that can be estimated and forecasted in a very fast way, like the proposed model with the corresponding estimation method. Furthermore, we take suitable representatives of the introduced two model classes. As we consider the lasso based model without (1) and with (2) day-of-the-week effects, we consider for the benchmark models a similar weekday modeling extension to keep comparability.

3.1 24 univariate AR(p) models

The model of 24 univariate AR(p) models is given by

$$P_{d,h} = \phi_{h,0} + \sum_{k=1}^{p} \phi_{h,k} (P_{d-k,h} - \phi_{h,0}) + \varepsilon_{d,h}$$
(7)

where $(\varepsilon_{d,h})_{d\in\mathbb{Z}}$ is i.i.d. with $\varepsilon_{d,h} \sim N(0, \sigma_h^2)$. It models the linear autoregressive dependence structure of each price to the corresponding hour well. We estimate the model by solving the Yule-Walker equations, which solution is guaranteed to be stationary. We select the model orders p_h by minimizing the Akaike information criteria (AIC) given a maximal upper bound of p_h by $p_{\text{max}} = 50$. For the model version with weekday effects we consider the equation

$$P_{d,h} = \phi_{h,0} + \sum_{j=0}^{6} \psi_{h,j} W_j(d) + \sum_{k=1}^{p} \phi_{h,k} \left(P_{d-k,h} - \sum_{j=0}^{6} \psi_{h,j} W_j(d-k) - \phi_{h,0} \right) + \varepsilon_{d,h}.$$
(8)

We estimate model (8) in a two-step approach. First we estimate the day-of-the-week coefficients $\psi_{h,j}$ by OLS and solve the Yule-Walker equations for the resulting residuals. As for model (7) we minimize the AIC with an upper bound for the order p_h of $p_{\text{max}} = 50$. Note that model (8) is just one possibility to include the day-of-the-week effects. Other ones are possible as well, but give similar results. We consider the chosen option as it allows for the application of the Yule-Walker equations as equivalently done for (7).

3.2 Autoregressive expert models

As mentioned in many applications, like in Weron and Misiorek (2008) or Maciejowska et al. (2015), simple autoregressive models with a fixed regression structure are used. As the assumed regression structure is usually build on some prior knowledge of experts, we refer these models as expert models. We consider two expert models as benchmarks, a simple expert model and a weekday based expert model. They are given by

$$P_{d,h} = \phi_{h,0} + \phi_{h,1}P_{d-1,h} + \phi_{h,2}P_{d-2,h} + \phi_{h,7}P_{d-7,h} + \varepsilon_{d,h},$$

$$P_{d,h} = \phi_{h,0} + \phi_{h,1}P_{d-1,h} + \phi_{h,2}P_{d-2,h} + \phi_{h,7}P_{d-7,h} + \psi_{0}W_{0}(d) + \psi_{1}W_{1}(d) + \psi_{6}W_{6}(d) + \varepsilon_{d,h}.$$
(9)
$$(10)$$

The latter one matches the first one but adds linear weekday effects on Sunday, Monday and Saturday. Here we assume for the error process again that $(\varepsilon_{d,h})_{d\in\mathbb{Z}}$ is i.i.d. with $\varepsilon_{d,h} \sim N(0, \sigma_h^2)$. We estimate the parameters by solving the corresponding least square problem. The chosen model of Weron and Misiorek (2008) is just a representative of an expert model. There are several other versions available as well, they tend to slightly differ in some model components. For example Karakatsani and Bunn (2008a) propose a model with a linear price dependency on a daily lag of order 1 and 7, but not at lag 2. Note that model (9) is nested in model (7), which is nested in the proposed model (1). Thus, we expect similar behavior, if the lags of the nested models capture the relevant features. These models with a simpler structure have the advantage of being easier to interpret and implement. Furthermore, the overestimation issue is negligible.

3.3 Univariate AR(p) models

This is a model of the second model class. Ziel et al. (2015a) and Ziel et al. (2015b) show that the univariate AR(p) model can be a powerful forecast method. It performs often even better than several modern models. Denote $P_t = P_{d,h}$ with t = 24d + j for $j \in \{0, \ldots, 23\}$ a univariate time series representation of electricity price at day d and hour h. The considered model and its weekday effect version are given by

$$P_{t} = \phi_{0} + \sum_{k=1}^{p} \phi_{k} (P_{t-k} - \phi_{0}) + \varepsilon_{t},$$
(11)

$$P_t = \phi_0 + \sum_{j=0}^{6} \psi_j W_j(\tilde{d}(t)) + \sum_{k=1}^{p} \left(\phi_k P_{t-k} - \sum_{j=0}^{6} \psi_j W_j(\tilde{d}(t) - k) - \phi_0 \right) + \varepsilon_t$$
(12)

where $(\varepsilon_{d,h})_{d\in\mathbb{Z}}$ is i.i.d. with $\varepsilon_{d,h} \sim N(0, \sigma_h^2)$ and $\tilde{d}(t) = \lfloor t/24 \rfloor$ gives the day d that corresponds to time point t. Again we estimate the model (11) by solving the Yule-Walker equations and minimize the AIC with respect to a maximal order of $p_{\max} = 700$. Note that a maximal oder larger than 168 is required to capture any weekly dependency structure. As for model (8) we estimate the weekday version (12) by the two-step approach. We solve the OLS regression for the day-of-the-week parameters and solve the Yule-Walker equations of the residuals by minimizing the AIC with maximal order of $p_{\max} = 700$.

3.4 PCA based autoregressive models

The model with 24 univariate AR(p) models is a special case of the 24-dimensional AR(p) process given by

$$\boldsymbol{P}_{d} = \boldsymbol{\phi}_{0} + \sum_{k=1}^{p} \boldsymbol{\Phi}_{k} (\boldsymbol{P}_{d-k} - \boldsymbol{\phi}_{0}) + \boldsymbol{\varepsilon}_{d}$$
(13)

with 24-dimensional intercept vector ϕ_0 , 24×24-dimensional autoregressive parameter matrices Φ_k , $P_d = (P_{d,0}, \ldots, P_{d,23})'$ and $\varepsilon_d \sim N_{24}(\mathbf{0}, \mathbf{\Sigma})$. Such a general 24-dimensional AR(p)-process has $24 + 24 \times 24 \times p = 24 + 576p$ parameters. This is usually quite a lot and leads quickly to models with too many parameters. Note that model (13) is able to capture the lagged cross-hour dependency structure as seen in Figure 1. The proposed model (1) is nested in (13) (and vice versa) with an appropriate choice of $\mathcal{I}(h, l)$. As the lasso estimation technique overcomes the overestimation problem the same features can be covered. Another approach that deals with this over-parametrization problem is principal component analysis (PCA) based.

The approach is based on a dimension reduction of the 24-dimensional process by PCA. Maciejowska and Weron (2013) and Maciejowska and Weron (2015) use it and a generalization to forecast the UK APX electricity spot prices. They show that the PCA approach can be suitable to capture the intraday relationship between the electricity prices.

The principal component decomposition of \boldsymbol{P}_d is given by

$$\boldsymbol{P}_d = \boldsymbol{\Gamma} \boldsymbol{F}_d \tag{14}$$

with Γ as 24×24-dimensional loading matrix and F_d as factor scores at day d. As usual in PCA analysis we assume that without loss of generality the columns of Γ are ordered by the absolute value of their corresponding eigenvalue. For the dimension reduction we consider only the first K principal components of the scores which we denote by $F_{K,d}$. We denote the corresponding $K \times 24$ -dimensional reduced loading matrix by Γ_K .

For forecasting the scores we employ a multivariate AR (VAR) model to the K-dimensional vector $\mathbf{F}_{K,d}$. Thus, we assume that the factor scores follow a process that is given by

$$\boldsymbol{F}_{K,d} = \boldsymbol{\phi}_{K,0} + \sum_{k=1}^{p} \boldsymbol{\Phi}_{K,k} (\boldsymbol{F}_{K,d-k} - \boldsymbol{\phi}_{K,0}) + \boldsymbol{\varepsilon}_{K,d}$$
(15)

with K-dimensional intercept vector $\phi_{K,0}$, $K \times K$ dimensional autoregressive parameter matrices $\Phi_{K,k}$ and $\varepsilon_{K,d}$ as K-dimensional zero-mean error term vector. Now the dimension reduced model has only $K + K \times K \times p$ parameters.

As it holds $P_d = \Gamma_K \phi_{K,0} + \sum_{k=1}^p \Gamma_K \Phi_{K,k} (F_{K,d-k} - \phi_{K,0}) + \Gamma_K \varepsilon_{K,d}$ model (15) is a special case of model (13), but also of the proposed model (1) if $\mathcal{I}(h, l)$ is chosen correctly.

Again we estimate model (15) by solving the multivariate Yule-Walker equations and minimizing the AIC with respect to a maximal order of $p_{\max} = 50$. We perform the estimation for $K \in \{2, ..., 12\}$. Given a forecast $\hat{F}_{K,d}$ of the factor scores for day d and an estimate $\hat{\Gamma}$ of the loading matrix we can easily get the price forecast by $\hat{P}_d = \hat{\Gamma}_K \hat{F}_{K,d}$ using the plug-in principle.

The day-of-the-week version of model (15) is given by

$$\boldsymbol{F}_{K,d} = \phi_{K,0} + \sum_{j=0}^{6} \psi_j W_j(d) + \sum_{k=1}^{p} \left(\boldsymbol{\Phi}_{K,k} \boldsymbol{F}_{K,d-k} - \sum_{j=0}^{6} \psi_j W_j(d-k) \right) + \boldsymbol{\varepsilon}_{K,d}$$
(16)

where ψ_j are the K-dimensional vectors of weekday parameter coefficient, and $\phi_{K,0}$ is again the K-dimensional intercept vector, $\Phi_{K,k}$ the $K \times K$ autoregressive parameter matrices and $\varepsilon_{K,d}$ as K-dimensional error process. As for models (8) and (12) we estimate the day-of-the-week model (12) by the two-step approach. First we solve the OLS problem for the weekday parameters and take the Yule-Walker equations solution of the residuals by minimizing the AIC with maximal possible order of $p_{\max} = 50$ and for $K \in \{2, \ldots, 12\}$.

4 Empirical study and results

We perform an out-of-sample forecast study by using a rolling window. As in-sample period we select always $D = 730 = 2 \times 365$ days (usually two years) of data for the estimation. Given the estimated model we perform an out-of-sample forecast for the next 24 values. The considered data range is from 17.12.2009 to 12.08.2014. Thus, the full out-of-sample period is from 17.12.2011 to 12.08.2014 which is exactly N = 969 days.

As forecasting measure we consider the overall MAE and RMSE, as they are also suitable on markets with negative prices. The MAE is a more robust measure, whereas the RMSE is the optimal measure for least square problems and more sensitive to outliers. They are given by

$$MAE = \frac{1}{24N} \sum_{n=1}^{N} \sum_{h=0}^{23} |P_{D+n,h} - \widehat{P}_{D+n,h}|, \qquad (17)$$

$$RMSE = \sqrt{\frac{1}{24N} \sum_{n=1}^{N} \sum_{h=0}^{23} |P_{D+n,h} - \widehat{P}_{D+n,h}|^2}$$
(18)

where $\widehat{P}_{D+n,h}$ denotes the forecasted price of day D+n. However, to better understand how the models perform over the day, we use the hourly MAE and RMSE at hour h, denoted by MAE_h and RMSE_h. They are defined through

$$MAE_{h} = \frac{1}{N} \sum_{n=1}^{N} |P_{D+n,h} - \widehat{P}_{D+n,h}|, \qquad (19)$$

$$RMSE_{h} = \sqrt{\frac{1}{N} \sum_{n=1}^{N} |P_{D+n,h} - \widehat{P}_{D+n,h}|^{2}}.$$
(20)

We conduct the out-of-sample study for a selected number of electricity markets and all considered models. For the PCA based models in (15) we only report the model with minimal MAE for each market. The selected electricity prices are the German/Austrian EXAA price, the German/Austrian and Swiss EPEX prices, the Belgian BELPEX price, the Dutch APX price, the Danish West, Danish East and Sweden(4) Nordpool prices, the Polish POLPX price and the Czech OTE price. A summary of the considered prices with the subsequently used abbreviations is given in Table 1. Note that all prices are hourly day-ahead electricity prices. For comparison issues we converted all prices in EUR/MWh. If the reported currency is not EUR we consider the corresponding end of day exchange rate of the local currency to convert the prices to EUR/MWh. A selection of different electricity prices is suitable to check if the intraday structure on different electricity markets is similarly or not. It also acts as a robustness check for the proposed lasso based estimation method.

For the PCA-based models the optimal number of factors K are between 3 and 10. They are also listed in Table 1. Note that Maciejowska and Weron (2015) get an optimal PCA-based model with 3 principal components for the half-hourly volume-weighted APX UK spot log-price. They also provide a nice interpretation for the factors.

Subsequently, we denote model (1) by lasso, (7) by 24d.AR, (9) by exp.AR, (11) by AR(p) and (15) by PCA^{*}. For the day-of-the-week model version we add the suffix '-wd' to the model abbreviation. The computed MAE and RMSE values are given in Table 2. There we also report the estimated standard deviations of MAE and RMSE which are computed using residual based bootstrap with a bootstrap sample size of 10000. These standard deviations can be used as significance criterion as in Ziel et al. (2015b). For example if the MAE and RMSE would be normally distributed, then the 1.96-sigma range around the mean matches the symmetric 95% confidence interval. Even though

Exchange	Region	Abbreviation	PCA^*	$\mathrm{PCA}^*\operatorname{-wd}$
EXAA	Germany&Austria	EXAA.DE&AT	7	8
EPEX	Germany&Austria	EPEX.DE&AT	6	10
EPEX	Switzerland	EPEX.CH	8	6
BELPEX	Belgium	BELPEX.BE	6	6
APX	Netherlands	APX.NL	6	8
Nordpool	Denmark (West)	NP.DK.West	7	3
Nordpool	Denmark (East)	NP.DK.East	4	4
Nordpool	Sweden (4)	NP.SW	6	4
POLPX	Poland (Auction I)	POLPX.PL	8	8
OTE	Czech Republic	OTE.CZ	7	9

Table 1: Summary of the considered electricity markets with abbreviations and optimal number of factors for the PCA based models.

the distribution is likely not normal we can use the Chebyshev inequality or extensions to make non-parametric statements about the confidence interval. In Table 2 we underlined all models that are not significantly worse than the best model, where we take the 2-sigma range of the best model as significance indicator.

In Table 2 we observe that concerning the MAE the lasso based approaches perform very well. The proposed model (lasso-wd) in equation (2) with weekdays effects is always the best model, except for the BELPEX and POLPX case. But even there it not significantly worse than the best models which are the lasso without weekday effects for BELPEX and the 24-dimensional AR model with weekday effects for the POLPX. It is remarkable that model (2) with weekdays effects is sometimes significantly better than all other considered models. In several cases (EPEX.AT&AT, EPEX.CH, APX.NL, NP.DK.East, OTE.CZ) there is only one model that is not significantly worse. In contrast, concerning the RMSE values the picture is not that clear. Still, the lasso-wd model (2) is most of the time the best model. However, as mentioned the RMSE is sensitive to outliers. So for example the Nordpool Denmark West price has 5 extreme prices between 1900EUR/MWh and 2000EUR/MWh on June 7, 2013. This leads to a drastic increase in the RMSE, as no (considered) model is able to capture these extreme price spikes. Concerning the RMSE the AR(p)-wd has a relatively low outlier sensitivity, as it is a member of the second model class which contains big single equation models and use all observations as input. Having a lot of observations helps these models to learn that the outliers are a rare event. In contrast, all models of the first class that use 24 small models (as the proposed lasso) have the problem that an outlier gets much more attention as less observations are available.

For the MAE_h and RMSE_h we discuss detailed results only for the German/Austrian EPEX spot price, which is the largest day-ahead electricity market in Europe (regarding the traded volume). The MAE_h and RMSE_h results are given in Figure 3. Results for the other markets are similar. The corresponding graphs are given in the supplementary material. There we also present the correlation plots and the weekly mean graphs as in Figures 1 and 2 for all markets.



Figure 3: Hourly performance measures for EPEX spot price of Germany and Austria.

MAE	lasso	lasso-wd	24d.AR	24d.AR-wd	exp.AR	exp.AR-wd	AR(p)	AR(p)-wd	PCA*	PCA*-wd
EVAA DEPAT	4.89	4.61	5.30	4.96	6.09	5.27	4.87	4.68	5.11	4.72
EAAA.DE&AI	(.033)	(.031)	(.035)	(.034)	(.041)	(.035)	(.033)	(.032)	(.034)	(.032)
FPFY DF&AT	6.46	6.01	6.73	6.32	7.43	6.69	6.39	6.17	6.62	6.09
EI EA.DE&AI	(.050)	(.048)	(.054)	(.052)	(.058)	(.054)	(.050)	(.049)	(.052)	(.050)
EPEX CH	5.44	5.15	5.82	5.36	6.68	5.65	5.37	5.23	5.93	5.38
	(.047)	(.045)	(.048)	(.045)	(.054)	(.047)	(.045)	(.043)	(.052)	(.048)
BELPEX BE	6.75	6.78	6.91	7.19	7.26	7.38	7.21	7.48	7.05	7.27
	(.052)	(.052)	(.053)	(.054)	(.056)	(.056)	(.053)	(.053)	(.052)	(.054)
APX NL	5.11	4.95	5.26	5.04	5.62	5.30	5.13	5.01	5.25	5.03
AI A.NL	(.034)	(.033)	(.035)	(.033)	(.038)	(.035)	(.034)	(.033)	(.034)	(.033)
NP DK Wost	6.89	6.62	7.21	6.95	8.03	7.86	7.61	7.42	6.83	6.78
	(.193)	(.191)	(.187)	(.186)	(.298)	(.328)	(.194)	(.194)	(.188)	(.187)
NP DK East	6.31	6.01	6.42	6.18	6.66	6.40	6.13	6.07	6.50	6.35
MI.DR.East	(.069)	(.067)	(.055)	(.054)	(.057)	(.057)	(.053)	(.052)	(.056)	(.055)
NP SWA	3.28	3.20	3.67	3.60	3.73	3.54	3.48	3.52	3.78	3.74
	(.038)	(.038)	(.039)	(.037)	(.039)	(.038)	(.039)	(.038)	(.039)	(.039)
POLPX PL	2.78	2.68	2.82	2.65	3.19	2.73	2.83	2.77	3.04	2.81
	(.030)	(.029)	(.027)	(.027)	(.030)	(.028)	(.027)	(.027)	(.030)	(.028)
OTE CZ	5.49	5.10	5.67	5.26	6.52	5.63	5.34	5.11	5.60	5.20
011.02	(.039)	(.036)	(.041)	(.039)	(.047)	(.042)	(.038)	(.037)	(.040)	(.038)
RMSE	lasso	lasso-wd	24d.AR	24d.AR-wd	exp.AR	exp.AR-wd	AR(p)	AR(p)-wd	PCA*	PCA*-wd
RMSE EXAA DE&AT	lasso 7.02	lasso-wd 6.63	24d.AR 7.52	24d.AR-wd 7.12	exp.AR 8.70	exp.AR-wd 7.58	AR(p) 7.01	AR(p)-wd <u>6.73</u>	PCA* 7.33	PCA*-wd 6.79
RMSE EXAA.DE&AT	lasso 7.02 (.070)	lasso-wd 6.63 (.067)	24d.AR 7.52 (.068)	24d.AR-wd 7.12 (.066)	exp.AR 8.70 (.081)	exp.AR-wd 7.58 (.068)	AR(p) 7.01 (.067)	$ \begin{array}{r} AR(p)-wd \\ $	PCA* 7.33 (.072)	PCA*-wd 6.79 (.070)
RMSE EXAA.DE&AT	lasso 7.02 (.070) 10.06	lasso-wd 6.63 (.067) 9.47	24d.AR 7.52 (.068) 10.70	24d.AR-wd 7.12 (.066) 10.24	exp.AR 8.70 (.081) 11.63	exp.AR-wd 7.58 (.068) 10.71	AR(p) 7.01 (.067) <u>10.02</u>	AR(p)-wd <u>6.73</u> (.066) <u>9.70</u>	PCA* 7.33 (.072) 10.45	PCA*-wd 6.79 (.070) <u>9.82</u>
RMSE EXAA.DE&AT EPEX.DE&AT	lasso 7.02 (.070) 10.06 (.275)	lasso-wd 6.63 (.067) 9.47 (.282)	24d.AR 7.52 (.068) 10.70 (.308)	24d.AR-wd 7.12 (.066) 10.24 (.323)	exp.AR 8.70 (.081) 11.63 (.284)	exp.AR-wd 7.58 (.068) 10.71 (.301)	$ \begin{array}{r} AR(p) \\ 7.01 \\ (.067) \\ \underline{10.02} \\ (.265) \\ \end{array} $	$\begin{array}{c} AR(p)\text{-wd} \\ \underline{6.73} \\ (.066) \\ \underline{9.70} \\ (.272) \end{array}$	PCA* 7.33 (.072) 10.45 (.296)	PCA*-wd 6.79 (.070) <u>9.82</u> (.324)
RMSE EXAA.DE&AT EPEX.DE&AT	lasso 7.02 (.070) 10.06 (.275) 9.03	lasso-wd 6.63 (.067) 9.47 (.282) <u>8.60</u>	24d.AR 7.52 (.068) 10.70 (.308) 9.35	24d.AR-wd 7.12 (.066) 10.24 (.323) <u>8.72</u>	exp.AR 8.70 (.081) 11.63 (.284) 10.58	exp.AR-wd 7.58 (.068) 10.71 (.301) 9.16	AR(p) 7.01 (.067) <u>10.02</u> (.265) <u>8.77</u>	AR(p)-wd <u>6.73</u> (.066) <u>9.70</u> (.272) 8.48	PCA* 7.33 (.072) 10.45 (.296) 9.93	PCA*-wd 6.79 (.070) <u>9.82</u> (.324) 9.18
RMSE EXAA.DE&AT EPEX.DE&AT EPEX.CH	lasso 7.02 (.070) 10.06 (.275) 9.03 (.220)	lasso-wd 6.63 (.067) 9.47 (.282) 8.60 (.227)	24d.AR 7.52 (.068) 10.70 (.308) 9.35 (.211)	24d.AR-wd 7.12 (.066) 10.24 (.323) <u>8.72</u> (.211)	exp.AR 8.70 (.081) 11.63 (.284) 10.58 (.212)	exp.AR-wd 7.58 (.068) 10.71 (.301) 9.16 (.201)	$\begin{array}{c} AR(p) \\ 7.01 \\ (.067) \\ \underline{10.02} \\ (.265) \\ \underline{8.77} \\ (.208) \end{array}$	AR(p)-wd 6.73 (.066) 9.70 (.272) 8.48 (.210)	PCA* 7.33 (.072) 10.45 (.296) 9.93 (.260)	PCA*-wd 6.79 (.070) <u>9.82</u> (.324) 9.18 (.267)
RMSE EXAA.DE&AT EPEX.DE&AT EPEX.CH BELPEX BE	lasso 7.02 (.070) 10.06 (.275) 9.03 (.220) 10.33	lasso-wd 6.63 (.067) 9.47 (.282) 8.60 (.227) <u>10.37</u>	24d.AR 7.52 (.068) 10.70 (.308) 9.35 (.211) <u>10.58</u>	24d.AR-wd 7.12 (.066) 10.24 (.323) <u>8.72</u> (.211) 10.98	exp.AR 8.70 (.081) 11.63 (.284) 10.58 (.212) 11.21	exp.AR-wd 7.58 (.068) 10.71 (.301) 9.16 (.201) 11.30	AR(p) 7.01 (.067) <u>10.02</u> (.265) <u>8.77</u> (.208) <u>10.79</u>	AR(p)-wd <u>6.73</u> (.066) <u>9.70</u> (.272) 8.48 (.210) 11.02	PCA* 7.33 (.072) 10.45 (.296) 9.93 (.260) <u>10.58</u>	PCA*-wd 6.79 (.070) <u>9.82</u> (.324) 9.18 (.267) 10.90
RMSE EXAA.DE&AT EPEX.DE&AT EPEX.CH BELPEX.BE	lasso 7.02 (.070) 10.06 (.275) 9.03 (.220) 10.33 (.239)	lasso-wd 6.63 (.067) 9.47 (.282) <u>8.60</u> (.227) <u>10.37</u> (.239)	$\begin{array}{c} 24 \text{d.AR} \\ 7.52 \\ (.068) \\ 10.70 \\ (.308) \\ 9.35 \\ (.211) \\ \underline{10.58} \\ (.234) \end{array}$	$\begin{array}{c} 24 \text{d.AR-wd} \\ \hline 7.12 \\ (.066) \\ 10.24 \\ (.323) \\ \hline 8.72 \\ (.211) \\ 10.98 \\ (.243) \end{array}$	exp.AR 8.70 (.081) 11.63 (.284) 10.58 (.212) 11.21 (.232)	exp.AR-wd 7.58 (.068) 10.71 (.301) 9.16 (.201) 11.30 (.231)	$\begin{array}{c} AR(p) \\ 7.01 \\ (.067) \\ \hline 10.02 \\ (.265) \\ \hline 8.77 \\ (.208) \\ \hline 10.79 \\ (.229) \end{array}$	AR(p)-wd <u>6.73</u> (.066) <u>9.70</u> (.272) 8.48 (.210) 11.02 (.222)	$\begin{array}{c} PCA^{*} \\ \hline 7.33 \\ (.072) \\ 10.45 \\ (.296) \\ 9.93 \\ (.260) \\ \hline 10.58 \\ (.210) \end{array}$	PCA*-wd 6.79 (.070) <u>9.82</u> (.324) 9.18 (.267) 10.90 (.215)
RMSE EXAA.DE&AT EPEX.DE&AT EPEX.CH BELPEX.BE	lasso 7.02 (.070) 10.06 (.275) 9.03 (.220) 10.33 (.239) 7.26	lasso-wd 6.63 (.067) 9.47 (.282) <u>8.60</u> (.227) <u>10.37</u> (.239) 7.02	24d.AR 7.52 (.068) 10.70 (.308) 9.35 (.211) <u>10.58</u> (.234) 7.44	24d.AR-wd 7.12 (.066) 10.24 (.323) <u>8.72</u> (.211) 10.98 (.243) 7.14	exp.AR 8.70 (.081) 11.63 (.284) 10.58 (.212) 11.21 (.232) 8.00	exp.AR-wd 7.58 (.068) 10.71 (.301) 9.16 (.201) 11.30 (.231) 7.50	$\begin{array}{c} AR(p) \\ 7.01 \\ (.067) \\ \hline 10.02 \\ (.265) \\ \hline 8.77 \\ (.208) \\ \hline 10.79 \\ (.229) \\ \hline 7.26 \end{array}$	AR(p)-wd <u>6.73</u> (.066) <u>9.70</u> (.272) 8.48 (.210) 11.02 (.222) <u>7.09</u>	PCA* 7.33 (.072) 10.45 (.296) 9.93 (.260) <u>10.58</u> (.210) 7.40	PCA*-wd 6.79 (.070) <u>9.82</u> (.324) 9.18 (.267) 10.90 (.215) <u>7.09</u>
RMSE EXAA.DE&AT EPEX.DE&AT EPEX.CH BELPEX.BE APX.NL	lasso 7.02 (.070) 10.06 (.275) 9.03 (.220) 10.33 (.239) 7.26 (.079)	lasso-wd 6.63 (.067) 9.47 (.282) 8.60 (.227) <u>10.37</u> (.239) 7.02 (.074)	24d.AR 7.52 (.068) 10.70 (.308) 9.35 (.211) <u>10.58</u> (.234) 7.44 (.077)	$\begin{array}{r} 24 \text{d.AR-wd} \\ 7.12 \\ (.066) \\ 10.24 \\ (.323) \\ \hline 8.72 \\ (.211) \\ 10.98 \\ (.243) \\ \hline 7.14 \\ (.073) \\ \end{array}$	exp.AR 8.70 (.081) 11.63 (.284) 10.58 (.212) 11.21 (.232) 8.00 (.087)	exp.AR-wd 7.58 (.068) 10.71 (.301) 9.16 (.201) 11.30 (.231) 7.50 (.076)	$\begin{array}{c} AR(p) \\ \hline 7.01 \\ (.067) \\ \hline 10.02 \\ (.265) \\ \hline 8.77 \\ (.208) \\ \hline 10.79 \\ (.229) \\ \hline 7.26 \\ (.078) \end{array}$	AR(p)-wd 6.73 (.066) <u>9.70</u> (.272) 8.48 (.210) 11.02 (.222) <u>7.09</u> (.076)	$\begin{array}{c} PCA^{*} \\ \hline 7.33 \\ (.072) \\ 10.45 \\ (.296) \\ \hline 9.93 \\ (.260) \\ \hline 10.58 \\ (.210) \\ \hline 7.40 \\ (.081) \end{array}$	$\begin{array}{c} PCA^{*}\text{-wd} \\ \hline 6.79 \\ (.070) \\ \hline 9.82 \\ (.324) \\ \hline 9.18 \\ (.267) \\ \hline 10.90 \\ (.215) \\ \hline 7.09 \\ (.079) \\ \end{array}$
RMSE EXAA.DE&AT EPEX.DE&AT EPEX.CH BELPEX.BE APX.NL	lasso 7.02 (.070) 10.06 (.275) 9.03 (.220) 10.33 (.239) 7.26 (.079) 30.21	lasso-wd 6.63 (.067) 9.47 (.282) <u>8.60</u> (.227) <u>10.37</u> (.239) 7.02 (.074) <u>29.94</u>	24d.AR 7.52 (.068) 10.70 (.308) 9.35 (.211) <u>10.58</u> (.234) 7.44 (.077) <u>29.47</u>	$\begin{array}{r} 24 \text{d.AR-wd} \\ 7.12 \\ (.066) \\ 10.24 \\ (.323) \\ \hline 8.72 \\ (.211) \\ 10.98 \\ (.243) \\ \hline 7.14 \\ (.073) \\ \hline 29.37 \\ \end{array}$	exp.AR 8.70 (.081) 11.63 (.284) 10.58 (.212) 11.21 (.232) 8.00 (.087) 46.27	exp.AR-wd 7.58 (.068) 10.71 (.301) 9.16 (.201) 11.30 (.231) 7.50 (.076) 50.67	$\begin{array}{c} AR(p) \\ \hline 7.01 \\ (.067) \\ \hline 10.02 \\ (.265) \\ \hline 8.77 \\ (.208) \\ \hline 10.79 \\ (.229) \\ \hline 7.26 \\ (.078) \\ \hline 30.69 \end{array}$	AR(p)-wd <u>6.73</u> (.066) <u>9.70</u> (.272) 8.48 (.210) 11.02 (.222) <u>7.09</u> (.076) <u>30.50</u>	$\begin{array}{c} PCA^{*} \\ \hline 7.33 \\ (.072) \\ 10.45 \\ (.296) \\ \hline 9.93 \\ (.260) \\ \hline 10.58 \\ (.210) \\ \hline 7.40 \\ (.081) \\ \hline 29.61 \end{array}$	PCA*-wd 6.79 (.070) <u>9.82</u> (.324) 9.18 (.267) 10.90 (.215) <u>7.09</u> (.079) 29.35
RMSE EXAA.DE&AT EPEX.DE&AT EPEX.CH BELPEX.BE APX.NL NP.DK.West	$\begin{array}{c} \text{lasso} \\ 7.02 \\ (.070) \\ 10.06 \\ (.275) \\ 9.03 \\ (.220) \\ \textbf{10.33} \\ (.239) \\ 7.26 \\ (.079) \\ \hline 30.21 \\ (5.845) \end{array}$	lasso-wd 6.63 (.067) 9.47 (.282) <u>8.60</u> (.227) <u>10.37</u> (.239) 7.02 (.074) <u>29.94</u> (5.884)	$\begin{array}{c} 24 \text{d.AR} \\ 7.52 \\ (.068) \\ 10.70 \\ (.308) \\ 9.35 \\ (.211) \\ \hline 10.58 \\ (.234) \\ 7.44 \\ (.077) \\ \hline 29.47 \\ (5.934) \end{array}$	$\begin{array}{r} 24 \text{d.AR-wd} \\ \hline 7.12 \\ (.066) \\ 10.24 \\ (.323) \\ \hline 8.72 \\ (.211) \\ 10.98 \\ (.243) \\ \hline 7.14 \\ (.073) \\ \hline 29.37 \\ (5.956) \\ \end{array}$	exp.AR 8.70 (.081) 11.63 (.284) 10.58 (.212) 11.21 (.232) 8.00 (.087) 46.27 (7.033)	$\begin{array}{c} \text{exp.AR-wd} \\ \hline 7.58 \\ (.068) \\ \hline 10.71 \\ (.301) \\ 9.16 \\ (.201) \\ \hline 11.30 \\ (.231) \\ \hline 7.50 \\ (.076) \\ \hline 50.67 \\ (8.386) \end{array}$	$\begin{array}{c} AR(p) \\ \hline 7.01 \\ (.067) \\ \hline 10.02 \\ (.265) \\ \hline 8.77 \\ (.208) \\ \hline 10.79 \\ (.229) \\ \hline 7.26 \\ (.078) \\ \hline 30.69 \\ (5.662) \end{array}$	$\begin{array}{c} AR(p)\text{-wd} \\ \hline 6.73 \\ (.066) \\ \hline 9.70 \\ (.272) \\ \hline 8.48 \\ (.210) \\ \hline 11.02 \\ (.222) \\ \hline 7.09 \\ (.076) \\ \hline 30.50 \\ (5.697) \\ \end{array}$	$\begin{array}{c} {\rm PCA}^{*} \\ \hline 7.33 \\ (.072) \\ 10.45 \\ (.296) \\ 9.93 \\ (.260) \\ \hline 10.58 \\ (.210) \\ \hline 7.40 \\ (.081) \\ \hline 29.61 \\ (6.004) \end{array}$	$\begin{array}{r} {\rm PCA^{*}\text{-wd}} \\ \hline 6.79 \\ (.070) \\ \hline 9.82 \\ (.324) \\ \hline 9.18 \\ (.267) \\ \hline 10.90 \\ (.215) \\ \hline 7.09 \\ (.079) \\ \hline 29.35 \\ (6.016) \\ \end{array}$
RMSE EXAA.DE&AT EPEX.DE&AT EPEX.CH BELPEX.BE APX.NL NP.DK.West	$\begin{array}{c} \text{lasso} \\ 7.02 \\ (.070) \\ 10.06 \\ (.275) \\ 9.03 \\ (.220) \\ \hline \textbf{10.33} \\ (.239) \\ 7.26 \\ (.079) \\ \hline \textbf{30.21} \\ (5.845) \\ 12.38 \end{array}$	lasso-wd 6.63 (.067) 9.47 (.282) <u>8.60</u> (.227) <u>10.37</u> (.239) 7.02 (.074) <u>29.94</u> (5.884) 11.92	$\begin{array}{r} 24 \text{d.AR} \\ 7.52 \\ (.068) \\ 10.70 \\ (.308) \\ 9.35 \\ (.211) \\ \hline 10.58 \\ (.234) \\ 7.44 \\ (.077) \\ \hline 29.47 \\ (5.934) \\ \hline 10.59 \end{array}$	$\begin{array}{r} 24 \text{d.AR-wd} \\ \hline 7.12 \\ (.066) \\ \hline 10.24 \\ (.323) \\ \hline 8.72 \\ (.211) \\ \hline 10.98 \\ (.243) \\ \hline 7.14 \\ (.073) \\ \hline 29.37 \\ (5.956) \\ \hline 10.26 \end{array}$	exp.AR 8.70 (.081) 11.63 (.284) 10.58 (.212) 11.21 (.232) 8.00 (.087) 46.27 (7.033) 11.07	exp.AR-wd 7.58 (.068) 10.71 (.301) 9.16 (.201) 11.30 (.231) 7.50 (.076) 50.67 (8.386) 10.83	$\begin{array}{c} {\rm AR(p)}\\ {\rm 7.01}\\ {\rm (.067)}\\ {\rm \underline{10.02}}\\ {\rm (.265)}\\ {\rm \underline{8.77}}\\ {\rm (.208)}\\ {\rm \underline{10.79}}\\ {\rm (.229)}\\ {\rm 7.26}\\ {\rm (.078)}\\ {\rm \underline{30.69}}\\ {\rm (5.662)}\\ {\rm \underline{10.34}} \end{array}$	AR(p)-wd <u>6.73</u> (.066) <u>9.70</u> (.272) 8.48 (.210) 11.02 (.222) <u>7.09</u> (.076) <u>30.50</u> (5.697) 10.06	$\begin{array}{c} {\rm PCA}^{*} \\ \hline 7.33 \\ (.072) \\ 10.45 \\ (.296) \\ 9.93 \\ (.260) \\ \hline 10.58 \\ (.210) \\ \hline 7.40 \\ (.081) \\ \hline 29.61 \\ \hline (6.004) \\ 10.77 \end{array}$	$\begin{array}{r} {\rm PCA^{*}\text{-wd}} \\ \hline 6.79 \\ (.070) \\ \hline 9.82 \\ (.324) \\ \hline 9.18 \\ (.267) \\ \hline 10.90 \\ (.215) \\ \hline 7.09 \\ (.079) \\ \hline 29.35 \\ (6.016) \\ \hline 10.52 \\ \end{array}$
RMSE EXAA.DE&AT EPEX.DE&AT EPEX.CH BELPEX.BE APX.NL NP.DK.West NP.DK.East	$\begin{array}{c} \text{lasso} \\ 7.02 \\ (.070) \\ 10.06 \\ (.275) \\ 9.03 \\ (.220) \\ \textbf{10.33} \\ (.220) \\ \textbf{10.33} \\ (.239) \\ 7.26 \\ (.079) \\ \hline 30.21 \\ (5.845) \\ 12.38 \\ (.917) \end{array}$	$\begin{array}{c} \text{lasso-wd} \\ \hline \textbf{6.63} \\ (.067) \\ \hline \textbf{9.47} \\ (.282) \\ \hline \textbf{8.60} \\ (.227) \\ \hline \underline{10.37} \\ (.239) \\ \hline \textbf{7.02} \\ (.074) \\ \hline \textbf{29.94} \\ (5.884) \\ \hline 11.92 \\ (.826) \\ \end{array}$	$\begin{array}{r} 24 \text{d.AR} \\ 7.52 \\ (.068) \\ 10.70 \\ (.308) \\ 9.35 \\ (.211) \\ \hline 10.58 \\ (.234) \\ 7.44 \\ (.077) \\ \hline 29.47 \\ (5.934) \\ \hline 10.59 \\ (.285) \end{array}$	$\begin{array}{r} 24 \text{d.AR-wd} \\ \hline 7.12 \\ (.066) \\ \hline 10.24 \\ (.323) \\ \hline 8.72 \\ (.211) \\ \hline 10.98 \\ (.243) \\ \hline 7.14 \\ (.073) \\ \hline 29.37 \\ (5.956) \\ \hline 10.26 \\ (.297) \\ \end{array}$	exp.AR 8.70 (.081) 11.63 (.284) 10.58 (.212) 11.21 (.232) 8.00 (.087) 46.27 (7.033) 11.07 (.281)	$\begin{array}{c} \text{exp.AR-wd} \\ \hline 7.58 \\ (.068) \\ \hline 10.71 \\ (.301) \\ 9.16 \\ (.201) \\ \hline 11.30 \\ (.231) \\ \hline 7.50 \\ (.076) \\ \hline 50.67 \\ (8.386) \\ \hline 10.83 \\ (.289) \\ \end{array}$	$\begin{array}{c} AR(p) \\ \hline 7.01 \\ (.067) \\ \hline 10.02 \\ (.265) \\ \hline 8.77 \\ (.208) \\ \hline 10.79 \\ (.229) \\ \hline 7.26 \\ (.078) \\ \hline 30.69 \\ (5.662) \\ \hline 10.34 \\ (.311) \end{array}$	AR(p)-wd <u>6.73</u> (.066) <u>9.70</u> (.272) 8.48 (.210) 11.02 (.222) <u>7.09</u> (.076) <u>30.50</u> (5.697) 10.06 (.311)	$\begin{array}{c} {\rm PCA}^{*} \\ \hline 7.33 \\ (.072) \\ \hline 10.45 \\ (.296) \\ \hline 9.93 \\ (.260) \\ \hline \underline{10.58} \\ (.210) \\ \hline 7.40 \\ (.081) \\ \hline \underline{29.61} \\ (6.004) \\ \hline 10.77 \\ (.288) \end{array}$	$\begin{array}{r} {\rm PCA^{*}\text{-wd}} \\ \hline 6.79 \\ (.070) \\ \hline 9.82 \\ (.324) \\ \hline 9.18 \\ (.267) \\ \hline 10.90 \\ (.215) \\ \hline 7.09 \\ (.079) \\ \hline 29.35 \\ (6.016) \\ \hline 10.52 \\ (.297) \\ \hline \end{array}$
RMSE EXAA.DE&AT EPEX.DE&AT EPEX.CH BELPEX.BE APX.NL NP.DK.West NP.DK.East	$\begin{array}{c} \text{lasso} \\ 7.02 \\ (.070) \\ 10.06 \\ (.275) \\ 9.03 \\ (.220) \\ \textbf{10.33} \\ (.220) \\ \textbf{10.33} \\ (.239) \\ 7.26 \\ (.079) \\ 30.21 \\ (5.845) \\ 12.38 \\ (.917) \\ \underline{6.76} \end{array}$	lasso-wd 6.63 (.067) 9.47 (.282) <u>8.60</u> (.227) <u>10.37</u> (.239) 7.02 (.074) <u>29.94</u> (5.884) 11.92 (.826) 6.64	$\begin{array}{r} 24 \text{d.AR} \\ 7.52 \\ (.068) \\ 10.70 \\ (.308) \\ 9.35 \\ (.211) \\ \hline 10.58 \\ (.234) \\ \hline 7.44 \\ (.077) \\ \hline 29.47 \\ (5.934) \\ \hline 10.59 \\ (.285) \\ \hline 6.97 \\ \end{array}$	$\begin{array}{r} 24 \text{d.AR-wd} \\ \hline 7.12 \\ (.066) \\ \hline 10.24 \\ (.323) \\ \hline 8.72 \\ (.211) \\ \hline 10.98 \\ (.243) \\ \hline 7.14 \\ (.073) \\ \hline 29.37 \\ (5.956) \\ \hline 10.26 \\ (.297) \\ \hline 6.72 \\ \end{array}$	exp.AR 8.70 (.081) 11.63 (.284) 10.58 (.212) 11.21 (.232) 8.00 (.087) 46.27 (7.033) 11.07 (.281) <u>7.06</u>	$\begin{array}{r} \text{exp.AR-wd} \\ \hline 7.58 \\ (.068) \\ \hline 10.71 \\ (.301) \\ 9.16 \\ (.201) \\ \hline 11.30 \\ (.231) \\ \hline 7.50 \\ (.076) \\ \hline 50.67 \\ (8.386) \\ \hline 10.83 \\ (.289) \\ \hline 6.82 \end{array}$	$\begin{array}{r} AR(p) \\ \hline 7.01 \\ (.067) \\ \hline 10.02 \\ (.265) \\ \hline 8.77 \\ (.208) \\ \hline 10.79 \\ (.229) \\ \hline 7.26 \\ (.078) \\ \hline 30.69 \\ (5.662) \\ \hline 10.34 \\ (.311) \\ \hline 6.90 \end{array}$	AR(p)-wd 6.73 (.066) 9.70 (.272) 8.48 (.210) 11.02 (.222) 7.09 (.076) 30.50 (5.697) 10.06 (.311) 6.78	$\begin{array}{c} {\rm PCA}^{*} \\ \hline 7.33 \\ (.072) \\ \hline 10.45 \\ (.296) \\ \hline 9.93 \\ (.260) \\ \hline 10.58 \\ (.210) \\ \hline 7.40 \\ (.081) \\ \hline 29.61 \\ (6.004) \\ \hline 10.77 \\ (.288) \\ \hline 7.08 \end{array}$	$\begin{array}{r} {\rm PCA^{*}\text{-wd}} \\ \hline 6.79 \\ (.070) \\ \hline 9.82 \\ (.324) \\ 9.18 \\ (.267) \\ \hline 10.90 \\ (.215) \\ \hline 7.09 \\ (.079) \\ \hline 29.35 \\ (6.016) \\ \hline 10.52 \\ (.297) \\ \hline 7.10 \\ \end{array}$
RMSE EXAA.DE&AT EPEX.DE&AT EPEX.CH BELPEX.BE APX.NL NP.DK.West NP.DK.East NP.SW4	$\begin{array}{c} \text{lasso} \\ 7.02 \\ (.070) \\ 10.06 \\ (.275) \\ 9.03 \\ (.220) \\ \textbf{10.33} \\ (.239) \\ 7.26 \\ (.079) \\ \overline{30.21} \\ (5.845) \\ 12.38 \\ (.917) \\ \overline{6.76} \\ (.223) \end{array}$	$\begin{array}{c} \text{lasso-wd} \\ \hline \textbf{6.63} \\ (.067) \\ \hline \textbf{9.47} \\ (.282) \\ \hline \textbf{8.60} \\ (.227) \\ \hline \textbf{10.37} \\ (.239) \\ \hline \textbf{7.02} \\ (.074) \\ \hline \textbf{29.94} \\ (5.884) \\ \hline \textbf{11.92} \\ (.826) \\ \hline \textbf{6.64} \\ (.226) \\ \end{array}$	$\begin{array}{c} 24 \text{d.AR} \\ 7.52 \\ (.068) \\ 10.70 \\ (.308) \\ 9.35 \\ (.211) \\ \hline 10.58 \\ (.234) \\ \hline 7.44 \\ (.077) \\ \hline 29.47 \\ (5.934) \\ \hline 10.59 \\ (.285) \\ \hline 6.97 \\ (.225) \\ \end{array}$	$\begin{array}{r} 24 \text{d.AR-wd} \\ \hline 7.12 \\ (.066) \\ \hline 10.24 \\ (.323) \\ \hline 8.72 \\ (.211) \\ \hline 10.98 \\ (.243) \\ \hline 7.14 \\ (.073) \\ \hline 29.37 \\ (5.956) \\ \hline 10.26 \\ (.297) \\ \hline 6.72 \\ (.229) \\ \end{array}$	exp.AR 8.70 (.081) 11.63 (.284) 10.58 (.212) 11.21 (.232) 8.00 (.087) 46.27 (7.033) 11.07 (.281) <u>7.06</u> (.217)	$\begin{array}{r} \text{exp.AR-wd} \\ \hline 7.58 \\ (.068) \\ \hline 10.71 \\ (.301) \\ 9.16 \\ (.201) \\ \hline 11.30 \\ (.231) \\ \hline 7.50 \\ (.076) \\ \hline 50.67 \\ (8.386) \\ \hline 10.83 \\ (.289) \\ \hline 6.82 \\ (.224) \\ \end{array}$	$\begin{array}{r} AR(p) \\ \hline 7.01 \\ (.067) \\ \hline 10.02 \\ (.265) \\ \hline 8.77 \\ (.208) \\ \hline 10.79 \\ (.229) \\ \hline 7.26 \\ (.078) \\ \hline 30.69 \\ (5.662) \\ \hline 10.34 \\ (.311) \\ \hline 6.90 \\ (.241) \\ \end{array}$	$\begin{array}{r} AR(p)\text{-wd} \\ \hline 6.73 \\ (.066) \\ \hline 9.70 \\ (.272) \\ \hline 8.48 \\ (.210) \\ \hline 11.02 \\ (.222) \\ \hline 7.09 \\ (.076) \\ \hline 30.50 \\ (5.697) \\ \hline 10.06 \\ (.311) \\ \hline 6.78 \\ (.243) \\ \hline \end{array}$	$\begin{array}{c} {\rm PCA}^{*} \\ \hline 7.33 \\ (.072) \\ \hline 10.45 \\ (.296) \\ \hline 9.93 \\ (.260) \\ \hline 10.58 \\ (.210) \\ \hline 7.40 \\ (.081) \\ \hline 29.61 \\ (6.004) \\ \hline 10.77 \\ (.288) \\ \hline 7.08 \\ (.220) \\ \end{array}$	$\begin{array}{r} {\rm PCA*-wd}\\ \hline 6.79\\ (.070)\\ \hline 9.82\\ (.324)\\ 9.18\\ (.267)\\ \hline 10.90\\ (.215)\\ \hline 7.09\\ (.079)\\ \hline 29.35\\ (6.016)\\ \hline 10.52\\ (.297)\\ \hline 7.10\\ (.221)\\ \end{array}$
RMSE EXAA.DE&AT EPEX.DE&AT EPEX.CH BELPEX.BE APX.NL NP.DK.West NP.DK.East NP.SW4 POLPX PL	$\begin{array}{c} \text{lasso} \\ 7.02 \\ (.070) \\ 10.06 \\ (.275) \\ 9.03 \\ (.220) \\ \textbf{10.33} \\ (.220) \\ \textbf{10.33} \\ (.239) \\ 7.26 \\ (.079) \\ \overline{30.21} \\ (5.845) \\ 12.38 \\ (.917) \\ \overline{6.76} \\ (.223) \\ \overline{5.37} \end{array}$	$\begin{array}{c} \text{lasso-wd} \\ \hline \textbf{6.63} \\ (.067) \\ \hline \textbf{9.47} \\ (.282) \\ \hline \textbf{8.60} \\ (.227) \\ \hline \textbf{10.37} \\ (.239) \\ \hline \textbf{7.02} \\ (.074) \\ \hline \textbf{29.94} \\ (5.884) \\ \hline \textbf{11.92} \\ (.826) \\ \hline \textbf{6.64} \\ (.226) \\ \hline \textbf{5.17} \end{array}$	$\begin{array}{r} 24 \text{d.AR} \\ 7.52 \\ (.068) \\ 10.70 \\ (.308) \\ 9.35 \\ (.211) \\ \hline 10.58 \\ (.234) \\ \hline 7.44 \\ (.077) \\ \hline 29.47 \\ (5.934) \\ \hline 10.59 \\ (.285) \\ \hline 6.97 \\ (.225) \\ \hline 5.06 \\ \end{array}$	$\begin{array}{r} 24 \text{d.AR-wd} \\ \hline 7.12 \\ (.066) \\ \hline 10.24 \\ (.323) \\ \hline 8.72 \\ (.211) \\ \hline 10.98 \\ (.243) \\ \hline 7.14 \\ (.073) \\ \hline 29.37 \\ (5.956) \\ \hline 10.26 \\ (.297) \\ \hline 6.72 \\ (.229) \\ \hline 4.94 \end{array}$	exp.AR 8.70 (.081) 11.63 (.284) 10.58 (.212) 11.21 (.232) 8.00 (.087) 46.27 (7.033) 11.07 (.281) 7.06 (.217) 5.61	$\begin{array}{r} \text{exp.AR-wd} \\ \hline 7.58 \\ (.068) \\ \hline 10.71 \\ (.301) \\ 9.16 \\ (.201) \\ \hline 11.30 \\ (.231) \\ \hline 7.50 \\ (.076) \\ \hline 50.67 \\ (8.386) \\ \hline 10.83 \\ (.289) \\ \hline 6.82 \\ (.224) \\ \hline 5.08 \end{array}$	$\begin{array}{r} AR(p) \\ \hline 7.01 \\ (.067) \\ \hline 10.02 \\ (.265) \\ \hline 8.77 \\ (.208) \\ \hline 10.79 \\ (.229) \\ \hline 7.26 \\ (.078) \\ \hline 30.69 \\ (5.662) \\ \hline 10.34 \\ (.311) \\ \hline 6.90 \\ (.241) \\ \hline 4.98 \end{array}$	AR(p)-wd <u>6.73</u> (.066) <u>9.70</u> (.272) 8.48 (.210) 11.02 (.222) <u>7.09</u> (.076) <u>30.50</u> (5.697) 10.06 (.311) <u>6.78</u> (.243) 4.92	$\begin{array}{c} {\rm PCA}^{*} \\ \hline 7.33 \\ (.072) \\ \hline 10.45 \\ (.296) \\ \hline 9.93 \\ (.260) \\ \hline 10.58 \\ (.210) \\ \hline 7.40 \\ (.081) \\ \hline 29.61 \\ (6.004) \\ \hline 10.77 \\ (.288) \\ \hline 7.08 \\ (.220) \\ \hline 5.52 \end{array}$	$\begin{array}{r} {\rm PCA}^{*}{\rm -wd} \\ \hline 6.79 \\ (.070) \\ \hline 9.82 \\ (.324) \\ 9.18 \\ (.267) \\ \hline 10.90 \\ (.215) \\ \hline 7.09 \\ (.079) \\ \hline 29.35 \\ (6.016) \\ \hline 10.52 \\ (.297) \\ \hline 7.10 \\ (.221) \\ \hline 5.19 \\ \end{array}$
RMSE EXAA.DE&AT EPEX.DE&AT EPEX.CH BELPEX.BE APX.NL NP.DK.West NP.DK.East NP.SW4 POLPX.PL	$\begin{array}{c} \text{lasso} \\ \hline 7.02 \\ (.070) \\ \hline 10.06 \\ (.275) \\ \hline 9.03 \\ (.220) \\ \hline 10.33 \\ (.220) \\ \hline 10.33 \\ (.239) \\ \hline 7.26 \\ (.079) \\ \hline 30.21 \\ (5.845) \\ \hline 12.38 \\ (.917) \\ \hline 6.76 \\ (.223) \\ \hline 5.37 \\ (.156) \end{array}$	$\begin{array}{c} \text{lasso-wd} \\ \hline \textbf{6.63} \\ (.067) \\ \hline \textbf{9.47} \\ (.282) \\ \hline \textbf{8.60} \\ (.227) \\ \hline \textbf{10.37} \\ (.239) \\ \hline \textbf{7.02} \\ (.074) \\ \hline \textbf{29.94} \\ (5.884) \\ \hline \textbf{11.92} \\ (.826) \\ \hline \textbf{6.64} \\ (.226) \\ \hline \textbf{5.17} \\ (.145) \\ \end{array}$	$\begin{array}{r} 24 \text{d.AR} \\ 7.52 \\ (.068) \\ 10.70 \\ (.308) \\ 9.35 \\ (.211) \\ \hline 10.58 \\ (.234) \\ \hline 7.44 \\ (.077) \\ \hline 29.47 \\ (5.934) \\ \hline 10.59 \\ (.285) \\ \hline 6.97 \\ (.225) \\ \hline 5.06 \\ (.127) \\ \end{array}$	$\begin{array}{r} 24 \text{d.AR-wd} \\ \hline 7.12 \\ (.066) \\ \hline 10.24 \\ (.323) \\ \hline 8.72 \\ (.211) \\ \hline 10.98 \\ (.243) \\ \hline 7.14 \\ (.073) \\ \hline 29.37 \\ (5.956) \\ \hline 10.26 \\ (.297) \\ \hline 6.72 \\ (.229) \\ \hline 4.94 \\ (.130) \\ \end{array}$	exp.AR 8.70 (.081) 11.63 (.284) 10.58 (.212) 11.21 (.232) 8.00 (.087) 46.27 (7.033) 11.07 (.281) 7.06 (.217) 5.61 (.133)	$\begin{array}{r} \text{exp.AR-wd} \\ \hline 7.58 \\ (.068) \\ \hline 10.71 \\ (.301) \\ 9.16 \\ (.201) \\ \hline 11.30 \\ (.231) \\ \hline 7.50 \\ (.076) \\ \hline 50.67 \\ (8.386) \\ \hline 10.83 \\ (.289) \\ \hline 6.82 \\ (.224) \\ \hline 5.08 \\ (.133) \\ \end{array}$	$\begin{array}{r} AR(p) \\ \hline 7.01 \\ (.067) \\ \hline 10.02 \\ (.265) \\ \hline 8.77 \\ (.208) \\ \hline 10.79 \\ (.229) \\ \hline 7.26 \\ (.078) \\ \hline 30.69 \\ (5.662) \\ \hline 10.34 \\ (.311) \\ \hline 6.90 \\ (.241) \\ \hline 4.98 \\ (.114) \\ \end{array}$	AR(p)-wd <u>6.73</u> (.066) <u>9.70</u> (.272) 8.48 (.210) 11.02 (.222) <u>7.09</u> (.076) <u>30.50</u> (5.697) 10.06 (.311) <u>6.78</u> (.243) 4.92 (.115)	$\begin{array}{c} {\rm PCA}^{*} \\ \hline 7.33 \\ (.072) \\ \hline 10.45 \\ (.296) \\ \hline 9.93 \\ (.260) \\ \hline 10.58 \\ (.210) \\ \hline 7.40 \\ (.081) \\ \hline 29.61 \\ (6.004) \\ \hline 10.77 \\ (.288) \\ \hline 7.08 \\ (.220) \\ \hline 5.52 \\ (.139) \end{array}$	$\begin{array}{r} {\rm PCA*-wd} \\ \hline 6.79 \\ (.070) \\ \hline 9.82 \\ (.324) \\ \hline 9.18 \\ (.267) \\ \hline 10.90 \\ (.215) \\ \hline 7.09 \\ (.079) \\ \hline 29.35 \\ (6.016) \\ \hline 10.52 \\ (.297) \\ \hline 7.10 \\ (.221) \\ \hline 5.19 \\ (.138) \\ \end{array}$
RMSE EXAA.DE&AT EPEX.DE&AT EPEX.CH BELPEX.BE APX.NL NP.DK.West NP.DK.East NP.SW4 POLPX.PL	$\begin{array}{c} \text{lasso} \\ 7.02 \\ (.070) \\ 10.06 \\ (.275) \\ 9.03 \\ (.220) \\ \textbf{10.33} \\ (.220) \\ \textbf{10.33} \\ (.239) \\ 7.26 \\ (.079) \\ \hline \textbf{30.21} \\ (5.845) \\ 12.38 \\ (.917) \\ \hline \textbf{6.76} \\ (.223) \\ \hline \textbf{5.37} \\ (.156) \\ \textbf{8.08} \end{array}$	lasso-wd 6.63 (.067) 9.47 (.282) <u>8.60</u> (.227) <u>10.37</u> (.239) 7.02 (.074) <u>29.94</u> (5.884) 11.92 (.826) 6.64 (.226) <u>5.17</u> (.145) 7.57	$\begin{array}{r} 24 \text{d.AR} \\ \hline 7.52 \\ (.068) \\ \hline 10.70 \\ (.308) \\ \hline 9.35 \\ (.211) \\ \hline 10.58 \\ (.234) \\ \hline 7.44 \\ (.077) \\ \hline 29.47 \\ (5.934) \\ \hline 10.59 \\ (.285) \\ \hline 6.97 \\ (.225) \\ \hline 5.06 \\ (.127) \\ \hline 8.46 \end{array}$	$\begin{array}{r} 24 \text{d.AR-wd} \\ \hline 7.12 \\ (.066) \\ \hline 10.24 \\ (.323) \\ \hline 8.72 \\ (.211) \\ \hline 10.98 \\ (.243) \\ \hline 7.14 \\ (.073) \\ \hline 29.37 \\ (5.956) \\ \hline 10.26 \\ (.297) \\ \hline 6.72 \\ (.229) \\ \hline 4.94 \\ (.130) \\ \hline 7.98 \\ \end{array}$	exp.AR 8.70 (.081) 11.63 (.284) 10.58 (.212) 11.21 (.232) 8.00 (.087) 46.27 (7.033) 11.07 (.281) 7.06 (.217) 5.61 (.133) 9.66	$\begin{array}{r} \text{exp.AR-wd} \\ \hline 7.58 \\ (.068) \\ \hline 10.71 \\ (.301) \\ 9.16 \\ (.201) \\ \hline 11.30 \\ (.231) \\ \hline 7.50 \\ (.076) \\ \hline 50.67 \\ (8.386) \\ \hline 10.83 \\ (.289) \\ \hline 6.82 \\ (.224) \\ \hline 5.08 \\ (.133) \\ \hline 8.54 \end{array}$	$\begin{array}{r} {\rm AR(p)}\\ {\rm 7.01}\\ (.067)\\ {\rm 10.02}\\ (.265)\\ {\rm 8.77}\\ (.208)\\ {\rm 10.79}\\ (.229)\\ {\rm 7.26}\\ (.078)\\ {\rm 30.69}\\ (5.662)\\ {\rm 10.34}\\ (.311)\\ {\rm 6.90}\\ (.241)\\ {\rm 4.98}\\ (.114)\\ {\rm 7.90}\\ \end{array}$	AR(p)-wd <u>6.73</u> (.066) <u>9.70</u> (.272) 8.48 (.210) 11.02 (.222) <u>7.09</u> (.076) <u>30.50</u> (5.697) 10.06 (.311) <u>6.78</u> (.243) 4.92 (.115) <u>7.58</u>	$\begin{array}{c} {\rm PCA}^{*} \\ \hline 7.33 \\ (.072) \\ \hline 10.45 \\ (.296) \\ \hline 9.93 \\ (.260) \\ \hline 10.58 \\ (.210) \\ \hline 7.40 \\ (.081) \\ \hline 29.61 \\ (6.004) \\ \hline 10.77 \\ (.288) \\ \hline 7.08 \\ (.220) \\ \hline 5.52 \\ (.139) \\ \hline 8.25 \end{array}$	$\begin{array}{r} {\rm PCA*-wd} \\ \hline 6.79 \\ (.070) \\ \hline 9.82 \\ (.324) \\ \hline 9.18 \\ (.267) \\ \hline 10.90 \\ (.215) \\ \hline 7.09 \\ (.079) \\ \hline 29.35 \\ (6.016) \\ \hline 10.52 \\ (.297) \\ \hline 7.10 \\ (.221) \\ \hline 5.19 \\ (.138) \\ \hline 7.81 \\ \end{array}$

Table 2: MAE and RMSE values for the considered markets and models. The corresponding estimated standard deviations are given in parenthesis. The best model is highlighted in bold font. All models that are not significantly worse than best (indicated by the 2-sigma range of the best model) are underlined.

In Figure 3 we observe that the daily pattern for the MAE_h and $RMSE_h$ are similar. In general we see that the model with weekday effects estimated by lasso performs as one of the best models for all hours over the day. It seems that during the morning hours it outperforms the other models most clearly. It is interesting that in the night hours from 0:00 to 5:00 the lasso based models perform similarly to the univariate AR(p). But this matches well the observation from the beginning that these hours are highly correlated with the last hours of the previous day. The univariate AR(p) and the lasso models can capture this behavior. Moreover, in the evening hours (esp. 21:00 to 23:00) the 24-dimensional AR model and the 24-dimensional autoregressive expert model perform as good as the lasso based models. Again, this coincides with the observation from Figure 1, that these hours are strongly correlated with the same hours of the previous days. The mentioned models are all able to cover this dependency well. Furthermore, we see a relatively well performance of the PCA*-wd model. We know from Table 2 that for the German/Austrian EPEX price only the PCA*-wd model performs not significantly worse. The behavior is similar as both core models (ignoring the weekday effects) are nested in (13). However, regarding the cross-country Table 2, we know that in most cases the lasso-wd model outperforms the PCA*-wd model. The reason is likely that cross-hour dependency is modeled directly instead of using the reduced PCA representation. It might happen that relevant information gets lost due to the dimension reduction technique.

Another interesting fact in Figure 3 is that it seems that in the morning hours around 6:00 to 7:00 the lasso-wd performs much better in the MAE_h score than in the $RMSE_h$. However, having a closer look at the graph it turns out the AR(p) process catches up in terms of $RMSE_h$. The reason might be again that the AR(p) based models are less

Furthermore, we can see in Figure 3 that it seems that the weekday dummies in the model (2) improve the performance significantly compared to model (1). We will understand this better when we analyze the intraday structure and the variable importance.

Hence, additionally to the out-of-sample study we have a closer look at the intraday structure of the electricity prices by using the proposed model (2). As the lasso estimator sparsifies and regularizes, we can study the finally selected parameters and their magnitude. For a measure of importance of a parameter $\beta_{h,i}$ resp. $\tilde{\beta}_{h,i}$ we consider $\iota_{h,i}$, the fraction of the absolute standardized parameter of a model to the sum of all absolute standardized parameters:

$$\iota_{h,i} = \frac{|\widetilde{\beta}_{h,i}|}{\sum_{j=1}^{m_h} |\widetilde{\beta}_{h,j}|} \tag{21}$$

Obviously, we have that $0 \leq \iota_{h,i} \leq 1$ and $\sum_{i=1}^{m_h} \iota_{h,i} = 1$. The larger $\iota_{h,i}$ the larger the relative impact to the electricity price of hour h. We estimate $\iota_{h,i}$ for model (2) which includes the weekday dummies for all considered markets by the corresponding plug-in estimator $\hat{\iota}_{h,i}$ using the estimated standardized parameter vector $\hat{\beta}_h$. Given the estimated $\hat{\iota}_{h,i}$ values, we can sort them decreasingly. Then the largest $\hat{\iota}_{h,i}$ value matches the most relevant parameter in the model for hour h.

Again, we only discuss the results for the German/Austrian EPEX spot price in detail. In Table 3 we present the 7 most important variables for each model of hour h and their $\hat{\iota}_{h,i}$ value. The corresponding tables for the other electricity prices are given in the supplementary material. In Table 3 we can observe an interesting pattern that we

	Importance: 1	Importance: 2	Importance: 3	Importance: 4	Importance: 5	Importance: 6	Importance: 7
0	23@1(70.6)	19@1(5.9)	1@1(5.7)	18@1 (4.1)	21@1(3.6)	4@1 (3.1)	Sat (2.2)
1	23@1 (67.3)	19@1(8.6)	3@1(7.1)	4@1(5.9)	Sun (4.7)	5@1(2.6)	6@1(2.0)
2	23@1 (56.3)	4@1(9.4)	19@1(8.9)	3@1(7.9)	Sun (6.6)	22@7(5.1)	2@25(1.6)
3	23@1 (45.0)	4@1(14.3)	3@1(12.2)	19@1(8.5)	Sun (7.3)	22@7(4.3)	3@6(2.5)
4	23@1 (44.8)	4@1(13.8)	19@1 (10.3)	Sun (10.2)	3@1(7.4)	22@7(2.5)	4@25(2.4)
5	23@1(34.1)	Sun (15.5)	5@1(9.7)	4@1(8.2)	Sat (7.4)	19@1 (7.2)	22@7(3.8)
6	Sun (20.4)	23@1 (20.3)	Sat (12.8)	6@1(9.4)	Mon (8.7)	20@1 (6.0)	23@2(4.1)
7	Sun (23.9)	Sat (15.9)	23@1(12.1)	Mon (7.8)	20@1 (6.3)	6@1 (5.3)	19@1 (4.5)
8	Sun~(28.3)	Sat (15.9)	20@1 (8.5)	23@1(7.9)	18@1(7.0)	Mon (5.4)	21@1(5.2)
9	Sun (25.9)	20@1(13.3)	Sat (13.3)	23@1(7.0)	22@1(6.3)	Mon (6.1)	18@1 (5.5)
10	Sun (21.8)	Sat (11.9)	22@1 (9.6)	20@1 (9.0)	Mon (6.6)	13@1(6.1)	23@1(5.6)
11	Sun (17.2)	22@1(12.3)	Sat (11.7)	13@1 (9.2)	Mon (8.0)	23@1 (5.8)	11@7(4.1)
12	Sun (15.6)	13@1 (12.0)	Sat (10.7)	22@1(10.2)	Mon (8.1)	23@1(8.0)	17@1(4.9)
13	Sun (18.6)	Sat (11.2)	$13@1\ (11.1)$	22@1(10.0)	Mon (7.9)	23@1(7.3)	17@1 (6.8)
14	Sun (18.7)	Sat (10.1)	Mon (8.2)	23@1(8.2)	16@1(8.1)	22@1(7.6)	13@1(4.2)
15	Sun (19.8)	16@1 (12.5)	Sat (8.9)	Mon (8.0)	23@1(7.1)	17@6(6.0)	17@1(4.9)
16	Sun (19.3)	17@1(12.5)	Sat (8.8)	16@1 (8.5)	Mon (8.4)	23@1(7.1)	17@6(6.7)
17	17@1(26.4)	Sun (16.1)	Mon (10.1)	17@6(10.0)	Sat (6.7)	17@7(6.3)	18@1 (6.0)
18	18@1 (33.8)	Sun (14.1)	Mon (9.9)	18@6(9.4)	Sat (6.0)	18@2(5.9)	18@5(4.6)
19	19@1 (34.0)	Mon (9.4)	19@6(8.5)	Sun(8.2)	19@2(7.6)	8@6(5.0)	19@7(4.7)
20	20@1 (23.9)	20@2(10.4)	Sat (9.5)	Mon (7.8)	21@1(7.5)	20@6(7.5)	Sun (7.0)
21	21@1(22.8)	Sat (10.9)	21@2(9.9)	Mon (6.2)	21@28(4.7)	Sun (4.6)	23@1(4.6)
22	22@1(30.6)	22@2(9.5)	Sat (7.7)	23@3(6.5)	22@28(6.1)	22@5(5.1)	22@35(5.0)
23	23@1(24.8)	22@1(17.5)	23@3(10.7)	22@2(9.6)	Sat (5.7)	0@5~(~5.0)	12@7(4.4)

Table 3: Most relevant coefficients in model (2) for each hour of the German/Austrian EPEX price. H@L represents the estimated parameter for $Y_{d-L,H}$ for a model on $Y_{d,h}$. Sun, Mon, ..., Sat represent the weekday dummies. The corresponding $\hat{\iota}_{h,i}$ value is given in parenthesis.

have similarly seen already in the introduction. The most relevant parameters for the first 6 hours of the day is the last observed value of the previous day (which corresponds to $P_{d-1,23}$). For the first hour this importance is very distinct. The last observation $P_{d-1,23}$ of the previous day contributes 70.6% to the full model. From the physical point of view this makes sense. Obviously, the physical delivery of the auction result for a specific hour is based on related processes like the electricity load or wind energy production. These situations should not differ that much between 23:00 and the next day 0:00. So a large impact of the 23:00 price to the price of the next day at hour 0:00 is plausible. This observation seems to be very stable across all countries. So we observe for all considered markets that the most relevant variable for the first 0:00 hour is always the last hour of the previous day. For all markets, except for the BELPEX.BE, this is even the case for the first 3 hours (0:00, 1:00 and 2:00) of the day. A similar result is recently established in load forecasting by Dudek (2016). However, for electricity price forecasting this is not that obvious as we observe all 24 values at once. The last observation of the previous day is not more recent than the other ones.

11

The fact that for the first hours the last values of the previous day are very important underlines that the univariate AR(p) and the lasso based models perform well in this time horizon.

For the last 7 hours of the day (from 17:00 to 23:00) the most relevant lag is always the value of the previous day at the same hour (which corresponds to $P_{d-1,h}$). This matches well the observed fact that the 24-dimensional autoregressive models have a good out-of-sample performance for these hours.

Moreover, we see that for the morning and afternoon hours from 6:00 to 16:00 the most important parameter is the weekday dummy on Sunday. So the weekday effect seems to be more important than the autoregressive impact. In general we observe that the weekday dummies have a larger impact during the working hours than in the night and evening hours. Thus, we can deduce that the weekly seasonal structure of the electricity price is more distinct in the working hours. This coincides with Figure 2a, where we see that the difference in the weekly mean is smaller during night and evening than the rest of the day.

Additionally, we observe that only the weekday dummies for Sunday, Monday and Saturday occur in Table 3. Here it is interesting that the considered expert model from Weron and Misiorek (2008) uses exactly the same dummies. However, regarding the weekly sample means in Figure 2a this is not obvious. From this graph we would expect that the impact of Monday dummies is similar to Friday dummies. Furthermore the Monday dummies appear only for the hours from 6:00 to 21:00 which is definitively not expectable from Figure 2a which suggests a Monday dummy only from 0:00 to 5:00.

Another fact that we can derive from Table 3 concerns the weekly lagged impact. In many models, e.g. the expert model from Weron and Misiorek (2008) a linear dependence on the weekly lagged electricity price is considered $P_{d-7,h}$. However, in Table 3 it never appears in the top 4 for any hour. Only for some hours it is estimated to be the 5th, 6th or 7th most important variable. This is an indication that given a proper short-term autoregressive and weekday modeling the modeled electricity price dependence from the weekly lagged price gets weaker. The last fact from Table 3 that we want to mention concerns the evening hours. We observe that only for the evening hours from 18:00 to 23:00 a dependence of electricity price of the day before yesterday (lag = 2) appears. Again, this dependency is always to the price at the same hour.

5 Summary and conclusion

We present a day-ahead electricity spot price model approach for hourly data that is able to capture the intraday dependency structure well, especially the time-varying cross-hour dependencies. Even though the considered regression model consists of 24 simple linear regression models it explains the variety in the price data. This relies heavily on the utilized lasso estimation technique. It automatically shrinks and sparsifies the model parameters. Thus, the estimated 24 single models can vary over the day and capture the dependency in the data.

We show that the dependency structure significantly changes over the days. For example in the German/Austrian EPEX market, the night hours from 0:00 to 5:00 have a strong dependency to the last price at 23:00 of the previous day. This result holds similarly across all considered European markets. In contrast, the price at the evening hours of from 18:00 to 23:00 of the German/Austrian EPEX market depends more on the price of the previous day at the same hour. We highlight that the majority of available electricity price model approaches are not designed to cover both features, whereas the lasso estimation technique captures both features in an efficient way.

The conducted out-of-sample forecasting study shows that the considered lasso based approach performs strong in terms of forecasting accuracy. This is remarkable, if we have in mind that the estimation and forecasting procedure is very fast. Hence, the model is a suitable candidate for forecast combination approaches as in Bordignon et al. (2013) or Maciejowska et al. (2015). As the out-of-sample study shows similar performance results for all 10 considered electricity markets, it underlines that the flexible model approach can cover the local market specific dependency behavior.

Future research can go in different directions. The likely most important issue is the covering of the non-linear behavior, especially those effects that induce price spikes. This should be combined with an approach that takes other sources like renewable energy feed-in or load into account. Especially the impact of wind power is very important to cover price spikes, see e.g. Keles et al. (2012). Furthermore, there should be more research done for probabilistic forecasting.

References

- Abedinia, O., Amjady, N., Shafie-khah, M., and Catalão, J. (2015). Electricity price forecast using combinatorial neural network trained by a new stochastic search method. *Energy Conversion and Management*, 105:642–654.
- Bordignon, S., Bunn, D. W., Lisi, F., and Nan, F. (2013). Combining day-ahead forecasts for British electricity prices. *Energy Economics*, 35:88–103.
- Catalao, J., Pousinho, H., and Mendes, V. (2011). Hybrid wavelet-pso-anfis approach for short-term electricity prices forecasting. *IEEE Transactions on Power Systems*, 26(1):137–144.

- Chaâbane, N. (2014). A hybrid arfima and neural network model for electricity price prediction. International Journal of Electrical Power & Energy Systems, 55:187–194.
- Conejo, A. J., Plazas, M., Espinola, R., Molina, A. B., et al. (2005). Day-ahead electricity price forecasting using the wavelet transform and ARIMA models. *Power Systems, IEEE Transactions on*, 20(2):1035–1042.
- Cruz, A., Munoz, A., Zamora, J. L., and Espínola, R. (2011). The effect of wind generation and weekday on spanish electricity spot price forecasting. *Electric Power Systems Research*, 81(10):1924–1935.
- Dudek, G. (2016). Pattern-based local linear regression models for short-term load forecasting. *Electric Power Systems Research*, 130:139–147.
- Friedman, J., Hastie, T., Höfling, H., and Tibshirani, R. (2007). Pathwise coordinate optimization. The Annals of Applied Statistics, 1(2):302–332.
- Friedman, J., Hastie, T., and Tibshirani, R. (2010). Regularization paths for generalized linear models via coordinate descent. Journal of statistical software, 33(1):1.
- Haldrup, N., Nielsen, F. S., and Nielsen, M. O. (2010). A vector autoregressive model for electricity prices subject to long memory and regime switching. *Energy Economics*, 32(5):1044–1058.
- Hastie, T., Tibshirani, R., and Wainwright, M. (2015). Statistical Learning with Sparsity: The Lasso and Generalizations. CRC Press.
- Hu, L. and Taylor, G. (2014). A novel hybrid technique for short-term electricity price forecasting in uk electricity markets. *Journal of International Council on Electrical Engineering*, 4(2):114–120.
- Karakatsani, N. V. and Bunn, D. W. (2008a). Forecasting electricity prices: The impact of fundamentals and timevarying coefficients. *International Journal of Forecasting*, 24(4):764–785.
- Karakatsani, N. V. and Bunn, D. W. (2008b). Intra-day and regime-switching dynamics in electricity price formation. Energy Economics, 30(4):1776–1797.
- Keles, D., Genoese, M., Möst, D., and Fichtner, W. (2012). Comparison of extended mean-reversion and time series models for electricity spot price simulation considering negative prices. *Energy Economics*, 34(4):1012–1032.
- Ludwig, N., Feuerriegel, S., and Neumann, D. (2015). Putting big data analytics to work: Feature selection for forecasting electricity prices using the lasso and random forests. *Journal of Decision Systems*, 24(1):19–36.
- Maciejowska, K., Nowotarski, J., and Weron, R. (2015). Probabilistic forecasting of electricity spot prices using Factor Quantile Regression Averaging. *International Journal of Forecasting*.
- Maciejowska, K. and Weron, R. (2013). Forecasting of daily electricity spot prices by incorporating intra-day relationships: Evidence form the UK power market. In European Energy Market (EEM), 2013 10th International Conference on the, pages 1–5. IEEE.
- Maciejowska, K. and Weron, R. (2015). Short-and mid-term forecasting of baseload electricity prices in the UK: The impact of intra-day price relationships and market fundamentals. *Power Systems, IEEE Transactions on*, PP(99):1–12.
- Nowotarski, J. and Weron, R. (2014). Computing electricity spot price prediction intervals using quantile regression and forecast averaging. *Computational Statistics*, pages 1–13.
- Paraschiv, F., Erni, D., and Pietsch, R. (2014). The impact of renewable energies on EEX day-ahead electricity prices. *Energy Policy*, 73:196–210.
- Paraschiv, F., Fleten, S.-E., and Schürle, M. (2015). A spot-forward model for electricity prices with regime shifts. *Energy Economics*, 47:142–153.
- Shafie-Khah, M., Moghaddam, M. P., and Sheikh-El-Eslami, M. (2011). Price forecasting of day-ahead electricity markets using a hybrid forecast method. *Energy Conversion and Management*, 52(5):2165–2169.
- Singhal, D. and Swarup, K. (2011). Electricity price forecasting using artificial neural networks. International Journal of Electrical Power & Energy Systems, 33(3):550–555.
- Tibshirani, R. (1996). Regression shrinkage and selection via the lasso. Journal of the Royal Statistical Society: Series B (Statistical Methodology), 58(1):267–288.
- Weron, R. (2014). Electricity price forecasting: A review of the state-of-the-art with a look into the future. *International Journal of Forecasting*, 30(4):1030–1081.

- Weron, R. and Misiorek, A. (2008). Forecasting spot electricity prices: A comparison of parametric and semiparametric time series models. *International Journal of Forecasting*, 24(4):744–763.
- Ziel, F. (2015). Iteratively reweighted adaptive lasso for conditional heteroscedastic time series with applications to ar-arch type processes. *Computational Statistics and Data Analysis (in press)*.
- Ziel, F., Steinert, R., and Husmann, S. (2015a). Efficient modeling and forecasting of electricity spot prices. Energy Economics, 47:98–111.
- Ziel, F., Steinert, R., and Husmann, S. (2015b). Forecasting day ahead electricity spot prices: The impact of the exaa to other european electricity markets. *Energy Economics*, 51:430–444.

Appendix

6 Figures

6.1 Correlation plots



Figure 4: Sample correlations of $\mathbb{C}or(P_{d,h}, P_{d-1,l})$ for $h, l = 0, \ldots, 23$ and selected markets from 17.12.2009 to 12.08.2014.

Figure 5: Sample correlations of $\mathbb{C}or(P_{d,h}, P_{d-1,l})$ for $h, l = 0, \ldots, 23$ and selected markets from 17.12.2009 to 12.08.2014.

6.2 Weekly sample mean plots

Figure 6: Weekly sample mean of $P_{d,h}$ for selected markets.

Figure 7: Weekly sample mean of $P_{d,h}$ for selected markets.

6.3 MAE_h plots

Figure 8: MAE_h for selected markets

6.4 \mathbf{RMSE}_h plots

Figure 10: RMSE_h for selected markets

7 Tables of parameter importance

EXAA Germany & Austria

	Importance: 1	Importance: 2	Importance: 3	Importance: 4	Importance: 5	Importance: 6	Importance: 7
0	23@1(65.3)	3@1(10.2)	Sun (6.6)	19@1 (4.2)	Sat (3.7)	4@1 (3.2)	18@1 (3.1)
1	23@1 (50.0)	4@1(10.3)	Sun (8.9)	3@1(7.7)	19@1(6.3)	23@2(2.5)	18@1(2.0)
2	23@1(39.1)	3@1(13.2)	Sun(8.1)	23@2(7.3)	4@1(6.9)	19@1(4.9)	22@7(3.1)
3	23@1 (31.9)	3@1 (10.6)	4@1(9.7)	23@2(8.3)	Sun(8.3)	22@7(4.4)	20@1(3.7)
4	23@1 (30.6)	4@1(19.7)	Sun (11.1)	23@2(5.8)	22@7(5.5)	4@6(5.1)	20@1(4.6)
5	23@1 (27.2)	Sun (14.9)	4@1(7.5)	20@1 (6.7)	5@7(6.5)	22@7(6.1)	5@1(4.6)
6	Sun (25.7)	20@1(17.6)	23@1(14.2)	Sat (12.6)	6@7(7.8)	Mon (5.8)	22@7(4.5)
7	Sun (26.0)	20@1(17.8)	Sat (14.6)	Mon (7.7)	23@1(7.0)	18@1 (5.0)	7@7(4.8)
8	Sun (23.6)	20@1 (15.0)	Sat (13.5)	23@1(8.8)	Mon (8.1)	7@1(5.4)	18@1 (5.1)
9	Sun (23.8)	20@1(15.4)	Sat (13.5)	Mon (9.8)	23@1(8.0)	11@1(5.1)	18@1 (5.0)
10	Sun (19.5)	11@1(13.9)	Sat (12.4)	Mon (10.9)	20@1(8.6)	23@1(6.7)	22@1 (6.5)
11	Sun (14.2)	Sat (11.5)	Mon (11.3)	11@1(10.6)	22@1(9.9)	12@1(6.9)	23@1(6.0)
12	12@1(12.1)	Mon (11.9)	Sun (11.4)	Sat (9.3)	23@1(8.9)	13@1(8.8)	22@1(7.6)
13	13@1 (15.0)	Sun (13.9)	Mon (11.9)	23@1(10.1)	Sat (10.0)	17@1(6.2)	14@7(5.9)
14	Sun (15.3)	14@1 (15.0)	Mon~(11.2)	23@1 (10.5)	Sat (9.9)	17@1(7.8)	14@7(6.2)
15	Sun (17.3)	15@1 (11.2)	Mon (10.9)	17@1(10.6)	Sat (9.4)	23@1(8.7)	14@7(5.9)
16	Sun (16.9)	17@1 (15.2)	Mon (8.6)	Sat (8.3)	23@1(7.7)	16@7(6.8)	17@6(6.2)
17	17@1(32.2)	Sun (13.0)	Mon (11.1)	17@6(10.3)	17@7(9.1)	Sat (5.9)	22@7(3.8)
18	18@1 (42.1)	Mon~(14.2)	Sun (9.6)	18@6(9.4)	18@7(8.8)	Sat (4.5)	17@6(2.8)
19	19@1 (35.2)	Mon (11.3)	Sun(8.8)	20@1(8.0)	19@6(7.4)	Sat (6.7)	18@6(4.4)
20	20@1 (42.6)	20@6~(11.5)	Mon (10.0)	Sat (9.1)	Sun (6.5)	23@2(2.7)	20@4(2.3)
21	21@1(38.5)	21@6(12.0)	Mon (9.7)	Sat (7.7)	22@1(5.8)	23@2(4.5)	21@35(3.3)
22	22@1(44.8)	22@6(9.2)	23@2(6.6)	Sat (6.0)	23@1(5.7)	Mon (5.5)	0@5(5.4)
23	23@1 (36.8)	22@1 (9.4)	23@2(7.5)	23@6(7.1)	23@4(5.4)	23@28(5.2)	Mon (4.8)

Table 4: Most relevant coefficients for each hour. H@L represents the estimated parameter for $Y_{d-L,H}$ for a model on $Y_{d,h}$. Sun, Mon, ..., Sat represent the weekday dummies. The corresponding $\hat{\iota}_{h,i}$ value is given in parenthesis.

EPEX Germany & Austria

	Importance: 1	Importance: 2	Importance: 3	Importance: 4	Importance: 5	Importance: 6	Importance: 7
0	23@1(70.6)	19@1(5.9)	1@1(5.7)	18@1 (4.1)	21@1(3.6)	4@1 (3.1)	Sat (2.2)
1	23@1 (67.3)	19@1(8.6)	3@1(7.1)	4@1(5.9)	Sun (4.7)	5@1(2.6)	6@1(2.0)
2	23@1(56.3)	4@1(9.4)	19@1(8.9)	3@1(7.9)	Sun(6.6)	22@7(5.1)	2@25(1.6)
3	23@1 (45.0)	4@1(14.3)	3@1(12.2)	19@1(8.5)	Sun (7.3)	22@7(4.3)	3@6~(~2.5)
4	23@1 (44.8)	4@1(13.8)	19@1 (10.3)	Sun (10.2)	3@1(7.4)	22@7(2.5)	4@25(2.4)
5	23@1(34.1)	Sun (15.5)	5@1(9.7)	4@1(8.2)	Sat (7.4)	19@1(7.2)	22@7(3.8)
6	Sun (20.4)	23@1 (20.3)	Sat (12.8)	6@1(9.4)	Mon (8.7)	20@1 (6.0)	23@2(4.1)
7	Sun (23.9)	Sat (15.9)	23@1(12.1)	Mon (7.8)	20@1 (6.3)	6@1(5.3)	19@1(4.5)
8	Sun (28.3)	Sat (15.9)	20@1 (8.5)	23@1(7.9)	18@1(7.0)	Mon (5.4)	21@1(5.2)
9	Sun (25.9)	20@1(13.3)	Sat (13.3)	23@1(7.0)	22@1(6.3)	Mon (6.1)	18@1 (5.5)
10	Sun (21.8)	Sat (11.9)	22@1 (9.6)	20@1 (9.0)	Mon (6.6)	13@1(6.1)	23@1 (5.6)
11	Sun (17.2)	22@1(12.3)	Sat (11.7)	13@1 (9.2)	Mon (8.0)	23@1(5.8)	11@7(4.1)
12	Sun (15.6)	$13@1\ (12.0)$	Sat (10.7)	22@1(10.2)	Mon (8.1)	23@1(8.0)	17@1(4.9)
13	Sun (18.6)	Sat (11.2)	$13@1\ (11.1)$	22@1(10.0)	Mon (7.9)	23@1(7.3)	17@1 (6.8)
14	Sun (18.7)	Sat (10.1)	Mon (8.2)	23@1(8.2)	16@1(8.1)	22@1(7.6)	13@1(4.2)
15	Sun (19.8)	16@1(12.5)	Sat (8.9)	Mon (8.0)	23@1(7.1)	17@6(6.0)	17@1(4.9)
16	Sun (19.3)	17@1(12.5)	Sat (8.8)	16@1 (8.5)	Mon (8.4)	23@1(7.1)	17@6(6.7)
17	17@1(26.4)	Sun (16.1)	Mon (10.1)	17@6(10.0)	Sat (6.7)	17@7(6.3)	18@1 (6.0)
18	18@1 (33.8)	Sun (14.1)	Mon (9.9)	18@6(9.4)	Sat (6.0)	18@2(5.9)	18@5(4.6)
19	19@1(34.0)	Mon (9.4)	19@6(8.5)	Sun(8.2)	19@2(7.6)	8@6(5.0)	19@7(4.7)
20	20@1(23.9)	20@2(10.4)	Sat (9.5)	Mon (7.8)	21@1(7.5)	20@6(7.5)	Sun (7.0)
21	21@1(22.8)	Sat (10.9)	21@2(9.9)	Mon (6.2)	21@28(4.7)	Sun (4.6)	23@1(4.6)
22	22@1(30.6)	22@2(9.5)	Sat (7.7)	23@3~(~6.5)	22@28(6.1)	22@5(5.1)	22@35(5.0)
23	23@1(24.8)	22@1(17.5)	23@3(10.7)	22@2(9.6)	Sat (5.7)	0@5(5.0)	12@7(4.4)

Table 5: Most relevant coefficients for each hour. H@L represents the estimated parameter for $Y_{d-L,H}$ for a model on $Y_{d,h}$. Sun, Mon, ..., Sat represent the weekday dummies. The corresponding $\hat{\iota}_{h,i}$ value is given in parenthesis.

EPEX Switzerland

	Importance: 1	Importance: 2	Importance: 3	Importance: 4	Importance: 5	Importance: 6	Importance: 7
0	23@1(50.8)	22@1(11.4)	1@1(6.1)	Sat (5.6)	6@1(5.2)	5@1(4.2)	18@1(3.7)
1	23@1(47.6)	5@1(13.6)	2@1(12.0)	18@1(6.5)	Sun (5.6)	Sat (4.4)	2@3(3.9)
2	23@1(32.6)	2@1(12.7)	4@1(10.6)	Sun(8.4)	18@1(8.1)	5@1(5.4)	3@7(3.5)
3	23@1(30.7)	4@1(23.9)	Sun (9.6)	18@1(6.0)	4@7(4.3)	4@4(4.2)	5@1(3.7)
4	23@1(22.6)	4@1(20.8)	Sun (12.0)	5@1(6.7)	22@1(6.6)	4@4(5.1)	19@5(4.8)
5	23@1 (25.9)	Sun (19.2)	5@1(13.4)	Sat (6.9)	5@7(5.6)	18@1(4.9)	4@1(4.0)
6	Sun (27.4)	23@1 (20.8)	Sat (12.0)	20@1 (9.5)	22@1(4.8)	6@6(4.8)	5@1(4.7)
7	Sun (28.1)	Sat (12.4)	22@1(12.0)	20@1(9.9)	23@1(8.5)	18@1(5.4)	Mon (4.5)
8	Sun (22.8)	22@1(11.3)	19@1 (10.3)	Sat (8.8)	Mon (6.7)	10@1 (6.5)	$8@6\ (\ 5.6)$
9	Sun (16.9)	10@1 (10.7)	22@1 (9.8)	19@1(8.2)	Mon (7.7)	Sat (7.0)	9@2(6.5)
10	Sun (15.8)	22@1(14.2)	Mon (10.1)	11@1(9.7)	19@1 (9.5)	Sat (7.8)	10@1 (6.2)
11	Sun(8.3)	22@1(6.9)	10@2(6.3)	14@1 (5.0)	Sat (4.7)	Mon (4.4)	13@7(4.3)
12	Sun (8.2)	10@2(7.6)	22@1(7.2)	13@7(7.1)	23@1(5.1)	13@1(4.8)	12@1(4.5)
13	12@1(16.8)	Sun (15.8)	22@1(11.5)	13@7(9.0)	23@1(9.0)	Sat (8.8)	Mon (8.4)
14	Sun (17.0)	22@1(14.5)	Sat (10.2)	Mon (9.9)	13@7(8.6)	12@1(8.3)	14@1(7.3)
15	Sun (12.7)	22@1(10.6)	10@2(6.9)	Mon (6.7)	Sat (6.2)	23@1(5.3)	13@7(5.3)
16	Sun (14.5)	22@1(13.3)	Mon (9.1)	17@1(8.9)	Sat (6.8)	14@1(6.1)	18@1 (5.1)
17	17@1(15.0)	18@1 (10.3)	Sun(8.4)	22@1(6.5)	10@2(5.6)	17@6(4.7)	Mon (4.7)
18	18@1 (40.6)	18@6(9.9)	Mon (8.9)	Sun(8.8)	Sat (5.8)	23@1(4.8)	22@1(4.4)
19	19@1(28.9)	10@2~(11.2)	14@2(11.0)	Sun (5.2)	22@1(4.9)	23@1(4.7)	18@6(4.2)
20	20@1(26.7)	23@1(7.7)	Mon (6.8)	22@1(6.5)	9@2(5.9)	15@2(5.5)	Sun (5.4)
21	22@1(18.1)	23@1(17.7)	21@1(15.9)	Mon (9.8)	20@1(9.7)	13@7(4.6)	6@6(4.1)
22	22@1(29.6)	23@1(22.1)	Mon (9.0)	13@6(6.1)	13@7(4.7)	Sat (4.0)	20@1 (3.8)
23	23@1(39.6)	22@1(20.8)	Mon (9.8)	13@7(6.3)	6@6(4.6)	18@1 (4.6)	0@2(4.3)

Table 6: Most relevant coefficients for each hour. H@L represents the estimated parameter for $Y_{d-L,H}$ for a model on $Y_{d,h}$. Sun, Mon, ..., Sat represent the weekday dummies. The corresponding $\hat{\iota}_{h,i}$ value is given in parenthesis.

BELPEX Belgium

	Importance: 1	Importance: 2	Importance: 3	Importance: 4	Importance: 5	Importance: 6	Importance: 7
0	23@1(29.3)	0@1(10.4)	2@1(8.8)	7@7(6.9)	7@8(5.1)	1@1(4.3)	0@6(4.3)
1	2@1(19.9)	0@1(12.2)	23@1(8.8)	9@1(6.4)	1@1(5.6)	21@1(5.4)	13@1 (5.3)
2	4@1(23.2)	2@1(22.9)	0@1(7.6)	1@1(7.1)	12@6(6.2)	10@6 (5.9)	17@8(3.2)
3	4@1(22.8)	2@1(11.9)	3@1(8.6)	10@6 (5.4)	0@1(5.4)	8@7(4.2)	1@1(3.6)
4	4@1(33.7)	5@1(10.7)	10@6~(10.4)	3@1(8.5)	0@1(6.1)	4@7(5.0)	1@1(4.2)
5	5@1(38.7)	4@1(12.2)	6@6(11.5)	5@7(8.6)	15@8(5.1)	10@6(5.1)	6@2(4.1)
6	5@1(40.0)	6@7(13.1)	6@1(11.2)	7@6(7.0)	6@14(5.9)	6@6(5.7)	7@7(5.5)
7	5@1(29.9)	7@7(16.1)	7@1(12.9)	7@14(9.9)	7@6(7.1)	7@21(4.6)	8@7(3.8)
8	7@1(12.0)	10@7~(11.5)	5@1(10.9)	8@7~(10.6)	8@14(7.7)	3@1(6.5)	2@1(6.2)
9	10@7(59.7)	4@1(21.9)	5@1(18.3)	0@1(0.0)	0@2(0.0)	0@3~(~0.0)	0@4(0.0)
10	10@7(17.1)	2@1(13.5)	10@14(7.4)	10@35(6.2)	4@1(6.0)	6@1(5.5)	10@1 (5.1)
11	2@1(11.6)	10@7 (9.1)	11@14(7.8)	12@7(6.6)	11@35(6.0)	11@21(4.4)	4@1(4.1)
12	2@1(15.0)	12@1(8.9)	12@14(8.4)	13@7(7.4)	12@35(6.3)	10@7 (4.8)	12@7(3.7)
13	2@1(20.3)	13@7(13.4)	13@14(8.6)	3@1(7.5)	13@1(7.0)	16@7(6.8)	13@35(6.7)
14	3@1(11.4)	2@1(10.6)	16@7(9.6)	13@1(8.1)	14@14(7.8)	14@21(6.4)	14@35(5.3)
15	2@1(11.3)	15@14(10.1)	3@1(10.0)	16@7(6.7)	15@21(5.9)	16@1 (5.6)	15@35(5.2)
16	2@1(13.4)	16@14 (10.6)	17@1(9.0)	3@1(8.1)	16@7(7.5)	17@7(4.9)	16@28(4.7)
17	17@1(14.2)	17@7(12.4)	2@1(10.4)	17@14(7.8)	17@6(5.7)	4@1(4.3)	7@1(3.8)
18	18@1 (26.2)	2@1(11.0)	18@7(7.1)	18@14(6.5)	17@7(6.0)	7@1(5.6)	21@1(4.4)
19	19@1(14.1)	2@1(13.4)	19@7(8.2)	12@7(7.1)	7@1(6.7)	19@14(4.9)	19@2(4.5)
20	20@1(19.0)	2@1(10.4)	12@7(5.6)	20@2(5.3)	20@6(5.3)	3@1(4.7)	7@1(4.3)
21	2@1(15.9)	21@7(15.3)	21@1(13.0)	20@1(7.4)	21@6(6.2)	4@1(5.7)	21@14(4.8)
22	2@1(18.6)	22@6(11.4)	22@1 (9.8)	4@1(8.9)	21@7(7.9)	22@2(4.8)	10@1 (4.6)
23	2@1(26.9)	4@1(17.3)	23@1(8.5)	23@2(7.0)	21@7(5.5)	12@1 (5.4)	12@7(5.2)

Table 7: Most relevant coefficients for each hour. H@L represents the estimated parameter for $Y_{d-L,H}$ for a model on $Y_{d,h}$. Sun, Mon, ..., Sat represent the weekday dummies. The corresponding $\hat{\iota}_{h,i}$ value is given in parenthesis.

APX Netherlands

	Importance: 1	Importance: 2	Importance: 3	Importance: 4	Importance: 5	Importance: 6	Importance: 7
0	23@1(52.2)	2@1(14.7)	Sat (5.8)	9@1(5.3)	22@1(5.0)	21@1 (2.7)	1@8(2.1)
1	23@1 (22.9)	2@1(12.0)	5@1(7.2)	16@1(5.3)	1@6(5.3)	19@1(5.0)	22@1(4.7)
2	23@1 (22.5)	2@1(20.8)	5@1(11.1)	19@1(7.8)	3@1(6.3)	16@1 (5.6)	22@1 (5.3)
3	3@1(20.3)	23@1 (16.5)	5@1(9.1)	4@6(5.6)	Sun(4.6)	4@1(4.5)	21@1(4.2)
4	4@1(12.3)	3@1(12.3)	23@1(12.2)	5@1(11.5)	Sun (9.3)	5@7(5.6)	21@1(5.6)
5	5@1(23.5)	Sun (14.9)	23@1 (10.5)	6@7(9.9)	21@1(8.1)	19@1(6.9)	5@6(6.7)
6	Sun (23.2)	6@7~(12.5)	Sat (12.4)	23@1(7.0)	19@1(6.9)	5@1(6.6)	6@1(5.9)
7	Sun (27.5)	Sat (17.3)	20@1(6.9)	19@1(6.6)	7@1(6.1)	Mon (5.5)	7@7(3.8)
8	Sun (27.2)	Sat (17.0)	7@1(6.9)	7@7(6.0)	Mon (5.7)	19@1(5.7)	20@1 (5.7)
9	Sun (26.3)	Sat (11.6)	23@1(5.2)	20@1(4.3)	18@1(4.1)	7@7(4.0)	21@1(3.7)
10	Sun (25.9)	9@7~(10.8)	12@1 (10.5)	18@1(8.0)	Sat (6.9)	22@1(5.4)	10@6(4.8)
11	Sun (23.4)	12@1(11.8)	Sat (9.6)	22@1(7.0)	Mon (5.4)	16@1 (5.2)	10@7(4.1)
12	Sun (14.3)	12@1(11.5)	12@6(7.6)	Sat (6.1)	Mon (5.8)	22@1(4.2)	12@14(4.2)
13	Sun (16.2)	12@1(11.2)	Mon (10.6)	Sat (10.0)	16@1 (9.8)	12@6(6.1)	22@1 (5.6)
14	Sun (17.8)	Sat (12.3)	16@1(10.1)	12@1(8.9)	Mon (8.5)	15@7(7.3)	22@1(6.2)
15	Sun (18.7)	16@1 (14.6)	Sat (11.3)	15@7(9.0)	Mon (6.5)	22@1(6.5)	15@6(5.7)
16	Sun (16.7)	16@1(13.7)	Sat (10.3)	Mon (7.1)	16@7(6.4)	17@2(6.1)	17@1(5.5)
17	17@1(27.3)	Sun (11.5)	17@7(10.1)	Sat (9.2)	17@6(7.5)	17@2(5.6)	Mon (5.4)
18	18@1 (34.5)	18@2(8.2)	Mon (7.9)	18@7(6.1)	18@3(6.0)	17@6(5.3)	18@6(4.4)
19	19@1 (44.1)	19@2(15.4)	19@6(8.6)	19@3(6.6)	Mon (5.2)	19@14(5.1)	7@6(3.9)
20	20@1 (33.8)	20@2(16.3)	20@6(8.2)	20@3(7.8)	20@13(5.2)	15@6(5.1)	6@6(4.3)
21	21@1(25.4)	21@2(15.0)	21@4(6.0)	21@13(5.9)	Sat (5.3)	21@6(5.1)	0@1(4.7)
22	22@1(21.3)	22@2(13.2)	23@1(12.7)	22@13(5.4)	22@3(4.7)	Sat (4.4)	23@8(4.2)
23	23@1(27.2)	22@1(15.0)	23@2(8.8)	22@2(6.1)	23@3(5.4)	23@8(5.4)	23@28 (5.0)

Table 8: Most relevant coefficients for each hour. H@L represents the estimated parameter for $Y_{d-L,H}$ for a model on $Y_{d,h}$. Sun, Mon, ..., Sat represent the weekday dummies. The corresponding $\hat{\iota}_{h,i}$ value is given in parenthesis.

Nordpool Denmark West

	Importance: 1	Importance: 2	Importance: 3	Importance: 4	Importance: 5	Importance: 6	Importance: 7
0	23@1(70.2)	4@1 (9.4)	17@1(5.4)	23@2(4.4)	4@2 (4.1)	15@1(2.6)	3@4(2.4)
1	23@1 (61.8)	4@1(15.3)	17@1(6.6)	4@2(4.3)	3@4(4.2)	16@6(2.7)	3@1(1.8)
2	23@1 (62.0)	4@1(19.1)	3@1(5.3)	17@1(4.4)	3@4(2.7)	Sun (2.4)	4@6(2.0)
3	23@1 (48.8)	4@1(21.0)	4@6(11.1)	17@1(3.9)	3@1(3.7)	5@1(3.7)	16@6(3.2)
4	23@1 (48.3)	4@1(23.8)	Sun (6.7)	5@1(6.2)	4@6(4.7)	3@1(4.0)	17@1(2.5)
5	23@1 (37.2)	5@1(16.1)	Sun (12.8)	4@1(9.9)	Sat (6.9)	6@7(5.2)	18@1(3.7)
6	Sun (16.0)	23@1 (14.6)	21@1(13.9)	6@1(12.0)	Sat (11.1)	Mon (7.2)	19@1 (6.4)
7	13@1 (40.0)	16@1 (11.2)	18@1 (10.5)	Sat (10.4)	Sun (8.9)	Mon (8.4)	Fri (6.3)
8	13@1 (45.7)	Sat (13.0)	Sun (12.5)	18@1(11.1)	Mon (10.2)	Fri (7.4)	0@1(0.0)
9	13@1 (41.2)	16@1 (25.2)	17@1(8.2)	Mon (6.7)	Fri (5.1)	18@1(4.7)	Sat (4.0)
10	16@1 (33.1)	13@1 (28.7)	17@1(13.0)	15@1(10.4)	Mon (5.3)	Fri (3.8)	Sat (2.1)
11	16@1(32.6)	13@1 (29.8)	17@1(13.1)	15@1 (9.6)	Mon (5.7)	Fri (3.5)	Sat (2.4)
12	22@1(13.0)	21@1(10.0)	Sun (9.7)	13@1(8.8)	Sat (8.1)	18@1(7.8)	Mon (6.0)
13	Sun (10.9)	22@1(10.4)	21@1(10.2)	Sat (8.8)	17@1(7.0)	13@7(6.3)	Mon (6.0)
14	Sun (12.3)	21@1(10.4)	Sat (9.1)	17@1 (9.0)	23@1(7.7)	14@28(7.0)	22@1(6.5)
15	Sun (13.1)	21@1(10.1)	17@1(9.4)	Sat (8.9)	23@1(8.6)	15@28(6.7)	16@7(6.6)
16	17@1 (16.9)	16@7~(11.3)	16@1 (10.8)	23@1(10.4)	Sun (9.6)	17@6(6.8)	Mon (6.7)
17	17@1(22.4)	18@1(12.1)	16@7(9.1)	17@6(8.8)	Mon (8.7)	Sun (8.2)	Sat (5.9)
18	18@1 (40.1)	Mon (8.0)	18@2(7.5)	19@1(7.3)	17@6(6.0)	18@6(4.9)	Sun(4.9)
19	19@1(47.7)	12@7(10.0)	19@2(8.0)	20@5(5.6)	18@6(5.4)	14@6(4.0)	20@1(3.5)
20	21@1(22.0)	19@1 (10.8)	20@1(7.7)	20@6(7.6)	19@2(6.9)	12@7(6.8)	20@5(5.4)
21	21@1(34.8)	12@7(7.0)	23@1(6.7)	22@1(6.6)	21@5(6.0)	6@6(5.7)	21@8(4.7)
22	22@1 (31.6)	22@5(11.5)	23@1(8.2)	22@2(8.1)	21@1(5.8)	12@7(5.3)	22@4(4.5)
23	23@1 (43.3)	22@2(8.8)	23@3(4.4)	21@1(4.4)	23@6(3.5)	5@4(3.0)	22@5(3.0)

Table 9: Most relevant coefficients for each hour. H@L represents the estimated parameter for $Y_{d-L,H}$ for a model on $Y_{d,h}$. Sun, Mon, ..., Sat represent the weekday dummies. The corresponding $\hat{\iota}_{h,i}$ value is given in parenthesis.

Nordpool Denmark East

	Importance: 1	Importance: 2	Importance: 3	Importance: 4	Importance: 5	Importance: 6	Importance: 7
0	23@1(68.12)	4@1 (8.53)	4@2(8.08)	19@1 (4.19)	0@1(3.25)	23@2(2.11)	4@5(1.73)
1	$23@1 \ (64.05)$	4@1(14.74)	4@2(7.74)	3@1(7.09)	4@5(3.61)	19@1(2.44)	4@6(0.33)
2	23@1 (55.15)	4@1(17.31)	3@1(12.03)	4@2(5.01)	4@5(3.51)	4@6(3.10)	Sun (1.65)
3	23@1 (43.57)	4@1(22.45)	3@1(12.87)	4@6(11.09)	4@2(2.93)	Sun (2.82)	4@5(2.73)
4	23@1(42.37)	4@1(28.57)	3@1 (8.57)	4@6(6.15)	Sun (3.94)	4@2(3.01)	4@5(2.87)
5	23@1(33.17)	4@1(15.31)	Sun (11.31)	5@1(10.39)	5@6(5.68)	Sat (5.64)	19@1 (5.58)
6	23@1 (19.80)	Sun (13.82)	6@1(10.20)	19@1 (9.52)	Sat (9.12)	22@1(6.43)	21@1(6.40)
7	19@1 (23.52)	23@1 (16.48)	Sun (12.43)	21@1(11.61)	Sat (9.63)	16@1 (8.89)	Mon (7.23)
8	19@1(23.30)	21@1(17.47)	23@1(14.63)	Sun (10.59)	Mon (7.90)	Sat (7.64)	16@1(7.44)
9	19@1 (24.02)	21@1(23.68)	23@1 (13.26)	Mon (10.61)	Sun (8.49)	8@7(5.58)	Sat (5.51)
10	21@1(33.92)	19@1 (18.90)	23@1(11.54)	Mon (11.25)	8@7(6.11)	16@1 (5.55)	22@1 (5.41)
11	21@1(22.86)	16@1(12.18)	19@1(10.54)	23@1(9.81)	Mon (8.30)	Sun (5.61)	8@1(4.87)
12	21@1(20.38)	16@1(13.66)	19@1(8.47)	Sun (6.78)	Mon (6.72)	Sat (6.44)	23@1(6.30)
13	21@1(23.12)	14@1(9.47)	Mon (7.04)	Sat (6.94)	19@1 (6.93)	Sun (6.88)	16@1 (6.79)
14	21@1(21.38)	16@1(14.27)	23@1(9.69)	Sun (7.40)	Sat (6.98)	17@6(6.70)	8@1(5.72)
15	21@1(17.41)	16@1(15.43)	23@1(8.86)	8@1(8.04)	Sun (6.61)	Mon (5.94)	Sat (5.70)
16	16@1(27.37)	23@1(8.70)	21@1(7.92)	Mon (6.52)	17@6(6.11)	17@7(5.18)	16@6(4.64)
17	16@6(34.01)	17@1(24.75)	16@1(24.74)	16@7(13.44)	23@1(2.41)	Sun (0.64)	0@1(0.00)
18	16@6(40.52)	19@1(30.44)	16@1 (19.99)	16@7 (9.05)	0@1(0.00)	0@2(0.00)	$0@3\ (\ 0.00)$
19	19@1 (39.15)	22@1 (9.71)	23@1(6.63)	19@2(6.60)	Mon (6.52)	17@6 (5.66)	16@7(3.73)
20	20@1(24.92)	22@1(14.63)	21@5(6.47)	19@1(6.00)	23@1 (5.52)	Sat (5.08)	17@6(4.95)
21	21@1(22.91)	22@1(15.72)	23@1(11.11)	21@5(9.02)	17@7 (5.04)	Sat (4.53)	Fri (3.43)
22	22@1(37.78)	23@1(17.21)	22@2(11.15)	22@5(10.50)	14@6(3.15)	22@6(2.97)	22@7(2.77)
23	23@1(52.37)	22@2(6.60)	0@1 (5.79)	22@5(5.78)	22@1(5.64)	0@5(3.39)	3@3(3.17)

Table 10: Most relevant coefficients for each hour. H@L represents the estimated parameter for $Y_{d-L,H}$ for a model on $Y_{d,h}$. Sun, Mon, ..., Sat represent the weekday dummies. The corresponding $\hat{\iota}_{h,i}$ value is given in parenthesis.

Nordpool Sweden(4)

	Importance: 1	Importance: 2	Importance: 3	Importance: 4	Importance: 5	Importance: 6	Importance: 7
0	23@1(74.57)	1@1(10.65)	4@1(9.78)	18@3(2.33)	17@2(0.96)	7@4(0.63)	Sat (0.50)
1	23@1 (46.44)	4@1(13.77)	1@1(9.20)	22@6(5.28)	22@2(4.92)	3@3(3.10)	21@2(2.90)
2	23@1 (41.80)	4@1(16.20)	22@2(6.91)	22@6(6.81)	3@1(6.64)	3@3(4.90)	22@3(3.90)
3	23@1 (35.98)	4@1(22.09)	22@6(8.95)	22@2(6.30)	22@3(4.94)	3@3(4.83)	4@5(3.39)
4	23@1 (27.78)	4@1(16.84)	22@2(5.81)	23@6(5.18)	4@5(4.80)	22@6(3.42)	5@6(2.89)
5	$23@1 \ (26.23)$	4@1(17.77)	21@1(10.58)	22@6(8.55)	22@7(5.45)	5@7(4.66)	5@6(4.40)
6	21@1(25.42)	23@1(14.07)	22@6(8.11)	8@7(7.58)	11@7(6.84)	11@1(6.48)	19@1 (5.70)
7	21@1(17.21)	8@7(16.37)	11@7 (9.53)	11@1 (9.26)	10@7 (5.11)	16@1 (5.00)	18@1(4.56)
8	21@1(24.41)	8@7~(15.96)	11@7(12.62)	11@1(7.48)	19@1 (4.13)	21@3(3.76)	16@1(3.73)
9	21@1(30.62)	8@7(16.67)	11@7(12.07)	19@1(4.19)	11@1(4.03)	21@3(4.00)	20@7(2.73)
10	21@1(26.27)	8@7(17.43)	11@7 (10.60)	18@1 (6.24)	11@1 (5.22)	10@7(3.11)	7@1(3.06)
11	21@1(30.07)	8@7~(13.90)	11@7 (9.53)	11@1(5.41)	6@7(4.19)	18@1(3.62)	16@1(3.40)
12	21@1(19.11)	22@1(11.14)	16@1(7.47)	8@7(5.79)	11@1 (5.62)	19@1(4.79)	7@2(4.32)
13	21@1(17.35)	8@7(7.06)	22@1 (5.96)	7@2(5.10)	8@1(4.23)	23@1(3.94)	18@1(3.45)
14	21@1(18.97)	8@7(5.70)	7@2(4.75)	18@1(4.40)	16@1 (4.23)	8@1(4.14)	11@1(3.88)
15	21@1(20.49)	16@1 (8.22)	8@1(7.50)	8@7(4.90)	11@7(4.86)	23@1(4.70)	22@1(4.44)
16	21@1(13.72)	16@1(12.11)	11@1(5.77)	7@2(5.64)	11@7(4.82)	22@6(4.33)	22@1(4.25)
17	21@1 (18.05)	8@7(12.49)	17@1(10.94)	9@1 (10.56)	11@7 (10.54)	11@1(4.33)	21@3(3.53)
18	21@1(19.43)	8@7(17.33)	11@7(10.77)	19@1(8.85)	18@1 (6.29)	11@1(5.24)	21@3(4.49)
19	19@1 (11.97)	21@1(10.90)	8@7(9.15)	11@7(8.18)	11@1(7.31)	20@1 (5.64)	7@5(5.00)
20	21@1(23.22)	8@7(9.37)	11@7 (8.54)	20@1(7.37)	22@1 (5.51)	19@1 (5.50)	7@5(4.95)
21	21@1(54.22)	22@1 (26.19)	23@1(7.75)	14@6(3.83)	Mon (2.46)	16@6(1.92)	Fri (1.51)
22	22@1 (66.34)	23@1 (10.96)	0@1(4.78)	22@5(3.76)	21@1(2.87)	9@4(2.06)	21@6(1.94)
23	23@1(54.20)	22@1(26.52)	23@2(6.36)	0@1(3.83)	1@3(2.82)	4@5(2.70)	23@5(1.72)

Table 11: Most relevant coefficients for each hour. H@L represents the estimated parameter for $Y_{d-L,H}$ for a model on $Y_{d,h}$. Sun, Mon, ..., Sat represent the weekday dummies. The corresponding $\hat{\iota}_{h,i}$ value is given in parenthesis.

POLPX Poland

	Importance: 1	Importance: 2	Importance: 3	Importance: 4	Importance: 5	Importance: 6	Importance: 7
0	23@1(23.8)	0@1(13.9)	7@1(8.5)	1@1(7.0)	23@2(6.8)	0@7(6.0)	22@1(3.5)
1	23@1(28.1)	1@1(23.1)	1@7(9.6)	3@1(8.9)	22@1 (5.8)	Sat (5.6)	6@1(4.5)
2	23@1(23.6)	3@1(15.2)	2@1(10.4)	22@1(8.9)	3@7(8.1)	4@1(7.7)	5@1(6.7)
3	4@1(23.3)	23@1(17.5)	3@1(15.3)	3@7(9.5)	22@1(7.7)	Sun (7.2)	Sat (4.2)
4	4@1(29.4)	23@1(13.2)	5@1(11.6)	Sun (10.4)	3@7(8.6)	22@1(8.2)	4@5(4.6)
5	5@1(32.6)	Sun (10.9)	22@1(10.6)	23@1(10.1)	5@7(8.7)	6@5(5.1)	22@2(3.4)
6	6@1(23.2)	Sun (11.4)	23@1(9.0)	6@7(8.2)	22@1(7.7)	Mon (4.8)	6@6(3.8)
7	Sun (17.8)	23@1 (14.5)	7@1(14.2)	7@7(12.0)	Mon (7.4)	Sat (6.7)	7@6~(~5.3)
8	8@1(17.9)	Sun (12.9)	22@1(11.3)	8@7~(10.5)	Mon (5.7)	22@8(4.1)	Sat (4.0)
9	9@1(20.1)	Sun (16.6)	22@1(7.5)	Mon (7.2)	Sat (5.3)	9@7(4.8)	9@14(4.5)
10	10@1 (20.4)	Sun (17.1)	Mon (10.1)	21@1(6.8)	22@1(6.1)	12@1 (5.2)	8@6(3.9)
11	12@1(19.7)	14@7(9.7)	Sun (7.4)	14@8(5.6)	22@1(4.7)	12@3(4.0)	Mon (3.9)
12	12@1(24.1)	14@7(9.1)	12@21(5.9)	12@14(5.2)	13@8(4.7)	Mon (4.1)	Sun (4.0)
13	12@1(19.1)	14@7(8.3)	13@14(5.6)	13@21(5.3)	15@2(4.0)	Mon (3.9)	13@8(3.8)
14	14@1(12.3)	14@7 (9.8)	12@1 (9.7)	11@8(6.5)	14@21(6.5)	14@14(5.1)	15@1(4.7)
15	15@1(26.2)	14@7~(11.5)	Sun (6.7)	14@8(6.3)	15@14(5.5)	Mon (5.4)	22@1(5.3)
16	16@1(14.5)	16@7(8.9)	17@1(7.8)	22@1(7.2)	Sun(6.5)	16@21(4.9)	15@1(4.7)
17	17@1(41.4)	17@6(10.2)	Sun (9.0)	18@1(7.9)	16@7(7.7)	Mon (7.5)	16@8(5.6)
18	18@1(51.5)	18@6(8.6)	Sun(7.0)	Mon (6.5)	16@7(4.8)	16@8(4.0)	18@8(3.8)
19	19@1 (46.2)	18@8(7.1)	18@6(5.1)	Sun(4.2)	Mon (4.1)	19@7(3.6)	20@1(3.4)
20	20@1(56.0)	Mon (5.5)	20@6(4.6)	22@8(4.5)	22@7(3.9)	Sun(2.9)	20@20(2.6)
21	21@1(43.1)	22@7(8.6)	20@1(5.6)	11@8(5.4)	21@6(5.2)	21@7(5.2)	Sat (4.8)
22	22@1(34.6)	22@7(20.7)	22@8(6.7)	22@6(5.5)	22@35(4.5)	22@21(4.1)	23@1(3.8)
23	23@1(51.4)	23@7(10.8)	23@6 (9.1)	Sat (5.6)	Mon (5.2)	23@21 (4.2)	1@3(2.4)

Table 12: Most relevant coefficients for each hour. H@L represents the estimated parameter for $Y_{d-L,H}$ for a model on $Y_{d,h}$. Sun, Mon, ..., Sat represent the weekday dummies. The corresponding $\hat{\iota}_{h,i}$ value is given in parenthesis.

OTE Czech

	Importance: 1	Importance: 2	Importance: 3	Importance: 4	Importance: 5	Importance: 6	Importance: 7
0	23@1(55.3)	21@1(10.1)	4@1 (9.2)	7@1 (8.2)	Sat (5.1)	Sun (3.9)	3@1(3.1)
1	23@1(43.4)	4@1(14.1)	Sun (11.7)	21@1(9.7)	7@1(9.4)	3@1(4.5)	19@1(2.4)
2	23@1(37.3)	4@1(18.5)	Sun (10.4)	20@1(7.3)	18@1(4.7)	3@6(4.5)	19@1(4.4)
3	23@1(33.3)	4@1(22.4)	Sun (11.9)	20@1 (9.2)	19@1(4.5)	3@1(4.1)	18@1(4.1)
4	23@1(28.4)	4@1(21.6)	Sun (15.3)	20@1(8.3)	3@6(5.1)	18@1(4.5)	19@1(4.2)
5	23@1 (25.5)	Sun (20.2)	20@1(11.1)	4@1(9.4)	5@6(7.9)	19@1(4.2)	6@7(3.1)
6	Sun (28.6)	23@1 (16.8)	20@1(15.4)	Sat (12.5)	Mon (7.6)	6@1(5.4)	19@1(4.5)
$\overline{7}$	Sun (25.7)	20@1(17.3)	Sat (14.1)	23@1(8.1)	Mon (7.0)	7@1(4.6)	18@1(4.4)
8	Sun (25.4)	20@1(16.4)	Sat (13.7)	Mon (7.7)	8@1(6.7)	23@1(5.9)	8@7(3.9)
9	Sun (22.4)	20@1(18.4)	Sat (11.0)	11@1(9.8)	Mon (9.5)	17@1(6.9)	8@7(6.4)
10	Sun (17.5)	20@1(12.0)	11@1 (10.5)	Mon (10.4)	22@1(9.4)	Sat (9.1)	11@7(5.3)
11	22@1(14.1)	Sun (12.6)	Mon (11.3)	11@1(11.2)	Sat (8.3)	20@1 (6.3)	12@1 (5.7)
12	Sun (13.0)	22@1(11.1)	Mon (10.3)	14@1(9.7)	Sat (8.0)	12@1(6.6)	14@7(4.5)
13	14@1(14.3)	Sun (14.1)	22@1(13.3)	Mon (12.1)	Sat (8.6)	14@7(7.1)	17@1(5.4)
14	Sun (16.3)	14@1(12.2)	Mon (10.6)	22@1(9.7)	Sat (8.2)	14@7(7.2)	16@1 (6.9)
15	Sun (17.7)	16@1(14.5)	Mon (11.1)	22@1(9.9)	Sat (7.6)	14@7(6.8)	17@1(6.7)
16	Sun (16.6)	16@1(14.9)	17@1(12.3)	Mon (10.3)	22@1(9.4)	16@7(7.1)	Sat (6.7)
17	17@1(35.0)	Sun (12.7)	Mon (12.0)	17@7(9.5)	17@6(5.7)	Sat (5.3)	22@1(3.8)
18	18@1 (30.6)	Mon (10.9)	18@6~(10.5)	Sun(9.4)	18@7(7.7)	17@1(7.1)	Sat (4.9)
19	19@1(39.4)	Mon (12.0)	19@6(7.5)	19@7(6.9)	20@1(6.7)	Sun (6.1)	Sat (5.8)
20	20@1 (45.8)	Mon (13.2)	20@6~(10.9)	Sat (6.5)	20@7(6.3)	22@1(4.2)	20@21(3.4)
21	21@1(30.6)	22@1(14.0)	21@6(10.1)	Mon (9.5)	Sat (6.2)	21@7(5.6)	20@1(4.9)
22	22@1(45.7)	22@6(8.6)	Mon (5.6)	22@4(5.5)	Sat (4.2)	22@21(3.9)	22@7(3.8)
23	22@1(30.6)	23@1(26.1)	23@6(7.9)	Mon (6.9)	22@2 (4.1)	23@4(3.7)	23@35(2.7)

Table 13: Most relevant coefficients for each hour. H@L represents the estimated parameter for $Y_{d-L,H}$ for a model on $Y_{d,h}$. Sun, Mon, ..., Sat represent the weekday dummies. The corresponding $\hat{\iota}_{h,i}$ value is given in parenthesis.