

# ON THE DETERMINATION OF NUMBER FIELDS BY ARTIN L-FUNCTIONS

JÜRGEN KLÜNERS AND FLORIN NICOLAE

**ABSTRACT.** Let  $k$  be a number field,  $K/k$  a finite Galois extension with Galois group  $G$ ,  $\chi$  a faithful character of  $G$ . We prove that the Artin L-function  $L(s, \chi, K/k)$  determines the Galois closure of  $K$  over  $\mathbb{Q}$ . In the special case  $k = \mathbb{Q}$  it also determines the character  $\chi$ .

*Key words:* Number fields, Galois extension, Artin L-function

*Mathematics Subject Classification:* 11R42

## 1. INTRODUCTION

Let  $k$  be a number field,  $K/k$  a finite Galois extension with Galois group  $G$ ,  $\chi$  a faithful character of  $G$ . In Theorem 6 we prove that the Artin L-function  $L(s, \chi, K/k)$  determines the Galois closure  $\tilde{K}$  of  $K$  over  $\mathbb{Q}$ . In the special case  $k = \mathbb{Q}$  we prove in Theorem 5 that the Artin L-function determines  $K$  and the (faithful) character  $\chi$ . We give examples that in the case  $k \neq \mathbb{Q}$  we cannot expect more, especially there exist non-isomorphic arithmetically equivalent fields which cannot be distinguished by Artin L-functions.

The restriction to faithful characters is natural: let  $K/k$  be a finite normal extension with  $\text{Gal}(K/k) = G$ , and let  $\chi$  be a character of  $G$  with  $\text{Ker}(\chi) \neq \{1\}$ . Let  $L$  be the fixed field of  $\text{Ker}(\chi)$ ,  $H := G/\text{Ker}(\chi)$  the Galois group of  $L/k$ ,  $\varphi : H \rightarrow \mathbb{C}$ ,  $\varphi(\sigma\text{Ker}(\chi)) := \chi(\sigma)$  for  $\sigma \in G$ . We have that

$$L(s, \chi, K/k) = L(s, \varphi, L/k),$$

and  $\varphi$  is faithful.

As particular cases we obtain that the Dedekind zeta function of a number field determines its normal closure ([4], Theorem 1, p. 345) and that a Galois number field is determined by any Artin L-function corresponding to a character which contains all irreducible characters of the Galois group, the result of [3].

## 2. PROPERTIES OF ARTIN L-FUNCTIONS

We recall fundamental properties of Artin L-functions needed in the sequel.

**Proposition 1.** *Let  $k$  be a number field,  $K/k$  a finite Galois extension with Galois group  $G$ ,  $\chi$  a character of  $G$ . Let  $L$  be a finite Galois extension of  $k$  which contains  $K$ ,  $U := \text{Gal}(L/k)$ ,  $V := \text{Gal}(L/K)$ . We identify the groups  $G$  and  $U/V$ . Let*

$$\tilde{\chi} : U \rightarrow \mathbb{C}, \quad \tilde{\chi}(\sigma) := \chi(\sigma V).$$

We have

$$L(s, \tilde{\chi}, L/k) = L(s, \chi, K/k).$$

*Proof.* This follows straightforward from the definition of L-functions: [1], p. 297, formula (8).  $\square$

**Proposition 2.** *Let  $k$  be a number field,  $K/k$  a finite Galois extension with Galois group  $G$ . Let  $k \subseteq L \subseteq K$  be an intermediate field,  $H := \text{Gal}(K/L)$ ,  $\chi$  a character of  $H$ , and  $\chi^G$  be the induced character on  $G$ . We have*

$$L(s, \chi^G, K/k) = L(s, \chi, K/L).$$

*Proof.* This is a deep property proved by Emil Artin: [1], p. 297, formula (9).  $\square$

**Proposition 3.** *Let  $K/\mathbb{Q}$  be a finite Galois extension with Galois group  $G$ ,  $\chi$  and  $\varphi$  characters of  $G$ . If*

$$L(s, \chi, K/\mathbb{Q}) = L(s, \varphi, K/\mathbb{Q})$$

*then*

$$\chi = \varphi.$$

*Proof.* This follows from the definition of Artin L-functions and Tschebotarev's density theorem. Alternately, it was proved in [2], p. 179, Theorem 1 that if  $\chi \neq \varphi$  then the functions  $L(s, \chi, K/\mathbb{Q})$  and  $L(s, \varphi, K/\mathbb{Q})$  are linearly independent over  $\mathbb{C}$ .  $\square$

We remark that Proposition 3 is not true for Artin L-functions in Galois extensions  $K/k$  with base  $k \neq \mathbb{Q}$ . E.g. let  $K/\mathbb{Q}$  be a normal  $S_3$ -extension and denote by  $k$  the unique quadratic subfield of  $K/\mathbb{Q}$ . Then  $\text{Gal}(K/k) = C_3$ , the cyclic group of order 3. Denote by  $\chi_2$  and  $\chi_3$  the two non-trivial characters of  $C_3$ . Then by Proposition 2 we get for  $j = 1, 2$ :

$$L(s, \chi_j^{S_3}, K/\mathbb{Q}) = L(s, \chi_j, K/k).$$

Using  $\chi_2^{S_3} = \chi_3^{S_3}$  we get that  $L(s, \chi_2, K/k) = L(s, \chi_3, K/k)$ , but  $\chi_2 \neq \chi_3$ .


## 3. RESULTS

**Proposition 4.** *Let  $k$  be a number field,  $K/k$  a finite Galois extension with Galois group  $G$ , and  $\chi$  be a faithful character of  $G$ . Let  $\tilde{K}$  be the Galois closure of  $K$  over  $\mathbb{Q}$ ,  $\tilde{G}$  the Galois group of  $\tilde{K}/\mathbb{Q}$ ,  $U := \text{Gal}(\tilde{K}/k)$ , and  $V := \text{Gal}(\tilde{K}/K)$ . We identify the groups  $G$  and  $U/V$ . Let*

$$\tilde{\chi} : U \rightarrow \mathbb{C}, \quad \tilde{\chi}(\sigma) := \chi(\sigma V),$$

*and let  $\tilde{\chi}^{\tilde{G}}$  be the induced character of  $\tilde{\chi}$  on  $\tilde{G}$ . Then  $\tilde{\chi}^{\tilde{G}}$  is faithful, and*

$$L(s, \chi, K/k) = L(s, \tilde{\chi}^{\tilde{G}}, \tilde{K}/\mathbb{Q}).$$



*Proof.* We have

$$\text{Ker}(\tilde{\chi}) = V,$$

since  $\chi$  is faithful. We have

$$\bigcap_{\sigma \in \tilde{G}} \sigma V \sigma^{-1} = 1,$$

since  $\tilde{K}$  is the Galois closure of  $K$  over  $\mathbb{Q}$ . We have

$$\text{Ker}(\tilde{\chi}^{\tilde{G}}) = \bigcap_{\sigma \in \tilde{G}} \sigma \text{Ker}(\tilde{\chi}) \sigma^{-1} = \bigcap_{\sigma \in \tilde{G}} \sigma V \sigma^{-1} = 1,$$

hence  $\tilde{\chi}^{\tilde{G}}$  is faithful. We have

$$L(s, \chi, K/k) = L(s, \tilde{\chi}, \tilde{K}/k) = L(s, \tilde{\chi}^{\tilde{G}}, \tilde{K}/\mathbb{Q}),$$

by Propositions 1 and 2. □

We prove that a Galois number field is determined by any Artin L-function corresponding to a faithful character of the Galois group.

**Theorem 5.** *Let  $K_1/\mathbb{Q}$ ,  $K_2/\mathbb{Q}$  be finite Galois extensions. For  $j = 1, 2$  let  $G_j$  be the Galois group of  $K_j/\mathbb{Q}$ ,  $\chi_j$  a faithful character of  $G_j$ . If*

$$L(s, \chi_1, K_1/\mathbb{Q}) = L(s, \chi_2, K_2/\mathbb{Q})$$

*then*

$$K_1 = K_2 \text{ and } \chi_1 = \chi_2.$$

*Proof.* Let  $L := K_1 \cdot K_2$ . For  $j = 1, 2$  let

$$\tilde{\chi}_j : \text{Gal}(L/\mathbb{Q}) \rightarrow \mathbb{C}, \quad \tilde{\chi}_j(\sigma) := \chi_j(\sigma \text{Gal}(L/K_j)),$$

where we identify  $G_j$  with the factor group  $\text{Gal}(L/\mathbb{Q})/\text{Gal}(L/K_j)$ . Then  $\tilde{\chi}_j$  is a character of  $\text{Gal}(L/\mathbb{Q})$ , and

$$\text{Ker}(\tilde{\chi}_j) = \text{Gal}(L/K_j),$$

since  $\chi_j$  is faithful. We have

$$L(s, \chi_j, K_j/\mathbb{Q}) = L(s, \tilde{\chi}_j, L/\mathbb{Q}),$$

by Proposition 1. If  $L(s, \chi_1, K_1/\mathbb{Q}) = L(s, \chi_2, K_2/\mathbb{Q})$  then

$$L(s, \tilde{\chi}_1, L/\mathbb{Q}) = L(s, \tilde{\chi}_2, L/\mathbb{Q}),$$

hence, by Proposition 3,

$$\tilde{\chi}_1 = \tilde{\chi}_2.$$

This implies

$$\text{Gal}(L/K_1) = \text{Ker}(\tilde{\chi}_1) = \text{Ker}(\tilde{\chi}_2) = \text{Gal}(L/K_2),$$

hence

$$K_1 = K_2$$

and, by Proposition 3,

$$\chi_1 = \chi_2.$$

□

In the final example of this paper we show that we cannot expect a similar result to Theorem 5 for normal extension  $K_j/k$  for  $k \neq \mathbb{Q}$  and  $j = 1, 2$ . In the next theorem we give a version for relative extensions.

**Theorem 6.** *Let  $k_1$  and  $k_2$  be number fields. For  $j = 1, 2$  let  $K_j/k_j$  be a finite Galois extension with the Galois group  $G_j$ ,  $\tilde{K}_j$  the normal closure of  $K_j$  over  $\mathbb{Q}$ ,  $\chi_j$  a faithful character of  $G_j$ ,  $U_j := \text{Gal}(\tilde{K}_j/k_j)$ ,  $V_j := \text{Gal}(\tilde{K}_j/K_j)$ . We identify the groups  $G_j$  and  $U_j/V_j$ . Let*

$$\tilde{\chi}_j : U_j \rightarrow \mathbb{C}, \quad \tilde{\chi}_j(\sigma) := \chi_j(\sigma V_j).$$

*Let  $\tilde{G}_j := \text{Gal}(\tilde{K}_j/\mathbb{Q})$ , and let  $\tilde{\chi}_j^{\tilde{G}_j}$  be the induced character of  $\tilde{\chi}_j$  on  $\tilde{G}_j$ . If*


$$L(s, \chi_1, K_1/k_1) = L(s, \chi_2, K_2/k_2)$$

then

$$\tilde{K}_1 = \tilde{K}_2, \tilde{G}_1 = \tilde{G}_2$$

and

$$\tilde{\chi}_1^{\tilde{G}_1} = \tilde{\chi}_2^{\tilde{G}_2}.$$



*Proof.* For  $j = 1, 2$  the character  $\tilde{\chi}_j^{\tilde{G}_j}$  is faithful and we have

$$L(s, \chi_j, K_j/k_j) = L(s, \tilde{\chi}_j^{\tilde{G}_j}, \tilde{K}_j/\mathbb{Q}),$$

by Proposition 4. If

$$L(s, \chi_1, K_1/k_1) = L(s, \chi_2, K_2/k_2)$$

then

$$L(s, \tilde{\chi}_1^{\tilde{G}_1}, \tilde{K}_1/\mathbb{Q}) = L(s, \tilde{\chi}_2^{\tilde{G}_2}, \tilde{K}_2/\mathbb{Q}),$$

hence

$$\tilde{K}_1 = \tilde{K}_2, \tilde{G}_1 = \tilde{G}_2$$

and

$$\tilde{\chi}_1^{\tilde{G}_1} = \tilde{\chi}_2^{\tilde{G}_2},$$

by Theorem 5. □

We obtain now the well-known result ([4], Theorem 1, p. 345) that the zeta function of an algebraic number field determines its normal closure.

**Corollary 7.** *Let  $K_1$  and  $K_2$  be number fields. For  $j = 1, 2$  let  $\tilde{K}_j$  be the normal closure of  $K_j$  over  $\mathbb{Q}$ ,  $V_j := \text{Gal}(\tilde{K}_j/K_j)$ ,  $\tilde{G}_j := \text{Gal}(\tilde{K}_j/\mathbb{Q})$ , and let  $1_{V_j}$  be the trivial character of  $V_j$ . If*

$$\zeta_{K_1}(s) = \zeta_{K_2}(s)$$

then

$$\tilde{K}_1 = \tilde{K}_2, \tilde{G}_1 = \tilde{G}_2$$

and

$$1_{V_1}^{\tilde{G}_1} = 1_{V_2}^{\tilde{G}_2}.$$

*Proof.* For  $j = 1, 2$  let  $k_j = K_j$ ,  $G_j = \{1\}$ ,  $\chi_j := 1_{G_j}$ ,  $U_j := V_j$ . The character  $\chi_j$  is faithful, and

$$L(s, \chi_j, K_j/k_j) = \zeta_{K_j}(s).$$

We have

$$\tilde{\chi}_j : U_j \rightarrow \mathbb{C}, \quad \tilde{\chi}_j(\sigma) := \chi_j(\sigma V_j) = \chi_j(1) = 1,$$

hence

$$\tilde{\chi}_j = 1_{V_j}.$$

If

$$\zeta_{K_1}(s) = \zeta_{K_2}(s)$$

then

$$L(s, \chi_1, K_1/k_1) = L(s, \chi_2, K_2/k_2),$$

hence

$$\tilde{K}_1 = \tilde{K}_2, \quad \tilde{G}_1 = \tilde{G}_2$$

and

$$1_{V_1}^{\tilde{G}_1} = 1_{V_2}^{\tilde{G}_2},$$

by Theorem 6. □

We finish with an example which shows that all Artin  $L$ -functions in the Galois closures of the corresponding fields coincide. Let  $k = k_1 = k_2 := \mathbb{Q}(\sqrt[4]{3})$ ,  $K_1 := \mathbb{Q}(\sqrt[8]{3})$ ,  $K_2 := \mathbb{Q}(\sqrt[8]{16 \cdot 3})$ . The extensions  $K_1/k$  and  $K_2/k$  are Galois with Galois groups  $G_1$  and  $G_2$  of order 2. The fields  $K_1$  and  $K_2$  are non-isomorphic and have the same zeta function ([4], p. 350):

$$L(s, \text{Reg}_{G_1}, K_1/k) = \zeta_{K_1}(s) = \zeta_{K_2}(s) = L(s, \text{Reg}_{G_2}, K_2/k).$$

For  $j = 1, 2$  we have

$$\text{Reg}_{G_j} = 1_{G_j} + \chi_j,$$

where  $\chi_j$  is the non-trivial irreducible character of  $G_j$ . We have

$$L(s, 1_{G_1}, K_1/k) = \zeta_k(s) = L(s, 1_{G_2}, K_2/k),$$

hence

$$L(s, \chi_1, K_1/k) = L(s, \chi_2, K_2/k).$$

It follows that the Artin  $L$ -functions of  $K_1/k$  are identical with the Artin  $L$ -functions of  $K_2/k$ . We have

$$\tilde{K}_1 = \tilde{K}_2, \quad \tilde{G}_1 = \tilde{G}_2 = G,$$

by Theorem 6, since  $\text{Reg}_{G_1}$  and  $\text{Reg}_{G_2}$  are faithful. Using the notation of Theorem 6, there is an outer automorphism  $\alpha$  of  $G$  such that

$$V_2 = \alpha(V_1).$$

The Artin L-functions of  $\tilde{K}_1/K_1$  are identical with the Artin L-functions of  $\tilde{K}_2/K_2$ . This shows that  $K_1$  and  $K_2$  cannot be distinguished by Artin L-functions of  $\tilde{K}_1/K_1$  and  $\tilde{K}_2/K_2$ , nor by Artin L-functions of  $K_1/k$  and  $K_2/k$ .

## REFERENCES

- [1] Artin, F.: Zur Theorie der L-Reihen mit allgemeinen Gruppencharakteren, Abh. Math. Sem. Univ. Hamburg 8 (1931), no. 1, 292-306
- [2] Nicolae, F.: On Artin's L-functions. I, J. Reine Angew. Math. 539 (2001), 179-184
- [3] Nicolae, F.: On the determination of number fields by L-functions, to appear in Journal de Théorie des Nombres de Bordeaux
- [4] Perlis, R.: On the equation  $\zeta_K(s) = \zeta_{K'}(s)$ , J. Number Theory 9 (1977), 342-360.

MATHEMATISCHES INSTITUT DER UNIVERSITÄT PADERBORN, Warburger Str. 100, 33098 PADERBORN, GERMANY

*E-mail address:* `klueners@math.uni-paderborn.de`

“SIMION STOILLOW” INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY, P.O.BOX 1-764, RO-014700 BUCHAREST, ROMANIA

*E-mail address:* `florin.nicolae@imar.ro`