# BÄCKLUND TRANSFORMATIONS FOR GELFAND-DICKEY FLOWS, REVISITED

CHUU-LIAN TERNG<sup> $\dagger$ </sup> AND ZHIWEI WU<sup>\*</sup>

### 1. INTRODUCTION

The *j*-th flow of  $GD_n$ -hierarchy (cf. [5]) is the following evolution equation

$$\frac{\partial L}{\partial t_j} = \left[ (L^{\frac{j}{n}})_+, L \right] \tag{1.1}$$

on the space

$$\mathcal{D}_n = \{ L = \partial^n - \sum_{i=1}^{n-1} u_i \partial^{i-1} \, | \, u_i \in C^{\infty}(\mathbb{R}, \mathbb{C}), 1 \le i \le n-1 \},\$$

where  $(L^{\frac{j}{n}})_+$  is the differential operator component of the pseudo-differential operator  $L^{\frac{j}{n}}$ ,  $\partial = \partial_x$ , and  $j \not\equiv 0 \pmod{n}$ .

The  $A_{n-1}^{(1)}$ -KdV hierarchy constructed by Drinfeld-Sokolov in [6] is the GD<sub>n</sub>-hierarchy with matrix valued Lax pairs. The  $n \times n$  KdV hierarchy constructed by the first author and Uhlenbeck in [8] is equivalent to the GD<sub>n</sub>-hierarchy. The  $n \times n$  mKdV hierarchy given in [6] and the KW-hierarchy in [7] are equivalent. Moreover, the GD<sub>n</sub> and the  $n \times n$  mKdV hierarchies are related by the Miura transform.

Bäcklund transformations (BTs) for the KdV (i.e.,  $GD_2$ ) hierarchy constructed from a system of compatible differential equations are well-known (cf. [1], [12]).

Adler in [2] used the Miura transform to construct a BT for (1.1) as follows: Suppose  $L = \partial^n - \sum_{i=1}^{n-1} u_i \partial^{i-1}$  is a solution of the *j*-th GD<sub>n</sub> flow (1.1), and L is factored as a product of first order operators

$$L = (\partial - q_n) \cdots (\partial - q_1)$$

such that  $q = \sum_{i=1}^{n-1} q_i e_{ii}$  is a solution of the  $n \times n$  mKdV flow. Let  $\tilde{L}$  be defined by

$$\tilde{L} := (\partial - q_{n-1}) \cdots (\partial - q_1)(\partial - q_n).$$

Then  $\tilde{L}$  is a solution of (1.1). Rational solutions are obtained by applying Adler's BT to the vacuum solution  $L = \partial^n$  (cf. [2]).

Note that operators L and L in Adler's result are related by

$$\tilde{L} = (\partial - q_n)^{-1} L(\partial - q_n).$$

<sup>&</sup>lt;sup>†</sup>Research supported in part by NSF Grant DMS-1109342.

<sup>\*</sup>Research supported in part by NSF of China under Grant No. 11401327.

Given a solution L of (1.1), it is not difficult to see that  $\tilde{L} = (\partial + h)^{-1}L(\partial + h)$ is a solution of (1.1) if and only if h satisfies a system of ordinary differential equations in x and t variables. One goal of this paper is to find all solutions of this system for h. To achieve this, we need to use Bäcklund transformations for the  $n \times n$  KdV hierarchy constructed in [8] and the equivalence of the  $n \times n$  KdV and the GD<sub>n</sub> hierarchies.

 $n \times n$  KdV and the GD<sub>n</sub> hierarchies. Assume  $L = \partial^n - \sum_{i=1}^{n-1} u_i \partial^{(i-1)}$  is a solution of (1.1). We prove the following results:

• There exists a system for  $h : \mathbb{R}^2 \to \mathbb{C}$ ,

$$\begin{cases} h_x^{(n)} = r_n(u, h), \\ h_t = \eta_{n,j}(u, h), \end{cases}$$
(1.2)

for some differential polynomials  $r_n(u, h)$  and  $\eta_{n,j}(u, h)$  such that  $\tilde{L} = (\partial + h)^{-1}L(\partial + h)$  is a solution of (1.1) if and only if h is a solution of (1.2).

- There exists a differential polynomial  $\xi_n(u,h)$  such that  $r_n(u,h) = (\xi_n(u,h))_x$ .
- If h is a solution of (1.2) then there exists a constant  $k \in \mathbb{C}$  such that h satisfies

$$(BT)_{u,k} \begin{cases} h_x^{(n-1)} = \xi_n(u,h) - k, \\ h_t = \eta_{n,j}(u,h). \end{cases}$$
(1.3)

- Systems (1.2) and (1.3) are solvable for h.
- $\tilde{L} = (\partial + h)^{-1}L(\partial + h)$  is a solution of (1.1) if and only if there exists a constant  $k \in \mathbb{C}$  such that h is a solution of  $(BT)_{u,k}$  (we use h \* Lto denote  $\tilde{L}$  and call  $L \mapsto h * L$  a BT of L with parameter k).
- There exist differential polynomials  $p_{j,1}(u,\lambda), \ldots, p_{j,n}(u,\lambda)$  such that the following linear system for  $\phi : \mathbb{R}^2 \to \mathbb{C}$ ,

$$\begin{cases} L\phi = \lambda\phi, \\ \phi_t = \sum_{i=1}^n p_{j,i}(u,\lambda)\phi_x^{(i-1)}, \end{cases}$$
(1.4)

is solvable for all parameter  $\lambda \in \mathbb{C}$ , where  $p_{j,i}(u,\lambda)$  has degree  $\left[\frac{j}{n}\right]$  in  $\lambda$ . Moreover, let  $\phi_1, \ldots, \phi_{n-1}$  be linearly independent solutions of (1.4) with  $\lambda = k$ , and  $W(\phi_1, \ldots, \phi_{n-1})$  the Wronskian of  $\phi_1, \ldots, \phi_{n-1}$ . Then

$$h = (\ln W(\phi_1, \dots, \phi_{n-1}))_x$$

is a solution of  $(BT)_{u,k}$  (1.3) and  $\tilde{L} = (\partial + h)^{-1}L(\partial + h)$  is a solution of (1.1). In fact, this construction gives all solutions of (1.3).

We give algorithms to compute differential polynomials  $r_n, \eta_{n,j}, \xi_n$ , and  $p_{j,i}$ 's. For example, the second GD<sub>3</sub> flow for  $L = \partial^3 - u_2 \partial - u_1$  is (3.15), (1.2) is (3.18), (BT)\_{u,k} is (3.17), and if h is a solution of (3.17) then  $\tilde{L} = (\partial + h)^{-1}L(\partial + h) = \partial^3 - \tilde{u}_2 \partial - \tilde{u}_1$  is a new solution of (3.15), where  $\tilde{u}_i$  is given by the formula (3.19). System (1.4) is (7.5).

Note that given a solution of (1.1) and a constant k, system (1.3) gives rise to (n-1)-parameter family of new solutions of (1.1). When we apply BT with constant k = 0 and  $k \neq 0$  to the vacuum solution  $L = \partial^n$  we obtain explicit rational solutions and soliton solutions respectively.

We also obtain the following results:

- (a) We give a permutability formula for our BTs of the  $GD_n$  flows.
- (b) Adler's BT is our BT with parameter k = 0.
- (c) We construct a simple and natural action of the order n cyclic group  $\mathbb{Z}_n$  on the space of solutions of the KW-hierarchy and show that this  $\mathbb{Z}_n$ -action gives Adler's BT for the GD<sub>n</sub>-hierarchy under the Miura transform.

This paper is organized as follows: We set up notations and review the constructions and relations of various KdV-type hierarchies in section 2, construct BTs for the  $A_{n-1}^{(1)}$ -KdV hierarchy in section 3, and give a Permutability formula for these BTs and a relation between BTs and scaling transform in section 4. We use BTs to construct explicit rational and soliton solutions for the *j*-th  $A_{n-1}^{(1)}$ -KdV flow in section 5, and give a natural  $\mathbb{Z}_n$ -action on the KW-hierarchy and show that Adler's BT arises from this action in section 6. In the last section, we construct BTs for the GD<sub>n</sub>-hierarchy and explain the relation between our BTs and Adler's BTs.

#### 2. VARIOUS KDV TYPE HIERARCHIES

In this section, we set up notations and give a brief review of the constructions of the  $A_{n-1}^{(1)}$ -KdV, the  $n \times n$  mKdV, the  $n \times n$  KdV, and the KW-hierarchies.

Let  $\mathcal{B}_n^+$ ,  $\mathcal{B}_n^-$ ,  $\mathcal{T}_n$ ,  $\mathcal{N}_n^+$ , and  $\mathcal{N}_n^-$  denote the sub-algebras of upper triangular, lar, lower triangular, diagonal, strictly upper triangular, and strictly lower triangular matrices in  $sl(n, \mathbb{C})$  respectively, and  $\mathcal{N}_n^+$  the subgroup of upper triangular matrices in  $SL(n, \mathbb{C})$  with 1 on the diagonal entries. Let

$$\mathcal{L} = \left\{ \xi(\lambda) = \sum_{i \le m_0} \xi_i \lambda^i \, \middle| \, \xi_i \in sl(n, \mathbb{C}), m_0 \text{ an integer} \right\}, \\ \mathcal{L}_+ = \{\xi(\lambda) = \sum_{i \ge 0} \xi_i \lambda^i \in \mathcal{L}\}, \\ \mathcal{L}_- = \{\xi(\lambda) = \sum_{i < 0} \xi_i \lambda^i \in \mathcal{L}\}.$$

Note that  $\mathcal{L}_+, \mathcal{L}_-$  are Lie subalgebras of  $\mathcal{L}$  and  $\mathcal{L} = \mathcal{L}_+ \oplus \mathcal{L}_-$  as a direct sum of linear subspaces (such pair  $(\mathcal{L}_+, \mathcal{L}_-)$  is called a *splitting* of  $\mathcal{L}$ ). Given  $\xi = \sum_i \xi_i \lambda^i \in \mathcal{L}$ , we will use  $\xi_{\pm}$  to denote the projection of  $\xi$  onto  $\mathcal{L}_{\pm}$  w.r.t.  $\mathcal{L} = \mathcal{L}_+ \oplus \mathcal{L}_-$ , i.e.,

$$\xi_+ = \sum_{i \ge 0} \xi_i \lambda^i, \quad \xi_- = \sum_{i < 0} \xi_i \lambda^i.$$

**2.1. The**  $A_{n-1}^{(1)}$ -KdV hierarchy [6] Let

$$J = e_{1n}\lambda + b, \quad b = \sum_{i=1}^{n-1} e_{i+1,i}.$$
 (2.1)

Given  $u \in C^{\infty}(\mathbb{R}, \mathcal{B}_n^+)$ , a direct computation shows (cf. [9]) that there exists a unique  $Q(u, \lambda) \in \mathcal{L}$  such that  $Q(0, \lambda) = J$  and

$$\begin{cases} [\partial_x + J + u, Q(u, \lambda)] = 0, \\ Q(u, \lambda) \text{ is conjugate to } J. \end{cases}$$
(2.2)

Write

$$Q^{j}(u,\lambda) = \sum_{i \le \left[\frac{j}{n}\right]+1} Q_{j,i}(u)\lambda^{i}.$$
(2.3)

Let

$$V_n = \bigoplus_{i=1}^{n-1} \mathbb{C}e_{in}.$$
 (2.4)

It was proved in [6] that given  $u \in C^{\infty}(\mathbb{R}, V_n)$  and  $j \not\equiv 0 \pmod{n}$ , there exists a unique  $\mathcal{N}_n^+$ -valued differential polynomial  $\eta_j(u)$  such that

$$[\partial_x + b + u, Q_{j,0}(u) + \eta_j(u)] \in C^{\infty}(\mathbb{R}, V_n).$$

Let

$$Z_{j}(u,\lambda) = (Q^{j}(u,\lambda))_{+} + \eta_{j}(u) = \sum_{i=0}^{\left[\frac{j}{n}\right]+1} Z_{j,i}(u)\lambda^{i}.$$
 (2.5)

So we have

$$Z_{j,i}(u) = \begin{cases} Q_{j,i}(u), & i > 0, \\ Q_{j,0}(u) + \eta_j(u), & i = 0. \end{cases}$$

The *j*-th flow in the  $A_{n-1}^{(1)}$ -KdV hierarchy is the following evolution equation on  $C^{\infty}(\mathbb{R}, V_n)$ ,

$$u_{t_j} = [\partial_x + b + u, Z_{j,0}(u)], \qquad (2.6)$$

where  $b = \sum_{i=1}^{n-1} e_{i+1,i}$ . Moreover, it was proved in [6] that  $u : \mathbb{R}^2 \to V_n$  is a solution of (2.6) if and only if

$$[\partial_x + J + u, \partial_{t_j} + Z_j(u, \lambda)] = 0$$
(2.7)

for all parameter  $\lambda \in \mathbb{C}$ . We call (2.7) the Lax pair of the solution u of the j-th  $A_{n-1}^{(1)}$ -KdV flow.

4

## Definition 2.2. Frame

Let  $\mathcal{O}$  be a connected open subset of  $\mathbb{C}$ . A map  $E : \mathbb{R}^2 \times \mathcal{O} \to GL(n, \mathbb{C})$ is called a *frame* of the solution u of the *j*-th  $A_{n-1}^{(1)}$ -KdV flow if  $E(x, t, \lambda)$  is holomorphic for  $\lambda \in \mathcal{O}$  and satisfies

$$\begin{cases} E^{-1}E_x = J + u, \\ E^{-1}E_t = Z_j(u, \lambda) \end{cases}$$

where J is defined by (2.1). In other words,  $E(\cdot, \cdot, \lambda)$  is a parallel frame for the Lax pair (2.7) of u. When u is a real solution, we also require  $\overline{\mathcal{O}} = \mathcal{O}$ and E satisfies the following reality condition,

$$E(x,t,\overline{\lambda}) = E(x,t,\lambda).$$

Since  $Q(0,\lambda) = J$ ,  $Z_j(0,\lambda) = J^j$ . So u(x,t) = 0 is a solution (the vacuum solution) of (2.6) and  $E(x,t,\lambda) = \exp(xJ + tJ^j)$  is the frame of u = 0 satisfying  $E(0,0,\lambda) = I_n$  (the vacuum frame).

The following Theorem was proved by Drinfeld-Sokolov.

**Theorem 2.3.** ([6])  $u = \sum_{i=1}^{n-1} u_i e_{in}$  is a solution of the *j*-th  $A_{n-1}^{(1)}$ -KdV flow (2.6) if and only if  $L_u = \partial^n - \sum_{i=1}^{n-1} u_i \partial^{i-1}$  is a solution of the *j*-th  $GD_n$  flow.

Hence the  $GD_n$  and the  $A_{n-1}^{(1)}$ -KdV hierarchies are the same, and (2.7) is a matrix valued Lax pair for the  $GD_n$ -hierarchy.

## **2.4.** The $n \times n$ mKdV hierarchy ([6])

The Drinfeld-Sokolov  $n \times n \ mKdV$  hierarchy is constructed from the following splitting of the loop algebra  $\mathcal{L}$ ,

$$\mathcal{W}_{+} = \{\xi(\lambda) = \xi_{0} + \sum_{i>0} \xi_{i}\lambda^{i} \in \mathcal{L} \,|\, \xi_{0} \in \mathcal{B}_{n}^{-}\},$$
$$\mathcal{W}_{-} = \{\xi(\lambda) = \xi_{0} + \sum_{i<0} \xi_{i}\lambda^{i} \in \mathcal{L} \,|\, \xi_{0} \in \mathcal{N}_{n}^{+}\}.$$

Let  $\pi_{bn}$  denote the projection of  $sl(n, \mathbb{C})$  onto  $\mathcal{B}_n^-$  with respect to  $sl(n, \mathbb{C}) = \mathcal{B}_n^- \oplus \mathcal{N}_n^+$ , and let  $\pi_+$  denote the projection of  $\mathcal{L}$  onto  $\mathcal{W}_+$  with respect to the direct sum  $\mathcal{L} = \mathcal{W}_+ \oplus \mathcal{W}_-$  of linear subspaces. Then

$$\pi_+\left(\sum_{i\leq m_0}\xi_i\lambda^i\right) = \sum_{i>0}\xi_i\lambda^i + \pi_{bn}(\xi_0).$$
(2.8)

It was proved in [6] that if  $q \in C^{\infty}(\mathbb{R}, \mathcal{T}_n)$ , then  $[\partial_x + b + q, \pi_{bn}(Q_{j,0}(q))]$  lies in  $C^{\infty}(\mathbb{R}, \mathcal{T}_n)$ . The *j*-th  $n \times n$  mKdV flow is the following flow on  $C^{\infty}(\mathbb{R}, \mathcal{T}_n)$ ,

$$q_{t_j} = [\partial_x + b + q, \pi_{bn}(Q_{j,0}(q))], \qquad (2.9)$$

where  $Q_{j,0}(u)$  is defined by (2.3). Moreover,  $q : \mathbb{R}^2 \to \mathcal{T}_n$  is a solution of (2.9) if and only if

$$[\partial_x + J + q, \partial_{t_j} + \pi_+(Q^j(q,\lambda))] = 0$$

for all  $\lambda \in \mathbb{C}$ , where  $\pi_+$  is the projection of  $\mathcal{L}$  onto  $\mathcal{W}_+$  defined by (2.8) and  $Q(q, \lambda)$  is defined by (2.2).

## **2.5.** The $n \times n$ KdV hierarchy ([8], [9])

The  $n \times n$  KdV hierarchy is constructed from an unusual splitting of  $\mathcal{L}$ . Let  $\alpha = e^{\frac{2\pi i}{n}}$ , and

$$\Lambda = \sum_{k=1}^{n-1} \frac{1 - \alpha^k}{1 - \alpha} e_{k,k+1}.$$
(2.10)

Then  $\{\Lambda^i b^{n-1} \Lambda^j \mid 0 \leq i, j \leq n-1\}$  is a basis of  $sl(n, \mathbb{C})$  and  $\{\Lambda^i b^{n-1} \Lambda^j \mid i+j < n-1\}$  is a basis of  $\mathcal{N}_n^-$ . Let  $B: sl(n, \mathbb{C}) \to \mathcal{N}_n^+$  be the linear map defined by

$$B(\Lambda^i b^{n-1} \Lambda^j) = \Lambda^i b^t \Lambda^j.$$

Note that  $\operatorname{Ker}(B) = \mathcal{B}_n^+$ . It was proved in [9] that

$$\mathcal{L}^B_- = \{\xi(\lambda) = B(\xi_{-1}) + \sum_{i < 0} \xi_i \lambda^i \in \mathcal{L}\}$$

is a subalgebra of  $\mathcal{L}$ , and

$$\mathcal{L} = \mathcal{L}_+ \oplus \mathcal{L}_-^B$$

is a direct sum of linear subspaces. Let  $\pi_B$  denote the projection of  $\mathcal{L}$  onto  $\mathcal{L}_+$  with respect to the direct sum  $\mathcal{L} = \mathcal{L}_+ \oplus \mathcal{L}_-^B$ . The *j*-th  $n \times n$  KdV flow is the following evolution equation on  $C^{\infty}(\mathbb{R}, \bigoplus_{i=1}^{n-1} \mathbb{C}\Lambda^i)$ ,

$$\xi_{t_j} = [\partial_x + b + \xi, Q_{j,0}(\xi) - B(Q_{j,-1}(\xi))], \qquad (2.11)$$

where  $Q_{j,i}(\xi)$  is defined by (2.3). Moreover,  $\xi : \mathbb{R}^2 \to \bigoplus_{i=1}^n \mathbb{C}\Lambda^i$  is a solution of (2.11) if and only if

$$[\partial_x + J + \xi, \partial_{t_j} + \pi_B(Q^j(\xi, \lambda))] = 0$$
(2.12)

for all parameter  $\lambda \in \mathbb{C}$ , where  $Q(\xi, \lambda)$  is defined by (2.2).

The  $n \times n$  KdV hierarchy can be also constructed from the following equivalent splitting: Let  $f_i = (1 + \alpha + \dots + \alpha^{i-1})^{-1} \Lambda^i$ , where  $\alpha = e^{\frac{2\pi i}{n}}$  and  $\Lambda$  is as in (2.10). Let  $\phi_n(z) = I + \sum_{i=1}^{n-1} f_i z^i$ ,  $\lambda = z^n$ , and  $T : \mathcal{L} \to \mathcal{L}$  the algebra homomorphism defined by:

$$T(A(z)) = \phi_n(z)A(z^n)\phi_n(z)^{-1}.$$
(2.13)

Let  $\mathcal{L}_T = T(\mathcal{L})$ . It was proved in [9] that

$$(\mathcal{L}_T)_+ := T(\mathcal{L}_+) = \mathcal{L}_T \cap \mathcal{L}_+, \quad (\mathcal{L}_T)_- := T(\mathcal{L}_-^B) = \mathcal{L}_T \cap \mathcal{L}_-,$$

and

$$T(e_{1n}z^n + b) = az + b,$$

where  $a = \text{diag}(1, \alpha, \dots, \alpha^{n-1})$ . Hence the Lax pair (2.12) becomes

$$[\partial_x + az + b + \xi, \partial_{t_j} + (\tilde{Q}^j(\xi, z))_+] = 0, \qquad (2.14)$$

where  $\tilde{Q}(\xi, z) = \phi_n(z)Q(\xi, z^n)\phi_n(z)^{-1}$ . We call (2.12) and (2.14) the Lax pair for  $\xi$  in  $\lambda$ -guage and z-gauge respectively, where  $\lambda = z^n$ .

We need the following Proposition (proved in [6]) to state the equivalence of the  $n \times n$  KdV and the  $A_{n-1}^{(1)}$ -KdV hierarchies.

**Proposition 2.6.** ([6]) Given  $v \in C^{\infty}(\mathbb{R}, \mathcal{B}_n^+)$ , then there exist a unique  $N_n^+$ -valued differential polynomial  $\triangle$  and  $V_n$ -valued differential polynomial u in v such that

$$\triangle(\partial_x + J + v)\triangle^{-1} = \partial_x + J + u, \qquad (2.15)$$

where  $V_n$  is defined by (2.4).

**Definition 2.7.** Let  $\Psi : C^{\infty}(\mathbb{R}, \mathcal{B}_n^+) \to C^{\infty}(\mathbb{R}, V_n)$  and  $\Gamma : C^{\infty}(\mathbb{R}, \mathcal{B}_n^+) \to C^{\infty}(\mathbb{R}, \mathcal{B}_n^+)$  $C^{\infty}(\mathbb{R}, N_n^+)$  be the maps defined by

$$\Psi(v) = u, \quad \Gamma(v) = \triangle$$

respectively, where v, u, and  $\triangle$  are related by (2.15).

Let

$$\mathcal{G}_j = \begin{cases} \bigoplus_{i=1}^{n-j} \mathbb{C}e_{i,i+j}, & j \ge 0, \\ \bigoplus_{i=1-j}^n \mathbb{C}e_{i,i+j}, & j < 0. \end{cases}$$

**Theorem 2.8.** ([10]) Let  $\xi$  be a solution of the *j*-th  $n \times n$  KdV flow (2.11),  $u(\cdot,t) = \Psi(\xi(\cdot,t)), \text{ and } \Delta(\cdot,t) = \Gamma(\xi(\cdot,t)), \text{ where } \Psi \text{ and } \Gamma \text{ are the maps}$ defined in Definition 2.7. Then we have the following results:

- (a) u is a solution of the j-th A<sup>(1)</sup><sub>n-1</sub>-KdV flow (2.6).
  (b) The map Ψ maps the space of solutions of (2.11) bijectively to the space of solutions of (2.6).
- (c) Write  $\triangle = I_n + \sum_{i=1}^{n-1} \triangle_i$  with  $\triangle_i \in \mathcal{G}_i$ , then  $\triangle_1 = 0$ . (d) If F is a frame of a solution  $\xi$  of (2.11), then  $E = F \triangle^{-1}$  is a frame of the solution  $u(\cdot, t)$  of the *j*-th  $A_{n-1}^{(1)}$ -KdV flow (2.6).

## **2.9. The KW-hierarchy** ([7])

Let

$$\tau = e_{21} + e_{32} + \dots + e_{n,n-1} + e_{1n},$$
  
$$\alpha = e^{\frac{2\pi i}{n}},$$
  
$$a = \operatorname{diag}(1, \alpha, \dots, \alpha^{n-1}).$$

The KW-hierarchy is obtained from the splitting  $\mathcal{L}^{KW}_{\pm}$  of the algebra  $\mathcal{L}^{KW}$ , where

$$\mathcal{L}^{KW} = \{ A \in \mathcal{L} \, | \, A(\alpha^{-1}z) = \tau A(z)\tau^{-1} \}, \mathcal{L}^{KW}_{\pm} = \mathcal{L}^{KW} \cap \mathcal{L}_{\pm}.$$

Given  $v = (v_1, \dots, v_{n-1})$ , let

$$P(v) = \sum_{i=1}^{n-1} v_i \tau^i.$$
 (2.16)

Let  $\hat{Q}(v,z) = \sum_{i \leq 1} \hat{Q}_{1,i}(v) z^i$  be the unique solution of

$$[\partial_x + az + P(v), \hat{Q}(v, z)] = 0$$

satisfying  $\hat{Q}(0,z) = az$  and  $\hat{Q}(v,z)$  is conjugate to az. Write

$$\hat{Q}^j(v,z) = \sum_{i \le j} \hat{Q}_{j,i}(v) z^i.$$

The *j*-th flow in the KW-hierarchy is the following evolution equation on  $C^{\infty}(\mathbb{R}, \bigoplus_{i=1}^{n-1} \mathbb{C}\tau^i),$ 

$$\frac{\partial P(v)}{\partial t_j} = \sum_{i=1}^{n-1} (v_i)_{t_j} \tau^i = [\partial_x + P(v), \hat{Q}_{j,0}(v)].$$
(2.17)

Moreover, v is a solution of (2.17) if and only if

$$[\partial_x + az + P(v), \partial_{t_j} + \left(\hat{Q}^j(v, z)\right)_+] = 0.$$

Next we review relations between the KW, the  $n \times n$  mKdV, and the  $A_{n-1}^{(1)}$ -KdV hierarchies.

**Theorem 2.10.** ([10]) Let  $\Phi : \mathbb{R}^{1 \times (n-1)} \to \mathcal{T}_n$  be the linear isomorphism defined by  $\Phi(v_1, \ldots, v_{n-1}) = q = \operatorname{diag}(q_1, \ldots, q_n)$ , where  $q_i = \sum_{k=1}^{n-1} \alpha^{k(i-1)} v_k$  for  $1 \leq i \leq n$  and  $\alpha = \exp(2\pi i/n)$ . Then we have the following.

- (i) v is a solution of the j-th KW flow (2.17) if and only if  $q = \Phi(v)$  is a solution of the j-th  $n \times n$  mKdV flow (2.9).
- (ii) Let  $V(z) = ((\alpha^{(i-1)}z)^{j-1}), \lambda = z^n$  and  $J = e_{1n}\lambda + b$  as in (2.1). Then

$$V(z)^{-1}(\partial_x + az + P(v))V(z) = \partial_x + J + q,$$

where  $P(v) = \sum_{i=1}^{n-1} v_i \tau^i$  is as defined by (2.16).

**Theorem 2.11.** ([6]) Given  $q = \sum_{i=1}^{n} q_i e_{ii} : \mathbb{R}^2 \to sl(n, \mathbb{C})$ , let  $u_1, \ldots, u_{n-1}$  be defined by

$$(\partial - q_n) \cdots (\partial - q_1) = \partial^{(n)} - \sum_{i=1}^{n-1} u_i \partial^{i-1}$$

Then we have the following.

- (i) if q is a solution of the j-th  $n \times n$  mKdV- flow (2.9), then  $u = \sum_{i=1}^{n-1} u_i e_{in}$  is a solution of the j-th  $A_{n-1}^{(1)}$ -KdV flow (2.6) (this is the Miura transform).
- (ii) Let  $\Gamma$  and  $\Psi$  be the maps defined in Definition 2.7,  $\triangle(\cdot, t) = \Gamma(q(\cdot, t))$ , and K a frame of the solution q of (2.9). Then  $u(\cdot, t) = \Psi(q(\cdot, t))$ and  $E = K \triangle^{-1}$  is a frame of the solution u of (2.6).

#### BT FOR $GD_n$ FLOW

# 3. BÄCKLUND TRANSFORMATION FOR $A_{n-1}^{(1)}$ -KdV

In this section, we prove the following results:

- (i) Assume that E is a frame of a solution u of the j-th  $A_{n-1}^{(1)}$ -KdV flow (2.6), and f is of the form  $J + hI_n + N$  for some complex valued function h and an  $\mathcal{N}_n^+$ -valued map N such that  $\tilde{E} = Ef^{-1}$  is also a frame of a solution of (2.6). Then the entries of N are differential polynomials of u and h, and h satisfies the system (1.2).
- (ii) If u is a solution of (2.6), then system (1.2) is solvable and a solution h gives rise to a new solution of (2.6).
- (iii) All solutions of (1.2) can be constructed from frames of the Lax pair of a solution u of (2.6).
- (iv) h is a solution of (1.2) if and only if there exists a constant  $k \in \mathbb{C}$ such that h is a solution of (1.3).

Bäcklund transformations for the  $n \times n$  KdV hierarchy is constructed in [8] from a loop group factorization with respect to the splitting in z-gauge. We use the map T defined by (2.13) to state this result in  $\lambda$ -gauge as follows:

**Theorem 3.1.** ([8]) Let  $j \not\equiv 0 \pmod{n}$ , and  $\Lambda$  defined by (2.10). Then there exist differential polynomials  $A_i(\xi, Y)$  and  $B_i(\xi, Y)$  such that if F is a frame of a solution  $\xi$  of the *j*-th  $n \times n$  KdV flow (2.11) then  $\tilde{F} = F(J+Y)^{-1}$ is a frame of a solution solution  $\tilde{\xi}$  of (2.11) if and only if Y satisfies the following first order system,

$$\begin{cases} Y_x = A_j(\xi, Y), \\ Y_t = B_j(\xi, Y), \end{cases}$$
(3.1)

where  $J = e_{1n}\lambda + b$  is as in (2.1) and  $Y = \sum_{i=0}^{n-1} y_i \Lambda^i$ . Moreover, if  $\xi$  is a solution of (2.11), then we have the following:

- (i) System (3.1) is solvable for  $Y \in C^{\infty}(\mathbb{R}, \bigoplus_{i=0}^{n-1} \mathbb{C}\Lambda^i)$ .
- (ii) If Y is a solution of (3.1), then there exists a constant  $k \in \mathbb{C}$  such that  $\det(J+Y) = (-1)^{n-1}(\lambda - k).$
- (iii) Given constants  $k \in \mathbb{C}$  and  $\mathbf{c}_0 \in \mathbb{C}^n \setminus 0$ , let

$$\zeta(x,t) = (\zeta_1, \cdots, \zeta_n)^t = F(x,t,k)^{-1}(\mathbf{c}_0),$$

Then there is a unique solution  $Y = \sum_{i=0}^{n-1} y_i \Lambda^i$  of (3.1) satisfying  $y_0 = -\frac{\zeta_{n-1}}{\zeta_n}$  and  $(ke_{1n} + b + Y)\zeta = 0$ . (iv) All solutions of (3.1) can be constructed as in (iii).

Note that if  $\xi$  is a solution of (2.11) and Y is a solution of (3.1), then

$$\tilde{\xi} = (J+Y)(J+\xi)(J+Y)^{-1} - Y_x(J+Y)^{-1} - J$$

is again a solution of (2.11). By Theorems 2.8 and 3.1, we obtain the following.

**Theorem 3.2.** Let  $\xi$  be a solution of the *j*-th  $n \times n$ -KdV flow (2.11),  $Y = \sum_{i=0}^{n-1} y_i \Lambda^i$  a solution of (3.1), and  $\tilde{\xi}$  the solution of (2.11) constructed from  $\xi$  and Y as in Theorem 3.1. Let  $u(\cdot,t) = \Psi(\xi(\cdot,t))$ ,  $\tilde{u} = \Psi(\tilde{\xi}(\cdot,t))$ ,  $\Delta(\cdot,t) = \Gamma(\xi(\cdot,t))$ , and  $\tilde{\Delta}(\cdot,t) = \Gamma(\tilde{\xi}(\cdot,t))$ , where  $\Psi$  and  $\Gamma$  are defined in Definition 2.7. Set

$$f = \tilde{\bigtriangleup}(J+Y)\bigtriangleup^{-1}.$$

Then we have the following:

- (a) Both u and  $\tilde{u}$  are solutions of the *j*-th  $A_{n-1}^{(1)}$ -KdV flow (2.6).
- (b) If F is a frame of  $\xi$ , then  $E = F \triangle^{-1}$  and  $\tilde{E} = Ef^{-1}$  are frame of solutions u and  $\tilde{u}$  of (2.6) respectively.
- (c) f is of the form  $J + y_0 I_n + N$ , where N is an  $\mathcal{N}_n^+$ -valued map.

*Proof.* From Theorem 2.8, both u and  $\tilde{u}$  are solutions of (2.6),  $E = F \triangle^{-1}$  and  $\tilde{E} = \tilde{F} \tilde{\triangle}^{-1}$  are frames of u and  $\tilde{u}$  respectively. It follows from a direct computation that

$$\tilde{F}\tilde{\bigtriangleup}^{-1} = F(J+Y)^{-1}\tilde{\bigtriangleup}^{-1} = E\bigtriangleup(J+Y)^{-1}\tilde{\bigtriangleup}^{-1} = Ef^{-1}.$$

This proves (a) and (b).

Let  $\mathcal{G}_i = \bigoplus_{j=1}^{n-i} \mathbb{C}e_{j,i+j}$ . Write  $\triangle = \sum_{i=0}^{n-1} \triangle_i$  and  $\tilde{\triangle} = \sum_{i=0}^{n-1} \tilde{\triangle}_i$  with  $\triangle_i$ ,  $\tilde{\triangle}_i \in \mathcal{G}_i$ . By Theorem 2.8,  $\triangle_1 = \tilde{\triangle}_1 = 0$ . So  $f = J + y_0 \mathbf{I}_n + N$  for some  $N(x,t) \in \mathcal{N}_n^+$ .

Next we prove that entries of N in Theorem 3.2 are differential polynomials of u and h. First we need a Lemma, which can be proved by a direct computation.

**Lemma 3.3.** Let  $u = \sum_{i=1}^{n-1} u_i e_{in}$ ,  $E \in C^{\infty}(\mathbb{R}, GL(n, \mathbb{C}))$  and  $\phi$  the first column of E. Then  $E_x = E(J+u)$  if and only if  $E = (\phi, \phi_x, \dots, \phi_x^{(n-1)})$  and  $\phi_x^{(n)} = \sum_{i=1}^{n-1} u_i \phi_x^{(i-1)} + \lambda \phi$ , where  $J = e_{1n}\lambda + b$  is defined by (2.1).

We say a differential polynomial  $\eta$  has order k in h if  $\eta$  is a polynomial in  $h, h_x, \ldots, h_x^{(k)}$ .

**Theorem 3.4.** Let E and  $\tilde{E}$  be frames of solutions  $u = \sum_{i=1}^{n-1} u_i e_{in}$  and  $\tilde{u} = \sum_{i=1}^{n-1} \tilde{u}_i e_{in}$  of (2.6) respectively, and  $\phi$ ,  $\tilde{\phi}$  the first column of E and  $\tilde{E}$  respectively. Suppose there exists h(x,t) such that  $\phi = (\partial_x + h)\tilde{\phi}$ . Then  $E(x,\lambda) = \tilde{E}(x,\lambda)f$ , where f is of the form  $J + hI_n + N$ , for some  $N(x,t) \in \mathcal{N}_n^+$ . Moreover,

(i) there are differential polynomial  $s_i(u, h)$  of order (n - i) in h such that

$$\tilde{u}_i = u_i + s_i(u, h), \quad 1 \le i \le n - 1,$$
(3.2)

(ii) the ij-th entry of N is

$$\begin{cases} N_{ij} = C_{j-1,i-1} h_x^{(j-i)}, & 1 \le i < j < n, \\ N_{in} = u_i + s_i(u,h) + C_{n-1,i-1} h_x^{(n-i)}, & 1 \le i \le n-1, \end{cases}$$
(3.3)

where  $C_{j,i} = \frac{j!}{i!(j-i)!}$ . (We will use  $f_{u,h}$  to denote such f), (iii) h satisfies

$$\begin{cases} h_x^{(n)} = r_n(u, h), \\ h_t = \eta_{n,j}(u, h), \end{cases}$$
(3.4)

for some differential polynomials  $r_n(u,h)$  of order (n-1) in h and  $\eta_{n,j}(u,h)$  of order j in h.

*Proof.* From  $E^{-1}E_x = J + u$ ,  $\tilde{E}^{-1}\tilde{E}_x = J + \tilde{u}$ , and  $E = \tilde{E}f$ , we have

$$f_x = f(J+u) - (J+\tilde{u})f.$$
 (3.5)

Compare the *nn*-th entry of (3.5) to get  $u_{n-1} = h_x + \tilde{u}_{n-1} + (n-1)h_x$ , i.e.,

$$\tilde{u}_{n-1} = u_{n-1} - nh_x. \tag{3.6}$$

Compare the (n-1, n)-entry of (3.5) to get

$$\tilde{u}_{n-2} = u_{n-2} - (u_{n-1})_x - \frac{(n-3)n}{2}h_{xx} + nhh_x.$$
(3.7)

Use induction and compare the (n-i, n)-th entry of (3.5) for  $1 \le i \le n-1$  to see that  $\tilde{u}_{n-i} = u_{n-i} + s_{n-i}(u, h)$  for some differential polynomial  $s_{n-i}(u, h)$  of order i in h. This proves (i).

Let  $f = \tilde{E}^{-1}E$ . By assumption, the first column of f is  $(h, 1, 0, ..., 0)^t$ . Lemma 3.3 implies that  $\tilde{\phi}_x^{(n)} = \lambda \tilde{\phi} + \sum_{i=1}^{n-1} \tilde{u}_i \tilde{\phi}_x^{(i-1)}$ . Use  $\phi = \tilde{\phi}_x + h \tilde{\phi}$  to compute  $\phi_x^{(i)}$  and use (3.2) to get (3.3). This proves (ii).

Compare the 1*n*-th entry of the constant term of (3.5) (as a polynomial in  $\lambda$ ) to get  $h_x^{(n)} = r_n(u, h)$  for some differential polynomial  $r_n(u, h)$  of order (n-1) in h. Since E and  $\tilde{E}$  are frames of u and  $\tilde{u}$ , we have  $E^{-1}E_t = Z_j(u, \lambda)$ and  $\tilde{E}^{-1}\tilde{E}_t = Z_j(\tilde{u}, \lambda)$ , where  $Z_j$  is as in (2.5). But  $E = \tilde{E}f$  implies that

$$f_t = fZ_j(u,\lambda) - Z_j(\tilde{u},\lambda)f.$$
(3.8)

Compare the 11-th entry of the constant term of (3.8) to see that  $h_t = \eta_{n,j}(u,h)$  for some differential polynomial  $\eta_{n,j}(u,h)$  of order j. This proves (iii).

The proof of Theorem 3.4 gives the following Proposition in x-variable:

**Proposition 3.5.** Let  $u = \sum_{i=1}^{n-1} u_i e_{in}$ ,  $\tilde{u} = \sum_{i=1}^{n-1} \tilde{u}_i e_{in} \in C^{\infty}(\mathbb{R}, V_n)$ ,  $E, \tilde{E} \in C^{\infty}(\mathbb{R}, GL(n, \mathbb{C}))$  satisfying  $E^{-1}E_x = J + u$  and  $\tilde{E}^{-1}\tilde{E}_x = J + \tilde{u}$ . Suppose  $E = \tilde{E}f$  for some f of the form  $J + hI_n + N$ , where  $h \in C^{\infty}(\mathbb{R}, \mathbb{C})$ and  $N \in C^{\infty}(\mathbb{R}, \mathcal{N}_n^+)$ . Then  $f = f_{u,h}$ ,  $\tilde{u}_i = u_i + s_i(u,h)$  for  $1 \leq i \leq n-1$ , and  $h_x^{(n)} = r_n(u,h)$  as in Theorem 3.4.

**Proposition 3.6.** If  $g(x) \in GL(n, \mathbb{C})$  and  $tr(g^{-1}g_x) = 0$ , then det(g(x)) is a constant.

*Proof.* Since 
$$\frac{d}{dx} \ln(\det(g)) = \operatorname{tr}(g^{-1}g_x), \det(g)$$
 is constant.

**Proposition 3.7.** Let  $u, \tilde{u}$  and  $E, \tilde{E}$  be as in Proposition 3.5. Then there exists a differential polynomial  $\xi_n(u,h)$  of order (n-2) in h such that

$$\det(f_{u,h}(x,\lambda)) = (-1)^{n-1} (\lambda + h_x^{(n-1)} - \xi_n(u,h)),$$
(3.9)

$$h_x^{(n)} - r_n(u,h) = (h_x^{(n-1)} - \xi_n(u,h))_x, \qquad (3.10)$$

where  $f_{u,h}$  and  $r_n(u,h)$  are as in Theorem 3.4.

*Proof.* Use (3.3) and a direct computation to get (3.9). Let  $f_{u,h} = f$ . The proof of Theorem 3.4 implies that

$$f_x - f(J+u) + (J+\tilde{u})f = (h_x^{(n)} - r_n(u,h))e_{1n}.$$
 (3.11)

Let  $w = h_x^{(n)} - r_n(u, h)$ . By (3.11), we get

$$(\ln(\det f))_x = \operatorname{tr}(f^{-1}f_x) = \operatorname{tr}((J+u) - f^{-1}(J+\tilde{u})f + f^{-1}we_{1n}).$$

Since  $\operatorname{tr}(J+u) = \operatorname{tr}(J+\tilde{u}) = 0$ ,

$$(\ln(\det f))_x = \operatorname{tr}(f^{-1}we_{1n}) = wf^{n1}$$

where  $f^{n1}$  is the (n1)-th entry of the inverse of f. By definition of f, we see that  $f^{n1}$  is equal to  $(-1)^{n-1}(\det f)^{-1}$ . This proves (3.10).

**Theorem 3.8.** Suppose  $E, \tilde{E}$  are frames of solutions  $u, \tilde{u}$  of the *j*-th  $A_{n-1}^{(1)}$ -KdV flow (2.6) respectively, and  $\tilde{E} = Ef_{u,h}^{-1}$  for some smooth function h. Then there exists a constant  $k \in \mathbb{C}$  such that  $\det(f_{u,h}(x,t,k)) = 0$  and h satisfies

$$(BT)_{u,k} \begin{cases} h_x^{(n-1)} = \xi_n(u,h) - k, \\ h_t = \eta_{n,j}(u,h), \end{cases}$$
(3.12)

where  $\xi_n(u,h)$  is as in Proposition 3.7 and  $\eta_{n,j}(u,h)$  is as in Theorem 3.4.

Proof. Recall that  $\operatorname{tr}(J+u) = \operatorname{tr}(J+\tilde{u}) = \operatorname{tr}(Z_j(u,\lambda)) = \operatorname{tr}(Z_j(\tilde{u},\lambda)) = 0$ . By Proposition 3.6, determinants of both  $E(x,t,\lambda)$  and  $\tilde{E}(x,t,\lambda)$  are independent of x and t. Hence  $\operatorname{det}(f_{u,h})$  is independent of x, t. But  $\operatorname{det}(f_{u,h})$  is a degree one polynomial in  $\lambda$ . So there exists a constant  $k \in \mathbb{C}$  such that  $\operatorname{det}(f_{u,h}(x,t,\lambda)) = (-1)^{n-1}(\lambda-k)$ . By (3.9), we have  $\operatorname{det}(f_{u,h}) = (-1)^{n-1}(\lambda+h_x^{(n-1)}-\xi_n(u,h))$ . Hence

$$h_x^{(n-1)} - \xi_n(u,h) = -k.$$

It follows from Theorem 3.4 that h satisfies the second equation of (3.12).  $\Box$ 

The following Theorem gives a method to construct solutions of  $(BT)_{u,k}$  from frames of u.

**Theorem 3.9.** Let E be a frame of a solution  $u = \sum_{i=1}^{n-1} u_i e_{in}$  of the *j*-th  $A_{n-1}^{(1)}$ -KdV flow (2.6),  $k \in \mathbb{C}$ ,  $\mathbf{c}_0 \in \mathbb{C}^n \setminus 0$  constants, and  $v(x,t) = (v_1, \cdots, v_n)^t(x,t) := E(x,t,k)^{-1}(\mathbf{c}_0)$ . If  $v_n \neq 0$ , then (i)  $h = -\frac{v_{n-1}}{v_n}$  is a solution of  $(\mathrm{BT})_{u,k}$ , BT FOR  $GD_n$  FLOW

- (ii)  $\tilde{E} = E f_{u,h}^{-1}$  is a frame of a new solution  $\tilde{u} = \sum_{i=1}^{n-1} \tilde{u}_i e_{in}$  of (2.6), where  $\tilde{u}_i$ 's are given by (3.2),
- (iii)  $f_{u,h}(x,t,k)(v(x,t)) = 0$  for all x, t.

*Proof.* By Theorem 2.8, there exist a unique solution  $\xi$  of the *j*-th  $n \times n$ KdV flow (2.11) such that  $\Psi(\xi(\cdot,t)) = u(\cdot,t)$  and  $\Delta(\cdot,t) = \Gamma(\xi(\cdot,t)) \in N_n^+$ , where  $\Psi$  and  $\Gamma$  are as in Definition 2.7. Then  $F = E \triangle$  is a frame for  $\xi$ .

Let  $\zeta = (\zeta_1, \cdots, \zeta_n)^t = \triangle^{-1} v$ , so  $\zeta = F(x, t, k)^{-1}(\mathbf{c}_0)$ . Let  $Y(\zeta)$  be as in Theorem 3.1 (iii). Then  $\tilde{F} = F(J+Y)^{-1}$  is a frame of a new solution  $\tilde{\xi}$  of (2.11),  $y_0 = -\frac{\zeta_{n-1}}{\zeta_n}$ , and  $(ke_{1n} + b + Y)\zeta = 0$ . Hence  $\det(J+Y)(x,t,k) = 0$ . By Theorem 3.2,  $\tilde{u} = \Psi(\tilde{\xi})$  is a solution of (2.6) and  $\tilde{E} = Ef^{-1}$  is a frame of  $\tilde{u}$ , where  $f = \tilde{\Delta}(J+Y)\Delta^{-1}$ . By Theorem 3.4,  $f = f_{u,y_0}$ . Therefore,  $\det(f_{u,y_0})(x,t,k) = 0. \text{ By Theorem (3.8), } y_0 \text{ is a solution of (3.12). Write} \\ \triangle = I_n + \sum_{i=1}^{n-1} \triangle_i \text{ and } \tilde{\triangle} = I_n + \sum_{i=1}^{n-1} \tilde{\triangle}_i \text{ with } \triangle_i \in \mathcal{G}_i = \bigoplus_{j=1}^{n-i} \mathbb{C}e_{j,j+i}.$ By Theorem 2.8 (b), we have  $\Delta_1 = \tilde{\Delta}_1 = 0$ . It follows from a direct computation that  $y_0 = -\frac{\zeta_{n-1}}{\zeta_n} = -\frac{v_{n-1}}{v_n} = h$ . So we have proved (i) and (ii). From  $(ke_{1n}+b+Y)(\zeta) = 0, \zeta = \Delta^{-1}v$  and  $f_{u,h} = \tilde{\Delta}(J+Y)\Delta^{-1}$ , we have

 $f_{u,h}(x,t,k)v(x,t,k) = 0.$ 

**Theorem 3.10.** Let u be a solution of (2.6), and  $k \in \mathbb{C}$  a constant. Then

- (i) all solutions of  $(BT)_{u,k}$  are obtained by the method given in Theorem 3.9.
- (ii)  $(BT)_{u,k}$ , *i.e.*, (3.12) with parameter k, is solvable,

*Proof.* Let E be the frame of a solution u of (2.6) such that  $E(0,0,\lambda) = I_n$  for all  $\lambda \in \mathbb{C}$ . Let  $\mathbf{c} = (c_1, \dots, c_{n-1}, 1)$  and  $v = (v_1, \dots, v_n) := E^{-1}(\cdot, \cdot, k)(\mathbf{c})$ . Then  $v_n \neq 0$  in an open neighborhood of (0,0) in  $\mathbb{R}^2$ . It also follows from  $E^{-1}E_x = J + u$  that

$$v_x = -(e_{1n}k + b + u)v. (3.13)$$

Let  $h = -v_{n-1}/v_n$ , and  $\Phi : \mathbb{C}^{n-1} \to \mathbb{C}^{n-1}$  the map defined by

$$\Phi(\mathbf{c}) = (h(0,0), h_x(0,0), \cdots, h_x^{(n-2)}(0,0)).$$

We claim that  $\Phi$  is onto. Let  $\phi = -\frac{1}{v_n}v$ , then  $h = \phi_{n-1}$ . Eq.(3.13) implies that

$$\begin{cases} \phi_x = -(ke_{1n} + b + u + hI_n)\phi, \\ \phi(0,0) = -\mathbf{c}. \end{cases}$$

Compare the entries of both sides, we have

$$\begin{aligned} (\phi_1)_x &= -\phi_1 \phi_{n-1} + k + u_1, \\ (\phi_i)_x &= -\phi_{i-1} - \phi_i \phi_{n-1} + u_i, \quad 2 \le i \le n-1. \end{aligned}$$

This shows that given any  $(c_1, \ldots, c_{n-1})$  there exists a solution h of (3.12) satisfying

$$h(0,0) = -c_{n-1}, \quad h_x(0,0) = c_{n-2} + u_{n-1}(0,0) - c_{n-1}^2, \quad \cdots$$

This proves (i). Since we have constructed solutions with all initial data, system (3.12) with parameter k is solvable.

**Corollary 3.11.** If  $u = \sum_{i=1}^{n-1} u_i e_{in}$  is a solution of (2.6) and h is a solution of (BT)<sub>u,k</sub> (3.12), then  $\tilde{u} = \sum_{i=1}^{n-1} \tilde{u}_i e_{in}$  defined by  $\tilde{u}_i = u_i + s_i(u, h)$  as in (3.2) is a solution of (2.6). Moreover, if E is a frame of u, then  $Ef_{u,h}^{-1}$  is a frame of  $\tilde{u}$  for  $\lambda \neq k$  and det $(f_{u,h}(x,t,\lambda)) = (-1)^{n-1}(\lambda - k)$ .

**Remark 3.12.** Given a solution u of (2.6), we can construct new solutions of (2.6) either by solving the compatible non-linear system  $(BT)_{u,k}$ , i.e., (3.12), for h or solve the frame E(x, t, k) by solving a linear system. When n = 2 this is the classical relation between two component linear systems and systems of Riccati equations.

It follows from (3.10) that if h is a solution of (3.4) then  $h_x^{(n-1)} - \xi_n(u, h) = k(t)$  for some function of t. We need the following lemma to prove that k(t) is a constant.

**Lemma 3.13.** Let  $K : \bigoplus_{i=0}^{n-1} \mathbb{C}\Lambda^i \to \mathbb{C}^n$  be the map defined by

$$K(\mathbf{c}) = (y_0(0,0), (y_0)_x(0,0), \cdots, (y_0)_x^{(n-1)}(0,0))^t,$$

where  $Y = \sum_{i=0}^{n-1} y_i \Lambda^i$  is the solution of (3.1) with initial data  $Y(0,0) = \mathbf{c}$ . Then K is onto.

*Proof.* Let F be a frame of  $\xi = \sum_{i=1}^{n-1} \xi_i \Lambda^i$  of the *j*-th  $n \times n$  KdV flow (2.11). By Theorem 3.1,  $\tilde{F} = F(J+Y)^{-1}$  is a frame for a new solution  $\tilde{\xi} = \sum_{i=1}^{n-1} \tilde{\xi}_i \Lambda^i$ . Hence we have

$$(J+Y)_x = (J+Y)(J+\xi) - (J+\xi)(J+Y).$$

The coefficients of  $\lambda$  of the both sides are zero. So the above equality is equivalent to

$$(b+Y)_x = (b+Y)(b+\xi) - (b+\xi)(b+Y).$$
(3.14)

Compare the (n, n)-th entry of both sides to get

$$y_1 = \alpha(y_0)_x + \xi_1$$
, where  $\alpha = e^{\frac{2\pi i}{n}}$ .

Let  $\mathcal{G}_i = \bigoplus_{j=1}^{n-i} \mathbb{C}e_{i,j+i}$ . Compare the  $\mathcal{G}_i$ -components of both sides of (3.14) for  $1 \leq i \leq n-2$ , and use the fact that

$$\Lambda^{k}b - b\Lambda^{k} = (\sum_{i=1}^{k-1} \alpha^{i})a\Lambda^{k-1}, \quad a = \operatorname{diag}(1, \alpha, \cdots, \alpha^{n-1}), \quad \alpha = e^{\frac{2\pi i}{n}}.$$

14

We obtain

$$(y_i)_x \Lambda^i = (y_{i+1} - \xi_{i+1}) \left( \sum_{j=1}^i \alpha^j \right) a \Lambda^i + (\xi_{i+1} - \tilde{\xi}_{i+1}) \Lambda^{i+1} b + \left( \sum_{k=0}^i y_k (\xi_{i-k} - \tilde{\xi}_{i-k}) \right) \Lambda^i, \quad (\xi_0 = \tilde{\xi}_0 = 0).$$

Compare the (n - i, n)-th entry of both sides and by induction to get

$$y_{i+1} = \frac{\alpha^{i+1}}{\sum_{j=1}^{i} \alpha^j} (y_i)_x + p_i(y_0, \cdots, y_{i-1}, \xi), \quad 1 \le j \le n-2,$$

where  $p_i$ 's are polynomials in  $y_0, \dots, y_{i-1}$  and  $\xi$ . Therefore, given  $(y_0)_x^{(i)}(0,0)$  for  $0 \le i \le n-1$ , we can write down  $y_i(0,0), 0 \le i \le n-1$  uniquely. So K is onto.

**Theorem 3.14.** Let u be a solution of the j-th  $A_{n-1}^{(1)}$ -KdV flow (2.6), and h a solution of (3.4). Let  $\tilde{u}_i = u_i + s_i(u,h)$  as in (3.2), and  $\tilde{u} = \sum_{i=1}^{n-1} \tilde{u}_i e_{in}$ . Then  $\tilde{u}$  is a solution of (2.6) and  $\tilde{E} = Ef_{u,h}^{-1}$  is a frame of  $\tilde{u}$ .

*Proof.* By Theorem 2.8, there exist a unique solution  $\xi$  of the *j*-th  $n \times n$  KdV flow (2.11) such that  $\Psi(\xi(\cdot,t)) = u(\cdot,t)$  and  $\triangle(\cdot,t) = \Gamma(\xi(\cdot,t)) \in N_n^+$  and  $F = E \triangle$  is a frame for  $\xi$ .

By Lemma 3.13, there exists  $\mathbf{c} \in \oplus_{i=0}^{n-1} \mathbb{C} \Lambda^i$  such that

$$K(\mathbf{c}) = (h(0,0), h_x(0,0), \dots, h_x^{(n-1)}(0,0))^t.$$

Let  $Y = \sum_{i=0}^{n-1} y_i \Lambda^i$  be the solution of (3.1) with initial data

$$\sum_{i=0}^{n-1} y_i(0,0)\Lambda^i = \mathbf{c}.$$

By Theorem 3.1,  $\hat{F} = F(J+Y)^{-1}$  is a frame of the new solution  $\hat{\xi}$  of (2.11). Let  $\hat{u}(\cdot,t) = \Psi(\hat{\xi}(\cdot,t))$ , and  $\hat{\bigtriangleup}(\cdot,t) = \Gamma(\hat{\xi}(\cdot,t))$ . By Theorem 2.8 again,  $\hat{u}$  is a solution of (2.6) and  $\hat{E} = \hat{F}\hat{\bigtriangleup}^{-1}$  is a frame of  $\hat{u}$ . Since  $F = E\hat{\bigtriangleup}$ , we see that

$$\hat{E} = E \triangle (J+Y)^{-1} \hat{\triangle}^{-1} = Eg^{-1},$$

where  $g = \hat{\Delta}(J+Y)\Delta^{-1}$ . By Theorem 3.2, g is of the form  $J + y_0\mathbf{I}_n + N$ for some  $\mathcal{N}_n^+$ -valued map. From Theorem 3.4,  $g = f_{u,y_0}$ . This shows that  $\hat{E}, E$  are frames of solutions  $u, \hat{u}$  of (2.6) and  $\hat{E} = Ef_{u,y_0}^{-1}$ . By Theorem 3.4,  $y_0$  satisfies (3.4). By assumption, h is a solution of (3.4). We have chosen the initial data of Y for system (3.1) so that  $(y_0)_x^{(i)}(0,0) = h_x^{(i)}(0,0)$  for  $0 \le i \le n-1$ . So h and  $y_0$  are solutions of (3.4) with the same initial values at (0,0). By Frobenius Theorem, there is only one solution with the same initial data. Hence  $y_0 = h$ . This proves  $\hat{E} = Ef_{u,h}^{-1}$  is a frame of the new solution  $\hat{u} = \sum_{i=1}^{n-1} \hat{u}_i e_{in}$  and  $\hat{u}_i = u_i + s_i(u, h)$ . **Corollary 3.15.** Let E be a frame of a solution u of (2.6). Then  $\tilde{E} = E f_{u,h}^{-1}$  is a frame of some solution of (2.6) if and only if h is a solution of (3.4).

**Corollary 3.16.** If u is a solution of (2.6) and h is a solution of (3.4), then there exists a constant  $k \in \mathbb{C}$  such that h is a solution of  $(BT)_{u,k}$  (3.12).

**Corollary 3.17.** If u is a solution of the *j*-th  $A_{n-1}^{(1)}$ -KdV flow (2.6), then

- (a) system (3.4) is solvable,
- (b) the space of solutions of system (3.4) is the union of the spaces of solutions of (BT)<sub>u,k</sub>, i.e., (3.12), for all k ∈ C.

Proof. From the proof of Theorem 3.14, given any constant  $(d_0, \dots, d_{n-1})^t \in \mathbb{C}^n$ , there is a solution h of (3.4) such that  $h_x^{(i)}(0,0) = d_i$  for  $0 \le i \le n-1$ . This proves (a). It follows from Theorem 3.8 and 3.14 that, if h is a solution of (3.4), then there exists a constant  $k \in \mathbb{C}$  such that  $h_x^{(n-1)} = \xi_n(u,h) - k$ . Hence h is a solution of (3.12) with parameter k.

**Definition 3.18.** Let u be a solution of (2.6), and h a solution of  $(BT)_{u,k}$ , i.e., (3.12). We use h \* u to denote the solution  $\tilde{u}$  constructed from u and h as in Corollary 3.11 and call  $u \mapsto h * u$  a Bäcklund transformation with parameter k.

**Example 3.19.** We use the algorithm given in section 2 to compute the second  $A_2^{(1)}$ -KdV flow and obtain

$$\begin{cases} (u_1)_t = (u_1)_{xx} - \frac{2}{3}(u_2)_{xxx} + \frac{2}{3}u_2(u_2)_x, \\ (u_2)_t = -(u_2)_{xx} + 2(u_1)_x. \end{cases}$$
(3.15)

Note that solutions of (3.15) give rise to solutions of the *Boussinesq equation*. In fact, take the derivative with respect to t on both sides of the second equation and use the first equation to see that  $u_2$  satisfies the Boussinesq equation:

$$(u_2)_{tt} = -\frac{1}{3}(u_2)_x^{(4)} + \frac{4}{3}(u_2)_x^2 + \frac{4}{3}u_2(u_2)_{xx}.$$
(3.16)

Note that the right hand side is equal to  $-\frac{1}{3}(u_2)_{xxx} + \frac{4}{3}(u_2(u_2)_x)_x$ .

We use the proof of Theorem 3.4 to compute differential polynomials  $s_i(u,h)$ ,  $r_n(u,h)$  and  $f_{u,h}$ . The system (3.12) for a solution  $u = u_1e_{13} + u_2e_{23}$  of (3.15) is

$$\begin{cases} h_{xx} = -u_1 + (u_2)_x + hu_2 - 3hh_x - h^3 - k, \\ h_t = \frac{2}{3}(u_2)_x - h_{xx} - 2hh_x, \end{cases}$$
(3.17)

and system (3.4) is

$$\begin{cases} h_x^{(3)} = (-u_1 + (u_2)_x + hu_2 - 3hh_x - h^3)_x, \\ h_t = \frac{2}{3}(u_2)_x - h_{xx} - 2hh_x. \end{cases}$$
(3.18)

The new solution  $h * u = \tilde{u}_1 e_{13} + \tilde{u}_2 e_{23}$  of (3.15) is given by

$$\begin{cases} \tilde{u}_1 = u_1 - (u_2)_x + 3hh_x, \\ \tilde{u}_2 = u_2 - 3h_x. \end{cases}$$
(3.19)

Moreover, if E is a frame for u, then  $Ef_{u,h}^{-1}$  is a frame for h \* u, where

$$f_{u,h}(x,t,\lambda) = e_{13}\lambda + \begin{pmatrix} h & h_x & u_1 - (u_2)_x + h_{xx} + 3hh_x \\ 1 & h & u_2 - h_x \\ 0 & 1 & h \end{pmatrix}$$

If u is a solution of (2.6) and h is a solution of  $(BT)_{u,k}$ , then by (3.9), det $(f_{u,h}(x,t,k)) = 0$ . So  $Ef_{u,h}^{-1}$  is not holomorphic at  $\lambda = k$ . However, we can multiply  $Ef_{u,h}^{-1}$  on the left by some  $C(\lambda)$  independent of x, t to get a new frame for h \* u that is holomorphic at  $\lambda = k$ .

**Theorem 3.20.** Suppose  $E(x, t, \lambda)$  is a frame of a solution u of (2.6) that is holomorphic for  $\lambda$  in an open subset  $\mathcal{O}$  of  $\mathbb{C}$ . Let  $k \in \mathcal{O}$  be a constant, and h the solution of (3.12) constructed from E(x, t, k) and  $\mathbf{c} = (c_1, \dots, c_n)^t \in \mathbb{C}^n \setminus \mathbf{0}$  as in Theorem 3.9. Then

- (i)  $\tilde{E} = E f_{u,h}^{-1}$  is a frame of the new solution h \* u and  $\tilde{E}$  is holomorphic for all  $\lambda \in \mathcal{O} \setminus \{k\}$ .
- (iv) Let  $C(\lambda) = e_{1n}(\lambda k) + b + \sum_{i=1}^{n-1} c_i e_{i+1,n}$ . Then

$$E(x,t,\lambda) = C(\lambda)E(x,t,\lambda)f_{u,h}(x,t,\lambda)^{-1}$$

is a frame for h \* u that is holomorphic for all  $\lambda \in \mathcal{O}$ .

Proof. By Corollary 3.17, h is a solution of (3.4). It follows from Theorem 3.14 that h \* u is a solution of (2.6) and  $Ef_{u,h}^{-1}$  is a frame of h \* u. By definition,  $f_{u,h}(x,t,\lambda)$  is holomorphic for all  $\lambda \in \mathcal{O}$ . Since  $\det(f_{u,h}(x,t,\lambda) = (-1)^{n-1}(\lambda-k)$ , if  $\lambda \neq k$  then  $Ef_{u,h}^{-1}$  is holomorphic at  $\lambda$ .

Let  $f = f_{u,h}$ , and  $f^{\sharp}$  the matrix whose *ij*-th entry is  $(-1)^{i+j} \det(M_{ji})$ , where  $M_{ij}$  is the  $(n-1) \times (n-1)$  matrix obtained from f by crossing out the *i*-th row and *j*-th column. Then  $ff^{\sharp} = (-1)^{n-1} (\lambda - k) I_n$ . So we have

$$\operatorname{Im}(f^{\sharp}(x,t,k)) \subset \operatorname{Ker}(f(x,t,k)) = \mathbb{C}E^{-1}(x,t,k)(\mathbf{c}).$$
(3.20)

Let

$$\begin{split} F(x,t,\lambda) &= C(\lambda)E(x,t,\lambda)f_{u,h}^{-1}(x,t,\lambda) \\ &= (-1)^{n-1}\frac{1}{\lambda-k}C(\lambda)E(x,t,\lambda)f^{\sharp}(x,t,\lambda). \end{split}$$

Then  $F(x,t,\lambda)$  is holomorphic for  $\lambda \in \mathcal{O} \setminus \{k\}$  and has a possible simple pole at  $\lambda = k$ . We claim that the residue of  $F(x,t,\lambda)$  at  $\lambda = k$  is zero. The residue of  $F(x,t,\lambda)$  at  $\lambda = k$  is equal to  $(-1)^{n-1}C(k)E(x,t,k)f^{\sharp}(x,t,k)$ . By (3.20), it is equal to  $\phi(x,t)C(k)(\mathbf{c})$  for some function  $\phi$ . A direct computation implies that  $C(k)(\mathbf{c}) = 0$ . This proves the claim and F is holomorphic at  $\lambda = k$ . The above Theorem will be used to construct Bäcklund transformations for the n-d central affine curve flow in [11].

## 4. Permutability and Scaling Transform

In this section, we

- (i) give a Permutability formula for Bäcklund transformations,
- (ii) prove that the conjugation of a Bäcklund transformation with parameter k = 1 by the *r*-scaling transform gives a BT with parameter  $k = r^{-n}$

for the  $A_{n-1}^{(1)}$ -KdV hierarchy.

**Theorem 4.1.** [Permutability for BT]

Let u be a solution of (2.6),  $k_1, k_2 \in \mathbb{C}$  constants,  $h_i$  solutions of  $(BT)_{u,k_i}$ (3.12), and  $h_i * u$  the solution of (2.6) construct from u and  $h_i$  for i = 1, 2. Suppose  $h_1 \neq h_2$ . Set

$$\begin{cases} \tilde{h}_1 = h_1 + \frac{(h_1 - h_2)_x}{h_1 - h_2}, \\ \tilde{h}_2 = h_2 + \frac{(h_1 - h_2)_x}{h_1 - h_2}. \end{cases}$$
(4.1)

Then

(i)  $\tilde{h}_1$  is a solution of  $(BT)_{h_2*u,k_1}$  and  $\tilde{h}_2$  is a solution of  $(BT)_{h_1*u,k_2}$ , (ii)  $\tilde{h}_1 * (h_2 * u) = \tilde{h}_2 * (h_1 * u)$ .

*Proof.* Let  $E(x, t, \lambda)$  be a frame of u. By Theorem 3.10, there exist constant vectors  $v_1^0, v_2^0 \in \mathbb{R}^n$  that give  $h_1, h_2$ , i.e.,

$$v_1 = (v_{1,1}, \cdots, v_{1,n})^t = E(\cdot, \cdot, k_1)^{-1} v_1^0,$$
  

$$v_2 = (v_{2,1}, \cdots, v_{2,n})^t = E(\cdot, \cdot, k_2)^{-1} v_2^0,$$
  

$$h_i = -\frac{v_{i,n-1}}{v_{i,n}}, \quad i = 1, 2.$$

From (3.13),

$$(v_i)_x = -E^{-1}E_x(x,t,k_i)v_i = -(J+u)|_{\lambda=k_i}v_i, \quad i=1,2.$$

In particular, we have

$$\begin{cases} (v_{i,n-1})_x = -v_{i,n-2} - u_{n-1}v_{i,n}, \\ (v_{i,n})_x = -v_{i,n-1}, \end{cases} \quad i = 1, 2.$$

Therefore,

$$(h_i)_x = \frac{v_{i,n-2}}{v_{i,n}} - h_i^2 + u_{n-1}, \quad i = 1, 2.$$
 (4.2)

By Theorem 3.9,  $E_i(x, t, \lambda) = E(x, t, \lambda) f_{u,h_i}^{-1}$  is a frame for  $h_i * u$  for i = 1, 2 respectively. Let

$$\tilde{v}_1 = E_2^{-1}(x, t, k_1)v_1^0, \quad \tilde{v}_2 = E_1(x, t, k_2)^{-1}v_2^0.$$

Now we compute

$$\tilde{h}_i = -\frac{\tilde{v}_{i,n-1}}{\tilde{v}_{i,n}}.$$

From

$$\tilde{v}_1 = E_2^{-1}(x, t, k_1)v_1^0 = f_{u,h_2}(k_1)E(x, t, k_1)^{-1}v_1^0 = f_{u,h_2}(k_1)v_1,$$

we get

$$\begin{cases} \tilde{v}_{1,n-1} = v_{1,n-2} + h_2 v_{1,n-1} + (u_{n-1} - (h_2)_x) v_{1,n}, \\ \tilde{v}_{1,n} = v_{1,n-1} + h_2 v_{1,n}. \end{cases}$$

Together with (4.2), we have

$$\tilde{h}_1 = -h_2 + \frac{v_{1,n-2}v_{2,n} - v_{2,n-2}v_{1,n}}{v_{2,n-1}v_{1,n} - v_{1,n-1}v_{2,n}} = h_1 + \frac{(h_1 - h_2)_x}{h_1 - h_2}.$$

Similarly,  $\tilde{h}_2 = h_2 + \frac{(h_1 - h_2)_x}{h_1 - h_2}$ .

 $(\gamma$ 

A direct computation implies 
$$f_{u_2,\tilde{h}_1}f_{u,h_2} = f_{u_1,\tilde{h}_2}f_{u,h_1}$$
. Hence  $\tilde{h}_2 * (h_1 * u) = \tilde{h}_1 * (h_2 * u)$ .

Next we review the scaling transform of the  $A_{n-1}^{(1)}$ -KdV hierarchy.

**Proposition 4.2.** ([3]) Let  $u = \sum_{i=1}^{n-1} u_i e_{in}$  be a solution of the *j*-th  $A_{n-1}^{(1)}$ -KdV flow, and  $r \in \mathbb{R} \setminus \{0\}$ . Let  $D(r) = \text{diag}(1, r, \dots, r^{n-1})$ , and

$$(x,t) = r^{n+1-i}u_i(rx,r^jt), \quad 1 \le i \le n-1.$$

Then

(i)  $(r \cdot u) = \sum_{i=1}^{n-1} (r \cdot u_i) e_{in}$  is a solution of the *j*-th  $A_{n-1}^{(1)}$ -KdV flow, (ii) if  $E(x, t, \lambda)$  is a frame of *u* then

$$\tilde{E}(x,t,\lambda) := D(r)^{-1}E(rx,r^jt,r^{-n}\lambda)D(r)$$

is a frame of  $r \cdot u$ .

So the multiplicative group  $\mathbb{R}\setminus\{0\}$  acts on the space of solutions of the *j*-th  $A_{n-1}^{(1)}$ -KdV flow. The following Theorem proves that the conjugation of BT with parameter k = 1 by a scaling transform gives BT with arbitrary non-zero real parameter.

**Theorem 4.3.** Let u be a solution of the j-th  $A_{n-1}^{(1)}$ -KdV flow,  $r \in \mathbb{R} \setminus \{0\}$ , and h a solution of  $(BT)_{r \cdot u, 1}$ . Then

(i)  $\hat{h}(x,t) = r^{-1}h(r^{-1}x,r^{-j}t)$  is a solution of  $(BT)_{u,r^{-n}}$ , (ii)  $r^{-1} \cdot (h * (r \cdot u)) = \hat{h} * u$ .

*Proof.* Let E be the frame of u with  $E(0,0,\lambda) = I_n$ . By Proposition 3.6,

$$\det E(x,t,\lambda) = 1. \tag{4.3}$$

From Proposition 4.2,

$$F(x,t,\lambda) = D(r)^{-1}E(rx,r^{j}t,r^{-n}\lambda)D(r)$$

is a frame for  $\tilde{u} = r \cdot u$ , where  $D(r) = \text{diag}(1, r, \cdots, r^{n-1})$ . It follows from Theorem 3.14 that

$$F_1(x,t,\lambda) = F(x,t,\lambda)f_{r\cdot u,h}^{-1}(x,t,\lambda)$$

is a frame for  $h * (r \cdot u)$ . Apply the scaling transform given by  $r^{-1}$  to  $h * \tilde{u}$ , we have

$$F_{2}(x,t,\lambda) = D(r)F_{1}(r^{-1}x,r^{-j}t,r^{n}\lambda)D(r)^{-1}$$
  
=  $E(x,t,\lambda)D(r)f_{\tilde{u},h}^{-1}(r^{-1}x,r^{-j}t,r^{n}\lambda)D(r)^{-1}.$  (4.4)

is a frame for  $\hat{u} = r^{-1} \cdot (h * (r \cdot u))$ . A direct computation implies that

$$D(r)f_{\tilde{u},h}(r^{-1}x,r^{-j}t,r^{n}\lambda)D(r)^{-1} = r(e_{1n}\lambda + b + r^{-1}h(r^{-1}x,r^{-j}t)\mathbf{I}_{n} + r^{-1}D(r)ND(r)^{-1}),$$

where  $f_{\tilde{u},h} = e_{1n}\lambda + b + hI_n + N$  and N is strictly upper triangular. So we have proved  $F_2(x,t,\lambda) = r^{-1}E(x,t,\lambda)g^{-1}(x,t,\lambda)$  is a frame of  $\hat{u} = r^{-1} \cdot (h * (r \cdot u)),$  where

$$g(x,t,\lambda) = e_{1n}\lambda + b + \hat{h}(x,t)I_n + r^{-1}D(r)N(r^{-1}x,r^{-j}t)D(r)^{-1}$$

Note that  $r^{-1}D(r)N(r^{-1}x,r^{-j}t)D(r)^{-1} \in \mathcal{N}_n^+$ , and

$$\hat{E}(x,t,\lambda) = rF_2(x,t,\lambda) = E(x,t,\lambda)g^{-1}(x,t,\lambda)$$
(4.5)

is a frame of  $\hat{u}$ . So  $g = f_{u,\hat{h}}$ .

By Theorem 3.8, we have  $det(f_{\tilde{u},h}) = (-1)^{n-1}(\lambda - 1)$ . It follows from (4.3), (4.4) and (4.5) that

$$\det(f_{u,\hat{h}}^{-1}) = \det(E(x,t,\lambda)) = r^{n}\det(F_{2}(x,t,\lambda))$$
$$= r^{n}\det(f_{\tilde{u},h}^{-1}(r^{-1}x,r^{-j}t,r^{n}\lambda))$$
$$= (-1)^{n-1}\frac{r^{n}}{r^{n}\lambda - 1} = \frac{(-1)^{n-1}}{\lambda - r^{-n}}.$$

Hence  $\det(f_{u,\hat{h}})=(-1)^{n-1}(\lambda-r^{-n}).$ 

#### 5. Explicit solutions

In this section we apply Theorem 3.20 to u = 0 to construct explicit solutions of (2.6).

Proposition 5.1. If we apply BTs with non-zero parameters to the vacuum solution u = 0 of the *j*-th  $A_{n-1}^{(1)}$ -KdV flow (2.6) repeatedly, then we obtain infinitely many families of explicit solutions of (2.6) that are rational functions of exponential functions.

*Proof.* Let  $\lambda = z^n$ ,  $\alpha = e^{\frac{2\pi i}{n}}$ ,

$$D(z) = \text{diag}(1, z, \dots, z^{n-1}),$$
  

$$\Xi = (\alpha^{(i-1)(j-1)}),$$
  

$$A_i(x, t, z) = \exp(\alpha^{i-1}zx + (\alpha^{i-1}z)^j t),$$
  

$$(m_1(x, t, z), \dots, m_n(x, t, z)) = (e^{A_1}, \dots, e^{A_n})\Xi.$$

A direct computation implies that the vacuum frame is

$$E(x,t,z^{n}) = e^{xJ+tJ^{j}} = \frac{1}{n}D(z)^{-1} \begin{pmatrix} m_{1} & m_{2} & \cdots & m_{n} \\ m_{n} & m_{1} & \cdots & m_{n-1} \\ \vdots & \vdots & \vdots & \vdots \\ m_{2} & m_{3} & \cdots & m_{1} \end{pmatrix} D(z).$$

Given constants  $k \neq 0$  and  $\mathbf{c} = (c_1, \cdots, c_n)^t \in \mathbb{C}^n$ , let

$$(v_1, v_2, \cdots, v_n)^t = E(x, t, k^n)^{-1}(\mathbf{c}), \quad h = -\frac{v_{n-1}}{v_n}.$$
 (5.1)

Note that

$$E(x,t,k^n)^{-1} = E(-x,-t,k^n)$$

Let  $\{e_1, \ldots, e_n\}$  be the standard basis of  $\mathbb{C}^{n \times 1}$ , then

$$(m_3, \dots, m_n, m_1, m_2) = (e^{A_1}, \dots, e^{A_n}) \Xi \pi_1,$$
  
 $(m_2, m_3, \dots, m_n, m_1) = (e^{A_1}, \dots, e^{A_n}) \Xi \pi_2.$ 

where  $\pi_1 = (e_3, ..., e_n, e_1, e_2), \pi_2 = (e_2, e_3, ..., e_n, e_1)$  (permutation matrices). Hence

$$v_{n-1}(x,t) = k^{-(n-2)} (e^{A_1(-x,-t,k)}, \dots, e^{A_n(-x,-t,k)}) \Xi \pi_1 D(k) \mathbf{c},$$
  
$$v_n(x,t) = k^{-(n-1)} (e^{A_1(-x,-t,k)}, \dots, e^{A_n(-x,-t,k)}) \Xi \pi_2 D(k) \mathbf{c}.$$

This implies that h given by (5.1) is a rational function of exponentials. Hence entries of the new solution h \* 0 are rational functions of exponential functions. It follows from Theorem 3.20 that, the entries of the frame of h \* 0,

$$E(x,t,\lambda) = C(\lambda)E(x,t,\lambda)f_{u,h}(x,t,\lambda),$$

are given by rational functions of exponentials, where  $C(\lambda) = e_{1n}(\lambda - k) + b + \sum_{i=1}^{n-1} c_i e_{i+1,n}$ . If we apply Theorem 3.20 to h \* 0 with parameter  $k_2 \neq 0$ , then we get another family of new solutions that are rational of exponential functions.

**Example 5.2** (Explicit solutions of the *j*-th  $A_{n-1}^{(1)}$ -KdV flow).

We use the same notation as in the proof of Proposition 5.1. First note that  $\Xi \pi_2 D(k)$  is invertible. So given any  $\mathbf{a} \in \mathbb{C}^{n \times 1}$ , we can find suitable  $\mathbf{c} \in \mathbb{C}^n$  such that  $\Xi \pi_2 D(k) \mathbf{c} = \mathbf{a}$ . We will write down some explicit solutions for the *j*-th  $A_{n-1}^{(1)}$ -KdV flow (2.6) below.

Choose **c** such that  $\Xi \pi_2 D(k) \mathbf{c} = e_1 + e_2$ . A simple computation implies that  $\Xi \pi_1 D(k) \mathbf{c} = e_1 + \alpha e_2$ . The function *h* defined by (5.1) is

$$h = -k \frac{e^{-(kx+k^{j}t)} + \alpha e^{-(\alpha kx + \alpha^{j}k^{j}t)}}{e^{-(kx+k^{j}t)} + e^{-(\alpha kx + \alpha^{j}k^{j}t)}}.$$
(5.2)

This is a solution of (3.12) with parameter  $k \in \mathbb{C}$ . Let A, B be defined by

$$\begin{cases} A+B = -(kx+k^jt), \\ A-B = -(\alpha kx+\alpha^j k^jt). \end{cases}$$

Then (5.2) becomes  $h = -k \frac{e^{A+B} + \alpha e^{A-B}}{e^{A+B} + e^{A-B}}$ , so we get

$$h = \frac{\kappa}{2} \left( (1 - \alpha) \tanh B - (1 + \alpha) \right), \tag{5.3}$$

where  $\alpha = e^{\frac{2\pi i}{n}}$  and  $B = \frac{1}{2}((1-\alpha)kx + (1-\alpha^j)k^jt)$ . Note that *h* defined by (5.3) is a solution of (3.12) with parameter *k*.

(i) If  $k = (1 + \bar{\alpha})\mu$  for some  $\mu \in \mathbb{R}$ , then h given by (5.3) becomes a real function,

$$h = -\mu \sin(\frac{2\pi}{n}) \tan \theta - (1 + \cos(\frac{2\pi}{n}))\mu,$$
 (5.4)

where  $\theta = \mu \sin(\frac{2\pi}{n})x + \sum_{i=0}^{j-1} C_{j,i} \mu^j \sin(\frac{2(j-i)\pi}{n})t$ . Hence  $h * 0 = 3h_x(he_{13} - e_{23})$ 

is a real solution of the *j*-th  $A_{n-1}^{(1)}$ -KdV flow. In particular, when j = 2, (5.4) becomes

 $h = -\mu \sin \gamma \tan(\mu \sin \gamma (x + 2(1 + \cos \gamma)\mu t)) - (1 + \cos \gamma)\mu, \qquad (5.5)$ 

where  $\gamma = \frac{2\pi}{n}$ , and  $\mu \in \mathbb{R}$ . Then  $h * 0 = 3h_x(he_{13} - e_{23})$  is a real solution of the second  $A_{n-1}^{(1)}$ -KdV flow.

(ii) If n = 2m and j is odd, we choose **c** such that  $\Xi \pi_2 D(k)$ **c** =  $e_1 + e_{m+1}$ . Then  $\Xi \pi_1 D(k)$ **c** =  $e_1 - e_{m+1}$ , and h defined by (5.1) is

$$h = k \tanh(kx + k^j t).$$

When  $k \in \mathbb{R}\setminus 0$ , we obtain soliton solutions for the *j*-th  $A_{2m-1}^{(1)}$ -KdV flow. For example, when m = 1 and j = 3, this give rise to the 1-soliton solutions of the KdV equation.

(iii) If n = 2m and j is even, we choose **c** such that  $\Xi \pi_2 D(k)\mathbf{c} = e_1 + e_{m+1}$ . Then  $\Xi \pi_1 D(k)\mathbf{c} = e_1 - e_{m+1}$ , and h defined by (5.1) is

$$h = k \tanh(kx).$$

This gives stationary smooth solutions h \* 0 for the *j*-th  $A_{2m-1}^{(1)}$ -KdV flow.

**Example 5.3** (Explicit solutions for the second  $A_2^{(1)}$ -KdV flow (3.15)). For n = 3, we have  $\alpha = e^{\frac{2\pi i}{3}}$ .

(i) Let 
$$\frac{k}{2}(1-\alpha) = -\mu$$
. Then *h* defined by (5.3) becomes  

$$h = \sqrt{3\mu} \tanh(\sqrt{3\mu}(x-2\mu it)) + \mu i.$$
(5.6)

Substitute (5.6) to (3.19) with u = 0 to see that

$$\begin{cases} u_1 = 9\mu^3 \operatorname{sech}^2(\sqrt{3}\mu(x - 2\mu i t))(\sqrt{3} \tanh(\sqrt{3}\mu(x - 2\mu i t)) + i), \\ u_2 = -9\mu^2 \operatorname{sech}^2(\sqrt{3}\mu(x - 2\mu i t)) \end{cases}$$

is a complex solution of (3.15). Note that  $u_2$  is the solution of the Boussinesq equation (3.16) obtained in [4].

(ii) h given by (5.5) is

$$h = -\frac{\sqrt{3}}{2}\mu \tan(\frac{\sqrt{3}}{2}\mu(x+\mu t)) - \frac{\mu}{2}$$

Let  $\nu = \frac{\mu}{2}$ . Then

$$h_{\nu} = -\sqrt{3}\nu \tan(\sqrt{3}\nu(x+2\nu t)) - \mu$$

is a real solution of (3.15) and  $u_1, u_2$  defined by

$$\begin{cases} u_1 = 9\nu^3 \sec^2(\sqrt{3}\nu(x+2\nu t))(1+\sqrt{3}\tan(\sqrt{3}\nu(x+2\nu t))), \\ u_2 = 9\nu^2 \sec^2(\sqrt{3}\nu(x+2\nu t)), \end{cases}$$

is the real solution of (3.15).

**Proposition 5.4.** If we apply BT with parameter k = 0 to the vacuum solution u = 0 of the *j*-th  $A_{n-1}^{(1)}$ -KdV flow (2.6) repeatedly, then we obtain infinitely many families of explicit rational solutions of (2.6)

*Proof.* Note that  $E(x,t,\lambda) = e^{xJ+tJ^j}$  is a frame of the solution u = 0 of (2.6). Since entries of  $E(x,t,0) = \exp(bx + b^{j}t)$  are polynomials in x and t, we see that the solution h of (3.12) constructed in Theorem 3.9 with k = 0is a rational solution and  $\tilde{E} = C(\lambda)Ef_{0,h}^{-1}$  is a frame of the solution h \* 0. Apply Theorem 3.20 with parameter k = 0 and a constant vector  $\mathbf{c}_1$  to h \* 0 to get another solution. But  $\tilde{E}(x,t,0) = C(0)E(x,t,0)f_{0,h}^{-1}(x,t,0)$  is rational in x, t. So the solution  $h_1$  of (3.12) with u = h \* 0 is rational. Hence  $h_1 * (h * 0)$  is a rational function. 

**Example 5.5.** [Rational solutions for the second  $A_2^{(1)}$ -KdV flow] The coefficients of the constant term and  $\lambda^{-1}$  of the frame  $E(x, t, \lambda) =$  $e^{xJ+tJ^2}$  of the vacuum solution u = 0 as a power series in  $\lambda$  are

$$E_{0}(x,t) = \exp(bx + b^{2}t) = \begin{pmatrix} 1 & 0 & 0 \\ x & 1 & 0 \\ \frac{x^{2}}{2} + t & x & 1 \end{pmatrix},$$

$$E_{1}(x,t) = \begin{pmatrix} xt + \frac{1}{6}x^{3} & t + \frac{1}{2}x^{2} & x \\ \frac{1}{2}t^{2} + \frac{1}{2}x^{2}t + \frac{1}{24}x^{4} & xt + \frac{1}{6}x^{3} & t + \frac{1}{2}x^{2} \\ \frac{1}{2}xt^{2} + \frac{1}{6}x^{3}t + \frac{1}{5!}x^{5} & \frac{1}{2}t^{2} + \frac{1}{2}x^{2}t + \frac{1}{24}x^{4} & xt + \frac{1}{6}x^{3} \end{pmatrix}.$$

We apply Theorem 3.20 to u = 0 and  $v_0 = (a_1, a_2, 1)^t$  to get new solutions:

(i)  $h = \frac{a_1 x - a_2}{1 + a_1(\frac{x^2}{2} - t) - a_2 x}$  is a solution of (3.18). (ii)  $h * 0 = u_1 e_{13} + u_2 e_{23}$  is a rational solution of (3.15), where  $\begin{cases} u_1 = -\frac{3(a_1 x - a_2)(\frac{1}{2}a_1^2 x^2 - a_1 a_2 x + a_1^2 t + a_2^2 - a_1)}{(1 + a_1(\frac{x^2}{2} - t) - a_2 x)^3}, \\ u_2 = \frac{3(\frac{1}{2}a_1^2 x^2 - a_1 a_2 x + a_1^2 t + ba_2^2 - a_1)}{(1 + a_1(\frac{x^2}{2} - t) - a_2 x)^2}. \end{cases}$ 

**Example 5.6.** [Rational solutions for the second  $A_{n-1}^{(1)}$ -KdV flow]

Recall that  $E(x,t,\lambda) = e^{xJ+tJ^2}$  is a frame of u = 0. The coefficients of the constant and  $\lambda$  of  $E(x,t,\lambda)$  as a power series in  $\lambda$  are  $E(x,t,0) = \exp(bx + b^2t)$  and  $E_1(x,t) = \frac{\partial}{\partial\lambda}|_{\lambda=0}\exp(xJ + tJ^2)$ . A direct computation implies that  $J^k = (b^t)^{n-k}\lambda + b^k$ ,  $J^n = \lambda I_n$ , and  $\frac{\partial}{\partial\lambda}|_{\lambda=0}J^k = (b^t)^{n-k}$  for  $1 \le k \le n-1$ . So entries of  $E_1(x,t)$  are polynomials in x,t. In fact, for n = 2m + 1, we get

$$E_{1}(x,t) = (I + \sum_{j=1}^{n-1} \frac{x^{j}}{j!} b^{j}) (\sum_{j=1}^{m} \frac{t^{j}}{j!} (b^{t})^{n-2j} + \sum_{j=1}^{m} \frac{t^{m+j}}{(m+j)!} b^{2j-1}) + (\sum_{j=1}^{n-1} \frac{x^{j}}{j!} (b^{t})^{j} + \frac{x^{n}}{n!} + \sum_{j=1}^{n-1} \frac{x^{n+j}}{(n+j)!} b^{j}) (I + \sum_{j=1}^{m} \frac{t^{j}}{j!} b^{2j}).$$

For n = 2m,

$$E_{1}(x,t) = \left(I + \sum_{j=1}^{n-1} \frac{x^{j}}{j!} b^{j}\right) \left(\sum_{j=1}^{m-1} \frac{t^{j}}{j!} (b^{t})^{n-2j} + \frac{t^{m}}{m!} + \sum_{j=1}^{m-1} \frac{t^{m+j}}{(m+j)!} b^{2j}\right) \\ + \left(\sum_{j=1}^{n-1} \frac{x^{j}}{j!} (b^{t})^{j} + \frac{x^{n}}{n!} + \sum_{j=1}^{n-1} \frac{x^{n+j}}{(n+j)!} b^{j}\right) \left(I + \sum_{j=1}^{m-1} \frac{t^{j}}{j!} b^{2j}\right).$$

Apply Theorem 3.20 repeatedly to construct infinitely many families of explicit rational solutions.

## 6. $Z_n$ -ACTION FOR THE KW FLOWS

In this section we give a natural action of the cyclic group  $Z_n$  of n elements on the space of solutions of the j-th KW flow and show that Adler's Bäcklund transformation comes from this  $Z_n$ -action.

**Theorem 6.1.** Let  $\alpha = e^{\frac{2\pi i}{n}}$ ,  $a = \text{diag}(1, \alpha, \dots, \alpha^{n-1})$ , and  $\tau = e_{21} + e_{32} + \dots + e_{n,n-1} + e_{1n}$  as in the KW-hierarchy. If  $v = (v_1, \dots, v_{n-1})$  is a solution of the *j*-th KW flow (2.17), then

$$\alpha \cdot v = (\alpha v_1, \dots, \alpha^{(n-1)} v_{n-1})$$

is a solution of (2.17). In particular, this defines an  $\mathbb{Z}_n$ -action on the space of solutions of (2.17).

*Proof.* Let  $\mathcal{L}^{KW} = \mathcal{L}^{KW}_+ \oplus \mathcal{L}^{KW}_-$  be the splitting that gives the KW-hierarchy in section 2.

First note that  $a\tau = \alpha \tau a$ . A direct computation implies that  $\operatorname{Ad}(a^{-1})$  leaves  $\mathcal{L}^{KW}_{\pm}$  invariant, i.e.,  $a^{-1}(\mathcal{L}^{KW}_{\pm})a \subset \mathcal{L}^{KW}_{\pm}$ . So

$$a^{-1}(\partial_x + az + P(v))a = \partial_x + az + P(\alpha \cdot v).$$

Recall that  $\hat{Q}(v, z)$  is defined by  $[\partial_x + az + P(v), \hat{Q}(v, z)] = 0$  and  $\hat{Q}(v, z)$  is conjugate to az. Hence  $\hat{Q}(\alpha \cdot v, z) = a^{-1}\hat{Q}(v, z)a$ , which implies that

$$(\hat{Q}^{j}(\alpha \cdot v, z))_{+} = a^{-1}(\hat{Q}^{j}(v, z))_{+}a.$$
(6.1)

Since v is a solution of (2.17), we have

$$[\partial_x + az + P(v), \partial_t + (\hat{Q}^j(v, z))_+] = 0.$$

Conjugate the above equation by  $a^{-1}$  to get

$$a[\partial_x + az + P(v), \partial_t + (Q^j(v, z))_+]a^{-1}$$
  
=  $[\partial_x + az + P(\alpha \cdot v), \partial_t + a^{-1}(\hat{Q}^j(v, z))_+a], \text{ by (6.1)},$   
=  $[\partial_x + az + P(\alpha \cdot v), \partial_t + (\hat{Q}^j(\alpha \cdot v, z))_+] = 0.$ 

This proves that  $\alpha \cdot v$  is a solution of (2.17).

Next we use Theorem 6.1 to show that Adler's BT arises naturally from the  $Z_n$ -action on the KW-flows.

**Corollary 6.2.** Let  $\Phi : \mathbb{R}^{1 \times (n-1)} \to \mathcal{T}_n$  be the linear isomorphism defined by  $\Phi(v) = \operatorname{diag}(q_1, \ldots, q_n)$  in Theorem 2.10, where  $q_i = \sum_{i=1}^{n-1} \alpha^{k(i-1)} v_k$ . Let  $q = \operatorname{diag}(q_1, \ldots, q_n)$  be a solution of the *j*-th  $n \times n$  mKdV flow, and  $v = \Phi^{-1}(q)$  the corresponding solution of the *j*-th KW flow. Then  $\Phi(\alpha^{n-1} \cdot v) = \operatorname{diag}(q_n, q_1, \ldots, q_{n-1})$  and is a solution of the *j*-th  $n \times n$  mKdV flow.

*Proof.* By Theorem 2.10, v is a solution of the *j*-th KW flow. By Theorem 6.1,  $\alpha \cdot v = (\alpha v_1, \ldots, \alpha^{n-1} v_{n-1})$  is a solution of the *j*-th KW flow. Hence  $\Phi(\alpha \cdot v)$  is a also a solution of the *j*-th  $n \times n$  mKdV flow. A simple computation implies that  $\Phi(\alpha \cdot v) = \text{diag}(q_2, \ldots, q_n, q_1)$ . So

$$\Phi(\alpha^{n-1} \cdot v) = \operatorname{diag}(q_n, q_1, \dots, q_{n-1})$$

is a solution of the *j*-th  $n \times n$  mKdV flow.

This Corollary gives a proof of Adler's result: Suppose

$$L = \partial^n - \sum_{i=1}^{n-1} u_i \partial^{(i-1)} = (\partial - q_1) \cdots (\partial - q_n)$$

such that L is a solution of the j-th  $\text{GD}_n$  flow (1.1) and  $q = \text{diag}(q_1, \ldots, q_n)$  is a solution of the j-th  $n \times n$  mKdV flow (2.9). By Corollary 6.2,

$$\operatorname{diag}(q_n, q_1, \ldots, q_{n-1})$$

is a solution of (2.9). It follows from Theorem 2.11 that  $L = (\partial - q_n)(\partial - q_1)\cdots(\partial - q_{n-1})$  is a solution of (1.1).

### 7. Relation between Adler's BT and our BT

In this section, we prove that if L is a solution of the *j*-th GD<sub>n</sub> flow (1.1), then  $\tilde{L} = (\partial + h)^{-1}L(\partial + h)$  is a solution of (1.1) if and only if h is a solution of (3.4). We also show that Adler's BT is our BT with parameter k = 0.

**Proposition 7.1.** Let L be a solution of the *j*-th  $GD_n$  flow (1.1) and h:  $\mathbb{R}^2 \to \mathbb{C}$ . Then  $\tilde{L} = (\partial + h)^{-1}L(\partial + h)$  is a n-th order differential operator and is a solution of (1.1) if and only if

$$\begin{cases} h_x = r_n(u, h), \\ h_t = w_{n,j}(u, h), \end{cases}$$
(7.1)

for some differential polynomial  $w_{n,j}(u,h)$ , where  $r_n(u,h)$  is as in Theorem 3.4.

*Proof.* It follows from Lemma 3.3 and the proof of Theorem 3.4 that  $\tilde{L}$  is in  $\mathcal{D}_n$  if and only if h satisfies  $h_x^{(n)} = r_n(u,h)$  and the coefficient  $\tilde{u}_i$  of  $\partial^{i-1}$  is given by (3.2). So  $\tilde{L}$  is a solution of (1.1) if and only if

$$\tilde{L}_t = [(\partial + h)^{-1} (L^{\frac{1}{n}})_+ (\partial + h) - (\partial + h)^{-1} h_t, \tilde{L}].$$

Note that  $\tilde{L}^{j/n} = (\partial + h)^{-1} L^{j/n} (\partial + h) = \partial^j + \cdots$ , by comparing the highest power of  $\partial$ , we have

$$(\partial + h)^{-1} (L^{\frac{2}{n}})_+ (\partial + h) - (\partial + h)^{-1} h_t = (\tilde{L}^{\frac{2}{n}})_+.$$

Hence we obtain

$$h_t = -(L^{\frac{j}{n}})_-(\partial + h) + (\partial + h)(\tilde{L}^{\frac{j}{n}})_-$$

Let  $p_{-1}$ ,  $\tilde{p}_{-1}$  denote the coefficient of  $\partial^{-1}$  of  $L^{j/n}$  and  $\tilde{L}^{j/n}$  respectively. Since coefficients of  $(L^{j/n})_+$  and  $(L^{\frac{j+n}{n}})_+$  are differential polynomials in u and  $\tilde{u}$  respectively,  $p_{-1}$  and  $\tilde{p}_{-1}$  are differential polynomials in u and h. So we have

$$h_t = w_{n,j}(u,h) := -p_{-1}(u,h) + \tilde{p}_{-1}(u,h).$$
(7.2)

Hence we proved the proposition.

**Theorem 7.2.** Let  $L = \partial^n - \sum_{i=1}^{n-1} u_i \partial^{i-1}$  be a solution of the *j*-th  $GD_n$  flow (1.1), and  $h \in C^{\infty}(\mathbb{R}^2, \mathbb{C})$ . Then the following are equivalent:

- (a)  $\tilde{L} = (\partial + h)^{-1}L(\partial + h)$  is a solution of the *j*-th  $GD_n$  flow (1.1),
- (b) h is a solution of (7.1),
- (c) h is a solution of (3.4),
- (d) h is a solution of  $(BT)_{u,k}$  (3.12) for some constant  $k \in \mathbb{C}$ .

*Proof.* The equivalence of (a) and (b) is given by Proposition 7.1, and the equivalence of (c) and (d) is given by Corollary 3.17.

We prove (a) implies (c). Suppose  $\tilde{L} = (\partial + h)^{-1}L(\partial + h)$  is a solution of (1.1). Write  $\tilde{L} = \partial^n - \sum_{i=1}^{n-1} \tilde{u}_i \partial^{i-1}$ . By Theorem 2.3, both  $u = \sum_{i=1}^{n-1} u_i e_{in}$  and  $\tilde{u} = \sum_{i=1}^{n-1} \tilde{u}_i e_{in}$  are solutions of (2.6). Let  $\tilde{E}(x, t, \lambda)$  be a frame of

the solution  $\tilde{u}$ , and  $\tilde{\phi} = (\tilde{\phi}_1, \ldots, \tilde{\phi}_n)^t$  the first column of  $\tilde{E}$ . It follows from Lemma 3.3 that  $\{\tilde{\phi}_1, \ldots, \tilde{\phi}_n\}$  are solutions of  $\tilde{L} - \lambda = 0$ . Since  $L - \lambda = (\partial + h)(\tilde{L} - \lambda)(\partial + h)^{-1}$ ,  $\phi_i = (\partial + h)\tilde{\phi}_i$  is a solution of  $L - \lambda = 0$  for  $1 \leq i \leq n$ . Let  $E = ((\phi_i)_x^{(j-1)})$ , and  $f = \tilde{E}^{-1}E$ . So  $E = \tilde{E}f$ . A direct computation implies that the first column of f is  $(h, 1, \ldots, 0)^t$  and  $\det(f(x, t, \lambda))$  is of degree one in  $\lambda$ . Since  $\tilde{E}$  is a frame for  $\tilde{u}$ ,  $\det(\tilde{E})$  is independent of x, t, is an analytic function  $c(\lambda)$ , and  $c(\lambda) \neq 0$  for all  $\lambda \in \mathbb{C}$ . So  $\det(E(x, t, \lambda)) \neq 0$ for generic  $(x, t, \lambda)$ , i.e.,  $\{\phi_1, \ldots, \phi_n\}$  is a basis of  $L - \lambda = 0$  for generic  $\lambda$ . Hence E is a frame of u. By Theorem 3.4,  $E = \tilde{E}f_{u,h}$  and h is a solution of (3.4).

Next we prove (d) implies (a). Suppose h is a solution of  $(BT)_{u,k}$  (3.12). By Theorem 3.14 we have

- (i)  $\tilde{u} = h * u$  is a solution of (2.6),
- (ii) if  $\tilde{E}$  is a frame of  $\tilde{u}$ , then  $E = \tilde{E} f_{u,h}$  is a frame for u for all  $\lambda \neq k$ .
- (iii) the *i*1-th entries of  $\phi_i E$  and  $\tilde{\phi}_i \tilde{E}$  are related by  $\phi_i = (\partial + h)\tilde{\phi}_i$ .

Assume  $\lambda \neq k$ . Lemma 3.3 implies that  $\{\phi_1, \ldots, \phi_n\}$  is a basis of  $L - \lambda = 0$ . This is also a basis of  $(\partial + h)(\tilde{L} - \lambda)(\partial + h)^{-1} = 0$ , where  $\tilde{L} = \partial^n - \sum_{i=1}^{n-1} \tilde{u}_i \partial^{(i-1)}$ . It follows from Lemma 3.3 that  $L - \lambda = (\partial + h)(\tilde{L} - \lambda)(\partial + h)^{-1}$ . Hence  $L = (\partial + h)\tilde{L}(\partial + h)^{-1}$ . Or equivalently,  $\tilde{L} = (\partial + h)^{-1}L(\partial + h)$ .

Although both  $\eta_{n,j}(u, h)$  and  $w_{n,j}(u, h)$  can be computed, these two differential polynomials are not in explicit closed form; hence we do not know whether they are the same. But systems (7.1) and (3.4) have the same set of solutions.

**Corollary 7.3.** Let  $L = \partial^n - \sum_{i=1}^{n-1} u_i \partial^{i-1}$  be a solution of the *j*-th  $GD_n$ flow (1.1),  $u = \sum_{i=1}^{n-1} u_i e_{in}$ , and  $k \in \mathbb{C}$  a constant. Let  $p_{j,i}(u, \lambda)$  denote the *i*1-th entry of  $Z_j(u,k)$ , where  $Z_j(u,\lambda)$  is defined by (2.5) for the *j*-th  $A_{n-1}^{(1)}$ -KdV flow. Then

$$\begin{cases} (L-k)\phi = 0, \\ \phi_t = \sum_{i=1}^n p_{j,i}(u,k)\phi_x^{(i-1)}, \end{cases}$$
(7.3)

is solvable for  $\phi : \mathbb{R}^2 \to \mathbb{C}$ , Moreover, if  $\phi_1, \ldots, \phi_{n-1}$  are linearly independent solutions of (7.3), then

 $h = (\ln(W(\phi_1, \cdots, \phi_{n-1})))_x,$ 

is a solution of  $(BT)_{u,k}$  (3.12) and  $\tilde{L} = (\partial + h)^{-1}L(\partial + h)$  is a solution of (1.1), where  $W(f_1, \ldots, f_{n-1}) = \det((f_i)_x^{(j-1)})$  is the Wronskian.

*Proof.* By Theorem 2.3, u is a solution of (2.6). Let  $g = (\phi, \phi_x, \dots, \phi_x^{(n-1)})$ . Then system (7.3) is equivalent to

$$\begin{cases} g^{-1}g_x = ke_{1n} + b + u, \\ g^{-1}g_t = Z_j(u,k). \end{cases}$$
(7.4)

This is the Lax pair of u with  $\lambda = k$ , so it is solvable. If  $\phi$  is a solution of (7.3), then g satisfies (7.4). Let  $(v_1, \ldots, v_n)^t = g^{-1}e_n$ . It follows from Theorem 3.9 that  $h = -\frac{v_{n-1}}{v_n}$  is a solution of (3.12) with parameter k. Use Cramer's rule to get the formula of h in terms of Wronskians.

For example, when n = 3, equation (7.3) is

$$\begin{cases} \phi_{xxx} - u_2\phi_x - u_1\phi = \lambda\phi, \\ \phi_t = -\frac{2}{3}u_2\phi + \phi_{xx}. \end{cases}$$
(7.5)

Note that  $\tilde{L}$  and L in Adler's Theorem satisfies the condition  $\tilde{L} = (\partial - q_n)^{-1}L(\partial - q_n)$ . So by Theorem 7.2,  $-q_n$  is a solution of (1.3) for some  $k \in \mathbb{C}$ . Next we show that k = 0. In other words, Adler's BT is our BT with parameter k = 0.

**Theorem 7.4.** Let  $L = \partial^n - \sum_{i=1}^{n-1} u_i \partial^{(i-1)} = (\partial - q_n) \cdots (\partial - q_1)$  be a solution of the *j*-th  $GD_n$  flow (1.1) such that  $q = \sum_{i=1}^n q_i e_{ii}$  is a solution of the *j*-th  $n \times n$  mKdV flow. Then  $-q_n$  is a solution of  $(BT)_{u,0}$ , *i.e.*, (3.12) with parameter k = 0.

*Proof.* Let K denote the parallel frame for the Lax pair of the solution q of the j-th  $n \times n$  mKdV flow at  $\lambda = 0$  with  $K(0,0) = I_n$ , i.e., K is the solution of

$$\begin{cases} K^{-1}K_x = b + q, \\ K^{-1}K_t = \pi_{bn}(Q_{j,0}(q)) \end{cases}$$

where  $Q_{j,0}(q)$  is defined in section 2,  $b = \sum_{i=1}^{n-1} e_{i+1,i}$ , and  $\pi_{bn}$  is the projection of  $sl(n, \mathbb{C})$  onto  $\mathcal{B}_n^-$  along  $\mathcal{N}_n^+$ . Since both b + q and  $\pi_{bn}(Q_{j,0}(q))$  are lower triangular and  $K(0,0) = I_n$ , we see that  $K(x,t) \in B_n^-$ , i.e. lower triangular, for all x, t. Let  $\phi = (\phi_1, \ldots, \phi_n)^t$  denote the first column of P. It follows from  $K_x = K(b+q)$  that the *j*-th column of K is

$$(\partial - q_{j-1}) \cdots (\partial - q_1)\phi,$$

and  $L\phi = (\partial - q_n) \cdots (\partial - q_1)\phi = 0$ . Since K is lower triangular, we obtain

$$(\partial - q_i) \cdots (\partial - q_1)\phi_i = 0, \tag{7.6}$$

for  $1 \leq i \leq n-1$ .

Let  $\triangle(\cdot, t) := \Gamma(q(\cdot, t))$ , where  $\Gamma$  is given in Definition 2.7. By Theorem 2.11,  $g = K \triangle^{-1}$  is a frame of the Lax pair of the solution u of (2.6) at  $\lambda = 0$ . Note that  $\triangle$  is in  $N_n^+$ , so the first column of g is also  $\phi$ . Use  $g^{-1}g_x = b + u$  to obtain that  $g = (\phi, \phi_x, \dots, \phi_x^{(n-1)})$ .

Let

$$R := (\partial - q_{n-1}) \cdots (\partial - q_1) = \partial^{n-1} - \sum_{i=1}^{n-1} \xi_i \partial^{i-1}.$$

By (7.6),  $R\phi_j = 0$  for all  $1 \le j \le n-1$ . So we have

$$\sum_{i=1}^{n-1} \xi_i \phi_j^{(i-1)} = \phi_j^{(n-1)}, \quad 1 \le j \le n-1.$$

By Cramer's rule,

$$\xi_{n-1} = \frac{\det(\eta, \dots, \eta_x^{(n-4)}, \eta_x^{(n-3)}, \eta_x^{(n-1)})}{\det(\eta, \eta_x, \dots, \eta_x^{(n-2)})},$$
(7.7)

where  $\eta = (\phi_1, \ldots, \phi_{n-1})^t$ . Recall that  $g = (\phi, \ldots, \phi_x^{(n-1)})$ . Let  $g^{ij}$  denote the ij-th entry of  $g^{-1}$ . Then the numerator and denominator of the right hand side of (7.7) are  $-\det(g)g^{n-1,n}$  and  $\det(g)g^{nn}$  respectively. So  $\xi_n = -\frac{g^{n-1,n}}{g^{nn}}$ . By Theorem 3.9,  $\xi_{n-1}$  is a solution of (1.3) with k = 0. Compare the coefficient of  $\partial^{n-1}$  of  $L = (\partial - q_n)R$  to obtain  $q_n = -\xi_{n-1}$ . Hence  $-q_n$  is a solution of (1.3) with k = 0.

#### References

- Adler, M., Moser, J., On a class of polynomials connected with the Korteweg-de Vries Equation, Comm. Math. Phys. 61 (1978), 1–30.
- [2] Adler, M., On the Bäcklund transformation for the Gel'fand-Dickey equations, Comm. Math. Phys. 80(4) (1981), 517–527.
- Beals, R., Sattinger, D.H., Integrable systems and isomonodromy deformations, Phys. D 65(1-2) (1993), 17-47.
- [4] Deift, P., Tomei, C., Trubowitz, E., Inversie scattering and the Boussinesq equation, Comm. Pure Appl. Math. 35(5) (1982), 567–628.
- [5] Dickey, L. A., Soliton equations and Hamiltonian systems, second edition, Advanced Series in Mathematical Physics 26 (2003), World Scientific Publishing Co. Inc., River Edge, NJ.
- [6] Drinfel'd, V.G., Sokolov, V.V., Lie algebras and equations of Korteweg-de Vries type, (Russian) Current problems in mathematics, 24 (1984), 81–180, Itogi Nauki i Tekhniki, Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow.
- [7] Kupershmidt, B.A., Wilson, G., Modifying Lax equations and the second Hamiltonian structure, Invent. Math. 62 (1981), 403–436
- [8] Terng, C.L., Uhlenbeck, K., Bäcklund transformations and loop group actions, Comm. Pure Appl. Math. 53 (2000), 1–75.
- [9] Terng, C.L., Uhlenbeck, K., The  $n \times n$  KdV flows, J. Fixed Point Theory and its Applications, **10** (2011) 37–61.
- [10] Terng, C.L., Uhlenbeck, K., Tau function and Virasoro action for the  $n \times n \ KdV$ hierarchy, arXiv:1406.5240v1.
- [11] Terng, C.L., Wu, Z., N-dimension central affine curve flows, arXiv:1411.2725v1.
- [12] Wahlquist, H., Estabrook, F., Bäcklund transformation for solutions of the Kortewegde Vries equation, Phys. Rev. Lett. **31** (1973), 1386–1390.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT IRVINE, IRVINE, CA 92697-3875. EMAIL: CTERNG@MATH.UCI.EDU

DEPARTMENT OF MATHEMATICS, NINGBO UNIVERSITY, NINGBO, ZHEJIANG, 315211, CHINA. EMAIL: WUZHIWEI@NBU.EDU.CN