Hirota bilinear approach to GUE, NLS, and Painlevé IV

Saburo Kakei

Department of Mathematics, Rikkyo University, 3-34-1 Nishi-ikebukuro, Toshima-ku, Tokyo 171-8501, Japan

Abstract

Tracy and Widom showed that the level spacing function of the Gaussian unitary ensemble is related to a particular solution of the fourth Painlevé equation. We reconsider this problem from the viewpoint of Hirota's bilinear method in soliton theory and present another proof. We also consider the asymptotic behavior of the level spacing function as $s \to \infty$, and its relation to the "Clarkson-McLeod solution" to the Painlevé IV equation.

To the memory of Professor Ryogo Hirota

1 Introduction

Random matrix theory has its origins in mathematical statistics and nuclear physics [27], and has been applied to many fields such as wireless communications [39]. In the paper [35], Tracy and Widom introduced a limiting distribution function associated with the largest eigenvalue of the Gaussian unitary ensemble (GUE). The Tracy-Widom distribution has been applied to many branch of mathematics such as combinatorics and statistics. (For review, see [38].) Furthermore, recent experimental studies reveal that the Tracy-Widom distribution appears in various physical systems such as slow combustion of paper [28], turbulent liquid crystals [33], fiber laser systems [16], evaporating drops of colloidal suspensions [40]. (For a survey of recent experimental studies, see [34].)

The GUE-Tracy-Widom distribution function $F_2(s)$ is given by

$$F_2(s) = \exp\left[-\int_s^\infty (x-s)q^2(x)dx\right],\tag{1.1}$$

where q(x) is the unique solution to a special case of the Painlevé II equation ("Hastings-McLeod solution" [17])

$$\frac{d^2q}{ds^2} = s\frac{dq}{ds} + 2q^3 \tag{1.2}$$

satisfying the boundary condition $q(s) \sim \operatorname{Ai}(s)$ as $s \to \infty$. The function $F_2(s)$ is a scaling limit of the distribution function of the largest eigenvalue of the Gaussian unitary ensemble (GUE). For the ensemble of $N \times N$ random Hermitian matrices, joint distribution of eigenvalues is given by [27]

$$\mathbb{P}_{2,N}(x_1, x_2, \dots, x_N) = \frac{1}{Z_N} \prod_{1 \le i < j \le N} |x_i - x_j|^2 \prod_{i=1}^N e^{-x_i^2},$$
(1.3)

where the normalization constant $1/Z_N$ is given by

$$Z_N = \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \prod_{1 \le i < j \le N} |x_i - x_j|^2 \prod_{i=1}^N e^{-x_i^2} dx_1 \cdots dx_N$$
$$= 2^{-N(N-1)/2} \pi^{N/2} \prod_{j=1}^N j!.$$
(1.4)

We will give a bilinear-theoretical proof of the identity (1.4) in Appendix A.

The distribution function of the largest eigenvalue λ_{max} is then expressed as

$$F_{2,N}(s) := \mathbb{P}_{2,N}(\lambda_{\max} \le s) = \int_{x_1 \le \dots \le x_N \le s} \mathbb{P}_{2,N}(x_1, x_2, \dots, x_N) dx_1 \cdots dx_N$$
$$= \frac{1}{N!} \int_{-\infty}^s \dots \int_{-\infty}^s \mathbb{P}_{2,N}(x_1, x_2, \dots, x_N) dx_1 \cdots dx_N,$$
(1.5)

and the Tracy-Widom distribution function $F_2(s)$ is obtained as a scaling limit of $F_{2,N}(s)$ [14, 35]:

$$F_2(s) = \lim_{N \to \infty} F_{2,N} \left(\sqrt{2N} + \frac{s}{\sqrt{2N^{1/6}}} \right).$$
(1.6)

As discussed in [36], the distribution function $F_{2,N}(s)$ is related to the fourth Painlevé equation

$$\frac{d^2w}{dz^2} = \frac{1}{2w} \left(\frac{dw}{dz}\right)^2 + \frac{3}{2}w^3 + 4zw^2 + 2(z^2 - \alpha)w + \frac{\beta}{w}.$$
(1.7)

Theorem 1 (Tracy-Widom [36]). $R(s) := \{\log F_{2,N}\}'$ is given explicitly by

$$F_{2,N}(s) = \exp\left[-\int_{s}^{\infty} \left\{Nw - \frac{z^{2}}{2}w - \frac{z}{2}w^{2} - \frac{w^{3}}{8} + \frac{1}{8w}\left(\frac{dw}{dz}\right)^{2}\right\} dz\right], \quad (1.8)$$

where w = w(z) is a solution to the Painlevé IV equation with the parameters $\alpha = 2N - 1$ and $\beta = 0$, satisfying the boundary condition

$$w(s) \sim 0 \quad as \quad s \to +\infty. \tag{1.9}$$

The proof of Theorem 1 in [36] is based on the Fredholm determinantal expression,

$$F_{2,N}(s) := \mathbb{P}_{2,N}(\lambda_{\max} \le s) = \det \left[I - K_{2,N} \right]_{L^2(s,\infty)}.$$
(1.10)

The kernel of the integral operator $K_{2,N}$ is given by

$$K_{2,N}(x,z) = \frac{e^{-(x^2+z^2)/2}}{2^N \sqrt{\pi}(N-1)!} \cdot \frac{H_N(x)H_{N-1}(z) - H_{N-1}(x)H_N(z)}{x-z},$$
(1.11)

where $H_k(x)$ is the k-th Hermite polynomials.

In what follows, we present an alternative proof based on Hirota's bilinear method in soliton theory [3, 19, 30]. The solutions to many soliton equations can be expressed in terms of determinants with Wronski-type or Gram-type structure, and the bilinear forms of the equations are reduced to algebraic identities of determinants [13, 19, 20, 30]. We show

that the function $F_{2,N}(s)$ can be obtained from "two-directional" Wronskian solutions [20] to the nonlinear Schrödinger (NLS) equation

$$i\frac{\partial q}{\partial t} = \frac{\partial^2 q}{\partial x^2} + 2|q|^2 q.$$
(1.12)

We also consider the asymptotic behavior of $F_{2,N}(s)$ as $s \to \pm \infty$, and its relation to the "Clarkson-McLeod solution" [6, 12, 21] to the Painlevé IV equation.

2 Nonlinear Schödinger equation and the Painlevé IV

In this section, we review Hirota bilinear formulation of the NLS equation (1.12), and its similarity reduction to the Painlevé IV equation (1.7).

The relation between NLS-type equations and the Painlevé IV (1.7) has been discussed from various perspectives [2, 3, 7, 10, 22, 24, 25, 29]. Among these works, our approach is based mainly on the work of Jimbo and Miwa [22], in which they introduced the " τ function" in a systematic manner.

Hereafter we will forget the complex structure and consider the following coupled system of nonlinear partial differential equations:

$$-\frac{\partial q}{\partial t_2} + \frac{\partial^2 q}{\partial t_1^2} + 2q^2r = 0, \quad \frac{\partial r}{\partial t_2} + \frac{\partial^2 r}{\partial t_1^2} + 2qr^2 = 0.$$
(2.1)

Setting $r = \bar{q}$, $t_1 = x$, $t_2 = it$ recovers the NLS equation (1.12). It is well-known that the NLS equation has infinite number of conserved currents [3]. The first three of the conserved currents are given as

$$\mathcal{J}_1 = qr, \tag{2.2}$$

$$\mathcal{J}_2 = \frac{\partial q}{\partial t_1} r - q \frac{\partial r}{\partial t_1},\tag{2.3}$$

$$\mathcal{J}_3 = \frac{\partial^2 q}{\partial t_1^2} r + q \frac{\partial^2 r}{\partial t_1^2} - 2 \frac{\partial q}{\partial t_1} \frac{\partial r}{\partial t_1} + 2q^2 r^2, \qquad (2.4)$$

which satisfy the continuity equations

$$\frac{\partial \mathcal{J}_1}{\partial t_2} = \frac{\partial \mathcal{J}_2}{\partial t_1}, \quad \frac{\partial \mathcal{J}_2}{\partial t_2} = \frac{\partial \mathcal{J}_3}{\partial t_1}.$$
(2.5)

These equations guarantee the existence of " τ -function" of the NLS equation (2.1) that satisfies [22]

$$\frac{\partial^2}{\partial t_1^2} \log \tau = \mathcal{J}_1, \quad \frac{\partial^2}{\partial t_1 \partial t_2} \log \tau = \mathcal{J}_2, \quad \frac{\partial^2}{\partial t_2^2} \log \tau = \mathcal{J}_3.$$
(2.6)

We introduce $\rho_1 = \rho_1(t_1, t_2)$ and $\rho_2 = \rho_2(t_1, t_2)$ as

$$\rho_1(t_1, t_2) = \tau(t_1, t_2)q(t_1, t_2), \quad \rho_2(t_1, t_2) = \tau(t_1, t_2)r(t_1, t_2), \tag{2.7}$$

which is the same dependent-variable transformation used in [19, 20] to bilinearize (2.1). Then one can rewrite the equations (2.1) and (2.6) as bilinear differential equations of Hirota-type:

$$(D_1^2 + D_2)\tau \cdot \rho_1 = 0, (2.8)$$

$$(D_1^2 + D_2)\rho_2 \cdot \tau = 0, \tag{2.9}$$

$$D_1^2 \tau \cdot \tau = 2\rho_1 \rho_2, \tag{2.10}$$

$$D_1 D_2 \tau \cdot \tau = 2 D_1 \rho_1 \cdot \rho_2, \tag{2.11}$$

$$D_2^2 \tau \cdot \tau = 2D_1^2 \rho_1 \cdot \rho_2, \tag{2.12}$$

where we have used the Hirota differential operators D_1 , D_2 defined by

$$D_1^m D_2^n f(t_1, t_2) \cdot g(t_1, t_2) = \left(\frac{\partial}{\partial t_1} - \frac{\partial}{\partial t_1'}\right)^m \left(\frac{\partial}{\partial t_2} - \frac{\partial}{\partial t_2'}\right)^n f(t_1, t_2) g(t_1', t_2') \bigg|_{\substack{t_1' = t_1 \\ t_2' = t_2}}.$$
 (2.13)

Straightforward calculation shows

Proposition 2. Let $\tau(t_1, t_2)$, $\rho_1(t_1, t_2)$, $\rho_1(t_1, t_2)$ be a solution to the bilinear equations (2.8)–(2.12). Let λ , c_1 , c_2 be constants and assume $\lambda \neq 0$. Define $\tilde{\tau}(t_1, t_2)$, $\tilde{\rho}_1(t_1, t_2)$, $\tilde{\rho}_1(t_1, t_2)$ as

$$\tilde{\tau}(t_1, t_2) = \lambda^{-1 + (c_1 + c_2)/2} \tau(\lambda t_1, \lambda^2 t_2), \qquad (2.14)$$

$$\tilde{\rho}_1(t_1, t_2) = \lambda^{c_1} \rho_1(\lambda t_1, \lambda^2 t_2),$$
(2.15)

$$\tilde{\rho}_2(t_1, t_2) = \lambda^{c_2} \rho_2(\lambda t_1, \lambda^2 t_2).$$
(2.16)

Then $\tilde{\tau}(t_1, t_2)$, $\tilde{\rho}_1(t_1, t_2)$, $\tilde{\rho}_1(t_1, t_2)$ also solve the equations (2.8)–(2.12).

Motivated from Proposition 2, we now impose similarity conditions on $\tau(t_1, t_2)$, $\rho_1(t_1, t_2)$, $\rho_1(t_1, t_2)$:

$$\tau(t_1, t_2) = \lambda^{-1 + (c_1 + c_2)/2} \tau(\lambda t_1, \lambda^2 t_2), \qquad (2.17)$$

$$\rho_1(t_1, t_2) = \lambda^{c_1} \rho_1(\lambda t_1, \lambda^2 t_2), \qquad (2.18)$$

$$\rho_2(t_1, t_2) = \lambda^{c_2} \rho_2(\lambda t_1, \lambda^2 t_2), \tag{2.19}$$

for all $\lambda \in \mathbb{C}^*$, where c_1 and c_2 are constants. Then $q(t_1, t_2)$ and $r(t_1, t_2)$ satisfy

$$q(t_1, t_2) = \lambda^{1 + (c_1 - c_2)/2} q(\lambda t_1, \lambda^2 t_2), \qquad (2.20)$$

$$r(t_1, t_2) = \lambda^{1 + (-c_1 + c_2)/2} r(\lambda t_1, \lambda^2 t_2).$$
(2.21)

Proposition 3 (cf. [22]). Suppose $\tau(t_1, t_2)$, $q(t_1, t_2)$ and $r(t_1, t_2)$ satisfy the differential equations (2.1), (2.6), and the similarity conditions (2.17), (2.20), (2.21). Then one can introduce F(s) as

$$\tau(t_1, t_2) = (2\sqrt{-t_2})^{(2-c_1-c_2)/2} F(s), \quad s = \frac{-t_1}{2\sqrt{-t_2}}.$$
(2.22)

Furthermore, if we define H(s) as

$$H(s) = \frac{d}{ds} \log F(s) = \frac{F'(s)}{F(s)},$$
(2.23)

then H(s) solves the following differential equation,

$$(H'')^2 - 4(H - sH')^2 + 4H' \left\{ (H')^2 - (c_1 - c_2)H' + 2(c_1 + c_2 - 2) \right\} = 0.$$
(2.24)

Proof. Setting $\lambda = 1/(2\sqrt{-t_2})$ and $F(s) = \tau(-s, -1/4)$ in (2.17), we obtain (2.22). Similarly we obtain

$$q(t_1, t_2) = (2\sqrt{-t_2})^{(-2-c_1+c_2)/2}u(s),$$

$$r(t_1, t_2) = (2\sqrt{-t_2})^{(-2+c_1-c_2)/2}v(s)$$
(2.25)

by setting u(s) = q(-s, -1/4) and v(s) = r(-s, -1/4). Substituting (2.25) for (2.1), we have

$$u'' = 2su' + 2u^{2}v + (c_{1} - c_{2} + 2)u,$$

$$v'' = -2sv' + 2uv^{2} + (c_{1} - c_{2} - 2)v.$$
(2.26)

In the same manner, substituting (2.22) and (2.25) for (2.6), we obtain

$$H' + uv = 0, \quad sH' + H + \frac{1}{2}(u'v - uv') = 0,$$

$$4s^{2}H' + 12sH + u''v - 2u'v' + uv'' - 2u^{2}v^{2} + 4(c_{1} + c_{2} - 2) = 0.$$
(2.27)

Eliminating u, v from the equations (2.27), one obtains (2.24).

Denote as $2(\theta_{\infty} \pm \theta_0)$ the roots of a quadratic polynomial $x^2 - (c_1 - c_2)x + 2(c_1 + c_2 - 2)$. Then (2.24) is written as

$$(H'')^{2} - 4(sH' - H)^{2} + 4H'(H' - 2\theta_{\infty} - 2\theta_{0})(H' - 2\theta_{\infty} + 2\theta_{0}) = 0.$$
(2.28)

Setting $\sigma = H - 2t (\theta_{\infty} + \theta_0)$, we obtain the σ -form of the Painlevé IV equation [22]:

$$\left(\sigma''\right)^{2} - 4\left(s\sigma' - \sigma\right)^{2} + 4\sigma'\left(\sigma' + 4\theta_{0}\right)\left(\sigma' + 2\theta_{\infty} + 2\theta_{0}\right) = 0, \qquad (2.29)$$

(cf. [7, 11, 15, 32]).

Proposition 4 ([7, 22]). Define Q = Q(s), P = P(s) as

$$Q(s) = -\frac{\sigma'(s)}{2} = -\frac{H'(s)}{2} + \theta_{\infty} + \theta_0, \qquad (2.30)$$

$$P(s) = \frac{-\sigma''(s) - 2\{s\sigma'(s) - \sigma(s)\}}{2\{\sigma'(s) + 2\theta_{\infty} + 2\theta_{0}\}} = \frac{-H''(s) - 2\{sH'(s) - H(s)\}}{2H'(s)}.$$
 (2.31)

Then H can be expressed in terms of Q and P:

$$H = \frac{2}{P}Q^{2} - \left(P + 2s + \frac{4\theta_{0}}{P}\right)Q + (\theta_{0} + \theta_{\infty})(P + 2s), \qquad (2.32)$$

and we have the Hamiltonian system associated with the Painlevé IV:

$$\frac{dQ}{ds} = P\frac{\partial H}{\partial P} = -\frac{2}{P}Q^2 + \left(-P + \frac{4\theta_0}{P}\right)Q + (\theta_0 + \theta_\infty)P, \qquad (2.33)$$

$$\frac{dP}{ds} = -P\frac{\partial H}{\partial Q} = -4Q + P^2 + 2sP + 4\theta_0.$$
(2.34)

Proof. Straightforward calculation shows that P of (2.31) satisfies

$$H'P^{2} + H''P - (H' - 2\theta_{0} - 2\theta_{\infty})(H' + 2\theta_{0} - 2\theta_{\infty}) = 0.$$
(2.35)

The differential equation

$$\frac{dQ}{ds} = -\frac{2}{P}Q^2 + \left(-P + \frac{4\theta_0}{P}\right)Q + (\theta_0 + \theta_\infty)P \tag{2.36}$$

follows from (2.35). The relation (2.32) follows from (2.30), (2.31) and (2.36). Differentiating (2.32), one obtains (2.34). $\hfill\square$ The following proposition can be verified by straightforward calculation.

Proposition 5. Suppose H = H(s) satisfies (2.28) and define P(s) as (2.31). Then w = P(z) satisfies the Painlevé IV (1.7) with $\alpha = 2\theta_{\infty} - 1$, $\beta = -8\theta_0^2$.

We remark that the Hamiltonian (2.32) can be written in terms of w(s) = P(s):

$$H(s) = \theta_{\infty}w - \frac{s^2}{2}w - \frac{s}{2}w^2 - \frac{1}{8}w^3 + \frac{(w')^2}{8w} - \frac{2\theta_0^2}{w} + 2\theta_{\infty}s.$$
 (2.37)

3 Two-directional Wronskian and GUE

We start considering a class of solution of the bilinear equations (2.8)–(2.12), which is expressed in terms of "two-directional" Wronski determinant:

$$\tau_N(t_1, t_2) = \det \left[\partial_1^{i+j-2} f(t_1, t_2) \right]_{i,j=1}^N.$$
(3.1)

Hereafter we use the abbreviation $\partial_1 = \partial/\partial t_1$, $\partial_2 = \partial/\partial t_2$.

Theorem 6 (cf. [20]). Suppose the function $f(t_1, t_2)$ satisfies

$$\partial_2 f(t_1, t_2) = \partial_1^2 f(t_1, t_2). \tag{3.2}$$

Then the τ -function (3.1) gives a class of solutions for the bilinear equations (2.8)–(2.12) with $\tau = \tau_N(t_1, t_2)$, $\rho_1 = \tau_{N+1}(t_1, t_2)$, $\rho_2 = \tau_{N-1}(t_1, t_2)$.

The equations (2.8), (2.9), (2.10) are reduced cases of the equations (4.10a), (4.10b), (4.10c) of [20], respectively. One can find a proof in §4 of [20] and we omit here. We shall give a proof for the remaining two equations in Appendix B.

Here we choose $f(t_1, t_2)$ as

$$f(t_1, t_2) = \int_{-\infty}^{0} e^{kt_1 + k^2 t_2} dk.$$
(3.3)

This function admits self-similarity as

$$f(\lambda t_1, \lambda^2 t_2) = \int_{-\infty}^0 e^{k\lambda t_1 + k^2\lambda^2 t_2} dk = \int_{-\infty}^0 e^{kt_1 + k^2 t_2} \frac{dk}{\lambda} = \lambda^{-1} f(t_1, t_2)$$
(3.4)

for $\lambda > 0$, and hence the two-directional Wronskian (3.1) also has self-similarity:

$$\tau_N(\lambda t_1, \lambda^2 t_2) = \lambda^{-N^2} \tau_N(t_1, t_2).$$
 (3.5)

Thus the scaling exponent for $\rho_1(t_1, t_2) = \tau_{N+1}(t_1, t_2)$ is $c_1 = (N+1)^2$, and that for $\rho_2(t_1, t_2) = \tau_{N-1}(t_1, t_2)$ is $c_2 = (N-1)^2$. It follows from Proposition 3 that

$$F(s) = \left(2\sqrt{-t_2}\right)^{-(2-c_1-c_2)/2} \tau_N(t_1, t_2) = \left(2\sqrt{-t_2}\right)^{N^2} \tau_N(t_1, t_2)$$
(3.6)

depends only on $s = -t_1/(2\sqrt{-t_2})$. Since $c_1 = (N+1)^2$, $c_2 = (N-1)^2$, H(s) of (2.23) satisfies the σ -form Painlevé IV (2.28) with $\theta_0 = 0$, $\theta_\infty = N$:

$$(H'')^{2} - 4(H - sH')^{2} + 4H' (H' - 2N)^{2} = 0.$$
(3.7)

The two-directional Wronskian τ_N under consideration coincides with the distribution function $F_{2,N}(s)$ as shown below. For this purpose, we assume $t_2 < 0$ and substitute $f(t_1, t_2)$ of (3.3) into (3.1):

$$\tau_{N} = \det \left[\int_{-\infty}^{0} k^{i+j-2} e^{kt_{1}+k^{2}t_{2}} dk \right]_{i,j=1}^{N}$$

$$= \frac{e^{Ns^{2}}}{(-t_{2})^{N^{2}/2}} \det \left[\int_{-\infty}^{s} (k-s)^{i+j-2} e^{-k^{2}} dk \right]_{i,j=1}^{N}$$

$$= \frac{e^{Ns^{2}}}{(-t_{2})^{N^{2}/2}} \det \left[\int_{-\infty}^{s} k^{i+j-2} e^{-k^{2}} dk \right]_{i,j=1}^{N}.$$
(3.8)

On the other hand, applying an integral identity [4, 9, 37]

$$\int \cdots \int \det \left[\phi_j(x_k)\right]_{j,k=1,\dots,N} \det \left[\psi_j(x_k)\right]_{j,k=1,\dots,N} dx_1 \cdots x_N$$
$$= N! \det \left[\int \phi_j(x)\psi_k(x)dx\right]_{j,k=1,\dots,N}$$
(3.9)

to (1.5), one can rewrite $F_{2,N}(s)$ as

$$F_{2,N}(s) = \frac{2^{N(N-1)/2} \pi^{-N/2}}{\prod_{j=1}^{N} j!} \det \left[\int_{-\infty}^{s} x^{j+k-2} e^{-x^2} dx \right]_{j,k=1,\dots,N}.$$
 (3.10)

Comparing (3.6), (3.8), and (3.10), we obtain

$$F(s) = 2^{N(N+1)/2} \pi^{N/2} e^{Ns^2} \left\{ \prod_{j=1}^{N-1} (j!) \right\} F_{2,N}(s),$$
(3.11)

and thus

$$H(s) = 2Ns + R(s), \quad R(s) := \frac{d}{ds} \log F_{2,N}(s).$$
 (3.12)

It follows from (2.37) and (3.12) that

$$\frac{d}{ds}\log F_{2,N}(s) = Nw - \frac{s^2}{2}w - \frac{s}{2}w^2 - \frac{w^3}{8} + \frac{(w')^2}{8w}.$$
(3.13)

Integrating once, we obtain (1.8).

4 Asymptotic behavior

Real solutions to the Painlevé IV equation in the case $\beta = 0$ and real α with the boundary condition (1.9) are called "Clarkson-McLeod solution" [12, 6, 21].

Theorem 7 (Clarkson-McLeod solution [1, 6, 26]). In the case $\beta = 0$ and real α , there exists a unique one-parameter family $w(x;k^2)$ ($k^2 \in \mathbb{R}$) of real solutions of the Painlevé IV equation (1.7) determined by asymptotic condition

$$w(s) \sim k^2 2^{(\alpha+2)/2} s^{\alpha-1} e^{-s^2} \quad as \quad s \to +\infty.$$
 (4.1)

Moreover, in the case $\beta = 0$ and real α , any solutions of the Painlevé IV equation (1.7) with the boundary condition (1.9) coincides with $w(x; k^2)$ for some constant $k^2 \in \mathbb{R}$.

The parameters k, α for the solution of the Painlevé IV associated with $F_{2,N}(s)$ have been evaluated by Tracy and Widom ([36], p.55).

Theorem 8 (Tracy-Widom [36]). $R(s) := \{\log F_{2,N}\}'$ is given explicitly by (1.8) where $w = w(z; k^2)$ is the Clarkson-McLeod solution (Theorem 7) to the Painlevé IV, with the parameters $\alpha = 2N - 1$, $\beta = 0$, and $k^2 = 2^{-3/2} \pi^{-1/2} / (N - 1)!$.

Since no explicit proof of Theorem 8 has been written in [36], we give a proof based on the determinant expression (3.10). Toward this purpose, we consider two integrals

$$I_n := \int_{-\infty}^{\infty} x^n e^{-x^2} dx, \quad J_n := \int_s^{\infty} x^n e^{-x^2} dx \quad (n = 0, 1, 2, ...)$$
(4.2)

and evaluate a determinant

$$T_N(s) := \det\left[\int_{-\infty}^s x^{i+j-2} e^{-x^2} dx\right]_{i,j=1}^N = \det\left[I_{m+n-2} - J_{m+n-2}\right]_{i,j=1}^N$$
(4.3)

for large s. Here J_n can be represented in terms of the incomplete gamma function,

$$\Gamma(a,z) = \int_{z}^{\infty} t^{a-1} e^{-t} dt, \qquad (4.4)$$

as

$$J_n = \frac{1}{2} \Gamma\left((n+1)/2, s^2\right).$$
(4.5)

Using the asymptotic series [5]

$$\Gamma(a,z) \sim z^{a-1} e^{-z} \left[1 + \frac{a-1}{z} + \cdots \right] \quad \left(\text{as } z \to \infty \text{ in } |\arg z| < \frac{3\pi}{2} \right), \tag{4.6}$$

we can estimate asymptotic behavior of $J_n(s)$ as

$$J_n \sim \frac{1}{2} s^{n-1} e^{-s^2} \left[1 + \frac{n-1}{2s^2} + \cdots \right] \quad (\text{as } s \to \infty) \,. \tag{4.7}$$

Applying (4.7) to (4.3), we have

$$T_N(s) \sim \det \left[I_{i+j-2} \right]_{i,j=1}^N - \frac{1}{2} \det \left[I_{i+j-2} \right]_{i,j=1}^{N-1} s^{2N-3} e^{-s^2} \quad (\text{as } s \to \infty).$$
(4.8)

It follows from (1.4) and (3.9) that

$$\det \left[I_{i+j-2} \right]_{i,j=1}^{N} = \frac{1}{N!} Z_N = 2^{-N(N-1)/2} \pi^{N/2} \prod_{j=1}^{N-1} j!, \tag{4.9}$$

and thus

$$T_N(s) \sim \frac{Z_N}{N!} - \frac{1}{2} \frac{Z_{N-1}}{(N-1)!} s^{2N-3} e^{-s^2} \quad (\text{as } s \to \infty).$$
 (4.10)

Applying (3.10), (4.3) and (4.10) to (3.12), we obtain

$$R_N(s) = \frac{d}{ds} \log T_N(s)$$

$$\sim N \frac{Z_{N-1}}{Z_N} s^{2N-2} e^{-s^2} \quad (\text{as } s \to \infty)$$

$$= \frac{2^{N-1}}{\pi^{1/2} (N-1)!} s^{2N-2} e^{-s^2}.$$
(4.11)

Proposition 9. Higher order expansion of R(s) is given as

$$R(s) \sim \frac{2^{N-1}}{\pi^{1/2}(N-1)!} s^{2N-2} e^{-s^2} \times \left[1 - \frac{(N-1)(N-3)}{2s^2} + \frac{(N-1)(N-2)(N^2 - 7N + 15)}{8s^4} + \cdots \right] \quad (as \ s \to \infty).$$

$$(4.12)$$

Proof. We remark that R(s) satisfies the differential equation

$$(R'')^{2} - 4(R - sR')^{2} + 4(R')^{2}(R' + 2N) = 0.$$
(4.13)

Consider the asymptotic expansion

$$R(s) \sim s^{2N-2} e^{-s^2} \left(c_0 + c_1 s^{-1} + c_2 s^{-2} + c_3 s^{-3} + \cdots \right)$$
(4.14)

with $c_0 = 2^{N-1} \pi^{-1/2} / (N-1)!$. Inserting (4.14) into (4.13), one can obtain c_1, c_2, c_3, \ldots , recursively, and the desirous result follows.

Proof of Theorem 8. From (2.31) and (3.12), we have

$$w(s) = \frac{-R'' - 2sR' + 2R}{2(R' + 2N)}.$$
(4.15)

Inserting the asymptotic series (4.12) to (4.15), we have the desirous result.

Appendices

Appendix A: Bilinear-theoretic derivation of (1.4)

In [27], the proof of the integral formula (1.4) is given by using the Selberg integral. Here we give another proof based on the determinant expression

$$Z_N = N! \det \left[I_{i+j-2} \right]_{i,j=1}^N, \quad I_n = \int_{-\infty}^{\infty} x^n e^{-x^2} dx, \tag{A.1}$$

which is a consequence of the identity (3.9). If n is odd then $I_n = 0$, and if n is even, the integral I_n is a special value of the Gamma function:

$$I_{2m} = \int_0^\infty t^{(2m-1)/2} e^{-t} dt = \Gamma\left(m + \frac{1}{2}\right) = \sqrt{\pi} \prod_{j=1}^m \left(m - \frac{1}{2}\right).$$
(A.2)

We recall the Desnanot-Jacobi identity for determinants [8, 19] (or "Lewis Carroll identity", "Dodgson condensation"). We consider a determinant D of degree N and its minors. We denote by $D\begin{bmatrix}i\\j\end{bmatrix}$ the minor determinant obtained by deleting the *i*th row and the *j*th column, and by $D\begin{bmatrix}i_1 & i_2\\j_1 & j_2\end{bmatrix}$ the minor obtained by deleting the i_1 th and i_2 th rows and the j_1 th and j_2 th columns.

Theorem 10 (Desnanot-Jacobi identity [8, 19]). For i_1 , i_2 , j_1 , j_2 that satisfy $1 \le i_1 < i_2 \le N$ and $1 \le j_1 < j_2 \le N$, the following identity holds:

$$D \cdot D \begin{bmatrix} i_1 & i_2 \\ j_1 & j_2 \end{bmatrix} = D \begin{bmatrix} i_1 \\ j_1 \end{bmatrix} D \begin{bmatrix} i_2 \\ j_2 \end{bmatrix} - D \begin{bmatrix} i_1 \\ j_2 \end{bmatrix} D \begin{bmatrix} i_2 \\ j_1 \end{bmatrix}.$$
(A.3)

For M = 0, 1, 2, ..., N = 1, 2, ..., define A_N^M as

$$A_N^M = \det \left[I_{M+i+j-2} \right]_{i,j=1}^N.$$
 (A.4)

From the Desnanot-Jacobi identity (A.3), we have

$$A_{N+2}^{M}A_{N}^{M+2} = A_{N+1}^{M}A_{N+1}^{M+2} - \left(A_{N+1}^{M+1}\right)^{2}.$$
(A.5)

Setting M = 2m and N = 2n, we obtain

$$A_{2n+2}^{2m}A_{2n}^{2m+2} = A_{2n+1}^{2m}A_{2n+1}^{2m+2}$$
(A.6)

where we have used $A_{2n+1}^{2m+1} = 0$. Similarly we have

$$A_{2n+1}^{2m}A_{2n-1}^{2m+2} = A_{2n}^{2m}A_{2n}^{2m+2} - \left(A_{2n}^{2m+1}\right)^2, \tag{A.7}$$

$$A_{2n+2}^{2m-1}A_{2n}^{2m+1} = -\left(A_{2n+1}^{2m}\right)^2,\tag{A.8}$$

$$0 = A_{2n}^{2m-1} A_{2n}^{2m+1} - \left(A_{2n}^{2m}\right)^2.$$
(A.9)

The values of A_N^M (M, N = 0, 1, 2, ...) are determined by the recursion relations (A.6), (A.7), (A.8), (A.9), together with the initial conditions

$$A_0^M = 1, \quad A_1^M = I_M \quad (M = 0, 1, 2, ...).$$
 (A.10)

Proposition 11. Define P(l; m, n) as

$$P(l;m,n) = \begin{cases} 1 & (m=0,n=1,2,\ldots) \\ \prod_{i=1}^{m} \prod_{j=1}^{n} \left(l+i+j-\frac{3}{2}\right) & (m,n=1,2,\ldots) \end{cases}$$
(A.11)

Then A^M_N is expressed in terms of P(l;m,n) as

$$A_{2n}^{2m} = \pi^n P(0;m,n) P(1;m,n) \prod_{j=1}^{2n-1} \frac{j!}{2},$$
(A.12)

$$A_{2n+1}^{2m} = \pi^{(2n+1)/2} P(0;m,n+1) P(1;m,n) \prod_{j=1}^{2n} \frac{j!}{2},$$
(A.13)

$$A_{2n}^{2m+1} = (-\pi)^n P(1;m,n)^2 \prod_{j=0}^{n-1} \left\{ \frac{(2j+1)!}{2^{2j+1}} \right\}^2,$$
(A.14)

$$A_{2n+1}^{2m+1} = 0 \qquad (m, n = 0, 1, 2, \ldots).$$
(A.15)

Proof. It is enough to show that (A.12), (A.13), (A.14) satisfy the recursion relations (A.6), (A.7), (A.8), (A.9), and the initial conditions (A.10), that can be checked by straightforward calculation.

The formula (1.4) follows from a special case of Proposition 11:

$$Z_N = N! A_N^0 = N! \cdot \pi^{N/2} \prod_{j=1}^{N-1} \frac{j!}{2} = 2^{-N(N-1)/2} \pi^{N/2} \prod_{j=1}^N j!.$$
(A.16)

Appendix B: Proof of Theorem 6

In this appendix, we show that the two-directional Wronskian (3.1) actually satisfies the bilinear equations (2.11) and (2.12) with $\tau = \tau_N(t_1, t_2)$, $\rho_1 = \tau_{N+1}(t_1, t_2)$, $\rho_2 = \tau_{N-1}(t_1, t_2)$. We first introduce the following notation, which is a generalization of the abbreviated notation due to Freeman and Nimmo [13, 19, 20, 30]:

$$\begin{vmatrix} m_1, m_2, \dots, m_N \\ n_1, n_2, \dots, n_N \end{vmatrix} := \det \left[\partial_1^{m_i + n_j} f(t_1, t_2) \right]_{i,j=1}^N.$$
(B.1)

The τ -function (3.1) and its derivatives are written as

$$\tau_N = \begin{vmatrix} 0, \dots, N-1 \\ 0, \dots, N-1 \end{vmatrix},\tag{B.2}$$

$$\partial_1 \tau_N = \begin{vmatrix} 0, \dots, N-1 \\ 0, \dots, N-2, N \end{vmatrix} = \begin{vmatrix} 0, \dots, N-2, N \\ 0, \dots, N-1 \end{vmatrix},$$
(B.3)

$$\partial_1^2 \tau_N = \begin{vmatrix} 0, \dots, N-2, N \\ 0, \dots, N-2, N \end{vmatrix} = \begin{vmatrix} 0, \dots, N-3, N-1, N \\ 0, \dots, N-1 \end{vmatrix} + \begin{vmatrix} 0, \dots, N-2, N+1 \\ 0, \dots, N-1 \end{vmatrix}.$$
(B.4)

Using (3.2), we have

$$\partial_2 \tau_N = - \begin{vmatrix} 0, \dots, N-3, N-1, N \\ 0, \dots, N-1 \end{vmatrix} + \begin{vmatrix} 0, \dots, N-2, N+1 \\ 0, \dots, N-1 \end{vmatrix},$$
(B.5)

$$\partial_1 \partial_2 \tau_N = - \begin{vmatrix} 0, \dots, N-3, N-1, N \\ 0, \dots, N-2, N \end{vmatrix} + \begin{vmatrix} 0, \dots, N-2, N+1 \\ 0, \dots, N-2, N \end{vmatrix},$$
(B.6)

$$\partial_2^2 \tau_N = \begin{vmatrix} 0, \dots, N-3, N-1, N \\ 0, \dots, N-3, N-1, N \end{vmatrix} - 2 \begin{vmatrix} 0, \dots, N-3, N-1, N \\ 0, \dots, N-2, N+1 \end{vmatrix} + \begin{vmatrix} 0, \dots, N-2, N+1 \\ 0, \dots, N-2, N+1 \end{vmatrix}.$$
 (B.7)

Using the differential formulas (B.3) and the identity (A.3), we obtain

$$(\partial_{1}\tau_{N+1})\tau_{N-1} = \begin{vmatrix} 0, \dots, N-1, N+1 \\ 0, \dots, N-2 \end{vmatrix} \begin{vmatrix} 0, \dots, N-2 \\ 0, \dots, N-2, N+1 \\ 0, \dots, N-2, N \end{vmatrix} \begin{vmatrix} 0, \dots, N-1 \\ 0, \dots, N-1 \end{vmatrix} \begin{vmatrix} 0, \dots, N-1 \\ 0, \dots, N-1 \end{vmatrix} \begin{vmatrix} 0, \dots, N-1 \\ 0, \dots, N-2, N \end{vmatrix} | \begin{vmatrix} 0, \dots, N-1 \\ 0, \dots, N-2, N \end{vmatrix} ,$$
(B.8)
$$\tau_{N+1} (\partial_{1}\tau_{N-1}) = \begin{vmatrix} 0, \dots, N \\ 0, \dots, N-1 \end{vmatrix} \begin{vmatrix} 0, \dots, N-3, N-1 \\ 0, \dots, N-2 \end{vmatrix} \\ = \begin{vmatrix} 0, \dots, N-3, N-1, N \\ 0, \dots, N-2, N \end{vmatrix} \begin{vmatrix} 0, \dots, N-1 \\ 0, \dots, N-1 \end{vmatrix} \begin{vmatrix} 0, \dots, N-1 \\ 0, \dots, N-2, N \end{vmatrix}$$
(B.9)

It is straightforward to show that $\tau = \tau_N$, $\rho_1 = \tau_{N+1}$, $\rho_2 = \tau_{N-1}$ solve (2.11) by using (B.8), (B.9), and the differential relations (B.3), (B.5), (B.6).

We use the Desnanot-Jacobi identity (A.3) again to obtain

$$\left(\partial_{1}^{2}\tau_{N+1}\right)\tau_{N-1} = \begin{vmatrix} 0, \dots, N-1, N+1 \\ 0, \dots, N-2 \end{vmatrix} \begin{vmatrix} 0, \dots, N-2, N+1 \\ 0, \dots, N-2 \end{vmatrix} \begin{vmatrix} 0, \dots, N-2, N+1 \\ 0, \dots, N-1 \end{vmatrix} \begin{vmatrix} 0, \dots, N-1 \\ 0, \dots, N-1 \end{vmatrix} \begin{vmatrix} 0, \dots, N-1 \\ 0, \dots, N-1 \end{vmatrix} \begin{vmatrix} 0, \dots, N-1 \\ 0, \dots, N-2, N+1 \\ 0, \dots, N-2, N+1 \end{vmatrix} \begin{vmatrix} 0, \dots, N-1 \\ 0, \dots, N-2, N+1 \\ 0, \dots, N-2, N+1 \end{vmatrix} \end{vmatrix},$$
(B.10)
$$\tau_{N+1} \left(\partial_{1}^{2}\tau_{N-1}\right) = \begin{vmatrix} 0, \dots, N \\ 0, \dots, N-3, N-1 \\ 0, \dots, N-3, N-1 \\ 0, \dots, N-3, N-1, N \end{vmatrix} \begin{vmatrix} 0, \dots, N-1 \\ 0, \dots, N-3, N-1 \\ 0, \dots, N-3, N-1, N \end{vmatrix} \begin{vmatrix} 0, \dots, N-1 \\ 0, \dots, N-3, N-1 \\ 0, \dots, N-3, N-1 \end{vmatrix} \end{vmatrix} (0, \dots, N-3, N-1, N) \end{vmatrix},$$
(B.11)
$$\left(\partial_{1}\tau_{N+1}\right) \left(\partial_{1}\tau_{N-1}\right) = \begin{vmatrix} 0, \dots, N-1, N+1 \\ 0, \dots, N-1 \\ 0, \dots, N-1 \end{vmatrix} \begin{vmatrix} 0, \dots, N-2 \\ 0, \dots, N-3, N-1 \\ 0, \dots, N-3, N-1 \end{vmatrix} \end{vmatrix} = \begin{vmatrix} 0, \dots, N-2, N+1 \\ 0, \dots, N-3, N-1 \\ 0, \dots, N-3, N-1 \end{vmatrix} \begin{vmatrix} 0, \dots, N-2 \\ 0, \dots, N-3, N-1 \\ 0, \dots, N-3, N-1 \end{vmatrix} \end{vmatrix} (0, \dots, N-1) \\ = \begin{vmatrix} 0, \dots, N-2, N+1 \\ 0, \dots, N-1 \\ 0, \dots, N-1 \end{vmatrix} \begin{vmatrix} 0, \dots, N-1 \\ 0, \dots, N-1 \\ 0, \dots, N-3, N-1, N \end{vmatrix} (0, \dots, N-1) \\ = \begin{vmatrix} 0, \dots, N-2, N+1 \\ 0, \dots, N-1 \\ \end{vmatrix} (0, \dots, N-3, N-1, N) \end{vmatrix} (0, \dots, N-1) \\ (0, \dots, N-3, N-1, N) \end{vmatrix} (0, \dots, N-1) \end{vmatrix}$$
(B.12)

These relations give (2.12) with $\tau = \tau_N$, $\rho_1 = \tau_{N+1}$, and $\rho_2 = \tau_{N-1}$.

Appendix C: Bilinearization of the Painlevé IV

As in Proposition 3, under the similarity conditions (2.17)–(2.19), one can introduce F(s) as (2.22) and $G_1(s)$, $G_2(s)$ as

$$\rho_1(t_1, t_2) = (2\sqrt{-t_2})^{-c_1} G_1(s), \quad \rho_2(t_1, t_2) = (2\sqrt{-t_2})^{-c_2} G_2(s).$$
(C.1)

Bilinear equations for F(s), $G_1(s)$ and $G_2(s)$ follow from (2.8)–(2.12):

$$\left(D_s^2 + 2sD_s - c_1 + c_2 - 2\right)F \cdot G_1 = 0, \tag{C.2}$$

$$\left(D_s^2 + 2sD_s - c_1 + c_2 + 2\right)G_2 \cdot F = 0, \tag{C.3}$$

$$D_s^2 F \cdot F = G_1 G_2, \tag{C.4}$$

$$\left(sD_s^2 + \frac{d}{ds}\right)F \cdot F = D_sG_1 \cdot G_2,\tag{C.5}$$

$$\left(2s^2D_s^2 + 4(c_1 + c_2 - 2) + 6s\frac{d}{ds}\right)F \cdot F = D_s^2G_1 \cdot G_2.$$
(C.6)

These give a bilinearization of the Painlevé IV, however, which are different from that used in [18, 23, 31].

Acknowledgments

The author acknowledges Takashi Imamura, Makoto Katori, and Tomohiro Sasamoto for discussions on the Tracy-Widom distributions. Thanks are also due to Kenji Kajiwara

and Ralph Willox for information on bilinear forms of the Painlevé equations. This work is partially supported by JSPS Grant-in-Aid for Scientific Research No. 23540252.

References

- A.S. Abdullaye, Justification of asymptotic formulas for the fourth Painlevé equation, Stud. Appl. Math. 99 (1997), 255–283.
- [2] M.J. Ablowitz, A. Ramani and H. Segur, A connection between nonlinear evolution equations and ordinary differential equations of P-type. II, J. Math. Phys. 21 (1980), 1006–1015.
- [3] M.J. Ablowitz and H. Segur, Solitons and the inverse scattering transform, SIAM, Philadelphia, 1981.
- [4] C. Andréief, Note sur une relation les intégrales définies des produits des fonctions, Mém. de la Soc. Sci., Bordeaux 2 (1883), 1–14.
- [5] P.J. Davis, Gamma function and related functions, In Handbook of mathematical functions with formulas: graphs, and mathematical tables, M. Abramowitz and I.A. Stegun (eds.), Dover, New York, 1965.
- [6] A.P. Bassom, P.A. Clarkson, A.C. Hicks and J.B. McLeod, Integral equations and exact solutions for the fourth Painlevé equation, *Proc. R. Soc. Lond.* A437 (1992), 1–24.
- [7] A.P. Bassom, P.A. Clarkson and A.C. Hicks, On the application of solutions of the fourth Painlevé equation to various physically motivated nonlinear partial differential equations, Adv. Differential Equations, 1 (1996), 175–198.
- [8] D.M. Bressoud, *Proofs and confirmations: the story of the alternating sign matrix conjecture*, MAA: Spectrum Series, Cambridge University Press, 1999.
- [9] N.G. de Bruijn, On some multiple integrals involving determinants, J. Indian Math. Soc. 19 (1955), 133–151.
- [10] M. Can, On the relation between nonlinear Schrödinger equation and Painlevé IV equation, Il Nuovo Cimento 106 (1991), 205–207.
- [11] P.A. Clarkson, The fourth Painlevé transcendent, In Differential Algebra and Related Topics II, Li Guo and W.Y. Sit (eds.), World Scientific, Singapore (2008). Also available at http://kar.kent.ac.uk/id/eprint/23090
- [12] P.A. Clarkson and J.B. McLeod, Integral equations and connection formulae for the Painlevé equations, In *Painlevé Transcendents: Their Asymptotics and Physical Applications*, P. Winternitz and D. Levi (eds.), NATO Adv. Sci. Inst. Ser. B Phys., vol. 278, Plenum, New York, 1992, pp. 131.
- [13] N.C. Freeman and J.J.C. Nimmo, Soliton solutions of the Korteweg-de Vries and Kadomtsev-Petviashvili equations: the Wronskian technique, *Phys. Lett. A* 95 (1983), 1–3.

- [14] P.J. Forrester, The spectrum edge of random matrix ensembles, Nuclear Physics B402 (1993), 709–728.
- [15] P.J. Forrester and N.S. Witte, Application of the τ -function theory of Painlevé equations to random matrices: PIV, PII and the GUE, *Commun. Math. Phys.* **219** (2001), 357–398.
- [16] M. Fridman, R. Pugatch, M. Nixon, A.A. Friesem, and N. Davidson, Measuring maximal eigenvalue distribution of Wishart random matrices with coupled lasers, *Phys. Rev. E* 85 (2012), R020101.
- [17] S.P. Hastings and J.B. McLeod, A boundary value problem associated with the second Painlevé transcendent and the Korteweg-de Vries equation, Arch. Rat. Mech. Anal. 73 (1980), 31–51.
- [18] J. Hietarinta and R. Willox, On the bilinear forms of Painlevé equations, in *Bilinear Integrable Systems: from Classical to Quantum, Continuous to Discrete*, L.D. Fadeev and P. van Moerbeke (eds.), Springer, 2006, pp. 375–390.
- [19] R. Hirota, The direct method in soliton theory, Cambridge Tracts in Mathematics, Vol. 155, Cambridge University Press, Cambridge, 2004.
- [20] R. Hirota, Y. Ohta and J. Satsuma, Wronskian structures of solutions for soliton equations, Prog. Theor. Phys. Supplement 94 (1988), 59–72.
- [21] A.R. Its and A.A. Kapaev, Connection formulae for the fourth Painlevé transcendent; Clarkson-McLeod solution, J. Phys. A: Math. Gen. 31 (1998), 4073–4113.
- [22] M. Jimbo and T. Miwa, Monodromy preserving deformations of linear ordinary differential equations with rational coefficients II, *Physica* 2D (1981), 407–448.
- [23] K. Kajiwara and Y. Ohta, Determinant structure of the rational solutions for the Painlevé IV equation, J. Phys. A: Math. Gen. 31 (1998), 2431–2446.
- [24] S. Kakei and T. Kikuchi, Affine Lie group approach to a derivative nonlinear Schrödinger equation and its similarity reduction, *International Mathematics Re*search Notices 2004:78 (2004), 4181–4209.
- [25] A.V. Kitaev, Self-similar solutions of the modified nonlinear Schrödinger equation, *Theor. Math. Phys.* 64 (1985), 878-894.
- [26] Y. Lu, On the asymptotic representation of the solutions to the fourth general Painlevé equation, Int. J. Math. Math. Sci. 2003(13) (2003), 845–851.
- [27] M.L. Mehta, Random matrices, 3rd edition, Elsevier, 2004.
- [28] L. Miettinen, M. Myllys, J. Merikoski and J. Timonen, Experimental determination of KPZ height-fluctuation distributions, *Eur. Phys. J.* B46 (2005), 55–60.
- [29] R. Nakach, Self-similar solutions of nonlinear evolution equations of physical significance, in *Plasma Physics: Nonlinear Theory and Experiments*, H. Wilhelmsson (ed.), Nobel Foundation Symposia, Vol. 36, Springer Science+Buisiness Media, New York, 1977, pp. 456–474.

- [30] J.J.C. Nimmo and J. Zhao, Determinant and Pfaffian solutions of soliton equations, *Phys. Scr.* 89 (2014), 038005.
- [31] M. Noumi and Y. Yamada, Symmetries in the fourth Painlevé equation and Okamoto polynomials, Nagoya Math. J. 153 (1999), 53–86.
- [32] K. Okamoto, Studies on the Painlevé equations III. Second and fourth Painlevé equations, P_{II} and P_{IV}, Math. Ann. 275 (1986), 221–255.
- [33] K.A. Takeuchi and M. Sano, Universal fluctuations of growing interfaces: evidence in turbulent liquid crystals, *Phys. Rev. Lett.* **104** (2010), 230601.
- [34] K.A. Takeuchi, Experimental approaches to universal out-of-equilibrium scaling laws: turbulent liquid crystal and other developments, J. Stat. Mech. (2014), P01006
- [35] C.A. Tracy and H. Widom, Level-spacing distributions and the Airy kernel, Commun. Math. Phys. 159 (1994), 151–174.
- [36] C.A. Tracy and H. Widom, Fredholm determinants, differential equations and matrix models, *Commun. Math. Phys.* 163 (1994), 33–72.
- [37] C.A. Tracy and H. Widom, Correlation functions, cluster functions and spacing distributions for random matrices, J. Stat. Phys. 92 (1998), 809–835.
- [38] C.A. Tracy and H. Widom, Distribution functions for largest eigenvalues and their applications, in *Proceedings of the International Congress of Mathematicians*, Vol. I (Beijing, 2002), Higher Education Press, Beijing, 2002, pp. 587–596.
- [39] A.M. Tulino and S. Verdu, Random matrix theory and wireless communications (Foundations and Trends in Communications and Information Theory) Now Publishers, 2004.
- [40] P.J. Yunker, M.A. Lohr, T. Still, A. Borodin, D.J. Durian, and A.G. Yodh, Effects of particle shape on growth dynamics at edges of evaporating drops of colloidal suspensions, *Phys. Rev. Lett.* **110** (2013), 035501.