# A GENERALIZED INTENSITY BASED FRAMEWORK FOR SINGLE-NAME CREDIT RISK

FRANK GEHMLICH AND THORSTEN SCHMIDT

Abstract. The intensity of a default time is obtained by assuming that the default indicator process has an absolutely continuous compensator. Here we drop the assumption of absolute continuity with respect to the Lebesgue measure and only assume that the compensator is absolutely continuous with respect to a general  $\sigma$ -finite measure. This allows for example to incorporate the Merton-model in the generalized intensity based framework. An extension of the Black-Cox model is also considered. We propose a class of generalized Merton models and study absence of arbitrage by a suitable modification of the forward rate approach of Heath-Jarrow-Morton (1992). Finally, we study affine term structure models which fit in this class. They exhibit stochastic discontinuities in contrast to the affine models previously studied in the literature.

## 1. INTRODUCTION

The two most common approaches to credit risk modelling are the *structural* approach, pioneered in the seminal work of Merton [\[23\]](#page-13-0), and the reduced-form approach which can be traced back to early works of Jarrow, Lando, and Turnbull [\[18,](#page-12-0) [22\]](#page-12-1) and to [\[1\]](#page-12-2).

The default of a company happens when the company is not able to meet its obligations. In many cases the debt structure of a company is known to the public, such that default happens with positive probability at times which are known a priori. This, however, is excluded in the intensity-based framework and it is the purpose of this article to put forward a generalisation which allows to incorporate such effects. Examples in the literature are, e.g., structural models like [\[23\]](#page-13-0) and [\[13,](#page-12-3) [14\]](#page-12-4). The recently missed coupon payment by Argentina is an example for such a credit event as well as the default of Greece on the [1](#page-0-0)st of July<sup>1</sup>.

It is a remarkable observation of [\[2\]](#page-12-5) that it is possible to extend the reduced-form approach beyond the class of intensity-based models. The authors study a class of first-passage time models under a filtration generated by a Brownian motion and show its use for pricing and modelling credit risky bonds. Our goal is to start with even weaker assumptions on the default time and to allow for jumps in the compensator of the default time at deterministic times. From this general viewpoint it turns out, surprisingly, that previously used HJM approaches lead to arbitrage: the whole term structure is absolutely continuous and can not compensate for points in time bearing a positive default probability. We propose a suitable extension with an additional term allowing for discontinuities in the term structure at certain random times and derive precise drift conditions for an appropriate no-arbitrage condition. The related article [\[12\]](#page-12-6) only allows for the special case of finitely many risky times, an assumption which is dropped in this article.

<span id="page-0-0"></span>Date: November 19, 2015.

<sup>1</sup>Argentina's missed coupon payment on \$29 billion debt was voted a credit event by the International Swaps and Derivatives Association, see the announcements in [\[16\]](#page-12-7) and [\[24\]](#page-13-1). Regarding the failure of 1.5 Billon EUR of Greece on a scheduled debt repayment to the International Monetary fund, see e.g. [\[9\]](#page-12-8).

The structure of the article is as follows: in Section [2,](#page-1-0) we introduce the general setting and study drift conditions in an extended HJM-framework which guarantee absence of arbitrage in the bond market. In Section [3](#page-8-0) we study a class of affine models which are stochastically discontinuous. Section [4](#page-11-0) concludes.

#### 2. A general account on credit risky bond markets

<span id="page-1-0"></span>Consider a filtered probability space  $(\Omega, \mathcal{A}, \mathbb{G}, P)$  with a filtration  $\mathbb{G} = (\mathscr{G}_t)_{t \geq 0}$  (the general filtration) satisfying the usual conditions, i.e. it is right-continuous and  $\mathscr{G}_0$  contains the Pnullsets  $N_0$  of A. Throughout, the probability measure P denotes the objective measure. As we use tools from stochastic analysis, all appearing filtrations shall satisfy the usual conditions. We follow the notation from [\[17\]](#page-12-9) and refer to this work for details on stochastic processes which are not laid out here.

The filtration G contains all available information in the market. The default of a company is public information and we therefore assume that the default time  $\tau$  is a G-stopping time. We denote the *default indicator process H* by

$$
H_t = \mathbb{1}_{\{t \geq \tau\}}, \qquad t \geq 0,
$$

such that  $H_t = \mathbb{1}_{\llbracket \tau, \infty \rrbracket}(t)$  is a right-continuous, increasing process. We will also make use of the *survival process*  $1 - H = 1_{[0, \tau]}$ . The following remark recalls the essentials of the well-known intensity based approach.

Remark 2.1 (The intensity-based approach). The intensity-based approach consists in two steps: first, denote by  $\mathbb{H} = (\mathcal{H}_t)_{t\geq0}$  the filtration generated by the default indicator,  $\mathscr{H}_t = \sigma(H_s : 0 \leq s \leq t) \vee N_0$ , and assume that there exists a sub-filtration F of G, i.e.  $\mathscr{F}_t \subset \mathscr{G}_t$  holds for all  $t \geq 0$  such that

<span id="page-1-2"></span>
$$
\mathscr{G}_t = \mathscr{F}_t \vee \mathscr{H}_t, \qquad t \geq 0.
$$

Viewed from this perspective,  $\mathbb G$  is obtained from the default information  $\mathbb H$  by a *progressive* enlargement<sup>[2](#page-1-1)</sup> with the filtration  $\mathbb{F}$ . This assumption opens the area for the largely developed field of enlargements of filtration with a lot of powerful and quite general results.

Second, the following key assumption specifies the default intensity: assume that there is an F-progressive process  $\lambda$ , such that

(2) 
$$
P(\tau > t | \mathscr{F}_t) = \exp\left(-\int_0^t \lambda_s ds\right), \quad t \geq 0.
$$

It is immediate that the inclusion  $\mathscr{F}_t \subset \mathscr{G}_t$  is strict under existence of an intensity, i.e.  $\tau$  is not a F-stopping time. Arbitrage-free pricing can be achieved via the following result: Let Y be a non-negative random variable. Then, for all  $t \geq 0$ ,

$$
E[\mathbb{1}_{\{\tau>t\}}Y|\mathscr{G}_t] = \mathbb{1}_{\{\tau>t\}}e^{\int_0^t \lambda_s ds}E[\mathbb{1}_{\{\tau>t\}}Y|\mathscr{F}_t].
$$

Of course, this results hold also when a pricing measure Q is used instead of P. For further literature and details we refer for example to [\[11\]](#page-12-10), Chapter 12, and to [\[3\]](#page-12-11).

<span id="page-1-1"></span><sup>&</sup>lt;sup>2</sup>Note that here  $G$  is right-continuous and *P*-complete by assumption which is a priori not guaranteed by [\(1\)](#page-1-2). One can, however, use the right-continuous extension and we refer to [\[15\]](#page-12-12) for a precise treatment and for a guide to the related literature.

2.1. The generalized intensity-based framework. The default indicator process  $H$  is a bounded, cádlág and increasing process, hence a submartingale of class  $(D)$ , that is the family  $(X_T)$  over all stopping times T is uniformly integrable. By the Doob-Meyer decomposition<sup>[3](#page-2-0)</sup>, the process

$$
(3) \t\t M_t = H_t - \Lambda_t, \quad t \geq 0
$$

is a true martingale where  $\Lambda$  denotes the dual F-predictable projection, also called compensator, of H. As 1 is an absorbing state,  $\Lambda_t = \Lambda_{t \wedge \tau}$ . To keep the arising technical difficulties at a minimum, we assume that there is an increasing process A such that

<span id="page-2-1"></span>(4) 
$$
\Lambda_t = \int_0^{t \wedge \tau} \lambda_s dA(s), \quad t \geq 0,
$$

with a non-negative and predictable process  $\lambda$ . The process  $\lambda$  is called *generalized intensity* and we refer to Chapter VIII.4 of [\[5\]](#page-12-13) for a more detailed treatment of generalized intensities (or, equivalently, dual predictable projections) in the context of point processes.

Note that with  $\Delta M \leq 1$  we have that  $\Delta \Lambda = \lambda_s \Delta A(s) \leq 1$ . Whenever  $\lambda_s \Delta A(s) > 0$ , there is a positive probability that the company defaults at time  $s$ . We call such times risky times, i.e. predictable times having a positive probability of a default occurring right at that time. Note that under our assumption [\(4\)](#page-2-1), all risky times are deterministic. The relationship between  $\Delta\Lambda(s)$  and the default probability at time s will be clarified in Example [3.1.](#page-8-1)

2.2. An extension of the HJM-approach. A credit risky bond with maturity  $T$  is a contingent claim promising to pay one unit of currency at T. The price of the bond with maturity T at time  $t \leq T$  is denoted by  $P(t,T)$ . If no default occurred prior to or at T we have that  $P(T, T) = 1$ . We will consider zero recovery, i.e. the bond loses its total value at default, such that  $P(t, T) = 0$  on  $\{t \geq \tau\}$ . The family of stochastic processes  $\{(P(t,T)_{0\leq t\leq T}), T\geq 0\}$  describes the evolution of the term structure  $T \mapsto P(., T)$  over time.

Besides the bonds there is a *numéraire*  $X^0$ , which is a strictly positive, adapted process. We make the weak assumption that  $\log X^0$  is absolutely continuous, i.e.  $X_t^0 = \exp$ pr<br>et  $\int_0^t r_s ds$ with a progressively measurable process  $r$ , called the short-rate. For practical applications one would use the overnight index swap  $(OIS)$  rate for constructing such a numéraire.

The aim of the following is to extend the HJM approach in an appropriate way to the generalized intensity-based framework in order to obtain arbitrage-free bond prices. First approaches in this direction were  $[25, 8]$  $[25, 8]$  $[25, 8]$  and a rich source of literature is again  $[3]$ . Absence of arbitrage in such an infinite dimensional market can be described in terms of no asymptotic free lunch (NAFL) or the more economically meaningful no asymptotic free lunch with vanishing risk, see  $[21]$  and  $[6]$ .

Consider a pricing measure  $Q^* \sim P$ . Our intention is to find conditions which render  $Q^*$  an equivalent local martingale measure. In the following, only occasionally the measure P will be used, such that from now on, all appearing terms (like martingales, almost sure properties, etc.) are to be considered with respect to  $Q^*$ .

To ensure that the subsequent analysis is meaningful, we make the following technical assumption.

<span id="page-2-0"></span> ${}^{3}$ See [\[20\]](#page-12-17), Theorem 1.4.10.

<span id="page-3-1"></span>**Assumption 2.1.** The generalized default intensity  $\lambda$  is non-negative, predictable and Aintegrable on  $[0, T^*]$ :  $\mathbf{r}^T$ 

$$
r^* \lambda_s dA(s) < \infty, \quad Q^* \text{-a.s.}
$$

Moreover, A has vanishing singular part, i.e.

<span id="page-3-0"></span>(5) 
$$
A(t) = t + \sum_{0 < s \leq t} \Delta A(s).
$$

0

The representation [\(5\)](#page-3-0) of A is without loss of generality: indeed, if the continuous part  $A<sup>c</sup>$ The representation (5) of A is without loss of generality: indeed, if the continuous part  $A^c$  is absolutely continuous, i.e.  $A^c(t) = \int_0^t a(s)ds$ , replacing  $\lambda_s$  by  $\lambda_s a(s)$  gives the compensator of H with respect to  $\tilde{A}$  whose continuous part is t.

Next, we aim at building an arbitrage-free framework for bond prices. In the generalized intensity-based framework, the (HJM) approach does allow for arbitrage opportunities at risky times. We therefore consider the following generalization: consider a σ-finite (deterministic) measure  $\nu$ . We could be general on  $\nu$ , allowing for an absolutely continuous, a singular continuous and a pure-jump part. However, for simplicity, we leave the singular continuous part aside and assume that

$$
\nu = \nu^{ac} + \nu^d
$$

where  $\nu^{ac}(ds) = ds$  and  $\nu^d$  distributes mass only to points, i.e.  $\nu^d(A) = \sum_{i \geq 1} w_i \delta_{u_i}(A)$ , for  $0 < u_1 < u_2 < \dots$  and positive weights  $w_i > 0$ ,  $i \geq 1$ ; here  $\delta_u$  denotes the Dirac measure at

*u*. Moreover, we assume that defaultable bond prices are given by  
\n
$$
P(t,T) = \mathbb{1}_{\{\tau > t\}} \exp\left(-\int_t^T f(t,u)\nu(du)\right)
$$
\n(6) 
$$
= \mathbb{1}_{\{\tau > t\}} \exp\left(-\int_t^T f(t,u)du - \sum_{i \geq 1} \mathbb{1}_{\{u_i \in (t,T]\}} w_i f(t,u_i)\right), \quad 0 \leq t \leq T \leq T^*.
$$

<span id="page-3-3"></span>The sum in the last line gives the extension over the (HJM) approach which allows us to deal with risky times in an arbitrage-free way.

The family of processes  $(f(t,T))_{0\leq t\leq T}$  for  $T \in [0, T^*]$  are assumed to be Itô processes satisfying

<span id="page-3-4"></span>(7) 
$$
f(t,T) = f(0,T) + \int_0^t a(s,T)ds + \int_0^t b(s,T) \cdot dW_s
$$

with an *n*-dimensional  $Q^*$ -Brownian motion W.

Denote by  $\beta$  the Borel  $\sigma$ -field over  $\mathbb R$ .

<span id="page-3-2"></span>Assumption 2.2. We require the following technical assumptions:

(i) the initial forward curve is measurable, and integrable on  $[0, T^*]$ :

$$
\int_0^{T^*} |f(0, u)| < \infty, \qquad Q^* \text{-a.s.},
$$

(ii) the *drift parameter*  $a(\omega, s, t)$  is R-valued  $\mathcal{O}\otimes\mathcal{B}$ -measurable and integrable on  $[0, T^*]$ :

$$
\int_0^{T^*} \int_0^{T^*} |a(s, u)| ds \, \nu(du) < \infty, \quad Q^* \text{-a.s.},
$$

(iii) the volatility parameter  $b(\omega, s, t)$  is  $\mathbb{R}^n$ -valued,  $\mathcal{O}\otimes\mathcal{B}$ -measurable, and

$$
\sup_{s,t\leqslant T^*}\parallel b(s,t)\parallel<\infty,\quad Q^*\text{-a.s.}
$$

(iv) it holds that

$$
0 \leq \lambda(u_i)\Delta A(u_i) < w_i, \quad i \geq 1.
$$

Set

(8)  
\n
$$
\bar{a}(t,T) = \int_{t}^{T} a(t,u)\nu(du),
$$
\n
$$
\bar{b}(t,T) = \int_{t}^{T} b(t,u)\nu(du),
$$
\n
$$
H'(t) = \int_{0}^{t} \lambda_{s}ds - \sum_{u_{i} \leq t} \log \left(\frac{w_{i} - \lambda_{u_{i}}\Delta A(u_{i})}{w_{i}}\right)
$$

The following proposition gives the desired drift condition in the generalized Merton models.

.

<span id="page-4-1"></span>**Theorem [2.1](#page-3-1).** Assume that Assumptions 2.1 and [2.2](#page-3-2) hold. Then  $Q^*$  is an ELMM if and only if the following conditions hold:  $\{s : \Delta A(s) \neq 0\} \subset \{u_1, u_2, \dots\}$ , and

<span id="page-4-0"></span>(9) 
$$
\int_0^t f(s,s)\nu(ds) = \int_0^t r_s ds + H'(t),
$$

<span id="page-4-3"></span>(10) 
$$
\bar{a}(t,T) = \frac{1}{2} \parallel \bar{b}(t,T) \parallel^{2},
$$

for  $0 \leq t \leq T \leq T^* dQ^* \otimes dt$ -almost surely on  $\{t < \tau\}.$ 

The first condition,  $(9)$ , can be split in the continuous and pure-jump part, such that  $(9)$ is equivalent to

$$
f(t, t) = r_s + \lambda_s
$$
  

$$
f(t, u_i) = \log \frac{w_i}{w_i - \lambda(u_i)\Delta A(u_i)} \ge 0.
$$

The second relation states explicitly the connection of the forward rate at a risky time  $u_i$ to the probability  $Q^*(\tau = u_i | \mathscr{F}_{u_i-})$ , given that  $\tau \geq u_i$ , of course. It simplifies moreover, if  $\Delta A(u_i) = w_i$  to

<span id="page-4-4"></span>(11) 
$$
f(t, u_i) = -\log(1 - \lambda(u_i)).
$$

For the proof we first provide the canonical decomposition of

$$
J(t,T) := \int_t^T f(t,u)\nu(du), \qquad 0 \leq t \leq T.
$$

<span id="page-4-2"></span>**Lemma [2.2](#page-3-2).** Assume that Assumption 2.2 holds. Then, for each  $T \in [0, T^*]$  the process  $(J(t,T))_{0\leqslant t\leqslant T}$  is a special semimartingale and

$$
J(t,T) = \int_0^T f(0, u)\nu(du) + \int_0^t \bar{a}(u,T)du + \int_0^t \bar{b}(u,T)dW_u - \int_0^t f(u,u)\nu(du).
$$

*Proof.* Using the stochastic Fubini Theorem (as in [\[26\]](#page-13-3)), we obtain ˆ ˙

$$
J(t,T) = \int_{t}^{T} \left( f(0, u) + \int_{0}^{t} a(s, u)ds + \int_{0}^{t} b(s, u)dW_{s} \right) \nu(du)
$$
  
\n
$$
= \int_{0}^{T} f(0, u)\nu(du) + \int_{0}^{t} \int_{s}^{T} a(s, u)\nu(du)ds + \int_{0}^{t} \int_{s}^{T} b(s, u)\nu(du)dW_{s}
$$
  
\n
$$
- \int_{0}^{t} f(0, u)\nu(du) - \int_{0}^{t} \int_{s}^{t} a(s, u)\nu(du)ds - \int_{0}^{t} \int_{s}^{t} b(s, u)\nu(du)dW_{s}
$$
  
\n
$$
= \int_{0}^{T} f(0, u)\nu(du) + \int_{0}^{t} \bar{a}(s, T)ds + \int_{0}^{t} \bar{b}(s, T)dW_{s}
$$
  
\n
$$
- \int_{0}^{t} \left( f(0, u) - \int_{0}^{u} a(s, u)ds - \int_{0}^{u} b(s, u)dW_{s} \right) \nu(du),
$$

and the claim follows.

$$
\sqcup
$$

Proof of Theorem [2.1.](#page-4-1) Set,  $E(t) = \mathbb{1}_{\{\tau > t\}}$ , and  $F(t,T) = \exp\left(-\int_t^T$  $\int_t^t f(t, u)\nu(du)$ , such that  $P(t, T) = E(t)F(t, T)$ . Integration by parts yields that

<span id="page-5-2"></span>(12) 
$$
dP(t,T) = F(t-,T) dE(t) + E(t-) dF(t,T) + d[E, F(.,T)]_t =: (1') + (2') + (3').
$$

In view of  $(1')$ , we obtain from  $(4)$ , that

(13) 
$$
E(t) + \int_0^{t \wedge \tau} \lambda_s dA(s) =: M_t^1
$$

is a martingale. Regarding  $(2')$ , note that from Lemma  $2.2$  we obtain by Itô's formula that

<span id="page-5-0"></span>(14)  
\n
$$
\frac{dF(t,T)}{F(t-,T)} = \left(f(t,t) - \bar{a}(t,T) + \frac{1}{2} \|\bar{b}(t,T)\|^2\right)dt + \sum_{i \geq 0} \left(e^{f(t,t)} - 1\right) w_i \delta_{u_i}(dt) + dM_t^2,
$$

with a local martingale  $M^2$ . For the remaining term  $(3')$ , note that

<span id="page-5-1"></span>(15) 
$$
\sum_{0 < s \le t} \Delta E(s) \Delta F(s, T) = \int_0^t F(s-, T)(e^{f(s,s)} - 1)\nu(\{s\})dE(s)
$$

$$
= \int_0^t F(s-, T)(e^{f(s,s)} - 1)\nu(\{s\})dM_s^1
$$

$$
- \int_0^{t \wedge \tau} F(s-, T)(e^{f(s,s)} - 1)\nu(\{s\})\lambda_s dA(s).
$$

Inserting  $(14)$  and  $(15)$  into  $(12)$  we obtain

$$
\frac{dP(t,T)}{P(t-,T)} = -\lambda_t dA(t)
$$
  
+ 
$$
\left(f(t,t) - \bar{a}(t,T) + \frac{1}{2} \|\bar{b}(t,T)\|^2\right)dt
$$
  
+ 
$$
\sum_{i\geq 0} \left(e^{f(t,t)} - 1\right) w_i \delta_{u_i}(dt)
$$
  
- 
$$
\int_{\mathbb{R}} \nu(\{t\}) (e^{f(t,t)} - 1) \lambda_t dA(t) + dM_t^3
$$

with a local martingale  $M^3$ . We obtain a  $Q^*$ -local martingale if and only if the drift vanishes. Next, we can separate between absolutely continuous and discrete part. The absolutely continuous part yields [\(10\)](#page-4-3) and  $f(t, t) = r_t + \lambda_t dQ^* \otimes dt$ -almost surely. It remains to compute the discontinuous part, which is given by

$$
\sum_{i: u_i \le t} P(u_i-,T) (e^{f(u_i, u_i)} - 1) w_i - \sum_{0 < s \le t} P(s-,T) e^{f(s,s)} \lambda_s \Delta A(s),
$$

for  $0 \leq t \leq T \leq T^*$ . This yields  $\{s : \Delta A(s) \neq 0\} \subset \{u_1, u_2, \dots\}$ . The discontinuous part vanishes if and only if

$$
\mathbb{1}_{\{u_i \le T^* \wedge \tau\}} e^{-f(u_i, u_i)} w_i = \mathbb{1}_{\{u_i \le T^* \wedge \tau\}} \Big( w_i - \lambda_{u_i} \Delta A(u_i) \Big), \qquad i \ge 1,
$$

which is equivalent to

$$
\mathbb{1}_{\{u_i \le T^* \wedge \tau\}} f(u_i, u_i) = -\mathbb{1}_{\{u_i \le T^* \wedge \tau\}} \log \frac{w_i - \lambda_{u_i} \Delta A(u_i)}{w_i}, \qquad i \ge 1.
$$

We obtain  $(9)$  and the claim follows.

<span id="page-6-0"></span>Example 2.1 (The Merton model). The paper [\[23\]](#page-13-0) considers a simple capital structure of a firm, consisting only of equity and a zero-coupon bond with maturity  $U > 0$ . The firm defaults at  $U$  if the total market value of its assets is not sufficient to cover the liabilities.

We are interested in setting up an arbitrage-free market for credit derivatives and consider a market of defaultable bonds  $P(t,T)$ ,  $0 \le t \le T \le T^*$  with  $0 < U \le T^*$  as basis for more complex derivatives. In a stylized form the Merton model can be represented by a Brownian motion W denoting the normalized logarithm of the firm's assets, a constant  $K > 0$  and the default time #

$$
\tau = \begin{cases} U & \text{if } W_U \leq K \\ \infty & \text{otherwise.} \end{cases}
$$

Assume for simplicity a constant interest rate r and let  $\mathbb F$  be the filtration generated by W. Then  $P(t,T) = e^{-r(T-t)}$  whenever  $T < U$  because these bonds do not carry default risk. On the other hand, for  $t < U \leq T$ ,

$$
P(t,T) = e^{-r(T-t)} E^* [\mathbb{1}_{\{\tau > T\}} | \mathscr{F}_t] = e^{-r(T-t)} E^* [\mathbb{1}_{\{\tau = \infty\}} | \mathscr{F}_t] = e^{-r(T-t)} \Phi\bigg(\frac{W_t - K}{\sqrt{U - t}}\bigg),
$$

where Φ denotes the cumulative distribution function of a standard normal random variable and  $E^*$  denotes the expectation with respect to  $Q^*$ . For  $t \to U$  we recover  $P(U, U)$  $1_{\{\tau=\infty\}}$ . The derivation of representation [\(6\)](#page-3-3) with  $\nu(du) := du + \delta_U(du)$  is straightforward. A simple calculation with

<span id="page-6-1"></span>(16) 
$$
P(t,T) = 1_{\{\tau > t\}} \exp \left(-\int_{t}^{T} f(t,u) du - f(t,U) 1_{\{t < U \le T\}}\right)
$$

yields  $f(t, T) = r$  for  $T \neq U$  and

$$
f(t, U) = -\log \Phi\left(\frac{W_t - K}{\sqrt{U - t}}\right).
$$

By Itô's formula we obtain

$$
b(t, U) = -\frac{\varphi\left(\frac{W_t - K}{\sqrt{U - t}}\right)}{\Phi\left(\frac{W_t - K}{\sqrt{U - t}}\right)} (U - t)^{-1/2},
$$

and indeed,  $a(t, U) = \frac{1}{2}b^2(t, U)$ . Note that the conditions for Proposition [2.1](#page-4-1) hold and, the market consisting of the bonds  $P(t, T)$  satisfies NAFL, as expected. More flexible models of arbitrage-free bond prices can be obtained if the market filtration  $\mathbb F$  is allowed to be more general, as we show in Section [3](#page-8-0) on affine generalized Merton models.

Example 2.2. (An extension of the Black-Cox model) The model suggested in [\[4\]](#page-12-18) uses a first-passage time approach to model credit risk. Default happens at the first time, when the firm value falls below a pre-specified boundary, the default boundary. We consider a stylized version of this approach and continue the example [2.1.](#page-6-0) Extending the original approach, we include a zero-coupon bond with maturity  $U$ . The reduction of the firm value at  $U$  is equivalent to considering a default boundary with an upward jump at that time. Hence, we consider a Brownian motion  $W$  and the default boundary

$$
D(t) = D(0) + K \mathbb{1}_{\{U \ge t\}}, \qquad t \ge 0,
$$

with  $D(0) < 0$ , and let default be the first time when W hits D, i.e.

$$
\tau = \inf\{t \geq 0 : W_t \leq D(t)\}
$$

with the usual convention that inf  $\varnothing = \infty$ . The following lemma computes the default probability in this setting and the forward rates are directly obtained from this result together with [\(16\)](#page-6-1). The filtration  $\mathbb{G} = \mathbb{F}$  is given by the natural filtration of the Brownian motion W after completion. Denote the random sets

$$
\Delta_1 := \{ (x, y) \in \mathbb{R}^2 : x\sqrt{T - U} \leq D(U) - (y\sqrt{U - t} + W_t), y\sqrt{U - t} + W_t > D(0) \}
$$
  

$$
\Delta_2 := \{ (x, y) \in \mathbb{R}^2 : x\sqrt{T - U} \leq D(U) - (y\sqrt{U - t} + 2D(0) - W_t),
$$
  

$$
y\sqrt{U - t} + D(0) - W_t > 0 \}.
$$

**Lemma 2.3.** Let  $D(0) < 0$ ,  $U > 0$  and  $D(U) \ge D(0)$ . For  $0 \le t < U$ , it holds on  $\{\tau > t\}$ , that

<span id="page-7-0"></span>(17) 
$$
P(\tau > T | \mathscr{F}_t) = 1 - 2\Phi\left(\frac{D(0) - W_t}{\sqrt{T - t}}\right) - \mathbb{1}_{\{T \ge U\}} 2(\Phi_2(\Delta_1) - \Phi_2(\Delta_2)),
$$

where  $\Phi_2$  is the distribution of a two-dimensional standard normal distribution and the sets  $\Delta_t = \Delta_t(D)$ ,  $t \geq U$  are given by

$$
\Delta_t = \{(x, y) \in \mathbb{R}^2 : x\sqrt{T - U} + y\sqrt{U} \leq -D(U), \}.
$$

For  $t \ge U$  it holds on  $\{\tau > t\}$ , that

$$
P(\tau > T | \mathcal{F}_t) = 1 - 2\Phi\left(\frac{D(U) - W_t}{\sqrt{T - t}}\right)
$$

.

*Proof.* The first part of  $(17)$  where  $T < U$  follows directly from the reflection principle and the property that W has independent and stationary increments. Next, consider  $0 \leq t$  $U \leq T$ . Then, on  $\{W_U > D(U)\},\$ 

<span id="page-7-1"></span>(18) 
$$
P(\inf_{[U,T]} W > D(U) | \mathscr{F}_U) = 1 - 2\Phi\left(\frac{D(U) - W_U}{\sqrt{T - U}}\right).
$$

Moreover, on  $\{W_t > D(0)\}\$ it holds for  $x > D(0)\$  that

$$
P(\inf_{[0,U]} W > D(0), W_U > x | \mathcal{F}_t) = P(W_U > x | \mathcal{F}_t) - P(W_U < x, \inf_{[0,U]} W \le D(0) | \mathcal{F}_t)
$$
  
= 
$$
\Phi\left(\frac{W_t - x}{\sqrt{U - t}}\right) - \Phi\left(\frac{2D(0) - x - W_t}{\sqrt{U - t}}\right).
$$

Hence,  $E[g(W_U)1_{\{\inf_{[0,U]} W > D(0)\}}|\mathscr{F}_t] = 1_{\{\inf_{[0,t]} W > D(0)\}} \int_{D(0)}^{\infty} g(x) f_t(x) dx$  with density " ⊤

$$
f_t(x) = 1_{\{x > D(0)\}} \frac{1}{\sqrt{U-t}} \left[ \phi \left( \frac{W_t - x}{\sqrt{U-t}} \right) - \phi \left( \frac{2D(0) - x - W_t}{\sqrt{U-t}} \right) \right].
$$

Together with [\(18\)](#page-7-1) this yields on  $\{\inf_{[0,t]} W > D(0)\}\$ 

$$
P(\inf_{[0,T]}(W-D) > 0 | \mathcal{F}_t) = \int_{D(0)}^{\infty} \left[1 - 2\Phi\left(\frac{D(U) - x}{\sqrt{T-U}}\right)\right] f_t(x) dx
$$
  
= 
$$
P(\inf_{[t,T]} W > D(0) | \mathcal{F}_t) - 2 \int_{D(0)}^{\infty} \Phi\left(\frac{D(U) - x}{\sqrt{T-U}}\right) f_t(x) dx.
$$

It remains to compute the integral. Regarding the first part, letting  $\xi$  and  $\eta$  be independent and standard normal, we obtain that

$$
\int_{D(0)}^{\infty} \Phi\left(\frac{D(U) - x}{\sqrt{T - U}}\right) \frac{1}{\sqrt{U - t}} \phi\left(\frac{x - W_t}{\sqrt{U - t}}\right) dx
$$
  
=  $P_t \left(\sqrt{T - U}\xi \le D(U) - (\sqrt{U - t}\eta + W_t), \sqrt{U - t}\eta + W_t > D(0)\right)$   
=  $\Phi_2(\Delta_1)$ ,

where we abbreviate  $P_t(\cdot) = P(\cdot | \mathcal{F}_t)$ . In a similar way,<br> $\int_0^\infty$   $\langle P(IU) - x \rangle = 1$   $\langle x - (2D(0) - W_0) \rangle$ 

$$
\int_{D(0)}^{\infty} \Phi\left(\frac{D(U) - x}{\sqrt{T - U}}\right) \frac{1}{\sqrt{U - t}} \phi\left(\frac{x - (2D(0) - W_t)}{\sqrt{U - t}}\right) dx
$$
  
=  $P_t \left(\sqrt{T - U} \xi \le D(U) - (\sqrt{U - t}\eta + 2D(0) - W_t), \sqrt{U - t}\eta + D(0) - W_t > 0\right)$   
=  $\Phi_2(\Delta_2)$ 

<span id="page-8-0"></span>and we conclude.  $\Box$ 

#### 3. Affine models in the generalized intensity-based framework

Affine processes are a well-known tool in the financial literature and one reason for this is their analytical tractability. In this section we closely follow [\[12\]](#page-12-6) and shortly state the appropriate affine models which fit the generalized intensity framework. For proofs, we refer the reader to this paper.

The main point is that affine processes in the literature are assumed to be stochastically continuous (see  $[7]$  and  $[10]$ ). Due to the discontinuities introduced in the generalized intensity-based framework, we propose to consider piecewise continuous affine processes.

<span id="page-8-1"></span>**Example 3.1.** Consider a non-negative integrable function  $\lambda$ , a constant  $\lambda' \geq 0$  and a deterministic time  $u > 0$ . Set

$$
K(t) = \int_0^t \lambda(s)ds + \mathbb{1}_{\{t \geqslant u\}} \kappa, \qquad t \geqslant 0.
$$

Let the default time  $\tau$  be given by  $\tau = \inf\{t \geq 0 : K_t \geq \zeta\}$  with a standard exponentialrandom variable  $\zeta$ . Then  $P(\tau = u) = 1 - e^{-\kappa} =: \lambda'$ . Considering  $\nu(ds) = ds + \delta_u(ds)$  with  $u_1 = u$  and  $w_1 = 1$ , we are in the setup of the previous section. The drift condition [\(9\)](#page-4-0) holds, if

$$
f(u, u) = -\log(1 - \lambda') = \kappa.
$$

Note, however, that  $K$  is not the compensator of  $H$ . Indeed, the compensator of  $H$  equals Note, how<br>  $\Lambda_t = \int_0^{t \wedge \tau}$  $\int_0^{t \wedge \tau} \lambda(s) ds + \mathbb{1}_{\{t \geq u\}} \lambda'$ , see [\[19\]](#page-12-21) for general results in this direction.

The purpose of this section is to give a suitable extension of the above example involving affine processes. Recall that we consider a  $\sigma$ -finite measure

$$
\nu(du) = du + \sum_{i \geq 1} w_i \delta_{u_i}(du),
$$

as well as  $A(u) = u + \sum_{i \geq 1} 1_{\{u \geq u_i\}}$ . The idea is to consider an affine process X and study arbitrage-free doubly stochastic term structure models where the compensator  $\Lambda$  of the default indicator process  $H = \mathbb{1}_{\{\cdot \leq \tau\}}$  is given by

<span id="page-9-0"></span>(19) 
$$
\Lambda_t = \int_0^t \left( \phi_0(s) + \psi_0(s)^\top \cdot X_s \right) ds + \sum_{i \geq 1} \mathbb{1}_{\{t \geq u_i\}} \left( 1 - e^{-\phi_i - \psi_i^\top \cdot X_{u_i}} \right).
$$

Note that by continuity of X,  $\Lambda_t(\omega) < \infty$  for almost all  $\omega$ . To ensure that  $\Lambda$  is non-decreasing we will require that  $\phi_0(s) + \psi_0(s)^\top \cdot X_s \geq 0$  for all  $s \geq 0$  and  $\phi_i + \psi_i^\top \cdot X_{u_i} \geq 0$  for all  $i \geq 1$ .

Consider a state space in canonical form  $\mathcal{X} = \mathbb{R}^m_{\geq 0} \times \mathbb{R}^n$  for integers  $m, n \geq 0$  with  $m + n = d$  and a d-dimensional Brownian motion W. Let  $\mu$  and  $\sigma$  be defined on X by

<span id="page-9-3"></span>(20) 
$$
\mu(x) = \mu_0 + \sum_{i=1}^d x_i \mu_i,
$$

<span id="page-9-4"></span>(21) 
$$
\frac{1}{2}\sigma(x)^{\top}\sigma(x) = \sigma_0 + \sum_{i=1}^d x_i \sigma_i,
$$

where  $\mu_0, \mu_i \in \mathbb{R}^d$ ,  $\sigma_0, \sigma_i \in \mathbb{R}^{d \times d}$ , for all  $i \in \{1, \ldots, d\}$ . We assume that the parameters  $\mu^i, \sigma^i$ ,  $i = 0, \ldots, d$  are admissible in the sense of Theorem 10.2 in [\[11\]](#page-12-10). Then the continuous, unique strong solution of the stochastic differential equation

(22) 
$$
dX_t = \mu(X_t)dt + \sigma(X_t)dW_t, \quad X_0 = x,
$$

is an *affine* process X on the state space  $\mathcal{X}$ , see Chapter 10 in [\[11\]](#page-12-10) for a detailed exposition.

We call a bond-price model *affine* if there exist functions  $A : \mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0} \to \mathbb{R}, B$ :  $\mathbb{R}_{\geqslant 0} \times \mathbb{R}_{\geqslant 0} \to \mathbb{R}^d$  such that

<span id="page-9-1"></span>(23) 
$$
P(t,T) = 1_{\{\tau > t\}} e^{-A(t,T) - B(t,T)^{\top} \cdot X_t},
$$

for  $0 \leq t \leq T \leq T^*$ . We assume that  $A(., T)$  and  $B(., T)$  are right-continuous. Moreover, we assume that  $t \mapsto A(t,.)$  and  $t \mapsto B(t,.)$  are differentiable from the right and denote by  $\partial_t^+$  the right derivative. For the convenience of the reader we state the following proposition giving sufficient conditions for absence of arbitrage in an affine generalized-intensity based setting. It extends [\[12\]](#page-12-6) where only finitely many risky times were treated.

<span id="page-9-5"></span>**Proposition 3.1.** Assume that  $\phi_0 : \mathbb{R}_{\geqslant 0} \to \mathbb{R}$ ,  $\psi_0 : \mathbb{R}_{\geqslant 0} \to \mathbb{R}^d$  are continuous,  $\psi_0(s)$  +  $\psi_0(s)^\top \cdot x \geq 0$  for all  $s \geq 0$  and  $x \in \mathcal{X}$  and the constants  $\phi_i \in \mathbb{R}$  and  $\psi_i \in \mathbb{R}^d$ ,  $i \geq 1$  satisfy  $\psi_0(s)$   $\cdot$   $x \ge 0$  for all  $s \ge 0$  and  $x \in \mathcal{X}$  and the constants  $\phi_i \in \mathbb{R}$  and  $\psi_i \in \mathbb{R}^a$ ,  $i \ge 1$  satisfy  $\phi_i + \psi_i^{\top} \cdot x \ge 0$  for all  $1 \le i \le n$  and  $x \in \mathcal{X}$  as well as  $\sum_{i \ge 1} |w_i|(|\phi_i| + |\psi_{i,1}| + \cdots +$ Moreover, let the functions  $A: \mathbb{R}_{\geqslant 0} \times \mathbb{R}_{\geqslant 0} \to \mathbb{R}$  and  $B: \mathbb{R}_{\geqslant 0} \times \mathbb{R}_{\geqslant 0} \to \mathbb{R}^d$  be the unique solutions of

<span id="page-9-2"></span>(24) 
$$
A(T, T) = 0
$$

$$
A(u_i, T) = A(u_i-, T) - \phi_i w_i
$$

$$
-\partial_t^+ A(t, T) = \phi_0(t) + \mu_0^{\top} \cdot B(t, T) - B(t, T)^{\top} \cdot \sigma_0 \cdot B(t, T),
$$

and

<span id="page-10-1"></span>(25) 
$$
B(T,T) = 0
$$

$$
B_k(u_i, T) = B_k(u_i-, T) - \psi_{i,k}w_i
$$

$$
-\partial_t^+ B_k(t,T) = \psi_{0,k}(t) + \mu_k^\top \cdot B(t,T) - B(t,T)^\top \cdot \sigma_k \cdot B(t,T),
$$

for  $0 \le t \le T$ . Then, the doubly-stochastic affine model given by [\(19\)](#page-9-0) and [\(23\)](#page-9-1) satisfies NAFL.

Proof. By construction,

$$
A(t,T) = \int_t^T a'(t,u)du + \sum_{i:u_i \in (t,T]} \phi_i w_i
$$

$$
B(t,T) = \int_t^T b'(t,u)du + \sum_{i:u_i \in (t,T]} \psi_i w_i
$$

with suitable functions a' and b' and  $a'(t,t) = \phi_0(t)$  as well as  $b'(t,t) = \psi_0(t)$ . A comparison of [\(23\)](#page-9-1) with [\(6\)](#page-3-3) yields the following: on the one hand, for  $T = u_i \in \mathcal{U}$ , we obtain  $f(t, u_i)$  $\phi_i + \psi_i^{\top} \cdot X_t$ . Hence, the coefficients  $a(t,T)$  and  $b(t,T)$  in [\(7\)](#page-3-4) for  $T = u_i \in \mathcal{U}$  compute to  $a(t, u_i) = \psi_i^{\top} \cdot \mu(X_t)$  and  $b(t, u_i) = \psi_i^{\top} \cdot \sigma(X_t)$ .

On the other hand, for  $T \notin \mathcal{U}$  we obtain that  $f(t,T) = a'(t,T) + b'(t,T)^{\top} \cdot X_t$ . Then, the coefficients  $a(t, T)$  and  $b(t, T)$  can be computed as follows: applying Itô's formula to  $f(t, T)$ and comparing with [\(7\)](#page-3-4) yields that

<span id="page-10-0"></span>(26) 
$$
a(t,T) = \partial_t a'(t,T) + \partial_t b'(t,T)^\top \cdot X_t + b'(t,T)^\top \cdot \mu(X_t)
$$

$$
b(t,T) = b'(t,T)^\top \cdot \sigma(X_t).
$$

Set  $\bar{a}'(t,T) = \int_t^T a'(t,u)du$  and  $\bar{b}'(t,T) = \int_t^T b'(t,u)du$  and note that,

$$
\int_t^T \partial_t a'(t, u) du = \partial_t \bar{a}'(t, T) + a'(t, t).
$$

As  $\partial_t^+ A(t,T) = \partial_t \bar{a}'(t,T)$ , and  $\partial_t^+ B(t,T) = \partial_t \bar{b}'(t,T)$ , we obtain from [\(26\)](#page-10-0) that

$$
\bar{a}(t,T) = \int_t^T a(t,u)\nu(du) = \int_t^T a(t,u)du + \sum_{u_i \in (t,T]} w_i \psi_i^\top \cdot \mu(X_t)
$$
  
\n
$$
= \partial_t^+ A(t,T) + a'(t,t) + \left(\partial_t^+ B(t,T) + b'(t,t)\right)^\top \cdot X_t + B(t,T)^\top \cdot \mu(X_t),
$$
  
\n
$$
\bar{b}(t,T) = \int_t^T b(t,u)\nu(du) = \int_t^T b(t,u)du + \sum_{u_i \in (t,T]} w_i \psi_i^\top \cdot \sigma(X_t)
$$
  
\n
$$
= B(t,T)^\top \cdot \sigma(X_t)
$$

for  $0 \le t \le T \le T^*$ . We now show that under our assumptions, the drift conditions [\(9\)](#page-4-0)-[\(10\)](#page-4-3) hold: Observe that, by equations  $(24)$ ,  $(25)$ , and the affine specification  $(20)$ , and  $(21)$ , the drift condition  $(10)$  holds. Moreover, from  $(11)$ ,

$$
\Delta H'(u_i) = \phi_i + \psi_i^{\top} \cdot X_{u_i}
$$

and  $\lambda_s = \phi_0(s) + \psi_0(s)^\top \cdot X_s$  by [\(19\)](#page-9-0). We recover  $\Delta \Lambda_{u_i} = 1 - \exp(-\phi_i - \psi_i^\top \cdot X_{u_i})$  taking values in  $[0, 1)$  by assumption. Hence, [\(9\)](#page-4-0) holds and the claim follows. **Example 3.2.** In the one-dimensional case we consider  $X$ , given as solution of

$$
dX_t = (\mu_0 + \mu_1 X_t)dt + \sigma \sqrt{X_t}dW_t, \qquad t \geq 0.
$$

Consider only one risky time  $u_1 = 1$  and let  $\phi_0 = \phi_1 = 0$ ,  $\psi_0 = 1$ , such that

$$
\Lambda = \int_0^t X_s ds + \mathbb{1}_{\{u \ge 1\}} (1 - e^{-\psi_1 X_1}).
$$

Hence the probability of having no default at time 1 just prior to 1 is given by  $e^{-\psi_1 X_1}$ , compare Example [3.1.](#page-8-1)

An arbitrage-free model can be obtained by choosing A and B according to Proposition [3.1](#page-9-5) which can be immediately achieved using Lemma 10.12 from [\[11\]](#page-12-10) (see in particular a Section 10.3.2.2 on the CIR short-rate model): denote  $\theta = \sqrt{\mu_1^2 + 2\sigma^2}$  and

$$
L_1(t) = 2(e^{\theta t} - 1),
$$
  
\n
$$
L_2(t) = \theta(e^{\theta t} + 1) + \mu_1(e^{\theta t} - 1),
$$
  
\n
$$
L_3(t) = \theta(e^{\theta t} + 1) - \mu_1(e^{\theta t} - 1),
$$
  
\n
$$
L_4(t) = \sigma^2(e^{\theta t} - 1).
$$

Then

$$
A_0(s) = \frac{2\mu_0}{\sigma^2} \log \left( \frac{2\theta e^{\frac{(\sigma - \mu_1)t}{2}}}{L_3(t)} \right), \quad B_0(s) = -\frac{L_1(t)}{L_3(t)}
$$

are the unique solutions of the Riccati equations  $B'_0 = \sigma^2 B_0^2 - \mu_1 B_0$  with boundary condition  $B_0(0) = 0$  and  $A'_0 = -\mu_0 B_0$  with boundary condition  $A_0(0) = 0$ . Note that with  $A(t, T) =$  $A_0(T - t)$  and  $B(t, T) = B_0(T - t)$  for  $0 \le t \le T < 1$ , the conditions of Proposition [3.1](#page-9-5) hold. Similarly, for  $1 \le t \le T$ , choosing  $A(t, T) = A_0(T - t)$  and  $B(t, T) = B_0(T - t)$  implies again the validity of [\(24\)](#page-9-2) and [\(25\)](#page-10-1). On the other hand, for  $0 \leq t < 1$  and  $T \geq 1$  we set  $u(T) = B(1, T) + \psi_1 = B_0(T - 1) + \psi_1$ , according to [\(25\)](#page-10-1), and let

$$
A(t,T) = \frac{2\mu_0}{\sigma^2} \log \left( \frac{2\theta e^{\frac{(\sigma - \mu_1)(1-t)}{2}}}{L_3(1-t) - L_4(1-t)u(T)} \right)
$$

$$
B(t,T) = -\frac{L_1(1-t) - L_2(1-t)u(T)}{L_3(1-t) - L_4(1-t)u(T)}.
$$

It is easy to see that [\(24\)](#page-9-2) and [\(25\)](#page-10-1) are also satisfied in this case, in particular  $\Delta A(1,T)$  $-\phi_1 = 0$  and  $\Delta B(1, T) = -\psi_1$ . Note that, while X is continuous, the bond prices are not even stochastically continuous because they jump almost surely at  $u_1 = 1$ . We conclude by Proposition [3.1](#page-9-5) that this affine model is arbitrage-free.  $\Diamond$ 

## 4. Conclusion

<span id="page-11-0"></span>In this article we studied a new class of dynamic term structure models with credit risk where the compensator of the default time may jump at predictable times. This framework was called generalized intensity-based framework. It extends existing theory and allows to include Merton's model, in a reduced-form model for pricing credit derivatives. Finally, we studied a class of highly tractable affine models which are only piecewise stochastically continuous.

#### GENERALIZED INTENSITY 13

### **REFERENCES**

- <span id="page-12-2"></span>[1] Artzner, P. and Delbaen, F. (1995), 'Default risk insurance and incomplete markets', Mathematical Finance 5(3), 187–195.
- <span id="page-12-5"></span>[2] Bélanger, A., Shreve, S. E. and Wong, D. (2004), 'A general framework for pricing credit risk', Mathematical Finance 14(3), 317–350.
- <span id="page-12-11"></span>[3] Bielecki, T. and Rutkowski, M. (2002), Credit Risk: Modeling, Valuation and Hedging, Springer Verlag. Berlin Heidelberg New York.
- <span id="page-12-18"></span>[4] Black, F. and Cox, J. C. (1976), 'Valuing corporate securities: some effects of bond indenture provisions', Journal of Finance 31, 351–367.
- <span id="page-12-13"></span>[5] Brémaud, P. (1981), *Point Processes and Queues*, Springer Verlag. Berlin Heidelberg New York.
- <span id="page-12-16"></span>[6] Cuchiero, C., Klein, I. and Teichmann, J. (2014), 'A new perspective on the fundamental theorem of asset pricing for large financial markets', ArXiv e-prints .
- <span id="page-12-19"></span>[7] Duffie, D., Filipović, D. and Schachermayer, W. (2003), 'Affine processes and applications in finance', Annals of Applied Probability 13, 984–1053.
- <span id="page-12-14"></span>[8] Duffie, D. and Singleton, K. (1999), 'Modeling term structures of defaultable bonds', Review of Financial Studies 12, 687–720.
- <span id="page-12-8"></span>[9] Eavis, P. (2015), 'Greece is placed in arrears, as the i.m.f. spells default', http://www.nytimes.com/2015/07/01/business/international/greece-is-placedin-arrears-as-the-imf-spells-default.html.
- <span id="page-12-20"></span>[10] Filipović, D. (2005), 'Time-inhomogeneous affine processes', Stochastic Processes and *their Applications* **115**(4), 639 – 659.
- <span id="page-12-10"></span>[11] Filipović, D. (2009), Term Structure Models: A Graduate Course, Springer Verlag. Berlin Heidelberg New York.
- <span id="page-12-6"></span>[12] Gehmlich, F. and Schmidt, T. (2015), 'Dynamic defaultable term structure modelling beyond the intensity paradigm', submitted .
- <span id="page-12-3"></span>[13] Geske, R. (1977), 'The valuation of corporate liabilities as compound options', Journal of Financial and Quantitative Analysis 12, 541–552.
- <span id="page-12-4"></span>[14] Geske, R. and Johnson, H. E. (1984), 'The valuation of corporate liabilities as compound options: A correction', Journal of Financial and Quantitative Analysis 19(2), 231–232.
- <span id="page-12-12"></span>[15] Guo, X. and Zeng, Y. (2008), 'Intensity process and compensator: a new filtration expansion approach and the Jeulin-Yor theorem', Ann. Appl. Probab.  $18(1)$ ,  $120-142$ .
- <span id="page-12-7"></span>[16] ISDA (2014), 'ISDA Americas credit derivatives determinations committee: Argentine republic failure to pay credit event', http://www2.isda.org/news/isda-americas-creditderivatives-determinations-committee-argentine-republic-failure-to-pay-credit-event.
- <span id="page-12-9"></span>[17] Jacod, J. and Shiryaev, A. (2003), Limit Theorems for Stochastic Processes, 2nd edn, Springer Verlag, Berlin.
- <span id="page-12-0"></span>[18] Jarrow, R. and Turnbull, S. (1995), 'Pricing options on financial securities subject to default risk', Journal of Finance 5, 53–86.
- <span id="page-12-21"></span>[19] Jeanblanc, M. and Rutkowski, M. (2000), Modeling default risk: an overview, in 'Mathematical Finance: theory and practice Fudan University', Modern Mathematics Series, High Education press, Beijing, pp. 171 – 269.
- <span id="page-12-17"></span>[20] Karatzas, I. and Shreve, S. E. (1988), Brownian Motion and Stochastic Calculus, Springer Verlag. Berlin Heidelberg New York.
- <span id="page-12-15"></span>[21] Klein, I., Schmidt, T. and Teichmann, J. (2015), 'When roll-overs do not qualify as numéraire: bond markets beyond short rate paradigms',  $arXiv$  1310.0032
- <span id="page-12-1"></span>[22] Lando, D. (1994), Three Essays on Contingent Claim Pricing, PhD thesis, Cornell University.
- <span id="page-13-0"></span>[23] Merton, R. (1974), 'On the pricing of corporate debt: the risk structure of interest rates', Journal of Finance 29, 449–470.
- <span id="page-13-1"></span>[24] Reuters (2014), 'Argentina debt talks must continue after default: U.S. judge', http://www.reuters.com/article/2014/08/01/us-argentina-debtidUSKBN0G13Z720140801.
- <span id="page-13-2"></span>[25] Schönbucher, P. (1998), 'Pricing credit risk derivatives', Working paper .
- <span id="page-13-3"></span>[26] Veraar, M. (2012), 'The stochastic Fubini theorem revisited', Stochastics 84(4), 543– 551.

Frank Gehmlich email: gehmlichfrank@gmail.com and Thorsten Schmidt, University of Freiburg, Dep. of Mathematics, Eckerstr 1, 79106 Freiburg, Germany email: thorsten.schmidt@stochastik.unifreiburg.de