EXISTENCE OF VERY WEAK SOLUTIONS TO ELLIPTIC SYSTEMS OF p-LAPLACIAN TYPE

MIROSLAV BULÍČEK AND SEBASTIAN SCHWARZACHER

ABSTRACT. We study vector valued solutions to non-linear elliptic partial differential equations with p-growth. Existence of a solution is shown in case the right hand side is the divergence of a function which is only q integrable, where q is strictly below but close to the duality exponent p'. It implies that possibly degenerate operators of p-Laplacian type are well posed in a larger class then the natural space of existence. The key novelty here is a refined a priori estimate, that recovers a duality relation between the right hand side and the solution in terms of weighted Lebesgue spaces.

1. Introduction

Let $\Omega \subset \mathbb{R}^d$ be a Lipschitz domain and $S: \Omega \times \mathbb{R}^{d \times N} \to \mathbb{R}^{d \times N}$ be a Carathéodory mapping. We investigate the existence of a very weak solution $u: \Omega \to \mathbb{R}^N$ with $N \in \mathbb{N}$ of the system

(1.1)
$$\operatorname{div} S(\cdot, \nabla u) = \operatorname{div} |f|^{p-2} f \quad \text{in } \Omega,$$
$$u = 0 \quad \text{on } \partial \Omega.$$

Here we assume growth, coercivity and monotonicity assumptions on S related to the exponent $p \in (1, \infty)$. Explicitly, that there exist constants $C_1, C_2 > 0$ and $C_3 \ge 0$, such that for all $z_1, z_2 \in \mathbb{R}^{d \times N}$ and almost all $x \in \Omega$ there holds

(1.2)
$$S(x, z_1) \cdot z_1 \ge C_1 |z_1|^p - C_3$$
, coercivity

(1.3)
$$|S(x, z_1)| \le C_2 |z_1|^{p-1} + C_3^{\frac{p-1}{p}}, \text{ boundedness}$$

(1.4)
$$(S(x, z_1) - S(x, z_2)) \cdot (z_1 - z_2) \ge 0$$
, monotonicity.

The model case is the p-Laplacian system

(1.5)
$$\operatorname{div} |\nabla u|^{p-2} \nabla u = \operatorname{div} |f|^{p-2} f \text{ in } \Omega,$$

$$u = 0 \text{ on } \partial \Omega.$$

It is well known, that if assumptions (1.2)–(1.4) are satisfied and $f \in L^p(\Omega; \mathbb{R}^{d \times N})$, then a solution of (1.1) exists with $\nabla u \in L^p(\Omega)$. It can for instance be shown by monotone operator theory. The starting point of our investigations is the following question.

Q: Does for $f \in L^q(\Omega; \mathbb{R}^{d \times N})$ and $q \neq p$ a distributional solution u of (1.1) exist, such that $\nabla u \in L^q(\Omega; \mathbb{R}^{d \times N})$?

Under general assumptions (1.2)–(1.4) the answer to this question is not affirmative for all $q \in (1, \infty)$. This is well known due to the various counterexamples even in the simplest case p = 2 and $f \equiv 0$. At the end of exponents smaller then 2 we mention the non-smooth solutions constructed by Serrin [19]. At the end of exponents larger than 2 the non-smooth

²⁰¹⁰ Mathematics Subject Classification. 35D99,35J57,35J60,35A01.

Key words and phrases. nonlinear elliptic systems, weighted estimates, existence, very weak solution, monotone operator, div-curl-biting lemma, weighted space, Muckenhoupt weights.

M. Bulíček's work is supported by the project LL1202 financed by the Ministry of Education, Youth and Sports, Czech Republic and by the University Centre for Mathematical Modelling, Applied Analysis and Computational Mathematics (Math MAC) S. Schwarzacher thanks the program PRVOUK P47, financed by Charles University in Prague. M. Bulíček is a member of the Nečas Center for Mathematical Modeling.

solutions constructed by Nečas [18]. The example of Serrin [19] implies also that there can be no hope for uniqueness in the large class of $W_0^{1,q}(\Omega;\mathbb{R}^N)$ with $q \in (1,p)$, unless stricter assumptions are available.

However, it turns out that closely around p the existence of solutions with natural integrability is true, even under the minimal assumptions (1.2)–(1.4).

More precisely, we are able to prove the following theorem.

Theorem 1.1. Let $\Omega \subset \mathbb{R}^d$ be a bounded Lipschitz domain and S satisfy (1.2)–(1.4). Then there is an ε depending on C_1, C_2, d, N , p and Ω , such that for all $q \in [p - \varepsilon, p]$ the following holds. If $f \in L^q(\Omega; \mathbb{R}^{d \times N})$, then there exists $u \in W_0^{1,q}(\Omega; \mathbb{R}^N)$ which is a distributional solution to (1.1).

Moreover, there is a constant c only depending on $C_1, C_2, p, q, d, N, \Omega$ and a constant c_1 depending additionally on C_3 , such that

(1.6)
$$\|\nabla u\|_{L^{q}(\Omega;\mathbb{R}^{d\times N})} \le c\|f\|_{L^{q}(\Omega;\mathbb{R}^{d\times N})} + c_{1}.$$

As was mentioned before, this result is optimal with respect to the generality of the assumptions (1.2)–(1.4). Let us briefly collect what was already known before.

In case q > p, the existence of a solution in $W_0^{1,p}(\Omega)$ is obvious. The integrability improvement follows by an argument known as Gehring's Lemma. With the minimal assumptions (1.2)–(1.4) it was proved in [16, Theorem 7.8]. See also the more classical result [11, Theorem 4.1] and for an overview and more details [17]. For the p-Laplacian, it is known that (1.6) holds also in the case of large exponents $q \in [p, \infty)$ [13, 5] and beyond [6, 9].

In the special case p=2, it was possible to show that there exists an ε depending on C_1, C_2 and Ω alone, such that existence, uniqueness and regularity are available in case $f \in L^q(\Omega)$ and $q \in [2-\varepsilon, 2+\varepsilon]$ [3]. However, it needed the stricter assumption, that S is Lipschitz continuous with respect to z. Existence, Uniqueness and Regularity for the full range $q \in (1, \infty)$ has recently been shown to hold, in case p=2 with S having additional Uhlenbeck type structure [4].

In the case of $p \neq 2$ and q < p very little is known about the existence of a distributional solution. The situation is quite delicate, since even for bounded domains the existence of any object of solution is not obvious. The only existence result available related to powers q < p for the p-Laplacian are restricted to the scalar case N = 1 and to a better structure of the right hand side. More precisely, when the right hand side is a function (or a Radon measure) [1]. But even for solutions to (1.5) the existence of solutions was not known in the case of $p \neq 2$ and q < p. This seems astonishing, since the a priori estimates (1.6) are available for solutions to (1.5) in case $\mathbb{R}^d = \Omega$ ever since the seminal work of Iwaniec [14] in 1992. There it is showed, that there is an $\varepsilon > 0$ that provided a distributional solution to the p-Laplace exists, it holds already (1.6) for $q \in [p - \varepsilon, p]$. Greco, Iwaniec and Sbordone could stretch the existence frame slightly by showing existence provided the right hand side is in the grande Lebesgue space, $f \in L^p(\Omega; \mathbb{R}^{d \times N})$ [12]. It is a space, that is slightly larger then $L^p(\Omega; \mathbb{R}^{d \times N})$ quantified in terms of logarithmic powers. Later the a priori estimates could be extended to the parabolic case in [15].

After having collected all above efforts we conclude that the existence was open ever since 1992 for q < p and could not be closed ever since. Therefore our main result is the existence of a distributional solution in case $q \in [p - \varepsilon, p]$ and not the estimate (1.6), although it is new in this general form.

The key observation is, that the L^q a priori estimate alone is not suitable to establish solutions to non-linear operators. This is due to the fact that the a priori estimates inherit only weak compactness or less and the only possible way to mach weak compactness and non-linearities is via convexity. In the setting here it is reflected via the use of the monotonicity, for instance via the Minty method. However, the method seemed lost the moment the weak limit is not a suitable test function anymore. Only recently a new point of view was

established, which allowed to regain a duality relation between f and ∇u [4]. The duality is gained by replacing L^q estimates with weighted L^p_ω estimates, where the weight is chosen in terms of the right hand side f. It has to be chosen in such a way that it belongs to the Muckenhoupt class \mathcal{A}_q . This is necessary since it is known, that many linear and sub-linear operators including the Laplace and the Maximal operator are bounded restrictively in these Muckenhoupt classes.

The key novelty, which is of its sovereign interest is the a priori estimate in the existence and regularity result below.

Theorem 1.2. Let $\Omega \subset \mathbb{R}^d$ be a bounded Lipschitz domain and S satisfy (1.2)–(1.3). Then there is an ε depending on $\frac{C_2^{p'}}{C_1}$, p, d, N and Ω , such that for all $q \in [p-\varepsilon, p]$ the following holds. If $f \in L^q(\Omega; \mathbb{R}^{d \times N})$, then there exists $u \in W_0^{1,q}(\Omega; \mathbb{R}^N)$ which is a distributional solution to (1.1).

Moreover, there is a constant c depending on C_1, C_2, p, q, d, N and Ω , such that

(1.7)
$$\int_{\Omega} |\nabla u|^p (M(f+1))^{q-p} \, \mathrm{d}x \le c \int_{\Omega} |f|^q \, \mathrm{d}x + c(C_3+1).$$

Let us point out that the above estimate measures ∇u more accurate in relation to the right hand side. For once we find by (5.10) that (1.7) implies (1.6). But it also implies for instance that non p-integrable singularities of ∇u , can only appear in areas, where f is large, quantified by the naturally related weight. We therefore believe, that estimates of the above type will be of increasing importance in the framework of the existence theory in many applications. Moreover, we wish to indicate its potential for numerical analysis, especial its use for adaptive schemes.

After little restrictions the method can easily be applied on unbounded domains.

Corollary 1.3. Let $\Omega \subset \mathbb{R}^d$ be a Lipschitz domain and S satisfy (1.2)-(1.3), with $C_3 = 0$. Then there is an ε depending on $\frac{C_2^{p'}}{C_1}$, p, d, N and Ω , such that for all $q \in [p-\varepsilon, p]$ the following holds. If $f \in L^q(\Omega; \mathbb{R}^{d \times N})$, then there exists $u \in D_0^{1,q}(\Omega; \mathbb{R}^N)^2$ which is a distributional solution to (1.1).

Moreover, there is a constant c only depending on C_1, C_2, p, q, d, N and Ω , such that

(1.8)
$$\int_{\Omega} |\nabla u|^q + |\nabla u|^p (Mf)^{q-p} \, \mathrm{d}x \le c \int_{\Omega} |f|^q \, \mathrm{d}x.$$

The structure of the paper is as follows. We introduce the necessary notation in the preliminary below. In Section 3 we introduce a truncation method of Sobolev functions, relative to an open set, which is the analytical highlight of this article. In Section 4 we deduce the a-priory estimates and in Section 5 we prove the existence.

2. Preliminary

Throughout the paper all cubes will have sides parallel to the axes. By c, C we denote a generic constant, i.e. its value may change at every appearance. Its dependencies are either stated in the results or are indicated in $C(\ldots)$. Let us recall the definition of the Hardy Littlewood maximal function. For any $f \in L^1_{loc}(\mathbb{R}^n)$ we define

$$Mf(x) := \sup_{\{Q \text{ a cube } : x \in Q\}} \langle |f| \rangle_Q \quad \text{with} \quad \langle |f| \rangle_Q := \int_Q |f(y)| \, \mathrm{d}y := \frac{1}{|Q|} \int_Q |f(y)| \, \mathrm{d}y.$$

It is standard, that the operator is sub linear and continuous from $L^s(\Omega) \to L^s(\Omega)$, for $s \in (1, \infty]$. Further, we say that $\omega : \mathbb{R}^d \to \mathbb{R}$ is a weight function if it is a measurable function

 $^{^1}$ Here M is the Hardy Littlewood maximal operator that is defined in Section 2.

 $^{{}^2}D_0^{1,q}(\Omega;\mathbb{R}^N) := \{ u \in L^1_{\mathrm{loc}}(\omega;\mathbb{R}^N) : \exists u^j \in C_0^{\infty}(\Omega;\mathbb{R}^N) \text{ s.t. } \nabla u^j \to \nabla u \text{ in } L^q(\Omega;\mathbb{R}^d) \}.$

that is almost everywhere finite and positive. For such a weight and arbitrary measurable $\Omega \subset \mathbb{R}^d$ we denote the space $L^p_\omega(\Omega; \mathbb{R}^N)$ with $p \in [1, \infty)$ as

$$L^p_\omega(\Omega,\mathbb{R}^d) := \bigg\{ f: \Omega \to \mathbb{R}^N, \text{ measurable }: \ \|f\|_{L^p_\omega} := \bigg(\int_\Omega |f(x)|^p \omega(x) \, \mathrm{d}x \bigg)^{\frac{1}{p}} < \infty \bigg\}.$$

We introduce the weighted Sobolev spaces, as

$$W^{1,q}_{\omega}(\Omega;\mathbb{R}^N) := \{ u \in W^{1,1}(\Omega;\mathbb{R}^N) : \nabla u \in L^q_{\omega}(\Omega;\mathbb{R}^N) \}$$
$$W^{1,q}_{0,\omega}(\Omega;\mathbb{R}^N) := W^{1,1}_0(\Omega;\mathbb{R}^N) \cap W^{1,q}_{\omega}(\Omega;\mathbb{R}^N).$$

Next, for $p \in [1, \infty)$, we say that a weight ω belongs to the Muckenhoupt class \mathcal{A}_p if and only if there exists a positive constant A such that for every cube $Q \subset \mathbb{R}^n$ the following holds

(2.1)
$$\left(\oint_{Q} \omega \, \mathrm{d}x \right) \left(\oint_{Q} \omega^{-(p'-1)} \, \mathrm{d}x \right)^{\frac{1}{p'-1}} \le A \qquad \text{if } p \in (1, \infty),$$

(2.2)
$$M\omega(x) \le A\omega(x)$$
 if $p = 1$.

In what follows, we denote by $A_p(\omega)$ the smallest constant A for which the inequality (2.1), resp. (2.2), holds.

The next result makes a very useful link between the maximal operator and \mathcal{A}_p -weights.

Lemma 2.1 (See pages 229–230 in [20] and page 5 in [21]). Let $f \in L^1_{loc}(\mathbb{R}^n)$ be such that $Mf < \infty$ almost everywhere in \mathbb{R}^n . Then for all $\alpha \in (0,1)$ we have $(Mf)^{\alpha} \in \mathcal{A}_1$. Furthermore, for all $p \in (1,\infty)$ and all $\alpha \in (0,1)$ there holds $(Mf)^{-\alpha(p-1)} \in \mathcal{A}_p$.

3. Relative Truncation

In this section we introduce a relative truncation. It is strongly influenced by the Lipschitz truncation method, developed in [10] see also [2, 7] where the concept was refined in the direction which can be adapted here. In contrast to the Lipschitz truncation, the *relative truncation* smoothens the function relative to a different independent function, or more explicit, it truncates the gradient on any given open set.

The first step is a suitable covering. We take the covering introduced in [8, Proposition 3.17] and proof some more properties needed for our situation.

Proposition 3.1. Let \mathcal{O} be an open subset of \mathbb{R}^d , with $\mathcal{O} \neq \mathbb{R}^d$. Then there exists a countable family Q_i of closed, dyadic cubes such that

- (a) $\bigcup_i Q_i = \mathcal{O}$ and all cubes Q_i have disjoint interiors.
- (b) $\operatorname{diam}(Q_i) < \operatorname{dist}(Q_i, \mathcal{O}^c) \leq 4\operatorname{diam}(Q_i)$
- (c) If $Q_i \cap Q_j \neq \emptyset$, then $\operatorname{diam}(Q_i) \leq 2\operatorname{diam}(Q_j) \leq 4\operatorname{diam}(Q_i)$.
- (d) For given Q_i , there exists at most $4^d 2^d$ cubes Q_j touching Q_i (boundaries intersect but not the interiors), we define A_i as the index set of all neighboring cubes of Q_i .
- (e) The family of cubes $\{\frac{3}{2}Q_i\}_{i\in\mathbb{N}}$ has finite intersection. The family can be split in 4^d-2^d pairwise disjoint subfamilies.
- (f) There is a partition of unity, $\psi_i \in C^{0,1}(\frac{9}{8}Q_i)$, such that $\chi_{\frac{1}{2}Q_i} \leq \psi_i \leq \chi_{\frac{9}{8}Q_i}$ and $\operatorname{diam}(Q_i)|\nabla \psi_i| \leq c(d)$ uniformly.

Proof. Since (a)–(d) are stated in [8, Proposition 3.17] we just proof the last two. First we show that the cubes $\frac{3}{2}Q_i$ have finite intersection. Firstly, by (b), we find that $\frac{3}{2}Q_i \subset \mathcal{O}$, which implies bu (a), that $\bigcup_i \frac{3}{2}Q_i = \bigcup_i Q_i$. Secondly, we find by (c) that $\frac{3}{2}Q_j$ only intersects its neighbors which implies by (d) that each $x \in Q_i$ is at most covered by $4^d - 2^d$ cubes of the family $\frac{3}{2}Q_j$. Moreover, by (c) we find that the $\frac{9}{8}Q_i$ does not intersect with $\frac{1}{2}Q_j$, for all $j \neq i$. Therefore it is standard to construct a partition of unity as requested.

Next we introduce the relative truncation of $u \in W_0^{1,p}(\Omega; \mathbb{R}^N)$, with respect to \mathcal{O} an open proper set. Since we do not assume, that $\mathcal{O} \subset \Omega$ we extend u by 0 outside Ω and define

$$u_{\mathcal{O}}(x) := \begin{cases} u(x) & x \in \mathbb{R}^d \setminus \mathcal{O}, \\ \sum_{i} \psi_i \bar{u}_i & x \in \mathcal{O}. \end{cases}$$

Where the \bar{u}_i are defined via the covering constructed in Proposition 3.1:

$$\bar{u}_i := \begin{cases} \int\limits_{\frac{9}{8}Q_i} u \, \mathrm{d}x & \text{if } \frac{9}{8}Q_i \subset \Omega, \\ 0 & \text{otherwise.} \end{cases}$$

At first we have to proof the following lemma, which is essential to show that the *relative* truncation is stable in Sobolev function spaces.

Lemma 3.2. Let Ω be a Lipschitz domain and let $u \in W_0^{1,p}(\Omega; \mathbb{R}^N)$, for $p \in [1, \infty)$. Then for any Q_i, Q_j which are members of the covering introduced in Proposition 3.1 with non-empty intersection, we have

(3.1)
$$\frac{|\bar{u}_j - \bar{u}_i|}{\operatorname{diam}(Q_i)} \le c(d, \Omega) \int_{\frac{3}{2}Q_i} |\nabla u| \, \mathrm{d}x + c(d, \Omega) \int_{\frac{3}{2}Q_j} |\nabla u| \, \mathrm{d}x$$

(3.2)
$$\oint_{\frac{9}{8}Q_i} \left| \frac{\bar{u} - \bar{u}_i}{\operatorname{diam}(Q_i)} \right|^p \le c(d, p, \Omega) \oint_{\frac{3}{2}Q_i} |\nabla u|^p \, \mathrm{d}x$$

Proof. The statement of (3.2), in case $\bar{u}_i \neq 0$ is just the Poincaré inequality. In case $\bar{u}_i = 0$ we enlarge the cube to $\frac{3}{2}Q_i$. Since we have a Lipschitz boundary and $\frac{9}{8}Q_i \cap \mathcal{O} \neq \emptyset$ we find that $|\frac{3}{2}Q_i| \leq c(\Omega)|\frac{3}{2}Q_i \cap \Omega^c|$. This implies, that we can apply Poincaré's inequality on u for the cube $\frac{3}{2}Q_i$, which finishes the proof of (3.2).

To proof (3.1) observe at first, that since Q_i and Q_j are neighbours, we find by (c) that $C_{i,j} = \frac{9}{8}Q_i \cap \frac{9}{8}Q_j$ have a comparible massure, to $|\frac{3}{2}Q_i|$ and $|\frac{3}{2}Q_j|$. Since in case $\bar{u}_i = \bar{u}_j = 0$, there is nothing to show, let us assume that $\bar{u}_i \neq 0$. By Poincaré we find

$$\begin{split} |\bar{u}_{j} - \bar{u}_{i}| &\leq \left| \bar{u}_{j} - \int_{C_{i,j}} u \, \mathrm{d}x \right| + \left| \int_{C_{i,j}} u - \langle u \rangle_{\frac{9}{8}Q_{i}} \, \mathrm{d}x \right| \\ &\leq \left| \bar{u}_{j} - \int_{C_{i,j}} u \, \mathrm{d}x \right| + c(d) \int_{\frac{9}{8}Q_{i}} |u - \langle u \rangle_{\frac{9}{8}Q_{i}} | \, \mathrm{d}x \leq \left| \bar{u}_{j} - \int_{C_{i,j}} u \, \mathrm{d}x \right| + c(d) \mathrm{diam}(Q_{i}) \int_{\frac{9}{8}Q_{i}} |\nabla u| \, \mathrm{d}x \end{split}$$

In case $\bar{u}_i \neq 0$ as well, then we find by symmetry

$$\left| \bar{u}_j - \oint_{C_{i,j}} u \, \mathrm{d}x \right| \le c(d) \mathrm{diam}(Q_j) \oint_{\frac{9}{8}Q_j} |\nabla u| \, \mathrm{d}x.$$

In case $\bar{u}_j = 0$ we enlarge the set $C_{i,j}$ to $\frac{3}{2}Q_j$ for which we know that $|\frac{3}{2}Q_j| \leq c(\Omega)|\frac{3}{2}Q_j \cap \Omega^c|$. Therefore, finally Poincaré's inequality implies

$$\oint_{C_{i,j}} |u| \, \mathrm{d}x \le c(d) \oint_{\frac{3}{2}Q_j} |u| \, \mathrm{d}x \le c(d,\Omega) \mathrm{diam}(Q_j) \oint_{\frac{3}{2}Q_j} |\nabla u| \, \mathrm{d}x.$$

This implies, that the *relative truncation* is stable in Sobolev spaces.

Lemma 3.3 (Stability). Let $u \in W_0^{1,p}(\Omega;\mathbb{R}^N)$, then $u_{\mathcal{O}} \in W_0^{1,p}(\Omega;\mathbb{R}^N)$. Moreover, the following estimate holds

$$\int_{\Omega} |\nabla (u - u_{\mathcal{O}})|^p dx \le c(d, p, \Omega) \int_{\mathcal{O} \cap \Omega} |\nabla u|^p dx.$$

Proof. It is enough to show the estimate since the zero-trace follows from the very definition of the relative truncation immediately. We define by A_i the index set of all j, such that $\frac{9}{8}Q_j \cap \frac{9}{8}Q_i \neq \emptyset$. Observe here, that $\#A_j \leq 4^n - 2^n$ by (d). Next we use (a), the fact that we have a partition of unity, (e), (c) and (f).

$$\int |\nabla (u - u_{\mathcal{O}})|^p \, \mathrm{d}x = \int_{\mathcal{O}} |\nabla (u - u_{\mathcal{O}})|^p \, \mathrm{d}x = \sum_i \int_{Q_i} \left| \nabla u - \sum_{j \in A_i} \nabla \psi_j \bar{u}_j \right|^p \, \mathrm{d}x$$

$$\leq \sum_i \int_{Q_i} \left| \sum_{j \in A_i} \nabla (u - \psi_j \bar{u}_j) \right|^p \, \mathrm{d}x \leq c(d, p) \sum_i \sum_{j \in A_i} \int_{Q_i} |\nabla (u - \psi_j \bar{u}_j)|^p \, \mathrm{d}x$$

$$\leq c(d, p) \sum_i \int_{Q_i} |\nabla u|^p \, \mathrm{d}x + c(d, p) \sum_i \sum_{j \in A_i} \int_{Q_i} \left| \frac{u - u_j}{\mathrm{diam}(Q_i)} \right|^p \, \mathrm{d}x$$

Now (a),(3.1) and (3.2) imply together with Jensen's inequality, (e) and (b) that

$$\int |\nabla (u - u_{\mathcal{O}})|^p dx \le c(d, p) \int_{\mathcal{O}} |\nabla u|^p dx + c(d, p, \Omega) \sum_i \int_{\frac{3}{2}Q_i} |\nabla u|^p dx \le c(d, p) \int_{\mathcal{O}} |\nabla u|^p dx.$$

4. Uniform a priori estimates

In this section, we will not mention the dependence of the constants on d, N anymore.

Proposition 4.1. Let $\Omega \subset \mathbb{R}^d$ be a bounded Lipschitz domain, S satisfy (1.2)–(1.3) and $h \in L^1(\Omega)$ not identically equal to zero. Then there exists $\varepsilon_0 > 0$ depending only on $\frac{C_2^{p'}}{C_1}$, p and Ω such that for any $\varepsilon \in (0, \varepsilon_0)$ the following holds. If $f \in L^p(\Omega; \mathbb{R}^{d \times N}) \cap L^p_{(Mh)^{-\varepsilon}}(\Omega; \mathbb{R}^{d \times N})$ and $u \in W_0^{1,p}(\Omega; \mathbb{R}^N)$ satisfies (1.1) in a weak sense, then $\nabla u \in L^p_{(Mh)^{-\varepsilon}}(\Omega; \mathbb{R}^{d \times N})$ as well. Moreover.

(4.1)
$$\int_{\Omega} \frac{|\nabla u|^p}{(Mh)^{\varepsilon}} dx \le C(C_1, C_2, p, \Omega) \int_{\Omega} \frac{|f|^p + C_3}{(Mh)^{\varepsilon}} dx.$$

Proof. First, we extend u and f with zero outside Ω . The function h, we will approximate by $g = h\chi_{\Omega} + \delta$. This implies, that $Mg > \delta$ and therefore $f, \nabla u \in L^p_{(Mg)^{-\varepsilon}}(\Omega; \mathbb{R}^{d \times N})$ a priori. Once the estimate is established for g with constants independent of δ the general result follows by letting $\delta \to 0$ and monotone convergence. For any $\lambda > 0$ we define $\mathcal{O}(\lambda) := \{x \in \mathbb{R}^d : Mg(x) > \lambda\}$. Since the maximal function is sub-linear it is an open set. In case $\mathcal{O}(\lambda) = \mathbb{R}^n$, we define $u_{\mathcal{O}(\lambda)} = 0$. Else, we are able to construct the relative truncation $u_{\mathcal{O}(\lambda)}$. Testing (1.1) with $u_{\mathcal{O}(\lambda)}$ which is possible due to Lemma 3.3 we get the identity

(4.2)
$$\int_{\{Mg \le \lambda\}} S \cdot \nabla u \, dx = \int_{\{Mg \le \lambda\}} |f|^{p-2} f \cdot \nabla u \, dx + \int_{\{Mg > \lambda\}} |f|^{p-2} f \cdot \nabla u_{\mathcal{O}(\lambda)} \, dx - \int_{\{Mg > \lambda\}} S \cdot \nabla u_{\mathcal{O}(\lambda)} \, dx.$$

Next, we focus on the estimates of integrals on the set where $Mg > \lambda$. Consider arbitrary $G \in L^{p'}(\Omega; \mathbb{R}^{d \times N})$ and arbitrary $\alpha \in (0,1)$. We get using (f)-the property of the partition of

unity, (3.1), (e) Hölder's inequality and (b)

$$\begin{split} &\int_{\{Mg>\lambda\}} |G\cdot\nabla u_{\mathcal{O}(\lambda)}| \,\mathrm{d}x \leq \sum_{i} \int_{Q_{i}} |G| \Big| \sum_{j\in A_{i}} \bar{u}_{j} \nabla \psi_{j} \Big| \,\mathrm{d}x \\ &= \sum_{i} \int_{Q_{i}} |G| \Big| \sum_{j\in A_{i}} (\bar{u}_{j} - \bar{u}_{i}) \nabla \psi_{j} \Big| \,\mathrm{d}x \\ &\leq c(\Omega) \sum_{i} \sum_{j\in A_{i}} |Q_{i}| \Big(\int_{Q_{i}} |G| \,\mathrm{d}x \Big) \Big(\int_{\frac{3}{2}Q_{i}} |\nabla u| \,\mathrm{d}x + \int_{\frac{3}{2}Q_{j}} |\nabla u| \,\mathrm{d}x \Big) \\ &= c(\Omega) \sum_{i} \sum_{j\in A_{i}} |Q_{i}| \Big(\int_{Q_{i}} \frac{|G|}{(Mg)^{\frac{\alpha}{p}}} (Mg)^{\frac{\alpha}{p}} \,\mathrm{d}x \Big) \Big(\int_{\frac{3}{2}Q_{i}} \frac{|\nabla u|}{(Mg)^{\frac{\alpha}{p'}}} (Mg)^{\frac{\alpha}{p'}} \,\mathrm{d}x + \int_{\frac{3}{2}Q_{j}} \frac{|\nabla u|}{(Mg)^{\frac{\alpha}{p'}}} (Mg)^{\frac{\alpha}{p'}} \,\mathrm{d}x \Big) \\ &\leq c(\Omega) \sum_{i} \sum_{j\in A_{i}} |Q_{i}| \Big(\int_{Q_{i}} \frac{|G|^{p'}}{(Mg)^{\frac{\alpha p'}{p}}} \,\mathrm{d}x \Big)^{\frac{1}{p'}} \Big(\Big(\int_{\frac{3}{2}Q_{i}} \frac{|\nabla u|^{p}}{(Mg)^{\frac{\alpha p}{p'}}} \,\mathrm{d}x \Big)^{\frac{1}{p}} + \Big(\int_{\frac{3}{2}Q_{i}} \frac{|\nabla u|^{p}}{(Mg)^{\frac{\alpha p}{p'}}} \,\mathrm{d}x \Big)^{\frac{1}{p}} \Big) \\ &\times \Big(\int_{5Q_{i}} (Mg)^{\alpha} \,\mathrm{d}x \Big). \end{split}$$

We estimate the last integral on the right hand side using two properties. First, since by Lemma 2.1 $(Mg)^{\alpha}$ is an \mathcal{A}_1 Muckenhoupt weight, $M(Mg)^{\alpha} \leq C(\alpha, d)(Mg)^{\alpha}$. Second, by (a) we know that $9Q_i \cap \mathcal{O}(\lambda)^c \neq \emptyset$. Consequently, for some $x_0 \in 9Q_i \cap \mathcal{O}(\lambda)^c$, we have

$$\int_{5Q_i} (Mg)^{\alpha} dx \le c \int_{9Q_i} (Mg)^{\alpha} dx \le cM(Mg)^{\alpha}(x_0)$$

$$\le c(\alpha)(Mg)^{\alpha}(x_0) \le c(\alpha)\lambda^{\alpha}.$$

We can use this to estimate (4.3) in the following way. For $i \in \mathbb{N}$ and $j \in A_i$, we deduce by Young's inequality

$$(4.4)$$

$$\lambda^{\alpha}|Q_{i}|\left(\int_{Q_{i}} \frac{|G|^{p'}}{(Mg)^{\frac{\alpha p'}{p}}} dx\right)^{\frac{1}{p'}} \left(\int_{\frac{3}{2}Q_{j}} \frac{|\nabla u|^{p}}{(Mg)^{\frac{\alpha p}{p'}}} dx\right)^{\frac{1}{p}}$$

$$\leq c(\Omega, \alpha)|Q_{i}|\left(\int_{Q_{i}} \frac{\lambda^{\frac{\alpha p'}{p}}|G|^{p'}}{(Mg)^{\frac{\alpha p'}{p}}} dx + \int_{\frac{3}{2}Q_{j}} \frac{\lambda^{\frac{\alpha p}{p'}}|\nabla u|^{p}}{(Mg)^{\frac{\alpha p}{p'}}} dx\right).$$

Therefore (4.3) implies using (e), that

$$(4.5) \qquad \int_{\{Mg>\lambda\}} |G \cdot \nabla u_{\mathcal{O}(\lambda)}| \, \mathrm{d}x \le c(\Omega, \alpha) \int_{\{Mg>\lambda\} \cap \Omega} \frac{\lambda^{\frac{\alpha p'}{p}} |G|^{p'}}{(Mg)^{\frac{\alpha p'}{p}}} \, \mathrm{d}x + \frac{\lambda^{\frac{\alpha p}{p'}} |\nabla u|^p}{(Mg)^{\frac{\alpha p}{p'}}} \, \mathrm{d}x.$$

Consequently, we get from (4.2) that

$$\int_{\{Mg \leq \lambda\}} S \cdot \nabla u \, \mathrm{d}x \leq \int_{\{Mg \leq \lambda\}} |f|^{p-1} |\nabla u| \, \mathrm{d}x + c(\Omega, \alpha) \int_{\{Mg > \lambda\}} \frac{\lambda^{\frac{\alpha p'}{p}} (|f|^p + |S|^{p'})}{(Mg)^{\frac{\alpha p'}{p}}} + \frac{\lambda^{\frac{\alpha p}{p'}} |\nabla u|^p}{(Mg)^{\frac{\alpha p}{p'}}} \, \mathrm{d}x.$$

We set $\overline{(p-1)} := \min((p-1), (p-1)^{-1})$ and take $\varepsilon \in (0, \alpha \overline{(p-1)})$. We multiply the above inequality by $\lambda^{-1-\varepsilon}$ and integrate over $\lambda \in (0, \infty)$ to deduce

$$(4.7) \qquad \int_{0}^{\infty} \int_{\{Mg \leq \lambda\}} \frac{S \cdot \nabla u}{\lambda^{1+\varepsilon}} \, \mathrm{d}x \, \mathrm{d}\lambda \leq \int_{0}^{\infty} \int_{\{Mg \leq \lambda\}} \frac{|f|^{p-1} |\nabla u|}{\lambda^{1+\varepsilon}} \, \mathrm{d}x \, \mathrm{d}\lambda + c(\Omega, \alpha) \int_{0}^{\infty} \int_{\{Mg > \lambda\}} \frac{\lambda^{\frac{\alpha p'}{p} - 1 - \varepsilon} (|f|^{p} + |S|^{p'})}{(Mg)^{\frac{\alpha p'}{p'}}} + \frac{\lambda^{\frac{\alpha p}{p'} - 1 - \varepsilon} |\nabla u|^{p}}{(Mg)^{\frac{\alpha p}{p'}}} \, \mathrm{d}x \, \mathrm{d}\lambda.$$

We get on the one hand using the Fubini theorem

$$\frac{1}{\varepsilon} \int_{\Omega} \frac{S \cdot \nabla u}{(Mg)^{\varepsilon}} \, \mathrm{d}x = \int_{\Omega} \int_{Mg}^{\infty} \frac{S \cdot \nabla u}{\lambda^{1+\varepsilon}} \, \mathrm{d}\lambda \, \mathrm{d}x = \int_{0}^{\infty} \int_{\{Mg < \lambda\}} \frac{S \cdot \nabla u}{\lambda^{1+\varepsilon}} \, \mathrm{d}x \, \mathrm{d}\lambda;$$

Thus on the other hand by (4.7) and the Fubini theorem

$$\begin{split} &\frac{1}{\varepsilon} \int_{\Omega} \frac{S \cdot \nabla u}{(Mg)^{\varepsilon}} \, \mathrm{d}x \leq \int_{0}^{\infty} \int_{\{Mg \leq \lambda\}} \frac{|f|^{p-1}|\nabla u|}{\lambda^{1+\varepsilon}} \, \mathrm{d}x \, \mathrm{d}\lambda \\ &+ c(\Omega, \alpha) \int_{0}^{\infty} \int_{\{Mg > \lambda\}} \frac{\lambda^{\frac{\alpha p'}{p} - 1 - \varepsilon} (|f|^{p} + |S|^{p'})}{(Mg)^{\frac{\alpha p'}{p}}} + \frac{\lambda^{\frac{\alpha p}{p'} - 1 - \varepsilon} |\nabla u|^{p}}{(Mg)^{\frac{\alpha p}{p'}}} \, \mathrm{d}\lambda \, \mathrm{d}x \\ &= \int_{\Omega} \int_{Mg}^{\infty} \frac{|f|^{p-1}|\nabla u|}{\lambda^{1+\varepsilon}} \, \mathrm{d}\lambda + c(\Omega, \alpha) \int_{0}^{Mg} \frac{\lambda^{\frac{\alpha p'}{p} - 1 - \varepsilon} (|f|^{p} + |S|^{p'})}{(Mg)^{\frac{\alpha p'}{p}}} + \frac{\lambda^{\frac{\alpha p}{p'} - 1 - \varepsilon} |\nabla u|^{p}}{(Mg)^{\frac{\alpha p}{p'}}} \, \mathrm{d}\lambda \, \mathrm{d}x \\ &= \frac{1}{\varepsilon} \int_{\Omega} \frac{|f|^{p-1}|\nabla u|}{(Mg)^{\varepsilon}} \, \mathrm{d}x + c(\Omega, \alpha) \int_{\Omega} \frac{1}{\frac{\alpha p'}{p} - \varepsilon} \frac{(|f|^{p} + |S|^{p'})}{(Mg)^{\varepsilon}} + \frac{1}{\frac{\alpha p}{p'} - \varepsilon} \frac{|\nabla u|^{p}}{(Mg)^{\varepsilon}} \, \mathrm{d}x \\ &\leq \frac{1}{\varepsilon} \int_{\Omega} \frac{|f|^{p-1}|\nabla u|}{(Mg)^{\varepsilon}} \, \mathrm{d}x + \frac{c(\Omega, \alpha)}{\alpha (p-1) - \varepsilon} \int_{\Omega} \frac{|f|^{p} + |S|^{p'} + |\nabla u|^{p}}{(Mg)^{\varepsilon}} \, \mathrm{d}x. \end{split}$$

By the assumptions (1.2),(1.3) and Young's inequality we deduce

$$(4.8) \qquad \int_{\Omega} \frac{|\nabla u|^p}{(Mg)^{\varepsilon}} \, \mathrm{d}x \le \frac{c(p)}{C_1} \int_{\Omega} \frac{|f|^p}{(Mg)^{\varepsilon}} \, \mathrm{d}x + \frac{\varepsilon C_2^{p'} c(\Omega, p)}{C_1 \alpha(\overline{(p-1)} - \varepsilon)} \int_{\Omega} \frac{|f|^p + |\nabla u|^p + C_3}{(Mg)^{\varepsilon}} \, \mathrm{d}x.$$

Thus, for $\varepsilon \in (0, \varepsilon_0)$, we can absorb the term with $|\nabla u|^p$ on the right hand side of (4.8) into the left hand side and we get (4.1) by letting $\delta \to 0$. Here

(4.9)
$$\varepsilon_0 = \frac{C_1 \overline{(p-1)}}{C_2^{p'} c(\Omega, p)},$$

since by this choice we can choose for $\varepsilon \in (0, \varepsilon_0)$ an α accordingly. Observe, that $\alpha \to 1$ (which is the non-stable limit), when $\varepsilon \to \varepsilon_0$.

5. Existence of a solution

The proof is making essential use of the following theorem that can be found in [4, Theorem 2.6].

Theorem 5.1 (weighted, biting div-curl lemma). Let $\Omega \subset \mathbb{R}^n$ be an open bounded set. Assume that for some $p \in (1, \infty)$ and given $\omega \in \mathcal{A}_p$ we have a sequence of vector-valued measurable functions $(a^k, b^k)_{k=1}^{\infty} : \Omega \to \mathbb{R}^n \times \mathbb{R}^n$ such that

(5.1)
$$\sup_{k \in \mathbb{N}} \int_{\Omega} |a^k|^p \omega + |b^k|^{p'} \omega \, \mathrm{d}x < \infty.$$

Furthermore, assume that for every bounded sequence $\{c^k\}_{k=1}^{\infty}$ from $W_0^{1,\infty}(\Omega)$ that fulfills

$$\nabla c^k \rightharpoonup^* 0$$
 weakly* in $L^{\infty}(\Omega)$

there holds

(5.2)
$$\lim_{k \to \infty} \int_{\Omega} b^k \cdot \nabla c^k \, \mathrm{d}x = 0,$$

(5.3)
$$\lim_{k \to \infty} \int_{\Omega} a_i^k \partial_{x_j} c^k - a_j^k \partial_{x_i} c^k \, \mathrm{d}x = 0 \qquad \qquad \text{for all } i, j = 1, \dots, n.$$

Then there exists a subsequence (a^k, b^k) that we do not relabel and there exists a nondecreasing sequence of measurable subsets $E_i \subset \Omega$ with $|\Omega \setminus E_j| \to 0$ as $j \to \infty$ such that

$$(5.4) a^k \rightharpoonup a weakly in L^1(\Omega; \mathbb{R}^n),$$

$$(5.5) b^k \rightharpoonup b weakly in L^1(\Omega; \mathbb{R}^n).$$

$$(5.6) a^k \cdot b^k \omega \rightharpoonup a \cdot b \omega weakly in L^1(E_j) for all j \in \mathbb{N}.$$

Proof of Theorem 1.2. As before, we extend every function by zero outside Ω , without further reference. We approximate a given $f \in L^q(\Omega; \mathbb{R}^{d \times N})$, by $f_k := \min\{k, |f|\}f/|f|$. Then $f_k \in L^q(\Omega; \mathbb{R}^{d \times N}) \cap L^\infty(\Omega; \mathbb{R}^{d \times N})$, we find $|f^k| \nearrow |f|$ and

(5.7)
$$f^k \to f$$
 strongly in $L^q(\mathbb{R}^n; \mathbb{R}^{n \times N})$.

For f^k we can use the standard monotone operator theory to find a solution $u^k \in W_0^{1,p}(\Omega;\mathbb{R}^N)$ fulfilling

(5.8)
$$\int_{\Omega} S(x, \nabla u^k) \cdot \nabla \varphi \, \mathrm{d}x = \int_{\Omega} f^k \cdot \nabla \varphi \, \mathrm{d}x \qquad \text{for all } \varphi \in W_0^{1,p}(\Omega; \mathbb{R}^N).$$

Hence, we fix $\varepsilon = p - q \in (0, \varepsilon_0)$. Then we find by (4.1) and the continuity of the maximal function that

(5.9)
$$\int_{\Omega} \frac{|\nabla u|^p}{(M(f+1))^{\varepsilon}} dx \leq C(C_1, C_2, p, \Omega) \int_{\Omega} \frac{|f_k|^p + C_3}{(M(f+1))^{\varepsilon}} dx \\ \leq c \int M(f)^q dx + cC_3 \leq c \int |f|^q dx + cC_3.$$

Moreover, by Young's inequality for the exponents $\frac{q}{p} + \frac{p-q}{p} = 1$, continuity of the maximal function and the previous we gain

(5.10)
$$\int_{\Omega} |\nabla u|^{q} dx = \int_{\Omega} \frac{|\nabla u|^{q} (M(f+1))^{\frac{(p-q)q}{p}}}{(M(f+1))^{\frac{(p-q)q}{p}}} dx$$
$$\leq c \int_{\Omega} \frac{|\nabla u|^{p}}{(M(f+1))^{\varepsilon}} dx + c \int_{\Omega} (M(f+1))^{q} dx \leq c \int |f|^{q} dx + c(C_{3}+1).$$

We define the weight $\omega: \frac{1}{(M(f+1))^{\varepsilon}}$. Using the a priori estimate, the reflexivity of the corresponding spaces and the growth assumption (1.3), we can pass to a subsequence (still denoted by u^k) such that

(5.11)
$$u^k \rightharpoonup u$$
 weakly in $W_0^{1,q}(\Omega; \mathbb{R}^N)$, (5.12) $\nabla u^k \rightharpoonup \nabla u$ weakly in $L^p_\omega \cap L^q(\Omega; \mathbb{R}^N)$, weakly in $L^p_\omega \cap L^q(\Omega; \mathbb{R}^N)$

(5.12)
$$\nabla u^k \rightharpoonup \nabla u \qquad \text{weakly in } L^p_\omega \cap L^q(\Omega; \mathbb{R}^{n \times N}),$$

(5.13)
$$S(x, \nabla u^k) \rightharpoonup \overline{S}$$
 weakly in $L^p_{\Omega} \cap L^q(\Omega; \mathbb{R}^{n \times N})$.

Hence by (5.10),(5.9) and 5.12 and the weak lower semicontinuity we obtain

(5.14)
$$\int_{\Omega} |\nabla u|^q + |\nabla u|^p \omega \, \mathrm{d}x \le c \int_{\Omega} |f|^q \, \mathrm{d}x + c,$$

which concludes the a priori estimate.

We still have to show that u is a distributional solution. Using (5.8), (5.7) and (5.13) it follows that

(5.15)
$$\int_{\Omega} \overline{S} \cdot \nabla \varphi \, \mathrm{d}x = \int_{\Omega} f \cdot \nabla \varphi \, \mathrm{d}x \quad \text{for all } \varphi \in C_0^{\infty}(\Omega; \mathbb{R}^N).$$

To complete the proof of Theorem 1.2, it remains to show that

(5.16)
$$\overline{S}(x) = S(x, \nabla u(x)) \quad \text{in } \Omega.$$

To do so, we use Theorem 5.1. We denote $a^k := \nabla u^k$ and $b^k := S(x, \nabla u^k)$. By using (5.14) and (1.3), we find that (5.1) is satisfied with the weight ω . Since, we assume by (4.9) that $\varepsilon < (p-1)$, we find by Lemma 2.1 that $\omega \in A_p$. Also the assumption (5.2) holds, which follows from 5.7, (5.8) and (5.15). Finally, (5.3) is valid trivially since a^k is a gradient. Therefore, Theorem 5.1 can be applied. Meaning, that we have a non-decreasing sequence of measurable sets E_j , such that $|\Omega \setminus E_j| \to 0$ and

$$S(x, \nabla u^k) \cdot \nabla u^k \omega \rightharpoonup \overline{S} \cdot \nabla u \omega$$
 weakly in $L^1(E_i)$.

This is enough, to apply some variant of the Minty trick. For any $G \in L^p_\omega(\Omega; \mathbb{R}^{n \times N})$ we get by 5.12 and 5.13

$$(S(x, \nabla u^k) - S(x, G)) \cdot (\nabla u^k - G) \omega \rightharpoonup (\overline{S} - S(x, G)) \cdot (\nabla u - G) \omega$$
 weakly in $L^1(E_j)$.

Due to the monotonicity condition (1.4) we find that the term on the left hand side is non-negative and consequently its weak limit is non-negative as well; especially $\int_{E_j} (\overline{S} - S(x, G)) \cdot (\nabla u - G) \omega \, dx \ge 0$ and

$$\int_{\Omega} (\overline{S} - S(x, G)) \cdot (\nabla u - G) \,\omega \, dx \ge \int_{\Omega \setminus E_j} (\overline{S} - S(x, G)) \cdot (\nabla u - G) \,\omega \, dx.$$

Letting $j \to \infty$ the Lebesgue dominated convergence theorem implies using the fact that $|\Omega \setminus E_j| \to 0$ as $j \to \infty$ we obtain

(5.17)
$$\int_{\Omega} (\overline{S} - S(x, G)) \cdot (\nabla u - G) \, \omega \, dx \ge 0 \quad \text{for all } G \in L^p_{\omega}(\Omega; \mathbb{R}^{n \times N}).$$

Hence, setting $G := \nabla u - \delta H$ where $H \in L^{\infty}(\Omega; \mathbb{R}^{n \times N})$ is an arbitrary function and dividing (5.17) by δ implies

$$\int_{\Omega} (\overline{S} - S(x, \nabla u - \delta H)) \cdot H \,\omega \,\mathrm{d}x \ge 0.$$

Finally, letting $\delta \to 0_+$ implies by the continuity assumption of S and dominated convergence

$$\int_{\Omega} (\overline{S} - S(x, \nabla u)) \cdot H \, \omega \, \mathrm{d}x \ge 0 \quad \text{for all } H \in L^{\infty}(\Omega; \mathbb{R}^{n \times N}).$$

Since ω is strictly positive almost everywhere in Ω , the relation (5.16) easily follows by choosing e.g.,

$$H := -\frac{\overline{S} - S(x, \nabla u)}{1 + |\overline{S} - S(x, \nabla u)|}.$$

Thus u is a distributional solution to (1.1).

Proof of Corollary 1.3. The proof of the a priori estimate (4.1) is exactly the same. Analogous to the estimate of (5.9), (5.10) one finds in case $C_3 = 0$, that

(5.18)
$$\int_{\Omega} |\nabla u_k|^q + |\nabla u_k|^p (Mf)^{q-p} \, \mathrm{d}x \le c \int_{\Omega} |f|^q.$$

The existence proof is also analogous. However, since Theorem 5.1 is only valid on bounded domains one has to use that \overline{S} satisfies (5.15) for an arbitrary bounded subset of Ω . This allows to show that $\overline{S} = S(\nabla u)$ in Ω' . Since it was arbitrarily chosen the existence follows. \square

References

- P. Bénilan, L. Boccardo, T. Gallouët, R. Gariepy, M. Pierre, and J. L. Vázquez. An L¹-theory of existence and uniqueness of solutions of nonlinear elliptic equations. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), 22(2):241–273, 1995.
- [2] D. Breit, L. Diening, and S. Schwarzacher. Solenoidal lipschitz truncation for parabolic pdes. Math. Models Methods Appl. Sci., 53(14):2671–2700, 2013.
- [3] M. Bulíček. On continuity properties of monotone operators beyond the natural domain of definition. *Manuscripta Math.*, 138(3-4):287–298, 2012.
- [4] M. Bulíček, L. Diening, and S. Schwarzacher. Existence, uniqueness and optimal regularity results for very weak solutions to nonlinear elliptic systems. Preprint MORE/2015/03, 2015.
- [5] L. A. Caffarelli and I. Peral. On $W^{1,p}$ estimates for elliptic equations in divergence form. Comm. Pure Appl. Math., 51(1):1-21, 1998.
- [6] E. DiBenedetto and J. Manfredi. On the higher integrability of the gradient of weak solutions of certain degenerate elliptic systems. Amer. J. Math., 115(5):1107–1134, 1993.
- [7] L. Diening. Function spaces and inequalities, lectures spring school on analysis, pasecky. *matfyzpress*, pages 1–23, 2013.
- [8] L. Diening and S. Fröschl. Extensions in spaces with variable exponents—the half space. In *Harmonic analysis and nonlinear partial differential equations*, RIMS Kôkyûroku Bessatsu, B22, pages 71–92. Res. Inst. Math. Sci. (RIMS), Kyoto, 2010.
- [9] L. Diening, P. Kaplický, and S. Schwarzacher. BMO estimates for the *p*-Laplacian. *Nonlinear Anal.*, 75(2):637–650, 2012.
- [10] L. Diening, M. Růžička, and J. Wolf. Existence of weak solutions for unsteady motions of generalized Newtonian fluids. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5), 9(1):1–46, 2010.
- [11] M. Giaquinta and E. Giusti. On the regularity of the minima of variational integrals. Acta Math., 148:31–46, 1982.
- [12] L. Greco, T. Iwaniec, and C. Sbordone. Variational integrals of nearly linear growth. Differential Integral Equations, 10(4):687–716, 1997.
- [13] T. Iwaniec. Projections onto gradient fields and L^p -estimates for degenerated elliptic operators. Studia Math., 75(3):293–312, 1983.
- [14] T. Iwaniec. p-harmonic tensors and quasiregular mappings. Ann. of Math. (2), 136(3):589-624, 1992.
- [15] J. Kinnunen and J. L. Lewis. Very weak solutions of parabolic systems of p-Laplacian type. $Ark.\ Mat.$, $40(1):105-132,\ 2002.$
- [16] J. Kristensen and G. Mingione. The singular set of minima of integral functionals. Arch. Ration. Mech. Anal., 180(3):331–398, 2006.
- [17] G. Mingione. Nonlinear aspects of Calderón-Zygmund theory. Jahresber. Dtsch. Math.-Ver., 112(3):159–191, 2010.
- [18] J. Nečas. Example of an irregular solution to a nonlinear elliptic system with analytic coefficients and conditions for regularity. In *Theory of nonlinear operators (Proc. Fourth Internat. Summer School, Acad. Sci., Berlin, 1975)*. Akademie-Verlag, Berlin, 1977.
- [19] J. Serrin. Isolated singularities of solutions of quasi-linear equations. Acta Math., 113:219–240, 1965.
- [20] A. Torchinsky. Real-variable methods in harmonic analysis, volume 123 of Pure and Applied Mathematics. Academic Press, Inc., Orlando, FL, 1986.
- [21] B. O. Turesson. Nonlinear potential theory and weighted Sobolev spaces, volume 1736 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, 2000.

Mathematical Institute, Faculty of Mathematics and Physics, Charles University in Prague Sokolovská 83, 186 75 Prague, Czech Republic

 $E ext{-}mail\ address: mbul8060@karlin.mff.cuni.cz}$

DEPARTMENT OF MATHEMATICAL ANALYSIS, FACULTY OF MATHEMATICS AND PHYSICS, CHARLES UNIVERSITY IN PRAGUE, SOKOLOVSKÁ 83, 186 75 PRAGUE, CZECH REPUBLIC

E-mail address: schwarz@karlin.mff.cuni.cz