# STABILITY PROPERTIES AND LARGE TIME BEHAVIOR OF VISCOSITY SOLUTIONS OF HAMILTON-JACOBI EQUATIONS ON METRIC SPACES

ATSUSHI NAKAYASU AND TOKINAGA NAMBA

ABSTRACT. We investigate asymptotic behaviors of a metric viscosity solution of a Hamilton-Jacobi equation defined on a general metric space in Gangbo-Święch sense. Our results include general stability and large time behavior of the solution.

### 1. INTRODUCTION

In this chapter we study stability of a solution of Hamilton-Jacobi equations on a general complete geodesic metric space space (X, d). Let H be a continuous function on  $X \times \mathbf{R}_+$  called a *Hamiltonian*. Consider a Cauchy problem of a Hamilton-Jacobi equation of the form

(1.1) 
$$\partial_t u + H(x, |Du|) = 0 \quad \text{in } (0, \infty) \times X,$$

(1.2) 
$$u|_{t=0} = u_t$$

with initial data  $u_0$  and a corresponding stationary equation of the form

(1.3) 
$$H(x, |Dv|) = c \quad \text{in } X$$

with some  $c \in \mathbf{R}$ .

The theory of Hamilton-Jacobi equations on generalized spaces have been developing in these years. For example, [17] and [3] study a stationary equations on topological networks and post-critically finite fractals including the Sierpinski gasket. In order to cover them a metric viscosity solution is posed, which means a theory of viscosity solutions on a general metric space. A notion of metric viscosity solution was first introduced by Giga-Hamamuki-Nakayasu [11] to the stationary equation (1.3) in a spirit of [17]. It was attempted to apply this idea to the evolutionary equation (1.2) in [15]. Afterwards some different notions of metric viscosity solutions were proposed by several authors; see, e.g., [1], [8] [9]. In particular, the metric viscosity solution by Gangbo-Święch [8], [9] is apparently compatible with stability argument. In fact, the authors of [8] construct a solution of (1.2) by Perron method while the other materials show a representation formula of a metric viscosity solution.

The main aim of the present work is to establish a general stability result for the Gangbo-Święch solutions. Roughly speaking, the stability is the proposition claiming that the semilimit of a family of viscosity solutions is a viscosity solution; see [2, Theorem A.2]. At least in the classical theory of viscosity solutions the stability is a fundamental property to derive some asymptotic behavior of the solution. Large time asymptotics of a viscosity solution of Hamilton-Jacobi equations is studied by

Date: June 26, 2018.

Namah-Roquejoffre [16] and Fathi [6] independently. Another aim of this chapter is to establish, as a consequence of the stability, a large time asymptotic behavior of the solution on a singular space such as the Sierpinski gasket. Based on the argument in [16] we will show that the solution u(t, x) + ct of (1.2) goes to a function v as  $t \to \infty$  and v is a solution of the stationary problem (1.3) with some constant  $c \in \mathbf{R}$ .

We restrict ourselves to the case when the metric space X is compact but the Sierpinski gasket can be handled. Let us extend H to  $X \times \mathbf{R}$  as an even function H(x, p) = H(x, |p|). The basic assumptions on the Hamiltonian are:

- (A1) H is continuous.
- (A2) H is convex in the second variable.
- (A3) H is coercive in the sense of

$$\lim_{p \to \infty} \inf_{x} H(x, p) = \infty.$$

(A4)  $\sup_x H(x,0) < \infty$ .

Set  $c := \sup_x H(x, 0)$ ; otherwise the stationary equation (1.3) has no solution. We first show that the stationary equation has at least one solution. Then, a standard barrier method implies that there exist upper and lower semilimits  $\overline{u}(x)$  and  $\underline{u}(x)$  of u(t, x) + ct as  $t \to \infty$  as real-valued functions. As a result the stability argument yields that  $\overline{u}$  and  $\underline{u}$  are a subsolution and a supersolution of the limit equation. Next note that for each  $x \in A := \{x \in X \mid H(x, 0) = \sup_x H(x, 0)\}$  the solution u(t, x) + ct is non-increasing since  $u_t + c \leq 0$  and so  $\overline{u} = \underline{u}$  on A by Dini's theorem. We see that  $\overline{u} = \underline{u}$  by a comparison principle for the stationary equation. This means that u(t, x) + ct converges to a solution  $\overline{u} = \underline{u}$  of the stationary equation locally uniformly.

In order to justify this argument we will establish solvability of (1.3) and a comparison principle for Gangbo-Święch solutions of (1.3), which are new. The authors of [16] invoke a result by Lions-Papanicolaou-Varadhan [14]. The argument is based on the ergodic theory but in this work we will follow a direct approach via Perron method by Fathi-Siconolfi [7].

# 2. Definition of Gangbo-Święch solutions

In this section we review the definition of metric viscosity solutions proposed by Gangbo and Święch; see [8] and [9]. Let (X, d) be a complete geodesic metric space.

For a real-valued function u on an open subset Q of the spacetime  $\mathbf{R} \times X$  define the *upper local slope* and *lower local slope* 

$$\begin{aligned} |\nabla^+ u|(t,x) &:= \limsup_{y \to x} \frac{[u(t,y) - u(t,x)]_+}{d(y,x)}, \\ |\nabla^- u|(t,x) &:= \limsup_{y \to x} \frac{[u(t,y) - u(t,x)]_-}{d(y,x)}. \end{aligned}$$

and the local slope

$$|\nabla u|(t,x) := \limsup_{y \to x} \frac{|u(t,y) - u(t,x)|}{d(y,x)}$$

It is easy to see that  $|\nabla^{-}u| = |\nabla^{+}(-u)|$ .

We next introduce smoothness classes for functions on a metric space.

**Definition 2.1.** We denote by  $\mathcal{C}(Q)$  the set of all functions u on Q such that u is locally Lipschitz continuous on Q and  $\partial_t u$  is continuous on Q. We also set

$$\overline{\mathcal{C}}^{1}(Q) := \{ u \in \mathcal{C}(Q) \mid |\nabla^{+}u| = |\nabla u| \text{ and they are continuous} \},\$$

$$\underline{\mathcal{C}}^{1}(Q) := \{ u \in \mathcal{C}(Q) \mid |\nabla^{-}u| = |\nabla u| \text{ and they are continuous} \}.$$

**Lemma 2.2.** Let  $u(t,x) := a(t)\phi(d(x,y)^2) + b(t)$  with  $y \in X$ ,  $\phi \in C^1(\mathbf{R}_+)$ ,  $\phi' \ge 0$ ,  $a, b \in C^1(\mathbf{R})$ . Then,  $u \in \underline{C}^1(\mathbf{R} \times X)$  and moreover

$$|\nabla^{-}u|(t,x) = |\nabla u|(t,x) = 2a(t)\phi'(d(x,y)^{2})d(x,y).$$

See [1] or [8] for the proof.

We consider a Hamilton-Jacobi equation of the form

(2.1) 
$$F(z, |Du|, \partial_t u) = 0 \quad \text{in } Q.$$

Here, z = (t, x), and  $F = F(z, p, q) \in C(Q \times \mathbf{R} \times \mathbf{R})$  is even and convex in p and strictly increasing in q. Set

$$F_r(z, p, q) := \begin{cases} \sup_{p' \in B} F(z, p + rp', q) & \text{if } r \ge 0\\ \inf_{p' \in B} F(z, p + rp', q) & \text{if } r \le 0 \end{cases}$$

for  $r \in \mathbf{R}$ , where B := [-1, 1]. Note that  $(z, p, q, r) \mapsto F_r(z, p, q)$  is continuous since  $(z, p, q, r, p') \mapsto F(z, p + rp', q)$  is continuous and B is compact. Also, it is easy to check  $r \mapsto F_r(z, p, q)$  is non-decreasing.

For a function u defined on Q with values in the extended real numbers  $\overline{\mathbf{R}} := \mathbf{R} \cup \{\pm \infty\}$ , we take its upper and lower semicontinuous envelope  $u^*$  and  $u_*$ .

**Definition 2.3** (Metric viscosity solutions of (2.1)). Let u be an **R**-valued function on Q.

We say that u is a *metric viscosity subsolution* (resp. *supersolution*) of (2.1) when for every  $\psi = \psi_1 + \psi_2$  with  $\psi_1 \in \underline{C}^1(Q)$  (resp.  $\psi_1 \in \overline{C}^1(Q)$ ) and  $\psi_2 \in \mathcal{C}(Q)$ , if  $u^* - \psi$  (resp.  $u_* - \psi$ ) attains a zero local maximum (resp. minimum) at a point  $z = (t, x) \in Q$ , i.e.  $(u^* - \psi)(z) = \max_{B_R(z)}(u^* - \psi) = 0$  (resp.  $(u_* - \psi)(z) = \min_{B_R(z)}(u_* - \psi) = 0$ ) for some R > 0, then

$$F_{-|\nabla\psi_{2}|^{*}(z)}(z, |\nabla\psi_{1}|(z), \partial_{t}\psi(z)) \leq 0 \text{ (resp. } F_{|\nabla\psi_{2}|^{*}(z)}(z, |\nabla\psi_{1}|(z), \partial_{t}\psi(z)) \geq 0.)$$

We say that u is a *metric viscosity solution* of (2.1) if u is both a metric viscosity subsolution and a metric viscosity supersolution of (2.1).

By a similar way we also define a notion of metric viscosity solutions for a stationary equation of the form

$$H(x, |Dv|) = 0 \quad \text{in } U$$

with  $U \subset X$  open. Here,  $H = H(x, p) \in C(U \times \mathbf{R})$  is even and convex in p. Note that one is able to define the local slopes  $|\nabla^{-}v|$ ,  $|\nabla^{+}v|$ ,  $|\nabla v|$  and smoothness  $\mathcal{C}(U)$ ,  $\overline{\mathcal{C}}^{1}(U)$ ,  $\underline{\mathcal{C}}^{1}(U)$  for a function v on U.

**Definition 2.4** (Metric viscosity solutions of (2.2)). Let v be an **R**-valued function on U.

We say that v is a metric viscosity subsolution (resp. supersolution) of (2.2) when for every  $\psi = \psi_1 + \psi_2$  with  $\psi_1 \in \underline{C}^1(U)$  (resp.  $\psi_1 \in \overline{C}^1(U)$ ) and  $\psi_2 \in \mathcal{C}(Q)$ , if  $v^* - \psi$  (resp.  $v_* - \psi$ ) attains a zero local maximum (resp. minimum) at a point  $x \in U$ , then

 $H_{-|\nabla \psi_2|^*(x)}(x, |\nabla \psi_1|(x)) \le 0 \text{ (resp. } H_{|\nabla \psi_2|^*(x)}(x, |\nabla \psi_1|(x)) \ge 0.)$ 

We say that v is a *metric viscosity solution* of (2.2) if v is both a metric viscosity subsolution and a metric viscosity supersolution of (2.2).

These notions satisfies the following natural propositions.

**Proposition 2.5** (Consistency). If u is a metric viscosity subsolution of (2.1) in  $I \times U$  with an open interval I and u is of the form u(t, x) = v(x), then v is a metric viscosity subsolution of (2.2) in U with H(x, p) := F(0, x, p, 0).

Conversely, if v is a metric viscosity subsolution of (2.2) in U, then u(t, x) := v(x) is a metric viscosity subsolution of (2.1) in  $\mathbf{R} \times U$  with F(t, x, p, a) := H(x, p).

**Proposition 2.6** (Transitive relation). Assume that F = F(z, p, q), G = G(z, p, q) satisfy  $G \leq F$  and let u be a metric viscosity subsolution of (2.1). Then, u is a metric viscosity subsolution of  $G(z, |Du|, \partial_t u) = 0$  in Q.

**Proposition 2.7** (Locality). Let  $Q_1$  and  $Q_2$  be two open subsets of  $(0, \infty) \times X$ . If u is a metric viscosity subsolution of (2.1) in  $Q_1$  and is a metric viscosity subsolution of (2.1) in  $Q_2$ , then u is a metric viscosity subsolution of (2.1) in  $Q = Q_1 \cup Q_2$ 

**Proposition 2.8** (Change of variable). Let  $\phi$  be a  $C^1$  diffeomorphism from an interval I to an interval J. If u is a metric viscosity subsolution of (2.1) in  $I \times U$ , then  $v(s, x) := u(\phi^{-1}(s), x)$  is a metric viscosity subsolution of

$$F(x, |Dv|, \phi'v_s) = 0$$
 in  $J \times U$ .

**Proposition 2.9** (Composition). Let a be a non-zero constant and b = b(t) be a  $C^1$  function on an interval I. If u is a metric viscosity subsolution of (2.1) in  $I \times U$ , then v(t, x) := au(t, x) + b(t) is a metric viscosity subsolution of

$$F(x, \frac{|Dv|}{a}, \frac{v_t - b'(t)}{a}) = 0 \quad in \ I \times U.$$

**Proposition 2.10** (Strong solutions). Let  $\psi_1 \in \overline{\mathcal{C}}^1(Q)$  and  $\psi_2 \in \mathcal{C}(Q)$ . If  $\psi = \psi_1 + \psi_2$  satisfies

$$F_{|\nabla\psi_2|^*(z)}(z, |\nabla\psi_1|(z), \partial_t\psi(z)) \le 0 \quad \text{for all } z \in Q,$$

then  $\psi$  is a metric viscosity subsolution of (2.1).

The proofs are straightforward so we omit them. For the proof of Proposition 2.10 see [9, Lemma 2.8].

# 3. Stability results

Let A be a topological space. For a family of functions  $\{u(\cdot; a)\}_{a \in A}$  defined on Q take its upper and lower semicontinuous envelopes

$$u^{*}(z;a) := \limsup_{(z',a') \to (z,a)} u(z';a'), \quad u_{*}(z;a) := \liminf_{(z',a') \to (z,a)} u(z';a').$$

The functions  $u^*(\cdot; a)$  and  $u^*(\cdot; a)$  are respectively called the upper and lower semilimit of  $\{u(\cdot; a)\}$  at  $a \in A$ . Also note that  $u^*(\cdot; a)$  is upper semicontinuous and that for each (z, a) there exists a sequence  $(z_j, a_j)$  such that

$$(z_j, a_j, u(z_j; a_j)) \rightarrow (z, a, u^*(z; a))$$

One of the main results of this section is:

**Lemma 3.1** (Stability). Let  $F = F(z, p, q; a) \in C(Q \times \mathbb{R} \times \mathbb{R} \times A)$  and let  $u = u(\cdot; a)$  be a family of metric viscosity subsolutions (resp. supersolution) of (2.1) with  $F = F(\cdot; a)$ . Assume that  $a \in A$  satisfies for each  $z \in Q$  (3.1)

 $\limsup_{a' \to a} \sup_{B_r(z)} u(\cdot; a') \le \sup_{B_r(z)} u^*(\cdot; a) \text{ (resp. } \liminf_{a' \to a} \inf_{B_r(z)} u(\cdot; a') \le \inf_{B_r(z)} u_*(\cdot; a))$ 

for all r > 0 small enough. Then, the upper (resp. lower) semilimit  $\overline{u} := u^*(\cdot, a)$ (resp.  $\underline{u} := u^*(\cdot, a)$ ) is a metric viscosity subsolution (resp. supersolution) of (2.1) with  $F = F(\cdot; a)$ .

Remark 3.2. An sufficient condition of the assumption (3.1) is that the metric space X is locally compact. Indeed, since  $B := \overline{B}_r(z)$  is compact for small r, we are able to take a sequence of maximum points  $z_{a'}$  of  $u(\cdot; a')$  and assume that  $z_{a'}$  converges to some  $\overline{z} \in B$  as  $a' \to a$  by taking a subsequence if necessary. Then,

$$\limsup_{a' \to a} \sup_{B} u(\cdot; a') = \limsup_{a' \to a} u(z_{a'}; a') \le u^*(\bar{z}; a) \le \sup_{B} u^*(\cdot; a).$$

We also point out that if the assumption is removed, then the lemma may be false in general; see [11].

A direct consequence of Lemma 3.1 is:

**Corollary 3.3** (Stability under extremum). Let  $F = F(z, p, q) \in C(Q \times \mathbf{R} \times \mathbf{R})$ . Let S be a family of metric viscosity subsolutions (resp. supersolutions) of (2.1). Then  $\overline{u}(z) := \sup_{v \in S} v(z)$  (resp.  $\underline{u}(z) := \inf_{v \in S} v(z)$ ) is a metric viscosity subsolution (resp. supersolutions) of (2.1).

*Proof.* Set A = S with the indiscrete topology and trivial families  $\{F\}_{v \in S}$  and  $\{U(\cdot; v) = v\}_{v \in S}$ . Note that  $U^*(z; v) = u^*(z)$  and  $\limsup_{v' \to v} \sup_{B_r(z)} U(\cdot; v') = \sup_{B_r(z)} U^*(\cdot; v) = \sup_{B_r(z)} u^*$ . Therefore, by applying Lemma 3.1 we see that u is a metric viscosity subsolution of (2.1).

Our proof of Lemma 3.1 is inspired by [8]. First recall Ekeland's variational principle of a classical version [4], [5].

**Lemma 3.4** (Ekeland's variational principle). Let (X, d) be a complete metric space and let  $F: X \to \overline{\mathbf{R}}$  be a upper semicontinuous function bounded from above (resp. below) satisfying  $D(F) := \{F > -\infty\}$  (resp.  $D(F) := \{F < +\infty\}$ ) is not empty. Then, for each  $\hat{x} \in D(F)$ , there exists  $\overline{x} \in X$  such that  $d(\hat{x}, \overline{x}) \leq 1$ ,  $F(\overline{x}) \geq F(\hat{x})$ (resp.  $F(\overline{x}) \leq F(\hat{x})$ ) and  $x \to F(x) - md(\overline{x}, x)$  attains a strict maximum (resp. minimum) at  $\overline{x}$  with  $m := \sup F - F(\hat{x})$  (resp.  $m := \inf F - F(\hat{x})$ ).

See [5] for the proof.

Proof of Lemma 3.1. Fix  $\psi = \psi_1 + \psi_2$  with  $\psi_1 \in \underline{\mathcal{C}}^1(Q)$  and  $\psi_2 \in \mathcal{C}(Q)$  such that  $\overline{u} - \psi$  attains a zero maximum at  $\hat{z} = (\hat{t}, \hat{x})$  over  $\overline{B}_R(\hat{z}) \subset Q$  with some R > 0, i.e.

(3.2) 
$$(\overline{u} - \psi)(\hat{z}) = \sup_{\overline{B}_R(\hat{z})} (\overline{u} - \psi) = 0$$

Set  $\tilde{\psi}_2(z) := \psi_2(z) + d(\hat{x}, x)^2 + (t - \hat{t})^2$  with z = (t, x) and  $\tilde{\psi} = \psi_1 + \tilde{\psi}_2$ .

Take a subsequence  $a_j \to a$  and a sequence of points  $z_j = (t_j, x_j) \in \overline{B}_R(\hat{z})$  such that  $z_j \to \hat{z}$  and  $u_j(z_j) = u(z_j; a_j) \to \overline{u}(\hat{z}) = u^*(\hat{z}; a)$ , where  $u_j := u(\cdot; a_j)$ . We see

by Ekeland's variational principle (Lemma 3.4) that there exists  $w_j = (s_j, y_j) \in \overline{B}_R(\hat{z})$  such that  $z = (t, x) \mapsto ((u_j)^* - \tilde{\psi})(z) - m_j d(y_j, x)$  attains maximum at  $w_j$  over  $\overline{B}_R(\hat{z})$  with

$$m_j := \sup_{\overline{B}_R(\hat{z})} ((u_j)^* - \tilde{\psi}) - ((u_j)^* - \tilde{\psi})(z_j) \ge 0$$

Note that

$$\limsup_{j \to \infty} m_j = \limsup_{j \to \infty} \sup_{\overline{B}_R(\hat{z})} ((u_j)^* - \tilde{\psi}) - (\overline{u} - \tilde{\psi})(\hat{z})$$

and so  $m_i \to 0$  by the assumptions (3.1) and (3.2). We also observe that

$$((u_j)^* - \tilde{\psi})(w_j) \ge (u_j - \tilde{\psi})(z_j) - m_j d(y_j, x_j) \ge (u_j - \tilde{\psi})(z_j) - 2Rm_j$$

and that the last term converges to  $(\overline{u} - \tilde{\psi})(\hat{z})$  as  $j \to \infty$ . Therefore,

$$\limsup_{j \to \infty} d(\hat{x}, y_j)^2 + (s_j - \hat{t})^2 \leq \limsup_{j \to \infty} ((u_j)^* - \psi)(w_j) - (\overline{u} - \tilde{\psi})(\hat{z})$$
$$\leq \limsup_{j \to \infty} \sup_{\overline{B}_R(\hat{z})} ((u_j)^* - \psi) - (\overline{u} - \tilde{\psi})(\hat{z})$$

and it follows from (3.1) and (3.2) that  $w_j = (s_j, y_j) \rightarrow \hat{z} = (\hat{t}, \hat{x})$ .

Now, since  $u_j$  is a metric viscosity subsolution,

$$F_{-r_j}(w_j, |\nabla \psi_1|(w_j), \partial_t \hat{\psi}(w_j); a_j) \le 0.$$

Here,  $r_j$  is some non-negative number such that

$$r_j \le |\nabla \psi_2|^*(w_j) + m_j + 2d(\hat{x}, y_j) + 2|s_j - \hat{t}|$$

and so  $\limsup r_j \leq |\nabla \psi_2|^*(\hat{z})$ . Since  $(z, p, q, r; a) \to F_r(z, p, q; a)$  is continuous and  $r \to F_r(z, p, q; a)$  is non-decreasing, we see that

$$F_{-|\nabla\psi_2|^*(\hat{z})}(\hat{z}, |\nabla\psi_1|(\hat{z}), \partial_t\psi(\hat{z}); a) \le 0$$

Therefore, u is a subsolution.

Another goal of this section is a principle to construct a metric viscosity solution by the Perron method.

**Proposition 3.5** (Perron method). Let  $F = F(z, p, q) \in C(Q \times \mathbb{R} \times \mathbb{R})$  and let g be an  $\mathbb{R}$ -valued function on  $\partial Q$ . Let S denote the set of all metric viscosity subsolutions (resp. supersolution) v of (2.1) satisfying  $v^* \leq g$  (resp.  $v_* \leq g$ ) on  $\partial Q$ . Then,  $u(z) := \sup_{v \in S} v(z)$  (resp.  $u(z) := \inf_{v \in S} v(z)$ ) is a metric viscosity solution of (2.1).

Perron method for construction of a viscosity solution to Hamilton-Jacobi equations was first presented by H. Ishii [13]. Actually, the authors of [8] have already established a similar result for metric viscosity solutions ([8, Theorem 7.6]). However, let us give a proof since we have slightly improved the result to apply it directly to construction of a solution of the limit equation (1.3). We remark that the function

$$\overline{u}(x) := \begin{cases} +\infty & \text{if } x \in Q \\ g(x) & \text{if } x \in \partial Q \end{cases}$$

is a supersolution of (2.1).

*Proof.* We only show that u is a supersolution since being a subsolution is due to Corollary 3.3. Fix  $\psi = \psi_1 + \psi_2$  with  $\psi_1 \in \overline{\mathcal{C}}^1(Q)$  and  $\psi_2 \in \mathcal{C}(Q)$  such that  $u_* - \psi$ attains a zero minimum at  $\hat{z} := (\hat{t}, \hat{x})$  over  $\overline{B}_R(\hat{z}) \subset Q$  with some R > 0. Set  $\tilde{\psi}_2(z) := \psi_2(z) - d(\hat{z}, z)^2$  and  $\tilde{\psi} = \psi_1 + \tilde{\psi}_2$ . Suppose by contradiction that

$$F_{|\nabla\psi_2|^*(\hat{z})}(\hat{z}, |\nabla\psi_1|(\hat{z}), \partial_t\psi(\hat{z})) < 0.$$

Since  $(z, p, q, r) \mapsto F_r(z, p, q)$  is continuous and  $r \mapsto F_r(z, p, q)$  is non-decreasing, we may assume that  $\tilde{\psi} = \psi_1 + \tilde{\psi}_2$  is a subsolution of

$$F_{|\nabla \tilde{\psi}_2|^*(z)}(z, |\nabla \psi_1|(z), \partial_t \tilde{\psi}(z)) \le 0 \quad \text{for all } z \in B_R(\hat{z})$$

by taking R small enough. Recalling Proposition 2.10, we see that  $\tilde{\psi}$  is a metric viscosity subsolution of (2.1) in  $B_R(\hat{z})$ . Now observe that

$$(u - \tilde{\psi})(z) \ge (u_* - \tilde{\psi})(z) \ge d(\hat{z}, z)^2 \ge \frac{R^2}{4} =: m > 0$$

for all  $z \in \overline{B}_R(\hat{z}) \setminus B_{R/2}(\hat{z})$ . Construct a new function

$$w(z) = \begin{cases} \max\{\tilde{\psi}(z) + m/2, u(z)\} & \text{if } z \in B_R(\hat{z}), \\ u(z) & \text{otherwise.} \end{cases}$$

Then, w is equal to u on  $Q \setminus B_{R/2}(\hat{z})$  and so it is a subsolution of (2.1) in  $Q \setminus B_{R/2}(\hat{z})$ . It follows from Proposition 2.9 and Corollary 3.3 that w is a subsolution of (2.1) in  $B_R(\hat{z})$ . Therefore, Proposition 2.7 shows that w is a subsolution of (2.1) in Q and so  $w \in \mathcal{S}$ . In particular,  $u(\hat{z}) \ge w(\hat{z})$  but  $w(\hat{z}) = \psi(\hat{z}) + m/2 = u(\hat{z}) + m/2$ . Since m > 0, we obtain a contradiction and conclude that u is a supersolution.  $\Box$ 

## 4. Application to large time behavior

We study large time asymptotic behaviors of solutions of the Hamilton-Jacobi equation (1.2) with a Hamiltonian H satisfying (A1)–(A4) on a compact geodesic metric space (X, d). First note uniqueness of the constant c such that (1.3) admits a solution.

**Proposition 4.1.** Assume (A1), (A2) and that X is compact. Let  $c \in \mathbf{R}$  be a constant such that (1.3) admits a real-valued continuous solution. Then,

(4.1) 
$$c = \max_{x \in Y} H(x, 0).$$

*Proof.* By the assumption (A2) it is enough to show that

(4.2) 
$$\sup_{x \in X} \inf_{p \in \mathbf{R}_+} H(x, p) \le c \le \sup_{x \in X} H(x, 0).$$

It is easy to show the second inequality of (4.2). Indeed, since a solution v attains a minimum at some point  $\hat{x} \in X$ , we have  $H_0(\hat{x}, 0) = H(\hat{x}, 0) \ge c$ , which implies  $\sup_{x \in X} H(x, 0) \ge c$ . In order to prove the first inequality of (4.2), fix  $\hat{x} \in X$ . Let us consider the function  $v(x) - nd(x, \hat{x})^2/2$  and take its maximum point  $x_n$  for each  $n = 0, \cdots$ . Now, since  $v(x_n) - nd(x_n, \hat{x})^2/2 \ge v(\hat{x})$ , we have  $d(x_n, \hat{x})^2 \le 2(\max v - v(\hat{x}))/n$ . Therefore,  $x_n \to \hat{x}$  as  $n \to \infty$ . Since u is a subsolution,  $H_0(x_n, nd(x_n, \hat{x})) = H(x_n, nd(x_n, \hat{x})) \le c$ . Hence,  $\inf_{p \in \mathbf{R}_+} H(x_n, p) \le c$  and sending  $n \to \infty$  yields  $\inf_{p \in \mathbf{R}_+} H(\hat{x}, p) \le c$ . We now obtained the inequalities (4.2).

Remark 4.2. It is a problem whether the inequalities (4.2) holds even if we remove the compactness assumption. One can show them by a similar argument to the proofs in Section 3 using Ekeland's variational principle provided  $p \mapsto \sup_{x \in X} H(x, p)$  is continuous.

In view of this proposition we hereafter define c by (4.1). Now we are able to state the main theorem of large time behavior.

**Theorem 4.3** (Large time behavior). Assume (A1)-(A4),  $u_0 \in Lip(X)$  and that X is compact. Let u be a Lipschitz continuous solution of (1.2) with (??) on  $[0,\infty) \times X$ . Then, u(t,x) + ct converges to a function v locally uniformly as  $t \to \infty$  in X and v is a solution of (1.3).

In order to prove this theorem we first establish regularity, existence and comparison results for the stationary equation (1.3). Set

$$A := \{ x \in X \mid H(x, 0) = c \}$$

**Proposition 4.4** (Lipschitz continuity of solutions of (1.3)). Assume (A1), (A3) and (A4). Then, real-valued continuous solutions of (1.3) are equi-Lipschitz continuous.

*Proof.* Note that there exists a constant  $L \in \mathbf{R}_+$  such that  $H(x,p) \geq c$  for all  $x \in X$  and  $p \geq L$  by (A3). Fix a real-valued continuous solution v. Consider the function  $v(x) - v(y) - 2L\sqrt{d(x,y)^2 + \varepsilon^2}$  for  $x, y \in X$  and take its maximum point  $x_{\varepsilon}$  with respect to x for each  $\varepsilon > 0$ . Note that  $x \mapsto v(y) + 2L\sqrt{d(x,y)^2 + \varepsilon^2}$  is of  $\underline{\mathcal{C}}^1(X)$  by Lemma 2.2. Hence, we see that

$$H_0\left(x_{\varepsilon}, \frac{2Ld(x_{\varepsilon}, y)}{\sqrt{d(x_{\varepsilon}, y)^2 + \varepsilon^2}}\right) = H\left(x_{\varepsilon}, \frac{2Ld(x_{\varepsilon}, y)}{\sqrt{d(x_{\varepsilon}, y)^2 + \varepsilon^2}}\right) \le c$$

Therefore, we see that  $2Ld(x_{\varepsilon}, y)/\sqrt{d(x_{\varepsilon}, y)^2 + \varepsilon^2} \leq L$  and so  $x_{\varepsilon} \to y$  as  $\varepsilon \to 0$ . Now, for each  $x, y \in X$ , we have

$$v(x) - v(y) - 2L\sqrt{d(x,y)^2 + \varepsilon^2} \le v(x_\varepsilon) - v(y) - 2L\sqrt{d(x_\varepsilon,y)^2 + \varepsilon^2}$$

Sending  $\varepsilon \to 0$  yields

$$v(x) - v(y) - 2Ld(x, y) \le 0,$$

which means that all subsolutions of (1.3) is 2L-Lipschitz continuous.

**Theorem 4.5** (Existence of a solution of (1.3)). Assume (A1), (A3) and (A4). Then, there exists at least one Lipschitz continuous solution of (1.3) whenever A is non-empty.

Proof. Define

 $S(x,y) := \sup\{w(x) \mid w \in C(X) \text{ is a subsolution of } (1.3) \text{ with } w(y) = 0\}.$ 

Note that the constant  $w \equiv 0$  is a subsolution of (1.3). Also Proposition 4.4 ensures that the solutions of (1.3) are equi-Lipschitz continuous and hence  $v := S(\cdot, y)$  is a Lipschitz continuous function on X. Now, Corollary 3.3 implies that v is a subsolution of (1.3) in X while Proposition 3.5 shows that v is a supersolution of (1.3) in  $X \setminus \{y\}$ . Since  $H(x, p) \ge H(x, 0) = c$  for  $x \in A$ , we see that  $v = S(\cdot, y)$  is a solution for every  $y \in A \neq \emptyset$ . **Theorem 4.6** (Comparison principle for (2.2)). Let U be an open subset of X such that  $\overline{U}$  is compact. Assume (A1), (A2) and that H(x,0) < 0 for all  $x \in U$ . Let u be a subsolution and v be a supersolution of (2.2) such that  $u^* < +\infty$  and  $v_* > -\infty$ . If  $u^* \leq v_*$  on  $\partial U$ , then  $u^* \leq v_*$  in U.

*Proof.* First note that we may assume  $u^*(x_0) \neq -\infty$  and  $v_*(x_0) \neq +\infty$  at some  $x_0 \in U$ ; otherwise the conclusion holds. Fix  $\theta \in (0, 1)$  and consider the upper semicontinuous function defined by

$$\Phi(x,y) := \theta u^*(x) - v_*(y) - \frac{1}{2\varepsilon} d(x,y)^2$$

for  $\varepsilon > 0$ . Thanks to the compactness of  $\overline{U}$ , we are able to take a maximum point  $(x_{\varepsilon}, y_{\varepsilon}) \in \overline{U} \times \overline{U}$  of  $\Phi$ . It follows from  $\Phi(x_{\varepsilon}, y_{\varepsilon}) \ge \Phi(x_0, x_0)$  that

$$\frac{1}{2\varepsilon}d(x_{\varepsilon}, y_{\varepsilon})^{2} \leq \theta u^{*}(x_{\varepsilon}) - v_{*}(y_{\varepsilon}) - \theta u^{*}(x_{0}) + v_{*}(x_{0})$$
$$\leq \theta \sup u^{*} - \inf u_{*} - \theta u^{*}(x_{0}) + v_{*}(x_{0}) < +\infty$$

Hence,  $d(x_{\varepsilon}, y_{\varepsilon}) \to 0$  and so we may assume that  $x_{\varepsilon}$  and  $y_{\varepsilon}$  converge to a same point  $\bar{x} \in \overline{U}$  by taking a subsequence. Let us consider the case when  $\bar{x} \in U$ . Then, since u and v are a subsolution and a supersolution,

$$H(x_{\varepsilon}, \frac{1}{\theta_{\varepsilon}}d(x_{\varepsilon}, y_{\varepsilon})) \le 0,$$
  
$$H(y_{\varepsilon}, \frac{1}{\varepsilon}d(x_{\varepsilon}, y_{\varepsilon})) \ge 0.$$

By the convexity of H the second inequality yields

$$(1-\theta)H(y_{\varepsilon},0)+\theta H(y_{\varepsilon},\frac{1}{\theta\varepsilon}d(x_{\varepsilon},y_{\varepsilon}))\geq 0$$

Hence,

$$(1-\theta)H(y_{\varepsilon},0)+\theta H(y_{\varepsilon},\frac{1}{\theta\varepsilon}d(x_{\varepsilon},y_{\varepsilon}))-\theta H(x_{\varepsilon},\frac{1}{\theta\varepsilon}d(x_{\varepsilon},y_{\varepsilon}))\geq 0.$$

Sending  $\varepsilon \to 0$  yields  $(1 - \theta)H(\bar{x}, 0) \ge 0$ . Since  $H(\bar{x}, 0) < 0$  and  $\theta < 1$ , we obtain a contradiction. Therefore,  $\bar{x} \in \partial U$ . We now observe that

$$\theta u^*(x_{\varepsilon}) - v_*(y_{\varepsilon}) \ge \Phi(x_{\varepsilon}, y_{\varepsilon}) \ge \sup_{x \in U} \Phi(x, x) = \sup_{U} (\theta u^* - v_*).$$

Hence, we see that  $\sup_U(\theta u^* - v_*) \leq (\theta u^* - v_*)(\bar{x}) \leq \sup_{\partial U}(\theta u^* - v_*)$ . Sending  $k \to 1$  implies  $\sup_U(u^* - v_*) \leq \sup_{\partial U}(u^* - v_*)$ .

**Corollary 4.7** (Comparison principle for (1.3)). Assume that X is compact. Let u be a subsolution and v be a supersolution of (1.3) such that  $u^* < +\infty$  and  $v_* > -\infty$ . If  $u^* \le v_*$  on A, then  $u^* \le v_*$  on X.

*Proof.* It follows from the definition of A that H(x, 0) - c < 0 for all  $x \in U := X \setminus A$ . Therefore, Theorem 4.6 implies  $u^* \leq v_*$  in  $X \setminus A$ .

We will also require a comparison principle for the evolution equation (1.2).

**Theorem 4.8** (Comparison principle for (1.2)). Assume (A1) and that X is compact. Let u be a subsolution and v be a supersolution of (1.2) such that  $u^* < +\infty$ and  $v_* > -\infty$ . If  $u^*|_{t=0} \le v_*|_{t=0}$ , then  $u^* \le v_*$  on  $(0,\infty) \times X$ . One is able to prove this theorem with the same idea as in [9, Proof of Proposition 3.3] and so we omit the proof.

Before starting the proof of Theorem 4.3, let us explain that the initial value problem (1.2), (??) admits a unique Lipschitz continuous solution. We will construct a solution by Perron method while the uniqueness is a direct consequence of the comparison principle (Theorem 4.8). Let  $\text{Lip}[u_0]$  denote the Lipschitz constant of  $u_0$  and set  $K = \max_{x \in X} |H(x, \text{Lip}[u_0])|$ . First note that  $\overline{u}(t, x) := u_0(x) + Kt$  and  $\underline{u}(t, x) := u_0(x) - Kt$  are a Lipschitz continuous supersolution and subsolution on  $[0, \infty) \times X$ , respectively. We then can construct a continuous solution u such that  $\underline{u} \leq \overline{u}$  by using Proposition 3.5 and Theorem 4.8. Take a constant  $L \in \mathbf{R}_+$  such that  $H(x, p) \geq c$  for all  $x \in X$  and  $p \geq L$ . Then, we see that  $|u(t, x) - u(s, y)| \leq K|t-s| + Ld(x, y)$  by a similar argument to the proof of Proposition:liplhj. Actually, this is a standard argument and we refer the reader to [10].

We are now able to prove the main theorem stated at the top of this section.

Proof of Theorem 4.3. Take the solution  $v_0$  of (1.3) in Theorem 4.5. Noting that  $u_0$  and  $v_0$  are bounded since X is compact, we are also able to see that  $v_0 - M \leq u_0 \leq v_0 + M$  for some large M > 0. Recall Propositions 2.5 and 2.9, which imply that  $v_0 - ct \pm M$  are solutions of (1.2). We then see by a comparison principle for (1.2) (Theorem 4.8) that  $v_0 - ct - M \leq u \leq v_0 - ct + M$ . Thus, the upper and lower semi-limits

$$\overline{v}(x) := \sup_{\substack{(t_j, x_j) \to (\infty, x) \\ (t_j, x_j) \to (\infty, x) }} \limsup_{j} \{u(t_j, x_j) + ct_j\},$$
$$\underline{v}(x) := \inf_{\substack{(t_j, x_j) \to (\infty, x) \\ j}} \liminf_{j} \{u(t_j, x_j) + ct_j\}$$

can be defined as a bounded function on X since  $v_0 - M \le \underline{v} \le \overline{v} \le v_0 + M$ . We next note that Propositions 2.8 and 2.9 show the function

$$w^\lambda(t,x):=u\left(\frac{t}{\lambda},x\right)+c\frac{t}{\lambda}$$

is a solution of

(4.3) 
$$\lambda \partial_t w^{\lambda} + H(x, |Dw^{\lambda}|) = c \quad \text{in } (0, \infty) \times X$$

for each  $\lambda > 0$ . Since

$$\overline{v}(x) = \sup_{\substack{(t_j, x_j, \lambda_j) \to (t, x, 0) \\ (t_j, x_j, \lambda_j) \to (t, x, 0)}} \limsup_j w^{\lambda_j}(t_j, x_j),$$
$$\underline{v}(x) = \inf_{\substack{(t_j, x_j, \lambda_j) \to (t, x, 0) \\ j}} \liminf_j w^{\lambda_j}(t_j, x_j)$$

for all t > 0 and  $x \in X$ , i.e.  $\overline{v}$  and  $\underline{v}$  are respectively nothing but the upper and lower semilimit of  $w^{\lambda}$  as  $\lambda \to 0$ , the stability result (Proposition 3.1) and Proposition 2.5 shows that  $\overline{v}$  and  $\underline{v}$  are a subsolution and a supersolution of (1.3), respectively.

We next claim that  $\overline{v} = \underline{v}$  on the set A. Indeed, for each  $x \in A$ , u(t, x) + ct converges to some v(x) since  $\partial_t u + c \leq 0$  and so it is a decreasing sequence. We also obtain that u is equi-Lipschitz continuous. By connecting these two facts, we see that  $\overline{v} \leq v \leq \underline{v}$  on A.

Finally, the comparison principle (Theorem 4.7) shows that  $\overline{v} \leq \underline{v}$  on the whole space X. Thus, we can conclude that u(t, x) + ct converges to some function  $v = \overline{v} = \underline{v}$  which is a solution of (1.3).

10

Remark 4.9. The convexity assumption (A2) is used only to guarantee a comparison principle holds for the stationary equation (1.3). It is possible to weaken the condition. For instance, let us consider the specific Hamiltonian  $H(x, p) = \sqrt{|p|}$ , which is not convex. One easily see that the equation (1.3) is equivalent to  $|Dv| = c^2$ . Since a comparison principle for the convex Hamiltonian  $|Dv| = c^2$  implies a comparison principle  $\sqrt{|Dv|} = c$ , the same behavior of the solution must occur to the Hamiltonian  $H(x, p) = \sqrt{|p|}$ . This scheme works for quasiconvex Hamiltonians H(x, p) = h(|p|) + f(x) with  $h: \mathbf{R}_+ \to \mathbf{R}$  such that  $h(p) - \lambda p$  is non-decreasing for some  $\lambda > 0$ .

Let us introduce the functions  $\phi_{-}, \phi_{\infty} \in C(X)$  by

$$\begin{split} \phi_{-}(x) &:= \inf_{t \geq 0} (u(x,t) + ct), \\ \phi_{\infty}(x) &:= \min\{S(x,y) + \phi_{-}(y) \mid y \in A\}, \end{split}$$

where the function S is defined in the proof of Theorem 4.5.

**Proposition 4.10.** Let  $u \in C([0,\infty) \times X)$  be a solution of (1.2). Then  $\phi_{-}$  and  $\phi_{\infty}$  are solutions of (1.3).

*Proof.* Note that u(x,t) + ct is supersolution of (1.3). Thus, by Corollary 3.3 and Proposition 3.5, which can be proved similarly for (1.3) as well, we immediately see that  $\phi_{-}$  is solution of (1.3). Since S is solution of (1.3) as we showed in the proof of Theorem 4.5, applying Corollary 3.3 and Proposition 3.5 again implies that  $\phi_{\infty}$  is solution of (1.3).

Theorem 4.11 (Asymptotic profile). We have

(4.4) 
$$\lim_{t \to \infty} (u(x,t) + ct) = \phi_{\infty}(x) \quad \text{for all } x \in X.$$

*Proof.* Denote the left-hand side of (4.4) by  $u_{\infty}(x)$ . Since

 $\phi_-(x) \leq u(x,t) + ct \quad \text{for all } (x,t) \in X \times [0,\infty),$ 

it follows that  $\phi_{-} \leq u_{\infty}$  in X. It also can be seen that  $\phi_{\infty} \leq \phi_{-}$  on  $\mathcal{A}$  and thus we have  $\phi_{\infty} \leq u_{\infty}$  on  $\mathcal{A}$  from above two inequalities. This fact and Theorem 4.7 leads us to the relationship that  $\phi_{\infty} \leq u_{\infty}$  on X.

We shall show the other inequality. In order to do, set  $v(x,t) := \inf_{s \ge t} (u(x,s) + cs)$  and then we can easily see  $v(x,0) = \phi_{-}(x)$ . Since  $\phi_{\infty} = \phi_{-}$  on A, Corollary 4.7 implies that  $\phi_{\infty} \ge \phi_{-} = v(\cdot,0)$  on X. It can be considered that  $\phi_{\infty}$  as a solution of (1.2) with  $\phi_{\infty} \ge u_{0}$  and thus by Theorem 4.8 we obtain  $v(x,t) \le \phi_{\infty}(x)$  for all  $x \in X$  and  $t \ge 0$ . Sending  $t \to \infty$  yields  $u_{\infty} \le \phi_{\infty}$  on X.

## References

- L. Ambrosio and J. Feng, On a class of first order Hamilton-Jacobi equations in metric spaces, J. Differential Equations 256 (2014), no. 7, 2194–2245.
- [2] G. Barles and B. Perthame, Discontinuous solutions of deterministic optimal stopping time problems, RAIRO Modél. Math. Anal. Numér. 21 (1987), no. 4, 557–579.
- [3] F. Camilli, R. Capitanelli and C. Marchi, Eikonal equations on the Sierpinski gasket, preprint.
- [4] I. Ekeland, On the variational principle, J. Math. Anal. Appl. 47 (1974), 324–353.
- [5] I. Ekeland, Nonconvex minimization problems, Bull. Amer. Math. Soc. (N.S.) 1 (1979), no. 3, 443–474.
- [6] A. Fathi, Sur la convergence du semi-groupe de Lax-Oleinik, C. R. Acad. Sci. Paris Sr. I Math. 327 (1998), no. 3, 267–270.

- [7] A. Fathi and A. Siconolfi, PDE aspects of Aubry-Mather theory for quasiconvex Hamiltonians, Calc. Var. Partial Differential Equations 22 (2005), no. 2, 185-228.
- [8] W. Gangbo and A. Święch, Optimal transport and large number of particles, Discrete Contin. Dyn. Syst. 34 (2014), no. 4, 1397–1441.
- [9] W. Gangbo and A. Święch, Metric viscosity solutions of Hamilton-Jacobi equations, preprint.
- [10] Y. Giga, N. Hamamuki, Hamilton-Jacobi equations with discontinuous source terms, Comm. Partial Differential Equations, 38 (2013), no. 2, 199-243.
- [11] Y. Giga, N. Hamamuki and A. Nakayasu, Eikonal equations in metric spaces, Trans. Amer. Math. Soc. to appear.
- [12] Y. Giga, Q. Liu and H. Mitake, Singular Neumann problem and large-time behaviour of solutions of noncoercive Hamilton-Jacobi equations Trans. Amer. Math. Soc. 366 (2013), 1905–1941.
- [13] H. Ishii, Perrons method for Hamilton-Jacobi equations Duke Math. J. 55 (1987), no. 2, 369-384.
- [14] P.-L. Lions, G. C. Papanicolaou and S. R. S. Varadhan, Homogenization of Hamilton-Jacobi equations, unpublished.
- [15] A. Nakayasu, On metric viscosity solutions for Hamilton-Jacobi equations of evolution type, preprint.
- [16] G., Namah and J.-M. Roquejoffre, Remarks on the long time behaviour of the solutions of Hamilton-Jacobi equations, Comm. Partial Differential Equations 24 (1999), no. 5–6, 883-893.
- [17] D., Schieborn and F. Camilli, Viscosity solutions of Eikonal equations on topological networks, preprint.

Graduate School of Mathematical Sciences, The University of Tokyo, 3-8-1 Komaba, Meguro, Tokyo, 153-8914 Japan

*E-mail address*: ankys@ms.u-tokyo.ac.jp

Graduate School of Mathematical Sciences, The University of Tokyo, 3-8-1 Komaba, Meguro, Tokyo, 153-8914 Japan

*E-mail address*: namba@ms.u-tokyo.ac.jp