

J. Number Theory 183 (2018), 146–171.

## ARITHMETIC PROPERTIES OF DELANNOY NUMBERS AND SCHRÖDER NUMBERS

ZHI-WEI SUN

Department of Mathematics, Nanjing University  
Nanjing 210093, People’s Republic of China  
zwsun@nju.edu.cn  
<http://maths.nju.edu.cn/~zwsun>

ABSTRACT. Define

$$D_n(x) = \sum_{k=0}^n \binom{n}{k}^2 x^k (x+1)^{n-k} \quad \text{for } n = 0, 1, 2, \dots$$

and

$$s_n(x) = \sum_{k=1}^n \frac{1}{n} \binom{n}{k} \binom{n}{k-1} x^{k-1} (x+1)^{n-k} \quad \text{for } n = 1, 2, 3, \dots$$

Then  $D_n(1)$  is the  $n$ -th central Delannoy number  $D_n$ , and  $s_n(1)$  is the  $n$ -th little Schröder number  $s_n$ . In this paper we obtain some surprising arithmetic properties of  $D_n(x)$  and  $s_n(x)$ . We show that

$$\frac{1}{n} \sum_{k=0}^{n-1} D_k(x) s_{k+1}(x) \in \mathbb{Z}[x(x+1)] \quad \text{for all } n = 1, 2, 3, \dots$$

Moreover, for any odd prime  $p$  and  $p$ -adic integer  $x \not\equiv 0, -1 \pmod{p}$ , we establish the supercongruence

$$\sum_{k=0}^{p-1} D_k(x) s_{k+1}(x) \equiv 0 \pmod{p^2}.$$

As an application we confirm Conjecture 5.5 in [S14a], in particular we prove that

$$\frac{1}{n} \sum_{k=0}^{n-1} T_k M_k (-3)^{n-1-k} \in \mathbb{Z} \quad \text{for all } n = 1, 2, 3, \dots,$$

where  $T_k$  is the  $k$ -th central trinomial coefficient and  $M_k$  is the  $k$ -th Motzkin number.

---

2010 *Mathematics Subject Classification*. Primary 11A07, 11B75; Secondary 05A10, 05A19, 11B65, 11C08, 33F99.

*Keywords*. Congruence, Polynomial, Delannoy number, Schröder number, Motzkin number  
Supported by the National Natural Science Foundation of China (grant 11571162).

## 1. INTRODUCTION

For  $m, n \in \mathbb{N} = \{0, 1, 2, \dots\}$ , the Delannoy number

$$D_{m,n} := \sum_{k \in \mathbb{N}} \binom{m}{k} \binom{n}{k} 2^k \quad (1.1)$$

in combinatorics counts lattice paths from  $(0, 0)$  to  $(m, n)$  in which only east  $(1, 0)$ , north  $(0, 1)$ , and northeast  $(1, 1)$  steps are allowed (cf. R. P. Stanley [St99, p. 185]). The  $n$ -th central Delannoy number  $D_n = D_{n,n}$  has another well-known expression:

$$D_n = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} = \sum_{k=0}^n \binom{n+k}{2k} \binom{2k}{k}. \quad (1.2)$$

For  $n \in \mathbb{Z}^+ = \{1, 2, 3, \dots\}$ , the  $n$ -th little Schröder number is given by

$$s_n := \sum_{k=1}^n N(n, k) 2^{k-1} \quad (1.3)$$

with the Narayana number  $N(n, k)$  defined by

$$N(n, k) := \frac{1}{n} \binom{n}{k} \binom{n}{k-1} \in \mathbb{Z}.$$

(See [Gr, pp. 268–281] for certain combinatorial interpretations of the Narayana number  $N(n, k) = N(n, n+1-k)$ .) For  $n \in \mathbb{N}$ , the  $n$ -th large Schröder number is given by

$$S_n := \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} \frac{1}{k+1} = \sum_{k=0}^n \binom{n+k}{2k} C_k, \quad (1.4)$$

where  $C_k$  denotes the Catalan number  $\binom{2k}{k}/(k+1) = \binom{2k}{k} - \binom{2k}{k+1}$ . It is well known that  $S_n = 2s_n$  for every  $n = 1, 2, 3, \dots$ . Both little Schröder numbers and large Schröder numbers have many combinatorial interpretations (cf. [St97] and [St99, pp. 178, 239–240]); for example,  $s_n$  is the number of ways to insert parentheses into an expression of  $n+1$  terms with two or more items within a parenthesis, and  $S_n$  is the number of lattice paths from the point  $(0, 0)$  to  $(n, n)$  with steps  $(1, 0)$ ,  $(0, 1)$  and  $(1, 1)$  which never rise above the line  $y = x$ .

Surprisingly, the central Delannoy numbers and Schröder numbers arising naturally in enumerative combinatorics, have nice arithmetic properties. In 2011 the author [S11b] showed that

$$\sum_{k=1}^{p-1} \frac{D_k}{k^2} \equiv (-1)^{(p-1)/2} 2E_{p-3} \pmod{p} \quad \text{and} \quad \sum_{k=1}^{p-1} \frac{S_k}{6^k} \equiv 0 \pmod{p}$$

for any prime  $p > 3$ , where  $E_0, E_1, E_2, \dots$  are the Euler numbers. In 2014 the author [S14a] proved that

$$\frac{1}{n^2} \sum_{k=0}^{n-1} (2k+1)D_k^2 \in \mathbb{Z} \quad \text{for all } n = 1, 2, 3, \dots,$$

and that

$$\sum_{k=0}^{p-1} D_k^2 \equiv \left(\frac{2}{p}\right) \pmod{p} \quad \text{for any odd prime } p,$$

where  $\left(\frac{\cdot}{p}\right)$  denotes the Legendre symbol.

*Definition 1.1.* We define

$$D_n(x) := \sum_{k=0}^n \binom{n}{k}^2 x^k (x+1)^{n-k} \quad \text{for } n \in \mathbb{N}, \quad (1.5)$$

and

$$s_n(x) := \sum_{k=1}^n N(n, k) x^{k-1} (x+1)^{n-k} \quad \text{for } n \in \mathbb{Z}^+. \quad (1.6)$$

Obviously  $D_n(1) = D_n$  for  $n \in \mathbb{N}$ , and  $s_n(1) = s_n$  for  $n \in \mathbb{Z}^+$ .

In this paper we obtain somewhat curious results involving the polynomials  $D_n(x)$  and  $s_n(x)$ . Our first theorem is as follows.

**Theorem 1.1.** (i) *For any  $n \in \mathbb{Z}^+$ , we have*

$$\frac{1}{n} \sum_{k=0}^{n-1} D_k(x) s_{k+1}(x) = W_n(x(x+1)), \quad (1.7)$$

where

$$W_n(x) = \sum_{k=1}^n w(n, k) C_{k-1} x^{k-1} \in \mathbb{Z}[x] \quad (1.8)$$

with

$$w(n, k) = \frac{1}{k} \binom{n-1}{k-1} \binom{n+k}{k-1} \in \mathbb{Z}. \quad (1.9)$$

(ii) *Let  $p$  be an odd prime. For any  $p$ -adic integer  $x$ , we have*

$$\begin{aligned} & \sum_{k=0}^{p-1} D_k(x) s_{k+1}(x) \\ \equiv & \begin{cases} p(1 - x(x+1)) \pmod{p^3} & \text{if } x \equiv 0, -1 \pmod{p}, \\ 2p^2 + \frac{2x+1}{x(x+1)} p^2 (x^2 q_p(x) - (x+1)^2 q_p(x+1)) \pmod{p^3} & \text{otherwise,} \end{cases} \end{aligned} \quad (1.10)$$

where  $q_p(z)$  denotes the Fermat quotient  $(z^{p-1} - 1)/p$  for any  $p$ -adic integer  $z \not\equiv 0 \pmod{p}$ .

*Remark 1.1.* It is interesting to compare the new numbers  $w(n, k)$  with the Narayana numbers  $N(n, k)$ .

Clearly, Theorem 1.1 in the case  $x = 1$  yields the following consequence.

**Corollary 1.1.** *For any positive integer  $n$ , we have*

$$\frac{1}{n} \sum_{k=0}^{n-1} D_k s_{k+1} = \sum_{k=1}^n w(n, k) C_{k-1} 2^{k-1} \equiv 1 \pmod{2}. \quad (1.11)$$

*Also, for any odd prime  $p$  we have*

$$\sum_{k=0}^{p-1} D_k s_{k+1} \equiv 2p^2(1 - 3q_p(2)) \pmod{p^3}. \quad (1.12)$$

*Remark 1.2.* For the prime  $p = 588811$ , we have  $q_p(2) \equiv 1/3 \pmod{p}$  and hence  $\sum_{k=0}^{p-1} D_k s_{k+1} \equiv 0 \pmod{p^3}$ . In 2016 J.-C. Liu [L] confirmed the author's conjecture (cf. [S11b, Conjecture 1.1]) that

$$\sum_{k=1}^{p-1} D_k S_k \equiv -2p \sum_{k=1}^{p-1} \frac{(-1)^k + 3}{k} \pmod{p^4} \quad \text{for any prime } p > 3.$$

From Theorem 1.1 we can deduce a novel combinatorial identity.

**Corollary 1.2.** *For any  $n \in \mathbb{Z}^+$ , we have*

$$\sum_{k=1}^n \binom{n}{k} \binom{n+k}{k-1} \frac{C_{k-1}}{(-4)^{k-1}} = \frac{\lfloor (n+1)/2 \rfloor}{4^{n-1}} \binom{n}{\lfloor n/2 \rfloor}^2. \quad (1.13)$$

*Remark 1.3.* If we let  $a_n$  denote the left-hand side of (1.13), then the Zeilberger algorithm (cf. [PWZ, pp. 101-119]) cannot find a closed form for  $a_n$ , and it only yields the following second-order recurrence relation:

$$(n+1)^2 a_n + (2n+3)a_{n+1} - (n+1)(n+3)a_{n+2} = 0 \quad \text{for } n = 1, 2, 3, \dots$$

Now we give our second theorem which can be viewed as a supplement to Theorem 1.1.

**Theorem 1.2.** *Let  $p$  be any odd prime. Then*

$$\sum_{k=0}^{p-1} k D_k(x) s_{k+1}(x) \equiv 2(x(x+1))^{(p-1)/2} \pmod{p}. \quad (1.14)$$

*In particular,*

$$\sum_{k=0}^{p-1} k D_k s_{k+1} \equiv 2 \left( \frac{2}{p} \right) \pmod{p}. \quad (1.15)$$

In the next section we are going to show Theorems 1.1-1.2 and Corollary 1.2. In Section 3 we will give applications of Theorems 1.1-1.2 to central trinomial coefficients, Motzkin numbers, and their generalizations. Section 4 contains two related conjectures.

Throughout this paper, for any polynomial  $P(x)$  and  $n \in \mathbb{N}$ , we use  $[x^n]P(x)$  to denote the coefficient of  $x^n$  in  $P(x)$ .

## 2. PROOFS OF THEOREMS 1.1-1.2 AND COROLLARY 1.2

Recall the following definition given in [S12a] motivated by the large Schröder numbers.

*Definition 2.1.* For  $n \in \mathbb{N}$  we set

$$S_n(x) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} \frac{x^k}{k+1} = \sum_{k=0}^n \binom{n+k}{2k} C_k x^k. \quad (2.1)$$

**Lemma 2.1.** *We have*

$$D_n(x) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k \quad \text{for } n \in \mathbb{N}. \quad (2.2)$$

Also, for any  $n \in \mathbb{Z}^+$  we have

$$D_{n+1}(x) - D_{n-1}(x) = 2x(2n+1)S_n(x) \quad (2.3)$$

and

$$(x+1)s_n(x) = S_n(x). \quad (2.4)$$

*Proof.* For  $k, n \in \mathbb{N}$ , we obviously have

$$\begin{aligned} [x^k]D_n(x) &= [x^k] \sum_{j=0}^n \binom{n}{j}^2 x^j (x+1)^{n-j} = \sum_{j=0}^k \binom{n}{j}^2 \binom{n-j}{k-j} \\ &= \binom{n}{k} \sum_{j=0}^k \binom{n}{j} \binom{k}{k-j} = \binom{n}{k} \binom{n+k}{k} \end{aligned}$$

with the help of the Chu-Vandermonde identity (cf. [G, (3.1)]). This proves (2.2).

Now fix  $n \in \mathbb{Z}^+$ . For  $k \in \mathbb{N}$ , by (2.2) we clearly have

$$\begin{aligned} & [x^{k+1}](D_{n+1}(x) - D_{n-1}(x)) \\ &= \binom{n+1+(k+1)}{2(k+1)} \binom{2(k+1)}{k+1} - \binom{n-1+(k+1)}{2(k+1)} \binom{2(k+1)}{k+1} \\ &= \frac{2n+1}{2k+1} \binom{n+k}{2k} \binom{2k+2}{k+1} = \frac{2(2n+1)}{k+1} \binom{n+k}{2k} \binom{2k}{k} \\ &= [x^{k+1}]2x(2n+1)S_n(x). \end{aligned}$$

So (2.3) follows.

For each  $k \in \mathbb{N}$ , it is apparent that

$$\begin{aligned}
[x^k](x+1)s_n(x) &= [x^k](x+1) \sum_{j=1}^n N(n, j) x^{j-1} (x+1)^{n-j} \\
&= \sum_{0 < j \leq k} N(n, j) \binom{n-j}{k-j} + \sum_{j=1}^{k+1} N(n, j) \binom{n-j}{k-j+1} \\
&= \frac{1}{n} \sum_{j=1}^{k+1} \binom{n}{j} \binom{n}{j-1} \binom{n-j+1}{k-j+1} \\
&= \frac{1}{n} \binom{n}{k} \sum_{j=0}^{k+1} \binom{n}{j} \binom{k}{k+1-j} \\
&= \frac{1}{n} \binom{n}{k} \binom{n+k}{k+1} = \binom{n}{k} \binom{n+k}{k} \frac{1}{k+1} = [x^k]S_n(x).
\end{aligned}$$

This proves (2.4).

The proof of Lemma 2.1 is now complete.  $\square$

*Remark 2.1.* Note that the Legendre polynomial of degree  $n$  is given by

$$P_n(x) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} \left(\frac{x-1}{2}\right)^k.$$

**Lemma 2.2.** *Let  $n \in \mathbb{Z}^+$ . Then*

$$n(n+1)S_n(x)^2 = \sum_{k=1}^n \binom{n+k}{2k} \binom{2k}{k} \binom{2k}{k+1} x^{k-1} (x+1)^{k+1} \quad (2.5)$$

and

$$\frac{D_{n-1}(x) + D_{n+1}(x)}{2} S_n(x) = \sum_{k=0}^n \binom{n+k}{2k} \binom{2k}{k}^2 \frac{2k+1}{(k+1)^2} x^k (x+1)^{k+1}. \quad (2.6)$$

*Remark 2.2.* The identities (2.5) and (2.6) are (2.1) and (3.6) of the author's paper [S12a] respectively.

**Lemma 2.3.** *For any  $m, n \in \mathbb{Z}^+$  with  $m \leq n$ , we have the identity*

$$\begin{aligned}
&\sum_{k=m}^n \binom{k+m}{2m} \left(2m+1 - m(m+1) \frac{2k+1}{k(k+1)}\right) \\
&= \frac{(n-m)(n+m+1)}{n+1} \binom{n+m}{2m}.
\end{aligned} \quad (2.7)$$

*Proof.* When  $n = m$ , both sides of (2.7) vanish.

Let  $m, n \in \mathbb{Z}^+$  with  $n \geq m$ . If (2.7) holds, then

$$\begin{aligned}
 & \sum_{k=m}^{n+1} \binom{k+m}{2m} \left( 2m+1 - m(m+1) \frac{2k+1}{k(k+1)} \right) \\
 &= \frac{(n-m)(n+m+1)}{n+1} \binom{n+m}{2m} \\
 & \quad + \binom{n+1+m}{2m} \left( 2m+1 - m(m+1) \frac{2(n+1)+1}{(n+1)(n+2)} \right) \\
 &= \binom{n+m+1}{2m} \left( \frac{(n-m)(n-m+1)}{n+1} + 2m+1 - \frac{m(m+1)(2n+3)}{(n+1)(n+2)} \right) \\
 &= \binom{(n+1)+m}{2m} \frac{(n+1-m)((n+1)+m+1)}{(n+1)+1}.
 \end{aligned}$$

In view of the above, we have proved Lemma 2.3 by induction.  $\square$

Let  $A$  and  $B$  be integers. The Lucas sequence  $u_n(A, B)$  ( $n = 0, 1, 2, \dots$ ) is defined by  $u_0(A, B) = 0$ ,  $u_1(A, B) = 1$ , and

$$u_{n+1}(A, B) = Au_n(A, B) - Bu_{n-1}(A, B) \quad \text{for } n = 1, 2, 3, \dots$$

It is well known that if  $\Delta = A^2 - 4B \neq 0$  then

$$u_n(A, B) = \frac{\alpha^n - \beta^n}{\alpha - \beta} \quad \text{for all } n \in \mathbb{N},$$

where  $\alpha$  and  $\beta$  are the two roots of the quadratic equation  $x^2 - Ax + B = 0$  (so that  $\alpha + \beta = A$  and  $\alpha\beta = B$ ). It is also known (see, e.g., [S10, Lemma 2.3]) that for any odd prime  $p$  we have

$$u_p(A, B) \equiv \left( \frac{\Delta}{p} \right) \pmod{p}, \quad \text{and } u_{p-\left(\frac{\Delta}{p}\right)}(A, B) \equiv 0 \pmod{p} \text{ if } p \nmid B.$$

In particular,  $F_p \equiv \left(\frac{5}{p}\right) \pmod{p}$  and  $p \mid F_{p-\left(\frac{5}{p}\right)}$  for any odd prime  $p$ , where  $F_n = u_n(1, -1)$  with  $n \in \mathbb{N}$  is the  $n$ -th Fibonacci number.

**Lemma 2.4.** *Let  $p$  be an odd prime, and let  $x$  be any integer not divisible by  $p$ . Then*

$$W_p(x) \equiv \frac{4x+1}{2x} \left( \left( \frac{4x+1}{p} \right) - 1 \right) \pmod{p}. \quad (2.8)$$

Moreover, if  $x \equiv -1/4 \pmod{p}$  then

$$W_p(x) \equiv 2p \pmod{p^2}, \quad (2.9)$$

otherwise we have

$$\begin{aligned}
W_p(x) &\equiv 2p + \frac{4x+1}{2x} \left( 1 - x^{p-1} + (p+1) \left( \left( \frac{4x+1}{p} \right) - 1 \right) \right) \\
&\quad - \frac{4x+1}{4x^{2-\left(\frac{4x+1}{p}\right)}} \left( 2x + \left( \frac{4x+1}{p} \right) \right) u_{p-\left(\frac{4x+1}{p}\right)}(2x+1, x^2) \pmod{p^2}.
\end{aligned} \tag{2.10}$$

*Proof.* Clearly,

$$\binom{2p-1}{p-1} = \prod_{k=1}^{p-1} \left( 1 + \frac{p}{k} \right) \equiv 1 + p \sum_{k=1}^{(p-1)/2} \left( \frac{1}{k} + \frac{1}{p-k} \right) \equiv 1 \pmod{p^2}.$$

(J. Wolstenholme [W] even showed that  $\binom{2p-1}{p-1} \equiv 1 \pmod{p^3}$  if  $p > 3$ .) Thus

$$w(p, p) = \frac{1}{p} \binom{2p}{p-1} = \frac{2}{p+1} \binom{2p-1}{p-1} \equiv 2(1-p) \pmod{p^2},$$

and

$$C_{p-1} = \frac{1}{p} \binom{2p-2}{p-1} = \frac{1}{2p-1} \binom{2p-1}{p-1} \equiv -(2p+1) \pmod{p^2}.$$

For each  $k = 1, \dots, p-1$ , clearly

$$\begin{aligned}
w(p, k) &= \frac{1}{k} \binom{p-1}{k-1} \binom{p+k-1}{k-1} \frac{p+k}{p+1} \\
&= \left( 1 + \frac{p}{k} \right) \frac{1}{p+1} \prod_{0 < j < k} \left( \frac{p-j}{j} \cdot \frac{p+j}{j} \right) \\
&\equiv \left( \frac{1}{p+1} + \frac{p}{k} \right) (-1)^{k-1} \equiv (-1)^{k-1} \left( 1 - p + \frac{p}{k} \right) \pmod{p^2}.
\end{aligned}$$



Therefore

$$\begin{aligned}
 W_p(x) &= w(p, p)C_{p-1}x^{p-1} + \sum_{k=1}^{p-1} w(p, k)C_{k-1}x^{k-1} \\
 &\equiv 2(1-p)C_{p-1}x^{p-1} + \sum_{k=1}^{p-1} \left(1-p + \frac{p}{k}\right) C_{k-1}(-x)^{k-1} \\
 &\equiv (2-2p)C_{p-1}x^{p-1} + (p+1) \left( \sum_{k=2}^p C_{k-1}(-x)^{k-1} + 1 - C_{p-1}(-x)^{p-1} \right) \\
 &\quad + p \sum_{k=1}^{p-1} \left( \frac{1}{k} - 2 \right) C_{k-1}(-x)^{k-1} \\
 &\equiv p+1 - (1-3p)(1+2p)x^{p-1} + (p+1) \sum_{k=1}^{p-1} C_k(-x)^k \\
 &\quad - \frac{p}{2} \sum_{k=1}^{p-1} \frac{\binom{2k}{k}}{k} (-x)^{k-1} \\
 &\equiv 2p+1 - x^{p-1} + (p+1) \sum_{k=1}^{p-1} \frac{C_k}{m^k} - \frac{p}{2} m \sum_{k=1}^{p-1} \frac{\binom{2k}{k}}{km^k} \pmod{p^2},
 \end{aligned}$$

where  $m$  is an integer with  $m \equiv -1/x \pmod{p^2}$ . By [S10, Theorem 1.1], we have

$$\begin{aligned}
 \sum_{k=1}^{p-1} \frac{C_k}{m^k} &\equiv m^{p-1} - 1 - \frac{m-4}{2} \left( \left( \frac{\Delta}{p} \right) - 1 + u_{p-\left(\frac{\Delta}{p}\right)}(m-2, 1) \right) \pmod{p^2} \\
 &\equiv -\frac{m-4}{2} \left( \left( \frac{\Delta}{p} \right) - 1 \right) \pmod{p},
 \end{aligned}$$

where

$$\Delta := m(m-4) \equiv -\frac{1}{x} \left( -\frac{1}{x} - 4 \right) = \frac{4x+1}{x^2} \pmod{p^2}.$$

So (2.8) follows.

If  $m \not\equiv 4 \pmod{p}$ , then by [S12b, Lemma 3.5] we have

$$\begin{aligned}
 \sum_{k=1}^{p-1} \frac{\binom{2k}{k}}{km^k} &\equiv \frac{(-m)^{p-1} - 1}{p} + \frac{m-4}{2} \left( \frac{\Delta}{p} \right) \frac{u_{p-\left(\frac{\Delta}{p}\right)}(2-m, 1)}{p} \\
 &= \frac{m^{p-1} - 1}{p} - \frac{m-4}{2} \left( \frac{\Delta}{p} \right) \frac{u_{p-\left(\frac{\Delta}{p}\right)}(m-2, 1)}{p} \pmod{p}
 \end{aligned}$$

and hence from the above we deduce that

$$\begin{aligned}
W_p(x) &\equiv 2p + 1 - x^{p-1} \\
&\quad + (p+1) \left( m^{p-1} - 1 - \frac{m-4}{2} \left( \left( \frac{\Delta}{p} \right) - 1 + u_{p-\left(\frac{\Delta}{p}\right)}(m-2, 1) \right) \right) \\
&\quad - \frac{p}{2} m \left( \frac{m^{p-1} - 1}{p} - \frac{m-4}{2} \left( \frac{\Delta}{p} \right) \frac{u_{p-\left(\frac{\Delta}{p}\right)}(m-2, 1)}{p} \right) \\
&\equiv 2p + 1 - x^{p-1} + \left( 1 - \frac{m}{2} \right) (m^{p-1} - 1) - (p+1) \frac{m-4}{2} \left( \left( \frac{4x+1}{p} \right) - 1 \right) \\
&\quad - \frac{m-4}{2} \left( 1 - \frac{m}{2} \left( \frac{4x+1}{p} \right) \right) u_{p-\left(\frac{4x+1}{p}\right)}(m-2, 1) \\
&\equiv 2p + 1 - x^{p-1} + \frac{2x+1}{2x} (1 - x^{p-1}) + (p+1) \frac{4x+1}{2x} \left( \left( \frac{4x+1}{p} \right) - 1 \right) \\
&\quad + \frac{4x+1}{2x} \left( 1 + \frac{1}{2x} \left( \frac{4x+1}{p} \right) \right) u_{p-\left(\frac{4x+1}{p}\right)}(m-2, 1) \pmod{p^2}
\end{aligned}$$

which is equivalent to (2.10) since

$$(-x)^{k-1} u_k(m-2, 1) = u_k(-x(m-2), (-x)^2) \equiv u_k(2x+1, x^2) \pmod{p^2}$$

for all  $k \in \mathbb{N}$ .

When  $m \equiv 4 \pmod{p}$  (i.e.,  $x \equiv -1/4 \pmod{p}$ ), we have

$$\sum_{k=1}^{p-1} \frac{\binom{2k}{k}}{k m^k} \equiv \sum_{k=1}^{p-1} \frac{\binom{2k}{k}}{k 4^k} \equiv 2q_p(2) \pmod{p}$$

by [ST, (1.12)], and hence

$$\begin{aligned}
W_p(x) &\equiv 2p + 1 - x^{p-1} + (p+1) \left( m^{p-1} - 1 + \frac{m-4}{2} \right) - \frac{p}{2} m 2q_p(2) \\
&\equiv 2p + 1 - x^{p-1} + m^{p-1} - 1 + \frac{m-4}{2} - 4(2^{p-1} - 1) \\
&\equiv 2p + 1 - x^{p-1} + 1 - x^{p-1} - \frac{4x+1}{2x} - 2(2^{2(p-1)} - 1) \\
&\equiv 2(p + (4x+1) + (1 - x^{p-1}) + (1 - 4^{p-1})) \\
&\equiv 2(p + (4x+1) + 1 - (4x+1 - 1)^{p-1}) \\
&\equiv 2(p + (4x+1) + (p-1)(4x+1)) \equiv 2p \pmod{p^2}.
\end{aligned}$$

Therefore (2.9) is valid.  $\square$

*Proof of Theorem 1.1.* (i) Fix  $n \in \mathbb{Z}^+$ . For each  $k \in \mathbb{Z}^+$ , by (2.3) and Lemma 2.2 we have

$$\begin{aligned}
 D_{k-1}(x) \frac{S_k(x)}{x+1} &= \frac{D_{k-1}(x) + D_{k+1}(x)}{2} \cdot \frac{S_k(x)}{x+1} - (2k+1) \frac{x}{x+1} S_k(x)^2 \\
 &= \sum_{j=0}^k \binom{k+j}{2j} \binom{2j}{j}^2 \frac{2j+1}{(j+1)^2} (x(x+1))^j \\
 &\quad - \frac{2k+1}{k(k+1)} \sum_{j=0}^k \binom{k+j}{2j} \binom{2j}{j} \binom{2j}{j+1} (x(x+1))^j \\
 &= \sum_{j=0}^k \binom{k+j}{2j} \binom{2j}{j}^2 \left( \frac{2j+1}{(j+1)^2} - \frac{2k+1}{k(k+1)} \cdot \frac{j}{j+1} \right) (x(x+1))^j \\
 &= \sum_{j=0}^k \binom{k+j}{2j} C_j^2 \left( 2j+1 - j(j+1) \frac{2k+1}{k(k+1)} \right) (x(x+1))^j.
 \end{aligned}$$

Combining this with (2.4) we obtain

$$D_{k-1}(x) s_k(x) = \sum_{j=0}^k \binom{k+j}{2j} C_j^2 \left( 2j+1 - j(j+1) \frac{2k+1}{k(k+1)} \right) (x(x+1))^j \quad (2.11)$$

for any  $k \in \mathbb{Z}^+$ . Therefore,

$$\begin{aligned}
 \sum_{k=1}^n D_{k-1}(x) s_k(x) &= \sum_{k=1}^n \sum_{j=0}^k \binom{k+j}{2j} C_j^2 \left( 2j+1 - j(j+1) \frac{2k+1}{k(k+1)} \right) (x(x+1))^j \\
 &= n + \sum_{j=1}^n C_j^2 (x(x+1))^j \sum_{k=j}^n \binom{k+j}{2j} \left( 2j+1 - j(j+1) \frac{2k+1}{k(k+1)} \right) \\
 &= \sum_{j=0}^{n-1} C_j^2 (x(x+1))^j \frac{(n-j)(n+j+1)}{n+1} \binom{n+j}{2j}
 \end{aligned}$$

with the help of Lemma 2.3. It follows that

$$\sum_{k=0}^{n-1} D_k(x) s_{k+1}(x) = \sum_{j=0}^{n-1} C_j(x(x+1))^j \frac{n}{j+1} \binom{n-1}{j} \binom{n+j+1}{j} = nW_n(x(x+1)).$$

For any  $k \in \mathbb{Z}^+$ , we have

$$w(n, k) = \frac{1}{n} \binom{n}{k} \binom{n+k}{k-1} = \frac{1}{n+1} \binom{n-1}{k-1} \binom{n+k}{k}$$

and hence  $w(n, k) = (n+1)w(n, k) - nw(n, k) \in \mathbb{Z}$ . So  $W_n(x) \in \mathbb{Z}[x]$ . This proves part (i) of Theorem 1.1.

(ii) Let  $x$  be any  $p$ -adic integer, and let  $y$  be an integer with  $y \equiv x(x+1) \pmod{p^2}$ . By (1.7) we have

$$\sum_{k=0}^{p-1} D_k(x) s_{k+1}(x) = pW_p(x(x+1)) \equiv pW_p(y) \pmod{p^3}. \quad (2.12)$$

If  $y \equiv 0 \pmod{p}$ , then

$$W_p(y) \equiv w(p, 1) + w(p, 2)y \equiv 1 - x(x+1) \pmod{p^2}.$$

If  $x \equiv -1/2 \pmod{p}$  (i.e.,  $y \equiv -1/4 \pmod{p}$ ), then  $W_p(y) \equiv 2p \pmod{p^2}$  by Lemma 2.4. Thus (1.10) holds for  $x \equiv 0, -1, -1/2 \pmod{p}$ .

Now assume that  $x \not\equiv 0, -1, -1/2 \pmod{p}$ , i.e.,  $y \not\equiv 0, -1/4 \pmod{p}$ . Then

$$\left(\frac{4y+1}{p}\right) = \left(\frac{(2x+1)^2}{p}\right) = 1$$

and

$$\begin{aligned} u_{p-1}(2y+1, y^2) &\equiv u_{p-1}(x^2 + (x+1)^2, x^2(x+1)^2) \\ &= \frac{((x+1)^2)^{p-1} - (x^2)^{p-1}}{(x+1)^2 - x^2} \\ &= \frac{(x+1)^{p-1} + x^{p-1}}{2x+1} ((x+1)^{p-1} - x^{p-1}) \\ &\equiv \frac{2}{2x+1} ((x+1)^{p-1} - x^{p-1}) \pmod{p^2}. \end{aligned}$$

Combining this with Lemma 2.4, we obtain

$$\begin{aligned} W_p(y) &\equiv 2p + \frac{(2x+1)^2}{2x(x+1)} (1 - x^{p-1}(x+1)^{p-1}) \\ &\quad - \frac{(2x+1)^2}{4x(x+1)} (2x(x+1) + 1) \frac{2}{2x+1} ((x+1)^{p-1} - x^{p-1}) \\ &\equiv 2p + \frac{(2x+1)^2}{2x(x+1)} (1 - x^{p-1} + 1 - (x+1)^{p-1}) \\ &\quad - \frac{2x+1}{2x(x+1)} (2x^2 + 2x + 1) ((x+1)^{p-1} - x^{p-1}) \\ &= 2p + \frac{2x+1}{x(x+1)} (x^2(x^{p-1} - 1) - (x+1)^2((x+1)^{p-1} - 1)) \pmod{p^2}. \end{aligned}$$

Therefore (1.10) holds in light of (2.12).

So far we have completed the proof of Theorem 1.1.  $\square$

*Proof of Corollary 1.2.* It is known (cf. [G, (3.133)]) that

$$D_k\left(-\frac{1}{2}\right) = F_k(0) = \begin{cases} (-1)^{k/2} \binom{k}{k/2} / 2^k & \text{if } 2 \mid k, \\ 0 & \text{otherwise.} \end{cases}$$

By (2.4) and [S11a, Lemma 4.3],

$$s_{k+1} \left( -\frac{1}{2} \right) = 2S_{k+1} \left( -\frac{1}{2} \right) = \begin{cases} (-1)^{k/2} C_{k/2} / 2^k & \text{if } 2 \mid k, \\ 0 & \text{otherwise.} \end{cases}$$

Therefore

$$\begin{aligned} & \sum_{k=0}^{n-1} D_k \left( -\frac{1}{2} \right) s_{k+1} \left( -\frac{1}{2} \right) \\ &= \sum_{\substack{0 \leq k < n \\ 2 \mid k}} \left( \frac{(-1)^{k/2}}{2^k} \right)^2 \binom{k}{k/2} C_{k/2} = \sum_{j=0}^{\lfloor (n-1)/2 \rfloor} \frac{\binom{2j}{j}^2}{(j+1)16^j}. \end{aligned}$$

By induction,

$$\sum_{j=0}^m \frac{\binom{2j}{j}^2}{(j+1)16^j} = \frac{(2m+1)^2}{(m+1)16^m} \binom{2m}{m}^2 \quad \text{for all } m \in \mathbb{N}.$$

So we have

$$\begin{aligned} & \sum_{k=0}^{n-1} D_k \left( -\frac{1}{2} \right) s_{k+1} \left( -\frac{1}{2} \right) \\ &= \frac{(2\lfloor (n-1)/2 \rfloor + 1)^2}{[\lfloor (n+1)/2 \rfloor] 16^{\lfloor (n-1)/2 \rfloor}} \left( \frac{2\lfloor (n-1)/2 \rfloor}{[\lfloor (n-1)/2 \rfloor]} \right)^2 = \frac{[\lfloor (n+1)/2 \rfloor]}{4^{n-1}} \binom{n}{\lfloor n/2 \rfloor}^2. \end{aligned}$$

On the other hand, by applying (1.7) with  $x = -1/2$  we obtain

$$\sum_{k=0}^{n-1} D_k \left( -\frac{1}{2} \right) s_{k+1} \left( -\frac{1}{2} \right) = nW_n \left( -\frac{1}{4} \right) = \sum_{k=1}^n \binom{n}{k} \binom{n+k}{k-1} \frac{C_{k-1}}{(-4)^{k-1}}.$$

Combining these we get the desired identity (1.13). This concludes the proof.  $\square$

**Lemma 2.5.** *Let  $p$  be any odd prime. For each  $j = 0, \dots, p$ , we have*

$$C_j^2 u_j \equiv \begin{cases} 2 \pmod{p} & \text{if } j = (p-1)/2, \\ 0 \pmod{p} & \text{otherwise,} \end{cases} \quad (2.13)$$

where

$$u_j := \sum_{j < k \leq p} (k-1) \binom{k+j}{2j} \left( 2j+1 - j(j+1) \frac{2k+1}{k(k+1)} \right). \quad (2.14)$$

*Proof.* Clearly,  $u_p = 0$  and

$$u_{p-1} = (p-1) \binom{p+(p-1)}{2(p-1)} \left( 2(p-1) + 1 - (p-1)p \frac{2p+1}{p(p+1)} \right) \equiv 0 \pmod{p}.$$

If  $(p-1)/2 < j < p-1$ , then  $C_j = (2j)!/(j!(j+1)!) \equiv 0 \pmod{p}$  and hence  $C_j^2 u_j \equiv 0 \pmod{p}$  since  $pu_j$  is a  $p$ -adic integer.

Note that

$$C_{(p-1)/2} = \frac{2}{p+1} \binom{p-1}{(p-1)/2} \equiv 2(-1)^{(p-1)/2} \pmod{p}.$$

As

$$\binom{k + (p-1)/2}{p-1} \equiv 0 \pmod{p} \quad \text{for all } k = \frac{p+1}{2}, \dots, p,$$

we have

$$\begin{aligned} u_{(p-1)/2} &\equiv \sum_{k=p-1}^p (k-1) \binom{k + (p-1)/2}{p-1} \left( 2 \cdot \frac{p-1}{2} + 1 - \frac{p-1}{2} \cdot \frac{p+1}{2} \cdot \frac{2k+1}{k(k+1)} \right) \\ &\equiv \frac{1}{4} \sum_{k=p-1}^p (k-1) \binom{k + (p-1)/2}{p-1} \frac{2k+1}{k(k+1)} \\ &= \frac{p-2}{4} \binom{p-1 + (p-1)/2}{(p-1)/2} \frac{2(p-1)+1}{(p-1)p} + \frac{p-1}{4} \binom{p + (p-1)/2}{(p+1)/2} \frac{2p+1}{p(p+1)} \\ &= \frac{p-2}{4} \cdot \frac{2p-1}{p-1} \cdot \frac{\prod_{1 < r \leq (p-1)/2} (p-1+r)}{((p-1)/2)!} \\ &\quad + \frac{p-1}{4} \cdot \frac{2p+1}{p+1} \cdot \frac{\prod_{r=1}^{(p-1)/2} (p+r)}{((p+1)/2)!} \\ &\equiv -\frac{1}{2} \cdot \frac{1}{(p-1)/2} - \frac{1}{4} \cdot \frac{1}{(p+1)/2} \equiv \frac{1}{2} \pmod{p}. \end{aligned}$$

Obviously,

$$u_0 = \sum_{k=1}^p (k-1) = \frac{p(p-1)}{2} \equiv 0 \pmod{p}.$$

Applying the Zeilberger algorithm via `Mathematica 9`, we find that

$$(j+2)u_j + 2(2j+1)u_{j+1} = \frac{f(p, j) \binom{p+j}{2j}}{2(j+1)(j+2)(2j+3)}$$

for all  $j = 0, \dots, p-1$ , where

$$f(p, j) := (p-j)(p+j+1) \left( (2j+3)^2 p^2 - (2j^2 + 8j + 7)p - (j+1)(j+2) \right).$$

This implies that for  $0 \leq j < (p-3)/2$  we have

$$u_j \equiv 0 \pmod{p} \implies u_{j+1} \equiv 0 \pmod{p}.$$

Thus  $u_j \equiv 0 \pmod{p}$  for all  $j = 0, \dots, (p-3)/2$ .

Combining the above, we immediately obtain the desired (2.13).  $\square$

*Proof of Theorem 1.2.* In view of (2.11),

$$\begin{aligned} & \sum_{k=1}^p (k-1) D_{k-1}(x) s_k(x) \\ &= \sum_{k=1}^p (k-1) \sum_{j=0}^k \binom{k+j}{2j} C_j^2 \left( 2j+1 - j(j+1) \frac{2k+1}{k(k+1)} \right) (x(x+1))^j \\ &= \sum_{k=1}^p (k-1) + \sum_{j=1}^p (x(x+1))^j C_j^2 u_j = \frac{p(p-1)}{2} + \sum_{j=1}^p C_j^2 u_j (x(x+1))^j, \end{aligned}$$

where  $u_j$  is given by (2.14). Thus, by applying Lemma 2.5 we find that

$$\frac{1}{p} \left( \sum_{k=0}^{p-1} k D_k(x) s_{k+1}(x) - 2(x(x+1))^{(p-1)/2} \right) \in \mathbb{Z}_p[x(x+1)], \quad (2.15)$$

where  $\mathbb{Z}_p$  denotes the ring of  $p$ -adic integers. Therefore (1.14) holds. (1.14) with  $x = 1$  gives (1.15). This concludes the proof.  $\square$

### 3. APPLICATIONS TO CENTRAL TRINOMIAL COEFFICIENTS AND MOTZKIN NUMBERS

Let  $n \in \mathbb{N}$  and  $b, c \in \mathbb{Z}$ . The  $n$ -th generalized central trinomial coefficient  $T_n(b, c)$  is defined to be  $[x^n](x^2 + bx + c)^n$ , the coefficient of  $x^n$  in the expansion of  $(x^2 + bx + c)^n$ . It is easy to see that

$$T_n(b, c) = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{2k} \binom{2k}{k} b^{n-2k} c^k. \quad (3.1)$$

Note that  $T_n(1, 1)$  is the central trinomial coefficient  $T_n$  and  $T_n(2, 1)$  is the central binomial coefficient  $\binom{2n}{n}$ . Also,  $T_n(3, 2)$  coincides with the central Delannoy number  $D_n$ . Sun [S14a] also defined the generalized Motzkin number  $M_n(b, c)$  by

$$M_n(b, c) = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{2k} C_k b^{n-2k} c^k. \quad (3.2)$$

Note that  $M_n(1, 1)$  is the usual Motzkin number  $M_n$  (whose combinatorial interpretations can be found in [St99, Ex. 6.38]) and  $M_n(2, 1)$  is the Catalan number  $C_{n+1}$ . Also,  $M_n(3, 2)$  coincides with the little Schröder number  $s_{n+1}$ . The author [S14a, S14b] deduced some congruences involving  $T_n(b, c)$  and  $M_n(b, c)$ , and

proposed in [S14b] some conjectural series for  $1/\pi$  involving  $T_n(b, c)$  such as

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{66k+17}{(2^{11}3^3)^k} T_k(10, 11^2)^3 &= \frac{540\sqrt{2}}{11\pi}, \\ \sum_{k=0}^{\infty} \frac{126k+31}{(-80)^{3k}} T_k(22, 21^2)^3 &= \frac{880\sqrt{5}}{21\pi}, \\ \sum_{k=0}^{\infty} \frac{3990k+1147}{(-288)^{3k}} T_k(62, 95^2)^3 &= \frac{432}{95\pi} (195\sqrt{14} + 94\sqrt{2}). \end{aligned}$$

Now we point out that  $T_n(b, c)$  and  $M_n(b, c)$  are actually related to the polynomials  $D_n(x)$  and  $s_{n+1}(x)$ .

**Lemma 3.1.** *Let  $b, c \in \mathbb{Z}$  with  $d = b^2 - 4c \neq 0$ . For any  $n \in \mathbb{N}$  we have*

$$T_n(b, c) = (\sqrt{d})^n D_n \left( \frac{b/\sqrt{d} - 1}{2} \right) \quad (3.3)$$

and

$$M_n(b, c) = (\sqrt{d})^n s_{n+1} \left( \frac{b/\sqrt{d} - 1}{2} \right). \quad (3.4)$$

*Proof.* In view of (3.1) and (3.2),

$$\frac{T_n(b, c)}{(\sqrt{d})^n} = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{2k} \binom{2k}{k} \left( \frac{b}{\sqrt{d}} \right)^{n-2k} \left( \frac{c}{d} \right)^k$$

and

$$\frac{M_n(b, c)}{(\sqrt{d})^n} = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{2k} C_k \left( \frac{b}{\sqrt{d}} \right)^{n-2k} \left( \frac{c}{d} \right)^k.$$

Note that

$$\left( \frac{b}{\sqrt{d}} \right)^2 - 4 \frac{c}{d} = 1.$$

So, it suffices to show the polynomial identities

$$\sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{2k} \binom{2k}{k} (2x+1)^{n-2k} (x(x+1))^k = D_n(x) \quad (3.5)$$

and

$$\sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{2k} C_k (2x+1)^{n-2k} (x(x+1))^k = s_{n+1}(x). \quad (3.6)$$



It is easy to verify (3.5) and (3.6) for  $n = 0, 1, 2$ . Let  $u_n(x)$  denote the left-hand side or the right-hand side of (3.5). By the Zeilberger algorithm (cf. [PWZ, pp. 101-119]), we have the recurrence

$$(n + 2)u_{n+2}(x) = (2x + 1)(2n + 3)u_{n+1}(x) - (n + 1)u_n(x) \quad \text{for } n = 0, 1, 2, \dots$$

Thus (3.5) is valid by induction. Let  $v_n(x)$  denote the left-hand side or the right-hand side of (3.6). By the Zeilberger algorithm, we have the recurrence

$$(n + 4)v_{n+2}(x) = (2x + 1)(2n + 5)v_{n+1}(x) - (n + 1)v_n(x) \quad \text{for } n = 0, 1, 2, \dots$$

So (3.6) also holds by induction.

The proof of Lemma 3.1 is now complete.  $\square$

With the help of Theorem 1.1, we are able to confirm Conjecture 5.5 of the author [S14a] by proving the following result.

**Theorem 3.1.** *Let  $b, c \in \mathbb{Z}$  and  $d = b^2 - 4c$ .*

(i) For any  $n \in \mathbb{Z}^+$ , we have

$$\frac{1}{n} \sum_{k=0}^{n-1} T_k(b, c) M_k(b, c) d^{n-1-k} = \sum_{k=1}^n w(n, k) C_{k-1} c^{k-1} d^{n-k} \in \mathbb{Z}. \quad (3.7)$$

Moreover, for any odd prime  $p$  not dividing  $cd$ , we have

$$\sum_{k=0}^{p-1} \frac{T_k(b, c) M_k(b, c)}{d^k} \equiv \frac{pb^2}{2c} \left( \left( \frac{d}{p} \right) - 1 \right) \pmod{p^2}, \quad (3.8)$$

and furthermore

$$\begin{aligned} & \sum_{k=0}^{p-1} \frac{T_k(b, c) M_k(b, c)}{d^k} \\ & \equiv \frac{pb^2}{2c} \left( \left( \frac{d}{p} \right) - 1 \right) + \frac{p^2}{2c} \left( b^2 \left( q_p(d) - q_p(c) + \left( \frac{d}{p} \right) \right) - d \right) \\ & \quad - \frac{pb^2}{4c^{2-\left(\frac{d}{p}\right)}} \left( 2c + d \left( \frac{d}{p} \right) \right) u_{p-\left(\frac{d}{p}\right)}(b^2 - 2c, c^2) \pmod{p^3}. \end{aligned} \quad (3.9)$$

(ii) For any odd prime  $p$  not dividing  $d$ , we have

$$\sum_{k=0}^{p-1} \frac{kT_k(b, c) M_k(b, c)}{d^k} \equiv 2 \left( \frac{cd}{p} \right) \pmod{p}. \quad (3.10)$$

*Proof.* (i) Let's first prove (3.7) for any  $n \in \mathbb{Z}^+$ .

We first consider the case  $d = 0$ , i.e.,  $c = b^2/4$ . In this case, for any  $k \in \mathbb{N}$  we have

$$T_k(b, c) = \left(\frac{b}{2}\right)^k T_k(2, 1) = \left(\frac{b}{2}\right)^k \binom{2k}{k}$$

and

$$M_k(b, c) = \left(\frac{b}{2}\right)^k M_k(2, 1) = \left(\frac{b}{2}\right)^k C_{k+1}.$$

Thus

$$\begin{aligned} & \frac{1}{n} \sum_{k=0}^{n-1} T_k(b, c) M_k(b, c) d^{n-1-k} \\ &= \frac{T_{n-1}(b, c) M_{n-1}(b, c)}{n} = \frac{1}{n} \left(\frac{b^2}{4}\right)^{n-1} \binom{2(n-1)}{n-1} C_n \\ &= c^{n-1} C_{n-1} C_n = w(n, n) C_{n-1} c^{n-1} \end{aligned}$$

and hence (3.7) is valid.

Now assume that  $d \neq 0$ . By Lemma 3.1 and (1.7), we have

$$\begin{aligned} & \frac{1}{n} \sum_{k=0}^{n-1} \frac{T_k(b, c) M_k(b, c)}{d^k} \\ &= \frac{1}{n} \sum_{k=0}^{n-1} D_k \left(\frac{b/\sqrt{d}-1}{2}\right) s_{k+1} \left(\frac{b/\sqrt{d}-1}{2}\right) \\ &= W_n \left(\frac{b/\sqrt{d}-1}{2} \cdot \frac{b/\sqrt{d}+1}{2}\right) = W_n \left(\frac{b^2/d-1}{4}\right) = W_n \left(\frac{c}{d}\right) \end{aligned}$$

and hence (3.7) holds in view of (1.8).

Below we suppose that  $p$  is an odd prime not dividing  $cd$ . From the above, we have

$$\sum_{k=0}^{p-1} \frac{T_k(b, c) M_k(b, c)}{d^k} = p W_p \left(\frac{c}{d}\right) \equiv p W_p(x) \pmod{p^3}, \quad (3.11)$$

where  $x$  is an integer with  $x \equiv c/d \pmod{p^2}$ . As  $p \nmid d$  and  $d(4x+1) \equiv 4c+d = b^2 \pmod{p^2}$ , we have

$$\left(\frac{4x+1}{p}\right) = \left(\frac{d^2(4x+1)}{p}\right) = \left(\frac{b^2 d}{p}\right).$$

In view of Lemma 2.4,

$$W_p(x) \equiv \frac{4c/d+1}{2c/d} \left(\left(\frac{4x+1}{p}\right) - 1\right) \equiv \frac{b^2}{2c} \left(\left(\frac{d}{p}\right) - 1\right) \pmod{p}.$$

Combining this with (3.11) we immediately obtain (3.8).

Now we show (3.9). If  $x \equiv -1/4 \pmod{p}$  (i.e.,  $p \mid b$ ), then by (2.9) we have

$$W_p(x) \equiv 2p \equiv \frac{p}{2c}(4c - b^2) \pmod{p^2}$$

and hence (3.9) holds by (3.11). Below we assume  $p \nmid b$ . Then

$$\left(\frac{(2x+1)^2 - 4x^2}{p}\right) = \left(\frac{4x+1}{p}\right) = \left(\frac{b^2d}{p}\right) = \left(\frac{d}{p}\right),$$

$p \mid u_{p-(\frac{d}{p})}(2x+1, x^2)$  and

$$\begin{aligned} & d^{p-1-(\frac{d}{p})} u_{p-(\frac{d}{p})}(2x+1, x^2) \\ &= u_{p-(\frac{d}{p})}(d(2x+1), d^2x^2) \\ &\equiv u_{p-(\frac{d}{p})}(2c+d, c^2) = u_{p-(\frac{d}{p})}(b^2-2c, c^2) \pmod{p^2}. \end{aligned}$$

So, applying (2.10) we get

$$\begin{aligned} W_p(x) &\equiv 2p + \frac{4c/d+1}{2c/d} \left(1 - \left(\frac{c}{d}\right)^{p-1} + (p+1) \left(\left(\frac{d}{p}\right) - 1\right)\right) \\ &\quad - \frac{4c/d+1}{4(c/d)^{2-(\frac{d}{p})}} \left(2\frac{c}{d} + \left(\frac{d}{p}\right)\right) d^{(\frac{d}{p})} u_{p-(\frac{d}{p})}(b^2-2c, c^2) \\ &\equiv 2p + \frac{b^2}{2c} \left(d^{p-1} - 1 - (c^{p-1} - 1) + (p+1) \left(\left(\frac{d}{p}\right) - 1\right)\right) \\ &\quad - \frac{b^2}{4c^{2-(\frac{d}{p})}} \left(2c+d\left(\frac{d}{p}\right)\right) u_{p-(\frac{d}{p})}(b^2-2c, c^2) \pmod{p^2}. \end{aligned}$$

This, together with (3.11), yields the desired (3.9).

(ii) Fix an odd prime  $p$  not dividing  $d$ . Let  $x = b/\sqrt{d} - 1$ . Then

$$x(x+1) = \frac{b/\sqrt{d}-1}{2} \cdot \frac{b/\sqrt{d}+1}{2} = \frac{b^2/d-1}{4} = \frac{c}{d}$$

is a  $p$ -adic integer. Thus, with the help of (2.15), we have

$$\sum_{k=0}^{p-1} kD_k(x)s_{k+1}(x) \equiv 2\left(\frac{c}{d}\right)^{(p-1)/2} \equiv 2\left(\frac{cd}{p}\right) \pmod{p}.$$

Combining this with Lemma 3.1, we immediately obtain (3.10).

In view of the above, we have proved Theorem 3.1.  $\square$

Let  $\omega$  denote the primitive cubic root  $(-1 + \sqrt{-3})/2$  of unity. Then  $\omega + \bar{\omega} = -1$  and  $\omega\bar{\omega} = 1$ . So,

$$u_n(-1, 1) = \frac{\omega^n - \bar{\omega}^n}{\omega - \bar{\omega}} = 0 \quad \text{and} \quad u_n(3, 9) = \frac{(-3\omega)^n - (-3\bar{\omega})^n}{(-3\omega) - (-3\bar{\omega})} = 0$$

for any  $n \in \mathbb{N}$  with  $3 \mid n$ . In view of this, Theorem 3.1 in the cases  $b = c \in \{1, 3\}$  yields the following consequence.

**Corollary 3.1.** *For any positive integer  $n$ , we have*

$$\frac{1}{n} \sum_{k=0}^{n-1} T_k M_k (-3)^{n-1-k} = \sum_{k=1}^n w(n, k) C_{k-1} (-3)^{n-k} \in \mathbb{Z} \quad (3.12)$$

and

$$\frac{1}{n} \sum_{k=0}^{n-1} \frac{T_k(3, 3) M_k(3, 3)}{(-3)^k} = \sum_{k=1}^n (-1)^{k-1} w(n, k) C_{k-1} \in \mathbb{Z}. \quad (3.13)$$

Moreover, for any prime  $p > 3$  we have

$$\sum_{k=0}^{p-1} \frac{T_k M_k}{(-3)^k} \equiv \frac{p}{2} \left( \binom{p}{3} - 1 \right) + \frac{p^2}{2} \left( q_p(3) + \binom{p}{3} + 3 \right) \pmod{p^3}, \quad (3.14)$$

$$\sum_{k=0}^{p-1} \frac{T_k(3, 3) M_k(3, 3)}{(-3)^k} \equiv \frac{3p}{2} \left( \binom{p}{3} - 1 \right) + \frac{p^2}{2} \left( 3 \binom{p}{3} + 1 \right) \pmod{p^3}, \quad (3.15)$$

$$\sum_{k=0}^{p-1} \frac{k T_k M_k}{(-3)^k} \equiv 2 \binom{p}{3} \pmod{p} \text{ and } \sum_{k=0}^{p-1} \frac{k T_k(3, 3) M_k(3, 3)}{(-3)^k} \equiv 2 \binom{-1}{p} \pmod{p}. \quad (3.16)$$

*Remark 3.1.* Let  $p > 3$  be a prime. In the case  $p \equiv 2 \pmod{3}$ , the author (cf. [S14a, Conjecture 5.6]) even conjectured that

$$\sum_{k=0}^{p-1} \frac{T_k(3, 3) M_k(3, 3)}{(-3)^k} \equiv p^3 - p^2 - 3p \pmod{p^4}$$

which is stronger than (3.15). The author's conjectural supercongruences (cf. [S14a, Conjecture 1.1(ii)])

$$\sum_{k=0}^{p-1} M_k^2 \equiv (2 - 6p) \binom{p}{3} \pmod{p^2}, \quad \sum_{k=0}^{p-1} k M_k^2 \equiv (9p - 1) \binom{p}{3} \pmod{p^2},$$

and

$$\sum_{k=0}^{p-1} T_k M_k \equiv \frac{4}{3} \binom{p}{3} + \frac{p}{6} \left( 1 - 9 \binom{p}{3} \right) \pmod{p^2}$$

remain open. We also observe that

$$\sum_{k=0}^{p-1} k T_k M_k \equiv \binom{-1}{p} - \frac{5}{3} \binom{p}{3} \pmod{p}.$$

The Lucas numbers  $L_0, L_1, L_2, \dots$  are given by

$$L_0 = 2, L_1 = 1, \text{ and } L_{n+1} = L_n + L_{n-1} \text{ for } n = 1, 2, 3, \dots$$

It is easy to see that  $L_n = 2F_{n+1} - F_n = 2F_{n-1} + F_n$  for all  $n \in \mathbb{Z}^+$ . Thus, for any odd prime  $p \neq 5$  we have

$$L_{p-\left(\frac{p}{5}\right)} = 2F_p - \left(\frac{p}{5}\right) F_{p-\left(\frac{p}{5}\right)} \equiv 2 \left(\frac{p}{5}\right) \pmod{p}$$

and hence

$$u_{p-\left(\frac{p}{5}\right)}(3, 1) = \frac{(\alpha^2)^{p-\left(\frac{p}{5}\right)} - (\beta^2)^{p-\left(\frac{p}{5}\right)}}{\alpha^2 - \beta^2} = F_{p-\left(\frac{p}{5}\right)} L_{p-\left(\frac{p}{5}\right)} \equiv 2 \left(\frac{p}{5}\right) F_{p-\left(\frac{p}{5}\right)} \pmod{p^2},$$

where  $\alpha = (1 + \sqrt{5})/2$  and  $\beta = (1 - \sqrt{5})/2$ . Note also that

$$u_n(3 \times 5, 5^2) = 5^{n-1} u_n(3, 1) = 5^{n-1} F_n L_n \quad \text{for any } n \in \mathbb{N}.$$

Thus Theorem 3.1 with  $(b, c) = (1, -1), (5, 5)$  leads to the following corollary.

**Corollary 3.2.** *For any  $n \in \mathbb{Z}^+$ , we have*

$$\frac{1}{n} \sum_{k=0}^{n-1} T_k(1, -1) M_k(1, -1) 5^{n-1-k} = \sum_{k=1}^n (-1)^{k-1} w(n, k) C_{k-1} 5^{n-k} \in \mathbb{Z} \quad (3.17)$$

and

$$\frac{1}{n} \sum_{k=0}^{n-1} \frac{T_k(5, 5) M_k(5, 5)}{5^k} = \sum_{k=1}^n w(n, k) C_{k-1} \in \mathbb{Z}. \quad (3.18)$$

Also, for any prime  $p \neq 2, 5$  we have the congruences

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{T_k(1, -1) M_k(1, -1)}{5^k} &\equiv \frac{p}{2} \left(1 - \left(\frac{p}{5}\right)\right) + \frac{p^2}{2} \left(5 - \left(\frac{p}{5}\right) - q_p(5)\right) \\ &\quad + \frac{p}{2} \left(5 - 2 \left(\frac{p}{5}\right)\right) F_{p-\left(\frac{p}{5}\right)} \pmod{p^3}, \end{aligned} \quad (3.19)$$

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{T_k(5, 5) M_k(5, 5)}{5^k} &\equiv \frac{5p}{2} \left(\left(\frac{p}{5}\right) - 1\right) + \frac{p^2}{2} \left(5 \left(\frac{p}{5}\right) - 1\right) \\ &\quad - \frac{5p}{2} \left(1 + 2 \left(\frac{p}{5}\right)\right) F_{p-\left(\frac{p}{5}\right)} \pmod{p^3}, \end{aligned} \quad (3.20)$$

and

$$\left(\frac{-5}{p}\right) \sum_{k=0}^{p-1} \frac{k T_k(1, -1) M_k(1, -1)}{5^k} \equiv \sum_{k=0}^{p-1} \frac{k T_k(5, 5) M_k(5, 5)}{5^k} \equiv 2 \pmod{p}. \quad (3.21)$$

## 4. TWO RELATED CONJECTURES

In view of (1.8) and (1.9), we can easily see that

$$\begin{aligned} W_1(x) &= 1, \quad W_2(x) = 2x + 1, \quad W_3(x) = 10x^2 + 5x + 1, \\ W_4(x) &= 70x^3 + 42x^2 + 9x + 1. \end{aligned}$$

Applying the Zeilberger algorithm (cf. [PWZ, pp.101-119]) via *Mathematica 9*, we obtain the following third-order recurrence with  $n \in \mathbb{Z}^+$ :

$$\begin{aligned} & (n+3)^2(n+4)(2n+3)W_{n+3}(x) \\ &= (n+3)(2n+5)(4x(2n+3)^2 + 3n^2 + 11n + 10)W_{n+2}(x) \\ & \quad - (n+1)(2n+3)(4x(2n+5)^2 + 3n^2 + 13n + 14)W_{n+1}(x) \\ & \quad + n(n+1)^2(2n+5)W_n(x). \end{aligned} \tag{4.1}$$

(This is a verified result, not a conjecture.)

For any  $n \in \mathbb{Z}^+$ , we clearly have  $w(n, n) = C_n$ . For the polynomial

$$w_n(x) := \sum_{k=1}^n w(n, k)x^{k-1}, \tag{4.2}$$

we have the relation

$$w_n(-1-x) = (-1)^{n-1}w_n(x) \tag{4.3}$$

since

$$\sum_{k=m}^n (-1)^{n-k} \binom{k-1}{m-1} w(n, k) = w(n, m) \quad \text{for all } m = 1, \dots, n, \tag{4.4}$$

which can be deduced with the help of the Chu-Vandermonde identity in the following way:

$$\begin{aligned} & \sum_{k=m}^n (-1)^{n-k} \binom{k-1}{m-1} w(n, k) \\ &= \binom{n-1}{m-1} \sum_{k=m}^n \frac{(-1)^{n-k}}{k} \binom{n-m}{n-k} \binom{-n-2}{k-1} (-1)^{k-1} \\ &= \frac{(-1)^n}{n+1} \binom{n-1}{m-1} \sum_{k=m}^n \binom{n-m}{n-k} \binom{-n-1}{k} \\ &= \frac{(-1)^n}{n+1} \binom{n-1}{m-1} \binom{-m-1}{n} = w(n, m). \end{aligned}$$

Via the Zeilberger algorithm we obtain the recurrence

$$(n+3)w_{n+2}(x) = (2x+1)(2n+3)w_{n+1}(x) - nw_n(x) \quad \text{for } n = 1, 2, 3, \dots \tag{4.5}$$

As  $w_2(x) = 2x+1$ , this recurrence implies that  $w_{2n}(-1/2) = 0$  and hence  $w_{2n}(x)/(2x+1) \in \mathbb{Z}[x]$  for all  $n \in \mathbb{Z}^+$ . We also note that

$$\sum_{n=1}^{\infty} w_n(x)y^n = \frac{1 - y - 2xy - \sqrt{(y-1)^2 - 4xy}}{2x(x+1)y}, \quad (4.6)$$

while

$$\sum_{n=0}^{\infty} S_n y^n = \frac{1 - y - \sqrt{y^2 - 6y + 1}}{2y}.$$

Now we pose two conjectures for further research.

**Conjecture 4.1.** *For any integer  $n > 1$ , all the polynomials*

$$w_{2n-1}(x), \quad \frac{w_{2n}(x)}{2x+1} \quad \text{and} \quad W_n(x)$$

*are irreducible over the field of rational numbers.*

**Conjecture 4.2.** (i) *For any  $n \in \mathbb{Z}^+$ , we have*

$$f_n(x) := \frac{1}{n} \sum_{k=0}^{n-1} D_k(x)R_k(x) \in \mathbb{Z}[x], \quad (4.7)$$

where

$$R_k(x) := \sum_{l=0}^k \binom{k}{l} \binom{k+l}{l} \frac{x^l}{2l-1} = \sum_{l=0}^k \binom{k+l}{2l} \binom{2l}{l} \frac{x^l}{2l-1}. \quad (4.8)$$

Also,  $f_2(x), f_3(x), \dots$  are all irreducible over the field of rational numbers, and

$$f_n(1) = \frac{1}{n} \sum_{k=0}^{n-1} D_k R_k \equiv (-1)^n \pmod{32}$$

for each  $n \in \mathbb{Z}^+$ , where  $R_k = R_k(1)$ .

(ii) *Let  $p$  be any odd prime. Then*

$$\sum_{k=0}^{p-1} D_k R_k \equiv \begin{cases} -p + 8p^2 q_p(2) - 2p^3 E_{p-3} \pmod{p^4} & \text{if } p \equiv 1 \pmod{4}, \\ -5p \pmod{p^3} & \text{if } p \equiv 3 \pmod{4}. \end{cases} \quad (4.9)$$

Also,

$$\sum_{k=1}^{p-1} \frac{D_k R_k}{k} \equiv \left( 4 - \left( \frac{-1}{p} \right) \right) q_p(2) \pmod{p}, \quad (4.10)$$

$$\sum_{k=1}^{p-1} k D_k R_k \equiv \frac{1}{2} + \frac{3}{2} p \left( 1 - 2 \left( \frac{-1}{p} \right) \right) \pmod{p^2}, \quad (4.11)$$

and

$$\sum_{k=1}^{p-1} k D_k(x) R_k(x) \equiv \frac{x^{p-1}}{2} \pmod{p}.$$

**Acknowledgment.** The author would like to thank the referee for helpful comments.

## REFERENCES

- [G] H. W. Gould, *Combinatorial Identities*, Morgantown Printing and Binding Co., 1972.
- [Gr] R. P. Grimaldi, *Fibonacci Numbers and Catalan Numbers: An Introduction*, John Wiley & Sons, New Jersey, 2012.
- [L] J.-C. Liu, *A supercongruence involving Delannoy numbers and Schröder numbers*, J. Number Theory **168** (2016), 117–127.
- [PWZ] M. Petkovšek, H. S. Wilf and D. Zeilberger, *A = B*, A K Peters, Wellesley, 1996.
- [St97] R. P. Stanley, *Hipparchus, Plutarch, Schröder, and Hough*, Amer. Math. Monthly **104** (1997), 344–350.
- [St99] R. P. Stanley, *Enumerative Combinatorics*, Vol. 2, Cambridge Univ. Press, Cambridge, 1999.
- [S10] Z.-W. Sun, *Binomial coefficients, Catalan numbers and Lucas quotients*, Sci. China Math. **53** (2010), 2473–2488.
- [S11a] Z.-W. Sun, *On congruences related to central binomial coefficients*, J. Number Theory **131** (2011), 2219–2238.
- [S11b] Z.-W. Sun, *On Delannoy numbers and Schröder numbers*, J. Number Theory **131** (2011), 2387–2397.
- [S12a] Z.-W. Sun, *On sums involving products of three binomial coefficients*, Acta Arith. **156** (2012), 123–141.
- [S12b] Z.-W. Sun, *On sums of binomial coefficients modulo  $p^2$* , Colloq. Math. **127** (2012), 39–54.
- [S14a] Z.-W. Sun, *Congruences involving generalized central trinomial coefficients*, Sci. China Math. **57** (2014), 1375–1400.
- [S14b] Z.-W. Sun, *On sums related to central binomial and trinomial coefficients*, in: M. B. Nathanson (ed.), *Combinatorial and Additive Number Theory: CANT 2011 and 2012*, Springer Proc. in Math. & Stat., Vol. 101, Springer, New York, 2014, pp. 257–312.
- [ST] Z.-W. Sun and R. Tauraso, *New congruences for central binomial coefficients*, Adv. in Appl. Math. **45** (2010), 125–148.
- [W] J. Wolstenholme, *On certain properties of prime numbers*, Quart. J. Appl. Math. **5** (1862), 35–39.