On a lower bound for sorting signed permutations by reversals

Andrei C. Bura^a, Ricky X. F. Chen^b, Christian M. Reidys^c

Biocomplexity Institute and Dept. of Mathematics, Virginia Tech, 1015 Life Sciences Circle, Blacksburg, VA 24061, USA

 a anbur12@vbi.vt.edu, b chen.ricky1982@gmail.com, c duck@santafe.edu

Abstract

Computing the reversal distances of signed permutations is an important topic in Bioinformatics. Recently, a new lower bound for the reversal distance was obtained via the plane permutation framework. This lower bound appears different from the existing lower bound obtained by Bafna and Pevzner through breakpoint graphs. In this paper, we prove that the two lower bounds are equal. Moreover, we confirm a related conjecture on skew-symmetric plane permutations, which can be restated as follows: let $p = (0, -1, -2, \ldots, n, n, n - 1, \ldots, 1)$ and let

$$\tilde{s} = (0, a_1, a_2, \dots a_n, -a_n, -a_{n-1}, \dots -a_1)$$

be any long cycle on the set $\{-n, -n+1, \ldots 0, 1, \ldots n\}$. Then, n and a_n are always in the same cycle of the product $p\tilde{s}$. Furthermore, we show the new lower bound via plane permutations can be interpreted as the topological genera of orientable surfaces associated to signed permutations.

Keywords: plane permutation; reversal; topological genus; fatgraph; breakpoint graph; involution

Mathematics Subject Classifications: 05A05; 92B05; 05C10

1 Introduction

In Bio-informatics, comparative study of genome sequences is very important to understand evolution. In particular, the problem of determining the minimum number of certain operations required to transform one of two given genome sequences into the other, is extensively studied. Combinatorially, this problem can be formulated as sorting a given permutation (or sequence) to the identity permutation, by certain operations, in a minimum number of steps, called the distance between the permutation to be sorted and

the identity, w.r.t. the operations chosen. The main operations studied are the transposition [1, 2, 5, 10], the block-interchange [5, 6] and the reversal [3, 13, 14], or some combination of them [4, 9].

In Chen and Reidys [7], a framework based on plane permutations was proposed in order to study these distance problems. As results, they give a general formulation for lower bounds of the transposition and block-interchange distance from which the existing lower bounds, obtained by Bafna and Pevzner [2], and Christie [6], can easily be derived. As to the reversal distance problem of signed permutations, the authors translate it into a block-interchange distance problem, by which they obtain a new lower bound. They observe that the new lower bound gives the exact reversal distance for most of the signed permutations.

This paper is mainly concerned with the reversal distances for signed permutations. A lower bound for the reversal distance was previously obtained in [3, 13] by studying breakpoint graphs associated to signed permutations. In [12], a topological framework for sorting signed permutations by reversals was proposed, where the topological genera of these orientable or non-orientable π -maps associated to signed permutations give a lower bound. These two existing lower bounds are equal [12]. However, it was not clear which is better, compared to the lower bound in [7]. In this paper, we will present another way of associating orientable fatgraphs to signed permutations whose genera give a lower bound for the reversal distance. Furthermore, we prove all these three approaches provide the same lower bound.

A brief outline of the paper is as follows: in Section 2, we review the approach to study the reversal distance via plane permutations proposed in [7], as well as a conjecture on skew-symmetric plane permutations to be confirmed in this paper. In Section 3, from each signed permutation, we construct a fatgraph and show that the genus of the fatgraph gives a lower bound for the reversal distance of the signed permutation. In Section 4, we will show the equivalence of the lower bound on the reversal distance obtained by Chen and Reidys [7] and the lower bound obtained by Bafna and Pevzner [2] through breakpoint graphs as well as the genus bound in Huang and Reidys [12].

2 Plane permutations and the reversal distance

Definition 2.1 (Plane permutation). A plane permutation on $[n] = \{1, 2, ..., n\}$ is a pair $\mathfrak{p} = (s, \pi)$ where $s = (s_i)_{i=0}^{n-1}$ is an *n*-cycle and π is an arbitrary permutation. The permutation $D_{\mathfrak{p}} = s \circ \pi^{-1}$ is called the diagonal of \mathfrak{p} .

Let $s = (s_0, s_1, \dots, s_{n-1})$. A plane permutation (s, π) can be represented by two aligned rows

$$(s,\pi) = \begin{pmatrix} s_0 & s_1 & \cdots & s_{n-2} & s_{n-1} \\ \pi(s_0) & \pi(s_1) & \cdots & \pi(s_{n-2}) & \pi(s_{n-1}) \end{pmatrix}$$
(1)

Note that $D_{\mathfrak{p}}$ is determined by the diagonal pairs, cyclically, in the two-line representation, i.e., $D_{\mathfrak{p}}(\pi(s_{i-1})) = s_i$ for 0 < i < n, and $D_{\mathfrak{p}}(\pi(s_{n-1})) = s_0$. In the following, by "the cycles of $\mathfrak{p} = (s, \pi)$ " we mean the cycles of π .

Given a plane permutation $\mathfrak{p} = (s, \pi)$ on [n] and a sequence h = (i, j, k, l), such that $i \leq j < k \leq l$ and $\{i, j, k, l\} \subset [n-1]$, let

$$s^h = (s_0, s_1, \dots, s_{i-1}, \underline{s_k, \dots, s_l}, s_{j+1}, \dots, s_{k-1}, s_i, \dots, s_j, s_{l+1}, \dots).$$

In other words s^h is the *n*-cycle obtained by transposing the blocks $[s_i, s_j]$ and $[s_k, s_l]$ in s. Let furthermore

$$\pi^h = D_{\mathfrak{p}}^{-1} \circ s^h.$$

This means, the derived plane permutation $\mathfrak{p}^h = (s^h, \pi^h)$ can be represented as

Note that the bottom row of the two-row representation of (s^h, π^h) is obtained by transposing the blocks $[\pi(s_{i-1}), \pi(s_{j-1})]$ and $[\pi(s_{k-1}), \pi(s_{l-1})]$ of the bottom row of (s, π) . In this way, each h not only determines a block-interchange on the s-sequence, but also it induces a block-interchange on the plane permutation (in two-row representation).

Definition 2.2. A signed permutation on [n] is a pair (a, w) where $a = a_1 a_2 \cdots a_n$ is a sequence on [n] while $w = w_1 w_2 \cdots w_n$ is a word of length n on the alphabet set $\{+, -\}$.

Usually, a signed permutation is represented by a single sequence $a_w = a_{w,1} a_{w,2} \cdots a_{w,n}$ where $a_{w,k} = w_k a_k$, i.e., each a_k carries a sign determined by w_k .

Given a signed permutation $a = a_1 a_2 \cdots a_{i-1} a_i a_{i+1} \cdots a_{j-1} a_j a_{j+1} \cdots a_n$ on [n], a reversal $\varrho_{i,j}$ acting on a will change a into

$$a' = \varrho_{i,j} \diamond a = a_1 a_2 \cdots a_{i-1} (-a_j) (-a_{j-1}) \cdots (-a_{i+1}) (-a_i) a_{j+1} \cdots a_n.$$

The reversal distance $d_r(a)$ of a signed permutation a on [n] is the minimum number of reversals needed to sort a into $e_n = 12 \cdots n$.

Example 2.1. The signed permutation -5, 1, -3, 2, 4 needs at least 4 steps to be sorted as illustrated below:

For a given signed permutation a, we associate to it the sequence s = s(a) as follows

$$s = s_0 s_1 s_2 \cdots s_{2n} = 0 a_1 a_2 \cdots a_n (-a_n) (-a_{n-1}) \cdots (-a_2) (-a_1).$$

In other words $s_0 = 0$ and $s_k = -s_{2n+1-k}$ for $1 \le k \le 2n$. Furthermore, such sequences will be referred to as skew-symmetric sequences since we have $s_k = -s_{2n+1-k}$. A sequence s is called negative if there exists $s_i < 0$ for some $1 \le i \le n$. Now we observe that the reversal distance of a is equal to the block-interchange distance of s into

$$e_n^{\sharp} = 012 \cdots n(-n)(-n+1) \cdots (-2)(-1),$$

where only certain block-interchanges are allowed. In other words, only block-interchanges h = (i, j, 2n + 1 - j, 2n + 1 - i) where $1 \le i \le j \le n$ are allowed. Hereafter, we will denote these particular block-interchanges on s as the reversals $\varrho_{i,j}$.

Let

$$\tilde{s} = (s) = (0, a_1, a_2, \dots, a_{n-1}, a_n, -a_n, -a_{n-1}, \dots, -a_2, -a_1),$$

 $p = (0, -1, -2, \dots, -n+1, -n, n, n-1, \dots, 2, 1).$

A plane permutation of the form (\tilde{s}, π) will be called skew-symmetric. Let $C(\pi)$ denote the number of cycles in the permutation π . Based on these definitions and notations, the new lower bound for the reversal distance of a obtained in [7] is

Theorem 2.3 (Chen and Reidys [7]).

$$d_r(a) \ge \frac{2n + 1 - C(p\tilde{s})}{2}.\tag{2}$$

The proof of the theorem relies on the observation that the increment of the number of cycles from \mathfrak{p} to \mathfrak{p}^h , for a given reversal h, is at most 2. Now start with $\mathfrak{p}=(\tilde{s},\pi_1)$ such that $D_{\mathfrak{p}}=p^{-1}$, and let us apply a sequence of reversals so that we arrive at $((e_n^{\natural}),\pi_2)$. Note that the starting plane permutation has $C(p(\tilde{s}))$ cycles, while the terminating plane permutation has $C(p(e_n^{\natural}))=2n+1$ cycles. Hence, at least $\frac{2n+1-C(p\tilde{s})}{2}$ reversals are needed for sorting s into e_n^{\natural} .

The effectiveness of the lower bound, when there exists a reversal increasing the number of cycles by exactly 2 (a 2-reversal for short), was analyzed in [7]. It was proved that there exists a 2-reversal for all skew-symmetric plane permutations but possibly two minor classes. One class is that of nonnegative permutations. For the other class, a 2-reversal exists if the following conjecture is true.

Conjecture 2.4. [7] Let $\mathfrak{p} = (\tilde{s}, \pi)$ be a skew-symmetric plane permutation on $[n]^{\pm} = \{-n, -n+1, \ldots, 0, 1, \ldots, n\}$, where $D_{\mathfrak{p}} = p^{-1}$. Then, n and s_n are in the same cycle of π .

3 Associating fatgraphs to signed permutations

Let G be a graph, with loops and multiple edges allowed. We call the ends of edges in G half-edges (or darts). An orientable fatgraph, also called *(combinatorial) map*, is a graph G with a specified cyclic order of half-edges around (i.e., incident to) each vertex of G. An orientable fatgraph can be viewed as a cell-complex, i.e., the underlying graph represents

the 1-skeleton while the (counterclockwise) cyclic order of half-edges indicate how to glue 2-cells on top of that to obtain a cell-complex. Accordingly, we can define the topological genus of the fatgraph to be the genus of the cell-complex. Hereafter, a fatgraph is alway orientable unless explicitly stated otherwise.

A fatgraph having n edges can be represented as a triple of permutations (α, β, γ) on [2n] where α is a fixed-point free involution and $\gamma = \alpha\beta$. This can be seen as follows: we label the half-edges of the fatgraph using the labels from the set [2n] so that each label appears exactly once. This induces two permutations α and β , where α is a fixed point free involution, whose cycles consist of the labels of the two half-edges of the same (untwisted) edge and β -cycles represent the counterclockwise cyclic arrangement of all half-edges incident to the same vertex. $\gamma = \alpha\beta$ -cycles are called the faces. The topological genus of a fatgraph (α, β, γ) satisfies

$$C(\beta) - C(\alpha) + C(\gamma) = 2 - 2q. \tag{3}$$

Conversely, from such a triple of permutations (α, β, γ) on [2n], a fatgraph can be constructed.

For the general case that α is not necessarily a fixed-point free involution, the pair (α, β, γ) satisfying $\gamma = \alpha\beta$ is called a hypermap.

Note a plane permutation $\mathfrak{p} = (s, \pi)$ can be interpreted as a hypermap $(D_{\mathfrak{p}}, \pi, s)$ [7, 8] and there is a connection between hypermaps and certain maps (called bipartite maps) observed by Walsh [16]. Motivated by this, we will associate an orientable fatgraph for a given signed permutation by transforming the skew-symmetric plane permutation for a signed permutation into a plane permutation whose diagonal is a fixed-point free involution. We next show that the topological genus of the fatgraph encoded by the plane permutation gives the lower bound for the reversal distance.

Given a plane permutation \mathfrak{p} on [n], w.l.o.g., we assume

$$\mathfrak{p} = (s,\pi) = \left(\begin{array}{cccc} 1 & 2 & \cdots & n-1 & n \\ \pi(1) & \pi(2) & \cdots & \pi(n-1) & \pi(n) \end{array}\right).$$

We obtain another plane permutation $(\hat{s}, \hat{\pi})$ on $[n] \cup [\bar{n}]$ having a fixed-point free involution diagonal, where $[\bar{n}] = \{\bar{1}, \bar{2}, \dots \bar{n}\}$, by the following construction:

step 1. put \bar{i} after i to obtain \hat{s} ;

step 2. we determine $\hat{\pi}$ as follows: firstly, when restricted to [n], $\hat{\pi}|_{[n]} = \pi$. Next, to guarantee the diagonal to be a fixed-point free involution, the image of \bar{i} (i.e., $\hat{\pi}(\bar{i})$) will be automatically determined. Namely, $\hat{\pi}(\bar{1}) = (\pi^{-1}(2))$, $\hat{\pi}(\bar{2}) = (\pi^{-1}(3))$, etc. In general, $\hat{\pi}(\bar{i}) = (\pi^{-1}(i+1))$ (in the sense of modulo n).

The following is an example to illustrate this construction, where n=6,

Lemma 3.1. When restricted to $[\bar{n}]$, $\hat{\pi}|_{[\bar{n}]}$ has the same number of cycles as $D_{\mathfrak{p}}$.

Proof. Note, identifying $\pi(i)$ and \bar{i} , $\hat{\pi}(\bar{i})$ and i+1, i.e., two elements on a same diagonalpair, will preserve cycle structure in the sense that

$$\hat{\pi}: \bar{i} \to \hat{\pi}(\bar{i}), \quad D_{\mathfrak{p}}: \pi(i) \to i+1.$$

Therefore, $\hat{\pi}|_{[\bar{n}]}$ has the same number of cycles as $D_{\mathfrak{p}}$.

For the example given above, we can check that $D_{\mathfrak{p}} = (153)(264)$ has two cycles, while $\hat{\pi}|_{\bar{5}} = (153)(264)$ has also two cycles.

Applying above construction to the associated skew-symmetric plane permutation $\mathfrak{p}=(\tilde{s},\pi)$ to a signed permutation a, where $D_{\mathfrak{p}}=p^{-1}$, we can associate a fatgraph \mathcal{F}_a to a. Let $g(\mathcal{F}_a)$ denote the genus of the fatgraph \mathcal{F}_a .

Theorem 3.1. The reversal distance $d_r(a)$ for a satisfies

$$d_r(a) \ge g(\mathcal{F}_a). \tag{4}$$

Proof. Let $\mathfrak{p}=(s,\pi)$ be the corresponding skew-symmetric plane permutation of a. Then, in \mathcal{F}_a , there are $C(\pi) + C(D_{\mathfrak{p}}) = C(ps) + 1$ vertices, 2n + 1 edges and one face. Thus,

$$2 - 2g(\mathcal{F}_a) = (C(ps) + 1) - (2n + 1) + 1$$

so that

$$g(\mathcal{F}_a) = \frac{2n + 1 - C(ps)}{2}.$$

Comparing with Theorem 2.3 completes the proof.

Example 3.1. Consider the signed permutation in Example 2.1. Its associated skewsymmetric plane permutation is

Applying the transformation above, we obtain

Applying the transformation above, we obtain
$$\begin{pmatrix} 0 & \bar{0} & -5 & \overline{-5} & 1 & \cdots & \bar{2} & 4 & \bar{4} & -4 & \overline{-4} & -2 & \overline{-2} & \cdots & -1 & \overline{-1} & 5 & \bar{5} \\ 5 & \bar{4} & 0 & \overline{-3} & -4 & \cdots & \overline{-1} & -5 & \bar{1} & -3 & \bar{3} & 2 & \bar{2} & \cdots & 4 & \bar{0} & -1 & \overline{-5} \end{pmatrix}.$$

and its associated fatgraph as illustrated in Figure 1. It can be seen that the fatgraph has 4 vertices, 11 edges (ribbons) and one face. So, the genus of the fatgraph is 4 by Euler's characteristic formula.

Compared to the fatgraphs associated to signed permutations in [12], the associated fatgraphs there are non-orientable in most cases while the associated fatgraphs in this paper are always orientable. The relation between the two fatgraphs (via these two different approaches) associated to the same signed permutation remains unclear and would be an interesting problem to investigate.

Remark. For the lower bounds for the transposition distances and the block-interchange distances of (unsigned) permutations, we can have analogous topological interpretations by transforming the associated plane permutations [7] into fatgraphs in the same way as we did here for the reversal distances of signed permutations.




Figure 1: The fatgraph associated to -5, 1, -3, 2, 4.

4 Comparison with the Bafna-Pevzner lower bound

In [12], it was shown that the topological-genus lower bound there and the lower bound given by breakpoint graphs in [3, 13] are equal. In this section, we will show that our new topological-genus lower bound equals the lower bound given by breakpoint graphs in [3, 13] as well.

The breakpoint graph for a given signed permutation $a = a_1 a_2 \cdots a_n$ on [n] can be obtained as follows: replace a_i with $(-a_i)a_i$, and add 0 at the beginning of the obtained sequence while adding -(n+1) at the end. In this way we obtain a sequence $b = b_0 b_1 b_2 \cdots b_{2n} b_{2n+1}$ on $[n]^{\pm} \cup \{-n-1\}$, where $b_0 = 0$, $b_{2n+1} = -n-1$. Draw a black edge between b_{2i} and b_{2i+1} , as well as a grey edge between i and -(i+1) for $0 \le i \le n$. The obtained graph is the breakpoint graph, BG(a), of a. Note that each vertex in BG(a) has degree two hence the graph can be decomposed into disjoint cycles. Denote the number of cycles in BG(a) by $C_{BG}(a)$. Then, the lower bound of the reversal distance via the breakpoint graph is

$$d_r(a) \ge n + 1 - C_{BG}(a). \tag{5}$$

In [7], an algebraic, equivalent formulation, of the breakpoint graph approach was obtained as follows: let θ_1 , θ_2 be the two involutions (without fixed points) determined by the black edges and grey edges in the breakpoint graph, respectively, i.e.

$$\theta_1 = (b_0, b_1)(b_2, b_3) \cdots (b_{2n}, b_{2n+1}),$$

 $\theta_2 = (0, -1)(1, -2) \cdots (n, -n-1).$

Then, we have

Proposition 4.1 (Chen and Reidys [7]).

$$d_r(a) \ge \frac{2n + 2 - C(\theta_1 \theta_2)}{2}.\tag{6}$$

To show the equivalence between our new topological-genus bound (equivalently, the lower bound in eq. (2)) and the lower bound given by the breakpoint graphs, it suffices to show, for any signed permutation a, there holds

$$C(p_r\tilde{s}) = C(\theta_1\theta_2) - 1.$$

Definition 4.1. Let σ be a permutation on the set $[n]^{\pm}$. We associate to σ the matrix $A_{\sigma} = [a_{ij}],$

$$a_{ij} = \begin{cases} 1 & \text{if } i = \sigma(j) \\ 0 & \text{if } i \neq \sigma(j) \end{cases}$$

where $i, j \in [n]^{\pm}$, index rows and columns, following the order $-n, -n+1, \ldots -1, 0, 1, \ldots, n$, respectively. We call A_{σ} the permutation matrix associated to σ and denote this by $\sigma \sim A_{\sigma}$.

Recall that $\sigma \sim A_{\sigma} \Leftrightarrow \sigma^{-1} \sim A_{\sigma}^{T}$, that if $\tau \sim A_{\tau}$ then $\sigma \circ \tau = A_{\sigma}A_{\tau}$, and that for id, the identity permutation, we have that $id \sim A_{id} = I_{2n+1}$ (i.e. the identity matrix).

Lemma 4.1. Let

$$p = (0, -1, -2, \dots - n, n, n - 1, \dots 1) \sim A_p = P,$$

$$\tilde{s} = (0, a_1, a_2, \dots a_n, -a_n, -a_{n-1}, \dots - a_1) \sim A_{\tilde{s}} = S.$$

Let $R = [r_{ij}]$ be the $(2n + 1) \times (2n + 1)$ unitary anti-diagonal matrix, also known as the exchange matrix. Namely,

$$r_{ij} = \begin{cases} 1 & \text{if } j = 2n - i + 2 \\ 0 & \text{if } j \neq 2n - i + 2 \end{cases}$$

where $i, j \in [2n+1]$ index rows and columns of R respectively. Then

$$PS = (PR)(RS), (7)$$

Proof. It suffices to check that $R = R^T$. Then, since R is a permutation matrix we have $R^2 = RR^T = I_{2n+1}$.

We next show that both permutations PR and RS are involutions with a unique fixed point.

Lemma 4.2. The permutation corresponding to the matrix PR is the involution

$$p_{invo} = (-n, n-1)(-n+1, n-2) \cdots (-1, 0)(n).$$

Proof. We compute

$$PR = \begin{bmatrix} 0 & 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \dots & 1 & 0 \\ 0 & 0 & 0 & 0 & \dots & 0 & 1 \\ 1 & 0 & 0 & 0 & \dots & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 & 0 & \dots & 0 & 1 \\ 0 & 0 & 0 & 0 & \dots & 1 & 0 \\ 0 & 0 & 0 & 0 & \dots & 1 & 0 \\ 0 & 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 1 & 0 & 0 & \dots & 0 & 0 \\ 1 & 0 & 0 & 0 & \dots & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & \dots & 0 & 1 & 0 \\ 0 & 0 & 0 & \dots & 0 & 1 & 0 \\ 0 & 0 & 0 & \dots & 1 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 1 & 0 & \dots & 0 & 0 & 0 \\ 1 & 0 & 0 & \dots & 0 & 0 & 0 \end{bmatrix}$$

Converting the matrix into a permutation completes the proof.

Lemma 4.3. The permutation corresponding to the matrix RS is the involution

$$s_{invo} = (0, -s_1)(s_1, -s_2) \cdots (s_{n-1}, -s_n)(s_n)$$

Proof. It is easy to check that left multiplication by the exchange matrix R reverses the order on the rows of the multiplied matrix $S = [a_{ij}], i, j \in [n]^{\pm}$. Since $a_{i,j} = 1 \iff i = s(j)$, when multiplied by the matrix R, the row indexed by i, in the matrix S, is sent to the row indexed by -i. This is due to the symmetry of $[n]^{\pm}$. However, this in turn means that in the permutation $s_{invo} \sim RS$, $j \xrightarrow{s_{invo}} -i$, $\forall j \in [n]^{\pm}$ such that s(j) = i. But now by virtue of the structure of the cycle $s = (0, s_1, s_2, \ldots s_n, -s_n, -s_{n-1}, \ldots -s_1)$, we have

$$s_{invo} = (0, -s_1)(s_1, -s_2) \cdots (s_{n-1}, -s_n)(s_n)$$

and the lemma follows.

We now know that both PR and RS are matrices corresponding to involutions, each with a unique fixed point n and s_n , respectively. We proceed to prove some results regarding the product of two involutions.

Lemma 4.4. Let σ_1 , σ_2 be two fixed-point free involutions on a set T, with |T| = 2n for $n \ge 1$. Then, there does not exist $k \ge 1$ such that $\sigma_1(\sigma_2\sigma_1)^k(x) = x$.

Proof. Assume by contradiction that there exists a $k \ge 1$ such that $\sigma_1(\sigma_2\sigma_1)^k(x) = x$. Then, since σ_1 is an involution, we have $\sigma_1(x) = (\sigma_2\sigma_1)^k(x)$.

Now if k = 1, we have $\sigma_1(x) = \sigma_2 \sigma_1(x)$, implying that $\sigma_1(x)$ is a fixed point of σ_2 , which is a contradiction.

If k > 1, since σ_2 is an involution, we have

$$\sigma_2\sigma_1(x) = (\sigma_1\sigma_2)^{k-1}(\sigma_1(x)) = \sigma_1(\sigma_2\sigma_1)^{k-1}(x).$$

Now, if k-1=1, we have that $\sigma_2\sigma_1(x)$ has to be a fixed point of σ_1 , which is again a contradiction. Otherwise, we can set $y=\sigma_2\sigma_1(x)$ and obtain $y=\sigma_1(\sigma_2\sigma_1)^{k-1}(y)$, and iterate the previous argument. In this way, we eventually obtain a fixed point, either for σ_1 or σ_2 , which contradicts our assumption, hence the lemma follows.

Lemma 4.5. Let σ_1 , σ_2 be two involutions on a set T such that each of them has a unique fixed point, a and b respectively. Then $\sigma_2\sigma_1$ has a cycle which contains both a and b.

Proof. There is nothing to prove if a = b, so we will assume $a \neq b$ in the following. Let now

$$\sigma_1 = (p_1, p_2)(p_3, p_4) \cdots (p_{2t-1}, p_{2t})(a),$$

$$\sigma_2 = (q_1, q_2)(q_3, q_4) \cdots (q_{2t-1}, q_{2t})(b),$$

$$\sigma'_1 = (p_1, p_2)(p_3, p_4) \cdots (p_{2t-1}, p_{2t})(a, x),$$

$$\sigma'_2 = (q_1, q_2)(q_3, q_4) \cdots (q_{2t-1}, q_{2t})(b, x),$$

where σ'_1 and σ'_2 are involutions on $T \cup \{x\}$, where $x \notin T$. We now compare the following two iterations:

$$a \to \sigma_{1}(a) \to \sigma_{2}\sigma_{1}(a) \to \sigma_{1}\sigma_{2}\sigma_{1}(a) \to (\sigma_{2}\sigma_{1})^{2}(a) \cdots (\sigma_{2}\sigma_{1})^{k_{1}}(a) = a,$$

$$x \to \sigma'_{1}(x) \to \sigma'_{2}\sigma'_{1}(x) \to \sigma'_{1}\sigma'_{2}\sigma'_{1}(x) \to (\sigma'_{2}\sigma'_{1})^{2}(x) \cdots \sigma'_{1}(\sigma'_{2}\sigma'_{1})^{k_{2}-1}(x) \to (\sigma'_{2}\sigma'_{1})^{k_{2}}(x) = x.$$

Note that $\sigma_1(a) = \sigma'_1(x) = a$, and that σ_1 and σ'_1 , excluding x, differ only at the image of a. Similarly, σ_2 and σ'_2 , excluding x, differ only at the image of b. Thus, the iterations starting with $\sigma_1(a)$ and $\sigma'_1(x)$ agree with each other until reaching a or b.

Claim 1. The iteration starting with $x \to \sigma_1'(x) = a$ will not reach a for a second time. This is because, otherwise, there must exists some $k \ge 1$, such that $\sigma_1'(\sigma_2'\sigma_1')^k(x) = a$ or $(\sigma_2'\sigma_1')^k(x) = a$. The former case can not happen, otherwise $(\sigma_2'\sigma_1')^k(x) = x$, which will close the iteration instead of continuing to a. By Lemma 4.4, the latter case, $(\sigma_2'\sigma_1')^k(x) = \sigma_2'(\sigma_1'\sigma_2')^{k-1}(a) = a$ cannot happen either. Hence, Claim 1 follows.

Claim 2. The iteration starting with $\sigma'_1(x) = a$ will reach b at least once. This is obvious since $\sigma'_1(\sigma'_2\sigma'_1)^{k_2-1}(x) = b$.

Now consider the first time the iteration starting with $\sigma'_1(x)$ reaches b. This must also be the first time the iteration starting with $\sigma_1(a)$ reaches b. There are two cases: either $(\sigma_2\sigma_1)^k(a) = b$ or $\sigma_1(\sigma_2\sigma_1)^k(a) = b$ for some $k \geq 1$. For the former case, we already have a and b as being in the same cycle of $\sigma_2\sigma_1$;

For the latter case, we have $(\sigma_2\sigma_1)^{k+1}(a) = \sigma_2(b) = b$, which also implies that a and b are in the same cycle of $\sigma_2\sigma_1$. This completes the proof.

Remark 4.2. Lemma 4.5 can be alternatively proved in the following approach: first, we show that there is a way to assign signs '+' and '-' to elements in the set T such that in both σ_1 and σ_2 , every 2-cycle has exactly one '+' element and one '-' element while a and b are positive, see a dihedral group action argument as in the Intersection-Theorem [15]. Then, we apply the Garsia-Milne Involution Principle [11] to explain that a and b are in the same cycle of the product $\sigma_1\sigma_2$.

Lemma 4.6. Let σ_1 , σ_2 , σ'_1 , σ'_2 be defined as in the proof of Lemma 4.5. Then,

$$C(\sigma_1'\sigma_2') - 1 = C(\sigma_1\sigma_2). \tag{8}$$

Proof. Following the discussion in the proof of Lemma 4.5, any cycle not containing a or b or x of $\sigma'_1\sigma'_2$ is also a cycle of $\sigma_1\sigma_2$. Thus, the difference $C(\sigma'_1\sigma'_2) - C(\sigma_1\sigma_2)$ equals the difference of the number of cycles containing a, b and x in $\sigma'_1\sigma'_2$ and the number of cycles containing a and b in $\sigma_1\sigma_2$.

On the one hand, we have already shown that a and b are in the same cycle of $\sigma_1\sigma_2$. On the other hand, it is clear that $\sigma'_1\sigma'_2(a) = b$; and by Claim 1 in the proof of Lemma 4.5, x and a are not in the same cycle of $\sigma'_1\sigma'_2$. Hence, the difference is exactly 1, completing the proof.

Theorem 4.3. For any given signed permutation a,

$$C(p\tilde{s}) = C(\theta_1 \theta_2) - 1. \tag{9}$$

П

Proof. By construction, the relation between the pair \tilde{s}_{invo} , p_{invo} and θ_1 , θ_2 is exactly the same as the pair σ_1 , σ_2 and σ'_1 , σ'_2 . Applying Lemma 4.6, we have

$$C(\theta_1\theta_2) - 1 = C(p_{invo}\tilde{s}_{invo}) = C(p\tilde{s}),$$

completing the proof.

Accordingly, the lower bound on the reversal distance obtained by Chen and Reidys [7] (and the topological-genus bound in the last section) is equivalent to the lower bound obtained by Bafna and Pevzner through breakpoint graphs [3, 13].

The discussion above has also implied Conjecture 2.4. Note that the conjecture can be equivalently formulated as follows: Let

$$p = (0, -1, -2, \dots - n, n, n - 1, \dots 1)$$
 and $s = (0, s_1, s_2, \dots s_n, -s_n, -s_{n-1}, \dots - s_1),$

both being long cycles on the set $[n]^{\pm}$, and let $\pi = p \circ s$. Then, the conjecture states that n and s_n are in the same cycle of π .

Corollary 4.1. The elements n and s_n are in the same cycle of the product $p \circ s$.

Proof. Applying Lemma 4.5 to the involution decomposition

$$p \circ s = p_{invo} \circ s_{invo}$$

the theorem follows.

References

- [1] L. Bulteau, G. Fertin, I. Rusu, Sorting by transpositions is difficult, SIAM J. Discrete Math. 26(3) (2012), 1148–1180.
- [2] V. Bafna, P. Pevzner, Sorting permutations by transpositions, Proc. 6th Symposium on Discrete Algorithms, ACM-SIAM, (1995), pp. 614-621.

- [3] V. Bafna, P. A. Pevzner, Genome rearragements and sorting by reversals, SIAM Journal on Computing, 25(2) (1996), 272–289.
- [4] M. Bader, E. Ohlebusch, Sorting by Weighted Reversals, Transpositions, and Inverted Transpositions, Journal of Computational Biology 14(5) (2007), 615–636.
- [5] D.A. Christie, Genome Rearrangement Problems, PhD. thesis, Glasgow University, 1998.
- [6] D.A. Christie, Sorting permutations by block-interchanges, Inf. Process. Lett., 60 (1996), 165–169.
- [7] R.X.F. Chen, C.M. Reidys, Plane permutations and applications to a result of Zagier–Stanley and distances of permutations, SIAM J. Discrete Math. 30(3) (2016), 1660–1684.
- [8] R.X.F. Chen, C.M. Reidys, On the local genus distribution of graph embeddings, J. Combin. Math. Combin. Comput. 101 (2017), 157–173.
- [9] Z. Dias, J. Meidanis, Genome rearrangements distance by fusion, fission, and transposition is easy, Proc. SPIRE 2001, pp. 250–253.
- [10] I. Elias, T. Hartman, A 1.375-approximation algorithm for sorting by transpositions, In Proc. of the 5th International Workshop on Algorithms in Bioinformatics (WABI05), volume 3692 of Lecture Notes in Computer Science, pp. 204–214.
- [11] A. Garsia and S. Milne, Method for constructing bijections for classical partition identities, Proc. Nat. Acad. Sci. U.S.A. 78 (1981), 2026–2028.
- [12] F.W.D. Huang, C.M. Reidys, A topological framework for signed permutations, arXiv:1410.4706.
- [13] S. Hannenhalli, P. Pevzner, To cut or not to cut (applications of comparative physical maps in molecular evolution), Proceedings of the seventh annual ACM-SIAM symposium on Discrete algorithms (1996), 304–313.
- [14] S. Hannenhalli, P. Pevzner, Transforming cabbage into turnip: Polynomial algorithm for sorting signed permutations by reversals, Journal of the ACM 46 (1999) 1–27.
- [15] C. M. Reidys, P. F. Stadler, P. Schuster, Generic properties of combinatory maps: Neural networks of RNA secondary structures, Bull. Math. Biol. 59 (1997), 339–397.
- [16] T.R.S. Walsh, Hypermaps versus bipartite maps, J. Comb. Theory, Ser. B 18 (1975), 155–163.