

DETERMINANTAL VARIETY AND NORMAL EMBEDDING

KARIN U. KATZ, MIKHAIL G. KATZ, DMITRY KERNER,
AND YEVGENY LIOKUMOVICH

ABSTRACT. The space GL_n^+ of matrices of positive determinant inherits an extrinsic metric space structure from \mathbb{R}^{n^2} . On the other hand, taking the infimum of the lengths of all paths connecting a pair of points in GL_n^+ gives an intrinsic metric. We prove bilipschitz equivalence between intrinsic and extrinsic metrics on GL_n^+ , exploiting the conical structure of the stratification of the space of $n \times n$ matrices by rank.

1. INTRODUCTION

Consider the group $GL^+(n, \mathbb{R})$ of $n \times n$ matrices of positive determinant. It is an open submanifold of the space \mathbb{R}^{n^2} of $n \times n$ matrices. Here $GL^+(n, \mathbb{R})$ carries two metrics: its extrinsic ambient Euclidean metric with distance function d_{ext} , and the induced intrinsic metric (i.e. least length of path) with distance function d_{int} . Our main result is the following.

Theorem 1.1. *There is a constant $C = C(n)$ such that $d_{int} < Cd_{ext}$.*

Thus the determinantal variety is *normally embedded*; in other words, the intrinsic and extrinsic metrics are *bilipschitz equivalent*. We prove Theorem 1.1 first for $n = 2, 3$ and then for general n . The proof uses in an essential way the conical structure of the stratification of the space \mathbb{R}^{n^2} of $n \times n$ matrices by rank. Indeed, the extrinsic and intrinsic metrics on a set as simple as $\{(x, y) : x^2 - y^3 > 0\}$ are clearly inequivalent, due to the fact that the curve $x^2 - y^3 = 0$ has a cusp.

Bilipschitz equivalence has been studied by a number of authors; see e.g., [1], [2], [4], [7]. For a study of the Lipschitz condition in an infinitesimal context see [3], [6].

2010 *Mathematics Subject Classification.* 53C23, 58A35, 32S60.

Key words and phrases. Determinantal variety, intrinsic metric, bilipschitz equivalence, conical stratification.

2. SOLUTION IN DIMENSION 2

For 2×2 matrices with coordinates given by the entries (a_{ij}) , the condition $a_{11}a_{22} - a_{12}a_{21} = 0$ for the determinantal variety in the space of matrices translates into

$$(a_{11} + a_{22})^2 - (a_{11} - a_{22})^2 - (a_{12} + a_{21})^2 + (a_{12} - a_{21})^2 = 0$$

or to simplify notation

$$x^2 + y^2 = z^2 + w^2. \quad (2.1)$$

Definition 2.1. Let $X_2 = \{A \in \mathbb{R}^{2^2} : \det(A) = 0\}$ denote the determinantal variety.

Lemma 2.2. *The intersection $X_2 \cap S^3$ with the unit sphere $S^3 \subseteq \mathbb{R}^{2^2}$ is a flat 2-torus, namely the Clifford torus $T \subseteq S^3$.*

Indeed, by (2.1) the complement $S^3 \setminus T$ is a union of two solid tori, consisting respectively of matrices of positive and negative determinant. Note that X_2 is a linear cone over $T \subseteq \mathbb{R}^{2^2}$.

Lemma 2.3. *The metrics d_{int} and d_{ext} on $T \subseteq \mathbb{R}^{2^2}$ are bilipschitz equivalent.*

Proof. By rescaling, we can assume that the furthest of the two points is at unit distance from the origin. By compactness, the only problem for a pair of points p, q can arise when the points collide, i.e., $d(p, q)$ tends to 0. By the smoothness of the Clifford torus, one has $\frac{d_{int}(p, q)}{d_{ext}(p, q)} \rightarrow 1$ as $d(p, q) \rightarrow 0$. \square

Lemma 2.4. *Bilipschitz equivalence holds for the intrinsic and the extrinsic metrics on X_2 .*

Proof. Let $p, q \in X_2$. If the apex $O \in X_2$ is one of the points p, q then the intrinsic and the Euclidean distances between them coincide by linearity of the cone.

Thus we can assume that $p \neq O$ and $q \neq O$. To connect a pair of points p, q , at different levels in the cone by a path lying in the cone, assume without loss of generality that p is further than q from the apex of the cone. We slide p along the ray toward the apex until it reaches the level of q , and then connect it to q by a shortest path contained in that level. The length of the combined path is clearly bilipschitz with the extrinsic distance in the ambient \mathbb{R}^{2^2} . \square

Proposition 2.5. *The intrinsic and the ambient metrics on the manifold of 2×2 matrices of positive determinant are bilipschitz equivalent.*

Proof. The closure of $GL^+(2, \mathbb{R})$ is a linear cone over the solid torus. Meanwhile, $GL^+(2, \mathbb{R})$ itself is the cone without the apex over the *interior* of the solid torus. We consider a straight line path in \mathbb{R}^{2^2} connecting a pair of points in $GL^+(2, \mathbb{R})$. The length of this path is the extrinsic distance by definition.

The path does not necessarily lie entirely inside $GL^+(2, \mathbb{R})$. As the path is straight and the determinantal variety X_2 is algebraic, the path splits into a finite number of segments satisfying the following:

- (1) the interior of each segment lies fully either inside $GL^+(2, \mathbb{R})$ or inside the component $GL^-(2, \mathbb{R})$;
- (2) the endpoints are in X_2 .

Applying Lemma 2.4, we replace every segment that lies in the component $GL^-(2, \mathbb{R})$ by an arc that lies in X_2 .

Finally, we push out the arc in X_2 into $GL^+(2, \mathbb{R})$, i.e., replace it by a nearby arc “just inside” $GL^+(2, \mathbb{R})$. We will explain the push-out procedure in detail since a similar argument will be used in the general case.

The vertex $O \in X_2$ is the unique singular point of the cone, i.e., the complement $X_2 \setminus \{O\}$ is a smooth manifold. Hence a path in X_2 disjoint from O can be pushed out infinitesimally into $GL^+(2, \mathbb{R})$ by following the normal direction, without significantly affecting its length.

Suppose a path joining $P, Q \in X_2$ passes through O . Using the linear structure of the cone, the path can be replaced by the shorter path given by the union of the straight line segments $PO \cup OQ \subseteq X_2$. Next, P and Q can be replaced by nearby points $P', Q' \in GL^+(2, \mathbb{R})$. We now form a path $P'O \cup OQ'$ lying entirely within $GL^+(2, \mathbb{R})$ except for the single point O .

Let $p \in OP'$ be the point of intersection of the segment with a sphere $S(O, \epsilon)$ of small radius $\epsilon > 0$ centered at O , and similarly for point $q \in OQ'$. The intersection $S(O, \epsilon) \cap GL^+(2, \mathbb{R})$ is an open solid torus and therefore connected. Hence there is a short path

$$\gamma \subseteq S(O, \epsilon) \cap GL^+(2, \mathbb{R})$$

joining p to q . The resulting path $P'p \cup \gamma \cup qQ'$ is only slightly longer than the original path joining P and Q . This completes the proof of the bilipschitz property for $GL^+(2, \mathbb{R})$. \square

3. SINGULARITIES IN DIMENSION 3

For 3×3 matrices, the singular locus of the determinantal variety $X_3 = \{A \in \mathbb{R}^{3^2} : \det A = 0\}$ consists of matrices of rank ≤ 1 . The

variety X_3 can be stratified as follows:

$$X_3 = X_{3,0} \cup X_{3,1} \cup X_{3,2}.$$

Here the stratum $X_{3,i}$ consists of matrices of rank i . The stratum $X_{3,1}$ lies in the closure of $X_{3,2}$, while $X_{3,0}$ lies in the closure of $X_{3,1}$. Each $X_{3,i}$ is smooth. Here $X_{3,2}$ is of codimension 1 in the space of matrices, while $X_{3,1}$ is of codimension 4 in the space of matrices, and $X_{3,0}$ is a single point.

The closure $\overline{X}_{3,2}$ is a cone on a smooth manifold away from $X_{3,1}$. Thus the only potential obstruction to bilipschitz equivalence is the singularity of $\overline{X}_{3,2}$ along $\overline{X}_{3,1}$, which we now analyze.

Lemma 3.1. *The 5-dimensional closure $\overline{X}_{3,1}$ is a linear cone over the smooth compact 4-manifold $(S^2 \times S^2)/\{\pm 1\}$.*

Proof. Here $S^2 \times S^2/\{\pm 1\}$ parametrizes the intersection of $X_{3,1}$ with the unit 8-sphere in the space of matrices. The manifold $S^2 \times S^2$ can be parametrized by pairs of unit column vectors $v, w \in S^2 \subseteq \mathbb{R}^3$ producing a rank 1 matrix

$$v {}^t w \in X_{3,1}.$$

The element -1 acts simultaneously on both factors of $S^2 \times S^2$ by the antipodal map. \square

Fix an element $x \in X_{3,1}$. We would like to understand the bilipschitz property for X_3 in a neighborhood of x . Exploiting the action of $GL(3, \mathbb{R}) \times GL(3, \mathbb{R})$ on $X_{3,1}$ by right and left multiplication, we may assume that

$$x = 1 \oplus \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}.$$

Definition 3.2. Let

$$L_x = \{1 \oplus A : A \in \mathbb{R}^{2^2}\} \subseteq \mathbb{R}^{3^2}$$

be a transverse slice to $X_{3,1} \subseteq \mathbb{R}^{3^2}$ at x , so that

$$L_x \cap \overline{X}_{3,2} = \{1 \oplus A \in L_x : \det(A) = 0\}. \quad (3.1)$$

Note that of course transversality is weaker than orthogonality. The ambient Euclidean metric on \mathbb{R}^9 plays no role here.

Lemma 3.3. *A transverse slice for $X_{3,1} \subseteq \overline{X}_{3,2}$ in local coordinates is a linear cone over a Clifford torus.*

Proof. The defining equation (3.1) of the slice is $\det(A) = 0$. By Lemma 2.2 this is a cone over a torus. \square

Now let $U_T \subseteq \mathbb{R}^5$ be the unit ball. Choose open sets $U_N \subseteq L_x$ and $U \subseteq \mathbb{R}^{3^2}$, each containing x , and a bilipschitz diffeomorphism

$$\phi: U \rightarrow U_T \times U_N \quad (3.2)$$

such that $\phi(U \cap \overline{X}_{3,1}) = U_T \times \{x\}$ and $\phi(U \cap \overline{X}_{3,2}) = U_T \times (U_N \cap \overline{X}_{3,2})$. This can be done using the $GL(3, \mathbb{R}) \times GL(3, \mathbb{R})$ action, as explained in [5, p. 138]. Hence Lemmas 3.3 and 2.4 imply that the intrinsic and extrinsic metrics on $U \cap \overline{X}_{3,2}$ are bilipschitz equivalent.

Corollary 3.4. *The intrinsic and extrinsic metrics on $\overline{X}_{3,2} = X_3$ are bilipschitz equivalent.*

This follows from Lemma 3.1 by a compactness argument.

Theorem 3.5. *The intrinsic and extrinsic metrics on $GL^+(3, \mathbb{R})$ are bilipschitz equivalent.*

Proof. The extrinsic distance between a pair of matrices of positive determinant is the length of the straight line path in \mathbb{R}^{3^2} joining them. We partition the path into finitely many segments, where the interior of each segment lies entirely in a connected component of $GL(3, \mathbb{R})$ while the endpoints are in $X_3 \subseteq \mathbb{R}^{3^2}$. Then we apply Corollary 3.4 to replace each segment belonging to the component $GL^-(3, \mathbb{R})$ by an arc in X_3 . If the arc in X_3 lies in the smooth part $X_{3,2} \subseteq X_3$ then it can be pushed out into $GL^+(3, \mathbb{R})$ by a small deformation in the normal direction, as in Section 2.

Unlike the case of Proposition 2.5, the determinantal variety is not a cone on a manifold, so that an additional argument is required to push the arc out of X_3 and into GL_3^+ while retaining bilipschitz control.

If an arc in X_3 joining points P, Q passes through the apex $O \in X_3$ then it can be replaced by the union $PO \cup OQ$ and pushed out into $GL^+(3, \mathbb{R})$ as in the proof of Proposition 2.5.

Otherwise we exploit the local product structure on $X_3 \setminus \{O\}$ as in (3.2). A path in $X = X_3$ that dips into the singular locus $X_{3,1} \subseteq X$ can be handled as follows. Given a path $\gamma: [0, 1] \rightarrow X$, let $a \in [0, 1]$ be the least parameter value t such that $\gamma(t)$ is contained in the singular locus $X_{3,1} \subseteq X$, and $b \in [0, 1]$ the greatest such value.

Step 1. Consider the restriction of the path γ to $[a, b] \subseteq [0, 1]$. We replace it by a path that lies entirely in the singular locus $X_{3,1}$. Section 4 proves that $X_{3,1}$ is embedded in X in a bilipschitz fashion. Hence this replacement can be performed in a bilipschitz-controlled way.

Step 2. Once the path $\gamma([a, b])$ is in the singular locus, we exploit a local trivialisation to push it in a constant direction as follows.

Choose $\delta > 0$ sufficiently small to be specified later. Then $\gamma(a - \delta)$ and $\gamma(b + \delta)$ are in the smooth part $X_{3,2} \subseteq X$. The path can easily be shortened to a smooth one, still denoted γ . Over a sufficiently small neighborhood of the smooth path, we can choose a smooth (and in particular bilipschitz) trivialisation of X over the path $\gamma([a, b])$. Let

$$I_{0,1}(t) = I_{0,1} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad (3.3)$$

be a constant section of the bundle over the path. Relative to the trivialisation of the bundle we can form a new path

$$\bar{\gamma}_\epsilon(t) = \gamma(t) + \epsilon I_{0,1}(t)$$

which pushes $\gamma([a, b])$ in the constant direction (3.3) for each t . By construction, the new path is contained in the nonsingular part $X_{3,2}$ of the determinantal variety, where $\epsilon > 0$ is chosen small enough so that the length of the path stays close to the original length of the path in $X_{3,1}$.

Step 3. It remains to check that the path can be patched up with the value of the path γ at the parameter values $a - \delta$ and $b + \delta$. Since the determinantal variety in the fiber is a cone over a connected space, the two values can be connected by an arbitrarily short path, provided ϵ and δ are chosen small enough.

Step 4. Once the path lies in the nonsingular part $X_{3,2} \subseteq X$ of the determinantal variety, it can be pushed out into $GL^+(3, \mathbb{R})$ by following the normal direction as in the case $n = 2$. \square

The higher dimensional case is treated inductively in Section 5.

4. THE GENERAL DETERMINANTAL VARIETY

Let $X_{n,k} \subseteq \mathbb{R}^{n^2}$ be the stratum of rank k matrices. Thus X_n is the closure of $X_{n,n-1}$ and each $X_{n,k}$ is a smooth connected manifold of dimension $n^2 - (n-k)^2$. For each $x \in X_{n,k}$ choose an open set $U_x \subseteq \mathbb{R}^{n^2}$ containing x , a metric ball $U_x^T \subseteq \mathbb{R}^{n^2 - (n-k)^2}$, an open set $U_x^N \subseteq \mathbb{R}^{(n-k)^2}$ containing 0, and a bilipschitz diffeomorphism $\phi = \phi_x: U_x \rightarrow U_x^T \times U_x^N$, such that

$$\phi(U_x \cap X_{n,k}) = U_x^T \times \{0\}, \quad \phi(U_x \cap X_n) = U_x^T \times (U_x^N \cap X_{n-k}). \quad (4.1)$$

See [5, p. 138] for the construction of the diffeomorphism ϕ_x . To ensure the bilipschitz property, it suffices to shrink slightly U_x , U_x^N and U_x^T .

Assume by induction that we have proved the bilipschitz equivalence of the intrinsic and extrinsic metrics on X_ℓ for $\ell < n$. Generalizing Lemma 3.1 to dimension n , we see $\bar{X}_{n,1}$ is a linear cone over a compact

smooth submanifold. So, the intrinsic and extrinsic metrics on $\overline{X}_{n,1}$ are bilipschitz equivalent. For each $x \in X_{n,1}$, the bilipschitz diffeomorphism ϕ_x and the induction hypothesis for X_{n-1} show the intrinsic and extrinsic metrics are bilipschitz equivalent on $U_x \cap X_n$. Then, a compactness argument shows the bilipschitz property holds for a sufficiently small open neighborhood $U_\epsilon(\overline{X}_{n,1})$ of $\overline{X}_{n,1}$ in X_n . More precisely, U_ϵ is defined to be a cone over an ϵ -neighborhood in the unit sphere.

Next we consider the stratum $X_{n,2}$. The complement $X_{n,2} \setminus U_\epsilon(\overline{X}_{n,1})$ is a cone on a compact smooth manifold with boundary. For each point $x \in X_{n,2} \setminus U_\epsilon(\overline{X}_{n,1})$, the bilipschitz diffeomorphism ϕ_x and the induction hypothesis for X_{n-2} show the intrinsic and extrinsic metrics are bilipschitz equivalent on $U_x \cap X_n$. Arguing by compactness as before, we obtain the bilipschitz property for a sufficiently small neighborhood $U_\epsilon(\overline{X}_{n,2})$ of $\overline{X}_{n,2}$ in X_n . Here ϵ may have to be chosen smaller than the one chosen for $X_{n,1}$.

We proceed in this way until we obtain the bilipschitz property for a neighborhood of $X_{n,n-2}$ in the determinantal variety X_n . By compactness, the bilipschitz property holds for X_n itself.

5. BILIPSCHITZ PROPERTY FOR THE SET OF MATRICES OF POSITIVE DETERMINANT

We prove Theorem 1.1 by pushing a path in the determinantal variety out into the component $GL^+(n, \mathbb{R})$, and mimicking the proof of Theorem 3.5.

If the path γ passes via the apex $O \in X$, it can be replaced by a pair of straight line segments and pushed out into C as in Section 2.

Otherwise let $k \geq 1$ be the least rank of a matrix $\gamma(t)$ along the path, and let $a_k \leq b_k \in [0, 1]$ be respectively the first and last occurrences of a matrix of rank k . As in Section 3, we push the path $\gamma([a_k, b_k])$ into $X_{n,k} \subseteq X$, by applying the results of Section 4. We then push it out into $X_{n,k+1}$ by following a constant direction, and patch it up at the endpoints as in Step 3 of the proof of Theorem 3.5.

The new path is in $X_{n,k+1}$. We now choose the corresponding parameter values $a_{k+1}, b_{k+1} \in [0, 1]$ and proceed inductively. Thus the path can be pushed out into $X_{n,n-1}$. Finally we follow the normal direction to push the path out into $GL^+(n, \mathbb{R})$.

ACKNOWLEDGMENTS

M. Katz was partially funded by the Israel Science Foundation grant no. 1517/12. D. Kerner was partially supported by the Israel Science Foundation grant 844/14. This paper answers a question posed by Asaf

Shachar at MO¹ and we thank him for posing the question. We are grateful to Yves Cornulier for a helpful comment posted there. We are grateful to Jake Solomon for providing the proof in Section 4 of the general case of the bilipschitz property for the determinantal variety. We thank Jason Starr for a helpful comment posted at MO.² We thank Amitai Yuval for pointing out a gap in an earlier version of the article, and Alik Nabutovsky and Kobi Peterzil for useful suggestions.

REFERENCES

- [1] Birbrair, L.; Fernandes, A.; Lê, D.; Sampaio, J. “Lipschitz regular complex algebraic sets are smooth.” *Proc. Amer. Math. Soc.* **144** (2016), no. 3, 983–987.
- [2] Birbrair, L.; Mostowski, T. “Normal embeddings of semialgebraic sets.” *Michigan Math. J.* **47** (2000), no. 1, 125–132.
- [3] Kanovei, V.; Katz, K.; Katz, M.; Nowik, T. “Small oscillations of the pendulum, Euler’s method, and adequacy.” *Quantum Studies: Mathematics and Foundations* **3** (2016), no. 3, 231–236.
See <http://dx.doi.org/10.1007/s40509-016-0074-x>
and <http://arxiv.org/abs/1604.06663>
- [4] Katz, K.; Katz, M. “Bi-Lipschitz approximation by finite-dimensional imbeddings.” *Geom. Dedicata* **150** (2011), 131–136.
- [5] MacPherson, R.; Procesi, C. “Making conical compactifications wonderful.” *Selecta Math. (N.S.)* **4** (1998), no. 1, 125–139.
- [6] Nowik, T., Katz, M. “Differential geometry via infinitesimal displacements.” *Journal of Logic and Analysis* **7:5** (2015), 1–44.
See <http://dx.doi.org/10.4115/jla.2015.7.5>
and <http://arxiv.org/abs/1405.0984>
- [7] Pedersen, H.; Ruas, M. “Lipschitz Normal Embeddings and Determinantal Singularities.” See <http://arxiv.org/abs/1607.07746>

K. KATZ, DEPARTMENT OF MATHEMATICS, BAR ILAN UNIVERSITY, RAMAT GAN 52900 ISRAEL
E-mail address: katzmik@math.biu.ac.il

M. KATZ, DEPARTMENT OF MATHEMATICS, BAR ILAN UNIVERSITY, RAMAT GAN 52900 ISRAEL
E-mail address: katzmik@macs.biu.ac.il

D. KERNER, DEPARTMENT OF MATHEMATICS, BEN GURION UNIVERSITY OF THE NEGEV, BE’ER SHEVA 8410501 ISRAEL
E-mail address: dmitry.kerner@gmail.com

Y. LIOKUMOVICH, IMPERIAL COLLEGE, LONDON
E-mail address: y.liokumovich@imperial.ac.uk

¹See <http://mathoverflow.net/questions/222162>

²See <http://mathoverflow.net/questions/230668>