# One-dimensional stable rings

### Bruce Olberding

 $Department\ of\ Mathematical\ Sciences,\ New\ Mexico\ State\ University,\ Las\ Cruces,\ NM\\ 88003-8001$ 

#### Abstract

A commutative ring R is stable provided every ideal of R containing a nonzerodivisor is projective as a module over its ring of endomorphisms. The class of stable rings includes the one-dimensional local Cohen-Macaulay rings of multiplicity at most 2, as well as certain rings of higher multiplicity, necessarily analytically ramified. The former are important in the study of modules over Gorenstein rings, while the latter arise in a natural way from generic formal fibers and derivations.

We characterize one-dimensional stable local rings in several ways. The characterizations involve the integral closure  $\overline{R}$  of R and the completion of R in a relevant ideal-adic topology. For example, we show: If R is a reduced stable ring, then there are exactly two possibilities for R: (1) R is a Bass ring, that is, R is a reduced Noetherian local ring such that  $\overline{R}$  is finitely generated over R and every ideal of R is generated by two elements; or (2) R is a bad stable domain, that is, R is a one-dimensional stable local domain such that  $\overline{R}$  is not a finitely generated R-module.

*Keywords:* local ring, Cohen-Macaulay ring, stable ideal, two-generated ideal 2010 MSC: 13H10, 13E15, 13A15

#### 1. Introduction

The class of stable rings has a long history dating back at least to the "Ubiquity" paper of Bass, where he showed that rings for which every ideal can be generated by two elements are stable [6, Corollary 7.3]. Following Lipman [21] and Sally and Vasconcelos [39, 40], we define an ideal I of a commutative ring R to be stable if I is projective over its ring of endomorphisms. A ring R is stable if every regular ideal of R (that is, every ideal containing a nonzerodivisor) is  $stable^1$ . Independently of Bass, Lipman [21] studied stable ideals in one-dimensional semilocal Cohen-Macaulay rings and showed how stability was

Email address: olberdin@nmsu.edu (Bruce Olberding)

<sup>&</sup>lt;sup>1</sup>Our definition of a stable ring differs slightly from Sally and Vasconcelos [39, 40] in that we require only that every regular ideal is stable, not that every ideal is stable. However, for a reduced one-dimensional Cohen-Macaulay ring, the two definitions agree [35, p. 260].

reflected in terms of invariants of the ring such as multiplicity, embedding dimension and the Hilbert function. (We recall the definitions of these terms in Section 2.1.) The terminology of "stable" ideal originates with Lipman; it reflects the stabilization of a certain chain of infinitely near local rings. Lipman in turn was motivated to introduce these ideals as a way to unify ideas from Arf and Zariski involving singularities of plane curves, and to produce the largest ring between a one-dimensional local Cohen-Macaulay ring R and its integral closure having the same multiplicity sequence as R [21, Corollary 3.10].

Sally and Vasconcelos, motivated by the work of Bass and Lipman, studied stable Noetherian local rings in detail in [39, 40], and showed that a reduced local Cohen-Macaulay ring with finite normalization is stable if and only if every ideal of R can be generated by two elements, thus substantiating a conjecture of Bass and proving a partial converse to his theorem mentioned above. Reduced Noetherian local rings having finite normalization and every ideal generated by two elements are known in the literature as Bass rings because of their importance in Bass's article [6]. Sally and Vasconcelos [40, Example 5.4] gave the first example of a Noetherian stable domain R of multiplicity > 2 (and hence without finite normalization). They used a construction of Ferrand and Raynaud that exhibits R as the preimage of a derivation over a specific field of characteristic 2. Heinzer, Lantz and Shah [15, (3.12)] showed that this technique can be modified to produce over this same field of characteristic 2 for each e > 2 an analytically ramified Noetherian local stable domain of multiplicity e. More recently, analytically ramified Noetherian stable local rings have proved to be useful theoretical tools for classifying the rank one discrete valuation rings (DVRs) that arise as the normalization of an analytically ramified onedimensional Noetherian local domain, as well as for describing properties of the generic formal fiber of a Noetherian local domain [31, 32]; see Remark 2.15 and Theorem  $2.16.^2$ 

The main goal of this article is to tie all these results together. The Noetherian assumption is not essential to most of our arguments, and thus we use general ideal-theoretic methods to work in a setting in which R is a one-dimensional local ring with regular maximal ideal.<sup>3</sup> For example, we show in Theorem 4.2 that R is stable if and only if the integral closure  $\overline{R}$  is a Dedekind ring with the property that every R-submodule containing R is a ring. This generalizes a result of Rush [36, Theorem 2.4] to rings that are not necessarily Noetherian. A finer classification is possible by distinguishing when the integral closure of R is a finitely generated R-module: We show in Theorems 4.5 and 4.7 that R is stable and has finite integral closure if and only if a suitable completion  $\widetilde{R}$  of R is a Bass ring, while R is stable without finite integral closure if and only if there is a nonzero prime ideal P of  $\widetilde{R}$  such that  $P^2 = 0$  and  $\widetilde{R}/P$  is a DVR. From

<sup>&</sup>lt;sup>2</sup>The definitions of normalization, finite normalization and analytically ramified rings appear at the end of this section and in Section 2.1.

 $<sup>^3</sup>$ We use the terminology of "local ring" for a ring with a unique maximal ideal. In particular, we do not require a local ring to be Noetherian.

this we deduce in Corollary 4.9 that R is reduced and stable if and only if R is a Bass ring or R is a stable domain without finite integral closure. The rings in the first class, the Bass rings, are classical, while the second class, the stable domains without finite integral closure, have a rather transparent structure, as indicated by Theorem 2.13.

We also revisit the two-generator property in Section 5. For R a onedimensional local ring with regular maximal ideal M, we show that  $M^n$  is two-generated (i.e., can be generated by two elements), for some  $n \geq 2$ , if and only if every M-primary ideal of R is two-generated. From this we deduce:

**Theorem 5.6.** Let R be a Cohen-Macaulay ring with maximal ideal M. Then these statements are equivalent.

- (1)  $M^n$  is two-generated, for some  $n \geq 2$ .
- (2) R has Krull dimension 1 and multiplicity at most 2.
- (3) R is one of the following:
  - (3a) a Bass ring;
  - (3b) a one-dimensional analytically ramified stable domain; or
  - (3c) a ring containing a nonzero principal prime ideal P such that  $P^2 = 0$  and R/P is a DVR.

We also generalize a theorem of Greither to rings that are not a priori Noetherian: If R is a local ring with regular maximal ideal and the integral closure of R is a Dedekind ring generated by two elements as an R-module, then R is a Bass ring (Theorem 5.8).

Conventions. All rings are commutative with 1. The Jacobson radical of the ring R is denoted Jac R. The total ring of quotients of R is denoted Q(R). The integral closure of R is the integral closure of R in Q(R) and is denoted  $\overline{R}$ . We say that R has finite integral closure if  $\overline{R}$  is a finitely generated R-module. When R is reduced,  $\overline{R}$  is the normalization of R, and R has finite normalization if  $\overline{R}$  is a finitely generated R-module. We write (R, M) for a local ring R with maximal ideal M.

### 2. Preliminaries and background

### 2.1. One-dimensional Cohen-Macaulay rings

We collect in this subsection properties of one-dimensional Noetherian local rings that are needed in later sections. Recall that a local ring (R, M) is analytically unramified provided that the M-adic completion  $\widehat{R}$  has no nontrivial nilpotent elements.

**Proposition 2.1.** ([11, Corollary 1.21] and [24, pp. 263-264]) Let (R, M) be a local ring with maximal ideal M and integral closure  $\overline{R}$ . Then

- (1) If R is reduced and one-dimensional, and there exists t > 0 such that  $M^t$  is finitely generated, then R is Noetherian.
- (2) If R is Noetherian, M is regular and  $\overline{R}$  is finitely generated as an R-module, then R is reduced.
- (3) If R is Noetherian and analytically unramified, then  $\overline{R}$  is finitely generated as an R-module.

The embedding dimension of a Noetherian local ring R is the minimal number of elements needed to generate its maximal ideal M. For sufficiently large values of n, the length of  $R/M^n$  is a polynomial in n of degree  $d = \dim R$ . The product of d! and the leading coefficient of this polynomial is the multiplicity of R. See [24] for more information on multiplicity.

**Proposition 2.2.** ([41, Propositions 11.1.10 and 11.2.1], [38, Theorem 1.1, p. 49] and [23, Theorem 10.2, p. 90]) Let (R, M) be a one-dimensional Cohen-Macaulay ring with maximal ideal M and integral closure  $\overline{R}$ . Then

- (1) If  $M^2 = mM$  for some  $m \in M$ , then the embedding dimension and multiplicity of R agree. In particular, R has minimal multiplicity.
- (2) Every ideal of R can be generated by e or fewer elements, where e is the multiplicity of R.
- (3) R has finite integral closure if and only if R is analytically unramified.

### 2.2. Completions of a one-dimensional local ring

For a one-dimensional local ring (R, M) that is not Noetherian, the M-adic completion  $\widehat{R}$  of R can be too coarse to contain useful information about R. For example, if  $M = M^2$ , then  $\widehat{R} = R/M$ . For this reason we work with another completion that agrees with the M-adic completion of R if R is one-dimensional and Noetherian.

**Notation 2.3.** Let R be a one-dimensional local ring with regular maximal ideal M, and let m be a nonzerodivisor in M. We denote by  $\widetilde{R}$  the mR-adic completion of R and by  $\widehat{R}$  the M-adic completion of R. Since R has Krull dimension 1 and m is a nonzerodivisor, mR is M-primary, and hence the ring  $\widetilde{R}$  is independent of the choice of nonzerodivisor  $m \in M$ .

**Remark 2.4.** Let R be a one-dimensional local ring with regular maximal ideal M. If (i) M is finitely generated, or (ii) M is stable, then  $\widetilde{R}$  can be identified with  $\widehat{R}$ . In case (i) some power of M is contained in a principal regular ideal mR, and so the mR-adic and M-adic topologies agree on R. For case (ii), Lemma 3.4(6) implies that  $M^2 = mM$  for some  $m \in M$ . Thus  $M^2 \subseteq mR$  and so again the two topologies coincide.

**Proposition 2.5.** ([23, Theorem 2.1(3), p. 11], [23, Theorem 2.1(4), p. 11] and [23, Theorem 2.8(1), p. 19]) Let R be a one-dimensional local ring with regular maximal ideal M.

- (1) If I is a regular ideal of R, then  $I\widetilde{R}$  is a regular ideal of  $\widetilde{R}$ .
- (2) If  $\lambda: R \to \widetilde{R}$  denotes the canonical mapping, then  $\widetilde{R} = \lambda(R) + x\widetilde{R}$  for all nonzerodivisors  $x \in R$ .
- (3) If I is a regular ideal of R, then the mapping  $\lambda$  in (2) induces an isomorphism  $R/I \cong \widetilde{R}/I\widetilde{R}$  and  $I = \lambda^{-1}(I\widetilde{R})$ .

*Proof.* Matlis proves the statements in the proposition for what he calls the "R-completion" H of a commutative ring R; that is,  $H = \varprojlim R/I$ , where I ranges over the regular ideals of R. Since in our case R is local of Krull dimension one and the maximal ideal M of R is regular, the R-completion of R is simply  $\widetilde{R}$ . Thus the cited references apply also to  $\widetilde{R}$ .

### 2.3. Quadratic extensions

Rush [36, Proposition 2.1] has shown that, if R is a stable ring, every R-submodule of  $\overline{R}$  that contains R is a ring. In this section we review some properties of extensions that share this property.

**Definition 2.6.** An extension  $R \subseteq S$  of rings is *quadratic* if  $xy \in xR + yR + R$  for all  $x, y \in S$ ; equivalently, every R-module between R and S is a ring.

**Remarks 2.7.** (1) Quadratic extensions were considered by Handelman [14] and Rush [36], and more recently in [31, 32, 33].

- (2) If an extension  $R \subseteq S$  of rings is quadratic, then, for every  $x \in S$ , we have  $x^2 \in xR + R$ ; that is, every  $x \in S$  is a root of a monic polynomial of degree at most 2 with coefficients in R. Thus every quadratic extension is an integral extension.
- (3) If  $R \subseteq S \subseteq T$  is an extension of rings and  $R \subseteq T$  is quadratic, then  $R \subseteq S$  is quadratic. Also, if I is an ideal of S that is also an ideal of R, then  $R/I \subseteq S/I$  is quadratic if and only if  $R \subseteq S$  is quadratic.

A key tool for analyzing quadratic rings is Handelman's classification of finite-dimensional algebras that are quadratic extensions of a base field.

**Lemma 2.8.** (Handelman [14, Lemma 5]) Let F be a field and let S be a finite-dimensional F-algebra such that  $F \subseteq S$  is a quadratic extension. Then S is isomorphic as an F-algebra to one of the following:

- (i) F,
- (ii) a field extension of F of degree 2,
- (iii) a local ring with square zero maximal ideal and residue field isomorphic to  ${\cal F},$
- (iv)  $F \times F$ , or
- (v)  $F \times F \times F$ , in which case  $F = \mathbb{F}_2$ .

**Remark 2.9.** If K/F is a finite field extension that is a quadratic extension of rings in the sense of Definition 2.6, then from Lemma 2.8 it follows that  $[K:F] \leq 2$ . Thus an extension of fields that is quadratic in the sense of Definition 2.6 agrees with the usual notion of a quadratic field extension.

The following proposition is deduced in [34] from Handelman's lemma.

**Proposition 2.10.** [34, Proposition 3.3] If  $R \subseteq S$  is a quadratic extension, then there are at most three prime ideals of S lying over any prime ideal of R.

#### 2.4. Stable rings

Noetherian stable rings have been studied in a number of contexts; cf. [2, 6, 9, 12, 14, 15, 21, 35, 39, 40, 42, 43]. Also, the related class of archimedean stable domains have been investigated recently by Gabelli and Roitman [10] and shown to generalize features of one-dimensional stable domains, and hence Noetherian stable domains, to a wider setting.

The following theorem summarizes some of the results discussed in the introduction.

**Theorem 2.11.** (cf. [6, Corollary 7.3], [9], [35, Theorem 1.4] and [39, Theorem 2.4]) Let R be a reduced Cohen-Macaulay ring. If every ideal of R can be generated by 2 elements (equivalently, R has Krull dimension  $\leq 1$  and multiplicity  $\leq 2$ ), then R is a stable ring. Conversely, if R is a reduced stable ring with finite normalization, then R is a Bass ring.

Thus in the case that R is an analytically unramified local Cohen-Macaulay ring, R is stable if and only if every ideal of R can be generated by two elements. The analytically ramified case is more complicated, as illustrated by results throughout the paper. We single out in the following definition a specific class of such rings.

**Definition 2.12.** A domain R is a *bad stable domain* if R is a one-dimensional stable local domain without finite normalization.

Bad stable domains were studied in [31, 32]. The appellation "bad" here is borrowed from the title of Nagata's appendix in [25], "Examples of bad Noetherian rings," and refers to the fact that the domain does not have finite normalization.

**Theorem 2.13.** [31, Proposition 2.1, Theorems 3.4 and 4.2 and Corollaries 3.5 and 4.3] The following are equivalent for a local domain R with quotient field F.

- (1) R is a bad stable domain.
- (2)  $\overline{R}$  is a DVR,  $R \subseteq \overline{R}$  is a quadratic extension and  $\overline{R}/R$  is a divisible R-module.
- (3) With  $\widetilde{R}$  as in Notation 2.3, there is a nonzero prime ideal P of  $\widetilde{R}$  such that  $P^2 = 0$  and  $\widetilde{R}/P$  is a DVR.

- (4)  $\overline{R}$  is a DVR and  $\overline{R}/R \cong \bigoplus_{i \in I} F/\overline{R}$  for some index set I.
- (5)  $\overline{R}$  is a DVR and  $\overline{R}/R$  is a direct sum of divisible Artinian uniserial R-modules.

Moreover, R is a bad Noetherian stable domain of multiplicity e if and only if the cardinality of the set I in (4) is e-1.

Theorem 2.13(3) and a theorem of Lech's can be used to guarantee the existence of bad stable domains with prescribed completions:

**Corollary 2.14.** The following are equivalent for a Noetherian local ring (R, M) that is complete in the M-adic topology.

- (1) The ring R is the completion of a bad Noetherian stable domain.
- (2) No nonzero integer of R is a zerodivisor and there is a prime ideal P of R such that  $P^2 = 0$  and R/P is a DVR.

*Proof.* That (1) implies (2) follows from Theorem 2.13(3). Conversely, assume (2). By a theorem of Lech [20, Theorem 1], the fact that  $M \neq 0$ , M is not an associated prime of R, and no nonzero integer of R is a zerodivisor implies that the ring R is the completion of a Noetherian local domain A. By Theorem 2.13(3), A is a bad Noetherian stable domain.

Remark 2.15. The archetypal example of a ring R such as in Corollary 2.14 is produced by Nagata idealization: Let V be a complete DVR, and let L be a nonzero finitely generated free V-module. Define V\*L as a V-module to be  $V\oplus L$ , and view V\*L as a ring with multiplication given by  $(v_1,\ell_1)(v_2,\ell_2)=(v_1v_2,v_1\ell_2+v_2\ell_1)$  for all  $v_1,v_2\in V$  and  $\ell_1,\ell_2\in L$ . With P=0\*L, the ring R=V\*L is a complete local ring meeting the hypotheses of Corollary 2.14, and hence R is the completion of a bad Noetherian stable domain A. Moreover, since the multiplicity of A is the same as that of its completion R, the multiplicity of A is  $e=1+\operatorname{rank} L$ .

To further motivate the class of bad stable domains, we mention three other contexts in which these rings arise. An extension  $U \subseteq V$  of DVRs is *immediate* if U and V have the same residue field and the maximal ideal of U extends to the maximal ideal of V. The **m**-adic completion of a DVR is an immediate extension.

**Theorem 2.16.** ([31, Corollary 5.7 and Theorem 5.9] and [32, Theorem 4.1])

- (1) If k is a field and A is an affine k-domain with Krull dimension d > 1 and quotient field F, then there exists a bad Noetherian stable domain R between A and F having multiplicity d and residue field finite over k.
- (2) If (A, M) is a Noetherian local domain and there exists a DVR V of A that birationally dominates A and satisfies V = A + MV, then the kernel of the exterior differential  $d_{V/A}: V \to \Omega_{V/A}$  is a bad stable domain between A and V.

(3) If  $U \subseteq V$  is an immediate extension of DVRs having quotient fields Q and F, respectively, then there is a one-to-one correspondence between bad stable domains R containing U and having normalization V and proper full V-submodules of the module  $\Omega_{F/Q}$  of Kähler differentials.

Remarks 2.17. (1) The ring R in Theorem 2.16(1) is constructed along the following lines: For an appropriate local A-algebra B contained in F such that B is essentially of finite type over k, teh ring R is defined to be  $R = F \cap (\widehat{B}/P^{(2)})$ , where  $\widehat{B}$  is the **m**-adic completion of B, P is a certain prime ideal of  $\widehat{B}$  and  $P^{(2)}$  is the second symbolic power of P. That R has multiplicity d depends on the fact that B is an excellent local ring [31, Corollary 5.6].

- (2) Every bad Noetherian stable domain of positive characteristic must arise as in Theorem 2.16(2) from a derivation [32, Theorem 6.4].
- (3) The one-to-one correspondence in Theorem 2.16(3) is used in [32, Theorem 7.6] to describe complete DVRs that are the normalization of an analytically ramified Noetherian local domain. For example, every equicharacteristic complete DVR is the normalization of both Noetherian and non-Noetherian bad stable domains.

### 3. Quadratic rings

In this section we introduce the notion of a quadratic ring and show that these rings exemplify some key technical features of stable local rings. We require first some preliminaries on stable ideals.

**Definitions 3.1.** Let R be a ring, and let Q(R) be its total ring of quotients.

- (1) An R-submodule I of Q(R) such that  $bI \subseteq R$  for some nonzerodivisor  $b \in R$  is called a *fractional ideal* of R. The fractional ideal I is *regular* if it contains a nonzerodivisor.
- (2) If I is a regular fractional ideal of the ring R, then I is invertible if I is a finitely generated fractional ideal such that  $IR_M$  is a principal fractional ideal of  $R_M$  for each maximal ideal M of R.
- (3) The ring R is finitely  $stable^4$  if every finitely generated regular ideal of R is stable.

**Remark 3.2.** Let I be a regular fractional ideal of a ring R. Then I is invertible if and only if IJ = R for some fractional ideal J [16, Proposition 2.3, p. 97]. If R has only finitely many maximal ideals, then an invertible ideal is principal [17, Exercise 2.11B, p. 33.].

**Notation 3.3.** Let I be a regular fractional ideal of a ring R. Then, by [34, (2.8)],  $\operatorname{End}(I) = \{q \in Q(R) : qI \subseteq I\}$ . To simplify notation we write  $E(I) = \operatorname{End}(I)$ .

 $<sup>^4\</sup>mathrm{The}$  rings we call "finitely stable" were called "stable" by Rush in [35, 36].

We collect in the next lemma some basic properties of a regular ideal I and its endomorphism ring E(I).

**Lemma 3.4.** With the terminology of Definition 3.1, let I be a regular fractional ideal of a ring R. Then

- (1) I is a regular fractional ideal of E(I).
- (2)  $I \subseteq \operatorname{Jac}(E(\operatorname{Jac} R))$ .
- (3) I is stable if and only if I is an invertible fractional ideal of E(I).
- (4) If  $I^2 = AI$  for some invertible fractional ideal A of R with  $A \subseteq I$ , then I = AE(I) and I is stable.
- (5) If there exists  $x \in I$  such that  $I^2 = xI$ , then I is stable.
- (6) Let R be finitely stable and local. Then I is stable if and only if  $I^2 = xI$  for some  $x \in I$ .

*Proof.* (1) This follows from the fact that every nonzerodivisor in R is also a nonzerodivisor in Q(R).

- (2) Let  $x \in \operatorname{Jac} R$  and  $w \in E(\operatorname{Jac} R)$ . Then  $wx \in \operatorname{Jac} R$ , and so 1 wx is a unit of R, hence also of  $E(\operatorname{Jac} R)$ . Since w was arbitrary in  $E(\operatorname{Jac} R)$ , we have  $x \in \operatorname{Jac} E(\operatorname{Jac} R)$ .
  - (3) This is proved in [34, Proposition 2.11].
  - (4) This is proved in [34, Proposition 2.13].
- (5) Since I is regular and  $I^2 \subseteq xR$ , the ideal xR is regular, hence invertible, and so this follows from (4).
- (6) Under the assumptions of (6), it is proved in [34, Corollary 5.7] that a regular ideal J of R is stable if and only if J is a principal ideal of E(I). The characterization in (6) now follows from (5).

**Notation 3.5.** Let R be a ring such that Jac R is a regular ideal. We associate to R a tower of rings in Q(R) by defining

$$R_0 := R$$
,  $R_i := E(\operatorname{Jac} R_{i-1})$  for every  $i \ge 1$ ,  $R_\infty := \bigcup_{i=1}^\infty R_i$ .

Lemma 3.4(1) implies that for each i, Jac  $R_i$  is a regular ideal of  $R_{i+1} = E(\operatorname{Jac} R_i)$ . In Lipman's terminology [21, Section 2], if R is a local Cohen-Macaulay ring and the Jacobson radical of each  $R_i$  is stable, then the ring  $R_{i+1}$  is the blow-up of  $R_i$  at its Jacobson radical and the localizations of the rings  $R_i$  at maximal ideals are the local rings "infinitely near" R.

**Definition 3.6.** Let R be a ring such that Jac R is a regular ideal. With Notation 3.5 we define R to be a quadratic ring if

- (1) R is local,
- (2)  $R \subseteq R_{\infty}$  is a quadratic extension, and
- (3) the maximal ideal of R is stable and regular.

As we show in Theorem 3.9, the relevance of this notion is that a finitely stable local ring with stable regular maximal ideal is a quadratic ring. Some technical properties of quadratic rings are elaborated in Lemmas 3.7 and 3.10 and Proposition 3.11. Parts of these results were proved for finitely stable local domains with stable maximal ideal in [29, Section 4], but the arguments given here are somewhat different and more general in that they permit zero divisors.

**Lemma 3.7.** Let R be a local ring with regular maximal ideal M and residue field F = R/M. Let  $R_1$  be as in Notation 3.5. Suppose that  $R \subseteq R_1$  is a quadratic extension with  $R \neq R_1$ .

- (1) If  $R_1$  is a local ring such that  $\operatorname{Jac} R_1 = M$ , then  $R_1 = R_2$  and  $R_1/M$  is a field extension of F of degree 2.
- (2) If  $R_1$  is a local ring such that  $\operatorname{Jac} R_1 \neq M$ , then  $R_1 = R + M_1$  and  $R_1/M_1 \cong F$ .
- (3) If  $R_1$  is not a local ring, then  $\operatorname{Jac} R_1 = M$ ,  $R_1 = R_2$  and  $R_1/M$  is isomorphic as an F-algebra to either
  - (i)  $F \times F$ , or
  - (ii)  $F \times F \times F$ , in which case  $F = \mathbb{F}_2$ .
- (4)  $(\operatorname{Jac} R_1)^2 \subseteq M$ .
- Proof. (1) Suppose that  $R_1$  is a local ring such that Jac  $R_1 = M$ . Then  $R_1/M$  is a field and Jac  $R = M = \operatorname{Jac} R_1$ . Hence  $R_1 = R_2$ . To see that  $R_1/M$  has degree 2 over F, let  $x,y \in R_1 \setminus R$ , and let S = xR + yR + R. Then S is a ring since  $R \subseteq R_1$  is a quadratic extension. Hence  $R/M \subseteq S/M$  is a quadratic extension, which by Remark 2.7 must be integral. Since  $R_1/M$  is a field, the subring S/M is an integral domain, and hence since S/M is integral over the field R/M, S/M is a field. Since  $R/M \subseteq S/M$ , Lemma 2.8 implies that S/M has degree 2 as a field extension of R/M. Since  $x,y \notin R$ , this forces S/M = (xR + R)/M = (yR + R)/M. Hence  $x \in yR + R$  for all  $x,y \in R_1 \setminus R$ , and so for each  $x \in R_1 \setminus R$ , we have  $R_1 = xR + R$ . Therefore the degree of  $R_1/M$  over R/M is 2.
- (2) Suppose that  $R_1$  is a local ring such that  $\operatorname{Jac} R_1 \neq M$ . Let  $M_1$  denote the maximal ideal of  $R_1$ . We claim that  $R_1 = R + M_1$ . By Lemma 3.4(2),  $M \subsetneq M_1$ . Let  $x \in M_1 \setminus M$ , let  $y \in R_1$ , and let S = R + xR + yR. Since  $x \in M_1 \cap S$ , we have  $M_1 \cap S \neq M$ . Because  $R \subseteq R_1$  is quadratic, S is a ring, and, by Remarks 2.7(2),  $R \subseteq R_1$  is an integral extension. Since  $R_1$  is local, so is S. Also, S/M cannot be a field since  $M \subsetneq M_1 \cap S \subsetneq S$ . Furthermore, since S is local, the ring S/M is not isomorphic to  $F \times F$  or  $F \times F \times F$ . By Lemma 2.8, S/M is a local ring with square zero maximal ideal and residue field F. If S is the maximal ideal of S, we have S is a local ring with the choice of S is a local ring with square zero maximal ideal and residue field S. If S is the maximal ideal of S, we have S is a local ring with the choice of S is a local ring with the choice of S is a local ring with the residue field S. If S is the maximal ideal of S, we have S is a local ring with the choice of S is a local ring with the residue field S. If S is the maximal ideal of S, we have S is a local ring with the residue field S. The choice of S is a local ring with the residue field S is a local ring with the residue field S.
- (3) Suppose that  $R_1$  is not a local ring, and let k denote the number of maximal ideals of  $R_1$ . By Lemma 3.4(2),  $M \subseteq \operatorname{Jac} R_1$ , and so  $R_1/M$  has k > 1 maximal ideals. Also, since  $R \subseteq R_1$  is a quadratic extension of rings,

 $R/M \subseteq R_1/M$  is a quadratic extension in which  $R_1/M$  has more than one maximal ideal. By [34, Lemma 5.2],  $R_1/M$  is isomorphic as an F-algebra to  $\prod_{i=1}^k F$ , and  $F = \mathbb{F}_2$  when k = 3. Also, since we have obtained a decomposition of  $R_1/M$  as a finite product of fields, Jac  $R_1 = M$  and hence  $R_1 = R_2$ .

(4) In light of (1) and (3), the only case in which (4) is not immediate is that of (2). Suppose that  $R_1$  is local and  $M \neq \operatorname{Jac} R_1$ . By Lemma 3.4(2),  $M \subsetneq \operatorname{Jac} R_1$ . Let  $a, b \in \operatorname{Jac} R_1$  with  $a, b \notin M$ . Define T = aR + bR + R. Then, as in the proof (2), Lemma 2.8 forces the square of the maximal ideal of the ring T to be contained in M. Thus  $ab \in M$ , and it follows that  $(\operatorname{Jac} R_1)^2 \subseteq M$ .  $\square$ 

For the next lemma, we recall that a ring R is  $Pr\ddot{u}fer$  if every finitely generated regular ideal is invertible.

**Lemma 3.8.** ([36, Proposition 2.1 and Theorem 2.3] and [34, Corollary 5.11])  $A \ ring \ R \ is finitely \ stable \ if \ and \ only \ if$ 

- (a)  $R \subseteq \overline{R}$  is a quadratic extension,
- (b)  $\overline{R}$  is a Prüfer ring, and
- (c) there are at most two maximal ideals of  $\overline{R}$  lying over each regular maximal ideal of R.

The next theorem shows that the class of quadratic rings contains the finitely stable local rings with stable maximal ideal.

**Theorem 3.9.** Let (R, M) be a finitely stable local ring with stable regular maximal ideal M, and let  $R_{\infty}$  be defined as in Notation 3.5. Then R is a quadratic ring for which  $R_{\infty}$  has at most two maximal ideals.

*Proof.* With  $\{R_i\}_{i=0}^{\infty}$  as in Notation 3.5, we prove the theorem by establishing four claims.

CLAIM 1. For each  $i \geq 0$ , if  $R_i$  is a local ring with maximal ideal  $M_i$ , then  $R_i \subseteq R_{i+1}$  is a quadratic extension.

We use the fact that  $R_i \subseteq R_{i+1}$  is a quadratic extension if and only if every finitely generated  $R_i$ -submodule of  $R_{i+1}$  containing  $R_i$  is a stable fractional ideal of  $R_i$ ; see [34, Proposition 3.5] or [35, proof of Lemma 2.1]. Thus we let A be a finitely generated  $R_i$ -submodule of  $R_{i+1}$  with  $R_i \subseteq A$ . Let  $m \in M$  be a nonzerodivisor. Then  $m \in M_i$  and

$$m \in mR_i \subseteq mA \subseteq mR_{i+1} \subseteq R_i$$

since  $R_{i+1} = \{q \in Q(R) : qM_i \subseteq M_i\}$ . Therefore mA is a finitely generated regular  $R_i$ -submodule of  $R_i$ , hence a finitely generated regular ideal of  $R_i$ . Every ring between a finitely stable ring and its total quotient ring is finitely stable [34, Proposition 5.1]. Hence  $R_i$  is finitely stable, and so mA is a stable ideal of  $R_i$  by Definition 3.1(3). Since  $mA \cong A$ , A is a stable fractional ideal of  $R_i$ . This proves Claim 1.

CLAIM 2. For each  $i \geq 0$  with  $R_i \subsetneq R_{i+1}$ ,  $R_i$  is local.

For i=0, the claim holds, since  $R=R_0$  is local. Suppose that i>0 and that  $R_i$  is local whenever  $R_i \subseteq R_{i+1}$ . If  $R_{i+1} \subseteq R_{i+2}$ , then also  $R_i \subseteq R_{i+1}$  and so  $R_i$  is local by induction. By Claim 1,  $R_i \subseteq R_{i+1}$  is a quadratic extension. Since  $R_{i+1} \neq R_{i+2}$ , Lemma 3.7(3) implies that  $R_{i+1}$  is a local ring, as desired for Claim 2.

CLAIM 3. (i) For each  $i \geq 0$ ,  $R_i \subseteq R_{i+1}$  is an integral extension. (ii) The extension  $R \subseteq R_{\infty}$  is integral.

The first statement follows from Claims 1 and 2 and Remark 2.7(2). The second statement follows from the first, since compositions of integral extensions are again integral [7, Proposition 6, p. 307].

Claim 4.  $R_{\infty}$  has at most two maximal ideals.

By Claim 3,  $R_{\infty} \subseteq \overline{R}$ , and by Lemma 3.8,  $\overline{R}$  has at most two maximal ideals. Since  $\overline{R}$  is an integral extension of  $R_{\infty}$ ,  $R_{\infty}$  also has at most two maximal ideals.

Claim 5. R is a quadratic ring.

By Lemma 3.8,  $R \subseteq \overline{R}$  is a quadratic extension, and Claim 3 implies  $R_{\infty} \subseteq \overline{R}$ . By Remark 2.7(3),  $R \subseteq R_{\infty}$  is a quadratic extension. By Definition 3.6, R is a quadratic ring.

By adding to Lemma 3.7 the assumption that the maximal ideal of R is stable, we obtain stronger results in the case where  $R_1$  is local. In particular,  $R_1$  inherits the property of having a stable maximal ideal.

**Lemma 3.10.** Let R be a local ring with stable regular maximal ideal M. Suppose that  $R \subseteq R_1$  is a quadratic extension and  $R_1$  is a local ring with maximal ideal  $M_1$ .

- (1) If  $R_1 \neq R_2$ , then  $M_1 = MR_2$ ,  $M_1$  is a stable ideal of  $R_1$  and  $R_1 = R + M_1$ .
- (2) If  $R_1 = R_2$ , then  $M_1$  is a principal ideal of  $R_1$  and  $R_1 = R + xR$  for every  $x \in R_1 \setminus R$ .

*Proof.* By Lemma 3.4(3), M is an invertible ideal of  $R_1$ . Since  $R_1$  is a local ring,  $M = mR_1$  for some nonzerodivisor  $m \in M$ . By Lemma 3.7(4),  $M_1^2 \subseteq M = mR_1$ , so that  $M_1^2 = mA$  for some ideal A of  $R_1$ .

CLAIM: (i) If  $A = R_1$ , then  $M_1$  is invertible in  $R_1$  and  $R_1 = R_2$ . (ii) If  $A \subsetneq R_1$ , then  $M_1^2 = mM_1$  and  $M_1 = mR_2 = MR_2$ .

For (i),  $A=R_1\Longrightarrow mR_1=M_1^2\Longrightarrow M_1(m^{-1}M_1)=R_1\Longrightarrow M_1$  is invertible. We have

$$R_2 = \{q \in Q(R_1) : qM_1 \subseteq M_1\} = \{q \in Q(R_1) : qM_1M_1^{-1} \subseteq M_1M_1^{-1}\} = R_1,$$

and so (i) holds.

For (ii),  $A \subsetneq R_1 \Longrightarrow M_1^2 = mA \subseteq mM_1 \subseteq M_1^2 \Longrightarrow M_1^2 = mM_1$ . By Lemma 3.4(4) we have

$$M_1 = mR_2 \subseteq MR_2 \subseteq M_1R_2 \subseteq mR_2 \Longrightarrow M_1 = mR_2 = MR_2$$

and so (ii) holds.

We return to the proof of Lemma 3.10. For (1),  $M_1 = MR_2$  and  $M_1$  is invertible in  $R_2$  by Claim (ii). Thus  $M_1$  is stable by Lemma 3.4(3). Since  $E(M) = R_1 \neq R_2 = E(M_1)$ , we have  $M \neq M_1$ . By Lemma 3.7(2),  $R_1 = R + M_1$ . This verifies (1).

For (2), we have  $R_1 = R_2$ . If  $M_1 = M$ , we have  $M_1 = mR_1$  is principal. By Lemma 3.7(1), the field  $R_1/M$  has degree 2 over R/M, and hence  $R_1 = R + xR$  for all  $x \in R_1 \setminus R$ , and so (2) holds if  $M_1 = M$ .

Thus we assume that  $M_1 \neq M$ . We have  $M_1$  is a principal ideal of  $R_1$  in either case  $A = R_1$  or  $A \subseteq R_1$ , using either (i) of the claim  $(M_1$  is invertible and so is a principal ideal of  $R_1$ ), or (ii) of the claim  $(M_1 = mR_2 = mR_1)$ . Thus the first part of (2) holds.

It remains to show  $R_1 = R + xR$ , for all  $x \in R_1$ , given that  $R_1 = R_2$ ,  $M_1 = nR_1$ , for some  $n \in R_1$ , and  $M_1 \neq M$ . We have also  $R \neq R_1$  and n is a nonzerodivisor.

By Lemma 3.7(4),  $M_1^2 \subseteq M$ . If  $mM_1 = M_1^2$ , then

$$mnR_1 = n^2R_1 \Longrightarrow M = mR_1 = nR_1 = M_1$$

a contradiction. Thus  $mM_1 \subsetneq M_1^2 \subseteq M = mR_1$ . Hence there exist  $c, d \in M_1$  with  $cd \in mR_1 \setminus mM_1$ ; say cd = um, where u is a unit of  $R_1$ . Then  $m = u^{-1}cd \in M_1^2$ , and

$$mR_1 \subseteq M_1^2 \subseteq M = mR_1 \Longrightarrow M_1^2 = M.$$

Therefore

$$M_1/M = M_1/M_1^2 \cong R_1/M_1 \cong R/M$$
,

by Lemma 3.7(2). Since R/M is a field, we have two equations.

(3.10a) 
$$M_1 = qR + M$$
,  $\forall q \in M_1 \setminus M$ . (3.10b)  $R_1 = R + M_1$ .

Let  $x \in R_1 \setminus R$ . By Equation 3.10b,  $x = r + m_1$ , for some  $r \in R$  and  $m_1 \in M_1$ , where  $m_1 \notin M$ , since  $x \notin R$ . Then, using Equations 3.10b and 3.10a, we have

$$R + xR \subseteq R_1 = R + M_1 \subseteq R + m_1 R = R + (r + m_1)R = R + xR.$$

This completes the proof of the lemma.

We collect now some of the main technical properties of quadratic rings that will be needed in the next section.

**Proposition 3.11.** Let R be a quadratic ring with maximal ideal M, let  $\{R_i\}_{i=0}^{\infty}$  and  $\{R_{\infty}\}$  be as in Notation 3.5, and let

$$n = \min\{k \in \{0, 1, 2, \dots, \infty\} : R_k = R_\infty\}.$$

Then the following statements hold for R.

- (1) There exists a nonzerodivisor  $m \in M$  such that  $M = mR_1$ .
- (2) For each i < n,  $R_i = R + M_i$  and  $R_i$  is a quadratic ring with stable maximal ideal  $M_i = mR_{i+1}$ .
- (3) If  $n = \infty$ , then  $R_{\infty}$  is a local ring with maximal ideal  $mR_{\infty}$  and  $R_{\infty} = R + mR_{\infty}$ .
- (4)  $R_{\infty}$  has at most 3 maximal ideals, each of which is principal.
- (5) If  $R_{\infty}$  is not local, then  $\operatorname{Jac} R_{\infty} = mR_{\infty}$ .
- (6) The R-module  $R_{\infty}$  is finitely generated if and only if  $n < \infty$ .
- (7) Suppose  $n < \infty$  and  $J := \bigcap_{i=1}^{\infty} M^i$  is a one-dimensional ideal of R. Then J is the intersection of the one-dimensional prime ideals of R, the number of which is not more than the number of maximal ideals of  $R_{\infty}$ . Moreover J is also the intersection of the one-dimensional prime ideals of  $R_{\infty}$ , and hence J is an ideal of  $R_{\infty}$ .
- *Proof.* (1) By Definition 3.6, M is stable. Since  $R \subseteq R_1$  is a quadratic extension, Proposition 2.10 implies that  $R_1$  has at most 3 maximal ideals. By Lemma 3.4(3), M is invertible in  $R_1$ . Since  $R_1$  is local,  $M = mR_1$ , for some nonzerodivisor  $m \in M$ .
- (2) The proof is by induction. If n=0 or i=0, the statement holds by (1). Suppose the result holds for some i with 0 < i < n-1. Then  $R_i$  is a quadratic ring with stable maximal ideal  $M_i = mR_{i+1}$  and  $R_i = R + M_i$ . Also  $R \subsetneq R_i \subsetneq R_{i+1} \subsetneq R_{i+2}$ , by the minimality of n. By Lemma 3.7(3),  $R_{i+1}$  is a local ring. By Lemma 3.7(1), the maximal ideal  $M_{i+1}$  of  $R_{i+1}$  properly contains  $M_i$ . By Lemma 3.10(1),  $M_{i+1} = M_iR_{i+2} = mR_{i+1}R_{i+2} = mR_{i+2}$ , and  $R_{i+1} = R_i + M_{i+1} = R + M_{i+1}$ . Since  $R \subseteq R_{\infty}$  is a quadratic extension, it is clear that  $R_{i+1} \subseteq R_{\infty}$  is also a quadratic extension, and so  $R_{i+1}$  is a quadratic ring.
- (3) If  $n = \infty$ , then by (2), each  $R_i$  is local with maximal ideal  $mR_{i+1}$  and  $R_i = R + M_i$ . Hence  $R_{\infty}$  is a local ring with maximal ideal  $mR_{\infty}$  and  $R_{\infty} = R + mR_{\infty}$ .
- (4) By Proposition 2.10,  $R_n$  has at most three maximal ideals. If  $n = \infty$ , then (4) follows from (3). If n = 0, then  $R = R_0$  is local and it has principal maximal ideal by (1), so that (4) holds.

Suppose that  $0 < n < \infty$ . Then  $R_{n-1} \neq R_n$ . By (2),  $R_{n-1}$  is a quadratic ring and the maximal ideal  $M_{n-1} = mR_n$  is stable. Also  $R_n = R_{n+1}$ . If  $R_n$  is local, then Lemma 3.10(2) implies that the maximal ideal  $M_n$  is principal, and so we have established (4).

- If  $R_n$  is not local, then Lemma 3.7(3) implies that Jac  $R_n = M_{n-1} = mR_n$ . Since  $R_n$  has at most three maximal ideals, we may denote them by  $N_1, N_2, \ldots, N_t$ , where  $2 \le t \le 3$ . Then  $mR_n = \bigcap_{i=1}^t N_i = \prod_{i=1}^t N_i$ , by [24, Theorem 1.3]. Then each  $N_i$  is invertible; e.g.,  $N_1$  has inverse  $m^{-1} \prod_{i=2}^t N_i$ . By Remark 3.2, each of the  $N_i$  is principal. Thus (4) holds.
- (5) Since  $R_{\infty}$  is not local, (3) implies that  $n < \infty$ . As noted in the proof of (4), we have Jac  $R_n = mR_n$ .
  - (6) If  $n = \infty$ , then clearly  $R_{\infty} = \bigcup_{i=1}^{\infty} R_i$  is not finitely generated.

For the converse, the case n = 0 is clear. Let  $0 < n < \infty$ . We show first that  $R_n$  is a finitely generated  $R_{n-1}$ -module.

Case 1: If  $R_n$  is local, then Lemma 3.10(2) implies that  $R_n = R_{n-1} + xR_{n-1}$ , for every  $x \in R_n \setminus R_{n-1}$ , and so  $R_n$  is finitely generated as an  $R_{n-1}$ -module if  $R_n$  is local.

Case 2: If  $R_n$  is not local, then, by (3),  $n < \infty$ . By (2),  $R_{n-1}$  is a quadratic ring, and so by Lemma 3.7(3),  $R_n/M_{n-1}$  has dimension at most three as a vector space over  $R_{n-1}/M_{n-1}$ . Therefore  $R_n$  is a finitely generated  $R_{n-1}$ -module.

If n = 1, then the proof is complete. If n > 1, the result will follow from the following claim (using induction).

CLAIM. Suppose  $0 \le k \le n-2$  and  $R_{k+2}$  is finitely generated as an  $R_{k+1}$ -module. Then  $R_{k+1}$  is finitely generated as an  $R_k$ -module.

Write  $R_{k+2} = b_1 R_{k+1} + \dots + b_t R_{k+1}$ , for some  $b_1, \dots, b_t \in R_{k+2}$ . By (2), we have  $M_{k+1} = m R_{k+2}$  and

$$R_{k+1} = R_k + mR_{k+2} = R_k + mb_1R_{k+1} + \dots + mb_tR_{k+1}$$
  
=  $R_k + mb_1(R_k + mR_{k+2}) + \dots + mb_t(R_k + mR_{k+2}).$ 

By Lemma 3.7(4),  $m(mR_{k+2}) \subseteq M_{k+1}^2 \subseteq M_k$ . Each  $mb_i \in R_{k+1}$ . Therefore

$$R_{k+1} \subseteq R_k + mb_1R_k + \dots + mb_tR_k + M_k \subseteq R_{k+1}$$
.

This proves the claim.

(7) By (4),  $R_{\infty}$  has at most 3 maximal ideals, say  $N_1,\ldots,N_r$ , all of which are regular and principal. For each  $j=1,\ldots,r$ , since  $N_j$  is a principal regular ideal of  $R_{\infty}$ , it follows that  $Q_j:=\bigcap_{k>0}N_j^k$  is a prime ideal and every prime ideal of  $R_{\infty}$  properly contained in  $N_j$  is contained in  $Q_j$  [3, Theorem 2.2]. Therefore each nonmaximal prime ideal of  $R_{\infty}$  is contained in one of the  $Q_j$ . Now  $R\subseteq R_{\infty}$  is a quadratic, hence integral, extension. By assumption  $J=\bigcap_{k>0}M^k$  is a one-dimensional ideal of R. Every prime ideal of  $R_{\infty}$  containing J also has dimension at most one. Since  $J\subseteq Q_1\cap\cdots\cap Q_r$ , each prime ideal  $Q_i$  has dimension one. Since also each nonmaximal prime ideal of  $R_{\infty}$  is contained in one of the  $Q_i$ , the set of one-dimensional prime ideals of  $R_{\infty}$  is  $\{Q_1,\ldots,Q_r\}$ . Since  $R_{\infty}$  is an integral extension of R, it follows that  $\{Q_1\cap R,\ldots,Q_r\cap R\}$  is the set of one-dimensional prime ideals of R. Now since  $\mathrm{Jac}R_{\infty}=N_1\cdots N_r=N_1\cap\cdots\cap N_r$ , we have that  $\bigcap_{k>0}(\mathrm{Jac}R_{\infty})^k=Q_1\cap\cdots\cap Q_r$ . In fact, since  $R_n=R_{\infty}$ , an inductive argument using Lemma 3.7(4) shows that some power of the ideal  $\mathrm{Jac}\,R_{\infty}$  is contained in M. Hence  $J=\bigcap_{k>0}M^k=\bigcap_{k>0}(\mathrm{Jac}\,R_{\infty})^k=Q_1\cap\cdots\cap Q_r$ .  $\square$ 

Corollary 3.12. Let R be a one-dimensional quadratic ring with maximal ideal M, let  $J = \bigcap_{i=1}^{\infty} M^i$ , let  $\overline{R}$  be the integral closure of R, and let  $\{R_i\}_{i=0}^{\infty}$  and  $R_{\infty}$  be as in Notation 3.5. Then

- (1)  $R_{\infty} = \overline{R}$ .
- (2) The R-module  $\overline{R}$  is finitely generated if and only if  $R_{\infty} = R_n$  for some  $n \geq 0$ .

- (3) The ring R/J has integral closure  $R_{\infty}/J$  and total quotient ring Q(R)/J.
- *Proof.* (1) Since  $R \subseteq R_{\infty}$  is a quadratic, hence integral, extension, and R has Krull dimension one,  $R_{\infty}$  also has Krull dimension one. Since there are only finitely many maximal ideals of the one-dimensional ring  $R_{\infty}$ , all of which are principal and regular, it follows that every regular ideal of  $R_{\infty}$  is principal, and hence the integral extension  $R_{\infty}$  of R is integrally closed in Q(R); cf. [19, Theorem 10.18, p. 237].
  - (2) This follows from (1) and Proposition 3.11(6).
- (3) By Proposition 3.11(1), there exists a nonzerodivisor  $m \in M$  such that  $M^2 = mM$ . Since R has dimension one, Q(R) = R[1/m]. Also, since  $M^2 = mM$ , it follows that  $J = (R :_R R[1/m])$ , and so J is an ideal of Q(R). Let  $m^*$  denote the image of m in R/J. Then  $m^*$  is a nonzerodivisor in the one-dimensional ring R/J. Hence  $Q(R)/J = R[1/m]/J = (R/J)[1/m^*] = Q(R/J)$ . By (1),  $\overline{R} = R_{\infty}$ , and so it follows that  $R_{\infty}/J$  is the integral closure of R/J in Q(R/J) = Q(R)/J.

By Proposition 3.11(4),  $R_{\infty}$  has at most three maximal ideals if R is a quadratic ring. In the next theorem we describe the quadratic rings R for which  $R_{\infty}$  has exactly three maximal ideals.

**Theorem 3.13.** Let R be a local ring with a regular maximal ideal. Then the following statements are equivalent.

- (1) R is a quadratic ring such that  $R_{\infty}$  has three maximal ideals.
- (2) There is a ring S between R and Q(R) such that
  - (i) S has exactly 3 maximal ideals,
  - (ii) each maximal ideal of S is principal and regular with residue field isomorphic to  $\mathbb{F}_2$ , and
  - (iii) R = k + Jac S, where k is the prime subring of R.

Proof. (1)  $\Rightarrow$  (2) Suppose that R is a quadratic ring and  $R_{\infty}$  has three maximal ideals. Since R is local,  $R \neq R_{\infty}$ , and hence  $R \neq R_1$ . If  $R_1 = R_{\infty}$ , then, by Lemma 3.7(3),  $M = \operatorname{Jac} R_1$  and the three maximal ideals of  $R_1$  each have residue field isomorphic to  $\mathbb{F}_2$ . Moreover, by Proposition 3.11(4), each maximal ideal of  $R_1$  is regular and principal. Since  $(k+M)/M \subseteq R/M$  and R/M has only 2 elements (because R/M is a subfield of  $R_1/M \cong \mathbb{F}_2 \times \mathbb{F}_2 \times \mathbb{F}_2$ ), it must be that k+M=R. Therefore to verify (2) it remains to show that  $R_1=R_{\infty}$  and choose  $S=R_1$ .

Since R is a quadratic ring, each  $R_i$  with  $R_i \neq R_{i+1}$  is a quadratic ring by Proposition 3.11(2). Denote the maximal ideal of each such  $R_i$  by  $M_i$ . Since  $R_{\infty}$  has 3 maximal ideals, Proposition 3.11(3) implies that there is n > 0 such that  $R_{n-1} \subsetneq R_n = R_{\infty}$ . We show that n = 1. Suppose by way of contradiction that n > 1. By Proposition 3.11(2),  $R_{n-2}$  is a quadratic ring with  $R_{n-2} \subsetneq R_{n-1} \subsetneq R_n = R_{\infty}$ , and, by Lemma 3.7(2),  $R_n/M_{n-1} \cong \mathbb{F}_2 \times \mathbb{F}_2 \times \mathbb{F}_2$ . To

show that such a case is impossible, we may after relabeling assume that n=2. We show the assumption that R is a quadratic ring with  $R \subsetneq R_1 \subsetneq R_2 = R_{\infty}$  and the fact that  $R_2/M_1 \cong \mathbb{F}_2 \times \mathbb{F}_2 \times \mathbb{F}_2$  lead to a contradiction of the fact that  $R \neq R_1$ .

CLAIM: The dimension of the R/M-vector space  $R_1/R$  is 1.

Let A be an R-module such that  $R \subseteq A \subsetneq R_1$ . It suffices to show that R = A. Since  $R \subseteq R_1$  is a quadratic extension,  $R \subseteq R_1$  is an integral extension and A is a ring. Thus  $A \subseteq R_1$  is an integral extension. Since  $R_1$  is local, A is local with maximal ideal, say N, lying over M and under  $M_1$ . By Proposition 3.11(1) and (2), there is a nonzerodivisor  $m \in M$  such that  $M = mR_1$  and  $M_1 = mR_2$ . Then  $mR_1 = M \subseteq N \subseteq M_1 = mR_2$ , so that  $R_1 \subseteq m^{-1}N \subseteq R_2$ . Since  $R_1 \subseteq R_2$  is a quadratic extension,  $m^{-1}N$  is a ring, and hence  $N = mA_1$ , where  $A_1 = E(N) = m^{-1}N$ . Moreover  $R_1 \subseteq A_1 \subseteq R_2$ . To prove that R = A, we show that  $R_1 = A_1$ . Once this is established, we have that A has maximal ideal  $N = mA_1 = mR_1 = M$ . Since by Lemma 3.10(1), R and  $R_1$  have the same residue field R/M, so does A. Thus the fact that N = M implies that A = R + N = R, as claimed.

For the claim it remains to show that with A and  $A_1$  as above,  $R_1 = A_1$ . Since  $R_1/M_1 \subseteq A_1/M_1 \subseteq R_2/M_1 \cong \mathbb{F}_2 \times \mathbb{F}_2 \times \mathbb{F}_2$ , and the last ring is reduced,  $A_1/M_1$  is also reduced. Hence  $M_1 = \operatorname{Jac} A_1$ . Since every element of the ring  $\mathbb{F}_2 \times \mathbb{F}_2 \times \mathbb{F}_2$  is an idempotent, either  $R_1 = A_1$  or  $A_1/M_1$  is a decomposable ring. Suppose that  $R_1 \neq A_1$ . Then  $A_1/M_1$  is a decomposable ring, and hence  $A_1$  has more than one maximal ideal. Since  $R \subseteq R_{\infty}$  is a quadratic extension and  $R \subseteq A \subseteq A_1 \subseteq R_{\infty}$ , we have that  $A \subseteq A_1$  is also a quadratic extension with  $A_1$  not local. By Lemma 3.7(3) (applied to  $A \subseteq A_1$ ),  $N = \operatorname{Jac} A_1$ . We have established above that  $\operatorname{Jac} A_1 = M_1$ , and so  $N = M_1$ . Therefore, by Lemma 3.7(1),  $R_1 = R + M_1 = R + N \subseteq A$ , contrary to the choice of A with  $A \subseteq R_1$ . This shows  $A_1/M_1$  cannot be a decomposable ring, which forces  $R_1 = A_1$  and proves that  $R_1/R$  has dimension 1 as an R/M-vector space. This verifies the claim.

We show now that the claim implies a contradiction to our assumption that  $R \neq R_1$ . Let  $t \in R_1 \setminus R$ . Since  $R_1/R$  has dimension 1, we have  $R_1 = R + tR$ , and hence  $M = mR_1 = mR + mtR$ , so that as an R/M-vector space,  $M/M^2$  has dimension at most 2. By Lemma 3.7(4),  $M_1^2 \subseteq M$ . Also,  $M_1^2/M^2$  is a proper subspace of  $M/M^2$ , and as such has dimension at most 1. (We have  $M_1^2 \subseteq M$  since  $M_1 = mR_2$ ,  $M = mR_1$  and m is a nonzerodivisor.) However,  $M_1^2/M^2 = m^2R_2/m^2R_1 \cong R_2/R_1$ , so that  $R_2/R_1$  has dimension 1 as an R/M-vector space. There is an exact sequence of R/M-vector spaces,

$$0 \to R_1/M_1 \to R_2/M_1 \to R_2/R_1 \to 0.$$

The left and right vector spaces have dimension 1, while the middle vector space has dimension 3, since  $R_2/M_1 \cong \mathbb{F}_2 \times \mathbb{F}_2 \times \mathbb{F}_2$ . This contradiction implies that the assumption  $R \subsetneq R_1 \subsetneq R_2$  is impossible under the conditions in (1). Therefore  $R_1 = R_2$ , from which it follows that  $R_1 = R_{\infty}$ . This completes the proof that (1) implies (2).

 $(2)\Rightarrow (1)$  By (2),  $M=\operatorname{Jac} S$  is the product of the three principal maximal ideals of S, so  $\operatorname{Jac} S$  is a principal regular ideal of S. Hence  $\operatorname{Jac} S$  is a stable maximal ideal of R. Moreover  $S=E(M)=R_1=R_\infty$  and  $R_1/M\cong \mathbb{F}_2\times\mathbb{F}_2\times\mathbb{F}_2$ . By Lemma 2.8,  $R/M\subseteq R_1/M$  is a quadratic extension, and hence  $R\subseteq R_1=R_\infty$  is also a quadratic extension by Remark 2.7. Therefore R is a quadratic ring and  $R_\infty$  has three maximal ideals.

### 4. Characterizations of one-dimensional stable rings

In this section we work in the following setting.

Setting 4.1. Let R be a one-dimensional local ring having a regular maximal ideal M, let  $\overline{R}$  denote the integral closure of R in its total quotient ring Q(R), and let  $m \in M$  be a nonzerodivisor. Let  $J = \bigcap_{i>0} m^i R$ . As in Notation 2.3, we denote by  $\widetilde{R}$  and  $\widehat{R}$  the completions of R in the mR-adic and M-adic topologies, respectively. Both J and  $\widetilde{R}$  are independent of the choice of nonzerodivisor m in M. Moreover, since m is a nonzerdivisor, R/J is a one-dimensional ring with regular maximal ideal M/J.

We first characterize a one-dimensional stable local ring in terms of its integral closure. Recall that a ring A is a  $Dedekind\ ring$  if every regular ideal I of A is invertible.

**Theorem 4.2.** With Setting 4.1, the following statements are equivalent.

- (1) R is a stable ring.
- (2) R is a finitely stable ring with stable maximal ideal.
- (3) R is a quadratic ring such that  $\overline{R}$  has at most two maximal ideals.
- (4)  $R \subseteq \overline{R}$  is a quadratic extension and  $\overline{R}$  is a Dedekind ring with at most two maximal ideals.
- (5) R/J is a stable ring.

*Proof.*  $(1) \Rightarrow (2)$  This is clear.

- $(2) \Rightarrow (3)$  This follows from Lemma 3.8 and Theorem 3.9.
- $(3) \Rightarrow (4)$  By Corollary 3.12(1),  $\overline{R} = R_{\infty}$ , and so  $R \subseteq \overline{R}$  is a quadratic extension. By (3),  $\overline{R}$  has at most 2 maximal ideals, and Proposition 3.11(4) implies the maximal ideals of  $\overline{R}$  are principal regular ideals. Since  $\overline{R}$  has dimension one, every regular ideal of  $\overline{R}$  is a principal regular ideal, and hence  $\overline{R}$  is a Dedekind ring.
- $(4) \Rightarrow (1)$  Let I be a regular ideal of R. Since  $\overline{R}$  has Krull dimension 1 and at most 2 maximal ideals, both of which are principal and regular (they are regular because they lie over M), it follows that every regular ideal of  $\overline{R}$  contains a power of Jac  $\overline{R}$ . Since  $\overline{R}$  is a Dedekind ring with only finitely many maximal ideals, every regular ideal is principal. In particular,  $I\overline{R}$  is a principal regular ideal; that is,  $I\overline{R} = x\overline{R}$  for some nonzerodivisor  $x \in \overline{R}$ . An argument of Rush [36, proof of Theorem 2.2], which is also given explicitly in [34, Proposition 3.6], shows that since  $\overline{R}$  has at most two maximal ideals, x can be chosen an element

- of I. Therefore  $R \subseteq x^{-1}I \subseteq \overline{R}$ . Since  $R \subseteq \overline{R}$  is a quadratic extension,  $x^{-1}I$  is a ring, and hence I = xE(I). This shows that every regular ideal of R is stable.
- $(1) \Rightarrow (5)$  Let I be an ideal of R such that I/J is a regular ideal in R/J. Then I is an M-primary ideal R and hence a regular ideal in R. Since R is stable, Lemma 3.4(6) implies that  $I^2 = xI$ , for some  $x \in I$ . Since  $I^2$  is M-primary, we have  $J \subseteq I^2 \subseteq xR$ , and hence  $I^2/J = (I/J)(xR/J)$ . Moreover, since  $I^2 \subseteq xR$ , the ideal xR is M-primary, hence regular in R. Therefore xR/J is M/J-primary in R/J and hence regular in R/J. Lemma 3.4(4) implies then that I/J is a stable ideal in R/J, which proves that R/J is a stable ring.
- $(5)\Rightarrow (1)$  Let I be a regular ideal of R. Then I is M-primary, and so there is k>0 such that  $m^k\in I$ . Since the image of  $m^k$  in R/J is a nonzerodivisor, I/J is regular. By (5) and Lemma 3.4(6) there exists  $x\in I$  such that  $I^2/J=(xI+J)/J$ . Thus  $I^2=xI+J$ , so that since J is an ideal of R[1/m], we have  $R[1/m]=I^2R[1/m]=xIR[1/m]+JR[1/m]=xR[1/m]+J$ . Therefore there exist  $\ell\geq 0$ ,  $r\in R$  and  $j\in J$  such that  $1=x(r/m^\ell)+j$ . Thus  $(1-j)m^\ell\in xR$ , and, since 1-j is a unit in R,  $m^\ell\in xR$ . Therefore  $J\subseteq m^{k+\ell}R\subseteq xI$ , so that  $I^2=xI+J=xI$ . By Lemma 3.4(5), I is stable.

**Remark 4.3.** Rush [35, Theorem 2.4] has proved the equivalence of (1) and (4) of Theorem 4.2 in the case where R is a Noetherian ring. Our proof that (4) implies (1) follows Rush's argument.

From the theorem we deduce a characterization of one-dimensional quadratic rings that shows these are stable except in a very special case.

Corollary 4.4. A one-dimensional local ring R is a quadratic ring if and only if either R is a stable ring or  $R = k + \operatorname{Jac} S$ , where k is the prime subring of R and S is a Dedekind ring between R and Q(R) with exactly three maximal ideals, each with residue field isomorphic to  $\mathbb{F}_2$ .

Proof. Suppose that R is a quadratic ring that is not a stable ring. By Theorem 4.2,  $\overline{R}$  has more than two maximal ideals. By Corollary 3.12(1),  $\overline{R} = R_{\infty}$ , and so, by Proposition 3.11(4),  $R_{\infty}$  has exactly three maximal ideals. By Theorem 3.13, there exists a ring S between R and Q(R) having exactly 3 maximal ideals, all of which are principal and have residue field isomorphic to  $\mathbb{F}_2$ , such that  $R = k + \operatorname{Jac} S$ . As noted in the proof of Theorem 3.13, we may take S to be  $R_{\infty}$ . Since the maximal ideals of the one-dimensional ring  $\overline{R} = R_{\infty}$  are principal and regular,  $\overline{R}$  is a Dedekind ring. The converse is a consequence of Theorems 3.9 and 3.13.

In Theorem 4.5 we characterize the one-dimensional stable local rings with finite integral closure; Theorem 4.7 similarly characterizes stable rings without finite integral closure.

**Theorem 4.5.** With Setting 4.1, the following statements are equivalent.

- (1) R is a stable ring with finite integral closure.
- (2) R/J is a Bass ring.

## (3) $\widetilde{R}$ is a Bass ring.

*Proof.* (1)  $\Rightarrow$  (2) By Theorem 4.2, R/J is a stable ring. Since also R has finite integral closure, Corollary 3.12(2) implies that  $R_n = R_{\infty}$  for some n. By Proposition 3.11(7), J is the nilradical of R. Therefore R/J is a reduced one-dimensional stable ring. To show that R/J is a Bass ring, it remains by Theorem 2.11 to show that R/J is a Noetherian ring with finite integral closure. By Proposition 3.11(7), since  $R_n = R_{\infty}$ , J is the intersection of the minimal prime ideals  $Q_1, \ldots, Q_r$  of  $R_{\infty}$ , and, as noted in the proof of the proposition, these prime ideals are comaximal. Now  $R_{\infty}$  is a one-dimensional ring with principal maximal ideals (Proposition 3.11(4)), and so for each  $i, R_{\infty}/Q_i$  is a principal ideal domain. Since  $Q_1, \ldots, Q_r$  are pairwise comaximal, the ring  $R_{\infty}/J$  is the product of the principal ideal domains  $R_{\infty}/Q_i$ . Therefore  $R_{\infty}/J$ is a principal ideal ring. By Proposition 3.11(6),  $R_{\infty}/J$  is a finitely generated R/J-module, and so the Eakin-Nagata Theorem [24, Theorem 3.7, p. 18] implies that R/J is a Noetherian ring. Moreover, by Corollary 3.12(3),  $R_{\infty}/J$  is the integral closure of R/J in its total quotient ring, and so R/J has finite integral closure. Therefore R/J is a Bass ring.

 $(2)\Rightarrow (3)$  Let M denote the maximal ideal of R. Since S:=R/J is reduced and has finite normalization, Proposition 2.2(3) implies that the M/J-adic completion  $\widehat{S}$  of S is also reduced, and hence, again by Proposition 2.2(3),  $\widehat{S}$  has finite normalization. Since S has multiplicity  $\leq 2$ , so does  $\widehat{S}$ . Thus  $\widehat{S}$  is a Bass ring. To show then that  $\widehat{R}$  is a Bass ring, we need only prove that  $\widehat{R}\cong \widehat{S}$ . Since mR/J contains a power of M/J, the (mR/J)-adic completion of S=R/J coincides with the (M/J)-adic completion  $\widehat{S}$  of S. For each i>0,  $J\subseteq m^iR$ , so that  $S/m^iS\cong R/m^iR$  as R-algebras. Since these isomorphisms are natural,

$$\widehat{S} = \underline{\lim} S/m^i S \cong \underline{\lim} R/m^i R = \widetilde{R},$$

which completes the proof of (3).

 $(3) \Rightarrow (2)$  As noted in the proof of (2) implies (3), the mR/J-adic completion of R/J coincides with  $\widetilde{R}$ , so we may assume without loss of generality that J=0 and show that R is a Bass ring. Since J=0, we may identify R with its image in  $\widetilde{R}$ . Also, since  $\widetilde{R}$  is a reduced ring, so is R.

We claim that R is Noetherian. Since R is reduced and has Krull dimension 1, it suffices by Proposition 2.1(1) to show that M is finitely generated. To this end, observe that since  $\widetilde{R}$  is a Noetherian ring, the  $\widetilde{R}$ -module  $M\widetilde{R}/m\widetilde{R}$  is finitely generated. By Proposition 2.5(2),  $\widetilde{R}=R+m\widetilde{R}$ , and so  $M\widetilde{R}/m\widetilde{R}$  is finitely generated as an R-module. By Proposition 2.5(3),  $M/mR\cong M\widetilde{R}/m\widetilde{R}$  as R-modules, and hence M/mR is a finitely generated R-module. We conclude that M is a finitely generated ideal of R and hence that R is a Noetherian ring.

Since R is Noetherian, the M-adic and mR-adic topologies agree on R. Therefore the M-adic completion  $\widehat{R}$  of the Noetherian ring R is a Bass ring. In particular, R is an analytically unramified Noetherian local ring. Since  $\widehat{R}$  has multiplicity  $\leq 2$ , so does R, which proves that R is a Bass ring.

 $(2) \Rightarrow (1)$  By Theorem 4.2, R is stable ring, and by Corollary 3.12(3),  $R_{\infty}/J$  is the integral closure of R/J. Since R/J is a Bass ring,  $R_{\infty}/J$  is a finitely generated R-module. Hence  $R_{\infty}$  is a finitely generated R-module, so that R has finite integral closure by Corollary 3.12(1).

A simple example shows that the conditions of the theorem are not in general equivalent to the condition that R is a Bass ring. The example, as well as later examples in this section, are constructed using Nagata idealization, which was discussed in Remark 2.15.

**Example 4.6.** Let V be a DVR with quotient field F. Then  $R = V \star F$  is a one-dimensional stable local ring with regular maximal ideal [31, Lemma 3.3]. The ring R has finite integral closure (it is in fact integrally closed), and R is not a Bass ring because it is not reduced, nor even Noetherian.

We turn next to the characterization of stable rings without finite integral closure.

**Theorem 4.7.** With Setting 4.1, the following statements are equivalent.

- (1) R is a stable ring without finite integral closure.
- (2) R/J is a stable ring without finite integral closure.
- (3) There exists a nonzero prime ideal P of  $\widetilde{R}$  such that  $P^2 = 0$  and  $\widetilde{R}/P$  is a DVR.
- *Proof.* (1)  $\Rightarrow$  (2) Suppose that R is a stable ring without finite integral closure. By Theorem 4.2, R/J is a stable ring. By Corollary 3.12(1) and (3),  $\overline{R}/J$  is the integral closure of R/J. Thus, since R does not have finite integral closure, neither does R/J.
- $(2)\Rightarrow (3)$  As observed in the proof of (2) implies (3) of Theorem 4.5, the (mR/J)-adic completion of R/J coincides with  $\widetilde{R}$ . We assume without loss of generality that J=0 and R is a stable ring without finite integral closure. Corollary 3.12(2) implies that  $R_i\neq R_\infty$ , for all i>0. By Proposition 3.11(3) there exists a nonzerodivisor  $x\in M$  such that  $xR_\infty$  is the maximal ideal of  $R_\infty$  and  $R_\infty=R+xR_\infty$ . To simplify notation, let  $S=R_\infty$  and Q=Q(R).

Claim 1: The inclusion mapping  $R \to S$  lifts to a surjective ring homomorphism  $\widetilde{R} \to \widetilde{S}$ , where  $\widetilde{S}$  denotes the completion of S in the mR-adic completion of S.

Consider the exact sequence

$$\operatorname{Hom}_R(Q/R,Q/R) \overset{\delta}{\to} \operatorname{Hom}_R(Q/R,Q/S) \to \operatorname{Ext}^1_R(Q/R,S/R).$$

Since x is a nonzerodivisor in R, the ideal xR is M-primary. Since R has Krull dimension one, this implies that some power of x is contained in every principal regular ideal of R. Consequently, since S = R + xS, it follows that S = R + rS for all nonzerodivisors  $r \in R$ ; that is, S/R is divisible with respect to the set of nonzerodivisors of R. This property, along with the fact that Q = R

R[1/m] is a countably generated R-module, implies that  $\operatorname{Ext}^1_R(Q/R,S/R)=0$  [4, Theorem 1.1], and hence the map  $\delta$  is a surjection. By [23, Theorem 2.2, p. 13],  $\operatorname{Hom}_R(Q/R,Q/R)$  can be identified with  $\widetilde{R}$ , while  $\operatorname{Hom}_R(Q/R,Q/S)$  can be identified with  $\widetilde{S}$ . Since these identifications are natural,  $\widetilde{R}$  maps onto  $\widetilde{S}$ . As discussed in the proof of [23, Theorem 2.9, p. 21], this mapping is a ring homomorphism.

Claim 2: There is a prime ideal P of  $\widetilde{R}$  such that  $\widetilde{R}/P$  is a DVR.

Since x is a nonzerodivisor in S, Proposition 2.5(1) implies that x is also a nonzerodivisor in  $\widetilde{S}$ . Since S has principal maximal ideal xS,  $\widetilde{S}$  has principal maximal ideal  $x\widetilde{S}$ . Thus, since x is a nonzerodivisor in  $\widetilde{S}$ , the ideal  $L = \bigcap_k x^k \widetilde{S}$  has residue ring  $\widetilde{S}/L$  that is a DVR. Since Claim 1 implies that  $\widetilde{R}$  maps onto  $\widetilde{S}/L$ , there is a prime ideal P of  $\widetilde{R}$  such that  $\widetilde{R}/P$  is a DVR.

CLAIM 3: The ideal P in Claim 2 has the property that  $P^2 = 0$ .

The ideal P in Claim 2 is the kernel of the induced map  $\widetilde{R} \to \widetilde{S}/\bigcap_k x^k \widetilde{S}$ . With m as in Setting 4.1, since mR is M-primary,  $\bigcap_k x^k \widetilde{S} = \bigcap_k m^k \widetilde{S}$ . Viewing  $\widetilde{R}$  and  $\widetilde{S}$  as subrings of  $\prod_{i>0} R/m^i R$  and  $\prod_{i>0} S/m^i S$ , respectively, we have

$$P = \{ \langle r_i + m^i R \rangle \in \widetilde{R} : \langle r_i + m^i S \rangle \in \bigcap_k m^k \widetilde{S} \}.$$

Let  $p, q \in P$ , and write  $p = \langle a_i + m^i R \rangle$  and  $q = \langle b_i + m^i R \rangle$ . Since  $p, q \in P$ , there exist  $c_i, d_i, s_i, t_i \in S$  such that  $a_i = m^{i+1}c_i + m^i s_i$  and  $b_i = m^{i+1}d_i + m^i t_i$ . Thus  $a_i, b_i \in m^i S \cap R$ . Since S = R + mS,  $R \subseteq S \subseteq R[1/m]$  and  $R \subseteq S$  is a quadratic extension, it follows that  $(m^i S \cap R)^2 \subseteq m^i R$  for all i > 0 [31, Lemma 3.2]. Therefore  $a_i b_i \in m^i R$  for all i > 0, which shows that pq = 0, and hence  $P^2 = 0$ .

 $(3) \Rightarrow (1)$  Since  $\widetilde{R}/P$  is a DVR and  $P^2 = 0$ ,  $\widetilde{R}$  is a one-dimensional stable ring [31, Lemma 3.3]. We claim that this implies R is a stable ring. To this end, we first make an observation about the canonical homomorphism  $\lambda : R \to \widetilde{R}$ .

CLAIM: If  $z \in \widetilde{R}$  is such that  $\lambda^{-1}(z\widetilde{R})$  is a regular ideal of R, then the principal ideal  $z\widetilde{R}$  is generated by the image of an element of  $\lambda^{-1}(z\widetilde{R})$ .

Let  $z \in \widetilde{R}$  be such that  $\lambda^{-1}(z\widetilde{R})$  is a regular ideal of R, and let x be a nonzerodivisor in  $\lambda^{-1}(z\widetilde{R})$ . Write  $\lambda(x) = zw$  for some  $w \in \widetilde{R}$ . If w is a unit in  $\widetilde{R}$ , then  $x\widetilde{R} = z\widetilde{R}$ , as claimed. Suppose that w is not a unit in  $\widetilde{R}$ . Since x is a nonzerodivisor in R, Proposition 2.5(2) gives that  $\widetilde{R} = \lambda(R) + x\widetilde{R}$ . Therefore  $z \in \lambda(R) + x\widetilde{R} = \lambda(R) + zw\widetilde{R}$ , and so, for some  $t \in \widetilde{R}$  and  $y \in R$ ,  $z(1-wt) = z - zwt = \lambda(y)$ . Since w is not a unit and  $\widetilde{R}$  is a local ring, 1-wt is a unit in  $\widetilde{R}$ . Since  $z(1-wt) = \lambda(y)$ , we have  $z\widetilde{R} = y\widetilde{R}$ . This shows that in all cases  $z\widetilde{R}$  is generated as a principal ideal by the image of an element of R. This proves the claim.

We use the claim now to verify that R is stable. Let I be a regular ideal of R. Since  $\widetilde{R}$  is a stable local ring, Lemma 3.4(6) implies that there exists  $z \in I\widetilde{R}$  such that  $I^2\widetilde{R} = zI\widetilde{R}$ . Since  $I^2\widetilde{R} \subseteq z\widetilde{R}$ , we have  $I^2 \subseteq \lambda^{-1}(z\widetilde{R})$ . Since  $I^2$  is a regular

ideal of R, the claim implies that there exists  $x \in \lambda^{-1}(z\widetilde{R})$  such that  $z\widetilde{R} = x\widetilde{R}$ . By Proposition 2.5(3),  $\lambda^{-1}(I\widetilde{R}) = I$ , and so  $x \in \lambda^{-1}(z\widetilde{R}) \subseteq \lambda^{-1}(I\widetilde{R}) = I$ . Furthermore,  $I^2\widetilde{R} = xI\widetilde{R}$ , so again by Proposition 2.5(3),  $I^2 = \lambda^{-1}(I^2\widetilde{R}) = \lambda^{-1}(xI\widetilde{R}) = xI$ . By Lemma 3.4(4), I is a stable ideal of R, which proves that R is stable. Finally, if R has finite integral closure, then, by Theorem 4.5,  $\widetilde{R}$  is a Bass ring. In particular  $\widetilde{R}$  is reduced, contrary to (3). Thus R does not have finite integral closure.

In the case where R is separated in the mR-adic topology (e.g., if R is Noetherian), we obtain the following classification of one-dimensional stable local rings.

**Corollary 4.8.** With Setting 4.1 and J=0, the ring R is stable if and only if R is a Bass ring or there is a nonzero prime ideal P of  $\widetilde{R}$  such that  $\widetilde{R}/P$  is a DVR and  $P^2=0$ .

*Proof.* Apply Theorems 4.5 and 4.7.

In particular, if R is complete in the mR-adic topology, then there are only two classes of (one-dimensional) stable rings: Bass rings and rings having a square zero prime ideal whose residue ring is a DVR. Thus, in the case in which R is mR-adically complete, R is stable and reduced if and only if R is a Bass ring. More generally, we have the following corollary.

**Corollary 4.9.** With Setting 4.1, the ring R is a reduced stable ring if and only if R is either a Bass ring or a bad stable domain.

Proof. Suppose that R is a reduced stable ring. If R has finite normalization, then by Corollary 3.12(2),  $R_i = R_{\infty}$ , for some i, and so, by Proposition 3.11(7), J = 0. In this case Theorem 4.5 implies that R is a Bass ring. Otherwise, suppose that R does not have finite normalization. By Corollary 3.12(2),  $R_i \neq R_{\infty}$ , for all i > 0, and so Proposition 3.11(3) implies  $R_{\infty}$  is local. In this case Proposition 3.11(7) implies that R has a unique minimal prime ideal, and hence, since R is reduced, R is a domain. By Theorem 4.7 there exists a nonzero prime ideal P of R such that R/P is a DVR and  $P^2 = 0$ . By Theorem 2.13, R is a bad stable domain. The converse is clear in light of Theorem 2.11.

The next two examples illustrate some complications with characterizing the non-reduced, non-complete one-dimensional stable local rings R such that  $\widetilde{R}$  has a prime ideal P with  $P^2=0$  and  $\widetilde{R}/P$  is a DVR. Such a ring R has a nonzero prime ideal Q such that  $Q^2=0$ . The first example shows that even if R is Noetherian, R/Q need not be a DVR. The second example shows that the maximal ideal of R can be two-generated but the ring R not be Noetherian.

**Example 4.10.** A local Cohen-Macaulay stable ring R having a nonzero prime ideal P such that  $P^2 = 0$  and R/P is a bad stable domain (and hence R/P is not a DVR). Let (A, M) be a bad Noetherian stable domain, let  $R = A \star (A/M)$  be the Nagata idealization of the A-module A/M (cf. Remark 2.15) and let

 $P=0\star(A/M)$ . Then  $P^2=0$  and  $R/P\cong A$  is a bad stable domain. Since A/M is a finitely generated A-module, R is a one-dimensional local Cohen-Macaulay ring. We claim that R is stable. Let m be a nonzerodivisor in M. Then  $\widetilde{R}=\widetilde{A}\star(\widetilde{A}/M\widetilde{A})$ , where  $\widetilde{R}$  is the mR-adic completion of R and  $\widetilde{A}$  is the mA-adic completion of A. Since A is a bad stable domain, Theorem 2.13 implies there is a nonzero prime ideal Q of  $\widetilde{A}$  such that  $\widetilde{A}/Q$  is a DVR and  $Q^2=0$ . Now  $L:=Q\star(\widetilde{A}/M\widetilde{A})$  is a prime ideal of  $\widetilde{R}$ , and  $L^2=0$  since  $Q^2=0$  and  $Q\cdot(\widetilde{A}/M\widetilde{A})=0$ . Also,  $\widetilde{R}/L\cong\widetilde{A}/Q$ , so that  $\widetilde{R}/L$  is a DVR. By Corollary 4.8, R is a stable ring. If A is chosen instead to be a non-Noetherian bad stable domain, then R is a non-Noetherian stable ring.

**Example 4.11.** A non-Noetherian separated stable local ring (R, M) of Krull dimension one whose maximal ideal can be generated by two elements. Let k be a field, let X be an indeterminate over k, and let  $V = k[X]_{(X)}$ . The (X)-adic completion of V is  $\widetilde{V} = k[[X]]$ . Define  $R = V \star \widetilde{V}$ . Since  $\widetilde{V}$  is not a finitely generated V-module, R is not a Noetherian ring. However, the maximal ideal  $M = XV \star \widetilde{V}$  of R is generated by two elements, (X,0) and (0,1), the first a nonzerodivisor in R and the second a zerodivisor. Since the ring R has a prime ideal whose square is zero and whose residue ring is a DVR, R is a stable ring [31, Lemma 3.3]. Moreover  $\bigcap_i M^i = \bigcap_i (m^i V \star m^i \widetilde{V}) = 0$ , and so R is separated in the M-adic topology.

### 5. The two-generator property

In this section we consider the two-generator property in our context. By a proper power of the ideal I, we mean an ideal of the form  $I^k$ , where k > 1. Sally [37, Proposition 1] has shown that if a proper power of a regular ideal (which, a priori, is not necessarily finitely generated) is two-generated, then so is every power of the ideal, including the ideal itself. We use this in the next proposition to connect the property of being two-generated with that of being stable.

**Proposition 5.1.** Let I be a regular ideal of the local ring R. Then, for some n > 1,  $I^n$  is two-generated if and only if I is two-generated with  $I^2 = aI$  for some  $a \in I$ .

Proof. Suppose that a proper power of I is a two-generated ideal. By the result of Sally discussed above, I is a two-generated ideal and we may write I=(x,y)R and  $I^2=(x^2,xy,y^2)R$ . The argument that follows is based on the proof of [40, Theorem 3.4]. Since  $I^2$  is two-generated, Nakayama's lemma implies that  $I^2=(x^2,xy)R=xI$ ,  $I^2=(xy,y^2)R=yI$  or  $I^2=(x^2,y^2)R$ . In the first two cases, we have that  $I^2=aI$  for some  $a\in I$ . Consider the last case,  $I^2=(x^2,y^2)R$ . Then  $xy=rx^2+sy^2$  for some  $r,s\in R$ . If r is a unit, then  $x^2\in(xy,y^2)R=yI$ , so that  $I^2=yI$ . Similarly, if s is a unit, then  $I^2=xI$ . If neither r nor s is a unit, then  $xy\in MI^2$ , with M the maximal ideal of R. Thus  $I^2=(x^2,y^2)R=(x^2-xy,xy-y^2)R+MI^2$ , and by Nakayama's lemma,  $I^2=(x^2-xy,xy-y^2)R=(x-y)I$ . This proves that in all cases  $I^2=aI$  for some  $a\in R$ . The converse of the proposition is clear.

**Corollary 5.2.** Let R be a reduced one-dimensional local ring with regular maximal ideal M. If, for some n > 1,  $M^n$  is two-generated, then every ideal of R is two-generated.

Proof. Suppose  $M^n$  is two-generated for some n > 1. By the result of Sally [37, Proposition 1] discussed above, M is two-generated. Also, by Proposition 5.1,  $M^2 = mM$  for some  $m \in M$ . Since R is reduced and one-dimensional, Proposition 2.1(1) implies that R is a Noetherian ring. Also, since  $M^2 = mM$  and M is two-generated, Proposition 2.2(1) implies that R has multiplicity at most 2. By Proposition 2.2(2) every ideal of R is two-generated.

Rush [36, Proposition 2.5] has shown that if R is a local ring for which every finitely generated regular ideal is two-generated, then R is finitely stable. Proposition 5.1 leads to a different proof of this result in slightly stronger form.

Corollary 5.3. If every finitely generated regular ideal of the local ring R has a power that is two-generated, then R is a finitely stable ring.

*Proof.* Let I be a finitely generated regular ideal of R. By assumption  $I^2$  has a power that is two-generated. Hence I has a proper power that is two-generated. By Proposition 5.1,  $I^2 = aI$  for some  $a \in I$ , and so I is stable by Lemma 3.4(5).

П

**Theorem 5.4.** With Setting 4.1, the following statements are equivalent.

- (1)  $M^n$  is two-generated, for some  $n \geq 2$ .
- (2)  $\tilde{R}$  is a Noetherian local ring of multiplicity at most 2.
- (3) Every regular ideal of R is two-generated.

Proof. (1)  $\Rightarrow$  (2) By Proposition 5.1, M is two-generated and  $M^2 = mM$  for some  $m \in M$ . Thus the maximal ideal  $M\widetilde{R}$  of  $\widetilde{R}$  is two-generated and  $M^2\widetilde{R} = mM^2\widetilde{R}$ . Since  $M^2 = mM$ , Remark 2.4 implies  $\widetilde{R} = \widehat{R}$ . Since  $\bigcap_i M^i\widehat{R} = 0$  and  $\widehat{R}$  is  $M\widehat{R}$ -adically complete with finitely generated maximal ideal, the ring  $\widehat{R}$  is Noetherian [8, Theorem 3]. By Proposition 2.5(1), m is a nonzerodivisor in  $\widehat{R}$  with  $M^2\widehat{R} = mM\widehat{R}$ , and so by Proposition 2.2(1) the multiplicity of  $\widehat{R} = \widetilde{R}$  is at most 2.

 $(2)\Rightarrow (3)$  Let I be a regular ideal of R. By Proposition 2.5(1),  $I\widetilde{R}$  is a regular ideal of  $\widetilde{R}$ . Since  $\widetilde{R}$  has Krull dimension 1 and multiplicity at most 2, Proposition 2.2(2) implies that every ideal of  $\widetilde{R}$  is two-generated. Since I is regular,  $m^k\in I$  for some k>0, and so  $I\widetilde{R}/m^{k+1}\widetilde{R}\cong I/m^{k+1}R$  by Proposition 2.5(3). By Proposition 2.5(2),  $\widetilde{R}=\lambda(R)+m^{k+1}\widetilde{R}$ , where  $\lambda:R\to\widetilde{R}$  is the canonical embedding. Since  $I\widetilde{R}/m^{k+1}\widetilde{R}$  can be generated by two elements as an  $\widetilde{R}$ -module,  $I/m^{k+1}R$  can be generated by two elements as an R-module. Therefore  $I=(x,y,m^{k+1})R$  for some  $x,y\in I$ . Since  $m^{k+1}\in MI$ , Nakayama's lemma implies that I can be generated by two elements.

$(3) \Rightarrow (1)$ This is clear.	
--------------------------------------	--

**Remark 5.5.** Example 4.11 is a non-Noetherian ring satisfying the conditions of Theorem 5.4.

Restricting to Noetherian rings, we have a stronger characterization.

**Theorem 5.6.** Let R be a Cohen-Macaulay ring with maximal ideal M. Then these statements are equivalent.

- (1)  $M^n$  is two-generated, for some  $n \geq 2$ .
- (2) R has Krull dimension 1 and multiplicity at most 2.
- (3) R is one of the following:
  - (3a) a Bass ring;
  - (3b) a bad stable domain; or
  - (3c) a ring containing a nonzero principal prime ideal P such that  $P^2 = 0$  and R/P is a DVR.

*Proof.* (1)  $\Rightarrow$  (2) Proposition 5.1 implies that every power of M is two-generated, and hence the Hilbert polynomial of M is constant. Since the degree of this polynomial is one less than the Krull dimension of R, the Hilbert-Samuel theorem [38, p. 4] implies that R has Krull dimension one. By Theorem 5.4,  $\widetilde{R}$  is a stable ring of multiplicity at most 2. Since R is Noetherian and has Krull dimension one, Remark 2.4 implies  $\widetilde{R} = \widehat{R}$ . Therefore, since  $\widehat{R}$  has multiplicity 2, so does R

 $(2) \Rightarrow (3)$  By Proposition 2.2(2) every ideal of R is two-generated, and hence R is a stable ring by Corollary 5.3. If  $\overline{R}$  is a finitely generated R-module, then R is a Bass ring by Theorem 4.5 and R satisfies (3a).

Suppose that  $\overline{R}$  is not a finitely generated R-module. Since R is one-dimensional and Noetherian, Remark 2.4 implies  $\widehat{R} = \widetilde{R}$ . By Theorem 4.7 there exists a nonzero prime ideal L of  $\widehat{R}$  such that  $L^2 = 0$  and  $\widehat{R}/L$  is a DVR. Viewing R as a subring of  $\widehat{R}$ , we consider two cases.

Case 1:  $L \cap R = 0$ .

In this case R is a domain. By Theorem 2.13, R is a bad stable domain and hence (3b) holds for R.

Case 2:  $P := L \cap R$  is a nonzero prime ideal of R.

We show that (3c) holds. Since  $L^2=0$ , we have that  $P^2=0$ . Since R is stable, Lemma 3.4(6) implies there exists  $m\in M$  such that  $M^2=mM$ . We claim that  $P\not\subseteq mR$ . Suppose to the contrary that  $P\subseteq mR$ . Since P is a prime ideal and  $m\not\in P$ , it follows that P=mP. Hence  $P\subseteq \bigcap_i m^iR=\bigcap_i M^i=0$ , a contradiction. Thus  $P\not\subseteq mR$ .

Next we claim that  $M^2 \subsetneq mR$ . Indeed, if  $M^2 = mR$ , then, since m is a nonzerodivisor, M is an invertible, hence principal, ideal of R. Thus R is a regular local ring, hence a domain. However,  $P^2 = 0$  and P is nonzero, so this contradiction shows that  $M^2 \subsetneq mR$ .

Now since  $M^2 \subsetneq mR$ ,  $P \not\subseteq mR$  and  $M/M^2$  is a vector space of dimension at most 2 over R/M, it follows that M=mR+P. Thus R/P is a DVR. Since M=mR+P and  $P^2=0$ , we have  $P/MP=P/(mP+P^2)=P/mP=P/(mR\cap P)\cong (P+mR)/mR=M/mR$ . Since  $M/M^2$  has dimension at most 2 as a vector space over R/M and  $M^2 \subsetneq mR$ , it follows that P/MP has dimension at most one as an R/M-vector space. By Nakayama's lemma, P is a principal ideal, and so (3c) is satisfied by R.

 $(3) \Rightarrow (1)$  It is clear that if R is a Bass ring, then every power of M is two-generated. Similarly, if R is a bad stable domain with multiplicity  $\leq 2$ , then, by Proposition 2.2(2), every power of M is two-generated. Finally, suppose there is a nonzero principal prime ideal P of R with R/P a DVR and  $P^2 = 0$ . Since R/P is a DVR and  $P^2 = 0$ , it follows that  $M^2 = mM$  for some  $m \in M$  [31, Lemma 3.3]. Since P is a principal ideal and R/P is a DVR, M is two-generated. Thus  $M^2 = mM$  is also two-generated, which verifies (1).

Greither [13, Theorem 2.1] has shown using multiplicity theory that a reduced Noetherian local ring R is a Bass ring if and only if  $\overline{R}$  can be generated by two elements as an R-module. Theorem 5.8, which uses a different method of proof, generalizes this result to rings that a priori need not be Noetherian.

**Lemma 5.7.** If  $R \subseteq S$  is an extension of rings such that S can be generated by two elements as an R-module, then  $R \subseteq S$  is a quadratic extension.

Proof. By [34, Lemma 3.1], it suffices to show that  $R_M \subseteq S_M$  is a quadratic extension for each maximal ideal M of R. Thus we may assume without loss of generality that R is a local ring. Since S can be generated by two elements, Nakayama's lemma implies that S = R + xR for some  $x \in S$ . Let  $s, t \in S$ . We claim that  $st \in sR + tR + R$ . Write s = a + xb and t = c + xd for some  $a, b, c, d \in R$ . Since S = R + xR and S is a ring,  $x^2 \in R + xR$ . Hence

$$st = ac + bcx + adx + bdx^{2} \in R + bxR + dxR + dx^{2}R$$
  
$$\subseteq R + bxR + dxR = R + sR + tR,$$

This proves  $R \subseteq S$  is a quadratic extension.

Recall that a generalized local ring is a local ring with finitely generated maximal ideal M such that  $\bigcap_i M^i = 0$ ; cf. [8].

**Theorem 5.8.** The following are equivalent for a one-dimensional local ring R with regular maximal ideal M.

- (1) R is a Bass ring.
- (2) R is a generalized local ring whose integral closure  $\overline{R}$  can be generated by two elements as an R-module.
- (3)  $\overline{R}$  is a Noetherian ring that can be generated by two elements as an Rmodule.

- *Proof.* (1)  $\Rightarrow$  (2) Suppose R is a Bass ring. Then R is a generalized local ring. Since  $\overline{R}$  is a finitely generated R-submodule of Q(R), there is a nonzerodivisor  $r \in R$  such that  $r\overline{R}$  is an ideal of R. Since R is a Bass ring,  $r\overline{R}$  is a two-generated ideal of R. Since r is a nonzerodivisor,  $\overline{R}$  and  $r\overline{R}$  are isomorphic as R-modules. Thus  $\overline{R}$  can be generated by two elements as an R-module.
- $(2)\Rightarrow (3)$  Since  $\overline{R}$  is a finitely generated R-submodule of Q(R) and M is a regular ideal, the ideal  $(R:\overline{R})$  contains a nonzerodivisor  $m\in M$ . Also, since R is a generalized local ring,  $\bigcap_i m^i R = \bigcap_i M^i = 0$ . We claim that R is reduced. The argument is essentially that of [24, pp. 263-264]. Let r be a nilpotent element of R. For each i>0,  $r/m^i\in \overline{R}$ . Since  $m\in (R:\overline{R})$ , we have  $m(r/m^i)\in R$  for all i>0. Thus  $r\in \bigcap_{i>1} m^{i-1}R=0$ , proving that r=0 and R is reduced. Since the maximal ideal of R is finitely generated, R is a Noetherian ring by Proposition 2.1(1). Therefore, since R is a reduced one-dimensional Noetherian local ring with regular maximal ideal,  $\overline{R}$  is a Noetherian ring [1, Theorem 3].
- $(3)\Rightarrow (1)$  Given (3), to prove that R is a Bass ring it suffices by Corollary 4.9 to show that R is a reduced stable ring. For this it is enough by Theorem 4.2 to prove that  $R\subseteq \overline{R}$  is a quadratic extension and  $\overline{R}$  is a reduced Dedekind ring with at most two maximal ideals. Since the R-module  $\overline{R}$  can be generated by two elements,  $\overline{R}/(\operatorname{Jac}\overline{R})$  has dimension at most 2 as an R/M-vector space. Thus  $\overline{R}$  has at most 2 maximal ideals. Since R is a one-dimensional local Noetherian ring with regular maximal ideal and  $\overline{R}$  is a Noetherian ring,  $\overline{R}$  is a finite product of Dedekind domains [1, Theorem 3]. Hence  $\overline{R}$  is a reduced Dedekind ring. By Lemma 5.7,  $R\subseteq \overline{R}$  is a quadratic extension, and so the proof is complete.  $\square$

ACKNOWLEDGMENT. I am grateful to the referee for a careful reading of the paper and many suggestions that have improved the presentation.

### References

- [1] D. D. Anderson and J. Huckaba, The integral closure of a Noetherian ring. II. Comm. Algebra 6 (1978), no. 7, 639–657.
- [2] D. D. Anderson, J. Huckaba and I. Papick, A note on stable domains, *Houston J. Math.* **13** (1987), no. 1, 13–17.
- [3] D. D. Anderson, J. Matijevic and W. Nichols, The Krull intersection theorem II, *Pacific J. Math.* **66** (1976), 15–22.
- [4] L. Angeleri Hügel, D. Herbera and J. Trlifaj, Divisible modules and localization. J. Algebra 294 (2005), no. 2, 519–551.
- [5] Y. Akizuki, Einige Bemerkungen über primäre Integrit?tsbereiche mit Teilerkettensatz, *Proc. Phys. Math. Soc. Japan* 17 (1935), 327–336.
- [6] H. Bass, On the ubiquity of Gorenstein rings, Math Z. 82 (1963), 8–28.

- [7] N. Bourbaki, *Commutative Algebra*, *Chapters 1–7*. Translated from the French. Reprint of the 1989 English translation. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 1998.
- [8] I. S. Cohen, On the structure and ideal theory of complete local rings. *Trans. Amer. Math. Soc.* **59**, (1946), 54–106.
- [9] J. A. Drozd and V. V. Kiričenko, On quasi-Bass orders, Math. USSR Izv. 6 (1972), 323–365.
- [10] S. Gabelli and M. Roitman, On finitely stable domains, preprint.
- [11] R. Gilmer, W. Heinzer and M. Roitman, Finite generation of powers of ideals. *Proc. Amer. Math. Soc.* **127** (1999), no. 11, 3141–3151.
- [12] H. P. Goeters, Noetherian stable domains, J. Algebra  ${\bf 202}$  (1998), no. 1, 343-356.
- [13] C. Greither, On the two generator problem for the ideals of a one-dimensional ring. J. Pure Appl. Algebra 24 (1982), no. 3, 265–276.
- [14] D. Handelman, Propinquity of one-dimensional Gorenstein rings, J. Pure Appl. Algebra 24 (1982), no. 2, 145–150.
- [15] W. Heinzer, D. Lantz and K. Shah, The Ratliff-Rush ideals in a Noetherian ring, *Comm. Algebra* **20** (1992), 491–622.
- [16] M. Knebusch and D. Zhang, Manis valuations and Prüfer extensions. I. A new chapter in commutative algebra. Lecture Notes in Mathematics, 1791. Springer-Verlag, Berlin, 2002.
- [17] T. Y. Lam, Exercises in Modules and Rings, Springer, 2007.
- [18] J. Lambek, Lectures on rings and modules. Second edition. Chelsea Publishing Co., New York, 1976.
- [19] M. Larsen and P. McCarthy, Multiplicative theory of ideals. Academic Press, 1971.
- [20] C. Lech, A method for constructing bad Noetherian local rings, Algebra, algebraic topology and their interactions (Stockholm, 1983), 241–247, Lecture Notes in Math., 1183, Springer, Berlin, 1986.
- [21] J. Lipman, Stable ideals and Arf rings, Amer. J. Math. 93 (1971) 649–685.
- [22] E. Matlis, *Torsion-free modules*. Chicago Lectures in Mathematics. The University of Chicago Press, Chicago-London, 1972.
- [23] E. Matlis, 1-dimensional Cohen-Macaulay rings. Lecture Notes in Mathematics, Vol. 327. Springer-Verlag, Berlin-New York, 1973.

- [24] H. Matsumura, Commutative Ring Theory, Cambridge University Press, 1986.
- [25] M. Nagata, Local rings. Interscience Tracts in Pure and Applied Mathematics, No. 13, John Wiley & Sons, New York-London, 1962.
- [26] B. Olberding, Globalizing local properties of Prüfer domains, J. Algebra 205 (1998), no. 2, 480–504.
- [27] B. Olberding, On the classification of stable domains, J. Algebra 243 (2001), no. 1, 177–197.
- [28] B. Olberding, Stability of ideals and its applications. *Ideal theoretic methods in commutative algebra (Columbia, MO, 1999)*, 319–341, Lecture Notes in Pure and Appl. Math., 220, Dekker, New York, 2001.
- [29] B. Olberding, On the structure of stable domains, Comm. Algebra 30 (2002), no. 2, 877–895.
- [30] B. Olberding, Noetherian rings without finite normalization, *Progress in commutative algebra 2*, 171–203, Walter de Gruyter, Berlin, 2012.
- [31] B. Olberding, Generic formal fibers and analytically ramified stable rings, Nagoya Math. J. 211 (2013), 109–135.
- [32] B. Olberding, One-dimensional bad Noetherian domains, *Trans. Amer. Math. Soc.* **366** (2014), 4067–4095.
- [33] B. Olberding, Prescribed subintegral extensions of local Noetherian domains, J. Pure Appl. Algebra, 218 (2014), 506–521.
- [34] B. Olberding, Finitely stable rings, in Commutative Algebra: Recent advances in Commutative rings, integer-valued polynomials, and polynomial functions, Springer, 2014, 269–291.
- [35] D. Rush, Rings with two-generated ideals, J. Pure Appl. Algebra 73 (1991), no. 3, 257–275.
- [36] D. Rush, Two-generated ideals and representations of abelian groups over valuation rings, *J. Algebra* **177** (1995), no. 1, 77–101.
- [37] J. D. Sally, On the number of generators of powers of an ideal, *Proc. Amer. Math. Soc.*, **53** (1975), 24–26.
- [38] J. D. Sally, Numbers of generators of ideals in local rings. Marcel Dekker, Inc., New York-Basel, 1978.
- [39] J. D. Sally and W. Vasconcelos, Stable rings and a problem of Bass, Bull. Amer. Math. Soc. **79** (1973), 575–576.
- [40] J. D. Sally and W. Vasconcelos, Stable rings, J. Pure Appl. Algebra 4 (1974), 319–336.

- [41] I. Swanson and C. Huneke, *Integral closure of ideals, rings, and modules*. London Mathematical Society Lecture Note Series, 336. Cambridge University Press, Cambridge, 2006.
- [42] P. Zanardo, The class semigroup of local one-dimensional domains, *J. Pure Appl. Algebra* **212** (2008), no. 10, 2259-?2270.
- [43] P. Zanardo, Algebraic entropy of endomorphisms over local one-dimensional domains, *J. Algebra Appl.* 8 (2009), no. 6, 759?-777.