

HERMITIAN SYMMETRIC SPACE, FLAT BUNDLE AND HOLOMORPHICITY CRITERION

HASSAN AZAD, INDRANIL BISWAS, C. S. RAJAN, AND SHEHRYAR SIKANDER

ABSTRACT. Let $K \backslash G$ be an irreducible Hermitian symmetric space of noncompact type and $\Gamma \subset G$ a closed torsionfree discrete subgroup. Let X be a compact Kähler manifold and $\rho : \pi_1(X, x_0) \rightarrow \Gamma$ a homomorphism such that the adjoint action of $\rho(\pi_1(X, x_0))$ on $\text{Lie}(G)$ is completely reducible. A theorem of Corlette associates to ρ a harmonic map $X \rightarrow K \backslash G / \Gamma$. We give a criterion for this harmonic map to be holomorphic. We also give a criterion for it to be anti-holomorphic.

1. INTRODUCTION

Let G be a noncompact simple Lie group of adjoint type and $K \subset G$ a maximal compact subgroup, such that $K \backslash G$ is an irreducible Hermitian symmetric space of noncompact type. The Lie algebra of G will be denoted by \mathfrak{g} . Let Γ be a closed torsionfree discrete subgroup of G . Take a compact connected Kähler manifold X ; fix a base point $x_0 \in X$. Let

$$\rho : \pi_1(X, x_0) \rightarrow \Gamma$$

be a homomorphism such that the adjoint action of $\rho(\pi_1(X, x_0))$ on \mathfrak{g} is completely reducible. This ρ produces a C^∞ principal G -bundle $E_G \rightarrow X$ equipped with a flat connection D . A reduction of structure group of E_G to K is given by a map $X \rightarrow K \backslash G / \Gamma$. Note that $K \backslash G / \Gamma$ is a Kähler manifold; it need not be compact.

A theorem of Corlette says that there is a C^∞ reduction of structure group of E_G to K such that corresponding map $H_D : X \rightarrow K \backslash G / \Gamma$ is harmonic [Co]. Our aim here is to address the following:

- When is H_D holomorphic?
- When is H_D anti-holomorphic?

Let $H : X \rightarrow K \backslash G / \Gamma$ be a C^∞ map giving a reduction of structure group of E_G to K . We give a criterion under which H is holomorphic or anti-holomorphic (see Theorem 4.3). Since a holomorphic or anti-holomorphic map between Kähler manifolds is harmonic, Theorem 4.3 gives criterion for H_D to be holomorphic or anti-holomorphic (see Corollary 4.4).

2010 *Mathematics Subject Classification.* 32M15, 32L05, 53C55.

Key words and phrases. Hermitian symmetric space, flat bundle, Hermitian structure, holomorphicity.

2. HARMONIC MAP TO G/K

A Lie group is called simple if its Lie algebra is so [He, page 131]. Let G be a connected real noncompact simple Lie group whose center is trivial. It is known that if K_1 and K_2 are two maximal compact subgroups of G , then there is an element $g \in G$ such that $K_1 = g^{-1}K_2g$ [He, page 256, Theorem 2.2(ii)]. In particular, any two maximal compact subgroups of G are isomorphic. Assume that G satisfies the condition that the dimension of the center of a maximal compact subgroup K of it is positive. This in fact implies that the center of K is isomorphic to $S^1 = \mathbb{R}/\mathbb{Z}$ [He, page 382, Proposition 6.2].

Fix a maximal compact subgroup $K \subset G$. Consider the left-translation action of K on G . The above conditions on G imply that the corresponding quotient $K \backslash G$ is an irreducible Hermitian symmetric space of noncompact type. Conversely, given any irreducible Hermitian symmetric space of noncompact type, there is a group G of the above type such that $K \backslash G$ is isometrically isomorphic to it [He, page 381, Theorem 6.1(i)]. In fact, this gives a bijection between the isomorphism classes of Lie groups of the above type and the holomorphic isometry classes of irreducible Hermitian symmetric spaces [He, page 381, Theorem 6.1(i)]. The Lie algebra of G will be denoted by \mathfrak{g} .

Fix a closed torsionfree discrete subgroup $\Gamma \subset G$. Therefore, the two-sided quotient

$$M_\Gamma := K \backslash G / \Gamma \tag{2.1}$$

is a connected Kähler manifold.

Let X be a compact connected orientable real manifold. Fix a base point $x_0 \in X$. Let

$$\beta : \tilde{X} \longrightarrow X \tag{2.2}$$

be the universal cover of X associated to x_0 . The right-action of the fundamental group $\pi_1(X, x_0)$ on \tilde{X} will be denoted by “ \cdot ”. Let

$$\rho : \pi_1(X, x_0) \longrightarrow \Gamma \tag{2.3}$$

be a homomorphism such that the adjoint action of $\rho(\pi_1(X, x_0)) \subset G$ on the Lie algebra \mathfrak{g} is completely reducible. Associated to ρ we have a principal G -bundle $E_G \longrightarrow X$ equipped with a flat connection D . We note that E_G is the quotient of $\tilde{X} \times G$ (see (2.2)) where two points $(x_1, g_1), (x_2, g_2) \in \tilde{X} \times G$ are identified if there is an element $z \in \pi_1(X, x_0)$ such that $x_2 = x_1 \cdot z$ and $g_2 = \rho(z)^{-1}g_1$. The natural projection $\tilde{X} \times G \longrightarrow \tilde{X}$ produces the projection $E_G \longrightarrow X$ from the quotient space, and the right-translation action of G on $\tilde{X} \times G$ produces the action of G on E_G . The trivial connection on the trivial principal G -bundle $\tilde{X} \times G \longrightarrow \tilde{X}$ descends to the connection D on the principal G -bundle E_G .

The above construction gives the following:

Corollary 2.1. *The pulled back principal G -bundle β^*E_G is identified with the trivial principal G -bundle $\tilde{X} \times G$ by sending any $(x, g) \in \tilde{X} \times G$ to (x, z) , where z is the equivalence class of (x, g) . This identification takes the connection β^*D on β^*E_G to the trivial connection on the trivial principal G -bundle $\tilde{X} \times G$.*

A Hermitian structure on E_G is a C^∞ reduction of structure group

$$E_K \subset E_G$$

to the maximal compact subgroup $K \subset G$. For such a reduction of structure group E_K , we have the reduction of structure group

$$\beta^* E_K \subset \beta^* E_G = \tilde{X} \times G \longrightarrow \tilde{X},$$

where β is the projection in (2.2). Therefore, a Hermitian structure on E_G is given by a $\pi_1(X, x_0)$ -equivariant C^∞ map

$$h : \tilde{X} \longrightarrow G/K \quad (2.4)$$

with $\pi_1(X, x_0)$ acting on the left of G/K via the homomorphism ρ .

Let $\iota : G/K \longrightarrow K \backslash G$ be the isomorphism that sends a coset gK to the coset Kg^{-1} . Let

$$E_K \subset E_G$$

be the reduction of structure group to K corresponding to a map h as in (2.4). Since h is $\pi_1(X, x_0)$ -equivariant, the composition $\iota \circ h$ descends to a map

$$H_D : X \longrightarrow M_\Gamma = K \backslash G / \Gamma \quad (2.5)$$

(see (2.1)).

Fix a Riemannian metric g_X on X . A theorem of Corlette says that there is a Hermitian structure $E_K \subset E_G$ such that the above map h is harmonic with respect to $\beta^* g_X$ and the natural invariant metric on G/K [Co, page 368, Theorem 3.4] (in the special case where $\dim X = 2$ and $G = \mathrm{PSL}(2, \mathbb{R})$, this was proved in [Do]). Note that h is harmonic if and only if H_D in (2.5) is harmonic with respect to g_X and the natural metric on M_Γ . If E'_K is another Hermitian structure such that corresponding map $\tilde{X} \longrightarrow G/K$ is also harmonic, then there is an automorphism δ of the principal G -bundle E_G such that

- δ preserves the flat connection D on E_G , and
- $\delta(E_K) = E'_K$.

In other words, if $h' : \tilde{X} \longrightarrow G/K$ is the $\pi_1(X, x_0)$ -equivariant map corresponding to this E'_K , then there is an element $g \in G$ such that

- g commutes with $\rho(\pi_1(X, x_0))$, and
- $h'(y) = gh(y)$ for all $y \in \tilde{X}$.

Note that the first condition that g commutes with $\rho(\pi_1(X, x_0))$ implies that the map $y \longmapsto gh(y)$ intertwines the actions of $\pi_1(X, x_0)$ on \tilde{X} and G/K .

3. CONSTRUCTIONS USING FLAT CONNECTION AND HERMITIAN STRUCTURE

Let G and K be as before. As before, the Lie algebra of G will be denoted by \mathfrak{g} . The Lie algebra of K will be denoted by \mathfrak{k} . We have the isotypical decomposition

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p} \quad (3.1)$$

for the adjoint action of the center of K [He, p. 208, Theorem 3.3(iii)]. Note that the adjoint action of K on \mathfrak{g} preserves this decomposition. Let

$$\phi : \mathfrak{g} \longrightarrow \mathfrak{k} \quad \text{and} \quad \psi : \mathfrak{g} \longrightarrow \mathfrak{p} \quad (3.2)$$

be the projections associated to the decomposition in (3.1).

Let Y be a connected complex manifold, F_G a C^∞ principal G -bundle on Y and D^Y a flat connection on the principal G -bundle F_G . Fix a C^∞ reduction of structure group of F_G

$$F_K \subset F_G$$

to the subgroup K . Let

$$\text{ad}(F_K) = F_K(\mathfrak{k}) := F_K \times^K \mathfrak{k} \longrightarrow Y \quad \text{and} \quad F_K(\mathfrak{p}) := F_K \times^K \mathfrak{p} \longrightarrow Y \quad (3.3)$$

be the vector bundles associated to this principal K -bundle F_K for the K -modules \mathfrak{k} and \mathfrak{p} respectively. Let $\text{ad}(F_G) = F_G \times^G \mathfrak{g} \longrightarrow Y$ be the adjoint vector bundle for F_G . From (3.1) it follows that

$$\text{ad}(F_G) = \text{ad}(F_K) \oplus F_K(\mathfrak{p}). \quad (3.4)$$

The connection D^Y on F_G is defined by a \mathfrak{g} -valued 1-form on the total space of F_G ; this form on F_G will also be denoted by D^Y . The restriction of this form D^Y to F_K will be denoted by D' . Consider the \mathfrak{k} -valued 1-form $\phi \circ D'$ on F_K , where ϕ is the projection in (3.2). It is K -equivariant and restricts to the Maurer–Cartan form on the fibers of the principal K -bundle F_K . Hence $\phi \circ D'$ is a connection on F_K ; we will denote by D_K this connection on F_K . Consider the \mathfrak{p} -valued 1-form $\psi \circ D'$ on F_K , where ψ is the projection in (3.2). It is K -equivariant and its restriction to any fiber of the principal K -bundle F_K vanishes identically. Therefore, $\psi \circ D'$ is a C^∞ section $F_K(\mathfrak{p}) \otimes T^*Y$, where $F_K(\mathfrak{p})$ is the vector bundle defined in (3.3), and T^*Y is the real cotangent bundle of Y . Let

$$D^{Y,\mathfrak{p}} \in C^\infty(Y, F_K(\mathfrak{p}) \otimes T^*Y) \quad (3.5)$$

be this $F_K(\mathfrak{p})$ -valued 1-form on Y .

Using the complex structure on Y we may decompose any $F_K(\mathfrak{p})$ -valued 1-form on Y into a sum of $F_K(\mathfrak{p})$ -valued forms of types $(1, 0)$ and $(0, 1)$. Let

$$D^{Y,\mathfrak{p}} = D^{1,0} + D^{0,1} \quad (3.6)$$

be the decomposition of the above $F_K(\mathfrak{p})$ -valued 1-form $D^{Y,\mathfrak{p}}$ into $(1, 0)$ and $(0, 1)$ parts.

Define

$$\omega_{D^Y} := \frac{1}{\sqrt{-1}}(D^{1,0} - D^{0,1}) \in C^\infty(Y, (\text{ad}(F_G) \otimes T^*Y) \otimes_{\mathbb{R}} \mathbb{C}),$$

where $D^{1,0}$ and $D^{0,1}$ are constructed in (3.6). From (3.6) it follows that

$$\omega_{D^Y} \in C^\infty(Y, F_K(\mathfrak{p}) \otimes T^*Y) \subset C^\infty(Y, (\text{ad}(F_G) \otimes T^*Y) \otimes_{\mathbb{R}} \mathbb{C}) \quad (3.7)$$

(see (3.4)). In other words, ω_{D^Y} is a real 1-form on Y with values in the real vector bundle $F_K(\mathfrak{p}) \subset \text{ad}(F_G) \otimes_{\mathbb{R}} \mathbb{C}$.

4. CRITERION FOR HOLOMORPHICITY

We now work in the set-up of Section 2. Assume that the manifold X is equipped with a complex structure. Identify two elements (g_1, g'_1) and (g_2, g'_2) of $(K \backslash G) \times G$ if there is an element $\gamma \in \Gamma$ such that $g_2 = g_1 \gamma^{-1}$ and $g'_2 = \gamma g'_1$. The corresponding quotient will be denoted by \mathcal{E}_G . The group G acts on \mathcal{E}_G ; the action of any $g \in G$ sends the equivalence class of $(g_1, g'_1) \in (K \backslash G) \times G$ to the equivalence class of $(g_1, g'_1 g)$. Consider the map

$$\mathcal{E}_G \longrightarrow M_\Gamma = K \backslash G / \Gamma$$

that sends the equivalence class of any $(g, g') \in (K \backslash G) \times G$ to the equivalence class of g . It, and the above action of G on \mathcal{E}_G , together make \mathcal{E}_G a principal G -bundle on M_Γ . Pull back the left invariant Maurer–Cartan form on G using the projection to the second factor $(K \backslash G) \times G \longrightarrow G$. This pulled back form descends to the quotient space \mathcal{E}_G . It is straight-forward to check that this descended form defines a connection on the principal G -bundle \mathcal{E}_G . This connection on \mathcal{E}_G will be denoted by \mathcal{D}^0 . The connection \mathcal{D}^0 is flat.

Consider the submanifold $\{(g^{-1}, g) \mid g \in G\} \subset G \times G$. Let

$$\mathcal{N} \subset (K \backslash G) \times G \tag{4.1}$$

be the image of it under the obvious quotient map. Let

$$\mathcal{E}_K \subset \mathcal{E}_G \tag{4.2}$$

be the image of \mathcal{N} in the quotient space \mathcal{E}_G (recall that \mathcal{E}_G is a quotient of $(K \backslash G) \times G$). It is straight-forward to check that the action of the subgroup $K \subset G$ on \mathcal{E}_G preserves \mathcal{E}_K . More precisely, \mathcal{E}_K is a reduction of structure group of the principal G -bundle $\mathcal{E}_G \longrightarrow M_\Gamma$ to the subgroup $K \subset G$.

Take ρ as in (2.3). As in Section 2, let (E_G, D) denote the associated flat principal G -bundle over the compact complex manifold X . Take any C^∞ reduction of structure group

$$E_K \subset E_G$$

to $K \subset G$.

Proposition 4.1. *The pulled back principal G -bundle $H_D^* \mathcal{E}_G$, where H_D is constructed in (2.5), is canonically identified with the principal G -bundle E_G . This identification between E_G and $H_D^* \mathcal{E}_G$ takes*

- (1) *the pulled back connection $H_D^* \mathcal{D}^0$ to the connection D on E_G , and*
- (2) *the reduction $H_D^* \mathcal{E}_K \subset H_D^* \mathcal{E}_G$ (see (4.2)) to the reduction $E_K \subset E_G$.*

Proof. Consider the pulled back reduction of structure group $\beta^* E_K \subset \beta^* E_G$, where β is the universal cover in (2.2). From Corollary 2.1 we know that $\beta^* E_G = \tilde{X} \times G$. Let

$$f' : \beta^* E_K \longrightarrow G$$

be the composition

$$\beta^* E_K \hookrightarrow \beta^* E_G \xrightarrow{\sim} \tilde{X} \times G \xrightarrow{\text{pr}_2} G,$$

where pr_2 is the projection to the second factor. Now define the map

$$f'' : \beta^* E_K \longrightarrow (K \backslash G) \times G, \quad z \longmapsto (\widetilde{f'(z)^{-1}}, f'(z)),$$

where $\widetilde{f'(z)^{-1}} \in K \backslash G$ is the image of $f'(z)^{-1}$ under the quotient map $G \longrightarrow K \backslash G$. The Galois group $\pi_1(X, x_0)$ for β has a natural right-action of the pullback $\beta^* E_K$ that lifts the right-action of $\pi_1(X, x_0)$ on \widetilde{X} . For any $z \in \beta^* E_K$ and $\gamma \in \pi_1(X, x_0)$, we have

$$f''(z \cdot \gamma) = (\widetilde{f'(z)^{-1}} \rho(\gamma), \rho(\gamma)^{-1} f'(z)),$$

where $z \cdot \gamma \in \beta^* E_K$ is the image of z under the action of γ . This, and the fact that $\text{image}(f'') \subset \mathcal{N}$ (defined in (4.1)), together imply that f'' descends to a map

$$\widetilde{d} : E_K \longrightarrow \mathcal{E}_K$$

(see (4.2)). This map \widetilde{d} is clearly K -equivariant, and the following diagram is commutative

$$\begin{array}{ccc} E_K & \xrightarrow{\widetilde{d}} & \mathcal{E}_K \\ \downarrow & & \downarrow \\ X & \xrightarrow{H_D} & M \end{array}$$

Therefore, we get an isomorphism of principal K -bundles

$$f : E_K \longrightarrow H_D^* \mathcal{E}_K. \quad (4.3)$$

Since E_G (respectively, \mathcal{E}_G) is the extension of structure group of E_K (respectively, \mathcal{E}_K) using the inclusion of K in G , the isomorphism f in (4.3) produces an isomorphism of principal G -bundles

$$\widehat{f} : E_G \longrightarrow H_D^* \mathcal{E}_G.$$

It is straight-forward to check that \widehat{f} takes the connection D to the connection $H_D^* \mathcal{D}^0$. \square

Define $\mathcal{E}_K(\mathfrak{p}) := \mathcal{E}_K \times^K \mathfrak{p} \longrightarrow M_\Gamma$ as in (3.3). Also, construct the associated vector bundle $E_K(\mathfrak{p})$ on X as done in (3.3).

Corollary 4.2. *The pulled back vector bundle $H_D^* \mathcal{E}_K(\mathfrak{p})$, where H_D is defined in (2.5), is canonically isomorphic to $E_K(\mathfrak{p})$.*

Proof. The isomorphism f in (4.3) between principal K -bundles produces an isomorphism between the associated vector bundles. \square

Let

$$\omega_{\mathcal{D}^0} \in C^\infty(M_\Gamma, \mathcal{E}_K(\mathfrak{p}) \otimes_{\mathbb{R}} T^* M_\Gamma) \quad (4.4)$$

be the real 1-form constructed as in (3.7) for the pair $(\mathcal{D}^0, \mathcal{E}_K)$. Using the isomorphisms in Proposition 4.1 and Corollary 4.2, together with the homomorphism

$$(dH_D)^* : H_D^* T^* M_\Gamma \longrightarrow T^* X,$$

the pulled back section $H_D^* \omega_{\mathcal{D}^0}$ produces a section

$$\widetilde{H_D^* \omega_{\mathcal{D}^0}} \in C^\infty(X, E_K(\mathfrak{p}) \otimes_{\mathbb{R}} T^* X). \quad (4.5)$$

Let

$$\omega_D \in C^\infty(X, E_K(\mathfrak{p}) \otimes T^*X) \quad (4.6)$$

be the real 1-form constructed as in (3.7) for the pair (D, E_K) .

Theorem 4.3. *The map H_D is holomorphic if and only if $\widetilde{H_D^* \omega_{\mathcal{D}^0}} = \omega_D$ (see (4.5) and (4.6)).*

The map H_D is anti-holomorphic if and only if $\widetilde{H_D^ \omega_{\mathcal{D}^0}} = -\omega_D$.*

Proof. The real tangent bundle TM_Γ of $M_\Gamma = K \backslash G / \Gamma$ is identified with the associated vector bundle $\mathcal{E}_K(\mathfrak{p})$, where \mathcal{E}_K is the principal K -bundle in (4.2). Also, we have $H_D^* \mathcal{E}_K = E_K$ by Proposition 4.1. Combining these we conclude that the vector bundle $E_K(\mathfrak{p}) \rightarrow X$ in (3.3) is identified with the pullback $H_D^* TM_\Gamma$. The $E_K(\mathfrak{p})$ -valued 1-form $D^{\mathfrak{p}} \in C^\infty(X, E_K(\mathfrak{p}) \otimes T^*X)$, obtained by substituting (X, D, E_K) in place of (Y, D^Y, F_K) in (3.5), coincides with the section given by the differential

$$dH_D : TX \rightarrow H_D^* TM_\Gamma.$$

From this it follows that ω_D in (4.6) is the section given by the homomorphism

$$TX \rightarrow H_D^* TM_\Gamma, \quad v \mapsto dH_D(J_X(v)),$$

where $J_X : TX \rightarrow TX$ is the almost complex structure on X .

Let

$$\mathcal{D}^{0,\mathfrak{p}} \in C^\infty(M_\Gamma, \mathcal{E}_K(\mathfrak{p}) \otimes T^*M_\Gamma)$$

be the section constructed just as $D^{Y,\mathfrak{p}}$ is constructed in (3.5) after substituting $(\mathcal{E}_G, \mathcal{D}^0, \mathcal{E}_K)$ in place of (Y, D^Y, F_K) . This $\mathcal{D}^{0,\mathfrak{p}}$ coincides with the section given by the identity map of TM_Γ (recall that TM_Γ is identified with $\mathcal{E}_K(\mathfrak{p})$). Therefore, the section $\omega_{\mathcal{D}^0}$ in (4.4) coincides with the section given by the almost complex structure

$$J_{M_\Gamma} : TM_\Gamma \rightarrow TM_\Gamma$$

on M_Γ . Consequently, $\widetilde{H_D^* \omega_{\mathcal{D}^0}} = \omega_D$ (respectively, $\widetilde{H_D^* \omega_{\mathcal{D}^0}} = -\omega_D$) if and only if the differential dH_D takes the almost complex structure J_X to J_{M_Γ} (respectively, $-J_{M_\Gamma}$). This completes the proof. \square

Now assume that the complex manifold X is Kähler. This means that X is equipped with a Hermitian structure g_X such that

- the almost complex structure J_X on X is orthogonal with respect to g_X , and
- the $(1, 1)$ on X associate to the pair (g_X, J_X) is closed.

If A and B are Kähler manifolds, then any holomorphic map $A \rightarrow B$ is harmonic [EL, page 1, § (1.2)(e)], [Li]. This implies that any anti-holomorphic map $A \rightarrow B$ is also harmonic; to see this simply replace the almost complex structure J_A of A by the almost complex structure $-J_A$. Therefore, Theorem 4.3 has the following corollary:

Corollary 4.4. *Take the Hermitian structure on E_G given by a map $H_D : X \rightarrow K \backslash G / \Gamma$. If $\widetilde{H_D^* \omega_{\mathcal{D}^0}} = \omega_D$, then the map*

$$X \rightarrow K \backslash G / \Gamma$$

given by a harmonic metric in [Co] is holomorphic. In that case H_D gives a harmonic metric. Conversely, if H_D gives a harmonic metric and $\widetilde{H_D^ \omega_{\mathcal{D}^0}} = \omega_D$, then H_D is holomorphic.*

Take the Hermitian structure on E_G given by a map $H_D : X \rightarrow K \backslash G / \Gamma$. If $\widetilde{H_D^ \omega_{\mathcal{D}^0}} = -\omega_D$, then the map $X \rightarrow K \backslash G / \Gamma$ given by a harmonic metric in [Co] is anti-holomorphic. In that case H_D gives a harmonic metric. Conversely, if H_D gives a harmonic metric and $\widetilde{H_D^* \omega_{\mathcal{D}^0}} = -\omega_D$, then H_D is anti-holomorphic.*

REFERENCES

- [Co] K. Corlette, Flat G -bundles with canonical metrics, *Jour. Diff. Geom.* **28** (1988), 361–382.
- [Do] S. K. Donaldson, Twisted harmonic maps and the self-duality equations, *Proc. London Math. Soc.* **55** (1987), 127–131.
- [EL] J. Eells and L. Lemaire, A report on harmonic maps, *Bull. London Math. Soc.* **10** (1978), 1–68.
- [He] S. Helgason, *Differential Geometry, Lie Groups, and Symmetric Spaces*, Graduate Studies in Mathematics, 34. American Mathematical Society, Providence, RI, 2001
- [Li] A. Lichnerowicz, Applications harmoniques et variétés kähleriennes, 1968/1969 Symposia Mathematica, Vol. III (INDAM, Rome, 1968/69), pp. 341–402, Academic Press, London.

DEPARTMENT OF MATHEMATICS AND STATISTICS, KING FAHD UNIVERSITY, SAUDI ARABIA

E-mail address: hassanaz@kfupm.edu.sa

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, BOMBAY 400005, INDIA

E-mail address: indranil@math.tifr.res.in

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, BOMBAY 400005, INDIA

E-mail address: rajan@math.tifr.res.in

INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS, STRADA COSTIERA 11, 34151 TRIESTE, ITALY

E-mail address: shehryar.sikander1@gmail.com