

Alice-Bob systems, P_s - T_d - C principles and multi-soliton solutions

S. Y. Lou^{1,2*}

¹ *Shanghai Key Laboratory of Trustworthy Computing, East China Normal University, Shanghai 200062, China,*

² *Ningbo Collaborative Innovation Center of Nonlinear Hazard System of Ocean and Atmosphere
and Faculty of Science, Ningbo University, Ningbo, 315211, China*

November 9, 2021

Abstract

To describe two-place physical problems, many possible models named Alice-Bob (AB) systems are proposed. To find and to solve these systems, the Parity (P), time reversal (T), charge conjugation (C), shifted-parity (P_s , parity with a shift), delayed time reversal (T_d , time reversal with a delay) and their possible combinations such as PT, PC, P_sC , P_sT_d and P_sT_dC etc. can be successively used. Especially, some special types of P_s - T_d - C group invariant multi-soliton solutions for the KdV-KP-Toda type, mKdV-sG type, NLS type and discrete H_1 type AB systems are explicitly constructed.

PACS numbers: 05.45.Yv 02.30.Ik 11.30.Er 11.10.Lm 03.50.Kk 02.30.Jr 02.30.Ks

1 Introduction

Usually, physical models are locally established around a single place (around a single space-time point), say, $\{x, t\}$. However, various physical problems may happen in two or more places which are linked each other, for instance, in quantum physics, two far-away particles (for simplicity, we call them Alice and Bob, respectively) may be entangled each other. Whence the state of Alice, $A(x, t)$, is given, the Bob's state,

$$B(x', t') = \hat{f}A = A^f, \quad (1)$$

for a suitable operator \hat{f} may be determined at the same time. Usually, $\{x', t'\}$ is not neighbour to $\{x, t\}$. Thus, the intrinsic two-place models, the Alice-Bob systems (ABSs), are nonlocal. Some special types of this kind of nonlocal models have been proposed, such as the nonlocal nonlinear Schrödinger equations (NLS) [1] with

$$\hat{f} = \hat{P}\hat{C}$$

where \hat{P} and \hat{C} are the usual parity and charge conjugation operators.

On the other hand, symmetries play an essential role in basic physical theories. For instance, the standard model is established by means of the $SU(3) \times SU(2) \times U(1)$ symmetry and all the predictions of the model have been completely proved. 18 Nobel prizes have been nominated to the scientists who made great achievements on the model. Symmetries are not only the basic principles to establish physical models but also the key points to solve complicated scientific problems. The parity, time reversal (\hat{T}), charge conjugation, space and time translations and their suitable combinations (say $\hat{P}\hat{C}\hat{T}$ in weak interaction processes) are conserved for most of basic physical problems. Therefore, almost all the important nonlinear models possess these (or some of these) symmetries. Though these symmetries exist in various physical models, they, especially the \hat{P} - \hat{C} - \hat{T} (parity and/or charge-conjugate and/or time-reversal) symmetries, are not directly used to solve nonlinear physical systems.

*Email: lousenyue@nbu.edu.cn

In this paper, we will reveal that these symmetries are not only very useful to established some types of ABSs but also quite effective to find exact $\hat{P}\hat{T}\hat{C}$ invariant solutions of ABSs. In section 2, the ABSs are defined in detail by introducing the AB-BA principle (AB-BA equivalence principle) and the $\hat{P}_s\hat{T}_d\hat{C}$ principle (shifted parity (\hat{P}_s) and/or delayed time reversal (\hat{T}_d) and/or charge conjugate invariance principle). Especially, various particular integrable ABSs, such as the AB-KdV systems, AB-MKdV systems, AB-KP systems, AB-NLS systems (the so-called nonlocal NLS models are included as special cases), AB-sine Gordon systems, AB-Toda systems and AB- H_1 systems are listed. In section 3, using the $\hat{P}_s\hat{T}_d\hat{C}$ principle, the $\hat{P}_s\hat{T}_d\hat{C}$ invariant multiple soliton solutions are explicitly obtained for various ABSs. The last section is a short summary and discussion.

2 AB systems

2.1 AB-BA equivalence principle (AB-BA principle).

It is reasonable that Alice and Bob are in the equivalent situation which implies that Bob's coordinate system $x' \equiv \{x'_i, i = 1, 2, \dots, n\}$ should be linked with that of Alice $x \equiv \{x_i, i = 1, 2, \dots, n\}$ as

$$x' = f(x) \equiv \hat{f}x. \quad (2)$$

Inversely, Alice's coordinate system is related to that of Bob's by

$$x = f^{-1}(x') \equiv \hat{f}^{-1}x'. \quad (3)$$

Our equivalence principle requires that the function f and its inverse f^{-1} should have the same form. In other words,

$$\hat{f}^2 = 1. \quad (4)$$

It is clear that there are infinitely many solutions of (4), since any solution of

$$F(x', x) \pm F(x, x') = 0 \quad (5)$$

solves (4) for an arbitrary F . However, if the spacial relation (2) (and then (3)) is further required to be only a linear transformation, then the solutions of (4) can be completely fixed.

For one dimensional transformation, there are only two linear transformation solutions of (4), the identity transformation

$$x' = x \text{ (for space } x), t' = t \text{ (for time } t), \quad (6)$$

and the shifted parity (\hat{P}_s) transformation for the spatial variable, the delayed time reversal (\hat{T}_d) for the time variable

$$x' = -x + x_0 \equiv \hat{P}_s x, \quad t' = -t + t_0 \equiv \hat{T}_d t. \quad (7)$$

When $x_0 = 0$ ($t_0 = 0$), \hat{P}_s (\hat{T}_d) is reduced back to the pure parity \hat{P} (time reversal \hat{T}).

In two dimensional case, the allowed linear transformations of (4) possess three cases, the identity transformation

$$\begin{pmatrix} x'_1 \\ x'_2 \end{pmatrix} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, \quad (8)$$

the shifted parity with arbitrary constants x_{10} and x_{20}

$$\begin{pmatrix} x'_1 \\ x'_2 \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} + \begin{pmatrix} x_{10} \\ x_{20} \end{pmatrix} \equiv \hat{P}_s^{x_1 x_2} x, \quad (9)$$

and the mixed shifted parity and rotation

$$\begin{pmatrix} x'_{i_1} \\ x'_{i_2} \end{pmatrix} = \begin{pmatrix} a_1 a_2 - 1 & a_1 \\ a_2(2 - a_1 a_2) & 1 - a_1 a_2 \end{pmatrix} \begin{pmatrix} x_{i_1} \\ x_{i_2} \end{pmatrix} - \begin{pmatrix} -x_{10} \\ a_2 x_{10} \end{pmatrix}, \quad (10)$$

where $\{i_1, i_2\} = \{1, 2\}$ or $\{2, 1\}$ and $\{a_1, a_2, x_{10}\}$ are arbitrary constants.

In three dimensional case, we have four cases, the identity transformation

$$\begin{pmatrix} x'_1 \\ x'_2 \\ x'_3 \end{pmatrix} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}, \quad (11)$$

the shifted parity with arbitrary constants x_{10} , x_{20} and x_{30}

$$\begin{pmatrix} x'_1 \\ x'_2 \\ x'_3 \end{pmatrix} = \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} + \begin{pmatrix} x_{10} \\ x_{20} \\ x_{30} \end{pmatrix} \equiv \hat{P}_s^{x_1 x_2 x_3} x, \quad (12)$$

the first type of mixed shifted parity and rotation ($A_1 \equiv -a_1 a_3 - a_2 a_4$)

$$\begin{pmatrix} x'_{i_1} \\ x'_{i_2} \\ x'_{i_3} \end{pmatrix} = \begin{pmatrix} A_1 - 1 & a_2 & a_3 \\ a_4(A_1 - 2) & 1 + a_2 a_4 & a_3 a_4 \\ a_1(A_1 - 2) & a_1 a_2 & 1 + a_1 a_3 \end{pmatrix} \begin{pmatrix} x_{i_1} \\ x_{i_2} \\ x_{i_3} \end{pmatrix} + \begin{pmatrix} x_{10} \\ a_4 x_{10} \\ a_1 x_{10} \end{pmatrix}, \quad (13)$$

and the second type of mixed shifted parity and rotation

$$\begin{pmatrix} x'_{i_1} \\ x'_{i_2} \\ x'_{i_3} \end{pmatrix} = \begin{pmatrix} a_1 a_3 - 1 & a_2 a_3 & a_3 \\ a_1 a_4 & a_2 a_4 - 1 & a_4 \\ a_1(A_1 + 2) & a_2(A_1 + 2) & A_1 + 1 \end{pmatrix} \begin{pmatrix} x_{i_1} \\ x_{i_2} \\ x_{i_3} \end{pmatrix} + \begin{pmatrix} -x_{10} \\ -x_{20} \\ a_1 x_{10} + a_2 x_{20} \end{pmatrix}. \quad (14)$$

In (13) and (14), $\{i_1, i_2, i_3\}$ are any permutations of $\{1, 2, 3\}$, i.e. $\{i_1, i_2, i_3\} = \{1, 2, 3\}, \{1, 3, 2\}, \{2, 1, 3\}, \{2, 3, 1\}, \{3, 1, 2\}$ or $\{3, 2, 1\}$, and $\{a_1, a_2, a_3, a_4, x_{10}, x_{20}\}$ are arbitrary constants.

Using the above AB-BA equivalence principle, our most general AB systems have the form

$$CAB \equiv \begin{cases} G_1(A, B) = 0, \\ G_2(A, B) = 0, \end{cases} \quad (15)$$

$$B = A^f = \hat{f}A \quad (16)$$

with the AB-BA equivalence condition

$$\hat{f}^2 = 1, \quad x' = \hat{f}x, \quad x = \hat{f}x' \quad (17)$$

for a suitable \hat{f} as listed in (7), (9), (10), (12), (13), (14) and so on.

Obviously, an ABS is a coupled system CAB (15) which is consistent under the correlated condition (16).

Because of the condition (17), the general coupled system (15) should be invariant under the transformation \hat{f} , i.e.,

$$\hat{f}CAB = CAB. \quad (18)$$

For physically significance, we only discuss the coupled system (15) known in natural science.

2.2 AB-KdV systems

In nonlinear physics, the KdV equation ($A = A(x, t)$) [2]

$$KdV \equiv A_t + A_{xxx} + 6AA_x = 0 \quad (19)$$

is one of the most important and famous models which can be applied in almost all physical fields especially in fluid related branches [3].

The most general coupled homogeneous KdV system may have the form

$$CKdV \equiv \begin{cases} A_t + a_1 A_{xxx} - a_2 B_{x'x'x'}^f + (a_3 A + a_4 B^f) A_x - (a_5 A + a_6 B^f) B_{x'}^f = 0, \\ B_{t'}^f - b_1 A_{xxx} + b_2 B_{x'x'x'}^f - (b_3 A + b_4 B^f) A_x + (b_5 A + b_6 B^f) B_{x'}^f = 0, \end{cases} \quad (20)$$

$$= \begin{cases} A_t + a_1 A_{xxx} - a_2 B_{x'x'x'}^f + (a_3 A + a_4 B^f) A_x - (a_5 A + a_6 B^f) B_{x'}^f = 0, \\ B_t - b_1 A_{x'x'x'}^f + b_2 B_{xxx} - (b_3 A^f + b_4 B) A_{x'}^f + (b_5 A^f + b_6 B) B_x = 0, \end{cases} \quad (21)$$

$$= \begin{cases} A_t + a_1 A_{xxx} + a_2 B_{1xxx} + (a_3 A + a_4 B_1) A_x + (a_6 A + a_7 B_1) B_{1x} = 0, \\ B_{1t} + b_1 A_{xxx} + b_2 B_{1xxx} + (b_3 A + b_4 B_1) A_x + (b_5 A + b_6 B_1) B_{1x} = 0, \end{cases} \quad (22)$$

where

$$\begin{aligned} A &= A(x, t), \quad A^f = \hat{f}A = A(x', t'), \quad B = B(x, t), \quad B^f = \hat{f}B = B(x', t') = B_1(x, t), \\ x' &= \hat{f}x = -x + x_0, \quad t' = \hat{f}t = -t + t_0, \quad \hat{f} = \hat{P}_s \hat{T}_d. \end{aligned} \quad (23)$$

The general coupled KdV system possesses three equivalent expressions (20), (21) and (22). However, the physical meanings of them are quite different. Equation (22) is a usual single place nonlinear model, (20) describes the two-place nonlinear interaction between $A(x, t)$ and $B(x', t')$, while (21) displays an interaction model between two-place quantities $\{A(x, t), B(x, t)\}$ and $\{A(x', t'), B(x', t')\}$.

The general coupled one-place KdV system (22) has been studied in many literatures [4, 5, 6, 7, 8, 9, 10]. Its possible integrable classification has been done by the general symmetry analysis, bi-Hamiltonian structure, Painlevé test and prolongation search on its Lax pairs. The model can also be derived from two layer fluids [4].

Due to the equivalence of (20), (21) and (22), it may be more interesting to find some possible two-place reductions which are not equivalent to some one-place models.

It is fortunate that the special case of (21)

$$SCKdV \equiv \begin{cases} A_t + a_1 A_{xxx} - a_2 B_{x'x'x'}^f + (a_3 A + a_4 B^f) A_x - (a_6 A + a_7 B^f) B_{x'}^f = 0, \\ B_t - a_2 A_{x'x'x'}^f + a_1 B_{xxx} - (a_7 A^f + a_6 B) A_{x'}^f + (a_4 A^f + a_3 B) B_x = 0, \end{cases} \quad (24)$$

allows an intrinsic two-place AB-KdV reduction

$$\begin{aligned} ABKdV &\equiv A_t + a_1 A_{xxx} + a_2 B_{xxx} + (a_3 A + a_4 A^f) A_x + (a_6 A + a_7 A^f) B_x = 0, \\ B &= A^f = A^{P_s T_d} = A(-x + x_0, -t + t_0), \end{aligned} \quad (25)$$

by directly taking $B = A$ from (24).

Here we list four special integrable realizations of (25).

The first type of integrable AB-KdV reduction has the form

$$ABKdV_1 \equiv A_t - \frac{1}{2} A_{xxx} + \frac{3}{2} B_{xxx} - 3(A - B)A_x + 6AB_x = 0, \quad B = A^{P_s T_d}. \quad (26)$$

It is integrable because it can also be obtained from the reduction of the Hirota-Satsuma system [11] and thus one can obtain its Lax pair [12]

$$\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_{xx} = \begin{pmatrix} -A & \lambda_1 \\ \lambda_2 & -B \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}, \quad (27)$$

$$\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_t = \begin{pmatrix} \frac{1}{2}(3B - A)_x + (A - 3B)\partial_x & -4\lambda_1\partial_x \\ -4\lambda_2\partial_x & \frac{1}{2}(3A - B)_x + (B - 3A)\partial_x \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \quad (28)$$

with two arbitrary spectral parameters λ_1 and λ_2 .

The second type of AB-KdV has the form

$$ABKdV_2 \equiv A_t + A_{xxx} + 3(A+B)A_x = 0, \quad B = \hat{P}_s \hat{T}_d A. \quad (29)$$

When $\{x', t'\}$ is changed as $\{x, t\}$, the $ABKdV_2$ is reduced back to the known KdV equation. It is not difficult to prove that the AB-KdV model (29) is integrable with the Lax pair

$$\begin{aligned} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_{xx} &= \begin{pmatrix} -\frac{1}{2}C - \lambda_1 & 0 \\ \lambda_2 - \lambda_4 A_x - \lambda_5 B_x + (\lambda_3 + \lambda_6 C)\partial_x & -\frac{1}{2}C - \lambda_1 \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}, \\ \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_t &= \begin{pmatrix} \frac{1}{2}C_x + \lambda_8 + (4\lambda_1 - C)\partial_x & 0 \\ E - D\partial_x & \frac{1}{2}C_x + \lambda_8 + (4\lambda_1 - C)\partial_x \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}, \end{aligned} \quad (30)$$

where

$$\begin{aligned} C &\equiv A + B, \\ D &\equiv \lambda_6 C_x + 2(\lambda_4 A_x + \lambda_5 B_x) + 4\lambda_2, \\ E &\equiv \lambda_7 + \lambda_4 A_{xx} + \lambda_5 B_{xx} + \frac{1}{2}\lambda_3 C - \frac{1}{4}C\lambda_6(C + 8\lambda_1). \end{aligned} \quad (31)$$

The third type of integrable AB-KdV system possesses the form

$$ABKdV_3 \equiv A_t + A_{xxx} + \frac{3}{4}[(A+B)^2]_x = 0, \quad B = A(-x + x_0, -t + t_0) \quad (32)$$

with the following Lax pair

$$\begin{aligned} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_{xx} &= \begin{pmatrix} -\frac{1}{2}C - \lambda_1 & 0 \\ \lambda_2 - C_{1x} + (\lambda_6 + \lambda_5 C - 2C_1)\partial_x & -\frac{1}{2}C - \lambda_1 \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}, \\ \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_t &= \begin{pmatrix} \frac{1}{2}C_x + \lambda_8 + (4\lambda_1 - C)\partial_x & 0 \\ E_1 - (4\lambda_2 + \lambda_5 C_x)\partial_x & \frac{1}{2}C_x + \lambda_8 + (4\lambda_1 - C)\partial_x \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}, \end{aligned} \quad (33)$$

where

$$\begin{aligned} C &\equiv A + B, \quad C_1 \equiv (\lambda_3 A + \lambda_4 B), \\ E_1 &\equiv C_{1xx} + 4\lambda_1 C_1 + \frac{1}{2}\lambda_6 C + \lambda_7 - \frac{1}{4}C[\lambda_5(C + 8\lambda_1) + C_1 - 3\lambda_4 A - 3\lambda_3 B]. \end{aligned} \quad (34)$$

The fourth type of the AB-KdV system can be written as

$$ABKdV_4 \equiv A_t + A_{xxx} + \frac{3}{2}(B + 2A)A_x + \frac{3}{2}AB_x = 0, \quad B = A^{P_s T_d}. \quad (35)$$

The integrability of the fourth AB-KdV (35) is guaranteed by the following Lax pair ($C_2 \equiv \lambda_3 A + \lambda_2 B$)

$$\begin{aligned} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_{xx} &= \begin{pmatrix} -\frac{1}{2}C - \frac{1}{4}\lambda_1 & 0 \\ -C_{2x} - C_2\partial_x & -\frac{1}{4}\lambda_1 \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}, \\ \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_t &= \begin{pmatrix} \frac{1}{2}C_x + \lambda_5 + (\lambda_1 - C)\partial_x & 0 \\ C_{2xx} + \frac{1}{2}C_2(\lambda_1 + 2C) - C_{2x}\partial_x & \lambda_5 + \lambda_1\partial_x \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}. \end{aligned} \quad (36)$$

It is also interesting that, the fourth type of the AB-KdV system (35) possesses another Lax pair:

$$\begin{aligned} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_{xx} &= \begin{pmatrix} -\frac{1}{2}C - \lambda_1 & 0 \\ \lambda_2 - C_{1x} + (\lambda_6 + \lambda_5 C - 2C_1)\partial_x & -\frac{1}{2}C - \lambda_1 \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}, \\ \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_t &= \begin{pmatrix} \frac{1}{2}C_x + \lambda_8 + (4\lambda_1 - C)\partial_x & 0 \\ E_2 - (4\lambda_2 + \lambda_5 C_x)\partial_x & \frac{1}{2}C_x + \lambda_8 + (4\lambda_1 - C)\partial_x \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}, \end{aligned} \quad (37)$$

with

$$E_2 \equiv C_{1xx} + 4\lambda_1 C_1 + \frac{1}{2}\lambda_6 C - \frac{1}{4}C[\lambda_5(C + 8\lambda_1) - 2C_1] + \lambda_7. \quad (38)$$

Finally, it is worth to emphasize that all the AB-KdV systems listed above possess the $\hat{P}_s \hat{T}_d = \hat{f}$ invariance and this invariance originates from the invariance of the coupled KdV (22)

$$\hat{P}_s \hat{T}_d C K dV = C K dV. \quad (39)$$

2.3 AB-NLS systems

Another ubiquitous nonlinear physical model is the so-called nonlinear Schrödinger (NLS) equation

$$NLS \equiv iA_t + A_{xx} + 2\sigma|A|^2 A = 0, \quad \sigma = \pm 1. \quad (40)$$

Similar to the AB-KdV systems as discussed in the last subsection, to extend the NLS equation to possible two-place AB models, we rewrite it as a special reduction of the AKNS system

$$AKNS \equiv \begin{cases} iA_t + A_{xx} + 2\sigma A^2 B_1 = 0, \\ -iB_{1t} + B_{1xx} + 2\sigma B_1^2 A = 0, \end{cases} \quad (41)$$

$$= \begin{cases} iA_t + A_{xx} + 2\sigma A^2 B^{P_s} = 0, & (B_1 = B^{P_s}), \\ -iB_t + B_{xx} + 2\sigma B^2 A^{P_s} = 0, \end{cases} \quad (42)$$

$$= \begin{cases} iA_t + A_{xx} + 2\sigma A^2 B^{P_s T_d C} = 0, & (B_1 = B^{P_s T_d C}), \\ -iB_t + B_{xx} + 2\sigma B^2 A^{P_s T_d C} = 0. \end{cases} \quad (43)$$

Similarly, though the systems (41), (42), and (43) are equivalent, their physical meanings are quite different. Especially, their reductions corresponding to $B = A^C = A^*$ (or $B_1 = A^C = A^*$ for (41)) are completely different. The reduction of (41) is just the well known NLS equation (40). The reduction of (42) has the form

$$ABNLS_1 \equiv iA_t + A_{xx} + 2\sigma A^2 B = 0, \quad B = A^{P_s C} = \hat{P}_s \hat{C} A = A^*(-x + x_0, t). \quad (44)$$

When $x_0 = 0$, the $ABNLS_1$ (44) is just the so-called nonlocal NLS which was first introduced by Ablowitz and Musslimani [1]. The model (44) without x_0 has been widely studied by many authors [13]. However, it is still a new model for $x_0 \neq 0$.

The reduction of (43) has the form

$$ABNLS_2 \equiv iA_t + A_{xx} + 2\sigma A^2 B = 0, \quad B = A^{P_s T_d} = \hat{P}_s \hat{T}_d A = A(-x + x_0, -t + t_0). \quad (45)$$

In our knowledge, the AB-NLS model (45) is also new. It is a really two-place nonlocal model though it is reduced from a one-place coupled system.

All the models in this subsection are P_s invariant, $T_d C$ invariant and $P_s T_d C$ invariant.

The integrability of the AB-NLS systems (44) and (45) is guaranteed by the integrability of the AKNS system (41) with the well known Lax pair

$$\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_x = \begin{pmatrix} \lambda & A \\ -\sigma B & -\lambda \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}, \quad (46)$$

$$\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_t = \begin{pmatrix} 2i\lambda^2 + i\sigma AB & 2i\lambda A + iA_x \\ -i\sigma(2\lambda B + B_x) & -2i\lambda^2 - i\sigma AB \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}, \quad (47)$$

where $B = A^{P_s C} = A^*(-x + x_0, t)$ for $ABNLS_1$ (44) and $B = A^{P_s T_d} = A(-x + x_0, -t + t_0)$ for $ABNLS_2$ (45).

2.4 AB-MKdV systems

The well known modified KdV equation (MKdV) can be written in the form

$$MKdV \equiv A_t + A_{xxx} + 6\sigma A^2 A_x = 0, \quad \sigma = \pm 1, \quad (48)$$

which is a natural reduction of the third order AKNS system,

$$AKNS3 \equiv \begin{cases} A_t + A_{xxx} + 6AB_1 A_x = 0, \\ B_{1t} + B_{1xxx} + 6AB_1 B_{1x} = 0, \end{cases} \quad (49)$$

by taking

$$B_1 = \sigma A.$$

It is clear that the one-place coupled model (49) possesses an equivalent two-place form

$$AKNS3 \equiv \begin{cases} A_t + A_{xxx} + 6AB^f A_x = 0, & B^f \equiv B(-x + x_0, -t + t_0) \\ B_t + B_{xxx} + 6A^f B B_x = 0, \end{cases} \quad (50)$$

by taking

$$B_1 = B^f.$$

Starting from the two-place form (50), one can readily obtain an intrinsic two-place AB-MKdV equation

$$ABMKdV \equiv A_t + A_{xxx} + 6\sigma ABA_x = 0, \quad B = A^f = \hat{P}_s \hat{T}_d A = A(-x + x_0, -t + t_0). \quad (51)$$

The model (51) is also a new model though a special case of (51) with $x_0 = 0$ and $t_0 = 0$ has been given by Ji and Zhu [14].

The ABMKdV (51) is integrable because it is a reduction of the third order AKNS system. Thus its Lax pair can be obtained from that of the AKNS system,

$$\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_x = \begin{pmatrix} -\lambda & A \\ -\sigma B & \lambda \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \quad (52)$$

$$\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_t = \begin{pmatrix} 4\lambda^3 + 2\sigma\lambda AB + \sigma AB_x - \sigma BA_x & -4\lambda^2 A + 2\lambda A_x - A_{xx} - 2\sigma A^2 B \\ 4\sigma\lambda^2 B + 2\sigma\lambda B_x + \sigma B_{xx} + 2B^2 A & -4\lambda^3 - 2\sigma\lambda AB - \sigma AB_x + BA_x \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}. \quad (53)$$

Many other types of intrinsic AB-MKdV system can be found from the coupled MKdV systems, say, those obtained from suitable Miura transforms of the AB-KdV systems [9].

2.5 AB-sine-Gordon equation

The well known sine-Gordon (sG) equation

$$sG \equiv A_{xt} = \sin(A) \quad (54)$$

is one of the most important integrable relativistic models which is widely used in the field theory, optics and condense matter physics. The sG model is also a special member of the AKNS hierarchy. Its general coupled AKNS form can be written as ($\sigma^2 = 1$)

$$ABsG \equiv \begin{cases} A_{xt} = \left(\sigma \sqrt{a - \sin(A) \sin(B)} + A_x A_t \right) \tan(A) + \frac{1}{2} \frac{\tan(A) B_t + \tan(B) A_t}{\tan(A) \tan(B) - a \sec(A) \sec(B)} A_x, \\ B_{xt} = \left(\sigma \sqrt{a - \sin(A) \sin(B)} + B_x B_t \right) \tan(B) + \frac{1}{2} \frac{\tan(A) B_t + \tan(B) A_t}{\tan(A) \tan(B) - a \sec(A) \sec(B)} B_x, \end{cases} \quad (55)$$

Though the coupled sG system (55) is one member of the AKNS hierarchy, it seems that the model have not yet been explicitly written down elsewhere. In addition to the known sG (or sinh-Gordon) reduction

(54) by taking $B = A$ and $a = 1$ (or $B = A = i\phi$ and $a = 1$), there exist some other simple interesting reductions which have not yet been studied. For instance, the tan-Gordon (TG) equation

$$A_{xt} = (c + A_x A_t) \tan(A), \quad (56)$$

the tanh-Gordon (ThG) equation

$$A_{xt} = (c + A_x A_t) \tanh(A), \quad (57)$$

and the two-place model, the AB-sG model ($B \equiv A^{P_s T_d} = A(-x + x_0, -t + t_0)$),

$$A_{xt} = \left(\sigma \sqrt{a - \delta \sin(A) \sin(B)} + A_x A_t \right) \tan(A) + \frac{1}{2} \frac{[\tan(A) B_t + \tan(B) A_t] A_x}{\tan(A) \tan(B) - a \sec(A) \sec(B)}, \quad (58)$$

can be reduced from (55) via $\{B = 0, \sigma \sqrt{a} = c\}$, $\{B = 0, A \rightarrow iA, \sigma \sqrt{a} = c\}$ and $\{B \rightarrow \delta A^{P_s T_d}, \delta^2 = 1\}$, respectively.

The Lax pair of the coupled sG system (54) and all their reductions can be written as

$$\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_x = \begin{pmatrix} -\lambda & \frac{1}{2} \frac{\sigma \cos(A) A_x}{\sqrt{a - \sin(A) \sin(B)}} \\ -\frac{1}{2} \frac{\sigma \cos(B) B_x}{\sqrt{a - \sin(A) \sin(B)}} & \lambda \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}, \quad (59)$$

$$\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_t = \frac{1}{4\lambda} \begin{pmatrix} -\sigma \sqrt{a - \sin(A) \sin(B)} & \sin(A) \\ \sin(B) & \sigma \sqrt{a - \sin(A) \sin(B)} \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \quad (60)$$

with the suitable selections of B and A for the corresponding models.

When $B = A$ and $a = 1$, the AB-sG system (58) returns back to the usual sG system (54). There are also many other types of AB-sG systems, for instance, the negative ones of the integrable AB-MKdV hierarchies.

2.6 AB-KP system

In 2+1 dimensions, one of the most important nonlinear systems is the so-called KP equation

$$KP \equiv A_{xt} = \left(\frac{1}{4} A_{xxx} + 6AA_x \right)_x + \frac{3}{4} A_{yy}. \quad (61)$$

In order to find some nontrivial AB-KP systems, one can rewrite some known coupled KP systems to their two-place forms. For instance, one of the coupled KP system has the form

$$CKP \equiv \begin{cases} A_{xt} = \left(\frac{1}{4} A_{xx} + 3AB + \frac{3}{2} A^2 - \frac{3}{2} B^2 \right)_{xx} + \frac{3}{4} A_{yy}, \\ B_{xt} = \left(\frac{1}{4} B_{xx} + 3AB + \frac{3}{2} B^2 - \frac{3}{2} A^2 \right)_{xx} + \frac{3}{4} B_{yy}, \end{cases} \quad (62)$$

$$= \begin{cases} A_{xt} = \left[\frac{1}{4} A_{xx} + 3AB^f + \frac{3}{2} A^2 - \frac{3}{2} (B^f)^2 \right]_{xx} + \frac{3}{4} A_{yy}, \\ B_{xt} = \left[\frac{1}{4} B_{xx} + 3A^f B + \frac{3}{2} B^2 - \frac{3}{2} (A^f)^2 \right]_{xx} + \frac{3}{4} B_{yy}, \end{cases} \quad (63)$$

with

$$A^f = \hat{P}_s^{xy} \hat{T}_d A = A(-x + x_0, -y + y_0, -t + t_0), \quad (64)$$

or

$$A^f = \hat{P}_s^x \hat{T}_d A = A(-x + x_0, y, -t + t_0), \quad (65)$$

or

$$A^f = \hat{P}_s^y A = A(x, -y + y_0, t). \quad (66)$$

It is straightforward to see that the coupled KP system (63) permits an intrinsic two-place AB-KP reduction,

$$ABKP \equiv A_{xt} = \left[\frac{1}{4}A_{xx} + 3AB + \frac{3}{2}A^2 - \frac{3}{2}B^2 \right]_{xx} + \frac{3}{4}A_{yy}, \quad B = A^f \quad (67)$$

with A^f given by (64), (65) or (66).

The coupled KP system (62) is Lax integrable with the Lax pair

$$\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_y = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_{xx} + \begin{pmatrix} 2(A+B) & -2(A-B) \\ 2(A-B) & 2(A+B) \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}, \quad (68)$$

$$\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_t = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}_{xxx} + \begin{pmatrix} \alpha_+ & -\alpha_- \\ \alpha_- & \alpha_+ \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}, \quad (69)$$

where

$$\alpha_{\pm} \equiv \frac{3}{2}(A \pm B)_x + \frac{3}{2} \int (A \pm B)_y dx + 3(A \pm B)\partial_x. \quad (70)$$

For the AB-KP equation (67), the Lax pair possesses the same form as (68) and (69) but with $B = A^f$ and (64), (65) or (66).

In fact, for the coupled KP system (62), there exist some other complicated AB reductions if other AB-BA equivalent coordinates are taken as those shown in (10), (13) and (14).

2.7 AB-Toda systems

The AB systems can also be established in discrete cases. One of the most important discrete integrable model is the so-called Toda lattice which is equivalent to

$$Toda \equiv (A_n + 2)A_{n,xx} - A_{n,x}^2 + \frac{1}{2}(A_n + 2)^2 E^2 A_{n-1} = 0, \quad (71)$$

where the difference operator E is defined as

$$EA_n = A_{n+1} - A_n. \quad (72)$$

In order to find some intrinsic two-place models, the AB-Toda systems, we should rewrite the *Toda* system (71) in the coupled forms.

One of the simplest coupled Toda system may have the form

$$CTD \equiv \begin{cases} [(A_n + 2)^2 + (B'_n + 2)^2]A_{nxx} = (A_n + 2) \left(A_{nx}^2 - B_{nx}'^2 \right) + 2(B'_n + 2)A_{nx}B'_{nx} \\ \quad + \frac{1}{4}[(A_n + 2)^2 + (B'_n + 2)^2] \left[(A_n + B'_n + 4)E^2 A_{n-1} + (A_n - B'_n)E^2 B'_{n-1} \right], \\ \{ (A_n + 2)^2 + (B'_n + 2)^2 \} B'_{nxx} = (B'_n + 2) \left(B_{nx}'^2 - A_{nx}^2 \right) + 2(A_n + 2)A_{nx}B'_{nx} \\ \quad + \frac{1}{4} \{ (A_n + 2)^2 + (B'_n + 2)^2 \} \left[(A_n + B'_n + 4)E^2 B'_{n-1} + (B'_n - A_n)E^2 A_{n-1} \right], \\ [(A_n + 2)^2 + (B_n^f + 2)^2]A_{nxx} = (A_n + 2) \left[A_{nx}^2 - (B_{nx}^f)^2 \right] + 2(B_n^f + 2)A_{nx}B_{nx}^f \\ \quad + \frac{1}{4}[(A_n + 2)^2 + (B_n^f + 2)^2] \left[(A_n + B_n^f + 4)E^2 A_{n-1} + (A_n - B_n^f)E^2 B_{n-1}^f \right], \\ [(A_n^f + 2)^2 + (B_n + 2)^2] B_{nxx} = (B_n + 2) \left[B_{nx}^2 - (A_{nx}^f)^2 \right] + 2(A_n^f + 2)A_{nx}^f B_{n,x} \\ \quad + \frac{1}{4}[(A_n^f + 2)^2 + (B_n + 2)^2] \left[(A_n^f + B_n + 4)E^2 B_{n-1} + (B_n - A_n^f)E^2 A_{n-1}^f \right], \end{cases} \quad (73)$$

$$= \begin{cases} [(A_n + 2)^2 + (B_n^f + 2)^2]A_{nxx} = (A_n + 2) \left[A_{nx}^2 - (B_{nx}^f)^2 \right] + 2(B_n^f + 2)A_{nx}B_{nx}^f \\ \quad + \frac{1}{4}[(A_n + 2)^2 + (B_n^f + 2)^2] \left[(A_n + B_n^f + 4)E^2 A_{n-1} + (A_n - B_n^f)E^2 B_{n-1}^f \right], \\ [(A_n^f + 2)^2 + (B_n + 2)^2] B_{nxx} = (B_n + 2) \left[B_{nx}^2 - (A_{nx}^f)^2 \right] + 2(A_n^f + 2)A_{nx}^f B_{n,x} \\ \quad + \frac{1}{4}[(A_n^f + 2)^2 + (B_n + 2)^2] \left[(A_n^f + B_n + 4)E^2 B_{n-1} + (B_n - A_n^f)E^2 A_{n-1}^f \right], \end{cases} \quad (74)$$

where

$$A_n^f = \hat{f}A_n = \hat{P}_s^x \hat{P}_s^n A_n = A(-x + x_0, -n + n_0), \quad (75)$$

or

$$A_n^f = \hat{f}A_n = \hat{P}_s^x A_n = A(-x + x_0, n), \quad (76)$$

or

$$A_n^f = \hat{f}A_n = \hat{P}_s^n A_n = A(x, -n + n_0). \quad (77)$$

It is interesting that in addition to the usual Toda reduction (71) from (73) via $B' = A$, there are three intrinsic two-place Toda, AB-Toda reductions,

$$\begin{aligned} [(A_n + 2)^2 + (B_n + 2)^2]A_{nxx} &= (A_n + 2) [A_{nx}^2 - (B_{nx})^2] + 2(B_n + 2)A_{nx}B_{nx} \\ &+ \frac{1}{4}[(A_n + 2)^2 + (B_n + 2)^2] [(A_n + B_n + 4)E^2 A_{n-1} + (A_n - B_n)E^2 B_{n-1}], \end{aligned} \quad (78)$$

by taking $B_n = A_n^f$ with A_n^f given by (75), (76) or (77).

The AB-Toda system (78) is Lax integrable with the following Lax pair

$$E \begin{pmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \\ \phi_4 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -\frac{1}{2}(A_n + B_n + 4) & 0 & \lambda_1 - u_n & 0 \\ \frac{1}{2}(B_n - A_n) & -\frac{1}{2}(A_n + B_n + 4) & \lambda_2 - v_n & \lambda_1 - u_n \end{pmatrix} \begin{pmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \\ \phi_4 \end{pmatrix} \quad (79)$$

$$\begin{pmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \\ \phi_4 \end{pmatrix}_t = \begin{pmatrix} 1 - \lambda_1 + u_{n-1} & 0 & 1 & 0 \\ v_{n-1} - \lambda_2 & 1 - \lambda_1 + u_{n-1} & 0 & 1 \\ -\frac{1}{2}(A_n + B_n + 4) & 0 & 1 & 0 \\ \frac{1}{2}(B_n - A_n) & -\frac{1}{2}(A_n + B_n + 4) & 0 & 1 \end{pmatrix} \begin{pmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \\ \phi_4 \end{pmatrix} \quad (80)$$

where u_n and v_n are related to A_n and B_n by

$$u_{nx} = \frac{1}{2}E(A_n + B_n), \quad v_{nx} = \frac{1}{2}E(A_n - B_n), \quad (81)$$

and λ_1 and λ_2 are two arbitrary constant spectral parameters.

2.8 AB- H_1 systems

The same idea can also be applied to difference-difference systems such as the H_1 , H_2 , H_3 , Q_1 , Q_2 , Q_3 and Q_4 [15]. Here, we discuss only some AB- H_1 extensions. The usual H_1 system possesses the form

$$(u - \hat{u})(\tilde{u} - \hat{u}) + q - p = 0, \quad u = u(n, m), \quad \hat{u} = u(n, m + 1), \quad \tilde{u} = u(n + 1, m), \quad \hat{\tilde{u}} = u(n + 1, m + 1) \quad (82)$$

with arbitrary constants p and q . One of its integrable coupled extension reads

$$(\hat{\hat{A}} - A)(\hat{A} - \tilde{A} + \hat{B} - \tilde{B})(\hat{A} - \tilde{A} - \hat{B} + \tilde{B}) = (p_1 - q_1)(\tilde{A} - \hat{A}) - (p_2 - q_2)(\tilde{B} - \hat{B}), \quad (83)$$

$$(\hat{\hat{B}} - B)(\hat{B} - \tilde{B} + \hat{A} - \tilde{A})(\hat{B} - \tilde{B} - \hat{A} + \tilde{A}) = (p_1 - q_1)(\tilde{B} - \hat{B}) - (p_2 - q_2)(\tilde{A} - \hat{A}) \quad (84)$$

with arbitrary constants p_1 , p_2 , q_1 and q_2 .

It is straightforwardly verified that both the H_1 equation (82) and the coupled H_1 system of equations(83) and (84) are $\hat{f}\epsilon\hat{F}$ invariant with

$$F = \{\hat{P}_s^n, \hat{P}_s^m, \hat{P}_s^u, \hat{P}_s^n \hat{P}_s^m, \hat{P}_s^n \hat{P}_s^u, \hat{P}_s^u \hat{P}_s^m, \hat{P}_s^n \hat{P}_s^m \hat{P}_s^u\} \quad (85)$$

where \hat{P}_s^u , \hat{P}_s^n and \hat{P}_s^m are defined by

$$\hat{P}_s^n n = -n + n_0, \quad \hat{P}_s^m m = -m + m_0, \quad \hat{P}_s^u \{u, A, B\} = \{-u + u_0, -A + u_0, -B + u_0\} \quad (86)$$

with arbitrary constants n_0 , m_0 and u_0 . Because of the $\hat{P}_s^n \hat{P}_s^m$ invariance, the coupled H_1 system (83) and (84) allows an AB- H_1 reduction

$$(\hat{A} - A)(\hat{A} - \tilde{A} + \hat{B} - \tilde{B})(\hat{A} - \tilde{A} - \hat{B} + \tilde{B}) = (p_1 - q_1)(\tilde{A} - \hat{A}) - (p_2 - q_2)(\tilde{B} - \hat{B}), \quad (87)$$

$$B = A^f = \hat{f}A, \quad \hat{f} \in F. \quad (88)$$

By fixing constants $p_2 = p_1$ and $q_2 = q_1$, the AB- H_1 system (87) can be reduced to

$$(\hat{A} - A)(\tilde{A} - \hat{A} - \hat{B} + \tilde{B}) = (p_1 - q_1), \quad (89)$$

$$B = A^f, \quad \hat{f} \in F. \quad (90)$$

The integrability of the AB- H_1 is integrable because of the existence of the Lax pair

$$\hat{\phi} = \alpha \begin{pmatrix} A & B & p_1 - r_1 - A\hat{A} - B\hat{B} & p_2 - r_2 - A\hat{B} - B\hat{A} \\ B & A & p_2 - r_2 - A\hat{B} - B\hat{A} & p_1 - r_1 - A\hat{A} - B\hat{B} \\ 1 & 0 & -\hat{A} & -\hat{B} \\ 0 & 1 & -\hat{B} & -\hat{A} \end{pmatrix} \phi, \quad (91)$$

$$\tilde{\phi} = \beta \begin{pmatrix} A & B & p_1 - r_1 - A\tilde{A} - B\tilde{B} & q_2 - r_2 - A\tilde{B} - B\tilde{A} \\ B & A & q_2 - r_2 - A\tilde{B} - B\tilde{A} & q_1 - r_1 - A\tilde{A} - B\tilde{B} \\ 1 & 0 & -\tilde{A} & -\tilde{B} \\ 0 & 1 & -\tilde{B} & -\tilde{A} \end{pmatrix} \phi, \quad (92)$$

where $\phi = (\phi_1, \phi_2, \phi_3, \phi_4)^T$ is the spectral function, α , β , r_1 and r_2 are arbitrary constant spectral parameters. When $p_2 = p_1$ and $q_2 = q_1$, (91) and (92) becomes the Lax pair of (89).

3 P_s - T_d - C invariant multi-soliton solutions of the AB-systems.

In the last section, we have listed many new integrable two-place ABSs. Only two special cases, the AB-NLS system (44) with $x_0 = 0$ and the complex AB-MKdV system (51) with $x_0 = t_0 = 0$, have been studied by some authors.

Ablowitz and Musslimani have obtained the singular multiple soliton solutions of the nonlocal focusing equation (44) with $x_0 = 0$ by using the inverse scattering transformation and Riemann-Hilbert problem [1]. Analytical dark and antidark soliton interaction solutions in the defocusing case of (44) with $x_0 = 0$ have been obtained by Li and Xu [13]. The complex soliton solutions of (51) with $x_0 = t_0 = 0$ have been obtained by Ji and Zhu with help of the Darboux transformations [14].

On the other hand, we know that symmetries of a nonlinear system can be used to find new solutions from known ones. The symmetry reduction method is used to find solutions which are invariant under the symmetry transformations.

In the last section, we have established some integrable systems via P_s - T_d - C invariant principles. Thus, a natural interesting question is that *can we find some exact solutions via the P_s - T_d - C symmetries?* Especially, *can we find P_s - T_d - C invariant solutions?* In the following subsections, we list some P_s - T_d - C group invariant solutions for general ABSs including the integrable ones proposed in the last section.

3.1 P_s - T_d -invariant multi-soliton solutions of the KdV-KP-Toda type AB systems

Here, we discuss only the P_s - T_d - C group invariant solutions, thus it is not necessary to restrict the ABSs integrable. We call the models

$$GAB \equiv G(A, B) = 0, \quad B = A^f \quad (93)$$

as the KdV-KP-Toda type ABSs if

$$KKT \equiv G(A, A) = \begin{cases} KdV, \\ KP, \\ Toda, \end{cases} \quad (94)$$

where *KdV*, *KP* and *Toda* are defined by (19), (61) and (71), respectively. Clearly, the AB-KdV, AB-KP and AB-Toda systems listed explicitly above are all special cases of (93) with the condition KKT (94).

Now we focus on the \hat{f} -invariant solutions, i.e.,

$$A^{\hat{f}} = A \quad (95)$$

with

$$\hat{f} = \hat{P}_s \hat{T}_d$$

for the KdV type equations,

$$\hat{f} = \hat{P}_s^{xy} \hat{T}_d, \text{ or } \hat{f} = \hat{P}_s^x \hat{T}_d, \text{ or } \hat{f} = \hat{P}_s^y$$

for the KP type equations, and

$$\hat{f} = \hat{P}_s^n \hat{P}_s^x, \text{ or } \hat{f} = \hat{P}_s^x, \text{ or } \hat{f} = \hat{P}_s^n$$

for the Toda-type equations. In other words, to search for the \hat{f} group invariant solutions of (93) with (94) is equivalent to finding the \hat{f} invariant solutions of the usual KdV, KP and Toda equations.

For the usual KdV, KP and Toda systems, various exact solutions have been found. Consequently, the only thing needs to do is to select the \hat{f} invariant solutions from the known ones of the usual KdV, KP and Toda systems. Here, only the multiple soliton solutions are considered.

It is known that for the KdV (19), KP (61) and Toda (71), their known N -soliton solutions can be uniformly written as

$$A = 2(\ln F)_{xx}, \quad (96)$$

$$F = \sum_{\mu=0,1} \exp \left(\sum_{j=1}^N \mu_j \xi_j + \sum_{1 \leq j < l}^N \mu_j \mu_l \theta_{jl} \right), \quad (97)$$

where the summation of μ should be done for all permutations of $\mu_i = 0, 1, i = 1, 2, \dots, N$ and

$$\xi_j = k_j x - k_j^3 t + \xi_{0j}, \quad \exp(\theta_{jl}) = \left(\frac{k_j - k_l}{k_j + k_l} \right)^2 \quad (98)$$

for the KdV equation,

$$\xi_j = k_j x + p_j y + k_j \left(k_j^2 + \frac{3}{4} p_j^2 \right) t + \xi_{0j}, \quad \exp(\theta_{jl}) = \frac{4(k_j - k_l)^2 - 3(p_j - p_l)^2}{4(k_j + k_l) - 3(p_j - p_l)^2} \quad (99)$$

for the KP equation and

$$\xi_j = k_j n + 2 \sinh \frac{k_j}{2} x + \xi_{0j}, \quad \exp(\theta_{jl}) = \frac{\sinh^2 \frac{1}{4}(k_j - k_l)}{\sinh^2 \frac{1}{4}(k_j + k_l)} \quad (100)$$

for the Toda equation.

It is clear that the multiple soliton solutions (96) with (97) are not explicitly \hat{f} -invariant. To select out the \hat{f} -invariant solutions from (96) with (97), we can use two facts, (i) $\{\xi_{0j}, j = 1, 2, \dots, n\}$ are arbitrary constants, (ii) the solution (96) is invariant under the transformation

$$F \rightarrow \beta \exp(Kx + \Omega t + X_0)F \quad (101)$$

with arbitrary constants β , K , Ω and X_0 .

Because of the arbitrariness of ξ_{0j} , we rewrite ξ_j as

$$\xi_j = \eta_j - \frac{1}{2} \sum_{i=1}^{j-1} \theta_{ij} - \frac{1}{2} \sum_{i=j+1}^N \theta_{ji}, \quad (102)$$

where

$$\eta_j \equiv k_j \left(x - \frac{1}{2}x_0 \right) - k_j^3 \left(t - \frac{1}{2}t_0 \right) + \eta_{0j} \quad (103)$$

for the KdV equation,

$$\eta_j = k_j \left(x - \frac{1}{2}x_0 \right) + p_j \left(y - \frac{1}{2}y_0 \right) + k_j \left(k_j^2 + \frac{3}{4}p_j^2 \right) \left(t - \frac{1}{2}t_0 \right) + \eta_{0j}, \quad (104)$$

for the KP equation, and

$$\eta_j = k_j \left(n - \frac{1}{2}n_0 \right) + 2 \sinh \frac{k_j}{2} \left(x - \frac{1}{2}x_0 \right) + \eta_{0j}, \quad (105)$$

for the Toda equation.

From the expressions (103), (104) and (105), we know that

$$\hat{f}\eta_j = -\eta_j + 2\eta_{0j}. \quad (106)$$

Because of the invariance (101), we take the transformation

$$F \rightarrow \prod_{i < j} K_{ij} \exp \left(\sum_{l=1}^N \eta_l \right) F \quad (107)$$

where

$$K_{ij} = k_i - k_j$$

for the KdV equation,

$$K_{ij} = \sqrt{4(k_i - k_j)^2 - 3(p_i - p_j)^2}$$

for the KP equation and

$$K_{ij} = \sinh \frac{1}{4}(k_i - k_j)$$

for the Toda equation.

By using the transformations (102) and (107), the N -soliton solution (96) becomes

$$A = 2 \left[\ln \sum_{\nu=1,-1} K_\nu \cosh \left(\frac{1}{2} \sum_{j=1}^N \nu_j \eta_j \right) \right]_{xx}, \quad (108)$$

where the summation of $\nu = \{\nu_1, \nu_2, \dots, \nu_N\}$ should be done for all non-dual permutations of $\nu_i = 1, -1, i = 1, 2, \dots, N$, and

$$K_\nu = \prod_{i>j} (k_i - \nu_i \nu_j k_j), \quad (109)$$

for the KdV equation,

$$K_\nu = \prod_{i>j} \sqrt{4(k_i - \nu_i \nu_j k_j)^2 - 3(p_j - \nu_i \nu_j p_l)^2} \quad (110)$$

for the KP equation, and

$$K_\nu = \prod_{i>j} \sinh \frac{1}{4} (k_i - \nu_i \nu_j k_j) \quad (111)$$

for the Toda equation.

In (108), we have used the summation for the non-dual permutation. A pair permutations, $\nu = \{\nu_1, \nu_2, \dots, \nu_N\}$ and $\nu' = \{\nu'_1, \nu'_2, \dots, \nu'_N\}$, are called dual each other if

$$\sum_{j=1}^N \nu_j \xi_j = - \sum_{j=1}^N \nu'_j \xi_j, \quad (112)$$

is true for arbitrary $\xi_j, j = 1, 2, \dots, N$.

From the expression (108) we know that

$$A_{\eta_{0j}=0}^f = \hat{f} A_{\eta_{0j}=0} = A_{\eta_{0j}=0}. \quad (113)$$

Thus the multiple soliton solution $A_{\eta_{0j}=0}$ solves all the KdV-KP-Toda type AB systems (93) with the conditions $\hat{f} = \hat{P}_s^{xy} \hat{T}_d$ for the AB-KP systems, and $\hat{f} = \hat{P}_s^n \hat{P}_s^x$ for the AB-Toda systems. In fact, all the other KdV-KP-Toda types systems including the whole KdV hierarchy, KP hierarchy, Toda hierarchy, Boussinesq hierarchy, 1+1 and 2+1 dimensional Sawada-Kortera hierarchies, 1+1 and 2+1 dimensional Kaup-Kupershmidt hierarchies, and so on, possess the same expression (108) with suitable selections of η_j and K_ν .

Though the expression (108) is equivalent to (96), it appears first time in literature. For the expression (96), some types of resonant singular solutions can be obtained by means of the limiting procedure

$$k_j \rightarrow \pm k_i. \quad (114)$$

However, for the expression (108), the direct limiting procedure of (114) will lead to $(N-1)$ soliton solution because the transformation (107) moves all the possible resonant solutions to ∞ . To recover the resonant solutions, one has to readjust the constants η_{0j} . In other words, if letting the constants η_{0j} depending on $k_i, i = 1, 2, \dots, N$, one may obtain quite different resonant solutions.

3.2 $P_s T_d$ -invariant multi-soliton solutions of the MKdV-sG type AB systems

The most general MKdV-sG system may also be written as (93) but with

$$MS \equiv G(A, A) = \begin{cases} MKdV, \\ sG, \end{cases} \quad (115)$$

where $MKdV$ and sG are defined by (48) and (54), respectively.

Using the same procedure as for the KdV-KP-Toda type AB systems, looking for the $\hat{f} = \hat{P}_s \hat{T}_d$ invariant multiple soliton solutions of the generalized MKdV-sG AB system (93) with (115) is equivalent to finding the \hat{f} -invariant solutions of the MKdV and sG equations.

It is known that the N soliton solutions of the MKdV system (48) and the sG equation (54) possess the common form

$$A = A^{MKdV} = \frac{1}{2} A_x^{sG} = \delta i \left(\ln \frac{F_-}{F_+} \right)_x, \quad \delta^2 = 1 \quad (116)$$

with

$$F_{\pm} = \sum_{\mu=1,0} \exp \left[\sum_{j=1}^N \mu_j \left(\xi_j \pm \frac{i\pi}{2} \right) + \sum_{l < j} \mu_l \mu_j \theta_{lj} \right] \quad (117)$$

where the summation of μ should be done for all permutations of $\mu_i = 0, 1, i = 1, 2, \dots, N$ and

$$\xi_j = k_j x - k_j^3 t + \xi_{0j}, \quad \exp(\theta_{jl}) = \left(\frac{k_j - k_l}{k_j + k_l} \right)^2 \quad (118)$$

for the MKdV equation and

$$\xi_j = k_j x + k_j^{-1} t + \xi_{0j}, \quad \exp(\theta_{jl}) = \left(\frac{k_j - k_l}{k_j + k_l} \right)^2 \quad (119)$$

for the sG equation.

From the expression (116), we know that the solution is invariant under the transformation

$$F_{\pm} \rightarrow \beta \exp(Kx + \Omega t + X_0) F_{\pm} \quad (120)$$

with arbitrary constants $K, \beta, \Omega,$ and X_0 .

Using the invariance (120) and the arbitrariness of ξ_{0j} , we can rewrite the solution (120) as

$$A = \begin{cases} \pm 2 \frac{\partial}{\partial x} \tan^{-1} \frac{\sum_{\nu_e} K_{\nu} \sinh(\sum_{j=1}^N \nu_j \eta_j)}{\sum_{\nu_o} K_{\nu} \cosh(\sum_{j=1}^N \nu_j \eta_j)}, \\ \pm 2 \frac{\partial}{\partial x} \tan^{-1} \frac{\sum_{\nu_o} K_{\nu} \sinh(\sum_{j=1}^N \nu_j \eta_j)}{\sum_{\nu_e} K_{\nu} \cosh(\sum_{j=1}^N \nu_j \eta_j)}, \\ \pm 2 \frac{\partial}{\partial x} \tan^{-1} \frac{\sum_{\nu_e} K_{\nu} \cosh(\sum_{j=1}^N \nu_j \eta_j)}{\sum_{\nu_o} K_{\nu} \sinh(\sum_{j=1}^N \nu_j \eta_j)}, \\ \pm 2 \frac{\partial}{\partial x} \tan^{-1} \frac{\sum_{\nu_o} K_{\nu} \cosh(\sum_{j=1}^N \nu_j \eta_j)}{\sum_{\nu_e} K_{\nu} \sinh(\sum_{j=1}^N \nu_j \eta_j)} \end{cases} \quad (121)$$

via the similar trick as in the KdV-KP-Toda case, where the summation of ν_o should be done for all non-dual odd permutations of $\nu_i = 1, -1, i = 1, 2, \dots, N$ with odd number of $\nu_i = 1$, the summation of ν_e should be done for all non-dual even permutations of $\nu_i = 1, -1, i = 1, 2, \dots, N$ with even number of $\nu_i = 1$,

$$K_{\nu} \equiv \prod_{i>j} (k_i - \nu_i \nu_j k_j), \quad (122)$$

and η_j being defined as

$$\eta_j = k_j \left(x - \frac{1}{2} x_0 \right) - k_j^3 \left(t - \frac{1}{2} t_0 \right) + \eta_{0j} \quad (123)$$

for the MKdV equation, and

$$\eta_j = k_j \left(x - \frac{1}{2}x_0 \right) + k_j^{-1} \left(t - \frac{1}{2}t_0 \right) + \eta_{0j} \quad (124)$$

for the sG equation. Four expressions of (121) are equivalent by selecting suitable arbitrary constants η_{0j} and using periodic property of the triangular functions and hyperbolic functions.

In (121), we have used summations on non-dual odd (even) permutations (the permutations with odd (even) numbers of $\nu_j = 1$). We should add the permutation with zero number of $\nu_j = 1$ to the even permutations. For odd N (for odd number of solitons), even permutations ν_e and odd permutations ν_o are dual each other, say, for $N = 3$,

$$\begin{aligned} \nu_e &= [\nu_1, \nu_2, \nu_3]_e = \{[-1, -1, -1], [1, 1, -1], [1, -1, 1], [-1, 1, 1]\} \\ \nu_o &= [\nu_1, \nu_2, \nu_3]_o = \{[1, 1, 1], [-1, -1, 1], [-1, 1, -1], [1, -1, -1]\} = \nu_e^{dual}. \end{aligned} \quad (125)$$

For even N , ν_e and ν_o are completely different and not dual. For example, when $N = 4$, the non-dual even and odd permutations possess the forms

$$\nu_o = [\nu_1, \nu_2, \nu_3, \nu_4]_o = \{[1, -1, -1, -1], [-1, 1, -1, -1], [-1, -1, 1, -1], [-1, -1, -1, 1]\}, \quad (126)$$

and

$$\nu_e = [\nu_1, \nu_2, \nu_3, \nu_4]_e = \{[1, 1, 1, 1], [1, 1, -1, -1], [1, -1, 1, -1], [1, -1, -1, 1]\}, \quad (127)$$

respectively. The remaining permutations are duals of (126) and (127).

From the expression (121) we know that

$$A_{\eta_{0j}=0}^f = -A_{\eta_{0j}=0}. \quad (128)$$

Therefore, the multiple soliton solutions $A_{\eta_{0j}=0}$ i.e., (121) with $\eta_{0j} = 0$, solve the MKdV-sG AB systems (93) with (115) including (51) with $\sigma = 1$ and (58) with $\{a = \delta = 1\}$.

3.3 $P_s T_d C$ -invariant and $P_s C$ -invariant multi-soliton solutions of the NLS type AB systems

We assume the generalized NLS-type AB systems have the following forms

$$G(A, B) = 0, \quad B = \hat{P}_s \hat{T}_d A = A(-x + x_0, -t + t_0), \text{ or } B = \hat{P}_s A = A(-x + x_0, t) \quad (129)$$

with

$$G(A, A^*) = NLS, \quad (130)$$

and NLS being defined by (40). It is clear that the integrable AB-NLS systems $ABNLS_1$ and $ABNLS_2$ expressed by (44) and (45) are just two explicit examples of (129).

From the condition (130), we know that if A is a $\hat{P}_s \hat{T}_d \hat{C}$ invariant solution of the usual NLS equation, then it is also a solution for all general NLS type AB systems (129) with $\hat{f} = \hat{P}_s \hat{T}_d$.

3.3.1 N-soliton solutions of the defocusing AB-NLS systems

The known N -soliton solutions for the defocusing NLS system (40) with $\sigma = -1$ possesses the form

$$A = \sqrt{2}\alpha \exp(-i\alpha^2 t + i\phi_0) \frac{\sum_{\mu} \exp\left(\sum_{j=1}^N \mu_j (\xi_j + 2i\theta_j) + \sum_{j<l} \mu_j \mu_l \theta_{jl}\right)}{\sum_{\mu} \exp\left[\sum_{j=1}^N \mu_j \xi_j + \sum_{j<l} \mu_j \mu_l \theta_{jl}\right]}, \quad (131)$$

where

$$\xi_j = \sqrt{2}\alpha \sin(\theta_j) \left(x + \sqrt{2}\alpha \cos(\theta_j)t \right) + \xi_{0j}, \quad (132)$$

and

$$\exp(\theta_{jl}) = \left(\frac{\sin \frac{1}{2}(\theta_j - \theta_l)}{\sin \frac{1}{2}(\theta_j + \theta_l)} \right)^2 \quad (133)$$

with arbitrary real constants α , ξ_{0j} , θ_j , $j = 1, 2, \dots, N$ and ϕ_0 . The summations on μ in the solution (131) should be done for all permutations $\mu_j = 0, 1, j = 1, 2, \dots, N$.

Similar to the KdV-KP-Toda case, the N-soliton solution (131) can also be re-written to an elegant form,

$$A = \sqrt{2}\alpha \exp \left[-i\alpha^2 \left(t - \frac{1}{2}t_0 \right) + i\phi'_0 \right] \frac{\sum_{\nu} A_{\nu} \cosh \left[\frac{1}{2} \sum_{j=1}^N \nu_j (\eta_j + 2i\theta_j) \right]}{\sum_{\nu} A_{\nu} \cosh \left(\frac{1}{2} \sum_{j=1}^N \nu_j \eta_j \right)}, \quad (134)$$

where the summations on ν should be done for all possible non-dual permutations $\nu_j = 1, -1, j = 1, 2, \dots, N$,

$$\eta_j = \sqrt{2}\alpha \sin(\theta_j) \left[x - \frac{1}{2}x_0 + \sqrt{2}\alpha \cos(\theta_j) \left(t - \frac{1}{2}t_0 \right) \right] + \eta_{0j}, \quad (135)$$

$$A_{\nu} = \prod_{l < j} \sin \frac{\theta_j - \nu_j \nu_l \theta_l}{2}, \quad (136)$$

and θ_j , η_{0j} , α and ϕ'_0 are arbitrary real constants.

The expression (134) with (135) and (136) is $\hat{P}_s \hat{T}_d \hat{C}$ invariant if $\phi'_0 = \eta_{0j} = 0$. Thus, from the condition (130) we know that $A_{\eta_{0j}=\phi'_0=0}$ solves all the defocusing AB-NLS systems (129) with $\hat{f} = \hat{P}_s \hat{T}_d$. However, for the defocusing AB-NLS (129) with $\hat{f} = \hat{P}_s \hat{C}$, we can not find nontrivial multi-soliton solutions from (134). The fact indicates that there may exist symmetry breaking multi-soliton solutions. It is remarkable that the $\hat{P}_s \hat{T}_d \hat{C}$ symmetry breaking multi-soliton solutions have been obtained for the defocusing AB-NLS system (44) with $x_0 = 0$ by means of the Daboux transformation [13].

3.3.2 N-soliton solutions of the focusing AB-NLS systems

The known N -soliton solution for the focusing NLS system (40) with $\sigma = 1$ possesses the form

$$A = \sqrt{2} \frac{\sum_{\mu >} \exp \left(\sum_{j=1}^{2N} \mu_j \xi_j + \sum_{j < l} \mu_j \mu_l \theta_{jl} \right)}{\sum_{\mu =} \exp \left(\sum_{j=1}^{2N} \mu_j \xi_j + \sum_{j < l} \mu_j \mu_l \theta_{jl} \right)}, \quad (137)$$

where

$$\xi_j = \begin{cases} k_j x + ik_j^2 t + \xi_{0j}, & j = 1, 2, \dots, N, \\ k_j^* x - ik_j^{*2} t + \xi_{0j}^*, & j = N+1, \dots, 2N, \end{cases} \quad (138)$$

$$\begin{aligned} \exp(\theta_{jl}) &= 2(k_j - k_l)^2, \quad j, l = 1, 2, \dots, N, \\ \exp(\theta_{j(N+l)}) &= \frac{1}{2(k_j + k_l^*)^2}, \quad j, l = 1, 2, \dots, N, \\ \theta_{(N+j)(N+l)} &= \theta_{jl}^*, \end{aligned} \quad (139)$$

with arbitrary complex constants $k_{0j}, \xi_{0j}, j = 1, 2, \dots, N$.

The summations on $\mu_=>$ and $\mu_>$ in the solution (137) should be done for all permutations $\mu_j = 0, 1$ with the conditions

$$\sum_{j=1}^N \mu_j = \sum_{j=1}^N \mu_{N+j}, \quad (140)$$

and

$$\sum_{j=1}^N \mu_j = 1 + \sum_{j=1}^N \mu_{N+j}, \quad (141)$$

respectively.

It is clear that the solution (137) is not $\hat{P}_s\hat{T}_d\hat{C}$ invariant for an arbitrary ξ_{0j} . Thus it is not a solution of the general focusing NLS type AB system (129) with $\sigma = 1$.

In a similar way, to find $\hat{P}_s\hat{T}_d\hat{C}$ invariant solutions from (137), the only thing to do is to rewrite ξ_j as

$$\begin{aligned} \xi_j &= k_j \left(x - \frac{1}{2}x_0 \right) + ik_j \left(t - \frac{1}{2}t_0 \right) + \eta_{0j} - \frac{1}{2} \sum_{i=1}^{i-1} \theta_{ji} - \frac{1}{2} \sum_{i=j+1}^{2n} \theta_{ij} \\ &= \eta_j - \frac{1}{2} \sum_{i=1}^{i-1} \theta_{ji} - \frac{1}{2} \sum_{i=j+1}^{2n} \theta_{ij}. \end{aligned} \quad (142)$$

One can prove that the solution (137) with (142) is P_sT_dC invariant for

$$\eta_{0j} = 0 \quad (143)$$

and P_sC invariant if

$$\eta_{0j} = k_j - k_j^* = 0. \quad (144)$$

Thus the solution $A_{\eta_{0j}=0}$ (i.e., (137) with (142) and (143)) solves the general focusing NLS AB system (129) with $\{B = A^{P_sT_d}, \sigma = 1\}$, and $A_{\eta_{0j}=k_j-k_j^*=0}$ (i.e., (137) with (142) and (144)) solves the general focusing NLS AB system (129) with $\{B = A^{P_s}, \sigma = 1\}$.

The expression (137) can also be expressed in terms of hyperbolic and triangle functions to reveal explicit $\hat{P}_s\hat{T}_d\hat{C}$ invariants. However, we do not write them down here because the final expression is much more ugly than (137).

As an example, we write down the $N = 2$ case to demonstrate the explicit $\hat{P}_s\hat{T}_d\hat{C}$ invariance. In this case, the solution (137) can be rewritten as ($k_1 \equiv k_{1r} + ik_{1i}, k_2 \equiv k_{2r} + ik_{2i}$)

$$A_{N=2} = \frac{2\sqrt{2}|k_1 - k_2||k_1 + k_2^*| [k_{1r}e^{i\eta_{1i}} \cos(\alpha + \beta + i\eta_{2r}) + k_{2r}e^{i\eta_{2i}} \cos(\alpha - \beta + i\eta_{1r})]}{|k_1 - k_2|^2 \cosh(\eta_{1r} + \eta_{2r}) + |k_1 + k_2^*|^2 \cosh(\eta_{1r} - \eta_{2r}) + 4k_{1r}k_{2r} \cos(\eta_{1i} - \eta_{2i} + 4\beta)}, \quad (145)$$

with

$$\begin{aligned} \eta_{1i} &= k_{1i} \left(x - \frac{1}{2}x_0 \right) + (k_{1r}^2 - k_{1i}^2) \left(t - \frac{1}{2}t_0 \right) + \eta_{01i} = -\frac{i}{2}(\eta_1 - \eta_1^*), \\ \eta_{2i} &= k_{2i} \left(x - \frac{1}{2}x_0 \right) + (k_{2r}^2 - k_{2i}^2) \left(t - \frac{1}{2}t_0 \right) + \eta_{02i}, \\ \eta_{1r} &= k_{1r} \left(x - \frac{1}{2}x_0 \right) - 2k_{1r}k_{1i} \left(t - \frac{1}{2}t_0 \right) + \eta_{01r} = \frac{1}{2}(\eta_1 + \eta_1^*), \\ \eta_{2r} &= k_{2r} \left(x - \frac{1}{2}x_0 \right) - 2k_{2r}k_{2i} \left(t - \frac{1}{2}t_0 \right) + \eta_{02r}, \end{aligned} \quad (146)$$

and

$$\begin{aligned}\alpha &= \arctan\left(\frac{k_{1i} - k_{2i}}{k_{1r} - k_{2r}}\right), \\ \beta &= \arctan\left(\frac{k_{1i} - k_{2i}}{k_{1r} + k_{2r}}\right).\end{aligned}\quad (147)$$

From (145) with (146) and (147), it is straightforward to find that $A_{N=2}$ is $\hat{P}_s \hat{T}_d \hat{C}$ invariant only for $\eta_{0j} = 0$, $j = 1, 2$, and $\hat{P}_s \hat{C}$ invariant if $\eta_{0j} = k_j - k_j^* = 0$, $j = 1, 2$.

3.4 $\hat{P}_s^u \hat{P}_s^n \hat{P}_s^m$ invariant multi-soliton solutions of a special H_1 system

A general AB- H_1 system may have the form

$$G(A, B) = 0, \quad B = \hat{f}A, \quad \hat{f} \in \hat{F} \quad (148)$$

with the condition

$$G(A, A) = (A - \hat{A})(\tilde{A} - \hat{A}) + q - p = 0 \quad (149)$$

for suitable $\hat{f} \in \hat{F}$.

It is clear that the special AB- H_1 system (89) with $\sigma = 1$, $q_1 = 2q$, $p_1 = 2p$ is just a special case of (148) and (149) with

$$\hat{f} = \hat{P}_s^u \hat{P}_s^n \hat{P}_s^m. \quad (150)$$

Therefore, to look for the $\hat{P}_s^u \hat{P}_s^n \hat{P}_s^m$ invariant solutions of the AB- H_1 equation (148) is equivalent to finding the $\hat{P}_s^u \hat{P}_s^n \hat{P}_s^m$ invariant solution of the usual H_1 system (149).

For the usual H_1 equation, its N -soliton solution possesses the form [16]

$$A = an + bm + \gamma - \frac{h}{g} \quad (151)$$

where $a = \sqrt{p}$ and $b = \sqrt{q}$,

$$\begin{cases} g = |\psi, E_3\psi, E_3^2\psi, \dots, E_3^{N-1}\psi|, \\ h = |\psi, E_3\psi, \dots, E_3^{N-2}\psi, E_3^N\psi|, \\ \psi = \psi(n, m, l) = (\psi_1, \dots, \psi_n)^T, \quad E_3\psi = \psi(n, m, l+1), \\ \psi_i = \psi_i(n, m, l) = \rho_i^+(a+k_i)^n (b+k_i)^m k_i^l + \rho_i^-(a-k_i)^n (b-k_i)^m (-k_i)^l, \end{cases} \quad (152)$$

ρ_i^\pm and k_i are arbitrary constant parameters.

In the solution (151) with (152), the arbitrary constants ρ_i^+/ρ_i^- indicate the positions of N solitons are arbitrary and these freedoms are introduced by the translation invariance of n and m . However, the AB- H_1 systems are not $\{n, m\}$ translation invariant. Thus, the solution (151) with (152) for arbitrary ρ_i^+/ρ_i^- is not a solution of the AB- H_1 system.

It is interesting that if we rewrite the constants ρ_i^+/ρ_i^- as

$$\frac{\rho_i^+}{\rho_i^-} = \left(\frac{a-k_i}{a+k_i}\right)^{\frac{n_0}{2}} \left(\frac{b-k_i}{b+k_i}\right)^{\frac{m_0}{2}} e^{\xi_{0i}} \quad (153)$$

with arbitrary ξ_{i0} , the solution (151) can be rewritten in an elegant form

$$\begin{aligned}A &= an + bm + \gamma - [\ln(g)]_n = an + bm + \gamma - \frac{g_n}{g} \\ &= an + bm + \gamma - \frac{\sum_\nu K_\nu \sum_{j=1}^N \nu_j k_j \sinh\left(\sum_{i=1}^N \frac{1}{2} \nu_i \xi_i\right)}{\sum_\nu K_\nu \cosh\left(\sum_{i=1}^N \frac{1}{2} \nu_i \xi_i\right)}\end{aligned}\quad (154)$$

where the summation of ν should be done for all non-dual permutations of $\nu_i = 1, -1, i = 1, 2, \dots, N$ while the sinh and cosh functions are defined by

$$\begin{aligned}\sinh \frac{\xi_i}{2} &= \left(\frac{a+k_i}{a-k_i} \right)^{\frac{n}{2}-\frac{n_0}{4}} \left(\frac{b+k_i}{b-k_i} \right)^{\frac{m}{2}-\frac{m_0}{4}} e^{\frac{\xi_{i0}}{2}} - \left(\frac{a+k_i}{a-k_i} \right)^{\frac{n_0}{4}-\frac{n}{2}} \left(\frac{b+k_i}{b-k_i} \right)^{\frac{m_0}{4}-\frac{m}{2}} e^{-\frac{\xi_{i0}}{2}}, \\ \cosh \frac{\xi_i}{2} &= \left(\frac{a+k_i}{a-k_i} \right)^{\frac{n}{2}-\frac{n_0}{4}} \left(\frac{b+k_i}{b-k_i} \right)^{\frac{m}{2}-\frac{m_0}{4}} e^{\frac{\xi_{i0}}{2}} + \left(\frac{a+k_i}{a-k_i} \right)^{\frac{n_0}{4}-\frac{n}{2}} \left(\frac{b+k_i}{b-k_i} \right)^{\frac{m_0}{4}-\frac{m}{2}} e^{-\frac{\xi_{i0}}{2}}, \\ \sinh(\xi_1 + \xi_2) &= \sinh(\xi_1) \cosh(\xi_2) + \sinh(\xi_2) \cosh(\xi_1), \\ \cosh(\xi_1 + \xi_2) &= \cosh(\xi_1) \cosh(\xi_2) + \sinh(\xi_2) \sinh(\xi_1).\end{aligned}\tag{155}$$

Now, it is straightforward to find that the solution (154) is $\hat{P}_s^u \hat{P}_s^n \hat{P}_s^m$ invariant if $\xi_{i0} = 0$ for all $i = 1, 2, \dots, N$. In other words, $A_{\xi_{i0}=0}$ expressed by (154) solves the AB- H_1 systems (148) (including (89)) for $p_1 = 2p, q = 2q$ with $\hat{f} = \hat{P}_s^u \hat{P}_s^n \hat{P}_s^m$.

4 Conclusions and discussions

In summary, various intrinsic two-place models, named Alice-Bob systems for convenience, are proposed via the AB-BA equivalence principle and $\hat{P}_s\text{-}\hat{T}_d\text{-}\hat{C}$ principle. All the ABSs are appeared first time in this paper except for two special cases, the AB-NLS (44) with $x_0 = 0$ and the AB-MKdV (51) with $x_0 = t_0 = 0$.

Though the concept of the ABSs is first proposed, all the concrete ABSs listed in this paper are reductions of known coupled systems. Here, many types of integrable systems such as the KdV, MKdV, KP, Toda, sG, NLS and H_1 equations are extended to some types of ABSs. However, in principle, any one place physical model can be extended to ABS no matter whether it is integrable or not.

The $\hat{P}\text{-}\hat{T}\text{-}\hat{C}$ principles are important in many physical fields and various physically important models possess P-T-C invariance. In this paper, the parity \hat{P} and time reversal \hat{T} are generalized to shifted parity \hat{P}_s and delayed time reversal \hat{T}_d . Thus, the $\hat{P}\text{-}\hat{T}\text{-}\hat{C}$ principles are extended to the $\hat{P}_s\text{-}\hat{T}_d\text{-}\hat{C}$ principles.

The $\hat{P}_s\text{-}\hat{T}_d\text{-}\hat{C}$ principles are used not only to establish ABSs but also to solve ABSs. Especially, some types of multi-soliton solutions of many ABSs have been obtained from well-known integrable models via the $\hat{P}_s\text{-}\hat{T}_d\text{-}\hat{C}$ principles. The $\hat{P}_s\text{-}\hat{T}_d\text{-}\hat{C}$ invariant multiple soliton solutions are valid also for many nonintegrable ABSs. For instance, the AB-KdV system (25), is not integrable for general a_i , however, its multiple soliton solution can be obtained in terms of (108) via a simple scaling procedure.

In addition to the $\hat{P}_s\text{-}\hat{T}_d\text{-}\hat{C}$ invariant multiple soliton solutions of many ABSs, more solutions can be obtained to be $\hat{P}_s\text{-}\hat{T}_d\text{-}\hat{C}$ symmetry breaking. For instance, for the defocusing AB-NLS system (44) with $\sigma = -1$ and $x_0 = 0$, the \hat{P} -invariant solutions of the usual defocusing NLS are also the solutions of the defocusing $ABNLS_1$. Though we have not yet found the \hat{P} invariant solution of $ABNLS_1$, there really exist other types of multiple soliton solutions which are not \hat{P} -invariant [1, 13].

In [13], Li and Xu have obtained the explicit form of non-parity invariant multiple soliton solutions of the defocusing $ABAKNS_1$ with $\sigma = -2$ (where 2 can be scaled to 1) and $x_0 = 0$ by means of the Darboux transformation, say, the two soliton solution possesses the form [13]

$$A = |\kappa| e^{2i|\kappa|^2 t + i\phi_0} \left\{ 1 - \frac{2\kappa_{1r} (\gamma + e^{-2\kappa_i(x+2\kappa_r t)}) (\kappa^* \gamma^* + \kappa e^{2\kappa_i(x-2\kappa_r t)})}{|\kappa|^2 |\gamma|^2 + |\kappa|^2 e^{-8\kappa_i \kappa_r t} + \kappa_r (\kappa \gamma e^{2\kappa_i(x-2\kappa_r t)} + \kappa^* \gamma^* e^{-2\kappa_i(x+2\kappa_r t)})} \right\}, \tag{156}$$

with complex constants $\kappa \equiv \kappa_r + i\kappa_i, \gamma$ and the real constant ϕ_0 .

Thus, the following important question should be studied in future: *Are there any other types of multiple soliton solutions which may be $P_s\text{-}T_d\text{-}C$ invariant or $P_s\text{-}T_d\text{-}C$ symmetry breaking for the AB systems listed in this paper?*

Another important question which should be studied further is *can we establish other types of AB systems with and without AB-BA equivalence principle?* The answer is YES. Here, we just write down a special AB-sG system by means of the AB-coordinate (10),

$$A_{xx} - A_{yy} = 2 \sin \frac{A}{2} \cos \frac{A^f}{2}, \quad (157)$$

with

$$A^f = A(x', y'), \quad (158)$$

$$x' = \coth(\theta)x + \operatorname{csch}(\theta)y + \exp\left(-\frac{1}{2}\theta\right)x_0, \quad (159)$$

$$y' = -\operatorname{csch}(\theta)x - \coth(\theta)y - \exp\left(\frac{1}{2}\theta\right)x_0.$$

The AB-sG equation (157) is also integrable. The Lax pair of (157) possesses the form

$$\begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \\ \psi_4 \end{pmatrix}_x = \begin{pmatrix} -\lambda - C_C & -iS_S & A_{x_1} + S_C & -iA_{x_1}^f - iC_S \\ iS_S & -\lambda - C_C & iA_{x_1}^f + iC_S & +A_{x_1} + S_C \\ S_C - A_{x_1} & iA_{x_1}^f - iC_S & \lambda + C_C & iS_S \\ iC_S - iA_{x_1}^f & S_C - A_{x_1} & -iS_S & \lambda + C_C \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \\ \psi_4 \end{pmatrix}, \quad (160)$$

and

$$\begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \\ \psi_4 \end{pmatrix}_y = \begin{pmatrix} -\lambda + C_C & iS_S & A_{x_1} - S_C & -iA_{x_1}^f + iC_S \\ -iS_S & -\lambda + C_C & iA_{x_1}^f - iC_S & A_{x_1} - S_C \\ -S_C - A_{x_1} & iA_{x_1}^f + iC_S & \lambda - C_C & -iS_S \\ -iC_S - iA_{x_1}^f & -S_C - A_{x_1} & iS_S & \lambda - C_C \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \\ \psi_4 \end{pmatrix}, \quad (161)$$

where

$$\begin{aligned} C_C &= \frac{1}{16\lambda} \cos \frac{A}{2} \cos \frac{A^f}{2}, & S_S &= \frac{1}{16\lambda} \sin \frac{A}{2} \sin \frac{A^f}{2}, \\ C_S &= \frac{1}{16\lambda} \cos \frac{A}{2} \sin \frac{A^f}{2}, & S_C &= \frac{1}{16\lambda} \sin \frac{A}{2} \cos \frac{A^f}{2}, \\ A_{x_1} &= \frac{1}{8}(A_x + A_y), & A_{x_1}^f &= \frac{1}{8}(A_x^f + A_y^f), \end{aligned} \quad (162)$$

and λ is a constant spectral parameter.

We do not discuss the model (157) further and leave the studies on the ABSs with different types of AB coordinates such as (10), (13) and (14) in future.

Acknowledgement. The authors is in debt to Profs. J. F. He, Z. N. Zhu, Y. Chen, J. Lin, D. J. Zhang, X. B. Hu, Q. P. Liu, Y. Q. Li and X. Y. Tang for their helpful discussions. The work was sponsored by the Global Change Research Program of China (No.2015CB953904), the National Natural Science Foundations of China (Nos. 11435005, 11175092, and 11205092), Shanghai Knowledge Service Platform for Trustworthy Internet of Things (No. ZF1213) and K. C. Wong Magna Fund in Ningbo University.

References

- [1] M. J. Ablowitz and Z. H. Musslimani, Phys. Rev. Lett. 110 (2013) 064105.

- [2] D. J. Korteweg and G. de Vries, *Philosophical Magazine* 39 (1985) 422.
- [3] D. G. Crighton, *Appl. Math.* 39 (1995) 39
- [4] S.Y. Lou, B. Tong, H.C. Hu, X.Y. Tang, *J. Phys. A: Math. Gen.* 39 (2006) 513527.
- [5] M. Antonowicz, A. P. Fordy, *Phys. D*, 28 (1987) 345.
- [6] A. Karasu, *J. Math. Phys.* 38 (1997) 3616.
- [7] M. V. Foursov, *J. Math. Phys.* 44 (2003) 3088.
- [8] A. G. Meshkov, M. J. Balakhnev, *SIGMA*, 4 (2008) 018; Z. Popowicz, *SIGMA*, 6 (2010) 018.
- [9] D. S. Wang, *Nonlinear Analysis*, 73 (2010) 270281; D. S. Wang, *Appl. Math. Lett.* 23 (2010) 665.
- [10] S. Baker, V.Z. Enolskii, A.P. Fordy, *Phys. Lett. A* 201 (1995) 167.
- [11] R. Hirota, and J. Satsuma, *Phys. Lett. A*, 85 (1981) 407;
- [12] R. Dodd and A. Fordy, *Phys. Lett. A*, 89 (1982) 168.
- [13] M. Li and T. Xu, *Phys. Rev. E*. 91 (2015) 033202.
- [14] J. L. Ji and Z. N. Zhu, *On a nonlocal modified Korteweg-de Vries equation: integrability, Darboux transformation and soliton solutions*, Preprint, (2015).
- [15] V. E. Adler, A. I. Bobenko and Y. B. Suris, *Commun. Math. Phys.* 233 (2002) 513.
- [16] J. Hietarinta and D. J. Zhang, *J. Phys. A: Math. Theor.* 42 (2009) 404006.