INVERTIBLE SUMS OF MATRICES

JUSTIN CHEN

ABSTRACT. We give an elementary proof of a Caratheodory-type result on the invertibility of a sum of matrices, due first to Facchini and Barioli. The proof yields a polynomial identity, expressing the determinant of a large sum of matrices in terms of determinants of smaller sums. Interpreting these results over an arbitrary commutative ring gives a stabilization result for a filtered family of ideals of determinants. Generalizing in another direction gives a characterization of local rings. An analogous result for semilocal rings is also given – interestingly, the semilocal case reduces to the case of matrices.

Let k be a field, and let $A_1, \ldots, A_m \in M_n(k)$ be square matrices over k, for fixed $n, m \in \mathbb{N}$. Write [m] for the index set $\{1, \ldots, m\}$, and for a subset $S \subseteq [m]$, write |S| for the cardinality of S.

Theorem 1. If $A_1 + \ldots + A_m$ is invertible, then there exists $S \subseteq [m]$ with $|S| \le n$ such that $\sum_{i \in S} A_i$ is invertible.

Notice that the upper bound $p \leq n$ is the best possible: for the *n* elementary matrices E_{ii} , $i = 1, \ldots, n$, $E_{11} + \ldots + E_{nn} = id_n$ is invertible, but any sum of $\leq n-1$ of the E_{ii} is not invertible.

In proving Theorem 1, we may assume m > n (if $m \le n$, then take S = [m]), which we do henceforth. With this, Theorem 1 is then a consequence of the following polynomial identity:

Lemma 2. Let $T := k[x_{i\beta\gamma} \mid 1 \le i \le m, 1 \le \beta, \gamma \le n]$ be a polynomial ring over k in mn^2 indeterminates, and let $M_i := (x_{i\beta\gamma})_{\beta,\gamma=1}^n \in M_n(T)$ be generic matrices, for i = 1, ..., m. If m > n, then as polynomials (i.e. elements of T),

$$\sum_{\subseteq [m]} (-1)^{|S|} \det\left(\sum_{i \in S} M_i\right) = 0.$$

In particular,

$$\det\left(\sum_{i=1}^{m} M_i\right) \in \left(\det\left(\sum_{i \in S} M_i\right) \mid S \subseteq [m], |S| \le n\right).$$

Lemma 2 implies Theorem 1: Suppose there exist $A_1, \ldots, A_m \in M_n(k)$ with $A_1 + \ldots + A_m$ invertible but every sum of $\leq n$ of the A_i 's is not invertible. Write $A_i =: (a_{i\beta\gamma})_{\beta,\gamma=1}^n$ for $i = 1, \ldots, m$ (so $a_{i\beta\gamma} \in k$). Let $\varphi : T \to k$, $\varphi(x_{i\beta\gamma}) = a_{i\beta\gamma}$ be the evaluation map, inducing $\widetilde{\varphi} : M_n(T) \to M_n(k)$, $\widetilde{\varphi}(M_i) = A_i$. Since det is a polynomial in the entries of a matrix, $\varphi(\det(\sum_{i \in S} M_i)) = \det(\widetilde{\varphi}(\sum_{i \in S} M_i)) = \det(\sum_{i \in S} A_i)$ for

S

²⁰¹⁰ Mathematics Subject Classification. 15A09, 13H99.

every $S \subseteq [m]$. Applying φ to the containment in Lemma 2 implies that the nonzero element det $\left(\sum_{i=1}^{m} A_i\right)$ is contained in the ideal $\left(\det\left(\sum_{i\in S} A_i\right) \mid S \subseteq [m], |S| \le n\right)$ of k, but each generator is 0, a contradiction. Thus no such A_i 's can exist. \Box

The polynomial identity of Lemma 2 follows in turn from a combinatorial identity:

Lemma 3. For $m, n \in \mathbb{N}$, m > n, let $\{z_{i,j} \mid 1 \leq i \leq m, 1 \leq j \leq n\}$ be a set of mn commuting indeterminates. Then (as polynomials in $\mathbb{Z}[z_{ij}]$)

$$\sum_{S \subseteq [m]} (-1)^{|S|} \prod_{j=1}^{n} \sum_{i \in S} z_{i,j} = 0.$$

Lemma 3 implies Lemma 2: By definition, for an $n \times n$ matrix $Y = (y_{\beta\gamma})_{\beta,\gamma=1}^n$,

$$\det Y = \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) \prod_{j=1}^n y_{j\sigma(j)}.$$

Hence

$$\sum_{S\subseteq[m]} (-1)^{|S|} \det\left(\sum_{i\in S} M_i\right) = \sum_{S\subseteq[m]} (-1)^{|S|} \sum_{\sigma\in S_n} \operatorname{sgn}(\sigma) \prod_{j=1}^n \sum_{i\in S} x_{ij\sigma(j)}$$
$$= \sum_{\sigma\in S_n} \operatorname{sgn}(\sigma) \sum_{S\subseteq[m]} (-1)^{|S|} \prod_{j=1}^n \sum_{i\in S} x_{ij\sigma(j)}$$

so for fixed $\sigma \in S_n$, setting $z_{i,j} := x_{ij\sigma(j)}$ in Lemma 3 gives the desired vanishing. The second statement of Lemma 2 follows from the first, by induction on m: the identity yields det $\left(\sum_{i=1}^{m} M_i\right) \in \left(\det\left(\sum_{i\in S} M_i\right) \mid S \subseteq [m], |S| < m\right)$, which implies both the base case (by taking m = n + 1) and the inductive step. \Box

Proof of Lemma 3: As every monomial in the sum is of the form $z_{i_1,1} \ldots z_{i_n,n}$ for some $i_j \in [m]$, it suffices to show that the coefficient of $z_{i_1,1} \ldots z_{i_n,n}$ is 0, for any fixed choice of $i_1, \ldots, i_n \in [m]$. Now $z_{i_1,1} \ldots z_{i_n,n}$ appears for a particular $S \subseteq [m]$ iff $\{i_1, \ldots, i_n\} \subseteq S$, and for such an S, $z_{i_1,1} \ldots z_{i_n,n}$ appears exactly once, with coefficient $(-1)^{|S|}$. Thus the coefficient of $z_{i_1,1} \ldots z_{i_n,n}$ is

$$\sum_{\{i_1,\dots,i_n\}\subseteq S\subseteq [m]} (-1)^{|S|} = \sum_{\substack{S'\subseteq [m]\setminus\{i_1,\dots,i_n\}}} (-1)^{|S'|+n}$$
$$= \sum_{l=0}^{m-n} \binom{m-n}{l} (-1)^{l+n}$$
$$= (-1)^n (1-1)^{m-n} = 0.$$

Remark. It has come to our attention that the statement of Theorem 1 has in fact appeared before, posed as a problem with accompanying solution in [4]. Also, a version of the first statement of Lemma 2 can be found in [3, Theorem 2.2]. The solution given in [4] follows a slightly different approach than the proof given here, as well as using different lemmas. We have chosen to present the reasoning here for its originality, and to emphasize the simple yet pleasing proof of Lemma 3.

It should be noted that the theorems of both [3] and [4] are stated only for fields. To the best of our knowledge, the ideal-theoretic results below have not been observed before. Other studies of determinants (and characteristic polynomials) of sums of matrices can be found in [1], [6].

Now let R be a ring (henceforth all rings, except for matrix rings, are always commutative with $1 \neq 0$). It is natural to ask: to what extent do the above results generalize to $M_n(R)$? We first give a generalization of Lemma 2:

Proposition 4. Let R be a ring, $n \in \mathbb{N}$, I an index set (possibly infinite). For any collection of matrices $\{A_i \mid i \in I\} \subseteq M_n(R)$, consider the R-ideals

$$I_j := \left(\det\left(\sum_{i \in S} A_i\right) \mid S \subseteq I, |S| \le j \right)$$

for $j \in \mathbb{N}$. Then $0 = I_0 \subseteq I_1 \subseteq \ldots \subseteq I_n = I_{n+1} = \ldots$ is an ascending chain of ideals which stabilizes at position n.

Proof. It is immediate from the definition that $I_j \subseteq I_{j+1}$ for all $j \in \mathbb{N}$, so it suffices to show that $I_m \subseteq I_n$ for m > n by induction. This follows from the proof of Lemma 2: since the identity in Lemma 2 only involves coefficients of ± 1 , it continues to hold in the polynomial ring $\mathbb{Z}[x_{i\beta\gamma}]$. Applying the universal map $\mathbb{Z} \to R$ shows that the identity holds also in $R[x_{i\beta\gamma}]$, and specializing to R gives the result. \Box

Theorem 5. The following are equivalent for a ring R:

(i) R is local, i.e. has a unique maximal ideal \mathfrak{m}

(ii) For all (equivalently some) $n \ge 1$ and $A_1, \ldots, A_m \in M_n(R)$ with $A_1 + \ldots + A_m$ invertible, there exists $S \subseteq [m]$ with $|S| \le n$ such that $\sum_{i \in S} A_i$ is invertible.

Proof. (i) \implies (ii): First, note that for any $A \in M_n(R)$, A is invertible iff det A is a unit in R. Form the ideals I_j for $j \in [m]$ as above. By hypothesis, I_m contains a unit, so by Proposition 4 so does I_n . Then one of the generators of I_n is a unit: if not, then each generator would be in \mathfrak{m} , hence I_n would be as well, a contradiction. (Alternate proof: applying $R \twoheadrightarrow R/\mathfrak{m}$ reduces to Theorem 1).

(ii) \implies (i): Suppose R has two distinct maximal ideals $\mathfrak{m}_1, \mathfrak{m}_2$. Then there exists $m_1 \in \mathfrak{m}_1, m_2 \in \mathfrak{m}_2$ with $m_1 + m_2 = 1$. For an n such that (ii) holds, let $A_i := \operatorname{diag}(0, \ldots, m_1, \ldots, 0) \in M_n(R)$ be the diagonal matrix with m_1 in the i^{th} spot and 0 elsewhere, for $i = 1, \ldots, n$, and $A_{n+1} := m_2 \cdot \operatorname{id}_n$. Then $A_1 + \ldots + A_{n+1} = \operatorname{id}_n$, but any sum of $\leq n$ of the A_i 's has determinant either 0, m_1^n , or m_2^j for some $1 \leq j \leq n$, hence is not invertible.

Example. (1): Proposition 4 implies that if $I_n \subseteq J$ for some *R*-ideal *J*, then so is I_m for all *m*; e.g. if $A_1, A_2, A_3, A_4 \in M_3(\mathbb{Z})$, and any sum of at most 3 has determinant divisible by 10, then $\det(A_1 + A_2 + A_3 + A_4)$ is also divisible by 10. In particular, taking $J = I_n$ generalizes Theorem 5: for any ring *R* and $A_1, \ldots, A_m \in M_n(R)$, if $I_n \neq R$ then no (distinct) sum of the A_i 's is invertible.

(2): For generic matrices, the ideals I_j quickly become infeasible to compute. The smallest nontrivial case is n = 2: here $R = k[x_{i\beta\gamma} \mid 1 \le i \le 3, 1 \le \beta, \gamma \le 2]$ is a polynomial ring in 12 variables over a field k, and $A_i = (x_{i\beta\gamma}) \in M_2(R)$, i = 1, 2, 3. Then I_1 is a complete intersection prime ideal of codimension 3, whereas I_2 has two minimal primes of codimension 5, and one embedded prime of codimension 7.

JUSTIN CHEN

Theorem 5 says, in some sense, that the question of when a large sum of matrices is invertible is determined by the 1×1 case (precisely) when the ring is local. Motivated by this, we now shift perspectives and ask: if the ring is semilocal (i.e. has only finitely many maximal ideals), when is a large sum of ring elements (= 1×1 matrices) invertible? The following result is interesting in that it follows from a result for (a specific class of) noncommutative rings, but the proof is not obtained by imposing commutativity verbatim!

Theorem 6. Let R be a ring with n maximal ideals (say mSpec $R = \{\mathfrak{m}_1, \ldots, \mathfrak{m}_n\}$), and let $a_1, \ldots, a_m \in R$ with $a_1 + \ldots + a_m \in R^{\times}$. If char $R/\mathfrak{m}_i = \operatorname{char} R/\mathfrak{m}_j$ for all i, j (e.g. if R contains a field), then there exists $S \subseteq [m]$ with $|S| \leq n$ such that $\sum_{i \in S} a_i \in R^{\times}$.

Proof. For any ring S and $a \in S$, $a \in S^{\times}$ iff $\bar{a} \in (S/\operatorname{Rad}(S))^{\times}$, where $\operatorname{Rad}(S)$ is the Jacobson radical of S, i.e. the intersection of all maximal ideals of S. For R as above, $\operatorname{Rad}(R) = \mathfrak{m}_1 \cap \ldots \cap \mathfrak{m}_n$ is a finite intersection, so by Chinese Remainder $R/\operatorname{Rad}(R) \cong R/\mathfrak{m}_1 \times \ldots \times R/\mathfrak{m}_n$. Thus we may assume R is a direct product of n fields k_1, \ldots, k_n .

Now by assumption char $k_i = \operatorname{char} k_j$ for all i, j, so there exists a large field K such that $k_i \hookrightarrow K$ for all i (e.g. any residue field of $k_1 \otimes_k \ldots \otimes_k k_n$, where k is the (common) prime field; cf. [2], Section V.2.4, Cor. to Prop. 4). There is a ring map $\varphi: R \to M_n(K)$, sending $(r_1, \ldots, r_n) \in k_1 \times \ldots \times k_n \mapsto \operatorname{diag}(r_1, \ldots, r_n) \in M_n(K)$. Then $r \in R$ is a unit iff $\varphi(r)$ is a unit in $M_n(K)$, so applying Theorem 1 to $\varphi(a_1), \ldots, \varphi(a_m) \in M_n(K)$ gives the result.

Remark. If n = 2, then Theorem 6 holds without assuming that $\operatorname{char} R/\mathfrak{m}_1 = \operatorname{char} R/\mathfrak{m}_2$: if $a_1, a_2 \in k_1 \times k_2$ are such that none of $a_1, a_2, 1 - a_1, 1 - a_2$ is a unit, then $a_1, a_2 \in \{(1,0), (0,1)\}$, so either $a_1 + a_2$ or $1 - (a_1 + a_2)$ is a unit.

In general though, the hypothesis of equal characteristics in Theorem 6 is crucial, as the following examples show:

Example. In $\mathbb{F}_2 \times \mathbb{F}_3 \times k$ (where k is any field), the elements

$$a_1 = (0, 1, 1),$$

 $a_2 = a_3 = (1, -1, 0),$
 $a_4 = 1 - (a_1 + a_2 + a_3)$

satisfy $a_1 + a_2 + a_3 + a_4 = 1$, but no subset of $\{a_1, a_2, a_3, a_4\}$ of size ≤ 3 sums to a unit (to mentally verify this, it suffices to check that any nonempty subsum of $\{a_1, a_2, a_3\}$ contains both a coordinate equal to 0 and a coordinate equal to 1).

There are also (many) such examples with all elements a_i distinct: in $\mathbb{F}_2 \times \mathbb{F}_3 \times k_1 \times k_2$ (where k_1, k_2 are any fields), the elements

$$a_1 = (0, 0, 0, 1),$$

$$a_2, a_3, a_4 = (1, -1, 0, *),$$

$$a_5 = 1 - (a_1 + a_2 + a_3 + a_4) = (0, 1, 1, \cdot)$$

also sum to a unit, although no proper subset of them does. Here each * can be taken to be any element in k_2 , so if $|k_2| > 2$, then the a_i can be chosen to be pairwise distinct.

4

Finally, we record some additional interesting consequences of Lemma 2. The key feature of the next proposition is nonemptyness of the subset S:

Proposition 7. Let R be a ring, $n \in \mathbb{N}$, and pick any $A_1, \ldots, A_n \in M_n(R)$. Then for any $B \in M_n(R)$ with det $B \neq 0$, there exists $\emptyset \neq S \subseteq [n]$ such that $det\left(\left(\sum_{i \in S} A_i\right) + B\right) - det\left(\sum_{i \in S} A_i\right) \neq 0.$

Proof. This follows from the identity

$$\sum_{\substack{\emptyset \neq S \subseteq [n]}} (-1)^{|S|} \left(\det\left(\sum_{i \in S} A_i\right) - \det\left(\left(\sum_{i \in S} A_i\right) + B\right) \right) = \det B$$

which results from applying Lemma 2 to A_1, \ldots, A_n, B .

We end with a geometric interpretation, in terms of additively generated point configurations on cones over projective hypersurfaces:

Proposition 8. Let k be a field, char k = 0, $n \in \mathbb{N}$, and let $X = V(\det_n) \subseteq \mathbb{A}_k^{n^2}$ be the affine cone over the determinantal hypersurface. Let $\Delta \subseteq \mathbb{A}^{n^2}$ be an n-simplex with first barycentric subdivision $\overline{\Delta}$. If all vertices of $\overline{\Delta}$ other than the centroid lie on X, then in fact the centroid of $\overline{\Delta}$ also lies on X.

Proof. Let p_0, \ldots, p_n be the vertices of Δ . Each vertex of $\overline{\Delta}$ is of the form $\frac{1}{|S|} \left(\sum_{i \in S} p_i \right)$ for some $\emptyset \neq S \subseteq [n]$, which (by homogeneity) lies on X iff $\sum_{i \in S} p_i$ does. Viewing each p_i as an $n \times n$ matrix over k and applying Lemma 2 gives the result. \Box

In fact, the proof of [4] shows that Lemma 2 holds for any homogeneous polynomial, so Proposition 8 actually holds for any cone X (in any affine space) cut out by a degree n polynomial (and thus also for any intersection of such cones).

Acknowledgements: The author would like to thank Joe Kileel for valuable discussions, in particular for contributing to the proof of Lemma 3; as well as David Eisenbud for helpful comments. The examples following Theorems 5 and 6 were carried out with the help of Macaulay2.

References

- S. A. Amitsur. On the Characteristic Polynomial of a Sum of Matrices. Linear and Multilinear Algebra 8(3), p. 177–182, 1980.
- [2] N. Bourbaki. Éléments de mathématique. Algèbre, chapitres 4 à 7. Masson, Paris, 1981.
- [3] R. S. Costas-Santos. On the elementary symmetric functions of a sum of matrices. Journal of Algebra, Number Theory: Advances and Applications 1(2), p. 99–112, 2009.
- [4] A. Facchini, F. Barioli, R. Chapman, and R. M. Carroll. Nonsingular Sums of Matrices: 10784. American Mathematical Monthly 109(7), p. 664–666, 2002.
- [5] D. R. Grayson and M. E. Stillman. Macaulay2, a software system for research in algebraic geometry. Available at http://www.math.uiuc.edu/Macaulay2/.
- [6] C. Reutenauer and M. Schützenberger. A Formula for the Determinant of a Sum of Matrices. Letters in Mathematical Physics 13, p. 299–302, 1987.

Department of Mathematics, University of California, Berkeley, California, 94720 $\rm U.S.A$

E-mail address: jchen@math.berkeley.edu