

Generally Covariant Maxwell Theory for Media with a Local Response: Progress since 2000

Friedrich W. Hehl*, Yakov Itin†, and Yuri N. Obukhov‡

* Univ. of Cologne, Germany, and Univ. of Missouri-Columbia, USA; e-mail: hehl@thp.uni-koeln.de

† Jerusalem College of Technology and Hebrew Univ. Jerusalem, Israel; email: itin@math.huji.ac.il

‡ Nuclear Safety Institute, Russian Academy of Sciences, Moscow, Russia; email: obukhov@ibrae.ac.ru

Abstract—In the recent decades, it became more and more popular for engineers, physicists, and mathematicians alike to put the Maxwell equations into a generally covariant form. This is particularly useful for understanding the fundamental structure of electrodynamics (conservation of electric charge and magnetic flux). Moreover, it is ideally suited for applying it to media with local (and mainly linear) response behavior. We try to collect the new knowledge that grew out of this development. We would like to ask the participants of EMTS 2016 to inform us of work that we may have overlooked in our review.

28 March 2016, file *EspooHehlLocLinMax_10.tex*

Premetric classical electrodynamics has been consistently formulated in 2003 by Hehl and Obukhov [18]; earlier references can be found in that book. We may add O’Dell [65] and Serdyukov et al. [76], which we had overlooked at the time. The premetric framework consists of the generally covariant Maxwell equations in the 4-dimensional or the (1+3)-decomposed version, see Figs. 1 and 2:

$$dH = J \begin{cases} \underline{d}\mathcal{D} = \rho, \\ \underline{D} = \underline{d}\mathcal{H} - j; \end{cases} \quad (1)$$

$$dF = 0 \begin{cases} \underline{d}B = 0, \\ \underline{B} = -\underline{d}E. \end{cases} \quad (2)$$

If the medium has a local and linear response behavior, the excitation $H = (\mathcal{H}, \mathcal{D})$ is local and linear in the field strength $F = (E, B)$. In components we have

$$\check{H}^{ij} = \frac{1}{2}\chi^{ijkl} F_{kl} \quad \text{with} \quad \chi^{ijkl} = -\chi^{jikl} = -\chi^{ijlk}. \quad (3)$$

Here $i, j, \dots = 0, 1, 2, 3$ and $\check{H}^{ij} := (1/2)\epsilon^{ijkl} H_{kl}$. The response tensor density χ has 36 independent components. It can be decomposed into three irreducible pieces with $36 = 20 + 15 + 1$ components, respectively. Split in (1+3) dimensions, with $a, b, \dots = 1, 2, 3$, we have (for details, see [18], [24])

$$D^a = (\epsilon^{ab} - \epsilon^{abc} n_c) E_b + (\gamma^a_b + s_b^a - \delta_b^a s_c^c) B^b + \alpha B^a,$$

$$H_a = (\mu_{ab}^{-1} - \epsilon_{abc} m^c) B^b + (-\gamma^b_a + s_a^b - \delta_a^b s_c^c) E_b - \alpha E_a.$$

Up to here, the metric of spacetime, that is, the gravitational potential, did not enter anywhere. We have a premetric




Fig. 1. The electromagnetic field: The interrelation between the excitation $H = (\mathcal{D}, \mathcal{H})$ and the field strength $F = (E, B)$.




Fig. 2. The Faraday-Schouten pictograms of the electromagnetic field in 3-dimensional space. The 1-forms are depicted by 2 parallel planes, the 2-forms by flux tubes.

framework. The different irreducible parts split as follows (see [45]):

20 components of principal part: $6 \epsilon^{ab}$, $6 \mu_{ab}^{-1}$, $8 \gamma^a_b$;

15 components of skewon part: $9 s_a^b$, $3 n_c$, $3 m^c$;

1 component of axion part: α .

Progress has been achieved since then by different groups, see the more recent books of Delphenich [11], Favaro [13] (Ph.D. thesis), Lindell [46], [47], Raab & de Lange [69], Russer [72], and Scheck [75], see also Cabral & Lobo [6]. In particular Scheck underlines that polarizability and magnetization (and thus \mathcal{D} and \mathcal{H}) are microscopically defined, “i.e. for a single atom or even an elementary particle.”

Premetric electrodynamics turned out to be a lively and active subject. Mainly the following aspects were developed (we order them roughly chronologically):

- 1) *Magnetic charge is alien to the Maxwellian structure:* A magnetic charge can be mathematically included in premetric electrodynamics in a consistent way, see Edelen [12], Kaiser [41], and Hehl & Obukhov [20].

As shown in [18], the structure of the Maxwell equations can be understood as arising from (i) electric charge conservation $dJ = 0$ and (ii) magnetic flux conservation $dF = 0$. This distinction between the electric and magnetic counterparts of the theory can be viewed as a reflection of the topology properties of the underlying manifold: The electric current J and the corresponding excitation H are twisted forms, whereas the field strength F is untwisted. If we modified the electromagnetic structure, following Dirac, by introducing a magnetic current K , with $dF = K$, the Maxwell equations would be “symmetrized,” but the distinction between the two basic fields H and F would be destroyed.

- 2) *Skewon part of the electromagnetic response tensor:* The general concept of the skewon part of the response tensor was introduced in [17], for its relation to dissipation, see [18]. The effects of the skewon on light propagation were studied by Itin, Ni, Obukhov, et al. [62], [36], [40], [53], [58], [38], [37], [56]. A theory of the skewon in interaction with the gravitational field was provided in [23].
- 3) *Quantum Hall effect is independent of the external gravitational field:* The QHE can be phenomenologically described in a $(1+2)$ -dimensional premetric electrodynamic framework; also the local, linear response tensor turns out to be premetric. Therefore, the QHE cannot depend on an external gravitational field, see [19].
- 4) *Electromagnetism can couple to a possible Cartan torsion of spacetime only nonminimally:* Solanki, Preuss, et al. [68], [77] pointed out that a non-minimal coupling of gravity to electromagnetism, in particular to the torsion field, is possible, see also Hehl and Obukhov [16], Rubilar et al. [71], and Itin et al. [25], [26].
- 5) *Signature of the metric can be derived from electrodynamics:* In [18] it has been pointed out that the signature of the metric can be derived from the Lenz rule and the positivity of the energy of the electromagnetic field. The formalism has been straightened out, improved, and enriched by Itin et al. [27], [33].
- 6) *Four incarnations of the Tellegen gyrotor:* The Tellegen gyrotor [43], [72], [78], [79] is an element in linear and




Fig. 3. The pseudoscalar or axion piece α of the response tensor χ^{ijkl} of Chromium Sesquioxide Cr_2O_3 ; in units of $Y_0 = 1/Z_0$ as a function of the temperature T in kelvin. Here Z_0 is the vacuum impedance which, in SI, is $\approx 377 \text{ ohm}$.

passive network theory, which “rotates” currents I into voltages V . Similarly, (i) the axion part α of the electromagnetic response tensor rotates excitations $(\mathcal{H}, \mathcal{D})$ into field strengths (B, E) [18, Eqs.(D.1.112/3)], and so does (ii) the PEMC (perfect electromagnetic conductor) of Lindell & Sihvola [51], [52], see [47, Eq.(7.72)]. (iii), in the QHE, the current/charge (j, ρ) is rotated into the field strength (E, B) , see [18, Eq.(B.4.64)].¹ (iv) The hypothetical axion particle of elementary particle physics, see Wilczek [82] and Weinberg [81, pp. 458–461] has the same behavior as (i) and (ii). In the gyrotor and additionally, in its four incarnations, the P-symmetry (parity) and the T-symmetry (time) are always violated. It has been experimentally confirmed [24], see Fig.3, that in single crystals of Chromium Sesquioxide Cr_2O_3 at low temperatures the axion part is non-vanishing and maximally of the order of $10^{-3} \times$ vacuum admittance. This falsifies the so-called Post constraint.

- 7) *Axion and light propagation:* Ni [54], [55] proposed, as a device for violation of the equivalence principle, to modify classical electrodynamics by introducing an axion field. Carroll et al. [7] used an axion field with a time-like gradient for violation of Lorentz and parity invariance. This model was extended and reformulated in the premetric formalism by Itin [28], [32], [31]. Various theoretical issues of this model were studied by Balakin [4], and Noble [60]. Recently Ni [59], [57] provided constraints on axion and dilaton from polarized/unpolarized laboratory/astrophysical/cosmic experiments/observations.
- 8) *Forbid birefringence in vacuo and find the light cone:*

¹In spite of the last mentioned fact, one cannot build a gyrotor out of the QHE, as pointed out to us by von Klitzing [42].

Wave propagation in electrodynamics with a local and linear response law exhibits the phenomenon of birefringence. In the geometric optics approximation, this is manifest in the fourth order Fresnel equation for the wave covector. Such a picture is typical for complex (anisotropic, magnetoelectric, moving) media. However, it is natural to assume that the physical vacuum is a non-birefringent continuum. This no-birefringence condition reduces the general Fresnel equation to the light cone [21], [44]; thus, the spacetime metric is recovered from the electromagnetic response tensor. More recent developments can be found in [8], [14], [29].

- 9) *Dimensional analysis of physical quantities*: In [18] we analyzed electrodynamics in terms of a consistent theory of physical dimension. In particular the operational verification of the physical quantities were at center stage. In [22], we extended these considerations in the context of a discussion with Okun. The vacuum admittance $\sqrt{\varepsilon_0/\mu_0} = e^2/(2h\alpha)$ can be expressed in terms of the fine structure constant α , the elementary charge e and Planck's constant h . The possible time variability of the vacuum impedance and the fine structure constant were discussed by Tobar [80] and us.
- 10) *New initial value formulation of Maxwell's theory by Perlick*: It was studied by Perlick [66] for the premetric version. He derived several conditions for the evolution equations to be hyperbolic, strongly hyperbolic, or symmetric hyperbolic. In particular, he characterized all response laws for which the evolution equations are symmetric hyperbolic. The latter property is sufficient, but not necessary, for well-posedness of the initial-value problem. Symbol and *hyperbolic polynomials* of the wave equation and their relation to premetric electrodynamics were discussed by Beig [5]. Itin [35] derived premetric covariant jump conditions for the initial value hypersurface, the light cone, and a boundary between two media.

- 11) *The Kummer tensor of electrodynamics, Fresnel wave surfaces unmasked as Kummer surfaces*: We found a fourth rank Kummer tensor density [3] that is cubic in the response tensor χ ,

$$\mathcal{K}^{ijkl} := \chi^{ajib} \diamond \chi_{abcd}^\diamond \chi^{ckdl}, \quad (4)$$

where \diamond denotes a premetric duality operator. The Fresnel surfaces are defined by $\mathcal{K}^{ijkl} q_i q_j q_k q_l = 0$; here q_i is the wave covector of the propagating light. Following earlier ideas of Bateman, Favaro et al. [15] were able to show that, for vanishing skewon part, the Fresnel surfaces turn out to be Kummer surfaces with up to 16 singular points. By a suitable choice of the electromagnetic moduli of a hypothetical metamaterial according to [15],

$$\begin{aligned} \varepsilon^{ab} &= \frac{1}{4} \varepsilon_0 \text{diag}(-1 - \sqrt{3}, -1 - \sqrt{3}, -4 + 2\sqrt{3}), \\ \mu_{ab}^{-1} &= \frac{1}{4} \mu_0^{-1} \text{diag}(1 + \sqrt{3}, 1 + \sqrt{3}, 4 - 2\sqrt{3}), \end{aligned} \quad (5)$$




Fig. 4. The Fresnel surface for a metamaterial with the moduli as displayed in equation (5). This surface turns out to be a Kummer surface of algebraic geometry.

$$\gamma^a_b = \frac{1}{4} \sqrt{\varepsilon_0 \mu_0^{-1}} \text{diag}(3 + \sqrt{3}, -3 - \sqrt{3}, 0),$$

it was possible to create such a surface, see Fig. 4.

Russer pointed out to us [73] that the 16-fold symmetry of the Fresnel-Kummer surface reminds him of a 16-fold solution manifold in the transmission-line matrix (TLM) method, see [72, Sec.14.3] and [74]. Since both phenomena refer to the propagation of electromagnetic signals, there may very well exist an interrelationship.

- 12) *Possible factorizations of the Fresnel equation (FE)*: Which media give rise to such simplifications? Four cases were solved: (i) The FE cannot be factorized in polynomials of lower degree. Materials with anisotropic permittivity and permeability, e.g., display this behavior [34]. (ii) The FE factorizes in irreducible quadratic polynomials. For research on the situation where the two factors are distinct and have Lorentzian signature, see [9], [48], [63]. Light then propagates along two distinct light cones. When the two quadratic factors are the same, up to a constant, a single light cone is obtained, so that there is no birefringence, see item 8). (iii) The FE factorizes in four linear polynomials. Extreme Magneto-Electric (EME) materials [50] are known to have this property. (iv) The FE vanishes trivially, i.e., it is satisfied by any wave covector $q_i = (\omega, -k_a)$. Media

corresponding to this scenario are described in [49].

- 13) *Photon propagator in premetric electrodynamics*: The Green tensor is the photon propagator in momentum space. It is important in quantum field theoretical extension of electrodynamics. It was derived and applied to axion and skewon modified electrodynamics by Itin [30], [32], [40]. In a similar way, Andrianov et al. [1] studied the propagation of photons and massive vector mesons in the presence of a medium, which violates Lorentz and a CPT invariance. Pfeifer and Siemssen [67] applied the premetric photon propagator for studying the causal structure and the quantization problems in electrodynamic models with linear response.
- 14) *Accelerated reference frames in coordinate free form*: Auchmann and Kurz [2], by using a fiber bundle formalism, developed a general theory for observers by splitting 4-dimensional electrodynamics in a covariant way into time and space. This is particularly useful for interpreting experiments in accelerated motion.
- 15) *Energy-momentum tensor of the electromagnetic field* [61], [70]: The form of the energy-momentum tensor of the electromagnetic field in a medium with non-trivial electric and magnetic properties is a problem that dates back to the century-old work of Minkowski and Abraham. The Minkowski vs. Abraham controversy can be resolved by the crucial observation that the electromagnetic field in a medium is an open physical system. The closed system of interacting electromagnetism plus matter is characterized by the total energy-momentum tensor that is defined unambiguously within the Lagrange-Noether framework. Both Minkowski and Abraham tensors may arise under appropriate conditions for special splits of the total energy-momentum tensor.
- 16) *Nonlinear electrodynamics*: So far, we only treated linear response laws. A nonlinear law naturally arises when the quantized electromagnetic field excites radiative vacuum corrections. Alternatively, one can consider a fundamentally nonlinear electrodynamics with a Born-Infeld type Lagrangian, see [18]. Wave propagation in nonlinear models manifest itself typically in the birefringence property. An analysis of geometric optics in nonlinear electrodynamics reveals the product structure of the Fresnel wave surface, which gives rise to two optical metrics, see [64]. It is also possible to formulate a Finsler electrodynamics in a premetric way, see [10] and [39].

ACKNOWLEDGMENTS

We are very grateful to Alberto Favaro (London) for many useful discussions and detailed remarks. We'd also like to thank Gerry Kaiser (Portland, OR) for a discussion on the possible existence of magnetic charge.

REFERENCES

[1] A. A. Andrianov, S. S. Kolevatov and R. Soldati, *Propagation of photons and massive vector mesons between a parity breaking medium and vacuum*, JHEP **1111**, 007 (2011) [arXiv:1109.3440]. 4

[2] B. Auchmann and S. Kurz, *Observers and splitting structures in relativistic electrodynamics*, J. Phys. A **47**, 435202 (2014) [arXiv:1411.0837]. 4

[3] P. Baekler, A. Favaro, Y. Itin, and F. W. Hehl, *The Kummer tensor density in electrodynamics and in gravity*, Annals Phys. (NY) **349**, 297–324 (2014) [arXiv:1403.3467]. 3

[4] A. B. Balakin, V. V. Bochkarev and N. O. Tarasova, *Gradient models of the axion-photon coupling*, Eur. Phys. J. C **72**, 1895 (2012) [arXiv:1201.3009]. 2

[5] R. Beig, *Concepts of hyperbolicity and relativistic continuum mechanics*, Lect. Notes Phys. **692**, 101–116 (2006) [arXiv:gr-qc/0411092]. 3

[6] F. Cabral and F. S. N. Lobo, *Electrodynamics and spacetime geometry I: Foundations*, arXiv:1602.01492 (2016). 2

[7] S.M. Carroll, G.B. Field, and R. Jackiw, *Limits on a Lorentz- and parity-violating modification of electrodynamics*, Phys. Rev. **D41**, 1231–1240 (1990). 2

[8] M. F. Dahl, *Determination of an electromagnetic medium from the Fresnel surface*, J. Phys. A **45**, 405203 (2012). 3

[9] M. F. Dahl, *Characterisation and representation of non-dissipative electromagnetic medium with two Lorentz null cones*, J. Math. Phys. **54**, 011501 (2013). 3

[10] D. H. Delphenich, *Nonlinear electrodynamics and QED*, arXiv:hep-th/0309108 (2003). 4

[11] D. H. Delphenich, *Pre-metric electromagnetism*, electronic book, 410 pages, <http://www.neo-classical-physics.info/> (2009). See also the articles in Ann. Phys. (Berlin) **14**, 347–377 (2005) and **16**, 207–217 (2007). 2

[12] D. G. B. Edelen, *A metric free electrodynamics with electric and magnetic charges*, Ann. Phys. (NY) **112**, 366–400 (1978). 2

[13] A. Favaro, *Recent Advances in Classical Electromagnetic Theory*. Ph.D. thesis, Imperial College London (2012). 2

[14] A. Favaro and L. Bergamin, *The non-birefringent limit of all linear, skewonless media and its unique light-cone structure*, Ann. Phys. (Berlin) **523**, 383–401 (2011). 3

[15] A. Favaro and F. W. Hehl, *Light propagation in local and linear media: Fresnel-Kummer wave surfaces with 16 singular points*, Phys. Rev. A **93**, 013844 (2016) [arXiv: 1510.05566]. 3

[16] F. W. Hehl and Yu. N. Obukhov, *How does the electromagnetic field couple to gravity, in particular to metric, nonmetricity, torsion, and curvature?* In: *Gyros, Clocks, Interferometers...: Testing Relativistic Gravity in Space*. C. Lämmerzahl et al., eds. Lecture Notes in Physics Vol. **562** (Springer: Berlin, 2001) pp. 479–504 [arXiv:gr-qc/0001010]. 2

[17] F. W. Hehl, Yu. N. Obukhov, and G. F. Rubilar, *On a possible new type of a T odd skewon field linked to electromagnetism*. In: *Developments in Mathematical and Experimental Physics*, A. Macias, F. Uribe, and E. Diaz, eds. Volume A: *Cosmology and Gravitation* (Kluwer Academic/Plenum Publishers: New York, 2002) pp.241–256 [arXiv:gr-qc/0203096]. 2

[18] F. W. Hehl and Yu. N. Obukhov, *Foundations of Classical Electrodynamics: Charge, flux, and metric*. Birkhäuser, Boston, MA (2003). 1, 2, 3, 4

[19] F. W. Hehl, Yu. N. Obukhov and B. Rosenow, *Is the Quantum Hall Effect influenced by the gravitational field?* Phys. Rev. Lett. **93**, 096804 (2004) [arXiv:cond-mat/0310281]. 2

[20] F. W. Hehl and Yu. N. Obukhov, *Electric/magnetic reciprocity in premetric electrodynamics with and without magnetic charge, and the complex electromagnetic field*, Phys. Lett. A **323**, 169–175 (2004) [arXiv:physics/0401083]. 2

[21] F. W. Hehl and Yu. N. Obukhov, *To consider the electromagnetic field as fundamental, and the metric only as a subsidiary field*, Found. Phys. **35**, 2007–2025 (2004) [arXiv:physics/0404101]. 3

[22] F. W. Hehl and Yu. N. Obukhov, *Dimensions and units in electrodynamics*, Gen. Relat. Grav. **37**, 733–749 (2005) [arXiv:physics/0407022]. 3

[23] F. W. Hehl, Y. N. Obukhov, G. F. Rubilar and M. Blagojevic, *On the theory of the skewon field: From electrodynamics to gravity*, Phys. Lett. A **347**, 14–24 (2005) [arXiv:gr-qc/0506042]. 2

[24] F. W. Hehl, Y. N. Obukhov, J. P. Rivera and H. Schmid, *Relativistic nature of a magnetoelectric modulus of Cr₂O₃ crystals: A four-dimensional pseudoscalar and its measurement*, Phys. Rev. A **77**, 022106 (2008) [arXiv:0707.4407]. 1, 2

[25] Y. Itin, *Noether currents and charges for Maxwell-like Lagrangians*, J. Phys. A **36**, 8867–8884 (2003) [arXiv:math-ph/0307003]. 2

[26] Y. Itin and F.W. Hehl, *Maxwell's field coupled nonminimally to quadratic torsion: Axion and birefringence*. Phys. Rev. D **68**, 127701 (2003) [arXiv:gr-qc/0307063]. 2

- [27] Y. Itin and F.W. Hehl, *Is the Lorentz signature of the metric of spacetime electromagnetic in origin?* Ann. Phys. (NY) **312**, 60–83 (2004) [arXiv:gr-qc/0401016]. **2**
- [28] Y. Itin, *Carroll-Field-Jackiw electrodynamics in the pre-metric framework*, Phys. Rev. D **70**, 025012 (2004) [arXiv:hep-th/0403023]. **2**
- [29] Y. Itin, *Nonbirefringence conditions for spacetime*, Phys. Rev. D **72**, 087502 (2005). **3**
- [30] Y. Itin, *A Generalized photon propagator*, J. Phys. A **40**, F737–F744 (2007) [arXiv:0705.0756]. **4**
- [31] Y. Itin, *Wave propagation in axion electrodynamics*, Gen. Rel. Grav. **40**, 1219–1238 (2008) [arXiv:0706.2991]. **2**
- [32] Y. Itin, *Photon propagator for axion electrodynamics*, Phys. Rev. D **76**, 087505 (2007) [arXiv:0709.1637]. **2, 4**
- [33] Y. Itin and Y. Friedman, *Backwards on Minkowski’s road. From 4D to 3D Maxwellian electromagnetism*, Ann. Phys. (Berlin) **17**, 769–786 (2008) [arXiv:0807.2625]. **2**
- [34] Y. Itin, *Dispersion relation for electromagnetic waves in anisotropic media*, Phys. Lett. A **374**, 1113–1116 (2010). **3**
- [35] Y. Itin, *Covariant jump conditions in electromagnetism*, Annals Phys. (NY) **327**, 359–375 (2012) [arXiv:1403.7700]. **3**
- [36] Y. Itin, *Skewon no-go theorem*, Phys. Rev. D **88**, 107502 (2013) [arXiv:1310.8314]. **2**
- [37] Y. Itin, *Electromagnetic media with Higgs-type spontaneously broken transparency*, Phys. Lett. A **379**, 619–622 (2014) [arXiv:1406.3442]. **2**
- [38] Y. Itin, *Skewon modification of the light cone structure*, Phys. Rev. D **91**, 085002 (2015) [arXiv:1407.6722]. **2**
- [39] Y. Itin, C. Lämmerzahl and V. Perlick, *Finsler-type modification of the Coulomb law*, Phys. Rev. D **90**, 124057 (2014) [arXiv:1411.2670]. **4**
- [40] Y. Itin, *Photon propagator in skewon electrodynamics*, Phys. Rev. D **93**, 025023 (2016) [arXiv:1511.01840]. **2, 4**
- [41] G. Kaiser, *Energy-momentum conservation in pre-metric electrodynamics with magnetic charges*, J. Phys. A **37**, 7163–7168 (2004) [arXiv:math-ph/0401028]. **2**
- [42] K. von Klitzing, private communication to fwh (November 2004). **2**
- [43] K. Küpfmüller, W. Mathis, A. Reibiger, *Theoretische Elektrotechnik*, 19th ed., Springer Vieweg, Berlin (2013). **2**
- [44] C. Lämmerzahl and F. W. Hehl, *Riemannian light cone from vanishing birefringence in premetric vacuum electrodynamics*, Phys. Rev. **D70**, 105022 (2004) [arXiv:gr-qc/0409072]. **3**
- [45] O. L. de Lange and R. E. Raab, *Multipole theory and the Hehl-Obukhov decomposition of the electromagnetic constitutive tensor*, J. Math. Phys. **56**, 053502 (2015). **1**
- [46] I. V. Lindell, *Differential Forms in Electromagnetics*, IEEE Press: Piscataway, NJ, and Wiley-Interscience (2004). **2**
- [47] I. V. Lindell, *Multiforms, Dyadics, and Electromagnetic Media*, Wiley, Hoboken, NJ [and IEEE Press] (2015). **2**
- [48] I. V. Lindell, L. Bergamin, and A. Favaro, *Decomposable medium conditions in four-dimensional representation*, IEEE Transactions on Antennas and Propagation **60**, 367–376 (2012). **3**
- [49] I. V. Lindell and A. Favaro, *Electromagnetic media with no dispersion equation*, Progress In Electromagnetics Research B **51**, 269–289 (2013). **4**
- [50] I. V. Lindell, A. H. Sihvola, and A. Favaro, *Plane wave propagation in extreme magnetoelectric (EME) medium*, Talk abstract, submitted to EMTS 2016 (Feb. 2016). **3**
- [51] I.V. Lindell and A.H. Sihvola, *Perfect electromagnetic conductor*, J. Electromagn. Waves Appl. **19**, 861–869 (2005). **2**
- [52] I.V. Lindell and A.H. Sihvola, *Transformation method for problems involving perfect electromagnetic conductor (PEMC) structures*, IEEE Trans. Antennas Propag. **53**, 3005–3011 (2005). **2**
- [53] H. H. Mei, W. T. Ni, W. P. Pan, L. Xu and S. d. S. Alighieri, *New constraints on cosmic polarization rotation from the ACTPol cosmic microwave background B-Mode polarization observation and the BICEP2 constraint update*, Astrophys. J. **805**, 107 (2015). **2**
- [54] W.-T. Ni, *A non-metric theory of gravity*. Dept. Physics, Montana State University, Bozeman. Preprint December 1973. The paper is available via <http://gravity5.phys.nthu.edu.tw/webpage/article4/index.html> **2**
- [55] W.-T. Ni, *Equivalence principles and electromagnetism*, Phys. Rev. Lett. **38**, 301–304 (1977). **2**
- [56] W.-T. Ni, *Skewon field and cosmic wave propagation*, Phys. Lett. A **378**, 1217–1223 (2014) [arXiv:1312.3056]. **2**
- [57] W.-T. Ni, *Dilaton field and cosmic wave propagation*, Phys. Lett. A **378**, 3413–3418 (2014) [arXiv:1410.0126]. **2**
- [58] W.-T. Ni, *Spacetime structure and asymmetric metric from the premetric formulation of electromagnetism*, Phys. Lett. A **379**, 1297–1303 (2015) [arXiv:1411.0460]. **2**
- [59] W.-T. Ni, *Searches for the role of spin and polarization in gravity: a five-year update*, Int. J. Mod. Phys. Conf. Ser. **40**, 1660010 (2016) [arXiv:1501.07696]. **2**
- [60] A. Noble, *Axion electrodynamics in a waveguide*, Phys. Lett. A **372**, 2346–2349 (2008). **2**
- [61] Y. N. Obukhov, *Momentum of light in complex media*, Talk abstract, submitted to EMTS 2016 (Feb. 2016). **4**
- [62] Y. N. Obukhov and F.W. Hehl, *Possible skewon effects on light propagation*, Phys. Rev. D **70**, 125015 (2004) [arXiv:physics/0409155]. **2**
- [63] Y. N. Obukhov, T. Ramos, and G. Rubilar, *Relativistic Lagrangian model of a nematic liquid crystal interacting with an electromagnetic field*, Phys. Rev. E **86**, 031703 (2012). **3**
- [64] Y. N. Obukhov, and G. F. Rubilar, *Fresnel analysis of the wave propagation in nonlinear electrodynamics*, Phys. Rev. D **66**, 024042 (2002) [arXiv:gr-qc/0204028]. **4**
- [65] T. H. O’Dell, *The Electrodynamics of Magneto-Electric Media*, North-Holland, Amsterdam (1970). **1**
- [66] V. Perlick, *On the hyperbolicity of Maxwell’s equations with a local constitutive law*, J. Math. Phys. **52**, 042903 (2011) [arXiv:1011.2536]. **3**
- [67] C. Pfeifer and D. Siemssen, *The potential in general linear electrodynamics: Causal structure, propagators and quantization*, arXiv:1602.00946 (2016). **4**
- [68] O. Preuss, M. P. Haugan, S. K. Solanki and S. Jordan, *An astronomical search for evidence of new physics: Limits on gravity-induced birefringence from the magnetic white dwarf RE J0317-853*, Phys. Rev. D **70**, 067101 (2004) [arXiv:gr-qc/0405068]. **2**
- [69] R. E. Raab and O. L. de Lange, *Multipole Theory in Electromagnetism. Classical, quantum, and symmetry aspects, with applications*, Clarendon Press, Oxford, UK (2005). **2**
- [70] T. Ramos, G. F. Rubilar and Yu. N. Obukhov, *First principles approach to the Abraham-Minkowski controversy for the momentum of light in general linear non-dispersive media*, J. Opt. **17**, 025611 (2015) [arXiv:1310.0518]. **4**
- [71] G. F. Rubilar, Yu.N. Obukhov and F.W. Hehl, *Torsion nonminimally coupled to the electromagnetic field and birefringence*. Class. Quant. Grav. **20**, L185–L191 (2003) [arXiv:gr-qc/0305049]. **2**
- [72] P. Russer, *Electromagnetics, Microwave Circuit and Antenna Design for Communications Engineering*, 2nd ed., Artech House, Norwood, MA (2006). **2, 3**
- [73] P. Russer, private communication to fwh (January 2016). **3**
- [74] P. Russer and J. A. Russer, *Some remarks on the transmission line matrix (TLM) method and its application to transient EM fields and to EMC problems*. In: Computational Electromagnetics—Retrospective and Outlook, In honor of W. J. R. Hoeyer, I. Achmed and Z. Chen (eds.) (2015) pp. 29–56. **3**
- [75] F. Scheck, *Classical Field Theory, on Electrodynamics, Non-Abelian Gauge Theories and Gravitation*, Springer, Heidelberg (2012). **2**
- [76] A. Serdyukov, I. Semchenko, S. Tretyakov, and A. Sihvola, *Electromagnetics of Bi-anisotropic Materials, Theory and Applications*, Gordon and Breach, Amsterdam (2001). **1**
- [77] S.K. Solanki, O. Preuss, M.P. Haugan, et al., *Solar constraints on new couplings between electromagnetism and gravity*, Phys. Rev. D **69**, 062001 (2004). **2**
- [78] B. D. H. Tellegen, *The gyrator, a new electric network element*, Philips Res. Rep. **3**, 81–101 (1948). **2**
- [79] B. D. H. Tellegen, *The gyrator, an electric network element*, Philips Technical Review **18**, 120–124 (1956/57). Reprinted in H. B. G. Casimir and S. Gradstein (eds.) *An Anthology of Philips Research*. Philips’ Gloeilampenfabrieken, Eindhoven (1966) pp. 186–190. **2**
- [80] M. E. Tobar, *Global representation of the fine structure constant and its variation*, Metrologia **42**, 129–133 (2005) [arXiv:hep-ph/0306230v5] (original version June 2003). **3**
- [81] S. Weinberg, *The Quantum Theory of Fields. Volume II: Modern Applications*, Cambridge Univ. Press, Cambridge, UK (1996). **2**
- [82] F. Wilczek, *Two applications of axion electrodynamics*, Phys. Rev. Lett. **58**, 1799–1802 (1987). **2**