# Direct evaluation of overlap integrals between Slater-type-orbitals

Michael J. Caola<sup>1, a)</sup>

6, Normanton Rd, Bristol BS8 2TY, U.K.

(Dated: 21 April 2016)

We derive direct single-stage numerical evaluation of the electronic overlap integral between arbitrary atomic orbitals (including STOs). Integration is over cartesian co-ordinates, and replaces previous sums over 'special' functions. The results, in Mathematica 10 and Maple 18, agree with the literature to  $\sim 8$  digits. We briefly discuss possible use in quantum chemistry, including accuracy, algorithmic suitability and operating-system machine-implementation as an intrinsic function.

## I. INTRODUCTION

Single-centre electronic atomic orbitals,

$$\psi_{nlm}(\mathbf{r}) = F_{nl}(r,\alpha) \,\mathcal{Y}_{ml}(\mathbf{r}),\tag{1}$$

are building blocks in the quantum description of atoms, molecules, crystals and, hence in general of matter. In Eq(1)  $F_{nl}(r)$  is a radial function,  $\mathcal{Y}_{ml} = r^l Y_{ml}$  is a *solid* harmonic, and  $Y_{ml}$  the familiar spherical (surface) harmonic. We state that vital parts of Molecular Quantum Mechanics can be built with the overlap integral

$$I(\mathbf{R}) = I(\mathbf{R}, n, l, m, n', l', m') = \int d\mathbf{r} \, \psi_{nlm}(\mathbf{r})^* \, \psi_{n'l'm'}(\mathbf{r} - \mathbf{R})$$
(2)

where vector **R** is the spatial separation of the two orbital centres. Important normalised  $F_{nl}(r)$  are the Gaussian-

$$F_{nl}(r) = \dots e^{-\beta r^2} \tag{3}$$

and Slater-type-orbitals (STOs)

$$F_{nl}(r,\alpha) = F_{nl}(x,y,z,\alpha) = \frac{(2\alpha)^{n+1/2}}{\sqrt{(2n)!}} r^{n-1-l} e^{-\alpha r},$$
  
$$r = r(x,y,z) = \sqrt{x^2 + y^2 + z^2}$$
  
(4)

where  $\alpha$  is a screening constant. The STO is accepted as physically superior to the GTO, but numerical evaluation of its  $I(\mathbf{R})$  is more difficult; both have previously used special and associated functions, including: Fourier, Bessel, Laguerre, Gegenbauer, Gaunt, Hobson, ... We shall next evaluate the  $I(\mathbf{R})$  for STOs, as a direct singlestage integration, with no summations over 'special' functions.

## **II. ANALYSIS**

From Eq(1) and Eq(2) we have

$$\psi_{n'l'm'}(\mathbf{r} - \mathbf{R}) = F_{n'l'}(|\mathbf{r} - \mathbf{R}|, \alpha') \mathcal{Y}_{m'l'}(\mathbf{r} - \mathbf{R}), \quad (5)$$

which is valid for arbitrary F(r) and, with cartesian vectors  $\mathbf{r}(x, y, z)$  and  $\mathbf{R}(X, Y, Z)$ , will use

$$|\mathbf{r} - \mathbf{R}| = r' = r'(x, y, z) = \sqrt{(x - X)^2 + (y - Y)^2 + (z - Z)^2}.$$
 (6)

Also, we use the *cartesian* solid-harmonic<sup>1,2</sup> in Eq(5)

$$\mathcal{Y}_{ml}(\mathbf{r}) = \mathcal{Y}_{ml}(x, y, z) = \left[\frac{(2l+1)(l+m)!(l-m)!}{4\pi}\right]^{1/2}$$
$$\sum_{k=0}^{[(l-m)/2]} \frac{(-x-iy)^{k+m}(x-iy)^k z^{l-m-2k}}{2^{2k+m}(k+m)!k!(l-m-2k)!},$$
$$l = 0, 1, 2, \dots; m = -l..+l$$
(7)

Thus with Eqs(5,6,7) in Eq(2) we have

$$I = I(X, Y, Z) =$$

$$\int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy \int_{-\infty}^{\infty} dz \ F_{nl}(x, y, z, \alpha) \ \mathcal{Y}_{ml}(x, y, z)$$

$$F_{n'l'}(x - X, y - Y, z - Z, \alpha') \ \mathcal{Y}_{m'l'}(x - X, y - Y, z - Z)$$
(8)

The I(X, Y, Z) of Eq(8) can be evaluated by direct numerical computation and is valid for *arbitrary* orbitals specified by  $F_{nl}()$ ; this is our desired solution.

For the case of an STO (8) becomes

$$I = I(X, Y, Z) = \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy \int_{-\infty}^{\infty} dz \frac{(2\alpha)^{n+1/2}}{\sqrt{(2n)!}} r^{n-1-l} e^{-\alpha r} \mathcal{Y}_{ml}(x, y, z) \frac{(2\alpha')^{n'+1/2}}{\sqrt{(2n')!}} r'^{n'-1-l'} e^{-\alpha' r'} \mathcal{Y}_{m'l'}(x - X, y - Y, z - Z),$$
$$= \sqrt{x^2 + y^2 + z^2} \quad r' = \sqrt{(x - X)^2 + (y - Y)^2 + (z - Z)^2}$$
(9)

#### **III. NUMERICAL RESULTS**

r

We use Mathematica 10 and Maple 18 to calculate Eq(9). Each integral in the table below contains a

<sup>&</sup>lt;sup>a)</sup>caolam@blueyonder.co.uk

comma (e.g. -0.117413789,53804531) whose left figures agree with literature values<sup>3-10</sup>: this is typically 8 digits. These data are collected in <sup>9,10</sup>.

## IV. DISCUSSION

Our overlap integrals for STOs Eq(9) agree with the literature to  $\sim 8$  digits We ask active experts (quantum chemists/physicists and computer-aware numerical-analysts) if our direct evaluation Eq(10) could be useful.

Present methods, (sums over special function, SS) e.g.  $^{3-10}$ , to calculate Eq(9) are acceptable, so our proposed direct integration (DI) should consider *inter alia*:

- What minimum accuracy is needed for quantum molecular calculations? If >8 digits, then SS and DI give different values and we must ask
- Which of SS and DI is more accurate (suitable)? It would be wrong to automatically assume that the established SS is more accurate: SS and DI are different methods needing expert comparison. Along with accuracy we would like DI to have suitable and natural notation for its purpose, so we ask
- How would DI handle/evaluate any of the several integrals (of which the overlap is but one) occurring in quantum molecular mechanics? We sketch evaluation of coulomb (ab|cd), 'the two-electron, four centre integral, one of the greatest problems in quantum chemistry'<sup>11</sup>:

$$(ab|cd) = (12|34) = \int \frac{d1 \, d2 \, \psi_a(\mathbf{r1}) \psi_b(\mathbf{r1})^* \psi_c(\mathbf{r2}) \psi_d(\mathbf{r2})^*}{r_{12}}$$
(10)

where

$$\psi_a(r1) = \psi_{n_a l_a m_a}(\alpha_a, x1, y1, z1), \quad d1 = dx1 \, dy1 \, dz1,$$

$$r_{12} = |\mathbf{R} + \mathbf{r2} - \mathbf{r1}| =$$

$$\sqrt{(X+x2-x1)^2+(Y+y2-y1)^2+(Z+z2-z1)^2},$$

etc., and is evaluated in Mathematica 10 in the same way used for overlap Eq(9).

### V. REFERENCES

- <sup>1</sup>Louck J S 1996 Atomic, Molecular, & Optical handbook, ed Drake G W F (New York: AIP Press) p 9
- <sup>2</sup>Caola M J 1978 J. Phys. A: Math. Gen. **11** L23-5
- <sup>3</sup>Barnett M P2003 Int. J. Quantum Chem. **95** 791-805
- <sup>4</sup>Ozdogan T 2004 Collect. Czech Chem. Commun. 69 279-91
- <sup>5</sup>Mamedov B A 2004 Chin. J. Phys. **42** 176-81
- <sup>6</sup>Romanowski Z, Jailbout A F (2009) J. Math. Chem. 45 97-107
- <sup>7</sup>Yavuz M et al. 2005 Commun. Theor. Phys. 43 151
- <sup>8</sup>Rico J F et al. 1988 J. Comput. Chem. **9** 790
- <sup>9</sup>Ozdogan T, Nalcaci A 2012 Int. J. Phys. Sciences 7 5378-90
- <sup>10</sup>Yukcu N 2012 AWERProc. Inf. Tech. Comput. Sci. 2 61-6
- <sup>11</sup>Atkins P, Friedman R 2011 Molecular Quantum Mechanics(Oxford: O. U. P.)

n	l	m	n'	l'	m'	$\alpha$	$\alpha'$	R	Θ	$\Phi$	integral $Eq(9)$	$\mathbf{ref}$
1	0	0	2	1	0	10	2	1.4	0	0	-0.117413789,53804531	3
2	1	0	5	2	0	2	0.3	1.4	0	0	-0.23323008, 22624455-2	3
3	2	0	3	2	0	7.5	2.5	5	$\pi/3$	$2\pi/3$	-0.68034002, 4312253-4	4
3	2	1	3	<b>2</b>	0	9.7	6.4	0.3	$\pi/9$	$3\pi/4$	$0.013735076,\!44$	5
8	0	0	8	0	0	5	1	1	0	0	0.0107437341, 23348333	6
10	7	1	8	1	1	3	3	10	0	0	0.23447835, 22183802-2	7
1	0	0	1	0	0	10	10	1.4	0	0	0.66799473, 05543532-4	8
2	1	0	2	1	0	2	2	1.4	0	0	$-0.10074038,\!66530121$	8