

# A Fast Algorithm for Lattice Hyperonic Potentials

Hidekatsu NEMURA,<sup>1,2</sup> Sinya AOKI,<sup>1,2,3</sup> Takumi DOI,<sup>2</sup> Shinya GONGYO,<sup>2,3</sup> Tetsuo HATSUDA,<sup>2,4</sup> Yoichi IKEDA,<sup>2</sup> Takashi INOUE,<sup>2,5</sup> Takumi IRITANI,<sup>2,6</sup> Noriyoshi ISHII,<sup>2,7</sup> Takaya MIYAMOTO,<sup>2,3</sup> Keiko MURANO,<sup>2,7</sup> and Kenji SASAKI<sup>1,2</sup>

<sup>1</sup> Center for Computational Sciences, University of Tsukuba, Ibaraki, 305-8577, Japan

<sup>2</sup> Theoretical Research Division, Nishina Center, RIKEN, Saitama, 351-0198, Japan

<sup>3</sup> Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto, 606-8502, Japan

<sup>4</sup> Kavli IPMU (WPI), The University of Tokyo, Chiba 277-8583, Japan

<sup>5</sup> Nihon University, College of Bioresource Sciences, Kanagawa 252-0880, Japan

<sup>6</sup> Department of Physics and Astronomy, Stony Brook University, Stony Brook, New York 11794-3800, USA

<sup>7</sup> Research Center for Nuclear Physics (RCNP), Osaka University, Osaka 567-0047, Japan

E-mail: nemura.hidekatsu.gb@u.tsukuba.ac.jp

We describe an efficient algorithm to compute a large number of baryon-baryon interactions from  $NN$  to  $\Xi\Xi$  by means of HAL QCD method, which lays the groundwork for the nearly physical point lattice QCD calculation with volume  $(96a)^4 \approx (8.2\text{fm})^4$ . Preliminary results of  $\Lambda N$  potential calculated with quark masses corresponding to  $(m_\pi, m_K) \approx (146, 525)\text{MeV}$  are presented.

**KEYWORDS:** nuclear force, hyperon-nucleon interaction, lattice QCD, high performance computing

## 1. Introduction

Nuclear force and hyperonic nuclear forces provide a useful starting point to figure out how the hypernuclear systems are bound, in which hyperons (or strange quarks) are embedded in normal nuclear systems as “impurities.” [1] From studies of few-body systems for  $s$ -shell  $\Lambda$  hypernuclei, for example, it has been pointed out that the coupled-channel  $\Lambda N - \Sigma N$  interaction plays a vital role to make a hypernucleus being bounded [2] although phenomenological hyperon-nucleon potentials are not well constrained from experimental data.

In the past several years, a new approach to study the hadronic forces from the lattice QCD has been proposed [3, 4]. In this approach, the interhadron potential is obtained by means of the lattice QCD measurement of the Nambu-Bethe-Salpeter (NBS) wave function. The observables such as the phase shifts and the binding energies are calculated through the resultant potential [5]. This approach has been further extended and applied to various problems. See Ref. [6] and references therein for the state-of-the-art outcomes. In addition, a large scale lattice QCD calculation is now in progress [7] to study the baryon interactions from  $NN$  to  $\Xi\Xi$  by measuring the NBS wave functions for 52 channels from the  $2 + 1$  flavor lattice QCD, which is founded on inconspicuous but vital work [8] that reveals a beneficial algorithm for computing a large number of NBS wave functions simultaneously; see Eqs. (34)-(38) in Ref. [8] for the specific channels of the above 52 NBS wave functions. The purpose of this report is to present a key aspect of the algorithm that performs efficiently a concurrent computation of such a lot of NBS wave functions for various baryon channels. As a preliminary snapshot of the ongoing work, effective  $\Lambda N$  potential at almost physical quark masses corresponding to  $(m_\pi, m_K) \approx (146, 525)\text{MeV}$  is also presented, which is obtained from the single correlator  $\langle p\Lambda p\Lambda \rangle$  by adopting a recipe in which the effects from the  $\Sigma N$  channel are effectively included.

## 2. Effective Baryon Block Algorithm

In the HAL QCD method, the interaction is obtained through the four-point correlator defined by

$$F_{\alpha_1\alpha_2,\alpha_3\alpha_4}^{\langle B_1B_2\overline{B_3B_4} \rangle}(\vec{r}, t - t_0) = \sum_{\vec{X}} \left\langle 0 \left| B_{1,\alpha_1}(\vec{X} + \vec{r}, t) B_{2,\alpha_2}(\vec{X}, t) \overline{\mathcal{J}_{B_3\alpha_3 B_4\alpha_4}(t_0)} \right| 0 \right\rangle, \quad (1)$$

where the summation over  $\vec{X}$  selects states with zero total momentum. The  $B_{1,\alpha_1}(x)$  and  $B_{2,\alpha_2}(y)$  denote the interpolating fields of the baryons such as

$$\begin{aligned} p &= \varepsilon_{abc} (u_a C \gamma_5 d_b) u_c, & n &= -\varepsilon_{abc} (u_a C \gamma_5 d_b) d_c, & \Sigma^+ &= -\varepsilon_{abc} (u_a C \gamma_5 s_b) u_c, & \Sigma^- &= -\varepsilon_{abc} (d_a C \gamma_5 s_b) d_c, \\ \Sigma^0 &= \frac{1}{\sqrt{2}} (X_u - X_d), & \Lambda &= \frac{1}{\sqrt{6}} (X_u + X_d - 2X_s), & \Xi^0 &= \varepsilon_{abc} (u_a C \gamma_5 s_b) s_c, & \Xi^- &= -\varepsilon_{abc} (d_a C \gamma_5 s_b) s_c, \end{aligned}$$

where

$$X_u = \varepsilon_{abc} (d_a C \gamma_5 s_b) u_c, \quad X_d = \varepsilon_{abc} (s_a C \gamma_5 u_b) d_c, \quad X_s = \varepsilon_{abc} (u_a C \gamma_5 d_b) s_c. \quad (2)$$

For simplicity, we have suppressed the explicit spinor indices and spatial coordinates in Eq. (2).  $\overline{\mathcal{J}_{B_3\alpha_3 B_4\alpha_4}(t_0)}$  is the source operator which creates  $B_3 B_4$  states. Hereafter, the explicit time dependence is suppressed for simplicity. The 4pt correlator is evaluated through considering the Wick's contraction together with defining the baryon blocks  $[B_{1,\alpha_1}^{(0)}](\vec{x}; \xi'_{P_1 P_2 P_3})$  and  $[B_{2,\alpha_2}^{(0)}](\vec{y}; \xi'_{P_4 P_5 P_6})$ ,

$$\begin{aligned} F_{\alpha_1\alpha_2,\alpha_3\alpha_4}^{\langle B_1B_2\overline{B_3B_4} \rangle}(\vec{r}) &= \sum_{\vec{X}} \sum_P \sigma_P [B_{1,\alpha_1}^{(0)}](\vec{X} + \vec{r}; \xi'_{P_1}, \xi'_{P_2}, \xi'_{P_3}) [B_{2,\alpha_2}^{(0)}](\vec{X}; \xi'_{P_4}, \xi'_{P_5}, \xi'_{P_6}) \\ &\quad \times \varepsilon_{c'_1 c'_2 c'_3} \varepsilon_{c'_4 c'_5 c'_6} (C \gamma_5)_{\alpha'_1 \alpha'_2} (C \gamma_5)_{\alpha'_4 \alpha'_5} \delta_{\alpha'_3 \alpha_3} \delta_{\alpha'_6 \alpha_4}, \end{aligned} \quad (3)$$

$$\begin{aligned} \text{with } [B_{1,\alpha_1}^{(0)}](\vec{x}; \xi'_{P_1 P_2 P_3}) &= [B_{1,\alpha_1}^{(0)}](\vec{x}; \xi'_{P_1}, \xi'_{P_2}, \xi'_{P_3}) = \langle B_{1,\alpha_1}(\vec{x}) \bar{q}'_{B_{1,3}}(\xi'_{P_3}) \bar{q}'_{B_{1,2}}(\xi'_{P_2}) \bar{q}'_{B_{1,1}}(\xi'_{P_1}) \rangle, \quad \text{and} \\ [B_{2,\alpha_2}^{(0)}](\vec{y}; \xi'_{P_4 P_5 P_6}) &= [B_{2,\alpha_2}^{(0)}](\vec{y}; \xi'_{P_4}, \xi'_{P_5}, \xi'_{P_6}) = \langle B_{2,\alpha_2}(\vec{y}) \bar{q}'_{B_{2,6}}(\xi'_{P_6}) \bar{q}'_{B_{2,5}}(\xi'_{P_5}) \bar{q}'_{B_{2,4}}(\xi'_{P_4}) \rangle, \end{aligned} \quad (4)$$

where  $\sigma_P$  and  $\{\xi'_{P_1}, \dots, \xi'_{P_6}\}$  are the sign factor and the set of permutated spin-color-space-time coordinates for each permutation  $P$  due to the Wick's contraction, respectively. Both 3-tuple sets of the quark fields  $\{\bar{q}'_{B_{1,1}}, \bar{q}'_{B_{1,2}}, \bar{q}'_{B_{1,3}}\}$  and  $\{\bar{q}'_{B_{2,4}}, \bar{q}'_{B_{2,5}}, \bar{q}'_{B_{2,6}}\}$  are ordered so as to create the  $B_1$  and  $B_2$  states properly. Taking the expression in Eq. (3), the number of the iterations to obtain a  $F_{\alpha_1\alpha_2,\alpha_3\alpha_4}^{\langle B_1B_2\overline{B_3B_4} \rangle}(\vec{r})$  except the spatial degrees of freedom reduces to  $(N_c! N_\alpha)^B \times N_u! N_d! N_s! \times 2^{N_\Lambda + N_{\Sigma^0} - B}$  from the number of iterations in naive counting that is  $(N_c! N_\alpha)^{2B} \times N_u! N_d! N_s!$ , where  $N_c = 3, N_\alpha = 4$  and  $N_\Lambda, N_{\Sigma^0}, N_u, N_d, N_s$  and  $B$  are the numbers of  $\Lambda, \Sigma^0$ , up-quark, down-quark, strange-quark and the baryons (i.e., always  $B = 2$  in the present study), respectively. Through the employment of Fast-Fourier-Transform (FFT) [9, 10] we attain the expression in terms of the effective baryon blocks [8]

$$F_{\alpha_1\alpha_2,\alpha_3\alpha_4}^{\langle B_1B_2\overline{B_3B_4} \rangle}(\vec{r}) = \sum_P \sigma_P \sum_{\vec{X}} \left( [B_{1,\alpha_1}^{(P)}](\vec{X} + \vec{r}) \times [B_{2,\alpha_2}^{(P)}](\vec{X}) \right)_{\alpha_3\alpha_4} = \frac{1}{L^3} \sum_{\vec{q}} \left( \sum_P \sigma_P \left( [B_{1,\alpha_1}^{(P)}](\vec{q}) \times [B_{2,\alpha_2}^{(P)}](\vec{-q}) \right)_{\alpha_3\alpha_4} \right) e^{i\vec{q}\cdot\vec{r}}.$$

The effective baryon blocks,  $\left( [B_{1,\alpha_1}^{(P)}](\vec{q}) \times [B_{2,\alpha_2}^{(P)}](\vec{-q}) \right)_{\alpha_3\alpha_4}$ , are obtained by multiplying tensorial factors with the normal baryon blocks; for example, the specific form of the 4pt correlator  $F_{\alpha_1\alpha_2,\alpha_3\alpha_4}^{\langle p\Lambda\overline{pX_u} \rangle}(\vec{r})$  of the  $\langle p\Lambda\overline{pX_u} \rangle$  channel is given by [8],

$$\begin{aligned} F_{\alpha_1\alpha_2,\alpha_3\alpha_4}^{\langle p\Lambda\overline{pX_u} \rangle}(\vec{r}) &= \frac{1}{L^3} \sum_{\vec{q}} \left( [\overline{p}_{\alpha_1\alpha_3}^{(1)}](\vec{q}) [\widetilde{\Lambda}_{\alpha_2\alpha_4}^{(1)}](\vec{-q}) - [\overline{p}_{\alpha_1\alpha_4}^{(2)}]_{c'_3, c'_6}(\vec{q}) [\widetilde{\Lambda}_{\alpha_2\alpha_3}^{(2)}]_{c'_3, c'_6}(\vec{-q}) \right. \\ &\quad - [\overline{p}_{\alpha_1\alpha_3}^{(3)}]_{c'_2, \alpha'_2, c'_4, \alpha'_4}(\vec{q}) [\widetilde{\Lambda}_{\alpha_2\alpha_4}^{(3)}]_{c'_2, \alpha'_2, c'_4, \alpha'_4}(\vec{-q}) + [\overline{p}_{\alpha_1\alpha_4}^{(4)}]_{c'_1, \alpha'_1, c'_5, \alpha'_5}(\vec{q}) [\widetilde{\Lambda}_{\alpha_2\alpha_3}^{(4)}]_{c'_1, \alpha'_1, c'_5, \alpha'_5}(\vec{-q}) \\ &\quad \left. + [\overline{p}_{\alpha_1\alpha_3\alpha_4}^{(5)}]_{c'_1, \alpha'_1, c'_6}(\vec{q}) [\widetilde{\Lambda}_{\alpha_2}^{(5)}]_{c'_1, \alpha'_1, c'_6}(\vec{-q}) - [\overline{p}_{\alpha_1\alpha_3\alpha_4}^{(6)}]_{c'_3, c'_5, \alpha'_5}(\vec{q}) [\widetilde{\Lambda}_{\alpha_2}^{(6)}]_{c'_3, c'_5, \alpha'_5}(\vec{-q}) \right) e^{i\vec{q}\cdot\vec{r}}, \end{aligned} \quad (5)$$

where

$$\begin{aligned} [\widetilde{p}_{\alpha_1\alpha_3}^{(1)}](\vec{q}) &= [\widetilde{p}_{\alpha_1}^{(0)}](\vec{q}; \xi'_{123})\varepsilon_{c'_1c'_2c'_3}(C\gamma_5)_{\alpha'_1\alpha'_2}\delta_{\alpha'_3\alpha_3}, \\ [\widetilde{\Lambda}_{\alpha_2\alpha_4}^{(1)}](-\vec{q}) &= [\widetilde{\Lambda}_{\alpha_2}^{(0)}](-\vec{q}; \xi'_{456})\varepsilon_{c'_4c'_5c'_6}(C\gamma_5)_{\alpha'_4\alpha'_5}\delta_{\alpha'_6\alpha_4}, \end{aligned} \quad (6)$$

$$\begin{aligned} [\widetilde{p}_{\alpha_1\alpha_4}^{(2)}]_{c'_3c'_6}(\vec{q}) &= [\widetilde{p}_{\alpha_1}^{(0)}](\vec{q}; \xi'_{126})\varepsilon_{c'_1c'_2c'_3}(C\gamma_5)_{\alpha'_1\alpha'_2}\delta_{\alpha'_6\alpha_4}, \\ [\widetilde{\Lambda}_{\alpha_2\alpha_3}^{(2)}]_{c'_3c'_6}(-\vec{q}) &= [\widetilde{\Lambda}_{\alpha_2}^{(0)}](-\vec{q}; \xi'_{453})\varepsilon_{c'_4c'_5c'_6}(C\gamma_5)_{\alpha'_4\alpha'_5}\delta_{\alpha'_3\alpha_3}, \end{aligned} \quad (7)$$

$$\begin{aligned} [\widetilde{p}_{\alpha_1\alpha_3}^{(3)}]_{c'_2\alpha'_2c'_4\alpha'_4}(\vec{q}) &= [\widetilde{p}_{\alpha_1}^{(0)}](\vec{q}; \xi'_{143})\varepsilon_{c'_1c'_2c'_3}(C\gamma_5)_{\alpha'_1\alpha'_2}\delta_{\alpha'_3\alpha_3}, \\ [\widetilde{\Lambda}_{\alpha_2\alpha_3}^{(3)}]_{c'_2\alpha'_2c'_4\alpha'_4}(-\vec{q}) &= [\widetilde{\Lambda}_{\alpha_2}^{(0)}](-\vec{q}; \xi'_{256})\varepsilon_{c'_4c'_5c'_6}(C\gamma_5)_{\alpha'_4\alpha'_5}\delta_{\alpha'_6\alpha_4}, \end{aligned} \quad (8)$$

$$\begin{aligned} [\widetilde{p}_{\alpha_1\alpha_4}^{(4)}]_{c'_1\alpha'_1c'_5\alpha'_5}(\vec{q}) &= [\widetilde{p}_{\alpha_1}^{(0)}](\vec{q}; \xi'_{146})\varepsilon_{c'_4c'_5c'_6}(C\gamma_5)_{\alpha'_4\alpha'_5}\delta_{\alpha'_6\alpha_4}, \\ [\widetilde{\Lambda}_{\alpha_2\alpha_3}^{(4)}]_{c'_1\alpha'_1c'_5\alpha'_5}(-\vec{q}) &= [\widetilde{\Lambda}_{\alpha_2}^{(0)}](-\vec{q}; \xi'_{253})\varepsilon_{c'_1c'_2c'_3}(C\gamma_5)_{\alpha'_1\alpha'_2}\delta_{\alpha'_3\alpha_3}, \end{aligned} \quad (9)$$

$$\begin{aligned} [\widetilde{p}_{\alpha_1\alpha_3\alpha_4}^{(5)}]_{c'_1\alpha'_1c'_6}(\vec{q}) &= [\widetilde{p}_{\alpha_1}^{(0)}](\vec{q}; \xi'_{326})\varepsilon_{c'_1c'_2c'_3}(C\gamma_5)_{\alpha'_1\alpha'_2}\delta_{\alpha'_3\alpha_3}\delta_{\alpha'_6\alpha_4}, \\ [\widetilde{\Lambda}_{\alpha_2}^{(5)}]_{c'_1\alpha'_1c'_6}(-\vec{q}) &= [\widetilde{\Lambda}_{\alpha_2}^{(0)}](-\vec{q}; \xi'_{451})\varepsilon_{c'_4c'_5c'_6}(C\gamma_5)_{\alpha'_4\alpha'_5}, \end{aligned} \quad (10)$$

$$\begin{aligned} [\widetilde{p}_{\alpha_1\alpha_3\alpha_4}^{(6)}]_{c'_3c'_5\alpha'_5}(\vec{q}) &= [\widetilde{p}_{\alpha_1}^{(0)}](\vec{q}; \xi'_{346})\varepsilon_{c'_4c'_5c'_6}(C\gamma_5)_{\alpha'_4\alpha'_5}\delta_{\alpha'_3\alpha_3}\delta_{\alpha'_6\alpha_4}, \\ [\widetilde{\Lambda}_{\alpha_2}^{(6)}]_{c'_3c'_5\alpha'_5}(-\vec{q}) &= [\widetilde{\Lambda}_{\alpha_2}^{(0)}](-\vec{q}; \xi'_{251})\varepsilon_{c'_1c'_2c'_3}(C\gamma_5)_{\alpha'_1\alpha'_2}, \end{aligned} \quad (11)$$


$$\begin{aligned} [p_{\alpha_1}^{(0)}](\vec{x}; \xi'_{123}) &= \varepsilon_{b_1b_2b_3}(C\gamma_5)_{\beta_1\beta_2}\delta_{\beta_3\alpha_1} \det \begin{vmatrix} \langle u(\xi_1)\bar{u}(\xi'_1) \rangle & \langle u(\xi_1)\bar{u}(\xi'_3) \rangle \\ \langle u(\xi_3)\bar{u}(\xi'_1) \rangle & \langle u(\xi_3)\bar{u}(\xi'_3) \rangle \end{vmatrix} \langle d(\xi_2)\bar{d}(\xi'_2) \rangle, \\ [p_{\alpha_2}^{(0)}](\vec{y}; \xi'_{456}) &= \frac{1}{\sqrt{6}}\varepsilon_{b_4b_5b_6} \left\{ (C\gamma_5)_{\beta_4\beta_5}\delta_{\beta_6\alpha_2} + (C\gamma_5)_{\beta_5\beta_6}\delta_{\beta_4\alpha_2} - 2(C\gamma_5)_{\beta_6\beta_4}\delta_{\beta_5\alpha_2} \right\} \\ &\quad \times \langle u(\xi_6)\bar{u}(\xi'_6) \rangle \langle d(\xi_4)\bar{d}(\xi'_4) \rangle \langle s(\xi_5)\bar{s}(\xi'_5) \rangle. \end{aligned} \quad (13)$$

NB A roll of coordinates,  $\{\xi'_{123}, \xi'_{456}\}$ , is now called while Ref. [8] has used  $\{\xi'_{142}, \xi'_{563}\}$ ; this is advantageous for generalizing the algorithm toward various (e.g., 52) baryon channels. By employing the effective block algorithm, the number of iterations to evaluate the r.h.s. of Eq. (5) except the momentum space degrees of freedom becomes  $1 + N_c^2 + N_c^2 N_\alpha^2 + N_c^2 N_\alpha^2 + N_c^2 N_\alpha + N_c^2 N_\alpha = 370$ , which is remarkably smaller than the number  $(N_c! N_\alpha)^B \times N_u! N_d! N_s! \times 2^{N_\Lambda + N_{\Sigma^0} - B} = 3456$  that is seen in Eq. (3). The manipulation for the expression of Eq. (5) can be automatically done once the set of the interpolating fields (i.e., the quantum numbers) of both sink and source parts is given.

### 3. Results

A large number of baryon-baryon potentials including the channels from  $NN$  to  $\Xi\Xi$  are studied through the nearly physical point lattice QCD calculation by means of HAL QCD method with  $N_f = 2 + 1$  dynamical clover fermion gauge configurations generated on a  $L^4 = 96^4$  lattice using K computer, where the actual computing jobs are launched with the unified contraction algorithm (UCA) [11]; see also Ref. [8] for the thoroughgoing consistency check in the numerical outputs and comparison at various occasions between the UCA and the present algorithm. Preliminary studies indicate that the physical volume is  $(aL)^4 \approx (8.2\text{fm})^4$  with the lattice spacing  $a \approx 0.085\text{fm}$  and  $(m_\pi, m_K) \approx (146, 525)\text{MeV}$ . See Ref. [12] for details on the generation of the gauge configuration.

Figure 1 shows the preliminary snapshots for the  $\Lambda N$  potential which implicitly includes effects from the  $\Sigma N$ . The statistics in the figure is only a fraction about 1/10 of the entire schedule in FY2015. The left (center) panel shows the central potential in  $^1S_0$  ( $^3S_1$ - $^3D_1$ ) channel. Short-range repulsion and medium-to-long-range attraction are found in the range  $t - t_0 = 7 - 9$  for both central potentials. The right panel shows the tensor potential in the  $^3S_1$ - $^3D_1$  channel. The results in the range  $t - t_0 = 7 - 9$  show weak tensor force in the  $\Lambda N$ - $\Lambda N$  diagonal channel. More comprehensive study which explicitly considers the effects from the  $\Sigma N$  with increasing the statistics will be presented near future.



**Fig. 1.** Left:  $\Lambda N$  central potential in the  $^1S_0$  channel calculated with nearly physical point lattice QCD calculation on a volume  $(96a)^4 \approx (8.2\text{fm})^4$  with the lattice spacing  $a \approx 0.085\text{fm}$  and  $(m_\pi, m_K) \approx (146, 525)\text{MeV}$ . Center:  $\Lambda N$  central potential in the  $^3S_1$ - $^3D_1$  channel. Right:  $\Lambda N$  tensor potential in the  $^3S_1$ - $^3D_1$  channel.

## 4. Summary

In this paper, we present an efficient algorithm to calculate a large number of NBS wave functions simultaneously. The effective block algorithm significantly reduces the numerical costs for the calculation of 52 two-baryon channels and the computer program implemented lays the foundation for the nearly physical point lattice QCD calculation with volume size  $(96a)^4 \approx (8.2\text{fm})^4$ .

The preliminary snapshots of the  $\Lambda N$  potential are presented. Central potentials in both  $^1S_0$  and  $^3S_1$ - $^3D_1$  channels show the short-range repulsion and medium-to-long-range attraction. The tensor potential of the  $\Lambda N$  -  $\Lambda N$  diagonal channel seems to be weaker than the tensor potential of  $NN$  interaction. Further studies explicitly considering  $\Sigma N$  channels together with employing larger statistics are under progress and will be reported in the near future.

## Acknowledgments

The lattice QCD calculations have been performed on the K computer at RIKEN, AICS (Nos. hp120281, hp130023, hp140209, hp150223), HOKUSAI FX100 computer at RIKEN, Wako (No. G15023) and HA-PACS at University of Tsukuba (Nos. 12b-13, 13a-25, 14a-25, 15a-33, 14a-20, 15a-30). We thank ILDG/JLDG [13] which serves as an essential infrastructure in this study. This work is supported in part by MEXT Grant-in-Aid for Scientific Research (25105505, 15K17667, 25287046, 26400281), and SPIRE (Strategic Program for Innovative REsearch) Field 5 project. We thank all collaborators in this project.

## References

- [1] Reviewed in O. Hashimoto and H. Tamura, Prog. Part. Nucl. Phys. **57**, 564 (2006).
- [2] H. Nemura, Y. Akaishi and Y. Suzuki, Phys. Rev. Lett. **89**, 142504 (2002) [arXiv:nucl-th/0203013].
- [3] N. Ishii, S. Aoki, T. Hatsuda, Phys. Rev. Lett. **99**, 022001 (2007).
- [4] S. Aoki, T. Hatsuda and N. Ishii, Prog. Theor. Phys. **123** (2010) 89.
- [5] S. Aoki *et al.* [HAL QCD Collaboration], PTEP **2012**, 01A105 (2012).
- [6] K. Sasaki *et al.* [HAL QCD Collaboration], PTEP **2015**, no. 11, 113B01 (2015).
- [7] T. Doi *et al.*, in these proceedings; arXiv:1512.04199 [hep-lat]; N. Ishii *et al.*, in these proceedings; K. Sasaki *et al.*, in these proceedings;
- [8] H. Nemura, arXiv:1510.00903 [hep-lat].
- [9] N. Ishii, S. Aoki and T. Hatsuda, PoS LATTICE **2008**, 155 (2008) [arXiv:0903.5497 [hep-lat]].
- [10] H. Nemura, N. Ishii, S. Aoki and T. Hatsuda [PACS-CS Collaboration], PoS LATTICE**2008**, 156 (2008) [arXiv:0902.1251 [hep-lat]].
- [11] T. Doi and M. G. Endres, Comput. Phys. Commun. **184**, 117 (2013).
- [12] K.-I. Ishikawa *et al.*, PoS LATTICE **2015**, 075 (2015) [arXiv:1511.09222 [hep-lat]].
- [13] See <http://www.lqcd.org/ildg> and <http://www.jldg.org>