

Heavy-flavored tetraquark states with the $QQ\bar{Q}\bar{Q}$ configuration

Jing Wu¹, Yan-Rui Liu^{1*}

¹*School of Physics and Key Laboratory of Particle Physics and Particle Irradiation (MOE), Shandong University, Jinan 250100, China*

Kan Chen^{2,3}, Xiang Liu^{2,3†}

²*School of Physical Science and Technology, Lanzhou University, Lanzhou 730000, China*

³*Research Center for Hadron and CSR Physics, Lanzhou University and Institute of Modern Physics of CAS, Lanzhou 730000, China*

Shi-Lin Zhu^{4,5,6‡}

⁴*School of Physics and State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, China*

⁵*Collaborative Innovation Center of Quantum Matter, Beijing 100871, China*

⁶*Center of High Energy Physics, Peking University, Beijing 100871, China*

In the framework of the color-magnetic interaction, we systematically investigate the mass spectrum of the tetraquark states composed of four heavy quarks with the $QQ\bar{Q}\bar{Q}$ configuration in this work. We also show their strong decay patterns. Stable or narrow states in the $bb\bar{c}\bar{c}$ and $bc\bar{b}\bar{c}$ systems are found to be possible. We hope the studies shall be helpful to the experimental search for heavy-full exotic tetraquark states.

PACS numbers: 14.40.Rt, 12.39.Pn

I. INTRODUCTION

Searching for exotic states is an interesting research field full of challenges and opportunities. Among the various exotic candidates, the multiquark state is a very popular hadronic configuration. In fact, the concept of the multiquark state was proposed long time ago in Refs. [1–3]. In the past decades, experimentalists have been trying to find them. The observations of the charmonium-like/bottomonium-like XYZ states [4–11] have provided valuable hints of the existence of the exotic states. Especially, these charged charmonium-like or bottomonium-like states like the $Z(4430)$ [12–15], the $Z_1(4050)$ [16], the $Z_2(4250)$ [16], the $Z_c(3900)$ [17–20], the $Z_c(3885)$ [21–23], the $Z_c(4020)$ [24, 25], the $Z_c(4025)$ [26, 27], the $Z_c(4200)$ [14], the $Z_b(10610)$ [28], and the $Z_b(10650)$ [28] have forced us to consider the existence of the multiquark matter very seriously. So many observations in the heavy quark sector in recent years are surprising, which provides new opportunities for us to understand the nature of strong interaction.

Experimentalists continue to surprise us after the observations of XYZ states. In 2015, the LHCb Collaboration reported the hidden-charm pentaquarks $P_c(4380)$ and $P_c(4450)$ in the invariant mass of ψp^1 . They are consistent with the pentaquark structures [29–37]. With a model-independent analysis, the LHCb confirms their previous model-dependent evidence for these states [38]. Very recently, a structure $X(5568)$ was announced by the DØ Collaboration [39]. This narrow state is about 200 MeV below the $B\bar{K}$ threshold and decays into $B_s^0\pi^\pm$. Therefore, the $X(5568)$ may be a typical tetraquark state composed of four different flavors [40–52] while the molecule state assignment to $X(5568)$ is not favored [53]. However, the LHCb Collaboration did not confirm the existence of the $X(5568)$ [54], which makes some theorists consider the difficulty of explaining the $X(5568)$ as a genuine resonance [55–63]. Although there exist different opinions of the $X(5568)$, the observation of the $X(5568)$ again ignites theorist’s enthusiasm of exploring exotic tetraquark states.

As discussed above, there exist possible candidates of the hidden-charm tetraquark states and the tetraquark with a heavy flavor quark and three light quarks. If the multiquark states indeed exist in nature, we have a strong reason to believe that there are more tetraquark states with other flavor configurations. In this work, we focus on the heavy-flavor tetraquark states with the $Q_1Q_2\bar{Q}_3\bar{Q}_4$ configuration, where Q_i is a c or b quark. Although these tetraquark states are still missing in experiment, it is time to carry out a systematic investigation of the mass spectrum of the

*Electronic address: yrliu@sdu.edu.cn

†Electronic address: xiangliu@lzu.edu.cn

‡Electronic address: zhusl@pku.edu.cn

¹ For convenience, here and in the following, ψ means J/ψ once the symbol is adopted.

$Q_1Q_2\bar{Q}_3\bar{Q}_4$ tetraquark system, which may provide important information to further experimental exploration.

In Ref. [64], Iwasaki studied the hidden-charm tetraquark state composed of a pair of charmed quark and anti-quark as well as a pair of light quark and anti-quark, which has the $c\bar{c}q\bar{q}$ configuration. The dimeson configuration ($Q^2\bar{q}^2$) is stable against strong decay into two mesons [65]. Chao performed a systematical investigation of the $cc\bar{c}\bar{c}$ tetraquark system in a quark-gluon model for the first time in Ref. [66]. In Ref. [67], the authors used the Born-Oppenheimer approximation for heavy quarks in the MIT bag and found that the heavy-quark system $c^2\bar{c}^2$ is stable against breakup into two $c\bar{c}$ pairs. But in a potential model calculation [68], the authors suggested that for identical quarks, there is no stable $QQ\bar{Q}\bar{Q}$ state. Similar opinions were shared by the authors of Ref. [69]. However, in Ref. [70], Lloyd and Vary obtained bound tetraquark states by adopting a parameterized Hamiltonian to compute the spectrum of the $cc\bar{c}\bar{c}$ tetraquark states. The tetraquark spectrum was also studied with a generalization of the hyperspherical harmonic formalism in Ref. [71]. In addition, the calculation of the chromomagnetic interaction for the $Q^2\bar{Q}^2$ system was performed in Ref. [72]. In understanding the nature of the $Y(4260)$ meson in Ref. [73], as a byproduct, a lattice study indicates that the $J^{PC} = 1^{--} cc\bar{c}\bar{c}$ state is possible. If the P-wave $cc\bar{c}\bar{c}$ state exists where the orbital angular momentum contributes some repulsion, the lower ground tetraquark states should also exist. Recently, there are further discussions about the properties of fully-heavy tetraquarks [74–84].

In this work, we calculate the mass splittings of the $Q_1Q_2\bar{Q}_3\bar{Q}_4$ tetraquark system in a simple quark model systematically. A typical feature for such tetraquarks is that the isospin is always zero and the flavor wave function is always symmetric for identical quarks. One may only focus on the color-spin part when the Pauli principle is employed to exclude some configurations. For the interaction between the heavy quarks, the short-range gluon exchange force is a dominant source. In the one-gluon-exchange potential model, the color-spin or color-magnetic interaction part is responsible for the mass splitting of the ground hadrons with the same flavor content. In the present study, we will adopt the color-magnetic interaction (CMI) to perform the calculations, which violates the heavy quark symmetry.

We organize the paper as follows. After the introduction, we present the formalism of calculation in Sec. II. We give the numerical results and discuss the possible decay modes in Sec. III. We summarize our results in the final section.

II. FORMALISM

The color-magnetic interaction in the model reads

$$H_{CM} = - \sum_{i,j} C_{ij} \lambda_i \cdot \lambda_j \sigma_i \cdot \sigma_j, \quad (1)$$

where λ_i 's are the Gell-Mann matrices and σ_i 's are the Pauli matrices. The above Hamiltonian was deduced from the one-gluon-exchange interaction [85]. The effective coupling constants C_{ij} incorporate effects from the spatial wave function and the quark masses, which depend on the system. We determine their values in the next section. This Hamiltonian leads to a mass formula for the studied system

$$M = \sum_i m_i + \langle H_{CM} \rangle, \quad (2)$$

where m_i is the effective mass of the i -th constituent quark, which includes the constituent quark mass and effects from other terms such as color-electric interaction and color confinement.

To calculate the matrix elements for the color-spin interaction, one may construct the *color* \otimes *spin* wave functions explicitly and calculate them by definition. In studying multi-quark systems, a simpler way was used in Refs. [86–88]. One just calculates the matrix elements in color space and in spin space separately with the Hamiltonians $H_C = - \sum_{i,j} C_{ij} \lambda_i \cdot \lambda_j$ and $H_S = - \sum_{i,j} C_{ij} \sigma_i \cdot \sigma_j$. Then $\langle H_{CM} \rangle$ is obtained by combining the results of $\langle H_C \rangle$ and $\langle H_S \rangle$ with the common coefficient C_{ij} .

In order to consider the constraint from the Pauli principle, we use a diquark-antidiquark picture to analyze the configurations. Here the notation diquark just means two quarks. In the spin space, the allowed wave functions read

$$\begin{aligned} \chi_1 &= |(Q_1Q_2)_1(\bar{Q}_3\bar{Q}_4)_1\rangle_2, & \chi_2 &= |(Q_1Q_2)_1(\bar{Q}_3\bar{Q}_4)_1\rangle_1, \\ \chi_3 &= |(Q_1Q_2)_1(\bar{Q}_3\bar{Q}_4)_1\rangle_0, & \chi_4 &= |(Q_1Q_2)_1(\bar{Q}_3\bar{Q}_4)_0\rangle_1, \\ \chi_5 &= |(Q_1Q_2)_0(\bar{Q}_3\bar{Q}_4)_1\rangle_1, & \chi_6 &= |(Q_1Q_2)_0(\bar{Q}_3\bar{Q}_4)_0\rangle_0, \end{aligned} \quad (3)$$

where the subscripts on the right hand side denote the spin of the Q_1Q_2 , spin of the $\bar{Q}_3\bar{Q}_4$, and that of the system, respectively. According to the $SU(3)$ group theory, the diquark in the color space belongs to the representation 6_c or $\bar{3}_c$, while the anti-diquark's representation is $\bar{6}_c$ or 3_c . Then one has two kinds of color-singlet state

$$\phi_1 = |(Q_1Q_2)^6(\bar{Q}_3\bar{Q}_4)^{\bar{6}}\rangle, \quad \phi_2 = |(Q_1Q_2)^{\bar{3}}(\bar{Q}_3\bar{Q}_4)^3\rangle. \quad (4)$$

Combine the spin and color wave functions, we get twelve possible $color \otimes spin$ wave functions for the $Q_1 Q_2 \bar{Q}_3 \bar{Q}_4$ system

$$\begin{aligned}
\phi_1 \chi_1 &= |(Q_1 Q_2)_1^6 (\bar{Q}_3 \bar{Q}_4)_1^{\bar{6}}\rangle_2 \delta_{12} \delta_{34}, & \phi_2 \chi_1 &= |(Q_1 Q_2)_1^{\bar{3}} (\bar{Q}_3 \bar{Q}_4)_1^3\rangle_2, \\
\phi_1 \chi_2 &= |(Q_1 Q_2)_1^6 (\bar{Q}_3 \bar{Q}_4)_1^{\bar{6}}\rangle_1 \delta_{12} \delta_{34}, & \phi_2 \chi_2 &= |(Q_1 Q_2)_1^{\bar{3}} (\bar{Q}_3 \bar{Q}_4)_1^3\rangle_1, \\
\phi_1 \chi_3 &= |(Q_1 Q_2)_1^6 (\bar{Q}_3 \bar{Q}_4)_1^{\bar{6}}\rangle_0 \delta_{12} \delta_{34}, & \phi_2 \chi_3 &= |(Q_1 Q_2)_1^{\bar{3}} (\bar{Q}_3 \bar{Q}_4)_1^3\rangle_0, \\
\phi_1 \chi_4 &= |(Q_1 Q_2)_1^6 (\bar{Q}_3 \bar{Q}_4)_0^{\bar{6}}\rangle_1 \delta_{12}, & \phi_2 \chi_4 &= |(Q_1 Q_2)_1^{\bar{3}} (\bar{Q}_3 \bar{Q}_4)_0^3\rangle_1 \delta_{34}, \\
\phi_1 \chi_5 &= |(Q_1 Q_2)_0^6 (\bar{Q}_3 \bar{Q}_4)_1^{\bar{6}}\rangle_1 \delta_{34}, & \phi_2 \chi_5 &= |(Q_1 Q_2)_0^{\bar{3}} (\bar{Q}_3 \bar{Q}_4)_1^3\rangle_1 \delta_{12}, \\
\phi_1 \chi_6 &= |(Q_1 Q_2)_0^6 (\bar{Q}_3 \bar{Q}_4)_0^{\bar{6}}\rangle_0, & \phi_2 \chi_6 &= |(Q_1 Q_2)_0^{\bar{3}} (\bar{Q}_3 \bar{Q}_4)_0^3\rangle_0 \delta_{12} \delta_{34},
\end{aligned} \tag{5}$$

where the used notation is $|(Q_1 Q_2)_{spin}^{color} (\bar{Q}_3 \bar{Q}_4)_{spin}^{color}\rangle_{spin}$. We insert a symbol δ_{ij} in the wave functions to reflect the constraint from the Pauli principle. When the i -th quark and the j -th quark are the same, $\delta_{ij} = 0$. Otherwise, $\delta_{ij} = 1$.

Replacing Q_i with c or b quark, one gets nine possibilities for the flavor content, six of which need to be studied: $bbbb$, $cc\bar{c}\bar{c}$, $bb\bar{c}\bar{c}$, $bb\bar{b}\bar{c}$, $cc\bar{c}\bar{b}$, and $bcb\bar{c}$. The other three cases $ccb\bar{b}$, $cbb\bar{b}$, and $bc\bar{c}\bar{c}$ correspond to the antiparticles of $bb\bar{c}\bar{c}$, $bb\bar{b}\bar{c}$, and $cc\bar{c}\bar{b}$, respectively. So their formulas are not independent. Here, considering the Pauli principle, one may categorize the six systems into three sets: (1) $bbbb$, $cc\bar{c}\bar{c}$, and $bb\bar{c}\bar{c}$, where $\delta_{12} = \delta_{34} = 0$; (2) $bb\bar{b}\bar{c}$ and $ccb\bar{c}$, where $\delta_{12} = 0, \delta_{34} = 1$; and (3) $bcb\bar{c}$ where $\delta_{12} = \delta_{34} = 1$. The number of independent wave functions for them is 4, 6, and 12, respectively. We present the resulting CMI matrix elements system by system. By the way, the usually mentioned ‘‘good’’ diquark (spin=0, color= $\bar{3}_c$) exists only in the systems containing the (bc) substructure since the most attractive $(bb)_{spin=0}^{\bar{3}_c}$ and $(cc)_{spin=0}^{\bar{3}_c}$ objects are forbidden by the Pauli principle.

A. The $bb\bar{b}\bar{b}$ and $cc\bar{c}\bar{c}$ systems

The color-spin structures of these two systems are the same. The only difference lies in the quark mass. So we put them together for discussions.

The $bb\bar{b}\bar{b}$ system is a neutral state. Its possible quantum numbers are $I^G(J^{PC}) = 0^+(2^{++})$, $0^-(1^{+-})$, or $0^+(0^{++})$. The number of states is constrained by the Pauli principle. For the case $J = 2$, the wave function is $\phi_2 \chi_1$ and the obtained $\langle H_{CM} \rangle$ is given by $\frac{16}{3}(C_{bb} + C_{\bar{b}\bar{b}})$. The color-magnetic interaction is certainly repulsive. For the case $J = 1$, the wave function is $\phi_2 \chi_2$ and $\langle H_{CM} \rangle = \frac{16}{3}(C_{bb} - C_{\bar{b}\bar{b}})$. Since the quark-quark interaction and the quark-antiquark interaction is related with the G-parity transformation, the CMI for this $J = 1$ state is expected to be weak. For the case $J = 0$, there are two possible wave functions $\phi_2 \chi_3$ and $\phi_1 \chi_6$. Although the color-spin structures are different, their mixing is allowed by the color-magnetic interaction. The obtained symmetric CMI matrix in the $(\phi_2 \chi_3, \phi_1 \chi_6)^T$ base is

$$\langle H_{CM} \rangle = \begin{pmatrix} \frac{16}{3}(C_{bb} - 2C_{\bar{b}\bar{b}}) & 8\sqrt{6}C_{\bar{b}\bar{b}} \\ & 8C_{\bar{b}\bar{b}} \end{pmatrix}. \tag{6}$$

The $cc\bar{c}\bar{c}$ system has similar expressions. For the case $J = 2$, $\langle H_{CM} \rangle = \frac{16}{3}(C_{cc} + C_{\bar{c}\bar{c}})$ with the color-spin wave function $\phi_2 \chi_1$. For the case $J = 1$, $\langle H_{CM} \rangle = \frac{16}{3}(C_{cc} - C_{\bar{c}\bar{c}})$ with the wave function $\phi_2 \chi_2$. For $J = 0$, the CMI matrix is

$$\langle H_{CM} \rangle = \begin{pmatrix} \frac{16}{3}(C_{cc} - 2C_{\bar{c}\bar{c}}) & 8\sqrt{6}C_{\bar{c}\bar{c}} \\ & 8C_{\bar{c}\bar{c}} \end{pmatrix}. \tag{7}$$

The above compact tetraquark system has the same quark content as the molecular states composed of two bottomonium (charmonium) mesons. However, the interaction between heavy quarks is dominantly through the short-range gluon exchange. Once the interaction is strong enough to bind the two mesons, the resulting object very probably tends to form a compact structure instead of a loosely-bound molecule.

In the extreme case that the molecules exist, the configuration mixing is possible. For the S -wave $\Upsilon\Upsilon$ ($\psi\psi$) state, the allowed quantum numbers are $I^G(J^{PC}) = 0^+(2^{++})$ or $0^+(0^{++})$ since the wave function in spin space should be symmetric for identical mesons. The quantum numbers for the S -wave state composed of two η_b (η_c) mesons are only $0^+(0^{++})$. For the state composed of one η_b (η_c) and one Υ (ψ), the quantum numbers are just $I^G(J^{PC}) = 0^-(1^{+-})$. Therefore, the allowed J^{PC} appear both in the molecule and diquark-antidiquark pictures. The number of states is also equivalent. Considering that the interaction between heavy quarks is through the gluon exchange force, one does not expect large mass difference between these two configurations. The compact structure may contribute significantly to the properties of the molecules with the same quantum numbers.

If the tetraquark states have larger masses, the quark rearrangements into the meson-meson channels with the same quantum numbers may happen. We will discuss such possible decay patterns after their masses are estimated.

B. The $bb\bar{c}\bar{c}$ and $cc\bar{b}\bar{b}$ systems

The two systems are related through the C -parity transformation and they have the same color-spin matrix elements. The possible quantum numbers of them are $I(J^P) = 0(2^+)$, $0(1^+)$, or $0(0^+)$. Again the Pauli principle results in four states for the $bb\bar{c}\bar{c}$ (or $cc\bar{b}\bar{b}$) system. The color-spin wave functions are $\phi_2\chi_1$ and $\phi_2\chi_2$ for the case of $J = 2$ and $J = 1$, respectively. The corresponding $\langle H_{CM} \rangle$'s are $\frac{8}{3}(C_{bb} + C_{cc} + 2C_{b\bar{c}})$ and $\frac{8}{3}(C_{bb} + C_{cc} - 2C_{b\bar{c}})$. For the case $J = 0$, the allowed wave functions are the same as those of the previous systems, $\phi_2\chi_3$ and $\phi_1\chi_6$. Consider their mixing, one gets

$$\langle H_{CM} \rangle = \begin{pmatrix} \frac{8}{3}(C_{bb} + C_{cc} - 4C_{b\bar{c}}) & 8\sqrt{6}C_{b\bar{c}} \\ & 4(C_{bb} + C_{cc}) \end{pmatrix}. \quad (8)$$

These systems have the same quark content as the $B_c^{(*)}B_c^{(*)}$ meson-meson states. The quantum numbers of $B_c^{*-}B_c^{*-}$ ($B_c^{*+}B_c^{*+}$) are $I(J^P) = 0(2^+)$ or $0(0^+)$, those of $B_c^-B_c^-$ ($B_c^+B_c^+$) are $0(0^+)$. There is only one $B_c^-B_c^{*-}$ ($B_c^+B_c^{*+}$) state with $I(J^P) = 0(1^+)$.

C. The $bb\bar{b}\bar{c}$ and $cb\bar{b}\bar{b}$ systems

The two systems have the same matrix elements. Their quantum numbers are also $I(J^P) = 0(2^+)$, $0(1^+)$, or $0(0^+)$. We here focus on the $bb\bar{b}\bar{c}$ system. Now the number of the vector states is 3 and that of the scalar states is 2. For the $J = 2$ case, the color-spin wave function is again $\phi_2\chi_1$. The resulting color-magnetic matrix element is $\langle H_{CM} \rangle = \frac{8}{3}(C_{bb} + C_{bc} + C_{b\bar{b}} + C_{b\bar{c}})$. For the case $J = 1$, three possible wave functions $\phi_2\chi_2$, $\phi_2\chi_4$, and $\phi_1\chi_5$ are allowed. The last one has different color structure from the other two. Although their mixing occurs, from the obtained matrix [base: $(\phi_2\chi_2, \phi_2\chi_4, \phi_1\chi_5)^T$]

$$\langle H_{CM} \rangle = \begin{pmatrix} \frac{8}{3}(C_{bb} + C_{bc} - C_{b\bar{b}} - C_{b\bar{c}}) & \frac{8\sqrt{2}}{3}(C_{b\bar{b}} - C_{b\bar{c}}) & 8(C_{b\bar{b}} - C_{b\bar{c}}) \\ & \frac{8}{3}(C_{bb} - 3C_{bc}) & -4\sqrt{2}(C_{b\bar{c}} + C_{b\bar{b}}) \\ & & \frac{4}{3}(3C_{bb} - C_{bc}) \end{pmatrix}, \quad (9)$$

one observes that the mixing strength for $\phi_2\chi_2$ and $\phi_2\chi_4$ and that for $\phi_2\chi_2$ and $\phi_1\chi_5$ may be both small. The remaining case is for $J = 0$, where possible wave functions are $\phi_2\chi_3$ and $\phi_1\chi_6$. Now, one has

$$\langle H_{CM} \rangle = \begin{pmatrix} \frac{8}{3}(C_{bb} + C_{bc} - 2C_{b\bar{b}} - 2C_{b\bar{c}}) & 4\sqrt{6}(C_{b\bar{b}} + C_{b\bar{c}}) \\ & 4(C_{bb} + C_{bc}) \end{pmatrix}. \quad (10)$$

The meson-meson systems with the quark content $bb\bar{b}\bar{c}$ are ΥB_c^- , ΥB_c^{*-} , $\eta_b B_c^-$, and $\eta_b B_c^{*-}$. Their quantum numbers are $I(J^P) = 0(1^+)$, $0([2, 1, 0]^+)$, $0(0^+)$, and $0(1^+)$, respectively.

D. The $cc\bar{c}\bar{b}$ and $bc\bar{c}\bar{c}$ systems

The situation is similar to the $bb\bar{b}\bar{c}$ and $cb\bar{b}\bar{b}$ systems. By exchanging b and c there, one easily gets relevant matrix elements. For comparison, we focus on the $cc\bar{c}\bar{b}$ system. For the case $J = 2$, one has $\langle H_{CM} \rangle = \frac{8}{3}(C_{cc} + C_{bc} + C_{b\bar{c}} + C_{c\bar{c}})$. For the case $J = 1$, the matrix for the color-spin interaction reads

$$\langle H_{CM} \rangle = \begin{pmatrix} \frac{8}{3}(C_{cc} + C_{bc} - C_{b\bar{c}} - C_{c\bar{c}}) & \frac{8\sqrt{2}}{3}(C_{c\bar{c}} - C_{b\bar{c}}) & 8(C_{c\bar{c}} - C_{b\bar{c}}) \\ & \frac{8}{3}(C_{cc} - 3C_{bc}) & -4\sqrt{2}(C_{c\bar{c}} + C_{b\bar{c}}) \\ & & \frac{4}{3}(3C_{cc} - C_{bc}) \end{pmatrix}. \quad (11)$$

For the case $J = 0$, the matrix is

$$\langle H_{CM} \rangle = \begin{pmatrix} \frac{8}{3}(C_{cc} + C_{bc} - 2C_{b\bar{c}} - 2C_{c\bar{c}}) & 4\sqrt{6}(C_{b\bar{c}} + C_{c\bar{c}}) \\ & 4(C_{cc} + C_{bc}) \end{pmatrix}. \quad (12)$$

The signs for the non-diagonal matrix elements seem to be inconsistent with the previous systems after the replacements $b \rightarrow c$ and $c \rightarrow b$. Actually they do not affect the final results. For detailed argument, one may consult Eq. (2) of Ref. [89] and relevant explanations there.

The meson-meson states that these tetraquarks can rearrange into are ψB_c , ψB_c^* , $\eta_c B_c$, and $\eta_c B_c^*$.

E. The $b\bar{c}\bar{b}$ system

The Pauli principle does not give any constraints on the wave functions now. The two wave functions for $J = 2$ and the four wave functions for $J = 0$ will mix, respectively. However, one should be careful in discussing the mixing with the six wave functions for $J = 1$ because the system is neutral and may have C -parity.

If $J = 2$, both the diquark and the antidiquark have angular momentum 1. The state should have definite C -parity + and the quantum numbers are $I^G(J^{PC}) = 0^+(2^{++})$. With the base $(\phi_1\chi_1, \phi_2\chi_1)^T$, one may get the CMI matrix

$$\langle H_{CM} \rangle = \begin{pmatrix} -\frac{2}{3}(4C_{bc} - 5C_{b\bar{b}} - 10C_{b\bar{c}} - 5C_{c\bar{c}}) & 2\sqrt{2}(C_{b\bar{b}} - 2C_{b\bar{c}} + C_{c\bar{c}}) \\ \frac{4}{3}(4C_{bc} + C_{b\bar{b}} + 2C_{b\bar{c}} + C_{c\bar{c}}) & \end{pmatrix}. \quad (13)$$

If $J = 0$, both the diquark and the antidiquark have the same angular momentum. The quantum numbers for the system are $I^G(J^{PC}) = 0^+(0^{++})$. The obtained CMI matrix is

$$\langle H_{CM} \rangle = \begin{pmatrix} -\frac{4}{3} \begin{pmatrix} 2C_{bc} + 5C_{b\bar{b}} \\ +10C_{b\bar{c}} + 5C_{c\bar{c}} \end{pmatrix} & -\frac{10}{\sqrt{3}}(C_{b\bar{b}} - 2C_{b\bar{c}} + C_{c\bar{c}}) & 4\sqrt{2}(C_{b\bar{b}} - 2C_{b\bar{c}} + C_{c\bar{c}}) & 2\sqrt{6}(C_{b\bar{b}} + 2C_{b\bar{c}} + C_{c\bar{c}}) \\ 8C_{bc} & & 2\sqrt{6}(C_{b\bar{b}} + 2C_{b\bar{c}} + C_{c\bar{c}}) & 0 \\ & & \frac{8}{3}(2C_{bc} - C_{b\bar{b}} - 2C_{b\bar{c}} - C_{c\bar{c}}) & -\frac{4}{\sqrt{3}}(C_{b\bar{b}} - 2C_{b\bar{c}} + C_{c\bar{c}}) \\ & & & -16C_{bc} \end{pmatrix}, \quad (14)$$

where the base is $(\phi_1\chi_3, \phi_1\chi_6, \phi_2\chi_3, \phi_2\chi_6)^T$.

If $J = 1$, the states $\phi_1\chi_2$ and $\phi_2\chi_2$ have negative C -parities. All the other four wave functions $\phi_1\chi_4$, $\phi_2\chi_4$, $\phi_1\chi_5$, and $\phi_2\chi_5$ are not invariant under C -parity transformation. But we may construct four states which are invariant under C -parity transformations. The basic procedure is similar to that given in Ref. [90]. Explicitly, the two $C = +$ states are

$$\begin{aligned} [\phi\chi]_+^{6\bar{6}} &= \frac{1}{\sqrt{2}}(\phi_1\chi_4 + \phi_1\chi_5), \\ [\phi\chi]_+^{\bar{3}3} &= \frac{1}{\sqrt{2}}(\phi_2\chi_4 + \phi_2\chi_5), \end{aligned} \quad (15)$$

and the two $C = -$ states are

$$\begin{aligned} [\phi\chi]_-^{6\bar{6}} &= \frac{1}{\sqrt{2}}(\phi_1\chi_4 - \phi_1\chi_5), \\ [\phi\chi]_-^{\bar{3}3} &= \frac{1}{\sqrt{2}}(\phi_2\chi_4 - \phi_2\chi_5). \end{aligned} \quad (16)$$

Only states with the same quantum numbers may mix. So we have two color-spin matrices. For the states with $I^G(J^{PC}) = 0^+(1^{++})$, the matrix is

$$\langle H_{CM} \rangle = \begin{pmatrix} \frac{2}{3}(4C_{bc} + 5C_{b\bar{b}} + 5C_{c\bar{c}} - 10C_{b\bar{c}}) & -2\sqrt{2}(C_{b\bar{b}} + C_{c\bar{c}} + 2C_{b\bar{c}}) \\ \frac{4}{3}(-4C_{bc} + C_{b\bar{b}} + C_{c\bar{c}} - 2C_{b\bar{c}}) & \end{pmatrix}, \quad (17)$$

with the base $([\phi\chi]_+^{6\bar{6}}, [\phi\chi]_+^{\bar{3}3})^T$. For the states with $I^G(J^{PC}) = 0^-(1^{+-})$, the matrix reads

$$\langle H_{CM} \rangle = \begin{pmatrix} -\frac{2}{3} \begin{pmatrix} 4C_{bc} + 5C_{b\bar{b}} \\ +5C_{c\bar{c}} + 10C_{b\bar{c}} \end{pmatrix} & 2\sqrt{2}(C_{b\bar{b}} + C_{c\bar{c}} - 2C_{b\bar{c}}) & \frac{20}{3}(C_{b\bar{b}} - C_{c\bar{c}}) & -4\sqrt{2}(C_{b\bar{b}} - C_{c\bar{c}}) \\ \frac{4}{3} \begin{pmatrix} 4C_{bc} - C_{b\bar{b}} \\ -C_{c\bar{c}} - 2C_{b\bar{c}} \end{pmatrix} & -4\sqrt{2}(C_{b\bar{b}} - C_{c\bar{c}}) & \frac{8}{3}(C_{b\bar{b}} - C_{c\bar{c}}) & \\ & \frac{2}{3} \begin{pmatrix} 4C_{bc} - 5C_{b\bar{b}} \\ -5C_{c\bar{c}} + 10C_{b\bar{c}} \end{pmatrix} & 2\sqrt{2}(C_{b\bar{b}} + C_{c\bar{c}} + 2C_{b\bar{c}}) & \\ & & & -\frac{4}{3} \begin{pmatrix} 4C_{bc} + C_{b\bar{b}} \\ +C_{c\bar{c}} - 2C_{b\bar{c}} \end{pmatrix} \end{pmatrix}, \quad (18)$$

where the base is $(\phi_1\chi_2, \phi_2\chi_2, [\phi\chi]_{-}^{66}, [\phi\chi]_{-}^{33})^T$.

There are two kinds of molecule configurations for the $b\bar{c}b\bar{c}$ system. In the bottomonium+charmonium case, the allowed quantum numbers are $I^G(J^{PC}) = 0^+(0^{++})$ for the $\eta_b\eta_c$ system, $0^-(1^{+-})$ for the $\eta_b\psi$ system, $0^-(1^{+-})$ for the $\Upsilon\eta_c$, and $0^+([2, 1, 0]^{++})$ for the $\Upsilon\psi$ system. In the meson-antimeson case, those for $B_c^-B_c^+$ are $0^+(0^{++})$, those for $B_c^-B_c^{*+} \pm B_c^{*-}B_c^+$ are $0^\pm(1^{+\pm})$, and those for $B_c^{*-}B_c^{*+}$ are $0^+([2, 0]^{++})$ or $0^-(1^{+-})$.

III. NUMERICAL RESULTS

A. Parameters

We need to determine six coefficients $C_{b\bar{b}}$, $C_{c\bar{c}}$, $C_{b\bar{c}}$, C_{bb} , C_{cc} , and C_{bc} in discussing the mass splittings for various $Q_1Q_2\bar{Q}_3\bar{Q}_4$ systems. Their masses may be further estimated with the Hamiltonian in Eq. (2) once the effective quark masses m_c and m_b are used.

From the mass splitting between Υ and η_b , $m_\Upsilon - m_{\eta_b} = [\frac{16}{3}C_{b\bar{b}}] - [-16C_{b\bar{b}}] = 61$ MeV [91], one extracts $C_{b\bar{b}} = 2.9$ MeV. Similarly, the value of $C_{c\bar{c}} = 5.3$ MeV is obtained from the mass splitting $m_{J/\psi} - m_{\eta_c} = 114$ MeV. Since the excited B_c^* meson has not been observed yet, we just estimate the value of $C_{b\bar{c}}$ to be 3.3 MeV from $m_{B_c^*} - m_{B_c} = 70$ MeV calculated with a quark model [92]. Although the Ξ_{cc}^{++} baryon was confirmed recently by the LHCb Collaboration [93] after the first observation at SELEX [94, 95], the available heavy baryon masses are still not enough for us to extract the value of C_{cc} . Here we perform our calculation with the approximation $C_{QQ} = C_{Q\bar{Q}}$ ($Q = c, b$). Since there is no dynamics in the present model, the choice of the approximation to determine C_{QQ} is not unique. However, the results induced by the change of C_{QQ} should not be large [96, 97]. For comparison, we also adopt the approximation $\frac{C_{QQ}}{C_{Q\bar{Q}}} = \frac{C_{nn}}{C_{n\bar{n}}} \approx \frac{2}{3}$ and check the extreme case $C_{QQ} = 0$, where $C_{nn} = 18.4$ MeV is extracted from the light baryon masses [98]. By using the mass difference between ρ and π , one gets $C_{n\bar{n}} = 29.8$ MeV. The latter approximation certainly gives a smaller C_{QQ} . If the interactions within the diquarks are effectively attractive (repulsive), the approximation $\frac{C_{QQ}}{C_{Q\bar{Q}}} \approx \frac{2}{3}$ should result in heavier (lighter) tetraquark states. Here, the effective interaction within diquarks reflects the repulsion or attraction effect from the enhancement or cancellation between the quark-quark (and antiquark-antiquark) interaction in the case of channel coupling. For comparison, we use these two approximations in our estimation. The values of the quark-quark coupling parameters estimated with them are listed in Table I.

TABLE I: Two sets of effective coupling parameters in units of MeV with different approximations.

	Set I ($C_{QQ} = C_{Q\bar{Q}}$)	Set II ($\frac{C_{QQ}}{C_{Q\bar{Q}}} = \frac{C_{nn}}{C_{n\bar{n}}}$)
C_{cc}	5.3	3.3
C_{bb}	2.9	1.9
C_{bc}	3.3	2.0

To determine the masses of the tetraquarks, we adopt two approaches in the present work: 1) One estimates the meson masses with the effective heavy quark masses $m_b = 5052.9$ MeV and $m_c = 1724.8$ MeV. These values were adopted in understanding the strange properties of tetraquark states [98, 99] and the pentaquark states [100]; 2) One calculates the masses from a meson-meson threshold, where the mass formula is $M = M_{th} - \langle H_{CM} \rangle_{th} + \langle H_{CM} \rangle$ and the relevant meson masses are: $m_\Upsilon = 9460.3$ MeV, $m_\psi = 3096.9$ MeV, and $m_{B_c} = 6275.1$ MeV [91]. The latter method has been used in estimating the mass of an exotic T_{cc} [101, 102].

B. The $b\bar{b}b\bar{b}$ and $c\bar{c}c\bar{c}$ systems

It is easy to get the numerical results for the CMI matrix elements with the above two sets of parameters. Adopting the approximation $C_{QQ} = C_{Q\bar{Q}}$, we obtain the CMI matrices, their eigenvalues and corresponding eigenvectors, and estimated masses with two different approaches. These results are presented in Table II. In the approximation $C_{QQ} = \frac{C_{nn}}{C_{n\bar{n}}}C_{Q\bar{Q}}$, the estimated masses of the tetraquark states are slightly different from those in the former approximation. In the following discussions, we mainly use the masses estimated with the parameters in Set I. Since the hadron masses estimated with the effective quarks are usually like an upper limit [98–100, 103], we focus on the results estimated with reference thresholds. We assume that these masses are all reasonable values. To have an impression for the

spectrum, we plot relative positions for the $bb\bar{b}\bar{b}$ ($cc\bar{c}\bar{c}$) states in Fig. 1 (a) [(b)]. The solid/black lines are for the approximation $C_{QQ} = C_{Q\bar{Q}}$ and the dashdotted/blue lines are for the approximation $C_{QQ} = \frac{C_{nn}}{C_{n\bar{n}}}C_{Q\bar{Q}}$. We also show the results in the extreme case $C_{QQ} = 0$ with the dashed/red lines. The uncertainty caused by the change of C_{QQ} is less than 20 (37) MeV in the $bb\bar{b}\bar{b}$ ($cc\bar{c}\bar{c}$) case.

TABLE II: Results for the $bb\bar{b}\bar{b}$ and $cc\bar{c}\bar{c}$ systems in units of MeV with the approximation $C_{QQ} = C_{Q\bar{Q}}$. The masses in the fifth column are estimated with $m_b = 5052.9$ MeV and $m_c = 1724.8$ MeV. The last column lists masses estimated from the $(\Upsilon\Upsilon)$ or $(J/\psi J/\psi)$ threshold. The base for the $J = 0$ case is $(\phi_2\chi_3, \phi_1\chi_6)^T$.

System	J^{PC}	$\langle H_{CM} \rangle$	Eigenvalue	Eigenvector	Mass	$(\Upsilon\Upsilon)/(\psi\psi)$
$(bb\bar{b}\bar{b})$	2^{++}	30.9	30.9	1	20243	18921
	1^{+-}	0.0	0.0	1	20212	18890
	0^{++}	$\begin{pmatrix} -15.5 & 56.8 \\ 56.8 & 23.2 \end{pmatrix}$	$\begin{bmatrix} 63.9 \\ -56.2 \end{bmatrix}$	$\begin{bmatrix} (0.58, 0.81) \\ (-0.81, 0.58) \end{bmatrix}$	$\begin{bmatrix} 20275 \\ 20155 \end{bmatrix}$	$\begin{bmatrix} 18954 \\ 18834 \end{bmatrix}$
$(cc\bar{c}\bar{c})$	2^{++}	56.5	56.5	1	6956	6194
	1^{+-}	0.0	0.0	1	6899	6137
	0^{++}	$\begin{pmatrix} -28.3 & 103.9 \\ 103.9 & 42.4 \end{pmatrix}$	$\begin{bmatrix} 116.8 \\ -102.6 \end{bmatrix}$	$\begin{bmatrix} (0.58, 0.81) \\ (-0.81, 0.58) \end{bmatrix}$	$\begin{bmatrix} 7016 \\ 6797 \end{bmatrix}$	$\begin{bmatrix} 6254 \\ 6035 \end{bmatrix}$

The mass splitting between the scalar tetraquarks is around 120 MeV for the bottom system and 220 MeV for the charmed system. One finds that the mixing between different color structures is important here, which enlarges the mass difference between these two states. If one does not consider the mixing, the masses for the bottom case are 18913 MeV and 18875 MeV. Both states are below the $\Upsilon\Upsilon$ threshold and above the $\eta_b\Upsilon$ threshold. Once the mixing is considered, the higher state (6_c bb dominates) becomes a state above the $\Upsilon\Upsilon$ threshold while the lower one (3_c bb dominates) becomes a state below the $\eta_b\Upsilon$ threshold. Certainly the mass shift affects decay properties. The charmed case is similar.

From the diagrams (a) and (b) in Fig. 1, the estimated tetraquark masses are all above the lowest meson-meson threshold. This observation is consistent with those in Refs. [74, 79, 82]. The lowest $bb\bar{b}\bar{b}$ mass in [79] and ours are similar. From these diagrams, the masses obtained with parameters in Set II are all lower than those in Set I. This means that the interactions within the diquarks are effectively repulsive, which can be verified from the Hamiltonian expressions. If stable multi-quark states need attractive diquarks, these $bb\bar{b}\bar{b}$ and $cc\bar{c}\bar{c}$ compact tetraquarks would tend to become meson-meson states because the quark-antiquark interaction is usually attractive (see the diagonal matrix elements in the Hamiltonian expressions) and these tetraquarks should not be stable.

If the studied states do exist, finding out their decay properties are helpful to the search at experiments. Possible rearrangement decay modes are easy to be understood from Fig. 1. Since the feature for the $cc\bar{c}\bar{c}$ system is very similar to the bottom case, we here only concentrate on the latter one. For the $J = 2$ tetraquark, the present model estimation gives a mass around the $\Upsilon\Upsilon$ threshold. If the approximation $C_{QQ} \approx \frac{2}{3}C_{Q\bar{Q}}$ is more appropriate, the state is below the threshold and it should have a relatively narrow width through D -wave decay into $\eta_b\eta_b$. It is a basic feature that high spin multi-quark states have dominantly D -wave decay modes and should not be very broad [99, 100, 103]. For the $J = 1$ tetraquark, its mass is 30 MeV above the $\eta_b\Upsilon$ threshold. From the quantum numbers, its rearrangement decay channel is only this $\eta_b\Upsilon$. For the $J = 0$ tetraquarks, the higher one can decay into both $\Upsilon\Upsilon$ and $\eta_b\eta_b$ through both S - and D -wave interactions while the lower one decays only into $\eta_b\eta_b$. If we use their masses to denote these states, probably the ordering of the widths is $18954 > 18890 \sim 18834 > 18921$.

From Fig. 1 (a)-(b), the feature that all the states can decay is consistent with the feature that the effective interaction within the diquarks is repulsive. If we want to find relatively stable compact tetraquarks in Fig. 1, good candidates should be those states for which dashed/red lines are above solid/black lines. Although the $bb\bar{b}\bar{b}$ and $cc\bar{c}\bar{c}$ systems do not satisfy this condition, we will see that such systems exist.

C. The $bb\bar{c}\bar{c}$ and $cc\bar{b}\bar{b}$ systems

We present the numerical results for the $cc\bar{b}\bar{b}$ states in Table III. The $bb\bar{c}\bar{c}$ states are antiparticles of the $cc\bar{b}\bar{b}$ states and have the same results. Now the mass splitting between different spins is less than 140 MeV. This number lies between the splittings for the $bb\bar{b}\bar{b}$ case and the $cc\bar{c}\bar{c}$ case and thus the mixing effect is in the middle of the two. To understand the decay properties easily, we plot the spectrum for the $cc\bar{b}\bar{b}$ system and relevant meson-meson thresholds in Fig. 1 (c).




FIG. 1: Relative positions for the considered tetraquark states. The solid (black) and dashdotted (blue) lines correspond to masses estimated with the approximations $C_{QQ} = C_{Q\bar{Q}}$ and $C_{QQ} = \frac{C_{n\bar{n}}}{C_{n\eta}} C_{Q\bar{Q}}$, respectively. The dashed (red) lines are for the case $C_{QQ} = 0$. The dotted lines indicate various meson-meson thresholds. When the quantum numbers in the subscript of the symbol for a meson-meson state are equal to the J^{PC} (or J^P) of an initial state, the decay for the initial state into that meson-meson channel through S- or D-wave is allowed. We adopt the masses estimated with the reference thresholds of (a) $\Upsilon\Upsilon$, (b) $\psi\psi$, (c) $B_c B_c$, (d) ΥB_c , (e) ψB_c , and (f) $B_c B_c$. The masses are all in units of MeV.

Basically, the behaviors for the rearrangement decays are similar to those for the $b\bar{b}\bar{b}$ and $c\bar{c}\bar{c}$ systems. The main difference lies in the C parity. The former states have definite C parities while the $c\bar{c}\bar{b}$ states not. Without the condition of C parity conservation, the 2^+ $c\bar{c}\bar{b}$ tetraquark can also decay into the $B_c^+ B_c^{*+}$ channel through D -wave.

Up to now, experiments confirm only the ground B_c meson. It means that the axial vector tetraquark decaying into $B_c B_c^*$ may not be observed in the near future. In the case that the B_c^* meson is confirmed with enough data, the search for the 1^+ $c\bar{c}\bar{b}$ (or $b\bar{b}\bar{c}$) tetraquark is also possible. However, the interactions within the diquarks are effectively repulsive and these tetraquarks should not be stable.

TABLE III: Results for the $b\bar{b}\bar{c}$ and $c\bar{b}\bar{b}$ systems in units of MeV with the approximation $C_{QQ} = C_{Q\bar{Q}}$. The masses in the fifth column are estimated with $m_b = 5052.9$ MeV and $m_c = 1724.8$ MeV. The last column lists masses estimated from the $(B_c B_c)$ threshold. The base for the $J = 0$ case is $(\phi_2\chi_3, \phi_1\chi_6)^T$.

J^P	$\langle H_{CM} \rangle$	Eigenvalue	Eigenvector	Mass	$(B_c B_c)$
2^+	39.5	39.5	1	13595	12695
1^+	4.3	4.3	1	13560	12660
0^+	$\begin{pmatrix} -13.3 & 64.7 \\ 64.7 & 32.8 \end{pmatrix}$	$\begin{bmatrix} 78.4 \\ -58.9 \end{bmatrix}$	$\begin{bmatrix} (0.58, 0.82) \\ (-0.82, 0.58) \end{bmatrix}$	$\begin{bmatrix} 13634 \\ 13496 \end{bmatrix}$	$\begin{bmatrix} 12734 \\ 12597 \end{bmatrix}$

D. The $b\bar{b}\bar{c}$ and $c\bar{b}\bar{b}$ systems

TABLE IV: Results for the $b\bar{b}\bar{c}$ and $c\bar{b}\bar{b}$ systems in units of MeV with the approximation $C_{QQ} = C_{Q\bar{Q}}$. The masses in the fifth column are estimated with $m_b = 5052.9$ MeV and $m_c = 1724.8$ MeV. The last column lists masses estimated from the (ΥB_c) threshold. The base for the $J = 1$ case is $(\phi_2\chi_2, \phi_2\chi_4, \phi_1\chi_5)^T$ and that for the $J = 0$ case is $(\phi_2\chi_3, \phi_1\chi_6)^T$.

J^P	$\langle H_{CM} \rangle$	Eigenvalue	Eigenvector	Mass	(ΥB_c)
2^+	33.1	33.1	1	16917	15806
1^+	$\begin{pmatrix} 0.0 & -1.5 & -3.2 \\ -1.5 & -18.7 & -35.1 \\ -3.2 & -35.1 & 7.2 \end{pmatrix}$	$\begin{bmatrix} -43.3 \\ 31.7 \\ 0.1 \end{bmatrix}$	$\begin{bmatrix} (-0.07, -0.82, -0.57) \\ (-0.06, -0.57, 0.82) \\ (1.00, -0.09, 0.01) \end{bmatrix}$	$\begin{bmatrix} 16840 \\ 16915 \\ 16884 \end{bmatrix}$	$\begin{bmatrix} 15729 \\ 15804 \\ 15773 \end{bmatrix}$
0^+	$\begin{pmatrix} -16.5 & 60.7 \\ 60.7 & 24.8 \end{pmatrix}$	$\begin{bmatrix} 68.3 \\ -60.0 \end{bmatrix}$	$\begin{bmatrix} (0.58, 0.81) \\ (-0.81, 0.58) \end{bmatrix}$	$\begin{bmatrix} 16952 \\ 16823 \end{bmatrix}$	$\begin{bmatrix} 15841 \\ 15713 \end{bmatrix}$

We show the results for the $b\bar{b}\bar{c}$ (and $c\bar{b}\bar{b}$) states in Table IV and the spectrum for $b\bar{b}\bar{c}$ in Fig. 1 (d). The maximum mass splitting is around 130 MeV. Comparing with the former systems, the Pauli principle works only for one diquark now, which leads to one more 1^+ tetraquark.

From Fig. 1 (d), we may easily understand the rearrangement decay behaviors for the $b\bar{b}\bar{c}$ states. The two scalar states have similar properties to the states in the former systems. For the $J = 2$ tetraquark, there is one more D -wave decay channel compared to the $c\bar{b}\bar{b}$ case because the violation of the heavy quark spin symmetry results in the non-degeneracy for the thresholds of $\eta_b B_c^*$ and ΥB_c . The interesting observation appears for the $J^P = 1^+$ states. The color-spin mixing affects the masses of the states relatively largely. The resulting observation is: the highest and the intermediate states are kinematically allowed to decay into ΥB_c and $\eta_b B_c^*$ channels, while the lowest state has no rearrangement decay channel. From the relative positions for the solid/black, dashdotted/blue, and dashed/red lines, the interactions within the diquarks are effectively attractive for the lowest 1^+ tetraquark, repulsive for the intermediate 1^+ state and also attractive for the highest 1^+ state. This feature is a result of balance between attraction/repulsion in bc , repulsion in bb , attraction between quarks and antiquarks, and channel coupling. From the effective interaction within the diquarks and the estimated mass, the lowest 1^+ state is a good candidate of stable tetraquarks. Apparently, this state looks like an excited B_c where a $b\bar{b}$ pair is excited (see discussions in Ref. [99] for other tetraquarks). By checking the CMI matrix elements, one finds that the stable state is possible mainly because of the attraction within the bc diquark. The situation is similar to the T_{cc} state ($cc\bar{q}\bar{q}$) where the attractive ud diquark contributes dominantly [101, 102].

With the observed bottomonium and B_c states, in principle, resonance structures above the $\eta_b B_c$ threshold can all be investigated. Once experiments collect enough B_c data, interesting states in the ΥB_c channel would probably be observed first.

E. The $c\bar{c}\bar{b}$ and $b\bar{c}\bar{c}$ systems

We may call such tetraquarks as ‘‘mirror’’ partners of the previous states. The color-spin structure is the same as the $b\bar{b}\bar{c}$ system but the decay feature relies on masses and may be different. We present the numerical results in Table V and draw the spectrum in Fig. 1 (e). The maximum mass splitting between the two scalar tetraquarks is around 180 MeV. Rearrangement decays for these tetraquarks are possible only when the mass is high enough.

The rearrangement decay properties for the $J = 2$ and $J = 0$ tetraquarks are similar to those for the $b\bar{b}\bar{c}$ states. For the three axial vector tetraquarks, the low mass one has only one decay channel $\eta_c B_c^*$, the intermediate one has

two $\eta_c B_c^*$ and ψB_c , and the high mass state has one more channel ψB_c^* . Contrary to the $bb\bar{b}\bar{c}$ case, the effective interaction within the diquarks in the lowest 1^+ state is repulsive, which is a result that the cc interaction is stronger than the bb interaction. If this state does exist, it should be less stable than the lowest $1^+ bb\bar{b}\bar{c}$.

Early experimental investigations on possible resonances in the $cc\bar{c}\bar{b}$ system should be through the channel ψB_c , which means that four tetraquarks, a high mass scalar, two axial vectors, and one tensor, could be observed first.

TABLE V: Results for the $cc\bar{c}\bar{b}$ and $bc\bar{c}\bar{c}$ systems in units of MeV with the approximation $C_{QQ} = C_{Q\bar{Q}}$. The masses in the fifth column are estimated with $m_b = 5052.9$ MeV and $m_c = 1724.8$ MeV. The last column lists masses estimated from the (ψB_c) threshold. The base for the $J = 1$ case is $(\phi_2\chi_2, \phi_2\chi_4, \phi_1\chi_5)^T$ and that for the $J = 0$ case is $(\phi_2\chi_3, \phi_1\chi_6)^T$.

J^P	$\langle H_{CM} \rangle$	Eigenvalue	Eigenvector	Mass	(ψB_c)
2^+	45.9	45.9	1	10273	9442
1^+	$\begin{pmatrix} 0.0 & 7.5 & 16.0 \\ 7.5 & -12.3 & -48.6 \\ 16.0 & -48.6 & 16.8 \end{pmatrix}$	54.4	$(0.16, -0.57, 0.81)$	10282	9451
		-53.1	$(0.28, -0.76, -0.59)$	10174	9343
		3.3	$(0.95, 0.32, 0.04)$	10231	9400
0^+	$\begin{pmatrix} -22.9 & 84.3 \\ 84.3 & 34.4 \end{pmatrix}$	94.7	$(0.58, 0.81)$	10322	9491
		-83.3	$(-0.81, 0.58)$	10144	9313

F. The $bc\bar{b}\bar{c}$ system

We show the results for the twelve states in Table VI. Now the maximum mass splitting (240 MeV) again occurs between the scalar tetraquarks. In estimating the tetraquark masses, there are two reference thresholds we can use, $\Upsilon\psi$ and $B_c^+ B_c^-$. As the investigations in other systems [99, 100], the former threshold leads to lighter masses that can be treated as a lower limit on the theoretical side. We here adopt the masses estimated with the latter threshold. Future measurements may answer whether the scheme is reasonable or not. Seven meson-meson channels are involved in discussing decay properties with the obtained results. We show the spectrum and these channels in Fig. 1 (f).

TABLE VI: Results for the $bc\bar{b}\bar{c}$ system in units of MeV with the approximation $C_{QQ} = C_{Q\bar{Q}}$. The masses in the fifth column are estimated with $m_b = 5052.9$ MeV and $m_c = 1724.8$ MeV. The last two columns list masses estimated from the $(\Upsilon\psi)$ and the $(B_c B_c)$ thresholds. The bases for the $J = 2$ and $J = 0$ cases are $(\phi_1\chi_1, \phi_2\chi_1)^T$ and $(\phi_1\chi_3, \phi_1\chi_6, \phi_2\chi_3, \phi_2\chi_6)^T$, respectively. If $J = 1$, the bases for the cases $C = +$ and $C = -$ are $([\phi\chi]_+^{66}, [\phi\chi]_+^{33})^T$ and $(\phi_1\chi_2, \phi_2\chi_2, [\phi\chi]_-^{66}, [\phi\chi]_-^{33})^T$, respectively.

J^{PC}	$\langle H_{CM} \rangle$	Eigenvalue	Eigenvector	Mass	$(\Upsilon\psi)$	$(B_c B_c)$
2^{++}	$\begin{pmatrix} 40.5 & 4.5 \\ 4.5 & 37.3 \end{pmatrix}$	43.7	$(-0.82, -0.58)$	13599	12557	12700
		34.1	$(0.58, -0.82)$	13590	12548	12690
0^{++}	$\begin{pmatrix} -107.5 & -9.2 & 9.1 & 72.5 \\ -9.2 & 26.4 & 72.5 & 0.0 \\ 9.1 & 72.5 & -21.9 & -3.7 \\ 72.5 & 0.0 & -3.7 & -52.8 \end{pmatrix}$	-159.4	$(0.82, 0.08, -0.11, -0.56)$	13396	12354	12496
		78.8	$(-0.02, 0.81, 0.58, -0.03)$	13634	12592	12735
		-72.7	$(0.08, -0.58, 0.80, -0.14)$	13483	12441	12583
		-2.4	$(0.57, -0.01, 0.08, 0.82)$	13553	12511	12653
1^{++}	$\begin{pmatrix} 14.1 & -41.9 \\ -41.9 & -15.5 \end{pmatrix}$	-45.1	$(0.58, 0.82)$	13510	12468	12611
		43.7	$(-0.82, 0.58)$	13599	12557	12700
1^{+-}	$\begin{pmatrix} -58.1 & 4.5 & -16.0 & 13.6 \\ 4.5 & -2.1 & 13.6 & -6.4 \\ -16.0 & 13.6 & 3.5 & 41.9 \\ 13.6 & -6.4 & 41.9 & -19.7 \end{pmatrix}$	-77.7	$(-0.73, 0.16, -0.43, 0.50)$	13478	12436	12578
		36.8	$(0.05, -0.18, -0.80, -0.56)$	13592	12550	12693
		-35.5	$(0.68, 0.19, -0.41, 0.58)$	13520	12478	12620
		-0.1	$(0.00, 0.95, 0.00, -0.31)$	13555	12513	12656

Since there is no constraint from the Pauli principle, the obtained spectrum is more complicated than other systems. All the masses are above the lowest meson-meson threshold and they should decay. At least four rearrangement decay channels for the 2^{++} states are D -wave channels. If the 2^{++} state is above the $B_c^{*+} B_c^{*-}$ threshold, the S -wave channel is also opened. For the channels with $J^{PC} = 1^{++}$, the S -wave $\Upsilon\psi$ is the lowest one. The decays of the two 1^{++} tetraquarks into this channel are both allowed. For the four 1^{+-} states, they all have S -wave decay channels $\Upsilon\eta_c$ and $\eta_b\psi$. There is at least one allowed rearrangement decay channel, $\eta_b\eta_c$, for the four 0^{++} tetraquarks through S -wave interaction. It seems that all the $bc\bar{b}\bar{c}$ tetraquarks are not stable.

On the other hand, one finds that the interactions within the diquarks in several states are effectively attractive. This indicates that in the competition between the interactions of QQ and $Q\bar{Q}$, the stronger attraction in QQ may make such states relatively stable. The lowest 0^{++} state satisfies this condition and it has only one rearrangement decay channel. Maybe this state has relatively narrow width. The lower 1^{++} state has also similar features. Its S -wave decay channel is $\Upsilon\psi$ and this tetraquark is worthwhile study. Because of the existence of possible tetraquark states, searching for exotic phenomena with $\Upsilon\psi$ may help us to understand the strong interactions between heavy quarks. In Ref. [82], Richard et al also observe that bound $b\bar{c}\bar{b}$ might be more favorable than $b\bar{b}\bar{b}$ and $c\bar{c}\bar{c}$.

IV. DISCUSSIONS AND SUMMARY

In the chiral quark model, the interaction between the light quarks may also arise from the exchange of Goldstone bosons. For the pure heavy systems, such an interaction is absent. Therefore, one needs to consider only the gluon-exchange interaction for the present $Q_1Q_2\bar{Q}_3\bar{Q}_4$ systems. As a short range force, it helps to form compact tetraquarks rather than meson-meson molecules if bound four-quark states do exist. We have studied the mass splittings of these tetraquark states with the simple color-magnetic interaction in this work.

For the case ($b\bar{c}\bar{b}$ system) without constraint from the Pauli principle, the number of color-singlet tetraquarks with 6_c diquark is equal to that with $\bar{3}_c$ diquark. The two color structures may couple through the color-magnetic interaction. From Table VI, the coupling for the 2^{++} states is weak while that for other states is stronger. For the case with constraint from the Pauli principle, the number of color-singlet tetraquarks with the $\bar{3}_c$ diquark is bigger than that with 6_c diquark. Their mixing is generally significant (see Tables II-V). In both cases, the tetraquarks with 6_c diquark do not exist independently.

After the configuration mixing effects are considered, both the heaviest tetraquark and the lightest tetraquark for a system are the scalar states. The maximum mass differences are around 120 MeV, 220 MeV, 140 MeV, 130 MeV, 180 MeV, and 240 MeV for the $b\bar{b}\bar{c}$, $c\bar{c}\bar{c}$, $b\bar{b}\bar{c}$, $b\bar{b}\bar{c}$, $c\bar{c}\bar{b}$, and $b\bar{c}\bar{b}$ systems, respectively. Other states fall in these mass difference ranges. Whether the tetraquarks decay through quark rearrangements and how many thresholds fall in these ranges rely on the tetraquark masses. For comparison, we estimate the masses with two approaches. We mainly discuss the tetraquark properties by using the masses estimated with the threshold of some meson-meson state, the second approach.

Since the parameter C_{QQ} cannot be extracted with experimental data, we use two approximations to determine them. We also show results in the extreme case $C_{QQ} = 0$. From the comparison study, we find that (1) the estimated tetraquark masses are affected slightly and (2) the plotted spectra can be easily used to judge which states contain effectively attractive diquarks. From the viewpoint that narrow compact tetraquarks should have attractive diquarks inside them and have as few S -wave decay channels as possible, we find that most $QQ\bar{Q}\bar{Q}$ states are not stable. However, a stable 1^+ $b\bar{b}\bar{c}$ state is observed and relatively narrow $b\bar{c}\bar{b}$ tetraquarks are also possible in the present model investigations. The latter system is also proposed to have bound states in Ref. [82]. Although the model we use is simple and it does not involve dynamics, the basic features of the obtained spectra might be somehow reasonable. To give more reliable results, the present model needs to be improved.

If experiments could observe one resonant state in the channel of two heavy quarkonia, its nature as a tetraquark is favored. More tetraquarks should also exist and searches for them are strongly called for. We hope the decay channels discussed in this paper are helpful for the experimental search.

To summarize, we have explored the mass splittings between the $Q_1Q_2\bar{Q}_3\bar{Q}_4$ tetraquarks with a simple model. The mixing between different color structures are considered. We have estimated their masses with two approaches and discussed possible rearrangement decay channels. Stable or narrow tetraquarks in the $b\bar{b}\bar{c}$ and $b\bar{c}\bar{b}$ systems are worthwhile study in heavy meson-meson channels such as $\Upsilon\psi$. Hopefully, the exotic tetraquark states composed of four heavy quarks may be observed at LHCb in the future.

Acknowledgments

YRL thanks F. Huang (UCAS, Beijing) for a discussion and people at Tokyo Institute of Technology for their hospitality where the updated draft was finalized. This project is supported by National Natural Science Foundation of China under Grants No. 11775132, No. 11222547, No. 11175073, No. 11261130311 and 973 program. XL is also

supported by the National Program for Support of Top-notch Young Professionals.

-
- [1] R. L. Jaffe, “Multi-Quark Hadrons. I. The Phenomenology of $Q^2\bar{Q}^2$ Mesons,” *Phys. Rev. D* **15**, 267 (1977).
- [2] R. L. Jaffe, “Multi-Quark Hadrons. II. Methods,” *Phys. Rev. D* **15**, 281 (1977).
- [3] R. L. Jaffe, “Perhaps a Stable Dihyperon,” *Phys. Rev. Lett.* **38**, 195 (1977) Erratum: [*Phys. Rev. Lett.* **38**, 617 (1977)].
- [4] E. S. Swanson, “The New heavy mesons: A Status report,” *Phys. Rept.* **429**, 243 (2006) [hep-ph/0601110].
- [5] S. L. Zhu, “New hadron states,” *Int. J. Mod. Phys. E* **17**, 283 (2008) [hep-ph/0703225].
- [6] M. B. Voloshin, “Charmonium,” *Prog. Part. Nucl. Phys.* **61**, 455 (2008) [arXiv:0711.4556 [hep-ph]].
- [7] N. Drenska, R. Faccini, F. Piccinini, A. Polosa, F. Renga and C. Sabelli, “New Hadronic Spectroscopy,” *Riv. Nuovo Cim.* **33**, 633 (2010) [arXiv:1006.2741 [hep-ph]].
- [8] H. X. Chen, W. Chen, X. Liu and S. L. Zhu, “The hidden-charm pentaquark and tetraquark states,” arXiv:1601.02092 [hep-ph].
- [9] R. Chen, X. Liu and S. L. Zhu, “Hidden-charm molecular pentaquarks and their charm-strange partners,” arXiv:1601.03233 [hep-ph].
- [10] A. Hosaka, T. Iijima, K. Miyabayashi, Y. Sakai and S. Yasui, “Exotic Hadrons with Heavy Flavors -X, Y, Z and Related States,” arXiv:1603.09229 [hep-ph].
- [11] A. Ali, J. S. Lange and S. Stone, “Exotics: Heavy Pentaquarks and Tetraquarks,” *Prog. Part. Nucl. Phys.* **97**, 123 (2017) [arXiv:1706.00610 [hep-ph]].
- [12] S. K. Choi *et al.* [Belle Collaboration], “Observation of a resonance-like structure in the $\pi^\pm\psi'$ mass distribution in exclusive $B \rightarrow K\pi^\pm\psi'$ decays,” *Phys. Rev. Lett.* **100**, 142001 (2008) [arXiv:0708.1790 [hep-ex]].
- [13] R. Mizuk *et al.* [Belle Collaboration], “Dalitz analysis of $B \rightarrow K\pi^+\psi'$ decays and the $Z(4430)^+$,” *Phys. Rev. D* **80**, 031104 (2009) [arXiv:0905.2869 [hep-ex]].
- [14] K. Chilikin *et al.* [Belle Collaboration], “Observation of a new charged charmoniumlike state in $\bar{B}^0 \rightarrow J/\psi K^-\pi^+$ decays,” *Phys. Rev. D* **90**, no. 11, 112009 (2014) [arXiv:1408.6457 [hep-ex]].
- [15] R. Aaij *et al.* [LHCb Collaboration], “Observation of the resonant character of the $Z(4430)^-$ state,” *Phys. Rev. Lett.* **112**, 222002 (2014) [arXiv:1404.1903 [hep-ex]].
- [16] R. Mizuk *et al.* [Belle Collaboration], “Observation of two resonance-like structures in the $\pi^+\chi_{c1}$ mass distribution in exclusive $\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}$ decays,” *Phys. Rev. D* **78**, 072004 (2008) [arXiv:0806.4098 [hep-ex]].
- [17] M. Ablikim *et al.* [BESIII Collaboration], “Observation of a Charged Charmoniumlike Structure in $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ at $\sqrt{s}=4.26$ GeV,” *Phys. Rev. Lett.* **110**, 252001 (2013) [arXiv:1303.5949 [hep-ex]].
- [18] Z. Q. Liu *et al.* [Belle Collaboration], “Study of $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ and Observation of a Charged Charmoniumlike State at Belle,” *Phys. Rev. Lett.* **110**, 252002 (2013) [arXiv:1304.0121 [hep-ex]].
- [19] T. Xiao, S. Dobbs, A. Tomaradze and K. K. Seth, “Observation of the Charged Hadron $Z_c^\pm(3900)$ and Evidence for the Neutral $Z_c^0(3900)$ in $e^+e^- \rightarrow \pi\pi J/\psi$ at $\sqrt{s}=4170$ MeV,” *Phys. Lett. B* **727**, 366 (2013) [arXiv:1304.3036 [hep-ex]].
- [20] M. Ablikim *et al.* [BESIII Collaboration], “Observation of $Z_c(3900)^0$ in $e^+e^- \rightarrow \pi^0\pi^0 J/\psi$,” *Phys. Rev. Lett.* **115**, 112003 (2015) [arXiv:1506.06018 [hep-ex]].
- [21] M. Ablikim *et al.* [BESIII Collaboration], “Observation of a charged $(D\bar{D}^*)^\pm$ mass peak in $e^+e^- \rightarrow \pi D\bar{D}^*$ at $\sqrt{s}=4.26$ GeV,” *Phys. Rev. Lett.* **112**, 022001 (2014) [arXiv:1310.1163 [hep-ex]].
- [22] M. Ablikim *et al.* [BESIII Collaboration], “Confirmation of a charged charmoniumlike state $Z_c(3885)^\mp$ in $e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$ with double D tag,” *Phys. Rev. D* **92**, 092006 (2015) [arXiv:1509.01398 [hep-ex]].
- [23] M. Ablikim *et al.* [BESIII Collaboration], “Observation of a Neutral Structure near the $D\bar{D}^*$ Mass Threshold in $e^+e^- \rightarrow (D\bar{D}^*)^0\pi^0$ at $\sqrt{s}=4.226$ and 4.257 GeV,” *Phys. Rev. Lett.* **115**, 222002 (2015) [arXiv:1509.05620 [hep-ex]].
- [24] M. Ablikim *et al.* [BESIII Collaboration], “Observation of a Charged Charmoniumlike Structure $Z_c(4020)$ and Search for the $Z_c(3900)$ in $e^+e^- \rightarrow \pi^+\pi^-h_c$,” *Phys. Rev. Lett.* **111**, 242001 (2013) [arXiv:1309.1896 [hep-ex]].
- [25] M. Ablikim *et al.* [BESIII Collaboration], “Observation of $e^+e^- \rightarrow \pi^0\pi^0h_c$ and a Neutral Charmoniumlike Structure $Z_c(4020)^0$,” *Phys. Rev. Lett.* **113**, 212002 (2014) [arXiv:1409.6577 [hep-ex]].
- [26] M. Ablikim *et al.* [BESIII Collaboration], “Observation of a charged charmoniumlike structure in $e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$ at $\sqrt{s}=4.26$ GeV,” *Phys. Rev. Lett.* **112**, 132001 (2014) [arXiv:1308.2760 [hep-ex]].
- [27] M. Ablikim *et al.* [BESIII Collaboration], “Observation of a neutral charmoniumlike state $Z_c(4025)^0$ in $e^+e^- \rightarrow (D^*\bar{D}^*)^0\pi^0$,” *Phys. Rev. Lett.* **115**, 182002 (2015) [arXiv:1507.02404 [hep-ex]].
- [28] A. Bondar *et al.* [Belle Collaboration], “Observation of two charged bottomonium-like resonances in $\Upsilon(5S)$ decays,” *Phys. Rev. Lett.* **108**, 122001 (2012) [arXiv:1110.2251 [hep-ex]].
- [29] J. J. Wu, R. Molina, E. Oset and B. S. Zou, Prediction of narrow N^* and Λ^* resonances with hidden charm above 4 GeV, *Phys. Rev. Lett.* **105**, 232001 (2010), [arXiv:1007.0573 [nucl-th]].
- [30] J. J. Wu, R. Molina, E. Oset and B. S. Zou, Dynamically generated N^* and Λ^* resonances in the hidden charm sector around 4.3 GeV, *Phys. Rev. C* **84**, 015202 (2011), [arXiv:1011.2399 [nucl-th]].
- [31] Z. C. Yang, Z. F. Sun, J. He, X. Liu and S. L. Zhu, The possible hidden-charm molecular baryons composed of anti-charmed meson and charmed baryon, *Chin. Phys. C* **36**, 6 (2012), [arXiv:1105.2901 [hep-ph]].
- [32] W. L. Wang, F. Huang, Z. Y. Zhang and B. S. Zou, $\Sigma_c\bar{D}$ and $\Lambda_c\bar{D}$ states in a chiral quark model, *Phys. Rev. C* **84**, 015203 (2011), [arXiv:1101.0453 [nucl-th]].

- [33] S. G. Yuan, K. W. Wei, J. He, H. S. Xu and B. S. Zou, Study of $qqqc\bar{c}$ five quark system with three kinds of quark-quark hyperfine interaction, *Eur. Phys. J. A* **48**, 61 (2012), [arXiv:1201.0807 [nucl-th]].
- [34] J. J. Wu, T.-S. H. Lee and B. S. Zou, Nucleon Resonances with Hidden Charm in Coupled-Channel Models, *Phys. Rev. C* **85**, 044002 (2012), [arXiv:1202.1036 [nucl-th]].
- [35] C. Garcia-Recio, J. Nieves, O. Romanets, L. L. Salcedo and L. Tolos, Hidden charm N and Δ resonances with heavy-quark symmetry, *Phys. Rev. D* **87**, 074034 (2013), [arXiv:1302.6938 [hep-ph]].
- [36] C. W. Xiao, J. Nieves and E. Oset, Combining heavy quark spin and local hidden gauge symmetries in the dynamical generation of hidden charm baryons, *Phys. Rev. D* **88**, 056012 (2013), [arXiv:1304.5368 [hep-ph]].
- [37] T. Uchino, W. H. Liang and E. Oset, “Baryon states with hidden charm in the extended local hidden gauge approach,” *Eur. Phys. J. A* **52**, 43 (2016) [arXiv:1504.05726 [hep-ph]].
- [38] R. Aaij *et al.* [LHCb Collaboration], “Model-independent evidence for $J/\psi p$ contributions to $\Lambda_b^0 \rightarrow J/\psi p K^-$ decays,” arXiv:1604.05708 [hep-ex].
- [39] V. M. Abazov *et al.* [DØ Collaboration], “Observation of a New $B_s^0\pi^\pm$ State,” [arXiv:1602.07588 [hep-ex]].
- [40] S. S. Agaev, K. Azizi and H. Sundu, “Mass and decay constant of the newly observed exotic $X(5568)$ state,” arXiv:1602.08642 [hep-ph].
- [41] Z. G. Wang, “Analysis of the $X(5568)$ as scalar tetraquark state in the diquark-antidiquark model with QCD sum rules,” arXiv:1602.08711 [hep-ph].
- [42] W. Wang and R. Zhu, “Can $X(5568)$ be a tetraquark state?,” arXiv:1602.08806 [hep-ph].
- [43] W. Chen, H. X. Chen, X. Liu, T. G. Steele and S. L. Zhu, “Investigation of the $X(5568)$ as a fully open-flavor $sub\bar{d}$ tetraquark state,” arXiv:1602.08916 [hep-ph].
- [44] C. M. Zanetti, M. Nielsen and K. P. Khemchandani, “A QCD sum rule study for a charged bottom-strange scalar meson,” arXiv:1602.09041 [hep-ph].
- [45] S. S. Agaev, K. Azizi and H. Sundu, “Width of the exotic $X_b(5568)$ state through its strong decay to $B_s^0\pi^+$,” arXiv:1603.00290 [hep-ph].
- [46] Y. R. Liu, X. Liu and S. L. Zhu, “ $X(5568)$ and its partner states,” *Phys. Rev. D* **93**, 074023 (2016) [arXiv:1603.01131 [hep-ph]].
- [47] S. S. Agaev, K. Azizi and H. Sundu, “Charmed partner of the exotic $X(5568)$ state and its properties,” arXiv:1603.01471 [hep-ph].
- [48] J. M. Dias, K. P. Khemchandani, A. M. Torres, M. Nielsen and C. M. Zanetti, “A QCD sum rule calculation of the $X^\pm(5568) \rightarrow B_s^0\pi^\pm$ decay width,” arXiv:1603.02249 [hep-ph].
- [49] Z. G. Wang, “Analysis of the strong decay $X(5568) \rightarrow B_s^0\pi^+$ with QCD sum rules,” arXiv:1603.02498 [hep-ph].
- [50] X. G. He and P. Ko, “Flavor $SU(3)$ Symmetry for the $X(5568)$ State,” arXiv:1603.02915 [hep-ph].
- [51] F. Stancu, “ $X(5568)$ as a $sub\bar{d}$ Tetraquark in a Simple Quark Model,” arXiv:1603.03322 [hep-ph].
- [52] L. Tang and C. F. Qiao, “ $X(5568)$ as Tetraquark State with Open Flavors and its Charmed Partners,” arXiv:1603.04761 [hep-ph].
- [53] S. S. Agaev, K. Azizi and H. Sundu, “Exploring $X(5568)$ as a meson molecule,” arXiv:1603.02708 [hep-ph].
- [54] The LHCb Collaboration [LHCb Collaboration], “Search for structure in the $B_s^0\pi^\pm$ invariant mass spectrum,” LHCb-CONF-2016-004, CERN-LHCb-CONF-2016-004.
- [55] T. J. Burns and E. S. Swanson, “Interpreting the $X(5568)$,” arXiv:1603.04366 [hep-ph].
- [56] F. K. Guo, U. G. Meissner and B. S. Zou, “How the $X(5568)$ challenges our understanding of QCD,” arXiv:1603.06316 [hep-ph].
- [57] Q. F. L and Y. B. Dong, “Masses of open charm and bottom tetraquark states in relativized quark model,” arXiv:1603.06417 [hep-ph].
- [58] A. Esposito, A. Pilloni and A. D. Polosa, “Hybridized Tetraquarks,” arXiv:1603.07667 [hep-ph].
- [59] M. Albaladejo, J. Nieves, E. Oset, Z. F. Sun and X. Liu, “Can $X(5568)$ be described as a $B_s\pi$, $B\bar{K}$ resonant state?,” arXiv:1603.09230 [hep-ph].
- [60] A. Ali, L. Maiani, A. D. Polosa and V. Riquer, “ B_c^\pm decays into tetraquarks,” arXiv:1604.01731 [hep-ph].
- [61] R. Albuquerque, S. Narison, A. Rabemananjara and D. Rabetiarivony, “Nature of the $X(5568)$: a critical Laplace sum rule analysis at N2LO,” arXiv:1604.05566 [hep-ph].
- [62] X. Chen and J. Ping, “Is the exotic $X(5568)$ a bound state?,” arXiv:1604.05651 [hep-ph].
- [63] Y. Jin and S. Y. Li, “Large Production Rate of New $B_s^0\pi^\pm$ and $D_s^\pm\pi^\pm$ States in High Energy Multi-Production Process,” arXiv:1603.03250 [hep-ph].
- [64] Y. Iwasaki, “A Possible Model for New Resonances-Exotics and Hidden Charm,” *Prog. Theor. Phys.* **54**, 492 (1975).
- [65] J. I. Ballot and J. M. Richard, “Four Quark States In Additive Potentials,” *Phys. Lett. B* **123**, 449 (1983).
- [66] K. T. Chao, “The $(cc)(\bar{c}\bar{c})$ (Diquark - anti-Diquark) States in e^+e^- Annihilation,” *Z. Phys. C* **7**, 317 (1981).
- [67] L. Heller and J. A. Tjon, “On Bound States of Heavy $Q^2\bar{Q}^2$ Systems,” *Phys. Rev. D* **32**, 755 (1985).
- [68] J. P. Ader, J. M. Richard and P. Taxil, “Do Narrow Heavy Multi-Quark States Exist?,” *Phys. Rev. D* **25**, 2370 (1982).
- [69] B. Silvestre-Brac and C. Semay, “Systematics of $L = 0$ $q^2\bar{q}^2$ systems,” *Z. Phys. C* **57**, 273 (1993).
- [70] R. J. Lloyd and J. P. Vary, “All charm tetraquarks,” *Phys. Rev. D* **70**, 014009 (2004) [hep-ph/0311179].
- [71] N. Barnea, J. Vijande and A. Valcarce, “Four-quark spectroscopy within the hyperspherical formalism,” *Phys. Rev. D* **73**, 054004 (2006) [hep-ph/0604010].
- [72] B. Silvestre-Brac, “Systematics of $Q^2\bar{Q}^2$ systems with a chromomagnetic interaction,” *Phys. Rev. D* **46**, 2179 (1992).
- [73] T. W. Chiu *et al.* [TWQCD Collaboration], “ $Y(4260)$ on the lattice,” *Phys. Rev. D* **73**, 094510 (2006) [hep-lat/0512029].
- [74] C. Hughes, E. Eichten and C. T. H. Davies, “The Search for Beauty-fully Bound Tetraquarks Using Lattice Non-

- Relativistic QCD,” [arXiv:1710.03236 [hep-lat]].
- [75] Y. Bai, S. Lu and J. Osborne, “Beauty-full Tetraquarks,” arXiv:1612.00012 [hep-ph].
- [76] M. N. Anwar, J. Ferretti, F. K. Guo, E. Santopinto and B. S. Zou, “Spectroscopy and decays of the fully-heavy tetraquarks,” arXiv:1710.02540 [hep-ph].
- [77] Z. G. Wang, “Analysis of the $QQ\bar{Q}\bar{Q}$ tetraquark states with QCD sum rules,” Eur. Phys. J. C **77**, no. 7, 432 (2017) [arXiv:1701.04285 [hep-ph]].
- [78] V. R. Debastiani and F. S. Navarra, “A non-relativistic model for the $[cc][\bar{c}\bar{c}]$ tetraquark,” arXiv:1706.07553 [hep-ph].
- [79] M. Karliner, S. Nussinov and J. L. Rosner, “ $QQ\bar{Q}\bar{Q}$ states: masses, production, and decays,” Phys. Rev. D **95**, no. 3, 034011 (2017) [arXiv:1611.00348 [hep-ph]].
- [80] E. Eichten and Z. Liu, “Would a Deeply Bound $b\bar{b}b\bar{b}$ Tetraquark Meson be Observed at the LHC?,” arXiv:1709.09605 [hep-ph].
- [81] R. Vega-Morales, “Diagnosing $2\ell + \Upsilon$ resonances at the LHC,” arXiv:1710.02738 [hep-ph].
- [82] J. M. Richard, A. Valcarce and J. Vijande, “String dynamics and metastability of all-heavy tetraquarks,” Phys. Rev. D **95**, no. 5, 054019 (2017) [arXiv:1703.00783 [hep-ph]].
- [83] W. Chen, H. X. Chen, X. Liu, T. G. Steele and S. L. Zhu, “Hunting for exotic doubly hidden-charm/bottom tetraquark states,” Phys. Lett. B **773**, 247 (2017) [arXiv:1605.01647 [hep-ph]].
- [84] P. Bicudo, M. Cardoso, O. Oliveira and P. J. Silva, “Lattice QCD static potentials of the meson-meson and tetraquark systems computed with both quenched and full QCD,” arXiv:1702.07789 [hep-lat].
- [85] A. De Rujula, H. Georgi and S. L. Glashow, “Hadron Masses in a Gauge Theory,” Phys. Rev. D **12**, 147 (1975).
- [86] H. Hogaasen and P. Sorba, “The Colour triplet $qq\bar{q}$ cluster and pentaquark models,” Mod. Phys. Lett. A **19**, 2403 (2004) [hep-ph/0406078].
- [87] F. Buccella, H. Hogaasen, J. M. Richard and P. Sorba, “Chromomagnetism, flavour symmetry breaking and S-wave tetraquarks,” Eur. Phys. J. C **49**, 743 (2007) [hep-ph/0608001].
- [88] H. Hogaasen, J. M. Richard and P. Sorba, “A Chromomagnetic mechanism for the $X(3872)$ resonance,” Phys. Rev. D **73**, 054013 (2006) [hep-ph/0511039].
- [89] Y. R. Liu and M. Oka, “ $\Lambda_c N$ bound states revisited,” Phys. Rev. D **85**, 014015 (2012) [arXiv:1103.4624 [hep-ph]].
- [90] Y. R. Liu, “Heavy quark spin selection rules in meson-antimeson states,” Phys. Rev. D **88**, 074008 (2013) [arXiv:1304.7467 [hep-ph]].
- [91] C. Patrignani *et al.* [Particle Data Group], “Review of Particle Physics,” Chin. Phys. C **40**, no. 10, 100001 (2016).
- [92] S. Godfrey and N. Isgur, Mesons in a Relativized Quark Model with Chromodynamics, Phys. Rev. D **32**, 189 (1985).
- [93] R. Aaij *et al.* [LHCb Collaboration], “Observation of the doubly charmed baryon Ξ_{cc}^{++} ,” Phys. Rev. Lett. **119**, no. 11, 112001 (2017) [arXiv:1707.01621 [hep-ex]].
- [94] M. Mattson *et al.* [SELEX Collaboration], “First observation of the doubly charmed baryon Ξ_{cc}^+ ,” Phys. Rev. Lett. **89**, 112001 (2002) [hep-ex/0208014].
- [95] A. Ocherashvili *et al.* [SELEX Collaboration], “Confirmation of the double charm baryon $\Xi_{cc}^+(3520)$ via its decay to pD^+K^- ,” Phys. Lett. B **628**, 18 (2005) [hep-ex/0406033].
- [96] S. H. Lee, S. Yasui, W. Liu and C. M. Ko, “Charmed exotics in Heavy Ion Collisions,” Eur. Phys. J. C **54**, 259 (2008) [arXiv:0707.1747 [hep-ph]].
- [97] S. H. Lee and S. Yasui, “Stable multi-quark states with heavy quarks in a diquark model,” Eur. Phys. J. C **64**, 283 (2009) [arXiv:0901.2977 [hep-ph]].
- [98] J. Wu, Y. R. Liu, K. Chen, X. Liu and S. L. Zhu, “ $X(4140)$, $X(4270)$, $X(4500)$ and $X(4700)$ and their $cs\bar{c}\bar{s}$ tetraquark partners,” Phys. Rev. D **94**, no. 9, 094031 (2016) [arXiv:1608.07900 [hep-ph]].
- [99] K. Chen, X. Liu, J. Wu, Y. R. Liu and S. L. Zhu, “Triply heavy tetraquark states with the $QQ\bar{Q}\bar{q}$ configuration,” Eur. Phys. J. A **53**, no. 1, 5 (2017) [arXiv:1609.06117 [hep-ph]].
- [100] J. Wu, Y. R. Liu, K. Chen, X. Liu and S. L. Zhu, “Hidden-charm pentaquarks and their hidden-bottom and B_c -like partner states,” Phys. Rev. D **95**, no. 3, 034002 (2017) [arXiv:1701.03873 [hep-ph]].
- [101] T. Hyodo, Y. R. Liu, M. Oka, K. Sudoh and S. Yasui, Production of doubly charmed tetraquarks with exotic color configurations in electron-positron collisions, Phys. Lett. B **721**, 56 (2013), [arXiv:1209.6207 [hep-ph]].
- [102] T. Hyodo, Y. R. Liu, M. Oka and S. Yasui, “Spectroscopy and production of doubly charmed tetraquarks,” arXiv:1708.05169 [hep-ph].
- [103] S. Q. Luo, K. Chen, X. Liu, Y. R. Liu and S. L. Zhu, “Exotic tetraquark states with the $qq\bar{Q}\bar{Q}$ configuration,” Eur. Phys. J. C **77**, no. 10, 709 (2017) [arXiv:1707.01180 [hep-ph]].