

BOOSTED-BOTTOM JET TAGGING AND BSM SEARCHES

ZACK SULLIVAN

*Department of Physics, Illinois Institute of Technology,
Chicago, Illinois 60616-3793, USA*



I present a new scheme for tagging boosted heavy flavor jets called “ μ_x tagging” and its application to TeV-scale physics beyond the Standard Model. Using muons from B hadron decay to define a particular combination “ x ” of angular information, and jet substructure variables, we identify a clean ($\epsilon_{\text{fake}}/\epsilon_b \sim 1/100$) good efficiency ($\epsilon_b = 14\%$) tag. I demonstrate the usefulness of this new scheme by showing the reach for discovery of leptophobic $Z' \rightarrow b\bar{b}$ and $tH^\pm \rightarrow t\bar{t}b$.

1 Introduction

As searches for new particles at the CERN Large Hadron Collider (LHC) shift to TeV-scale energies, observation of their decays into jets becomes challenging. Dijet resonances are typically smaller than the QCD background unless top or bottom tags are applied. Unfortunately, current b -tagging efficiencies degrade (28–15%) around 1–2 TeV for light-jet fake rates of 1–2%¹ (producing low purity $\epsilon_{\text{fake}}/\epsilon_b \sim 1/10$). At this conference, I presented a new method for flavor tagging at TeV-scale energies called “ μ_x boosted-bottom-jet tagging.”² This method is derived from kinematic first principles, and provides a 14% efficiency for b -tagging, with a factor of 10 improvement in fake rejection over existing tags ($\epsilon_{\text{fake}}/\epsilon_b \sim 1/100$). In Sec. 2 I summarize the μ_x definition and cuts, and plot its transverse momentum- and pseudorapidity-dependent efficiencies. In Sec. 3 I briefly describe the reach provided by μ_x boosted- b tagging in an analysis for discovery of a leptophobic $Z' \rightarrow b\bar{b}$. Following the recent provocative proposals to measure $H^\pm \rightarrow tb$ in tbH^\pm -associated production at the LHC, I use the μ_x tag in Sec. 4 to provide a realistic estimate of the reach at a 14 TeV machine. I summarize our results in Sec. 5.

2 μ_x boosted- b tag

In Ref. 2 we introduced the μ_x boosted- b tag, a high purity b tag for use with boosted jets ($p_T > 500$ GeV) based on the kinematics of semi-muonic B hadron decay and jet substructure. At large momentum, the boost γ_B of the B hadron compresses its decay products into a narrow

subject at high energy. We define a “smart” angular lab frame observable

$$x \equiv \gamma_B \tan \theta_{\text{lab}}, \quad (1)$$

where θ_{lab} is the angle between the muon and the direction of the B hadron in the jet. For sufficiently boosted B hadrons ($\gamma_B \geq 3$) the lab frame distribution of the muon count N vs. x follows a universal shape. We define the μ_x boosted- b tag by first demanding $x < 3$ to capture 90% of muons from the B decays. Then we demand

$$f_{\text{subject}} \equiv \frac{p_{T\text{subject}}}{p_{T\text{jet}}} \geq 0.5, \quad (2)$$

to account for the observation that the hard fragmentation function for b quarks leads to the B hadron subject carrying a large fraction f_{subject} of the total jet momentum.

Both cuts in the μ_x tag depend on identification of the subject containing the B hadron. While exact identification of the B is not possible, an effective proxy can be found by taking standard anti- k_T clustered jets with $R = 0.4$, and reclustering the muon and calorimeter towers using a smaller $R = 0.04$. Following the detailed reconstruction algorithm of Ref. 2, we combine an identified probable charm subject remnant, with double the muon momentum (as a boosted neutrino proxy) to provide the input to γ_B and $p_{T\text{subject}}$. A custom μ_x tagging module `MuXboostedBTagging` for DELPHES¹ is available on GitHub³.

In Fig. 1 we see μ_x tagging efficiencies as a function of p_T and η for bottom jets, charm jets, light-light jets (where the muon came from a light-flavor hadron), and light-heavy jets (where a gluon split to $b\bar{b}/c\bar{c}$ — producing heavy-flavor hadrons in the final state). The kinematic nature of the tagging variables leads to nearly flat p_T efficiencies when $p_T > 500$ GeV. The η distribution is also flat except for B hadrons from gluon splitting. This leads to the intriguing possibility that the $g \rightarrow b\bar{b}$ contribution to jets in the Monte Carlo could be calibrated using the rapidity dependence of these highly-boosted jets.




Figure 1 – μ_x tagging efficiency vs. (left) jet p_T and (right) η_{jet} . Solid (dashed) lines include $\mu = 0$ (40) pileup events.

3 Leptophobic $Z' \rightarrow b\bar{b}$ at the LHC

New neutral vectors, generically called Z' bosons, appear in many BSM models. In cases where the decay to leptons is suppressed, we look to tag heavy flavors to overcome the QCD dijet background. We examine the reach² at a 13 TeV LHC for a leptophobic Z' decaying to $b\bar{b}$ or $c\bar{c}$ using a $U(1)'_B$ Lagrange density⁴

$$\mathcal{L} = \frac{g_B}{6} Z'_{B\mu} \bar{q} \gamma^\mu q, \quad (3)$$

with flavor-independent coupling to quarks.

The signal and backgrounds are simulated using a MLM-matched MadEvent sample⁵ and CT14ll0 PDFs⁶ fed through PYTHIA 8⁷ into DELPHES. We demand one or two μ_x tags, $|\eta_j| < 2.7$, and $\Delta\eta_{jj} < 1.5$. We reconstruct a dijet mass out of the two leading- p_T jets, and look for a resonance in the mass window $[0.85, 1.25] \times M_{Z'_B}$.

We see the reach for 5σ discovery of this leptophobic Z' in Fig. 2 for a two-tag, and one-tag inclusive sample,⁸ compared to current exclusion limits from Ref. 4. In 100 fb^{-1} of integrated luminosity at 13 TeV, a two b -tag analysis could discover a Z' of 3 TeV if the universal coupling $g_B \sim 2.5$. The single-tag inclusive search is even more effective — allowing for discovery up to 1 TeV above mass limits from Run I. In the absence of a discovery, the one-tag search would set a 95% C.L. exclusion that can access g_B couplings a factor of 2 smaller than current limits, and masses up to 2 TeV higher.




Figure 2 – 5σ discovery reach for a leptophobic Z' with universal coupling and one or two boosted- b tags at a 13 TeV LHC compared to exclusion limits from Ref. 4. Also shown is the 95% C.L. exclusion reach of the one-tag analysis.

4 Associated top-charged Higgs production $tH^\pm \rightarrow ttb$

In the MSSM, associated production of charged Higgs with a top quark produces a final state rich in b jets. Recent excitement was generated by a claim⁹ that the “wedge region” in $\tan\beta$ ($\tan\beta \sim 6$ where the h^0 shares equal coupling to top and bottom) could be explored up to 2 TeV in H^\pm mass at a 14 TeV LHC through the channel $tbH^\pm \rightarrow tb(tb)$. In contrast, others¹⁰ found that even 500 GeV could not be probed. We explore¹¹ whether the ~ 2 TeV limit can be reached in the wedge region, and how μ_x tagging performs in this final state.

Using MLM-matched samples for $tH^\pm \rightarrow ttb$ generated in MadEvent, showered in PYTHIA, and reconstructed in DELPHES, we look for final states involving one boosted-top tag, one μ_x boosted- b tag, and a fully reconstructed $t \rightarrow bl\nu$ decay (with a normal low-energy b tag). The background is dominated by fake tags from $t\bar{t}j$ and tjj that we take as measured from CMS data. After cuts on the relative p_T and angle of the two leading jets, we find $S/B \sim 1/10$. A preliminary estimate of the reach in $H^\pm tb$ Yukawa coupling y_{tb} and $\tan\beta$ are shown vs. H^\pm mass in Fig. 3. Our analysis appears to extend the results of Ref. 10 up to 2 TeV — meaning the wedge remains. It appears the wedge region will need to wait until a 100 TeV machine.




Figure 3 – 95% C.L. exclusion limits that can be reached as a function of H^\pm mass at a 14 TeV LHC for (left) $H^\pm tb$ Yukawa coupling y_{tb} and (right) $\tan \beta$.

5 Conclusions

At Moriond 2016 I presented the new μ_x boosted-bottom jet tag and its applications to some important searches for new physics at the LHC. Combining angular information x from B hadron decay with jet substructure f_{subjet} in TeV-scale jets allows for clean extraction of signals for Z' and MSSM charged Higgs above backgrounds. We find that the reach for leptophobic Z' discovery at a 13 TeV LHC is about 1 TeV higher than current limits. If a Z' is not found, 95% C.L. exclusion limits can be set up to 2 TeV higher, or g_B couplings a factor of 2 smaller, than the current limits.

Despite recent excitement, the search for MSSM charged Higgs in the mid- $\tan \beta$ “wedge” region in $tH^\pm \rightarrow t(tb)$ will remain elusive. The signal appears to be too small when realistic tagging efficiencies are applied. A 100 TeV collider is likely needed to fill this region. On the other hand, the μ_x tag could be used to immediately improve the existing searches for $W' \rightarrow t\bar{b}$ in the boosted-top and boosted-bottom channel.¹² The μ_x boosted-bottom jet tag is a powerful new tool in the exploration for physics beyond the Standard Model.

Acknowledgments

This work was supported by the U.S. Department of Energy under award No. DE-SC0008347.

References

1. J. de Favereau *et al.* (DELPHES 3 Collaboration), J. High Energy Phys. **02**, 057 (2014).
2. K. Pedersen and Z. Sullivan, Phys. Rev. D **93**, 014014 (2016).
3. <https://github.com/keith-pedersen/delphes/tree/MuXboostedBTagging>
4. B.A. Dobrescu and F. Yu, Phys. Rev. D **88**, 035021 (2013); Erratum: Phys. Rev. D **90**, 079901 (2014).
5. J. Alwall *et al.*, J. High Energy Phys. **07**, 079 (2014).
6. S. Dulat *et al.*, Phys. Rev. D **93**, 033006 (2016).
7. T. Sjostrand, S. Mrenna, and P.Z. Skands, Comput. Phys. Commun. **178**, 852 (2008).
8. Z. Sullivan and K. Pedersen, “Flavor tagging TeV jets for physics beyond the Standard Model,” to appear in *Proceedings of the XLVth International Symposium on Multiparticle Dynamics (ISMD 2015)*.
9. J. Hajer, Y. Y. Li, T. Liu and J. F. H. Shiu, J. High Energy Phys. **11**, 124 (2015).
10. N. Craig, F. D’Eramo, P. Draper, S. Thomas and H. Zhang, J. High Energy Phys. **06**, 137 (2015).
11. K. Pedersen and Z. Sullivan, in production.
12. D. Duffy and Z. Sullivan, Phys. Rev. D **90**, 015031 (2014).