

HEISENBERG UNIQUENESS PAIRS FOR SOME ALGEBRAIC CURVES AND SURFACES

D.K. GIRI AND R.K. SRIVASTAVA

ABSTRACT. Let $X(\Gamma)$ be the space of all finite Borel measure μ in \mathbb{R}^2 which is supported on the curve Γ and absolutely continuous with respect to the arc length of Γ . For $\Lambda \subset \mathbb{R}^2$, the pair (Γ, Λ) is called a Heisenberg uniqueness pair for $X(\Gamma)$ if any $\mu \in X(\Gamma)$ satisfies $\hat{\mu}|_{\Lambda} = 0$, implies $\mu = 0$. We explore the Heisenberg uniqueness pairs corresponding to the cross, exponential curves, and surfaces. Then, we prove a characterization of the Heisenberg uniqueness pairs corresponding to finitely many parallel lines. We observe that the size of the determining sets Λ for $X(\Gamma)$ depends on the number of lines and their irregular distribution that further relates to a phenomenon of interlacing of certain trigonometric polynomials.

1. INTRODUCTION

The notion of the Heisenberg uniqueness pair has been first introduced by Hedenmalm and Montes-Rodríguez as a version of the uncertainty principle (see [9]). We would like to mention that Heisenberg uniqueness pair up to some extent is similar to an annihilating pair of Borel measurable sets of positive measure as described by Havin and Jöricke (see [8]). To describe this, consider a pair of Borel measurable sets $\Gamma, \Lambda \subseteq \mathbb{R}$. Then (Γ, Λ) form a mutually annihilating pair if any $\varphi \in L^2(\Gamma)$ whose Fourier transform $\hat{\varphi}$ supported on Λ , implies $\varphi = 0$.

Heisenberg uniqueness pair has a close relation with the long-standing problem of determining the *exponential types* for finite measure which is eventually about exploring the density of the set $\{e^{i\lambda t} : \lambda \in \Lambda\}$ in $L^2(\mu)$. For more details, we refer to Poltoratski [15, 16]. In particular, the question of HUP can be thought as a dual problem of *gap problem* (see [15]). Let μ be a finite Borel measure which is supported on a closed set $\Gamma \subset \mathbb{R}$ and

$$G_{\Gamma} = \sup \{a > 0 : \exists \mu \neq 0, \text{ supp } \mu \subset \Gamma \text{ satisfies } \hat{\mu}|_{[0,a]} = 0\}.$$

Let $\Lambda \subset \mathbb{R}$. Then (Γ, Λ) would be a HUP as long as the determining set Λ intersect $[0, G_{\Gamma}]$ at most on a set of measure zero.

Date: August 7, 2021.

2000 Mathematics Subject Classification. Primary 42A38; Secondary 44A35.

Key words and phrases. Bessel function, convolution, Fourier transform, Wiener's lemma, symmetric polynomial.

In addition, the concept of determining the Heisenberg uniqueness pair for a class of finite measures has also a significant similarity with the celebrated result due to M. Benedicks (see [4]). That is, support of a function $f \in L^1(\mathbb{R}^n)$ and its Fourier transform \hat{f} both cannot be of finite measure simultaneously. Later, various analogues of the Benedicks theorem have been investigated in different set ups, including the Heisenberg group and Euclidean motion groups (see [14, 17, 19]).

Further, the concept of Heisenberg uniqueness pair has a sharp contrast to the known results about determining sets for measures by Sitaram et al. [3, 18], due to the fact that the determining set Λ for the function $\hat{\mu}$ has also been considered a thin set. In particular, if Γ is compact, then $\hat{\mu}$ is a real analytic function having exponential growth and it can vanish on a very delicate set. Hence, in this case, finding the Heisenberg uniqueness pairs becomes little easier. However, this question becomes immensely difficult when the measure is supported on a non-compact curve.

In the article [9], Hedenmalm and Montes-Rodríguez have shown that the pair (hyperbola, some discrete set) is a Heisenberg uniqueness pair. As a dual problem, a weak* dense subspace of $L^\infty(\mathbb{R})$ has been constructed to solve the Klein-Gordon equation. Further, a complete characterization of the Heisenberg uniqueness pairs corresponding to any two parallel lines has been given by Hedenmalm and Montes-Rodríguez (see [9]).

Afterward, a considerable amount of work has been done pertaining to the Heisenberg uniqueness pair in the plane as well as in the Euclidean spaces.

Recently, Lev [12] and Sjölin [20] have independently shown that circle and certain system of lines are HUP corresponding to the unit circle S^1 . Further, Vieli [23] has generalized HUP corresponding to circle in the higher dimension and shown that a sphere whose radius does not lie in the zero set of the Bessel functions $J_{(n+2k-2)/2}$; $k \in \mathbb{Z}_+$, the set of non-negative integers, is a HUP corresponding to the unit sphere S^{n-1} . In a recent article [22], the second author has shown that a cone is a Heisenberg uniqueness pair corresponding to sphere as long as the cone does not completely lay on the level surface of any homogeneous harmonic polynomial on \mathbb{R}^n . Thereafter, a sense of evidence emerged that the exceptional sets for the HUPs corresponding to the sphere are eventually contained in the zero sets of the spherical harmonics and the Bessel functions, though we yet resolve it (see [6, 22]).

Further, Sjölin [21] has investigated some of the Heisenberg uniqueness pairs corresponding to the parabola. Subsequently, Babot [2] has given a characterization of the Heisenberg uniqueness pairs corresponding to a certain system of three parallel lines. Thereafter, the authors in [6] have given some necessary and sufficient conditions for the Heisenberg uniqueness pairs corresponding to a system of four parallel lines. However, an exact analogue for the finitely many parallel lines is still unsolved.

In a major development, Jaming and Kellay [10] have given a unifying proof for some of the Heisenberg uniqueness pairs corresponding to the hyperbola, polygon, ellipse and graph of the functions $\varphi(t) = |t|^\alpha$, whenever $\alpha > 0$. Further, Gröchenig and Jaming [7] have worked out some of the Heisenberg uniqueness pairs corresponding to the quadratic surface.

Let Γ be a finite disjoint union of smooth curves in \mathbb{R}^2 . Let $X(\Gamma)$ be the space of all finite complex-valued Borel measure μ in \mathbb{R}^2 which is supported on Γ and absolutely continuous with respect to the arc length measure on Γ . For $(\xi, \eta) \in \mathbb{R}^2$, the Fourier transform of μ is defined by

$$\hat{\mu}(\xi, \eta) = \int_{\Gamma} e^{-i\pi(x \cdot \xi + y \cdot \eta)} d\mu(x, y).$$

In the above context, the function $\hat{\mu}$ becomes a uniformly continuous bounded function on \mathbb{R}^2 . Thus, we can analyze the pointwise vanishing nature of the function $\hat{\mu}$.

Definition 1.1. Let Λ be a set in \mathbb{R}^2 . The pair (Γ, Λ) is called a Heisenberg uniqueness pair for $X(\Gamma)$ if any $\mu \in X(\Gamma)$ satisfies $\hat{\mu}|_{\Lambda} = 0$, implies $\mu = 0$.

Since the Fourier transform is invariant under translation and rotation, one can easily deduce the following invariance properties about the Heisenberg uniqueness pair.

- (i) Let $u_o, v_o \in \mathbb{R}^2$. Then the pair (Γ, Λ) is a HUP if and only if the pair $(\Gamma + u_o, \Lambda + v_o)$ is a HUP.
- (ii) Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be an invertible linear transform whose adjoint is denoted by T^* . Then (Γ, Λ) is a HUP if and only if $(T^{-1}\Gamma, T^*\Lambda)$ is a HUP.

Now, we state first known results on the Heisenberg uniqueness pair due to Hedenmalm and Montes-Rodríguez [9]. Then we briefly summarize the recent progress on this problem.

Theorem 1.2. [9] *Let $\Gamma = L_1 \cup L_2$, where L_j ; $j = 1, 2$ are any two parallel straight lines and Λ a subset of \mathbb{R}^2 such that $\overline{\pi(\Lambda)} = \mathbb{R}$. Then (Γ, Λ) is a Heisenberg uniqueness pair if and only if the set*

$$(1.1) \quad \tilde{\Lambda} = \pi_1^a(\Lambda) \cup [\pi_1^b(\Lambda) \setminus \pi_1^c(\Lambda)]$$

is dense in \mathbb{R} .

Here we avoid mentioning the notations appeared in (1.1) as they are bit involved, however, we have written down the same notations as in the article [9]. Though, their main features can be perceived in Section 3.

Theorem 1.3. [9] *Let Γ be the hyperbola $x_1x_2 = 1$ and $\Lambda_{\alpha, \beta}$ a lattice-cross defined by*

$$\Lambda_{\alpha, \beta} = (\alpha\mathbb{Z} \times \{0\}) \cup (\{0\} \times \beta\mathbb{Z}),$$

where α, β are positive reals. Then $(\Gamma, \Lambda_{\alpha, \beta})$ is a Heisenberg uniqueness pair if and only if $\alpha\beta \leq 1$.

For $\xi \in \Lambda$, defining a function e_ξ on Γ by $e_\xi(x) = e^{i\pi x \cdot \xi}$. As a dual problem to Theorem 1.3, Hedenmalm and Montes-Rodríguez [9] have proved the following density result which in turn solve the one-dimensional Klein-Gordon equation.

Theorem 1.4. [9] *The pair (Γ, Λ) is a Heisenberg uniqueness pair if and only if the set $\{e_\xi : \xi \in \Lambda\}$ is a weak* dense subspace of $L^\infty(\Gamma)$.*

Let Γ denote a system of three parallel lines in the plane that can be expressed as $\Gamma = \mathbb{R} \times \{0, 1, p\}$, for some $p \in \mathbb{N}$ with $p \geq 2$. The following characterization for the Heisenberg uniqueness pairs corresponding to the above mentioned three parallel lines has been given by D.B. Babot [2].

Theorem 1.5. [2] *Let $\Gamma = \mathbb{R} \times \{0, 1, p\}$ and $\Lambda \subset \mathbb{R}^2$ a closed set which is 2-periodic with respect to the second variable. Then (Γ, Λ) is a HUP if and only if the set*

$$(1.2) \quad \tilde{\Lambda} = \Pi^3(\Lambda) \cup [\Pi^2(\Lambda) \setminus \Pi^{2*}(\Lambda)] \cup [\Pi^1(\Lambda) \setminus \Pi^{1*}(\Lambda)]$$

is dense in \mathbb{R} .

For the notations appeared in (1.2), we refer the article [2], as those notations are quite involved. However, they are the prototype of few notations those will appear in Section 3.

Afterward, the authors have given the following characterization for the Heisenberg uniqueness pairs corresponding to a certain system of four parallel lines (see [6]). Let Γ denotes a system of four parallel straight lines in \mathbb{R}^2 which can be represented by $\Gamma = \mathbb{R} \times \{\alpha, \beta, \gamma, \delta\}$, where $\alpha < \beta < \gamma < \delta$ with $(\gamma - \alpha)/(\beta - \alpha) = 2$ and $(\delta - \alpha)/(\beta - \alpha) \in \mathbb{N}$. By the invariance properties of HUP, one can assume that $\Gamma = \mathbb{R} \times \{0, 1, 2, p\}$ for some $p \geq 3$.

Theorem 1.6. [6] *Let $\Gamma = \mathbb{R} \times \{0, 1, 2, p\}$ and $\Lambda \subset \mathbb{R}^2$ be a closed set which is 2-periodic with respect to the second variable. Suppose $\Pi(\Lambda)$ is dense in \mathbb{R} . If (Γ, Λ) is a Heisenberg uniqueness pair, then the set*

$$\tilde{\Pi}(\Lambda) = \Pi^4(\Lambda) \bigcup_{j=0}^2 [\Pi^{(3-j)}(\Lambda) \setminus \Pi^{(3-j)*}(\Lambda)]$$

is dense in \mathbb{R} . Conversely, if the set

$$(1.3) \quad \tilde{\Pi}_p(\Lambda) = \Pi^4(\Lambda) \bigcup_{j=2}^3 [\Pi^j(\Lambda) \setminus \Pi_{j*}^p(\Lambda)] \cup [\Pi^1(\Lambda) \setminus \Pi^{1*}(\Lambda)]$$

is dense in \mathbb{R} , then (Γ, Λ) is a Heisenberg uniqueness pair.

For the notations appeared in (1.3), we refer the article [6]. However, the nature of their occurrence can be understood in the beginning of Section 3 when we formulate the Heisenberg uniqueness pairs for finitely many parallel lines.

Remark 1.7. The above problem becomes immensely deeper when we consider the p -th line at large that we shall elaborate in Section 3, Remark 3.1.

2. SOME INDIVIDUAL RESULTS FOR HEISENBERG UNIQUENESS PAIRS

In this section, we will explore some of the individual results for the Heisenberg uniqueness pairs corresponding to the cross and certain exponential curves in the plane as well as exponential surfaces in the Euclidean spaces.

Let $\Gamma = (\mathbb{R} \times F_n) \cup (F_n \times \mathbb{R})$, where $F_n = \{0, 1, \dots, n\}$. Consider a subset $\mathcal{C} \subset X(\Gamma)$ such that for any $\mu \in \mathcal{C}$, there exist $f_j \in L^1(\mathbb{R})$; $j \in F_n$ satisfying

$$(2.1) \quad d\mu(x, y) = \sum_{k=0}^n (f_k(x)dx d\delta_k(y) - f_k(y)d\delta_k(x)dy),$$

where δ_k denotes the point mass measure at k . Then the pair (Γ, Λ) is said to be a \mathcal{C} -HUP if any measure $\mu \in \mathcal{C}$ satisfies $\hat{\mu}|_{\Lambda} = 0$, implies μ is identically zero.

For $n = 0$, let $\alpha_j \in \mathbb{R}$; $j = 1, 2$ be independent over \mathbb{Q} , the set of rational numbers. Consider the lines $L_{\alpha_j} = \{(\xi, \eta) \in \mathbb{R}^2 : \eta - \xi = \alpha_j\}$; $j = 1, 2$.

Theorem 2.1. *Let $\Gamma = (\mathbb{R} \times \{0\}) \cup (\{0\} \times \mathbb{R})$ and $\Lambda = L_{\alpha_1} \cup L_{\alpha_2}$, where α_j ; $j = 1, 2$ are independent over \mathbb{Q} . Then (Γ, Λ) is a \mathcal{C} -HUP.*

Proof. For $\mu \in \mathcal{C}$ there exist $f \in L^1(\mathbb{R})$ such that

$$d\mu(x, y) = f(x)dx d\delta_o(y) - f(y)d\delta_o(x)dy.$$

By taking the Fourier transform of both the sides, we get

$$\begin{aligned} \hat{\mu}(\xi, \eta) &= \int_{\Gamma} e^{-i\pi(x\xi + y\eta)} d\mu(x, y) \\ &= \int_{\mathbb{R}} e^{-i\pi x\xi} f(x)dx - \int_{\mathbb{R}} e^{-i\pi y\eta} f(y)dy \\ &= \hat{f}(\xi) - \hat{f}(\eta). \end{aligned}$$

Now, $\hat{\mu}|_{L_{\alpha_1}} = 0$ implies that $\hat{f}(t) = \hat{f}(t + \alpha_1)$ for all $t \in \mathbb{R}$. Using the invariance property (ii), we can assume that $\alpha_j > 0$; $j = 1, 2$. Hence, \hat{f} is a periodic function with some period $p_o > 0$. By the simple recursions we can derive that $\hat{f}(t) = \hat{f}(t + m\alpha_1)$ for all $m \in \mathbb{Z}$ and $t \in \mathbb{R}$. Similarly, by the condition $\hat{\mu}|_{L_{\alpha_2}} = 0$, it follows that $\hat{f}(t) = \hat{f}(t + n\alpha_2)$ for all $n \in \mathbb{Z}$ and $t \in \mathbb{R}$. Thus, we infer that

$$\hat{f}(t) = \hat{f}(t + m\alpha_1 + n\alpha_2)$$

whenever, $m, n \in \mathbb{Z}$ and $t \in \mathbb{R}$. By Kröneckers approximation theorem (see [1]), we can always find $m_o, n_o \in \mathbb{Z}$ such that $|m_o\alpha_1 + n_o\alpha_2| < p_o$, which contradicts the fact that p_o is the period of f . This, in turn, implies that \hat{f} must be a constant function. Thus, by Riemann-Lebesgue lemma $f = 0$. \square

Remark 2.2. (i). Let $\Gamma_o = (\mathbb{R} \times \{0\}) \cup (\{0\} \times \mathbb{R})$ and $\Lambda = \{(\xi, \eta) \in \mathbb{R}^2 : \eta = \alpha\xi, |\alpha| < 1\}$. Then by Riemann-Lebesgue lemma, it can be easily deduced that (Γ_o, Λ) is a \mathcal{C} -HUP. On the other hand, if $\Lambda \subseteq \{(\xi, \eta) \in \mathbb{R}^2 : \eta = \xi\}$, then (Γ_o, Λ) is not a \mathcal{C} -HUP. Next, for $\Lambda = \{(\xi, \eta) \in \mathbb{R}^2 : \eta = \xi^2\}$, one can easily verify that (Γ_o, Λ) is a \mathcal{C} -HUP.

(ii). Consider $\Gamma = (\mathbb{R} \times F_n) \cup (F_n \times \mathbb{R})$ and $\Lambda = \{(\xi, \eta) \in \mathbb{R}^2 : \eta = \xi\}$. Then (Γ, Λ) is not a \mathcal{C} -HUP. For this, we can choose non-zero functions $f_j \in L^1(\mathbb{R})$; $j \in F_n$ such that

$$(2.2) \quad \hat{\mu}(\xi, \eta) = \sum_{k=0}^n \left(e^{-ik\pi\eta} \hat{f}_k(\xi) - e^{-ik\pi\xi} \hat{f}_k(\eta) \right)$$

and hence $\hat{\mu}(\xi, \eta) = 0$, whenever $\xi = \eta$. For $n = 1$, if Λ is the ξ -axis, then $(\hat{f}_0 + \hat{f}_1)(\xi) = \hat{f}_0(0) + e^{-i\pi\xi} \hat{f}_1(0)$ for all $\xi \in \mathbb{R}$. By Riemann-Lebesgue lemma, it is enough to consider $f_1 = -f_0$ and $\hat{f}_0(0) = 0$. Thus, (Γ, Λ) is not a \mathcal{C} -HUP.

(iii). In view of injectivity of the Fourier transform on $L^1(\mathbb{R})$, for a pair (Γ_o, Λ) to be a \mathcal{C} -HUP it is necessary that at least one of the orthogonal projections $\pi_j(\Lambda)$; $j = 1, 2$ on the axes must be dense in \mathbb{R} . It would be an interesting question to get a sufficient condition for the \mathcal{C} -Heisenberg uniqueness pairs corresponding to Γ_o .

Next, we will work out some of the Heisenberg uniqueness pairs corresponding to certain exponential curve and surfaces in the Euclidean spaces.

Let μ be a finite Borel measure having support on $\Gamma = \{(t, e^{t^2}) : t \in \mathbb{R}\}$ which is absolutely continuous with respect to the arc length on Γ . Then there exists $f \in L^1(\mathbb{R})$ such that $d\mu = g(t)dt$, where $g(t) = f(t)\sqrt{1 + 4t^2e^{2t^2}}$. Hence by finiteness of μ , it follows that $g \in L^1(\mathbb{R})$ and

$$(2.3) \quad \hat{\mu}(x, y) = \int_{\mathbb{R}} e^{-i\pi(xt + ye^{t^2})} g(t) dt.$$

Then $\hat{\mu}$ satisfies the PDE

$$(2.4) \quad (1 + \mathcal{T}_x) \hat{\mu} = \alpha \partial_y \hat{\mu}$$

in the distributional sense, where $\mathcal{T}_x = \sum_{n=0}^{\infty} \left(\sum_{k=1}^4 \frac{\alpha^{4n+k}}{(4n+k)!} \partial_x^{4n+k} \right)$ and $\alpha = \frac{i}{\pi}$.

Theorem 2.3. Let $\Gamma = \{(t, \alpha(t)) : t \in \mathbb{R}\}$, where $\alpha : \mathbb{R} \rightarrow \mathbb{R}_+$ be defined by $\alpha(t) = e^{t^2}$.

(i). Let Λ be a straight line. Then (Γ, Λ) is a Heisenberg uniqueness pair if and only if Λ is parallel to the x -axis.

(ii). Let L_j ; $j = 1, 2$ be two parallel lines which are not parallel to either of the axes. Then $(\Gamma, L_1 \cup L_2)$ is a Heisenberg uniqueness pair.

In order to prove Theorem 2.3, we need the following results. First, we state a result which can be found in Havin and Jöricke [8] at p. 36.

Lemma 2.4. [8] *Let $\varphi \in L^1[0, \infty)$. If $\int_{\mathbb{R}} \log |\hat{\varphi}| \frac{dx}{1+x^2} = -\infty$, then $\varphi = 0$.*

Next, we state the following form of Radon-Nikodym derivative theorem (see [5], p.91).

Proposition 2.5. *Let ν be a σ -finite signed measure which is absolutely continuous with respect to a σ -finite measure μ on the measure space (X, \mathcal{M}) . If $g \in L^1(\nu)$, then $g \frac{d\nu}{d\mu} \in L^1(\mu)$ and $\int g d\nu = \int g \frac{d\nu}{d\mu} d\mu$.*

As a consequence of Lemma 2.4 and Proposition 2.5, we prove the following result. Let $h_c : \mathbb{R} \rightarrow \mathbb{R}$ defined by $h_c(t) = (t + \frac{1}{2c})^2 + e^{t^2} - t^2$, where c is a non-zero constant. Let $|E|$ denotes the Lebesgue measure of the set $E \subset \mathbb{R}$.

Lemma 2.6. *Let $g \in L^1(\mathbb{R})$. Suppose $E \subset \mathbb{R}$ and $|E| > 0$. Then*

$$(2.5) \quad \int_{\mathbb{R}} e^{-i\pi cx h_c(t)} g(t) dt = 0$$

for all $x \in E$ if and only if $\psi_c(u) = g \circ h_c^{-1}(u^2) \frac{2u}{h'_c \circ h_c^{-1}(u^2)}$ is an odd function.

Proof. By Proposition 2.5, we can write the left-hand side of (2.5) as

$$(2.6) \quad \begin{aligned} I &= \int_{\mathbb{R}} e^{-i\pi cx h_c(t)} g(t) dt \\ &= \int_{\mathbb{R}} e^{-i\pi cx u^2} g \circ h_c^{-1}(u^2) \frac{2u du}{h'_c \circ h_c^{-1}(u^2)} \\ &= \int_{\mathbb{R}} e^{-i\pi cx u^2} \psi_c(u) du. \end{aligned}$$

Hence by Proposition 2.5 it follows that $\psi_c \in L^1(\mathbb{R})$ and we get

$$(2.7) \quad \begin{aligned} I &= \int_0^{\infty} e^{-i\pi cx u^2} \psi_c(u) du + \int_0^{\infty} e^{-i\pi cx u^2} \psi_c(-u) du \\ &= \int_0^{\infty} e^{-i\pi cx u^2} F_c(u) du, \end{aligned}$$

where $F_c(u) = \psi_c(u) + \psi_c(-u)$ for all $u > 0$. Clearly $F_c \in L^1(0, \infty)$ and by the change of variables $u^2 = v$, we have

$$(2.7) \quad I = \int_0^{\infty} e^{-i\pi cx v} F_c(\sqrt{v}) \frac{dv}{2\sqrt{v}}.$$

Let $\varphi(v) = F_c(\sqrt{v})/2\sqrt{v} \chi_{(0, \infty)}(v)$. Then $\varphi \in L^1(\mathbb{R})$ and from (2.7) we obtain $I = \hat{\varphi}(cx) = 0$ for all $x \in E$. That is, $\hat{\varphi}$ vanishes on the set cE of positive measure. Thus, by Lemma 2.4 we conclude that $\varphi = 0$ a.e. Hence, it follows that $F_c = 0$ a.e. on $[0, \infty)$.

Conversely, if ψ_c is an odd function, then (2.5) trivially holds. □

As a corollary to Lemma 2.6, we can easily derive the following result.

Corollary 2.7. *Let $g \in L^1(\mathbb{R})$ and $\alpha : \mathbb{R} \rightarrow \mathbb{R}_+$ be defined by $\alpha(t) = e^{t^2}$. Suppose $E \subset \mathbb{R}$ and $|E| > 0$. Then*

$$(2.8) \quad \int_{\mathbb{R}} e^{-i\pi x \alpha(t)} g(t) dt = 0$$

for all $x \in E$ if and only if g is an odd function.

Proposition 2.8. *Suppose $E \subset \mathbb{R}$ and $|E| > 0$. Assume $c, d \in \mathbb{R}$ with $c, d \neq 0$. Then*

(i) $\hat{\mu}(x, cx) = 0$ for all $x \in E$ if and only if ψ_c is an odd function.

(ii) $\hat{\mu}(x, cx+d) = 0$ for all $x \in E$ if and only if $\phi_c(u) = \chi \circ h_c^{-1}(u^2) \frac{2u}{h'_c \circ h_c^{-1}(u^2)}$

is an odd function of u , where $\chi(t) = e^{-i\pi d e^{t^2}} g(t)$.

Proof. (i). From (2.3) we can express

$$\hat{\mu}(x, cx) = \int_{\mathbb{R}} e^{-i\pi x(t+ce^{t^2})} g(t) dt = e^{i\pi x/4c} \int_{\mathbb{R}} e^{-i\pi cx h_c(t)} g(t) dt.$$

By Lemma 2.6, ψ_c is odd if and only if $\hat{\mu}(x, cx) = 0$ for all $x \in E$.

(ii). By a simple computation, we get

$$\hat{\mu}(x, cx+d) = \int_{\mathbb{R}} e^{-i\pi x(t+ce^{t^2})} \chi(t) dt = e^{i\pi x/4c} \int_{\mathbb{R}} e^{-i\pi cx h_c(t)} \chi(t) dt.$$

As similar to the above case, ϕ_c is odd if and only if $\hat{\mu}(x, cx+d) = 0$ for all $x \in E$. \square

Proof of Theorem 2.3. (i). In view of the invariance property (i), we can assume that Λ is the x -axis. Recall that $\hat{\mu}$ satisfies

$$\hat{\mu}(x, y) = \int_{\mathbb{R}} e^{-i\pi(xt+ye^{t^2})} g(t) dt.$$

Hence $\hat{\mu}|_{\Lambda} = 0$ implies that $\hat{g}(x) = 0$ for all $x \in \mathbb{R}$. Thus, we conclude that $\mu = 0$.

Conversely, suppose Λ is not parallel to the x -axis. If Λ is parallel to the y -axis, then by Corollary 2.7, it follows that (Γ, Λ) is not a HUP. Hence we can assume that Λ of the form $y = cx$, where $c \neq 0$. Choose a non-zero odd function $\varphi \in L^1(\mathbb{R})$ and let $g(t) = \frac{\varphi(\sqrt{h_c(t)} h'_c(t))}{2\sqrt{h_c(t)}}$. Then by Proposition 2.8, it follows that (Γ, Λ) is not a Heisenberg uniqueness pair.

(ii). Let $L_1 = \{(x, cx) : x \in \mathbb{R}\}$ and $L_2 = \{(x, cx+d) : x \in \mathbb{R}\}$, where $c, d \neq 0$. Since $\hat{\mu}|_{L_j} = 0$; $j = 1, 2$, by Proposition 2.8 it follows that ψ_c and ϕ_c

are odd functions. Let u_+ and u_- be the square roots of h_c . Since ϕ_c is an odd function, it implies that

$$\left[e^{i\pi d \left\{ e^{(h_c^{-1}(u_-)^2)} - e^{(h_c^{-1}(u_+)^2)} \right\}} - 1 \right] \psi_c(u) = 0.$$

That is, $\psi_c = 0$ a.e. and hence $g \circ h_c^{-1}(u^2) = 0$ almost all u . Thus, the pair $(\Gamma, L_1 \cup L_2)$ is a Heisenberg uniqueness pair. \square

Remark 2.9. Let $\alpha : \mathbb{R} \rightarrow \mathbb{R}_+$ be an even smooth function having finitely many local extrema and consider $\Gamma = \{(t, \alpha(t)) : t \in \mathbb{R}\}$. Then the conclusions of Theorem 2.3 would also hold.

Theorem 2.10. *Let Γ be the surface $x_{n+1} = e^{x_1^2} + \dots + e^{x_n^2}$ in \mathbb{R}^{n+1} and Λ an affine hyperplane in \mathbb{R}^{n+1} of dimension n . Then (Γ, Λ) is a Heisenberg uniqueness pair if and only if Λ is parallel to the hyperplane $x_{n+1} = 0$.*

For $u = (u_1, \dots, u_n)$, denoting $\varphi(u) = e^{u_1^2} + \dots + e^{u_n^2}$. Let μ be a finite Borel measure which is supported on $\Gamma = \{(u, \varphi(u)) : u \in \mathbb{R}^n\}$ and absolutely continuous with respect to the surface measure on Γ . Then by Radon-Nikodym theorem, there exists a measurable function f on \mathbb{R}^n such that $d\mu = g(u)du$, where $g(u) = f(u)\sqrt{1 + |\text{grad } \varphi(u)|^2}$. Then by the finiteness of μ , it follows that $g \in L^1(\mathbb{R}^n)$. Denote $u' = (u_2, \dots, u_n)$ and $x' = (x_2, \dots, x_n)$. Then the Fourier transform of μ can be expressed as

$$(2.9) \quad \hat{\mu}(x) = \int_{\mathbb{R}^n} e^{-\pi i(x' \cdot u' + x_{n+1}\varphi(u))} g(u) du$$

for $x \in \mathbb{R}^{n+1}$.

Proof of Theorem 2.10. Since Λ is an affine hyperplane in \mathbb{R}^{n+1} of dimension n , by the invariance properties of HUP, we can assume that Λ is a linear subspace of \mathbb{R}^{n+1} which can be considered as either $x_{n+1} = cx_1$, where $c \in \mathbb{R}$ or $x_1 = 0$.

If $\Lambda = \{(x_1, \dots, x_{n+1}) \in \mathbb{R}^{n+1} : x_{n+1} = 0\}$, then by the hypothesis, $\hat{\mu}|_{\Lambda} = 0$ implies $\hat{g} = 0$ on \mathbb{R}^n . Thus, it follows that (Γ, Λ) is a HUP.

Conversely, suppose Λ is not parallel to the hyperplane $x_{n+1} = 0$. Consider a non-zero compactly supported odd function $\psi \in L^1(\mathbb{R})$ together with a non-zero compactly supported function $h \in L^1(\mathbb{R}^{n-1})$. Then we have the following two cases.

(i). If $\Lambda = \{(x_1, \dots, x_{n+1}) \in \mathbb{R}^{n+1} : x_1 = 0\}$ and $g(u) = \psi(u_1)h(u')$. Then for $x \in \Lambda$, we have

$$\begin{aligned} \hat{\mu}(x) &= \int_{\mathbb{R}^n} e^{-\pi i(x'.u' + x_{n+1}\varphi(u))} g(u) du \\ &= \int_{\mathbb{R}^n} e^{-\pi i(x'.u' + x_{n+1}\varphi(u))} \psi(u_1)h(u') du \\ &= \int_{\mathbb{R}^{n-1}} e^{-\pi i\{x'.u' + x_{n+1}(\varphi(u) - e^{u_1^2})\}} \left(\int_{\mathbb{R}} e^{-\pi i x_{n+1} e^{u_1^2}} \psi(u_1) du_1 \right) h(u') du' \\ &= 0. \end{aligned}$$

Thus, (Γ, Λ) is not a Heisenberg uniqueness pair.

(ii). Suppose $\Lambda = \{(x_1, \dots, x_{n+1}) \in \mathbb{R}^{n+1} : x_{n+1} = cx_1, c \neq 0\}$. Consider a function $\tau(t) = \frac{\psi(\sqrt{h_c(t)})h'_c(t)}{2\sqrt{h_c(t)}}$ and write $g(u) = \tau(u_1)h(u')$. If we denote $x'' = (x_1, \dots, x_n)$. Then for $x \in \Lambda$, we have

$$\begin{aligned} \hat{\mu}(x) &= \int_{\mathbb{R}^n} e^{-\pi i(x''.u + cx_1\varphi(u))} g(u) du \\ &= \int_{\mathbb{R}^n} e^{-\pi i(x''.u + cx_1\varphi(u))} \tau(u_1)h(u') du \\ &= \int_{\mathbb{R}^{n-1}} e^{-\pi i\{x''.u' + cx_1(\varphi(u) - e^{u_1^2})\}} \left(\int_{\mathbb{R}} e^{-\pi i(x_1 u_1 + cx_1 e^{u_1^2})} \tau(u_1) du_1 \right) h(u') du'. \end{aligned}$$

By Lemma 2.6, it follows that

$$\begin{aligned} \int_{\mathbb{R}} e^{-\pi i x_1 (u_1 + c e^{u_1^2})} \tau(u_1) du_1 &= \int_{\mathbb{R}} e^{-\pi i c x_1 h_c(u_1)} \tau(u_1) du_1 \\ &= \int_{\mathbb{R}} e^{-\pi i c x_1 t^2} \psi(t) dt \\ &= 0. \end{aligned}$$

Thus, we conclude that (Γ, Λ) is not a Heisenberg uniqueness pair. \square

3. HEISENBERG UNIQUENESS PAIRS CORRESPONDING TO THE FINITE NUMBER OF PARALLEL LINES

A characterization of the Heisenberg uniqueness pairs corresponding to two parallel lines was being done by Hedenmalm and Montes-Rodríguez (see [9]). Afterward, Babot [2] worked out an analogous result for a certain system of three parallel lines. Further, authors had extended the above result to a system of four parallel lines (see [6]). In this section, we shall prove a characterization of the Heisenberg uniqueness pairs corresponding to a certain system of finitely many parallel lines. In the latter case, we observe the phenomenon of interlacing of the finitely many totally disconnected sets.

Denote $F_s = \{0, 1, \dots, s\}$. Let Γ_o be a system of $n + 2$ parallel lines that can be expressed as $\Gamma_o = \mathbb{R} \times \{\alpha_o, \dots, \alpha_{n+1}\}$, where $\alpha_o < \dots < \alpha_{n+1}$ and $p_j = (\alpha_j - \alpha_o)/(\alpha_1 - \alpha_o)$ with

$$p_j = \begin{cases} j & \text{if } 2 \leq j \leq n, \\ p \in \mathbb{N} \setminus F_n & \text{if } j = n + 1. \end{cases}$$

If (Γ_o, Λ_o) is a HUP, then by the invariance property (i), (Γ_o, Λ_o) can be thought as $(\Gamma_o - (0, \alpha_o), \Lambda_o)$. Since scaling in \mathbb{R}^2 can be viewed as a diagonal matrix, by the invariance property (ii), $(\Gamma_o - (0, \alpha_o), \Lambda_o)$ can be considered as $(T^{-1}(\Gamma_o - (0, \alpha_o)), T^*\Lambda_o)$, where $T = \text{diag}\{(\alpha_1 - \alpha_o), (\alpha_1 - \alpha_o)\}$. Let $\Gamma = T^{-1}(\Gamma_o - (0, \alpha_o))$ and $\Lambda = T^*\Lambda_o$. Then $\Gamma = \mathbb{R} \times (F_n \cup \{p\})$, where $p \in \mathbb{N}$ with $p \geq n + 1$. In view of the above facts, (Γ_o, Λ_o) is a HUP if and only if (Γ, Λ) is a HUP.

To state our main result of this section, we need to set up some necessary notations and few auxiliary results.

Let $\mu \in X(\Gamma)$. Then there exist $f_k \in L^1(\mathbb{R})$; $k \in F_{n+1}$ such that

$$(3.1) \quad d\mu(x, y) = \sum_{j=0}^n f_j(x) dx d\delta_j(y) + f_{n+1}(x) dx d\delta_p(y),$$

where δ_t denotes the point mass measure at t . For $(\xi, \eta) \in \mathbb{R}^2$, the Fourier transform of μ can be written as

$$(3.2) \quad \hat{\mu}(\xi, \eta) = \sum_{j=0}^n e^{j\pi i \eta} \hat{f}_j(\xi) + e^{p\pi i \eta} \hat{f}_{n+1}(\xi).$$

Since μ is supported on $\Gamma = \{(x, y) \in \mathbb{R}^2 : P(y) \equiv \prod_{k=0}^n (y - k)(y - p) = 0\}$, the function $\hat{\mu}$ will satisfy the PDE

$$(3.3) \quad P\left(\frac{1}{\pi i} \frac{\partial}{\partial \eta}\right) \hat{\mu} = 0$$

in the distributional sense with the initial condition $\hat{\mu}|_{\Lambda} = 0$. On the other hand, if (3.3) holds, then $\text{supp } \mu \subset \Gamma$.

Remark 3.1. Notice that for each fixed $(\xi, \eta) \in \Lambda \subset \mathbb{R}^2$, the right-hand side of (3.2) is a trigonometric polynomial of degree p that could have preferably some missing terms. Therefore, it is an interesting question to find out the smallest determining set Λ for the above trigonometric polynomial. We observe that the size of Λ depends on the choice of number of lines as well as irregular separation among themselves. That is, larger number of lines or value of p would force smaller size of Λ . Eventually, the problem would become more difficult for large value of p .

Observe that $\hat{\mu}$ is a 2-periodic function in the second variable. Thus, for any set $\Lambda \subset \mathbb{R}^2$, it is enough to consider the set

$$\mathcal{L}(\Lambda) = \{(\xi, \eta) : (\xi, \eta + 2k) \in \Lambda, \text{ for some } k \in \mathbb{Z}\}$$

for the purpose of HUP. Also, it is easy to verify that (Γ, Λ) is a HUP if and only if $(\Gamma, \overline{\mathcal{L}(\Lambda)})$ is a HUP, where $\overline{\mathcal{L}(\Lambda)}$ denotes the closure of $\mathcal{L}(\Lambda)$ in \mathbb{R}^2 . Thus, it is enough to work with the closed set $\Lambda \subset \mathbb{R}^2$ which is 2-periodic in the second variable.

Remark 3.2. Further, it is evident from the Riemann-Lebesgue lemma that the exponential functions, which appeared in (3.2), can not be expressed as the Fourier transform of functions in $L^1(\mathbb{R})$. However, by Wiener's lemma, they can be locally agreed with the Fourier transform of functions in $L^1(\mathbb{R})$.

Hence, in view of Remark 3.2 and the condition $\hat{\mu}|_\Lambda = 0$, we can classify these exponential functions in the form of the following annihilating spaces which will create dispensable sets in the determining set Λ .

Given $E \subset \mathbb{R}$ and a point $\xi \in E$, let I_ξ be an interval containing ξ . We define

(A1). $L_{loc}^{E,\xi} = \{\psi : E \rightarrow \mathbb{C} \text{ such that there is an interval } I_\xi \text{ and a function } \varphi \in L^1(\mathbb{R}) \text{ satisfying } \psi = \hat{\varphi} \text{ on } I_\xi \cap E\}$.

Next, for $2 \leq m \leq n$, we define

(A2). $P^{1,m}[L_{loc}^{E,\xi}] = \{\psi : E \rightarrow \mathbb{C} \text{ such that there is an interval } I_\xi \text{ and functions } \varphi_j \in L^1(\mathbb{R}); j \in F_{m-1} \text{ satisfying } \psi^m + \hat{\varphi}_{m-1}\psi^{m-1} + \dots + \hat{\varphi}_1\psi + \hat{\varphi}_0 = 0 \text{ on } I_\xi \cap E\}$.

Finally, we define the p -th space by

(A3). $P^{1,p}[L_{loc}^{E,\xi}] = \{\psi : E \rightarrow \mathbb{C} \text{ such that there is an interval } I_\xi \text{ and functions } \varphi_j \in L^1(\mathbb{R}); j \in F_n \text{ satisfying } \psi^p + \hat{\varphi}_n\psi^n + \dots + \hat{\varphi}_1\psi + \hat{\varphi}_0 = 0 \text{ on } I_\xi \cap E\}$.

We frequently need the following Wiener's lemma that will be a key ingredient for most of the proofs in the sequel.

Lemma 3.3. [11] *Let $\psi \in L_{loc}^{E,\xi}$ and $\psi(\xi) \neq 0$. Then $1/\psi \in L_{loc}^{E,\xi}$.*

For more details see [11], p.57.

Next, we establish a relation among the sets defined by **(A1)**, **(A2)** and **(A3)** that will help in figure out the dispensable sets.

Lemma 3.4. *The following inclusions hold.*

$$(3.4) \quad L_{loc}^{E,\xi} \subset P^{1,2}[L_{loc}^{E,\xi}] \subset \dots \subset P^{1,n}[L_{loc}^{E,\xi}] \subset P^{1,p}[L_{loc}^{E,\xi}].$$

For proving Lemma 3.4, we need the following result which is appeared in [6].

Lemma 3.5. [6] *For $p \geq 3$, the following inclusions hold.*

$$(3.5) \quad L_{loc}^{E,\xi} \subset P^{1,2}[L_{loc}^{E,\xi}] \subset P^{1,p}[L_{loc}^{E,\xi}].$$

Proof of Lemma 3.4. (a). By Lemma 3.5, it follows that $P^{1,2}[L_{loc}^{E,\xi}] \subset P^{1,3}[L_{loc}^{E,\xi}]$. Next, we assume that $P^{1,m-1}[L_{loc}^{E,\xi}] \subset P^{1,m}[L_{loc}^{E,\xi}]$, whenever $3 \leq m < n + 1$. If

$\psi \in P^{1,m}[L_{loc}^{E,\xi}]$, then there exists I_ξ containing ξ and $h_j \in L^1(\mathbb{R})$; $j \in F_{m-1}$ such that

$$(3.6) \quad \psi^m + \hat{h}_{m-1}\psi^{m-1} + \cdots + \hat{h}_1\psi + \hat{h}_0 = 0$$

on $I_\xi \cap E$. Now, consider a function $\chi \in L^1(\mathbb{R})$ such that $I_\xi \subset \text{supp } \hat{\chi}$. After multiplying (3.6) by ψ and $\hat{\chi}$ separately and adding the resultant equations, we can write

$$\psi^{m+1} + \left(\hat{\chi} + \hat{h}_{m-1}\right)\psi^m + \cdots + \left(\hat{\chi}\hat{h}_1 + \hat{h}_0\right)\psi + \hat{\chi}\hat{h}_0 = 0.$$

Hence, for the appropriate choice of φ_j ; $j \in F_m$, we have

$$\psi^{m+1} + \hat{\varphi}_m\psi^m + \cdots + \hat{\varphi}_1\psi + \hat{\varphi}_0 = 0$$

on $I_\xi \cap E$. Thus $\psi \in P^{1,m+1}[L_{loc}^{E,\xi}]$.

(b). It only remains to show that $P^{1,n}[L_{loc}^{E,\xi}] \subset P^{1,p}[L_{loc}^{E,\xi}]$. If $\psi \in P^{1,n}[L_{loc}^{E,\xi}]$, then there exists I_ξ containing ξ and $\varphi_j \in L^1(\mathbb{R})$; $j \in F_n$ such that

$$(3.7) \quad \psi^{n+1} + \hat{\varphi}_n\psi^n + \cdots + \hat{\varphi}_1\psi + \hat{\varphi}_0 = 0$$

on $I_\xi \cap E$. After multiplying (3.7) by ψ and $\hat{\varphi}_n$ separately, we can write

$$\psi^{n+2} + (\hat{\varphi}_{n-1} - \hat{\varphi}_n\hat{\varphi}_n)\psi^n + \cdots + (\hat{\varphi}_0 - \hat{\varphi}_1\hat{\varphi}_n)\psi + (-\hat{\varphi}_0\hat{\varphi}_n) = 0.$$

Hence, for the appropriate choice of $\varphi_j \in L^1(\mathbb{R})$; $j \in F_n$ and induction hypothesis, we conclude that

$$\psi^p + \hat{\varphi}_n\psi^n + \cdots + \hat{\varphi}_1\psi + \hat{\varphi}_0 = 0$$

on $I_\xi \cap E$. Thus $\psi \in P^{1,p}[L_{loc}^{E,\xi}]$. \square

Let $\Pi(\Lambda)$ be the projection of Λ on the ξ - axis. For partitioning the projection $\Pi(\Lambda)$, we introduce the notion of symmetric polynomials. Since the level surface of a complete homogeneous symmetric polynomial consists of only finitely many those points which can be represented by an upper triangular matrix. More details will be given in **Appendix A**.

For each $k \in \mathbb{Z}_+$, the complete homogeneous symmetric polynomial h_k of degree k is the sum of all monomials of degree k . That is,

$$h_k(x_1, \dots, x_s) = \sum_{\alpha_1 + \cdots + \alpha_s = k} x_1^{\alpha_1} \cdots x_s^{\alpha_s}.$$

Given $\xi \in \Pi(\Lambda)$, we denote the corresponding images on the η - axis by

$$\Sigma_\xi = \{\eta \in [0, 2) : (\xi, \eta) \in \Lambda\}.$$

Since it is expected that the set Σ_ξ may consist more than one images which depend upon the order of winding in Λ around the line $\{\xi\} \times \mathbb{R}$, the set $\Pi(\Lambda)$ can be decomposed into the following disjoint sets.

(**P**₁). $\Pi^1(\Lambda) = \{\xi \in \Pi(\Lambda) : \text{there is unique } \eta_0 \in \Sigma_\xi\}$.

Next, for $2 \leq m \leq n$ we define the following $n - 1$ partitioning sets.

(\mathbf{P}_m). $\Pi^m(\Lambda) = \{\xi \in \Pi(\Lambda) : \text{there are exactly } m \text{ distinct } \eta_j \in \Sigma_\xi; j \in F_{m-1}\}$.

For $n + 2$ distinct images $\eta_j \in [0, 2)$, denote $a_j = e^{\pi i \eta_j}$; $j \in F_{n+1}$. Then, in view of **Appendix A**, the remaining partitioning sets can be described as follows.

(\mathbf{P}_{n+1}). $\Pi^{n+1}(\Lambda) = \{\xi \in \Pi(\Lambda) : \text{there are at least } n + 1 \text{ distinct } \eta_j \in \Sigma_\xi \text{ for } j \in F_n \text{ and if there is another } \eta_{n+1} \in \Sigma_\xi, \text{ then } h_{p-n}(a_0, \dots, a_{n-1}, a_n) = h_{p-n}(a_0, \dots, a_{n-1}, a_{n+1})\}$.

(\mathbf{P}_{n+2}). $\Pi^{n+2}(\Lambda) = \{\xi \in \Pi(\Lambda) : \text{there are at least } n + 2 \text{ distinct } \eta_j \in \Sigma_\xi \text{ for } j \in F_{n+1} \text{ satisfies}$

$$(3.8) \quad h_{p-n}(a_0, \dots, a_{n-1}, a_n) \neq h_{p-n}(a_0, \dots, a_{n-1}, a_{n+1})\}.$$

In this way, we have a decomposition $\Pi(\Lambda) = \bigcup_{j=1}^{n+2} \Pi^j(\Lambda)$.

Next, we derive some of the auxiliary results related to $\Pi^m(\Lambda)$; $2 \leq m \leq n+1$. For m distinct images $\eta_j \in [0, 2)$, we write $a_j = e^{\pi i \eta_j}$; $j \in F_{m-1}$. Consider the system of equations

$$(3.9) \quad A_\xi^m X = B_\xi^m,$$

where $b_s = (a_o^s, \dots, a_{m-1}^s)$; $0 \leq s \leq m-1$ are the column vectors of A_ξ^m and $B_\xi^m = -(a_o^m, \dots, a_{m-1}^m)$. Since $\det A_\xi^m = \prod_{0 \leq i < j \leq m-1} (a_j - a_i) \neq 0$, the solution $X_\xi = (\tau_o, \dots, \tau_{m-1})$ can be expressed as

$$\tau_k(\xi) = \begin{cases} (-1)^m \prod_{i=0}^{m-1} a_i & \text{if } k = 0, \\ (-1)^{m-1} \sum_{i=0}^{m-1} \prod_{j \neq i} a_j & \text{if } k = 1, \\ \vdots & \vdots \\ \sum_{0 \leq i < j \leq m-1} a_i a_j & \text{if } k = m-2, \\ - \sum_{i=0}^{m-1} a_i & \text{if } k = m-1. \end{cases}$$

Observe that τ_k are constant multiples of the elementary symmetric polynomials in a_j ; $j \in F_{m-1}$. Therefore, by Wiener's lemma, τ_k can be locally agreed with the Fourier transform of some functions in $L^1(\mathbb{R})$.

Thus, the following sets in the projection $\Pi(\Lambda)$ will be dispensable in the process of getting the determining sets Λ for $X(\Gamma)$. Eventually, dispensable sets are those sets in $\Pi(\Lambda)$ for which we could not solve (3.2).

\mathbf{P}_{1^*} . Since each $\xi \in \Pi^1(\Lambda)$ has unique image in Σ_ξ , we can define a function a_o on $\Pi^1(\Lambda)$ by $a_o(\xi) = e^{\pi i \eta_0}$, where $\eta_0 = \eta_0(\xi) \in \Sigma_\xi$. Now, the first dispensable

set can be described by

$$\Pi^{1*}(\Lambda) = \left\{ \xi \in \Pi^1(\Lambda) : a_o \in P^{1,p}[L_{loc}^{\Pi^1(\Lambda),\xi}] \right\}.$$

Next, for $2 \leq m \leq n+1$ we define the m -th dispensable set $\Pi^{m*}(\Lambda)$ as follows. Given $\xi \in \Pi^m(\Lambda)$, let $a_j(\xi) = e^{\pi i \eta_j}$, where $\eta_j = \eta_j(\xi) \in \Sigma_\xi$; $j \in F_{m-1}$.

P_{m*}¹. Since each $\xi \in \Pi^m(\Lambda)$ has m distinct images in Σ_ξ , we define m many functions τ_j on $\Pi^m(\Lambda)$; $j \in F_{m-1}$ such that $X_\xi = (\tau_o, \dots, \tau_{m-1})$ is the solution of (3.9). Thus, the m -th dispensable set can be defined by

$$\Pi^{m*}(\Lambda) = \left\{ \xi \in \Pi^m(\Lambda) : \tau_j \in L_{loc}^{\Pi^2(\Lambda),\xi}; j \in F_{m-1} \right\}.$$

Now, for $m \leq k \leq n+1$, the corresponding dispensable sets can be defined by

P_{m*}². Further, we define k many functions τ_j on $\Pi^m(\Lambda)$; $j \in F_{k-1}$ such that $X_\xi = (\tau_o, \dots, \tau_{k-1})$ satisfies $A_\xi^{m,k} X_\xi = B_\xi^{m,k}$, where $b_s = (a_o^s, \dots, a_{m-1}^s)$; $0 \leq s \leq k-1$ are the column vectors of $A_\xi^{m,k}$ and $B_\xi^{m,k} = -(a_o^k, \dots, a_{m-1}^k)$. Hence, an auxiliary dispensable set can be described by

$$\Pi_{m*}^k(\Lambda) = \left\{ \xi \in \Pi^m(\Lambda) : \tau_j \in L_{loc}^{\Pi^m(\Lambda),\xi}; j \in F_{k-1} \right\}.$$

P_{m*}³. Finally, we define $n+1$ many functions τ_j on $\Pi^m(\Lambda)$; $j \in F_n$ such that $X_\xi = (\tau_o, \dots, \tau_n)$ satisfies $A_\xi^{m,p} X_\xi = B_\xi^{m,p}$, where $b_s = (a_o^s, \dots, a_{m-1}^s)$; $0 \leq s \leq n$ are the column vectors of $A_\xi^{m,p}$ and $B_\xi^{m,p} = -(a_o^p, \dots, a_{m-1}^p)$. Thus, the p -th dispensable set can be defined by

$$\Pi_{m*}^p(\Lambda) = \left\{ \xi \in \Pi^m(\Lambda) : \tau_j \in L_{loc}^{\Pi^2(\Lambda),\xi}; j \in F_n \right\}.$$

Now, we prove the following two results that speak a sharp contrast in the pattern of dispensable sets as compared to dispensable sets corresponding to two or three lines results. That is, a larger value of p will increase the size of dispensable sets in case of more than three finitely many parallel lines.

Lemma 3.6. *For $2 \leq m \leq n$, the following inclusion holds.*

$$\Pi^{m*}(\Lambda) \subset \Pi_{m*}^{m+1}(\Lambda) \subset \dots \subset \Pi_{m*}^n(\Lambda) \subset \Pi_{m*}^p(\Lambda).$$

Proof. If $\xi_o \in \Pi^{m*}(\Lambda)$, then $\tau_j \in L_{loc}^{\Pi^m(\Lambda),\xi_o}$; $j \in F_{m-1}$. Hence there exists an interval I_{ξ_o} containing ξ_o and $\varphi_j \in L^1(\mathbb{R})$ such that $\tau_j = \hat{\varphi}_j$; $j \in F_{m-1}$ and satisfy

$$\hat{\varphi}_o + \hat{\varphi}_1 a_j + \dots + \hat{\varphi}_{m-1} a_j^{m-1} + a_j^m = 0$$

on $I_{\xi_o} \cap \Pi^m(\Lambda)$ for $j \in F_{m-1}$. By similar iterations as in the proof of Lemma 3.4(a), we infer that there exist a common set of $\psi_j \in L^1(\mathbb{R})$; $j \in F_m$ such that

$$\hat{\psi}_o + \hat{\psi}_1 a_j + \dots + \hat{\psi}_m a_j^m + a_j^{m+1} = 0$$

on $I_{\xi_o} \cap \Pi^m(\Lambda)$; $j \in F_{m-1}$. If we denote $\hat{\psi}_j = \tau_j$, then $\xi_o \in \Pi_{m^*}^{m+1}(\Lambda)$. By induction it follows that if $\xi_o \in \Pi_{m^*}^k(\Lambda)$, then $\xi_o \in \Pi_{m^*}^{k+1}(\Lambda)$ for $m+1 \leq k \leq n$. Since $\xi_o \in \Pi_{m^*}^{n+1}(\Lambda)$, there exist a common set of functions $\psi_j \in L^1(\mathbb{R})$; $j \in F_n$ such that

$$\hat{\psi}_o + \hat{\psi}_1 a_j + \cdots + \hat{\psi}_n a_j^n + a_j^{n+1} = 0$$

on $I_{\xi_o} \cap \Pi^m(\Lambda)$; $j \in F_{m-1}$. By similar iterations as in the proof of Lemma 3.4(b), we infer that there exist a common set of $\psi_j \in L^1(\mathbb{R})$; $j \in F_n$ such that

$$\hat{\psi}_o + \hat{\psi}_1 a_j + \cdots + \hat{\psi}_n a_j^n + a_j^p = 0$$

on $I_{\xi_o} \cap \Pi^m(\Lambda)$; $j \in F_{m-1}$. If we denote $\hat{\psi}_j = \tau_j$, then $\xi_o \in \Pi_{m^*}^p(\Lambda)$. \square

Lemma 3.7. *The following inclusion holds.*

$$\Pi^{(n+1)^*}(\Lambda) \subseteq \Pi_{(n+1)^*}^p(\Lambda).$$

Moreover, equality holds for $p = n + 1$.

Proof. If $\xi_o \in \Pi^{(n+1)^*}(\Lambda)$, then $\tau_j \in L_{loc}^{\Pi^{n+1}(\Lambda), \xi_o}$. Hence there exists an interval I_{ξ_o} containing ξ_o and $\varphi_j \in L^1(\mathbb{R})$ such that $\tau_j = \hat{\varphi}_j$; $j \in F_n$ satisfy

$$\hat{\varphi}_o + \hat{\varphi}_1 a_j + \cdots + \hat{\varphi}_n a_j^n + a_j^{n+1} = 0$$

on $I_{\xi_o} \cap \Pi^{n+1}(\Lambda)$ for $j \in F_n$. By similar iterations as in the proof of Lemma 3.4(b), we infer that there exists the common set of $\psi_j \in L^1(\mathbb{R})$; $j \in F_n$ such that

$$\hat{\varphi}_o + \hat{\varphi}_1 a_j + \cdots + \hat{\varphi}_n a_j^n + a_j^p = 0$$

on $I_{\xi_o} \cap \Pi^{n+1}(\Lambda)$; $j \in F_n$. If we denote $\hat{\psi}_j = \tau_j$, then it follows that $\xi_o \in \Pi_{(n+1)^*}^p(\Lambda)$. \square

Now, we state our main result of this section.

Theorem 3.8. *Let $\Gamma = \mathbb{R} \times (F_n \cup \{p\})$ and $\Lambda \subset \mathbb{R}^2$ be a closed set which is 2-periodic with respect to the second variable. Suppose $\Pi(\Lambda)$ is dense in \mathbb{R} . If (Γ, Λ) is a Heisenberg uniqueness pair, then the set*

$$\tilde{\Pi}(\Lambda) = \Pi^{n+2}(\Lambda) \bigcup_{j=0}^n \left[\Pi^{(n-j+1)}(\Lambda) \setminus \Pi^{(n-j+1)^*}(\Lambda) \right]$$

is dense in \mathbb{R} . Conversely, if the set

$$\tilde{\Pi}_p(\Lambda) = \Pi^{n+2}(\Lambda) \bigcup_{j=0}^{n-1} \left[\Pi^{(n-j+1)}(\Lambda) \setminus \Pi_{(n-j+1)^*}^p(\Lambda) \right] \cup \left[\Pi^1(\Lambda) \setminus \Pi^{1^*}(\Lambda) \right]$$

is dense in \mathbb{R} , then (Γ, Λ) is a Heisenberg uniqueness pair.

Remark 3.9. If Γ is a system of $n + 2$ consecutive parallel lines, then we get $p = n + 1$ and from Lemma 3.7 it follows that $\Pi^{(n+1)^*}(\Lambda) = \Pi_{(n+1)^*}^{n+1}(\Lambda)$. Thus, the necessary and sufficient conditions in Theorem 3.8 will be nearly identical. However, In view of Lemma 3.6, it follows that $\Pi^{m^*}(\Lambda)$ is properly contained

in $\Pi_m^{n+1}(\Lambda)$; $2 \leq m \leq n$. Hence, $\tilde{\Pi}_{n+1}(\Lambda)$ is a proper subset of $\tilde{\Pi}(\Lambda)$. Thus, a unique necessary and sufficient condition corresponding to finitely many lines is still open.

We need the following two lemmas to prove the necessary part of Theorem 3.8. The main purpose of these lemmas is to pull down an interval from some of the partitioning sets of the projection $\Pi(\Lambda)$. The above argument helps to negate the assumption that $\tilde{\Pi}(\Lambda)$ is not dense in \mathbb{R} . Now, we prove the following lemmas for $2 \leq m \leq n+1$.

Lemma 3.10. *Suppose I is an interval such that $I \cap \Pi^{m*}(\Lambda)$ is dense in I . Then there exists an interval $I' \subset I$ such that $I' \subset \Pi^{m*}(\Lambda) \bigcup_{j=m+1}^{n+2} \Pi^j(\Lambda)$.*

Proof. If $\xi_o \in I \cap \Pi^{m*}(\Lambda)$, then there exist $\tau_j \in L_{loc}^{\Pi^m(\Lambda), \xi_o}$; $j \in F_{m-1}$ such that $X_{\xi_o} = (\tau_0, \dots, \tau_{m-1})$ is the solution of (3.9). For $\xi \in \Pi^m(\Lambda)$, consider the matrix

$$\mathcal{M}_\xi^{m, n+1} = [b_o \ \dots \ b_{m-2} \ b_{n+1}],$$

where $b_s = (a_o^s, \dots, a_{m-1}^s)$; $s \in F_{m-2} \cup \{n+1\}$ are the column vectors. Now, we define a function ρ on $\Pi^m(\Lambda)$ by $\rho(\xi) = (\det \mathcal{M}_\xi^{m, n+1})^2$.

Since τ_j ; $j \in F_{m-1}$ are constant multiples of the elementary symmetric polynomials, by the fundamental theorem of symmetric polynomials, ρ can be expressed as a polynomial in τ_j ; $j \in F_{m-1}$. Moreover, $\rho(\xi_o) \neq 0$. Hence it follows that $\rho \in L_{loc}^{\Pi^m(\Lambda), \xi_o}$. By hypothesis, $I \cap \Pi^{m*}(\Lambda)$ is dense in I , there exists an interval $I_{\xi_o} \subset I$ containing ξ_o such that ρ can be continuously extended on I_{ξ_o} . Thus, by continuity of ρ on I_{ξ_o} , there exists an interval $J \subset I_{\xi_o}$ containing ξ_o such that $\rho(\xi) \neq 0$ for all $\xi \in J$.

Consequently, $J \cap \Pi^{m*}(\Lambda)$ is dense in J and hence for $\xi \in J$, there exists a sequence $\xi_n \in J \cap \Pi^{m*}(\Lambda)$ such that $\xi_n \rightarrow \xi$. Thus, the corresponding image sequences $\eta_j^{(n)} \in \Sigma_{\xi_n} \subseteq [0, 2)$ will have convergent subsequences, say $\eta_j^{(n_k)}$ which converge to η_j ; $j \in F_{m-1}$. Since the set Λ is closed, $(\xi, \eta_j) \in \Lambda$ for $j \in F_{m-1}$.

Next, we claim that all of η_j ; $j \in F_{m-1}$ are distinct. On the contrary, suppose $\eta_j = \eta_k$ for some $j \neq k$; $j, k \in F_{m-1}$. Then by continuity of ρ on J , it follows that $\rho(\xi) = 0$, which contradicts the fact that $\rho(\xi) \neq 0$ for all $\xi \in J$.

Hence we infer that $J \subset \bigcup_{j=m}^{n+2} \Pi^j(\Lambda)$. Further, using the facts that $\tau_j \in L_{loc}^{\Pi^m(\Lambda), \xi_o}$

and $J \cap \Pi^{m*}(\Lambda)$ is dense in J , the functions τ_j can be extended continuously on an interval $I' \subset J$ containing ξ_o such that $\tau_j(\xi) \neq 0$ for all $\xi \in I'$. That is, if $\xi \in I' \cap \Pi^m(\Lambda)$, then $\tau_j \in L_{loc}^{\Pi^m(\Lambda), \xi}$ and hence $\xi \in \Pi^{m*}(\Lambda)$. Thus, we conclude

that $I' \subset \Pi^{m*}(\Lambda) \bigcup_{j=m+1}^{n+2} \Pi^j(\Lambda)$. □

Lemma 3.11. *Suppose I be an interval such that $I \cap \Pi_{(n+1)^*}^p(\Lambda)$ is dense in I . Then there exists an interval $I' \subset I$ such that $I' \subset \bigcup_{j=2}^{n+2} \Pi^j(\Lambda)$.*

Proof. If $\xi_o \in I \cap \Pi_{(n+1)^*}^p(\Lambda)$, then there exist $\tau_j \in L_{loc}^{\Pi^{n+1}(\Lambda), \xi_o}$; $j \in F_n$ such that $X_{\xi_o} = (\tau_o, \dots, \tau_n)$ is the solution of $A_{\xi_o}^{n+1} X_{\xi_o} = B_{\xi_o}^{n+1, p}$. Let $a_j = e^{\pi i \eta_j}$; $j \in F_n$. Then by a simple calculation, it follows that τ_n satisfies the estimate

$$|\tau_n(\xi_o)| = |h_{p-n}(a_o, \dots, a_n)| < C(p, n),$$

whenever $\xi_o \in I \cap \Pi_{(n+1)^*}^p(\Lambda)$. Since $I \cap \Pi_{(n+1)^*}^p(\Lambda)$ is dense in I , τ_n can be extended continuously on a small neighborhood containing ξ_o . Thus, there exists an interval $I' \subset I$ containing ξ_o such that

$$(3.10) \quad |\tau_n(\xi)| < C(p, n)$$

for all $\xi \in I'$.

Consequently, $I' \cap \Pi_{(n+1)^*}^p(\Lambda)$ is dense in I' and hence for $\xi \in I'$, there exists a sequence $\xi_n \in I' \cap \Pi_{(n+1)^*}^p(\Lambda)$ such that $\xi_n \rightarrow \xi$. Thus, the corresponding image sequences $\eta_j^{(n)} \in \Sigma_{\xi_n} \subseteq [0, 2)$ have a convergent subsequence, say $\eta_j^{(n_k)}$ which converges to η_j ; $j \in F_n$. Since the set Λ is closed, we get $(\xi, \eta_j) \in \Lambda$; $j \in F_n$.

We show that all of η_j ; $j \in F_n$ cannot be identical. If not, write $a_j^{(k)} = e^{\pi i \eta_j^{(n_k)}}$; $j \in F_n$, then

$$|\tau_n(\xi_{n_k})| = |h_{p-n}(a_o^{(k)}, \dots, a_n^{(k)})| \rightarrow C(p, n).$$

Further, by the continuity of τ_n on I' it follows that $|\tau_n(\xi_{n_k})| \rightarrow |\tau_n(\xi)|$. That is, $|\tau_n(\xi)| = C(p, n)$ which contradicts (3.10). Thus, $I' \subset \bigcup_{j=2}^{n+2} \Pi^j(\Lambda)$. \square

4. PROOF OF THEOREM 3.8

Proof of Theorem 3.8. We first prove the sufficient part of Theorem 3.8. Suppose the set $\tilde{\Pi}_p(\Lambda)$ is dense in \mathbb{R} . Then we show that (Γ, Λ) is a Heisenberg uniqueness pair. We claim that $\hat{f}_k|_{\tilde{\Pi}_p(\Lambda)} = 0$ for all $k \in F_{n+1}$. Since \hat{f}_k is continuous function which vanishing on a dense set $\tilde{\Pi}_p(\Lambda)$, it follows that $\hat{f}_k \equiv 0$ for all $k \in F_{n+1}$. Thus $\mu = 0$.

As $\Pi(\Lambda)$ decomposed into $n+2$ disjoint sets, the proof of the above assertion will be carried out in the following $n+2$ steps.

(S_{n+2}). If $\xi \in \Pi^{n+2}(\Lambda)$, then there exist at least $n+2$ distinct $\eta_j \in \Sigma_\xi$ for $j \in F_{n+1}$. Since $\hat{\mu}$ is vanishes on Λ , $\hat{\mu}(\xi, \eta_j) = 0$ for all $j \in F_{n+1}$. Hence $\hat{f}_k(\xi)$ for $k \in F_{n+1}$ satisfies a homogeneous system of $n+2$ equations. Since $\xi \in \Pi^{n+2}(\Lambda)$ and $h_{p-n}(a_0, \dots, a_{n-1}, a_n) \neq h_{p-n}(a_0, \dots, a_{n-1}, a_{n+1})$, it follows that $\hat{f}_k(\xi) = 0$ for all $k \in F_{n+1}$.

Next, for $2 \leq m \leq n+1$, we have the following cases.

(**S_m**). If $\xi \in \Pi^m(\Lambda)$, then there exist m distinct $\eta_j \in \Sigma_\xi$ such that $\hat{\mu}(\xi, \eta_j) = 0$; $j \in F_{m-1}$. That is,

$$(4.1) \quad \hat{f}_0(\xi) + \cdots + a_j^n(\xi) \hat{f}_n(\xi) + a_j^p(\xi) \hat{f}_{n+1}(\xi) = 0,$$

where $a_j(\xi) = e^{\pi i \eta_j}$; $j \in F_{m-1}$. In case if $\hat{f}_{n+1}(\xi) \neq 0$, then by applying the Wiener's lemma to (4.1), we conclude that $\xi \in \Pi_{m^*}^p(\Lambda)$. Thus, for $\xi \in \Pi^m(\Lambda) \setminus \Pi_{m^*}^p(\Lambda)$, we have $\hat{f}_{n+1}(\xi) = 0$.

Now, we continue the above argument for $m \leq k \leq n$.

Suppose $\hat{f}_k(\xi) \neq 0$ and $\hat{f}_s(\xi) = 0$; $s = k+1, \dots, n+1$, then by applying the Wiener's lemma to (4.1), we get $\xi \in \Pi_{m^*}^k(\Lambda)$. Thus, for $\xi \in \Pi^m(\Lambda) \setminus \Pi_{m^*}^k(\Lambda)$, we infer that $\hat{f}_k(\xi) = 0$.

Hence for $\xi \in \Pi^m(\Lambda) \setminus \Pi_{m^*}^p(\Lambda)$, we infer that $\hat{f}_k(\xi) = 0$ for all $k \in F_{n+1}$.

(**S₁**). If $\xi \in \Pi^1(\Lambda)$, then there exists unique $\eta_0 \in \Sigma_\xi$ such that $\hat{\mu}(\xi, \eta_0) = 0$. That is,

$$(4.2) \quad \hat{f}_0(\xi) + \cdots + a_o^n(\xi) \hat{f}_n(\xi) + a_o^p(\xi) \hat{f}_{n+1}(\xi) = 0,$$

where $a_o(\xi) = e^{\pi i \eta_0}$. In case if $\hat{f}_{n+1}(\xi) \neq 0$, then by applying Wiener's lemma to (4.2), we conclude that $a_o \in P^{1,p}[L_{loc}^{\Pi^1(\Lambda), \xi}]$. Thus for $\xi \in \Pi^1(\Lambda) \setminus \Pi^{1^*}(\Lambda)$, we have $\hat{f}_{n+1}(\xi) = 0$.

We repeat the above argument for $1 \leq k \leq n$.

Suppose $\hat{f}_k(\xi) \neq 0$ and $\hat{f}_s(\xi) = 0$; $s = k+1, \dots, n+1$, then by applying the Wiener's lemma to (4.2), we conclude that $a_o \in P^{1,k}[L_{loc}^{\Pi^1(\Lambda), \xi}]$. Hence, in view of Lemma 3.4, it follows that $\xi \in \Pi^{1^*}(\Lambda)$. Thus, for $\xi \in \Pi^1(\Lambda) \setminus \Pi^{1^*}(\Lambda)$, we infer that $\hat{f}_k(\xi) = 0$ for all $k \in F_{n+1}$. This completes the proof of sufficient condition. \square

Now, we prove the necessary part of Theorem 3.8. Suppose (Γ, Λ) is a Heisenberg uniqueness pair. Then we claim that the set $\tilde{\Pi}(\Lambda)$ is dense in \mathbb{R} . We observe that this is possible if the dispensable sets $\Pi^{j^*}(\Lambda)$; $j \in F_{n+1} \setminus \{0\}$ interlace to each other, though these sets are disjoint among themselves.

If possible, suppose $\tilde{\Pi}(\Lambda)$ is not dense in \mathbb{R} . Then there exists an open interval $I_o \subset \mathbb{R}$ such that $I_o \cap \tilde{\Pi}(\Lambda)$ is empty. This, in turn, implies that

$$(4.3) \quad \Pi(\Lambda) \cap I_o = \left(\bigcup_{j=1}^{n+1} \Pi^{j^*}(\Lambda) \right) \cap I_o.$$

From (4.3), we conclude that I_o intersects only the dispensable sets. Now, the remaining part of the proof of Theorem 3.8 is a consequence of the following two lemmas which provide the interlacing property of the dispensable sets $\Pi^{j^*}(\Lambda)$; $j \in F_{n+1} \setminus \{0\}$.

Lemma 4.1. *There does not exist any interval $J \subset I_o$ such that $\Pi(\Lambda) \cap J$ is contained in $\Pi^{j^*}(\Lambda)$; $j \in F_{n+1} \setminus \{0\}$.*

Proof. On the contrary, suppose there exists an interval $J \subset I_o$ such that $\Pi(\Lambda) \cap J \subset \Pi^{j^*}(\Lambda)$ for some $j \in F_{n+1} \setminus \{0\}$. Since $\Pi^{j^*}(\Lambda); j \in F_{n+1} \setminus \{0\}$ are disjoint, there could be following $n + 1$ possibilities.

(a). If $\xi_o \in \Pi(\Lambda) \cap J \subset \Pi^1(\Lambda)$, then $a_0 \in P^{1,p}[L_{loc}^{\Pi^1(\Lambda), \xi_o}]$ and hence there exists an interval $I_{\xi_o} \subset J$ containing ξ_o and $\varphi_k \in L^1(\mathbb{R}); k \in F_n$ such that

$$a_o^p + \hat{\varphi}_n a_o^n + \cdots + \hat{\varphi}_1 a_o + \hat{\varphi}_o = 0$$

on $I_{\xi_o} \cap \Pi^1(\Lambda)$. Now, consider a function $f_{n+1} \in L^1(\mathbb{R})$ such that $\hat{f}_{n+1}(\xi_o) \neq 0$ and $\text{supp } \hat{f}_{n+1} \subset I_{\xi_o}$. Let $f_k = f_{n+1} * \varphi_k; k \in F_n$, then we can construct a non-zero finite Borel measure μ which is supported on Γ that satisfies

$$\hat{\mu}(\xi, \eta) = \hat{f}_0(\xi) + a_0(\xi) \hat{f}_1(\xi) + \cdots + a_0^n(\xi) \hat{f}_n(\xi) + a_0^p(\xi) \hat{f}_{n+1}(\xi) = 0$$

for all $\xi \in I_{\xi_o} \cap \Pi^1(\Lambda)$, where $\eta \in \Sigma_\xi$. In addition, from (4.3) we conclude that $I_\xi \cap \Pi(\Lambda) = I_\xi \cap \Pi^1(\Lambda)$. This, in turn, implies $\hat{\mu}|_\Lambda = 0$. However, μ is a non-zero measure, which contradicts the fact that (Γ, Λ) is a HUP.

Next, for $2 \leq m \leq n$ we have the following cases.

(b). If $\xi_o \in \Pi(\Lambda) \cap J \subset \Pi^{m^*}(\Lambda)$, then by Lemma 3.6, $\xi_o \in \Pi_{m^*}^p(\Lambda)$ and hence there exists an interval $I_{\xi_o} \subset J$ containing ξ_o and $\varphi_k \in L^1(\mathbb{R}); k \in F_n$ such that

$$a_j^p + \hat{\varphi}_n a_j^n + \cdots + \hat{\varphi}_1 a_j + \hat{\varphi}_o = 0$$

on $I_{\xi_o} \cap \Pi^m(\Lambda)$ for $j \in F_{m-1}$. Let $f_{n+1} \in L^1(\mathbb{R})$ be such that $\hat{f}_{n+1}(\xi_o) \neq 0$ and $\text{supp } \hat{f}_{n+1} \subset I_{\xi_o}$. Consider $f_k = f_{n+1} * \varphi_k; k \in F_n$. Then we can construct a non-zero finite Borel measure μ which satisfies

$$\hat{\mu}(\xi, \eta_j) = \hat{f}_0(\xi) + a_j(\xi) \hat{f}_1(\xi) + \cdots + a_j^n(\xi) \hat{f}_n(\xi) + a_j^p(\xi) \hat{f}_{n+1}(\xi) = 0$$

for all $\xi \in I_{\xi_o} \cap \Pi^{m^*}(\Lambda)$, where $\eta_j \in \Sigma_\xi; j \in F_{m-1}$. Since $I_\xi \cap \Pi(\Lambda) = I_\xi \cap \Pi^{m^*}(\Lambda)$, it follows that $\hat{\mu}|_\Lambda = 0$. This contradicts the fact that (Γ, Λ) is a HUP.

(c). If $\xi_o \in \Pi(\Lambda) \cap J \subset \Pi^{(n+1)^*}(\Lambda)$, then by Lemma 3.7, $\xi_o \in \Pi_{(n+1)^*}^p(\Lambda)$ and hence there exists an interval $I_{\xi_o} \subset J$ containing ξ_o and $\varphi_k \in L^1(\mathbb{R}); k \in F_n$ such that

$$a_j^p + \hat{\varphi}_n a_j^n + \cdots + \hat{\varphi}_1 a_j + \hat{\varphi}_o = 0$$

on $I_{\xi_o} \cap \Pi^{n+1}(\Lambda)$ for $j \in F_n$. Let $f_{n+1} \in L^1(\mathbb{R})$ be such that $\hat{f}_{n+1}(\xi_o) \neq 0$ and $\text{supp } \hat{f}_{n+1} \subset I_{\xi_o}$. Let $f_k = f_{n+1} * \varphi_k; k \in F_n$. Then we can construct a non-zero finite Borel measure μ which satisfies

$$\hat{\mu}(\xi, \eta_j) = \hat{f}_0(\xi) + a_j(\xi) \hat{f}_1(\xi) + \cdots + a_j^n(\xi) \hat{f}_n(\xi) + a_j^p(\xi) \hat{f}_{n+1}(\xi) = 0$$

for all $\xi \in I_{\xi_o} \cap \Pi^{(n+1)^*}(\Lambda)$, where $\eta_j \in \Sigma_\xi; j \in F_n$. Since $I_\xi \cap \Pi(\Lambda) = I_\xi \cap \Pi^{(n+1)^*}(\Lambda)$, it follows that $\hat{\mu}|_\Lambda = 0$. This contradicts the fact that (Γ, Λ) is a HUP.

□

Next, we prove that there does not exist an interval $J \subset I_o$ which intersects more than one dispensable sets. Let $2 \leq s \leq n + 1$.

Lemma 4.2. *There does not exist any interval $J \subset I_o$ such that $\Pi(\Lambda) \cap J$ is contained in $\bigcup_{k=1}^s \Pi^{n_k^*}(\Lambda)$; $n_k \in F_{n+1} \setminus \{0\}$ with $n_1 < \dots < n_s$.*

Proof. (i). For $s = 2$, suppose there exists an interval $J \subset I_o$ such that $\Pi(\Lambda) \cap J \subset \Pi^{n_1^*}(\Lambda) \cup \Pi^{n_2^*}(\Lambda)$. Then, from (4.3) we get

$$(4.4) \quad J \cap \Pi(\Lambda) = J \cap (\Pi^{n_1^*}(\Lambda) \cup \Pi^{n_2^*}(\Lambda)).$$

We claim that $J \cap \Pi^{n_2^*}(\Lambda)$ is dense in J . If not, suppose there exists an interval $I \subset J$ such that $\Pi^{n_2^*}(\Lambda) \cap I = \emptyset$. Then from (4.3), we have $I \cap \Pi(\Lambda) = I \cap \Pi^{n_1^*}(\Lambda) \subset \Pi^{n_1^*}(\Lambda)$ which contradicts Lemma 4.1. Hence, by Lemma 3.10, there exists an interval $I' \subset J$ such that $I' \subset \Pi^{n_2^*}(\Lambda) \bigcup_{j=n_2+1}^{n+2} \Pi^j(\Lambda)$ which contradicts that I_o intersects only the dispensable sets.

(ii). Let $2 < s \leq n + 1$. Then for $2 \leq m < s$, we assume that there does not exist any interval $J \subset I_o$ such that

$$(4.5) \quad \Pi(\Lambda) \cap J \subset \bigcup_{k=1}^m \Pi^{n_k^*}(\Lambda); \quad n_k \in F_{n+1} \setminus \{0\}.$$

We claim that, there does not exist any interval $J \subset I_o$ such that

$$(4.6) \quad \Pi(\Lambda) \cap J \subset \bigcup_{k=1}^{m+1} \Pi^{n_k^*}(\Lambda); \quad n_k \in F_{n+1} \setminus \{0\}.$$

If possible, suppose there exists an interval $J \subset I_o$ such that (4.6) holds. Then $J \cap \Pi^{n_{m+1}^*}(\Lambda)$ is dense in J . On contrary, suppose there exists an interval $I \subset J$ such that $\Pi^{n_{m+1}^*}(\Lambda) \cap I = \emptyset$. Then by (4.3), it follows that $I \cap \Pi(\Lambda) \subset \bigcup_{k=1}^m \Pi^{n_k^*}(\Lambda)$, which in turn contradicts the assumption. Hence, by Lemma 3.10,

there exists an interval $I' \subset J$ such that $I' \subset \Pi^{n_{m+1}^*}(\Lambda) \bigcup_{j=n_{m+1}+1}^{n+2} \Pi^j(\Lambda)$ which contradict the assumption that I_o intersects only the dispensable sets. This completes the proof of Theorem 3.8. \square

5. APPENDIX A. CONSTRUCTION OF PARTITIONS OF $\Pi(\Lambda)$

In this section, we elaborate the use of symmetric homogeneous polynomials in reducing the following matrix to an upper triangular matrix that has crucial role in considering the p -th line at large.

Consider the homogeneous system of linear equations $AX = 0$, where entries of A are satisfying

$$a_{ij} = \begin{cases} \beta_i^j & \text{if } (i, j) \in F_{n+1} \times F_n, \\ \beta_i^p & \text{if } j = n + 1 \end{cases}$$

with $|\beta_i| = 1$; $i \in F_{n+1}$. Next, we reduce the matrix A to an upper triangular matrix by using the notion of the symmetric polynomial.

For each $k \in \mathbb{Z}_+$, the complete homogeneous symmetric polynomial h_k of degree k is the sum of all monomials of degree k which can be derived from the generating function identity

$$\sum_{k=0}^{\infty} h_k(x_1, \dots, x_s) t^k = \prod_{i=1}^s \frac{1}{1 - x_i t}.$$

For more details, we refer to [13]. Notice that

$$\begin{aligned} & \prod_{i=1}^s \left(\frac{1}{1 - x_i t} \right) \left(\frac{1}{1 - xt} \right) - \prod_{i=1}^s \left(\frac{1}{1 - x_i t} \right) \left(\frac{1}{1 - yt} \right) \\ &= t(x - y) \prod_{i=1}^s \left(\frac{1}{1 - x_i t} \right) \left(\frac{1}{1 - xt} \right) \left(\frac{1}{1 - yt} \right). \end{aligned}$$

By equating the coefficients of t^k in both sides, we get the identity

$$(5.1) \quad h_k(\bar{x}, x) - h_k(\bar{x}, y) = (x - y)h_{k-1}(\bar{x}, x, y),$$

where $\bar{x} = (x_1, \dots, x_s)$. By applying the chain of alternative row operations:

$$\begin{aligned} R'_i &= R_i - R_0; \quad i \in F_{n+1} \setminus \{0\}, \\ R''_i &= R'_i - \left(\frac{\beta_i - \beta_0}{\beta_1 - \beta_0} \right) R'_1; \quad i \in F_{n+1} \setminus \{0, 1\} \end{aligned}$$

and the identity (5.1), we obtain an upper triangular matrix with $(n+1, n+1)^{\text{th}}$ entry equals to

$$(h_{p-n}(\beta_0, \dots, \beta_{n-1}, \beta_n) - h_{p-n}(\beta_0, \dots, \beta_{n-1}, \beta_{n+1})) \prod_{j=0}^{n-1} (\beta_{n+1} - \beta_j).$$

6. APPENDIX B. CONCLUDING REMARKS

(a). We observed that for $\xi \in \Pi(\Lambda)$, if $\text{card}(\Sigma_\xi) \geq p + 1$ then $\xi \in \Pi^{n+2}(\Lambda)$. For this, given $c \in \mathbb{C}$, notice that there exists at most $p - n$ different images $\eta_o^{(k)} \in [0, 2)$ such that $\eta_o^{(k)} \neq \eta_s$; $s \in F_n \setminus \{0\}$ with all of η_s are distinct and satisfies

$$h_{p-n}(a_0^{(k)}, \dots, a_{n-1}, a_n) = c,$$

where $a_j = e^{\pi i \eta_j}$; $j \in F_n$.

In particular, if we consider $n + 2$ consecutive lines $\Gamma = \mathbb{R} \times F_{n+1}$, then $p = n + 1$ and condition (3.8) reduces to $a_n \neq a_{n+1}$. Hence, the projections $\Pi^{n+j}(\Lambda)$; $j = 1, 2$ regain to their simplest form. That is,

$$\begin{aligned}\Pi^{n+1}(\Lambda) &= \{\xi \in \Pi(\Lambda) : \text{card}(\Sigma_\xi) = n + 1\}, \\ \Pi^{n+2}(\Lambda) &= \{\xi \in \Pi(\Lambda) : \text{card}(\Sigma_\xi) \geq n + 2\}.\end{aligned}$$

Thus, we infer that for a system of consecutive parallel lines, Theorem 3.8 can be re-stated and could has a nearly unique necessary and sufficient condition.

(b). Let $\Gamma = \mathbb{R} \times F_{n+1}$ and $\Lambda = \mathbb{R} \times \{\frac{2k}{n+1}; k \in F_n\}$. Then for any $\xi \in \mathbb{R}$, $\text{card}(\Sigma_\xi) = n + 1$. Thus, $\xi \in \Pi^{n+1}(\Lambda)$ that in turn implies $\Pi^{(n+1)*}(\Lambda) = \mathbb{R}$. For this, let $\xi \in \Pi^{n+1}(\Lambda)$ and denote $a_k = e^{\pi i \eta_k}$, where $\eta_k = \frac{2k}{n+1}$; $k \in F_n$. Since $a_k^{n+1} = 1$, by a simple calculation, it follows that $X_\xi = ((-1)^{n+1}, 0, \dots, 0)$ is the solution of $A_\xi^{n+1} X_\xi = B_\xi^{n+1}$. Thus, $\xi \in \Pi^{(n+1)*}(\Lambda)$. By Theorem 3.8, we conclude that (Γ, Λ) is not a HUP.

(c). We observe a phenomenon of interlacing of $n + 1$ totally disconnected disjoint dispensable sets $\Pi^{(n-j+1)*}(\Lambda)$; $j \in F_n$ which are essentially derived from zero sets of $n + 2$ trigonometric polynomials. If Lemma 3.11 could be modified such that there exists $I' \subset I$ with $I' \subset \Pi_{(n+1)*}^p(\Lambda) \cup \Pi^{(n+2)}(\Lambda)$, then the necessary condition of Theorem 3.8 would be minimized. Hence a characterization of Λ for finitely many lines might be obtained that would be closed to three lines result. However, an exact analogue of three lines result for a large number of lines is still open.

(d). Finally, if we consider countably many parallel lines, then whether the projection $\Pi(\Lambda)$ would be still relevant after deleting the countably many dispensable sets, seems to be a reasonable question. However, since the dispensable sets are totally disconnected, one can think of analyzing countably many lines for HUP in terms of the Hausdorff dimension of the dispensable sets. We leave this question open for the time being.

Acknowledgements:

The authors extend a special thanks to Mr. Hiranmoy Pal for his contribution to **Appendix A** about the notion of symmetric polynomials. Finally, we gratefully acknowledge the support provided by IIT Guwahati, Government of India.

REFERENCES

- [1] T.M. Apostol, *Modular Functions and Dirichlet Series in Number Theory*, 2nd ed. New York: Springer-Verlag (1997).
- [2] D.B. Babot, *Heisenberg uniqueness pairs in the plane, Three parallel lines*, Proc. Amer. Math. Soc. 141 (2013), no. 11, 3899-3904.

- [3] S.C. Bagchi and A. Sitaram, *Determining sets for measures on \mathbb{R}^n* , Illinois J. Math. 26 (1982), no. 3, 419-422.
- [4] M. Benedicks, *On Fourier transforms of functions supported on sets of finite Lebesgue measure*, J. Math. Anal. Appl. 106 (1985), no.1, 180-183.
- [5] G.B. Folland, *Real analysis. Modern techniques and their applications*, Second edition, John Wiley and Sons, Inc., New York, 1999.
- [6] D.K. Giri and R. K. Srivastava, *Heisenberg uniqueness pairs for some algebraic curves in the plane*, Adv. Math., 310 (2017), 993-1016.
- [7] K. Gröchenig and P. Jaming, *The Cramér-Wold theorem on quadratic surfaces and Heisenberg uniqueness pairs*, arXiv:1608.06738.
- [8] V. Havin, and B. Jöricke, *The Uncertainty Principle in Harmonic Analysis*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], 28. Springer-Verlag, Berlin, 1994.
- [9] H. Hedenmalm, A. Montes-Rodríguez, *Heisenberg uniqueness pairs and the Klein-Gordon equation*, Ann. of Math. (2) 173 (2011), no. 3, 1507-1527.
- [10] P. Jaming and K. Kellay, *A dynamical system approach to Heisenberg uniqueness pairs*, J. Anal. Math., Springer, 2015, <http://link.springer.com/journal/11854>
- [11] J.P. Kahane, *Séries de Fourier absolument convergentes*, (French) Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 50 Springer-Verlag, Berlin-New York 1970
- [12] N. Lev, *Uniqueness theorem for Fourier transform*. Bull. Sc. Math., 135(2011), 134-140.
- [13] I.G. Macdonald, *Symmetric Functions and Hall Polynomials*, Second ed., Oxford University Press, New York, 1995.
- [14] E.K. Narayanan and P.K. Ratnakumar, *Benedick's theorem for the Heisenberg group*, Proc. Amer. Math. Soc. 138 (2010), no. 6, 2135-2140.
- [15] A. Poltoratski, *Spectral gaps for sets and measures*, Acta Math. 208 (2012), no. 1, 151-209.
- [16] A. Poltoratski, *A problem on completeness of exponentials*, Ann. of Math. (2) 178 (2013), no. 3, 983-1016.
- [17] F.J. Price and A. Sitaram, *Functions and their Fourier transforms with supports of finite measure for certain locally compact groups*, J. Funct. Anal. 79 (1988), no. 1, 166-182.
- [18] R. Rawat, A. Sitaram, *Injectivity sets for spherical means on \mathbb{R}^n and on symmetric spaces*, Jour. Four. Anal. Appl., 6 (2000), 343-348.
- [19] A. Sitaram, M. Sundari and S. Thangavelu, *Uncertainty principles on certain Lie groups*, Proc. Indian Acad. Sci. Math. Sci. 105 (1995), no. 2, 135-151.
- [20] P. Sjölin, *Heisenberg uniqueness pairs and a theorem of Beurling and Malliavin*, Bull. Sc. Math., 135(2011), 125-133.
- [21] P. Sjölin, *Heisenberg uniqueness pairs for the parabola*. Jour. Four. Anal. Appl., 19(2013), 410-416.
- [22] R.K. Srivastava, *Non-harmonic cones are Heisenberg uniqueness pairs for the Fourier transform on \mathbb{R}^n* , DOI: [arXiv:1507.02624](https://arxiv.org/abs/1507.02624).
- [23] F.J.G. Vieli, *A uniqueness result for the Fourier transform of measures on the sphere*, Bull. Aust. Math. Soc. 86(2012), 78-82.
- [24] F.J.G. Vieli, *A uniqueness result for the Fourier transform of measures on the paraboloid*, Matematicki Vesnik, Vol. 67(2015), No. 1, 52-55.

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY, GUWAHATI, INDIA 781039.

E-mail address: deb.giri@iitg.ac.in, rksri@iitg.ac.in