

Storage enhanced nonlinearities in a cold atomic Rydberg ensemble

E. Distante,^{1,*} A. Padrón-Brito,¹ M. Cristiani,¹ D. Paredes-Barato,¹ and H. de Riedmatten^{1,2}

¹*ICFO-Institut de Ciències Fotoniques, The Barcelona Institute of Science and Technology, 08860 Castelldefels (Barcelona), Spain*

²*ICREA-Institució Catalana de Recerca i Estudis Avançats, 08015 Barcelona, Spain*
(Dated: November 10, 2018)

The combination of electromagnetically induced transparency (EIT) with the nonlinear interaction between Rydberg atoms provides an effective interaction between photons. In this paper, we investigate the storage of optical pulses as collective Rydberg atomic excitations in a cold atomic ensemble. By measuring the dynamics of the stored Rydberg polaritons, we experimentally demonstrate that storing a probe pulse as Rydberg polaritons strongly enhances the Rydberg mediated interaction compared to the slow propagation case. We show that the process is characterized by two time scales. At short storage times, we observe a strong enhancement of the interaction due to the reduction of the Rydberg polariton group velocity down to zero. For longer storage times, we observe a further, weaker enhancement dominated by Rydberg induced dephasing of the multiparticle components of the state. In this regime, we observe a non-linear dependence of the Rydberg polariton coherence time with the input photon number. Our results have direct consequences in Rydberg quantum optics and enable the test of new theories of strongly interacting Rydberg systems.

PACS numbers: 32.80.Ee, 42.50.Nn, 42.50.Gy

The possibility to control the interaction between photons provided by highly nonlinear media is a key ingredient to the goal of quantum information processing (QIP) using photons and a unique tool to study the dynamics of many-body correlated systems [1]. Many different systems showing high nonlinear optical response at the single-photon level have been studied during the past years ranging from resonators coupled to single atoms [2–6], atomic ensembles [7], to artificial two-level atoms [8, 9].

A promising strategy to perform different QIP tasks using photons as carriers is the combination of electromagnetically induced transparency (EIT) [10–13] and Rydberg atoms [14] (see for example [15–34]). Using EIT one maps the state of the photons into atomic coherence in the form of Rydberg dark-state polaritons (DSPs) by means of an auxiliary coupling field. The strong Rydberg dipole-dipole (DD) interaction between neighboring excitations shifts the multiply-excited states from being resonantly coupled when these excitations are closer than a certain length called the *blockade radius*, r_b [25]. This way, only a single excitation can be created inside of a blockaded volume of the atomic cloud (so-called *super-atom*). This phenomenon, known as *Rydberg blockade*, has been used in combination with EIT to generate quantum states of light [27–29], single-photon switches and transistors [30–32] as well as a π phase shift controlled with single-photon level pulse [34]. These experiments typically require very high atomic densities and high-lying Rydberg states. By switching off and back on the coupling field, photons can be stored as Rydberg excitations and retrieved at later time [26, 29]. In this case the DD interaction dephases the collective emission of the multiparticle components of the stored photonic states [35, 36]. This feature was used to implement a determin-

istic single-photon source [26].

The key point of all these experiments is the strong nonlinear response arising from the DD interaction between high-lying Rydberg states. In the present paper, we experimentally demonstrate that storing the input photons as Rydberg excitations strongly enhances the nonlinear interaction when compared to the propagation case. The profound difference between propagating and storing Rydberg DSPs and its application in many-body Rydberg physics and QIP has been recently theoretically discussed [35, 37]. We show experimentally that the underlying many-body dynamics of strongly interacting DSPs during storage is characterized by two different time scales. A strong enhancement of the interaction happens at time scales shorter than what can be measured in this experiment. At longer time scales, the dynamics is dominated by the dephasing of multiparticle components of the input states. We confirm the latter by measuring for the first time the nonlinear dependence of the coherence time of stored Rydberg DSPs with respect to the input photon number [36]. Our results have a direct consequence in Rydberg quantum optics, demonstrating that the regime of strongly interacting DSPs required in most of the protocols can be achieved by storing the light even for a very short time. Moreover our experiment is a step forward in understanding the complicated many-body physics of the strongly interacting DSPs during storage [37].

Our measurement can be summarized as follows: we send coherent probe pulses with varying intensity and measure the number of emerging photons in the slow-light and in the storage case. The Rydberg DD interaction causes a non-linear input-output intensity relation, eventually leading to the saturation of the output photon number. Stronger, non-linear interactions lead to a




FIG. 1. (a) Counterpropagating probe (red) and coupling (blue) beams are focused using aspheric lenses onto a cold cloud of ^{87}Rb atoms. Probe and coupling beams are combined and separated using dichroic mirrors. Probe photons are detected using a fiber coupled single-photon APD. (b) Simplified atomic level scheme. (c) Probe transmission traces when coupling beam is off (red triangles) and on, showing the typical EIT transparency window (blue circles). Solid lines are fits to the data [38]. (d) Normalized and background subtracted counts of an input Gaussian probe (red area) when propagating as slow-light DSP (black line) and when stored for $t_s = 600$ ns (orange area). Here the input photon number is $N_{\text{in}} = 23.2 \pm 1.2$, with efficiencies $\eta = 0.336 \pm 0.006$ and $\eta = 0.078 \pm 0.002$ for the slow- and stored light respectively. Dashed orange line represents the leaked pulse during the storage process. (e) N_{out} normalized by the linear process efficiency T as a function of the input photon number N_{in} for the slow-light case (black triangles) and for two storage times. Solid curves represent fits with Eq.(1). Straight lines represent the linear behavior $N_{\text{out}}/T = N_{\text{in}}$ (oblique) and the saturation level N_{max} (horizontal). The Rydberg state used in these plots is $|70S_{1/2}\rangle$.

reduction of the maximum number of photons sustained by the medium [31]. In a first experiment, we measure N_{max} , the saturation plateau normalized by the linear process efficiency, T . We show that storage leads to a strong suppression of N_{max} compared to the slow-light case, thus demonstrating a strong enhancement of the Rydberg mediated photon interaction. The dependence of N_{max} on the storage time t_s shows that strong suppression happens at a short time scale. In a second experiment, we measure the coherence time of the storage process as a function of the input photon number. We show that higher intensity input fields suffer from stronger dephasing due to the Rydberg DD interaction.

Experiment In Fig.1 a schematic of this system is shown. We probe a cold cloud of ^{87}Rb atoms using 780 nm light ($\mathcal{E}$) with a detuning δ with respect to the $|g\rangle \leftrightarrow |e\rangle$ transition, where $|g\rangle = |5S_{1/2}, F=2\rangle$ and $|e\rangle = |5P_{3/2}, F=2\rangle$. The atomic sample is obtained using a magneto-optical trap, which generates a cloud of size $\sigma \sim 0.8$ mm with a peak density $\rho_0 = 3.2 \cdot 10^{10} \text{ cm}^{-3}$, and a temperature $T \sim 87.5 \mu\text{K}$ (measured by fluorescence imaging). A strong coupling field at 480 nm light is sent counterpropagating with respect to the probe. Using an excited-state locking scheme [39], we lock the

coupling beam resonantly to the $|e\rangle \leftrightarrow |r\rangle = |nS_{1/2}\rangle$ transition, where n is the principal quantum number. The probe and coupling laser fields are focused to waist radii $(w_p, w_c) \approx (7 \mu\text{m}, 13 \mu\text{m})$. This geometry gives $\approx 3.9 \times 10^4$ atoms in the interacting region. The optical depth (OD) of the cloud and the coupling Rabi frequency Ω_c are extracted by fitting the transmission of the probe as a function of the probe detuning δ with respect to the $|5S_{1/2}, F=2\rangle \leftrightarrow |5P_{3/2}, F=2\rangle$ transition using the model presented in [38]. We set them to be $\text{OD} \sim 6.2$ and $\Omega_c = (4.38 \pm 0.04) \text{ MHz}$ (see Fig.1(c)). The probe and the coupling beam are opposite circularly polarized [40] and the magnetic field is set to zero at the position of the cloud [41].

When $\delta = 0$ the presence of the coupling field converts the probe photons into propagating Rydberg DSPs [11–13]. These DSPs travel at reduced group velocity $v_g \sim |\Omega_c|^2 / (g^2 \rho_0)$, where g is the coupling strength between the probe and the $|g\rangle \leftrightarrow |e\rangle$ transition [11, 42]. By adiabatically switching off the coupling beam we store the state of the input field as an atomic coherence. The stored excitation is retrieved after a storage time t_s by switching the coupling beam back on.

We send a Gaussian, coherent probe pulse with a dura-




FIG. 2. Normalized efficiency η/T as a function of the input photon number N_{in} (a) For fixed storage time $t_s = 400$ ns comparison between non-interacting low-lying Rydberg state $26S_{1/2}$ (black triangles) with stronger interacting, higher n Rydberg states. (b) For fixed Rydberg state $|70S_{1/2}\rangle$, comparison between slow-light case (black triangles) with and the storage case for different storage times. In both plots, lines are fits with the Eq.(1).

tion of 410 ns (FWHM_{in}) and average number of photons N_{in} through the cold atomic gas. The light is detected after the ensemble with a single-photon APD and the counts are background subtracted and corrected for detection efficiency. Initially we calibrate N_{in} by measuring the transmitted pulse without loading the atoms. Then we measure N_{out} either when the probe pulse propagates as slow Rydberg DSPs or when they are stored in the $|nS_{1/2}\rangle$ state (see Fig.1(e)). In the absence of Rydberg interaction, the average number of photons N_{out} in the emerging pulse increases linearly with N_{in} , $N_{\text{out}} = TN_{\text{in}}$. Here $T < 1$ represents imperfect process efficiency. During slow-light propagation, this is caused by the decoherence of the ground Rydberg transition, which includes the natural lifetime of the Rydberg state, atomic collisions, coupling with external fields and laser linewidth. The storage process efficiency is further limited by imperfect pulse compression inside the medium (due to low OD) and by the dephasing of the collective state during the storage time, which is dominated by coupling with external fields and atomic motion. When the number of input photons is increased a nonlinear dependence arises, eventually leading to saturation of N_{out} . To quantify the effective interaction we fit our data with the model proposed in [31]. In that model the input-output relation is




FIG. 3. Maximum number of retrieved photons normalized by the process efficiency N_{max} (extracted from the fit with Eq.(1)) as a function of the storage time t_s for different Rydberg states. Comparison with the slow-light case (points at $t_s = 0$ ns) shows a strong reduction of N_{max} when storage is performed. Dotted lines are exponential fits to the storage data. (Inset) N_{max} as a function of the principal quantum number n for slow-light. Solid line is fit with the function $N_{\text{max}} = \alpha n^{-\gamma}$ giving $\gamma = 5.3 \pm 0.2$ (see main text).

described by:

$$N_{\text{out}} = N_{\text{max}}T(1 - e^{-N_{\text{in}}/N_{\text{max}}}) \quad (1)$$

where T represents the linear efficiency of the process at low photon number and N_{max} is the maximum number of photons that can emerge from the medium when unitary efficiency $T = 1$ is considered. As explained in [31], N_{max} decreases for stronger Rydberg interaction and can be used to quantify the effective blockade of the output field. Fig.1(e) reports an example of the data for $n = 70$ together with the fit using Eq.(1).

In Fig.2(b) a re-scaled efficiency η/T (being $\eta = N_{\text{out}}/N_{\text{in}}$) is shown for the high-lying state $|70S_{1/2}\rangle$ at different storage times. The data show that η/T tails off at lower N_{in} for longer storage times as a consequence of stronger nonlinearity. Similar data have been taken for a variety of Rydberg states (see e.g. Fig.2(a) for the results with $t_s = 400$ ns) and the fit results are shown in Fig.3. One could argue that saturation may arise as a result of medium saturation, when the density of photons and atoms inside the medium are comparable. However, in Fig.2(a) we observe that the response of the medium is linear (that is, the efficiency does not depend on N_{in}) at low-lying Rydberg states, where the interaction is negligible.

In Fig.3 we show N_{max} for different Rydberg levels, both in the slow-light or in the storage case. As expected, when n is increased N_{max} is reduced, due to stronger Rydberg interaction. In the propagation case this can be understood as a consequence of the blockade effect. When the density of photons in the medium becomes comparable to the density of *super-atoms* $\rho_{\text{SA}} = 3/4\pi r_b^3$, the medium saturates [43]. Since $r_b \sim n^{11/6}$, this condition

is reached at a lower number of photons for a higher Rydberg state. Following this, the maximum number of photons supported by the medium scales as $N_{\max} \sim n^{-11/2}$. The inset in Fig.3 shows a fit for the slow-light case with the function $N_{\max} = \alpha n^{-\gamma}$ which gives $\gamma = 5.3 \pm 0.2$.

When the probe pulse is stored, saturation occurs at a lower number of photons (an order of magnitude difference for $t_{\text{st}} = 2 \mu\text{s}$), as shown in Fig.3. Data show that $N_{\max}$ is strongly reduced soon after storage is performed. The two time scales of the process are evident when noticing that even an exponential fit of $N_{\max}$ in the storage case (dotted lines in Fig.3) fails to include the data of the slow-light case, represented by the $t_s = 0$ ns points in Fig.3. In the ideal limit of zero decoherence between the ground and the Rydberg state, the blockade radius increases without bounds when Ω_c goes to zero, according to the naive formula for the blockade radius $r_b = \sqrt[6]{C_6/\delta_{\text{EIT}}}$ in terms of the EIT bandwidth $\delta_{\text{EIT}} \propto \Omega_c^2$. This is not consistent with our data nor with other experimental results [29, 44]. A recent description by Moos *et al.* [37] shows that Ω_c has to be replaced with $\Omega_{\text{eff}}^2 = g^2 \rho_0 + \Omega_c^2$ upon storage. According to this new description, the blockade radius during storage becomes $r_b = \sqrt[6]{C_6 \Gamma / g^2 \rho_0}$. As a consequence, the blockaded volume would not increase significantly during the storage process and it could not be used to understand the data. Nevertheless Moos *et al.* suggest that the strongly interacting regime is achievable when the ratio between the Rydberg interaction and the kinetic energy of the DSP is strongly increased; this regime is achieved during the storage process when v_g is reduced to zero. This theory also suggests other specific effects (such as a quasicrystalline density of stored photons) which are interesting but not within the reach of our current setup.

At longer time scales, the Rydberg DD interaction acts as an extra source of dephasing for the many-body components of the stored DSP effectively blocking the collective emission of such components in the retrieved mode. This effect [35] has been observed before and it has been exploited to generate single photons deterministically [26, 36]. Here we show the first detailed time-dependent study. We measure the storage efficiency η versus the storage time t_s for a variety of input photon numbers N_{in} . The inset of Fig.4 reports an example of the efficiency data for the $|70S_{1/2}\rangle$ state for two different N_{in} . The results are summarized in Fig.4 where we show how the $1/e$ coherence time τ depends on N_{in} for two different Rydberg states $|60S_{1/2}\rangle$ and $|70S_{1/2}\rangle$ [45]. At low photon numbers, we observe larger dephasing at higher principal quantum number. At higher N_{in} , the interaction between Rydberg states introduces another source of dephasing, resulting in a reduction of τ . Both Rydberg states show similar dependence of τ (when normalized at low photon numbers) with respect to N_{in} . At first surprising, this result can be understood by noticing that the system starts to evolve from




FIG. 4. (Inset) Storage efficiency η as a function of storage time t_s for input photon number $N_{\text{in}} \sim 90$ (squares) and $N_{\text{in}} \sim 1730$ (diamonds). Solid lines represent fits using an exponential function. (Main Plot) Coherence time τ extracted from the fit as a function of the input photon number N_{in} for Rydberg levels $n = 60, 70$ (empty triangles and circles respectively). Filled squares and diamonds in the $n = 70$ set represent the fit of the data shown in the inset.

a partially blockaded configuration, contrary to the situation studied in [26, 35]. Following theory presented in [35], the interaction between Rydberg states induces a phase shift on the m -body component of the storage state $\phi_{\mu_1 \dots \mu_m} = -t \sum_{1 \leq i < j \leq m} V_{\mu_i \mu_j} / \hbar$. Here $V_{\mu_i \mu_j}$ is the Van der Waals potential describing the interaction between two Rydberg excitations μ_i and μ_j , which is strongly state dependent. Nevertheless, due to the blockade effect, two excitations cannot be closer than r_b . At this distance, the dipole potential is fixed by the EIT linewidth: $V(r_b) = \hbar \delta_{\text{EIT}}$. Since δ_{EIT} is similar in the two Rydberg states considered in our experiment, we expect both states to present similar dephasing rates.

Conclusions We have performed the first extensive measurement of the dynamics of stored Rydberg DSPs. Our data clearly demonstrate that storing photons as a Rydberg DSP enhances the Rydberg mediated interaction when compared to the slowly propagating case. This result may open the door to the obtention of strong photon-photon interactions at moderate atomic densities and lower Rydberg states. This would facilitate photonic QIP using Rydberg atoms by relaxing the strong requirements of high densities [46, 47] and high Rydberg levels. We have discussed the many-body dynamics of the process, showing that two different time scales are present. We suggest that a recent theory proposed by Moos *et al.* in [37] might explain our results at short time scales. At long time scales, we have presented the first time-dependent measurement of the dephasing induced by the Rydberg DD interaction and we have shown its clear dependence on the input photon number. In the future, our data might allow to test more detailed models of inter-

acting Rydberg DSPs, casting light on the strongly interacting many-body physics with Rydberg atoms. On the experimental side, reducing the lasers linewidth and the cloud temperature would enable the study of the dephasing at longer storage times and at higher Rydberg levels. Finally, these results can be extended to show nonlinearities at the single-photon level by increasing the density of the cloud and by reducing the size of the sample.

We thank M. Fleischhauer, M. Moos, R. Unanyan, C. Simon and M. Khazali for useful discussions. DPB has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 658258. We acknowledge financial support by the ERC starting grant QuLIMA, by the Spanish Ministry of Economy and Competitiveness (MINECO) Severo Ochoa through grant SEV-2015-0522, by AGAUR via 2014 SGR 1554 and by Fundació Privada Cellex.

* emanuele.distante@icfo.es

- [1] D. E. Chang, V. Vuletić, and M. D. Lukin, *Nature Photonics* **8**, 685 (2014).
- [2] B. Dayan, A. S. Parkins, T. Aoki, E. P. Ostby, K. J. Vahala, and H. J. Kimble, *Science* **319**, 22 (2008).
- [3] A. Reiserer, S. Ritter, and G. Rempe, *Science* **342**, 1349 (2013), 1311.3625.
- [4] A. Reiserer, N. Kalb, G. Rempe, and S. Ritter, *Nature* **508**, 237 (2014), 1404.2453.
- [5] T. G. Tiecke, J. D. Thompson, N. P. de Leon, L. R. Liu, V. Vuletić, and M. D. Lukin, *Nature* **508**, 241 (2014).
- [6] I. Shomroni, S. Rosenblum, Y. Lovsky, O. Bechler, G. Guendelman, and B. Dayan, *Science* **345**, 903 (2014).
- [7] W. Chen, K. M. Beck, R. Bücker, M. Gullans, M. D. Lukin, H. Tanji-Suzuki, and V. Vuletić, *Science* **341**, 768 (2013).
- [8] I. Fushman, D. Englund, A. Faraon, N. Stoltz, P. Petroff, and J. Vuckovic, *Science* **320**, 769 (2008).
- [9] T. Volz, A. Reinhard, M. Winger, A. Badolato, K. J. Hennessy, E. L. Hu, and A. Imamoglu, *Nature Photonics* **6**, 607 (2012), 1111.2915.
- [10] L. V. Hau, S. E. Harris, Z. Dutton, and C. H. Behroozi, *Nature* **397**, 594 (1999).
- [11] M. Fleischhauer and M. D. Lukin, *Phys. Rev. Lett.* **84**, 5094 (2000).
- [12] M. Fleischhauer and M. D. Lukin, *Phys. Rev. A* **65**, 022314 (2002).
- [13] M. Fleischhauer, A. Imamoglu, and J. P. Marangos, *Reviews of Modern Physics* **77**, 633 (2005).
- [14] T. Gallagher, *Rydberg Atoms* (Cambridge University Press, 2005).
- [15] I. Friedler, D. Petrosyan, M. Fleischhauer, and G. Kurizki, *Phys. Rev. A* **72**, 043803 (2005).
- [16] E. Shahmoon, G. Kurizki, M. Fleischhauer, and D. Petrosyan, *Phys. Rev. A* **83**, 033806 (2011).
- [17] A. V. Gorshkov, J. Otterbach, E. Demler, M. Fleischhauer, and M. D. Lukin, *Phys. Rev. Lett.* **105**, 060502 (2010).
- [18] A. V. Gorshkov, J. Otterbach, M. Fleischhauer, T. Pohl, and M. D. Lukin, *Phys. Rev. Lett.* **107**, 133602 (2011).
- [19] B. He, A. MacRae, Y. Han, A. I. Lvovsky, and C. Simon, *Phys. Rev. A* **83**, 022312 (2011).
- [20] A. V. Gorshkov, R. Nath, and T. Pohl, *Phys. Rev. Lett.* **110**, 153601 (2013).
- [21] D. Paredes-Barato and C. S. Adams, *Phys. Rev. Lett.* **112**, 040501 (2014).
- [22] B. He, A. V. Sharypov, J. Sheng, C. Simon, and M. Xiao, *Phys. Rev. Lett.* **112**, 133606 (2014).
- [23] M. Khazali, K. Heshami, and C. Simon, *Phys. Rev. A* **91**, 030301 (2015).
- [24] S. Das, A. Grankin, I. Iakoupov, E. Brion, J. Borregaard, R. Boddeda, I. Usmani, A. Ourjoumtsev, P. Grangier, and A. S. Sørensen, *arXiv:1506.04300 [quant-ph]* (2015).
- [25] J. D. Pritchard, D. Maxwell, A. Gauguier, K. J. Weatherill, M. P. A. Jones, and C. S. Adams, *Phys. Rev. Lett.* **105**, 193603 (2010).
- [26] Y. O. Dudin and A. Kuzmich, *Science* **336**, 887 (2012).
- [27] T. Peyronel, O. Firstenberg, Q. Y. Liang, S. Hofferberth, A. V. Gorshkov, T. Pohl, M. D. Lukin, and V. Vuletić, *Nature (London)* **488**, 57 (2012).
- [28] O. Firstenberg, T. Peyronel, Q.-Y. Liang, A. V. Gorshkov, M. D. Lukin, and V. Vuletić, *Nature* **502**, 71 (2013).
- [29] D. Maxwell, D. J. Szwer, D. Paredes-Barato, H. Busche, J. D. Pritchard, A. Gauguier, K. J. Weatherill, M. P. A. Jones, and C. S. Adams, *Phys. Rev. Lett.* **110**, 103001 (2013).
- [30] D. Tiarks, S. Baur, K. Schneider, S. Dürr, and G. Rempe, *Phys. Rev. Lett.* **113**, 053602 (2014).
- [31] S. Baur, D. Tiarks, G. Rempe, and S. Dürr, *Phys. Rev. Lett.* **112**, 073901 (2014).
- [32] H. Gorniaczyk, C. Tresp, J. Schmidt, H. Fedder, and S. Hofferberth, *Phys. Rev. Lett.* **113**, 053601 (2014).
- [33] R. Boddeda, I. Usmani, E. Bimbard, A. Grankin, A. Ourjoumtsev, E. Brion, and P. Grangier, *arXiv:1512.08480* (2015), arXiv: 1512.08480.
- [34] D. Tiarks, S. Schmidt, G. Rempe, and S. Dürr, *arXiv:1512.05740 [quant-ph]* (2015).
- [35] F. Bariani, Y. O. Dudin, T. A. B. Kennedy, and A. Kuzmich, *Phys. Rev. Lett.* **108**, 030501 (2012).
- [36] F. Bariani, P. M. Goldbart, and T. A. B. Kennedy, *Phys. Rev. A* **86**, 041802 (2012).
- [37] M. Moos, M. Höning, R. Unanyan, and M. Fleischhauer, *Phys. Rev. A* **92**, 053846 (2015).
- [38] M. Xiao, Y.-Q. Li, S.-Z. Jin, and J. Gea-Banacloche, *Phys. Rev. Lett.* **74**, 666 (1995).
- [39] R. P. Abel, A. K. Mohapatra, M. G. Bason, J. D. Pritchard, K. J. Weatherill, U. Raitzsch, and C. S. Adams, *Appl. Phys. Lett.* **94**, 071107 (2009).
- [40] R. Löw, H. Weimer, J. Nipper, J. B. Balewski, B. Butscher, H. P. Büchler, and T. Pfau, *J. Phys. B* **45**, 113001 (2012).
- [41] B. Albrecht, *Ph.D. thesis*, ICFO - The Institute of Photonics Sciences, Barcelona (2015), see chapter 4.1.3.
- [42] C. Liu, Z. Dutton, C. H. Behroozi, and L. V. Hau, *Nature* **409**, 490 (2001).
- [43] D. Petrosyan, J. Otterbach, and M. Fleischhauer, *Phys. Rev. Lett.* **107**, 213601 (2011).
- [44] D. Maxwell, D. J. Szwer, D. Paredes-Barato, H. Busche, J. D. Pritchard, A. Gauguier, M. P. A. Jones, and C. S. Adams, *Phys. Rev. A* **89**, 043827 (2014).

- [45] In state $|70S_{1/2}\rangle$ the function used is $\eta = \eta_0 \exp(-t/\tau)$, but for $|60S_{1/2}\rangle$ we have to use the function $\eta = \eta_0 \exp(-t/\tau) |p + (1-p) \exp(-i2\pi\Delta_{hf}t)|^2$ to account for the hyperfine splitting of the state Δ_{hf} . Here p is the probability to excite the $F = 1$ state.
- [46] J. B. Balewski, A. T. Krupp, A. Gaj, S. Hofferberth, R. Löw, and T. Pfau, [New Journal of Physics](#) **16**, 063012 (2014).
- [47] O. Firstenberg, C. S. Adams, and S. Hofferberth, [arXiv:1602.06117](#) (2016).