

Non-commutative algebraic geometry of semi-graded rings

Oswaldo Lezama

jolezamas@unal.edu.co

Edward Latorre

Seminario de Álgebra Constructiva - SAC²

Departamento de Matemáticas

Universidad Nacional de Colombia, Sede Bogotá

Abstract

In this paper we introduce the semi-graded rings, which extend graded rings and skew PBW extensions. For this new type of non-commutative rings we will discuss some basic problems of non-commutative algebraic geometry. In particular, we will prove some elementary properties of the generalized Hilbert series, Hilbert polynomial and Gelfand-Kirillov dimension. We will extend the notion of non-commutative projective scheme to the case of semi-graded rings and we generalize the Serre-Artin-Zhang-Verevkin theorem. Some examples are included at the end of the paper.

Key words and phrases. Non-commutative algebraic geometry, graded rings and modules, Hilbert series and Hilbert polynomial, Gelfand-Kirillov dimension, non-commutative schemes, skew *PBW* extensions.

2010 *Mathematics Subject Classification.* Primary: 16S38. Secondary: 16W50, 16S80, 16S36.

1 Introduction

Finitely graded algebras and skew *PBW* extensions cover many important classes of non-commutative rings and algebras coming from quantum mechanics. For example, quantum polynomials are examples of graded algebras, and universal enveloping algebras of finite-dimensional Lie algebras are examples of skew *PBW* (Poincaré-Birkhoff-Witt) extensions (see [2], [3], [9], [15] and [13]). There exists recent interest in developing the non-commutative projective algebraic geometry for finitely graded algebras (see [8], [11] and [15]). However, for non \mathbb{N} -graded algebras, in particular, for many important classes of skew *PBW* extensions, only few works have been realized (see [6]). In this paper we present an introduction to non-commutative algebraic geometry for non \mathbb{N} -graded algebras and rings, defining a new class of rings: the *semi-graded rings*. As we will see, the semi-graded rings generalize the finitely graded algebras and the skew *PBW* extensions. We will discuss the most basic problems on non-commutative algebraic geometry for semi-graded rings. The problems to be discussed are around the following topics: generalized Hilbert series and Hilbert polynomial, generalized Gelfand-Kirillov dimension, non-commutative schemes associated to semi-graded rings, and we will extend the Serre-Artin-Zhang-Verevkin theorem on projective schemes to semi-graded rings satisfying some natural restrictions.

In the present section we will recall the definition of finitely graded algebras and the Serre-Artin-Zhang-Verevkin theorem (see [3], [15], [17], [18]). We will also recall the definition of skew *PBW* extensions and some important examples of this type of non-commutative rings of polynomial type. In the second section we introduce the semi-graded rings and modules and we will prove some elementary properties of

them. The third and fourth sections are dedicated to give a generalization of the Hilbert series, Hilbert polynomial and the classical Gelfand-Kirillov dimension. These three notions are very important in any approach to non-commutative algebraic geometry. The purpose of the fourth section is to extend the notion of non-commutative projective scheme to the case of semi-graded rings. In the last section we present a generalization of the Serre-Artin-Zhang-Verevkin theorem about non-commutative schemes of finitely graded algebras to the case of semi-graded rings of special type. This main result can be applied to study the non-commutative algebraic geometry of some important particular examples of quantum algebras, probably not considered before in the literature.

Definition 1.1. *Let K be a field. It is said that a K -algebra A is finitely graded if the following conditions hold:*

- A is \mathbb{N} -graded: $A = \bigoplus_{n \geq 0} A_n$.
- A is connected, i.e., $A_0 = K$.
- A is finitely generated as K -algebra.

The most remarkable examples of finitely graded algebras for which the non-commutative projective algebraic geometry has been developed are the quantum plane, the Jordan plane, the Sklyanin algebra and the quantum polynomial ring in several variables (see [15]). Next we review the most basic facts of the non-commutative algebraic geometry of finitely graded algebras following the approach given by M. Artin and J. J. Zhang in [3], by A. B. Verevkin in [17], [18], and by Rogalski in [15].

Let A be a finitely graded K -algebra and $M = \bigoplus_{n \in \mathbb{Z}} M_n$ be a \mathbb{Z} -graded A -module which is finitely generated. Then,

- (i) For every $n \in \mathbb{Z}$, $\dim_K M_n < \infty$.
- (ii) The *Hilbert series* of M is defined by

$$h_M(t) := \sum_{n \in \mathbb{Z}} (\dim_K M_n) t^n.$$

In particular,

$$h_A(t) := \sum_{n=0}^{\infty} (\dim_K A_n) t^n.$$

- (iii) It says that A has *Hilbert polynomial* if there exists a polynomial $p_A(t) \in \mathbb{Q}[t]$ such that for all n sufficiently large, $p_A(n) = \dim_K A_n$. In this case $p_A(t)$ is called the Hilbert polynomial of A . Thus,

$$\dim_K A_n = p_A(n), \text{ for all } n \gg 0.$$

- (iv) The *Gelfand-Kirillov dimension* of A is defined by

$$\text{GKdim}(A) := \sup_V \overline{\lim}_{n \rightarrow \infty} \log_n \dim_K V^n, \tag{1.1}$$

where V ranges over all frames of A and $V^n := {}_K \langle v_1 \cdots v_n \mid v_i \in V \rangle$ (a *frame* of A is a finite dimensional K -subspace of A such that $1 \in V$; since A is a K -algebra, then $K \hookrightarrow A$, and hence, K is a frame of A of dimension 1).

- (v)

$$\text{GKdim}(A) = \overline{\lim}_{n \rightarrow \infty} \log_n \left(\sum_{i=0}^n \dim_K A_i \right).$$

(vi) If $p_A(t)$ exists, then

$$\text{GKdim}(A) = \deg(p_A(t)) + 1.$$

(vii) A famous Serre's theorem on commutative projective algebraic geometry states that the category of coherent sheaves over the projective n -space \mathbb{P}^n is equivalent to a category of noetherian graded modules over a graded commutative polynomial ring. The study of this equivalence for non-commutative finitely graded noetherian algebras is known as the non-commutative version of Serre's theorem and is due to Artin, Zhang and Vervekin. In the next numerals we give the ingredients needed for the formulation of this theorem (see [3], [17]).

(viii) Suppose that A is left noetherian. Let $\text{gr} - A$ be the abelian category of finitely generated \mathbb{Z} -graded left A -modules. It is defined the abelian category $\text{qgr} - A$ in the following way: The objects are the same as the objects in $\text{gr} - A$, and we let $\pi : \text{gr} - A \rightarrow \text{qgr} - A$ be the identity map on objects. The morphisms in $\text{qgr} - A$ are defined in the following way:

$$\text{Hom}_{\text{qgr} - A}(\pi(M), \pi(N)) := \varinjlim \text{Hom}_{\text{gr} - A}(M_{\geq n}, N/T(N)),$$

where the direct limit is taken over maps of abelian groups

$$\text{Hom}_{\text{gr} - A}(M_{\geq n}, N/T(N)) \rightarrow \text{Hom}_{\text{gr} - A}(M_{\geq n+1}, N/T(N))$$

induced by the inclusion homomorphism $M_{\geq n+1} \rightarrow M_{\geq n}$; $T(N)$ is the torsion submodule of N and an element $x \in N$ is torsion if $A_{\geq n}x = 0$ for some $n \geq 0$. The pair $(\text{qgr} - A, \pi(A))$ is called the *non-commutative projective scheme* associated to A , and denoted by $\text{qgr} - A$. Thus, $\text{qgr} - A$ is a quotient category, $\text{qgr} - A = \text{gr} - A / \text{tor} - A$.

(ix) (*Serre's theorem*) Let A be a commutative finitely graded K -algebra generated in degree 1. Then, there exists an equivalence of categories

$$\text{qgr} - A \simeq \text{coh}(\text{proj}(A)).$$

In particular,

$$\text{qgr} - K[x_0, \dots, x_n] \simeq \text{coh}(\mathbb{P}^n).$$

(x) Suppose that A is left noetherian; let $i \geq 0$; it is said that A satisfies the χ_i condition if for every finitely generated \mathbb{Z} -graded A -module M , $\dim_K(\text{Ext}_A^j(K, M)) < \infty$ for any $j \leq i$; the algebra A satisfies the χ condition if it satisfies the χ_i condition for all $i \geq 0$.

(xi) (*Artin-Zhang-Vervekin theorem*) If A is left noetherian and satisfies χ_1 , then $\pi(A)$ is left noetherian and there exists an equivalence of categories

$$\text{qgr} - A \simeq \text{qgr} - \Gamma(\pi(A))_{\geq 0},$$

where $\Gamma(\pi(A))_{\geq 0} := \bigoplus_{d=0}^{\infty} \text{Hom}_{\text{qgr} - A}(\pi(A), s^d(\pi(A)))$ and s is the autoequivalence of $\text{qgr} - A$ defined by the shifts of degrees.

Now we recall the definition of skew *PBW* extension defined firstly in [12]; many important algebras coming from mathematical physics are particular examples of skew *PBW* extensions: $\mathcal{U}(\mathcal{G})$, where \mathcal{G} is a finite dimensional Lie algebra, the algebra of q -differential operators, the algebra of shift operators, the additive analogue of the Weyl algebra, the multiplicative analogue of the Weyl algebra, the quantum algebra $\mathcal{U}'(\mathfrak{so}(3, K))$, 3-dimensional skew polynomial algebras, the dispin algebra, the Woronowicz algebra, the q -Heisenberg algebra, are particular examples of skew *PBW* extensions (see [13]).

Definition 1.2. Let R and A be rings. We say that A is a skew PBW extension of R (also called a σ -PBW extension of R) if the following conditions hold:

(i) $R \subseteq A$.

(ii) There exist finitely many elements $x_1, \dots, x_n \in A$ such A is a left R -free module with basis

$$\text{Mon}(A) := \{x^\alpha = x_1^{\alpha_1} \cdots x_n^{\alpha_n} \mid \alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n\}, \text{ with } \mathbb{N} := \{0, 1, 2, \dots\}.$$

The set $\text{Mon}(A)$ is called the set of standard monomials of A .

(iii) For every $1 \leq i \leq n$ and $r \in R - \{0\}$ there exists $c_{i,r} \in R - \{0\}$ such that

$$x_i r - c_{i,r} x_i \in R. \quad (1.2)$$

(iv) For every $1 \leq i, j \leq n$ there exists $c_{i,j} \in R - \{0\}$ such that

$$x_j x_i - c_{i,j} x_i x_j \in R + R x_1 + \cdots + R x_n. \quad (1.3)$$

Under these conditions we will write $A := \sigma(R)\langle x_1, \dots, x_n \rangle$.

Associated to a skew PBW extension $A = \sigma(R)\langle x_1, \dots, x_n \rangle$, there are n injective endomorphisms $\sigma_1, \dots, \sigma_n$ of R and σ_i -derivations, as the following proposition shows.

Proposition 1.3. Let A be a skew PBW extension of R . Then, for every $1 \leq i \leq n$, there exist an injective ring endomorphism $\sigma_i : R \rightarrow R$ and a σ_i -derivation $\delta_i : R \rightarrow R$ such that

$$x_i r = \sigma_i(r) x_i + \delta_i(r),$$

for each $r \in R$.

Proof. See [12], Proposition 3. □

A particular case of skew PBW extension is when all derivations δ_i are zero. Another interesting case is when all σ_i are bijective and the constants $c_{i,j}$ are invertible. We recall the following definition (cf. [12]).

Definition 1.4. Let A be a skew PBW extension.

(a) A is quasi-commutative if the conditions (iii) and (iv) in Definition 1.2 are replaced by

(iii') For every $1 \leq i \leq n$ and $r \in R - \{0\}$ there exists $c_{i,r} \in R - \{0\}$ such that

$$x_i r = c_{i,r} x_i. \quad (1.4)$$

(iv') For every $1 \leq i, j \leq n$ there exists $c_{i,j} \in R - \{0\}$ such that

$$x_j x_i = c_{i,j} x_i x_j. \quad (1.5)$$

(b) A is bijective if σ_i is bijective for every $1 \leq i \leq n$ and $c_{i,j}$ is invertible for any $1 \leq i < j \leq n$.

Observe that quasi-commutative skew PBW extensions are \mathbb{N} -graded rings but arbitrary skew PBW extensions are semi-graded rings as we will see below. Actually, the main motivation for constructing the non-commutative algebraic geometry of semi-graded rings is due to arbitrary skew PBW extensions.

Many properties of skew PBW extensions have been studied in previous works (see [1], [13]). For example, the global, Krull and Goldie dimensions of bijective skew PBW extensions were estimated in [13]. The next theorem establishes two classical ring theoretic results for skew PBW extensions.

Theorem 1.5. *Let A be a bijective skew PBW extension of a ring R .*

- (i) (*Hilbert Basis Theorem*) *If R is a left (right) Noetherian ring then A is also left (right) Noetherian.*
- (ii) (*Ore's theorem*) *If R is a left Ore domain R . Then A is also a left Ore domain.*

We conclude this introductory section fixing some notation: If not otherwise noted, all modules are left modules; B will denote a non-commutative ring; K will be a field; $A := \sigma(R)\langle x_1, \dots, x_n \rangle$ will represent a skew PBW extension.

2 Semi-graded rings and modules

In this section we introduce the semi-graded rings and modules, we prove some elementary properties of them, and we will show that graded rings, finitely graded algebras and skew PBW extensions are particular cases of this new type of non-commutative rings.

Definition 2.1. *Let B be a ring. We say that B is semi-graded (SG) if there exists a collection $\{B_n\}_{n \geq 0}$ of subgroups B_n of the additive group B^+ such that the following conditions hold:*

- (i) $B = \bigoplus_{n \geq 0} B_n$.
- (ii) For every $m, n \geq 0$, $B_m B_n \subseteq B_0 \oplus \dots \oplus B_{m+n}$.
- (iii) $1 \in B_0$.

The collection $\{B_n\}_{n \geq 0}$ is called a semi-graduation of B and we say that the elements of B_n are homogeneous of degree n . Let B and C be semi-graded rings and let $f : B \rightarrow C$ be a ring homomorphism, we say that f is homogeneous if $f(B_n) \subseteq C_n$ for every $n \geq 0$.

Definition 2.2. *Let B be a SG ring and let M be a B -module. We say that M is a \mathbb{Z} -semi-graded, or simply semi-graded, if there exists a collection $\{M_n\}_{n \in \mathbb{Z}}$ of subgroups M_n of the additive group M^+ such that the following conditions hold:*

- (i) $M = \bigoplus_{n \in \mathbb{Z}} M_n$.
- (ii) For every $m \geq 0$ and $n \in \mathbb{Z}$, $B_m M_n \subseteq \bigoplus_{k \leq m+n} M_k$.

We say that M is positively semi-graded, also called \mathbb{N} -semi-graded, if $M_n = 0$ for every $n < 0$. Let $f : M \rightarrow N$ be an homomorphism of B -modules, where M and N are semi-graded B -modules; we say that f is homogeneous if $f(M_n) \subseteq N_n$ for every $n \in \mathbb{Z}$.

As for the case of rings, the collection $\{M_n\}_{n \in \mathbb{Z}}$ is called a semi-graduation of M and we say that the elements of M_n are homogeneous of degree n .

Let B be a semi-graded ring and let M be a semi-graded B -module, let N be a submodule of M , let $N_n := N \cap M_n$, $n \in \mathbb{Z}$; observe that the sum $\sum_n N_n$ is direct. This induces the following definition.

Definition 2.3. *Let B be a SG ring and M be a semi-graded module over B . Let N be a submodule of M , we say that N is a semi-graded submodule of M if $N = \bigoplus_{n \in \mathbb{Z}} N_n$.*

Note that if N is semi-graded, then $B_m N_n \subseteq \bigoplus_{k \leq m+n} N_k$, for every $n \in \mathbb{Z}$ and $m \geq 0$: In fact, let $b \in B_m$ and $z \in N_n$, then $bz \in B_m M_n \subseteq \bigoplus_{k \leq m+n} M_k$ and $bz = z_1 + \dots + z_l$, with $z_i \in N_{n_i} \subseteq M_{n_i}$, but since the sum is direct, then $n_i \leq m + n$ for every $1 \leq i \leq l$.

Finally, we introduce an important class of semi-graded rings that includes finitely graded algebras and skew PBW extensions.

Definition 2.4. Let B be a ring. We say that B is finitely semi-graded (FSG) if B satisfies the following conditions:

- (i) B is SG.
- (ii) There exists finitely many elements $x_1, \dots, x_n \in B$ such that the subring generated by B_0 and x_1, \dots, x_n coincides with B .
- (iii) For every $n \geq 0$, B_n is a free B_0 -module of finite dimension.

Moreover, if M is a B -module, we say that M is finitely semi-graded if M is semi-graded, finitely generated, and for every $n \in \mathbb{Z}$, M_n is a free B_0 -module of finite dimension.

Remark 2.5. Observe if B is FSG, then $B_0 B_p = B_p$ for every $p \geq 0$, and if M is finitely semi-graded, then $B_0 M_n = M_n$ for all $n \in \mathbb{Z}$.

From the definitions above we get the following conclusions.

Proposition 2.6. Let $B = \bigoplus_{n \geq 0} B_n$ be a SG ring and I be a proper two-sided ideal of B semi-graded as left ideal. Then,

- (i) B_0 is a subring of B . Moreover, for any $n \geq 0$, $B_0 \oplus \dots \oplus B_n$ is a $B_0 - B_0$ -bimodule, as well as B .
- (ii) B has a standard \mathbb{N} -filtration given by

$$F_n(B) := B_0 \oplus \dots \oplus B_n. \quad (2.1)$$

- (iii) The associated graded ring $Gr(B)$ satisfies

$$Gr(B)_n \cong B_n, \text{ for every } n \geq 0 \text{ (isomorphism of abelian groups).}$$

- (iv) Let $M = \bigoplus_{n \in \mathbb{Z}} M_n$ be a semi-graded B -module and N a submodule of M . The following conditions are equivalent:

- (a) N is semi-graded.
- (b) For every $z \in N$, the homogeneous components of z are in N .
- (c) M/N is semi-graded with semi-graduation given by

$$(M/N)_n := (M_n + N)/N, \quad n \in \mathbb{Z}.$$

- (v) B/I is SG.

- (vi) If B is FSG and $I \cap B_n \subseteq IB_n$ for every n , then B/I is FSG.

Proof. (i) and (ii) are obvious. For (iii) observe that $Gr(B)_n = F_n(B)/F_{n-1}(B) \cong B_n$ for every $n \geq 0$ (isomorphism of abelian groups); in addition, note how acts the product: let $z := \overline{b_0 + \dots + b_m} \in Gr(B)_m$, $z' := \overline{c_0 + \dots + c_n} \in Gr(B)_n$, then

$$zz' = \overline{b_m c_n} = \overline{d_0 + \dots + d_{m+n}} = \overline{d_{m+n}} \in Gr(B)_{m+n} \cong B_{m+n}.$$

- (iv) (a) \Leftrightarrow (b) is obvious.

(b) \Rightarrow (c): Let $\overline{M}_n := (M/N)_n := (M_n + N)/N$, $n \in \mathbb{Z}$, then $\overline{M} := M/N = \bigoplus_{n \in \mathbb{Z}} \overline{M}_n$. In fact, let $z \in M$, then $\overline{z} \in \overline{M}$ can be written as $\overline{z} = \overline{z_1 + \dots + z_l} = \overline{z_1} + \dots + \overline{z_l}$, with $z_k \in M_{n_k}$, $1 \leq k \leq l$, thus, $\overline{z} \in \sum_{n \in \mathbb{Z}} \overline{M}_n$, and hence, $\overline{M} = \sum_{n \in \mathbb{Z}} \overline{M}_n$. This sum is direct since if $\overline{z_1} + \dots + \overline{z_l} = \overline{0}$, then $z_1 + \dots + z_l \in N$, so by (b) $z_k \in N$, i.e., $\overline{z_k} = \overline{0}$ for every $1 \leq k \leq l$. Now, let $b_m \in B_m$ and $\overline{z_n} \in \overline{M}_n$, then

$b_m \overline{z_n} = \overline{b_m z_n} = \overline{d_1 + \cdots + d_p}$, with $d_i \in M_{n_i}$ and $n_i \leq m + n$, so $\overline{b_m z_n} = \overline{d_1} + \cdots + \overline{d_p} \in \bigoplus_{k \leq m+n} \overline{M}_k$.

We have proved that \overline{M} is semi-graded.

(c) \Rightarrow (b): Let $z = z_1 + \cdots + z_l \in N$, with $z_i \in M_{n_i}$, $1 \leq i \leq l$, then $\overline{0} = \overline{z_1} + \cdots + \overline{z_l} \in \overline{M} = \bigoplus_{n \in \mathbb{Z}} \overline{M}_n$, therefore, $\overline{z_i} = \overline{0}$, and hence $z_i \in N$ for every i .

(v) The proof is similar to (b) \Rightarrow (c) in (iv).

(vi) By (v), \overline{B} is *SG*. Let $x_1, \dots, x_n \in B$ such that the subring generated by B_0 and x_1, \dots, x_n coincides with B , then it is clear that the subring of \overline{B} generated by \overline{B}_0 and $\overline{x_1}, \dots, \overline{x_n}$ coincides with \overline{B} . Let $n \geq 0$ and $\{z_1, \dots, z_l\}$ be a basis of the free left B_0 -module B_n , then $\{\overline{z_1}, \dots, \overline{z_l}\}$ is a basis of \overline{B}_n : in fact, let $\overline{z} \in \overline{B}_n$ with $z \in B_n$, then $z = c_1 z_1 + \cdots + c_l z_l$, with $c_i \in B_0$, $1 \leq i \leq l$, and hence, $\overline{z} = \overline{c_1} \overline{z_1} + \cdots + \overline{c_l} \overline{z_l}$, i.e., \overline{B}_n is generated by $\overline{z_1}, \dots, \overline{z_l}$ over \overline{B}_0 ; now, if $\overline{c_1} \overline{z_1} + \cdots + \overline{c_l} \overline{z_l} = \overline{0}$, with $c_i \in B_0$, then $c_1 z_1 + \cdots + c_l z_l \in I \cap B_n \subseteq IB_n$ and hence we can write

$$c_1 z_1 + \cdots + c_l z_l = d_1 z_1 + \cdots + d_l z_l, \text{ with } d_i \in I, 1 \leq i \leq l.$$

From this we get that $c_i = d_i$, so $\overline{c_i} = \overline{0}$ for every i . □

Note that the condition imposed to I in (vi) is of type Artin-Rees (see [14]).

Proposition 2.7. (i) *Any \mathbb{N} -graded ring is *SG*.*

(ii) *Let K be a field. Any finitely graded K -algebra is a *FSG* ring.*

(iii) *Any skew PBW extension is a *FSG* ring.*

Proof. (i) and (ii) follow directly from the definitions.

(iii) Let $A = \sigma(R)\langle x_1, \dots, x_n \rangle$ be a skew PBW extension, then $A = \bigoplus_{k \geq 0} A_k$, where

$$A_k := {}_R \langle x^\alpha \in \text{Mon}(A) \mid \deg(x^\alpha) =: \alpha_1 + \cdots + \alpha_n = k \rangle.$$

Thus, A_k is a free left R -module with

$$\dim_R A_k = \binom{n+k-1}{k} = \binom{n+k-1}{n-1}. \quad (2.2)$$

We are assuming that R is an *IBN* ring (*Invariant basis number*), and hence, A also satisfies this condition, see [7]. □

Remark 2.8. (i) Note that the class of *SG* rings includes properly the class of \mathbb{N} -graded rings: In fact, the enveloping algebra of finite-dimensional Lie algebra (see Example 6.14 below) proves this statement.

(ii) The example in (i) proves also that the class of *FSG* rings includes properly the class of finitely graded algebras.

(iii) Finally, the class of *FSG* rings includes properly the class of skew PBW extensions: For this consider the Artin-Schelter regular algebra of global dimension 3 defined by the following relations:

$$yx = xy + z^2, \quad zy = yz + x^2, \quad zx = xz + y^2.$$

Observe that this algebra is a particular case of a Sklyanin algebra wick in general are defined by the following relations:

$$axy + bxy + cz^2 = 0, \quad azy + byz + cx^2 = 0, \quad axz + bzx + cy^2 = 0, \quad a, b, c \in K.$$

3 Generalized Hilbert series and Hilbert polynomial

In this section we introduce the notion of generalized Hilbert series and generalized Hilbert polynomial for semi-graded rings. As in the classical case of finitely graded algebras over fields, these notions depends on the semi-graduation, in particular, they depend on the ring B_0 . We will compute these tools for skew *PBW* extensions.

Definition 3.1. Let $B = \sum_{n \geq 0} \oplus B_n$ be a FSG ring and $M = \bigoplus_{n \in \mathbb{Z}} M_n$ be a finitely semi-graded B -module. The generalized Hilbert series of M is defined by

$$Gh_M(t) := \sum_{n \in \mathbb{Z}} (\dim_{B_0} M_n) t^n.$$

In particular,

$$Gh_B(t) := \sum_{n=0}^{\infty} (\dim_{B_0} B_n) t^n.$$

We say that B has a generalized Hilbert polynomial if there exists a polynomial $Gp_B(t) \in \mathbb{Q}[t]$ such that

$$\dim_{B_0} B_n = Gp_B(n), \text{ for all } n \gg 0.$$

In this case $Gp_B(t)$ is called the generalized Hilbert polynomial of B .

Remark 3.2. (i) Note that if K is a field and B is a finitely graded K -algebra, then the generalized Hilbert series coincides with the habitual Hilbert series, i.e., $Gh_B(t) = h_B(t)$; the same is true for the generalized Hilbert polynomial.

(ii) Observe that if a semi-graded ring B has another semi-graduation $B = \bigoplus_{n \geq 0} C_n$, then its generalized Hilbert series and its generalized Hilbert polynomial can change, i.e., the notions of generalized Hilbert series and generalized Hilbert polynomial depend on the semi-graduation, in particular on B_0 . For example, consider the habitual real polynomial ring in two variables $B := \mathbb{R}[x, y]$, then $Gh_B(t) = \frac{1}{(1-t)^2}$ and $Gp_B(t) = t + 1$; but if we view this ring as $B = (\mathbb{R}[x])[y]$ then $C_0 = \mathbb{R}[x]$, its generalized Hilbert polynomial series is $\frac{1}{1-t}$ and its generalized Hilbert polynomial is 1.

For skew *PBW* extensions the generalized Hilbert series and the generalized Hilbert polynomial can be computed explicitly.

Theorem 3.3. Let $A = \sigma(R)\langle x_1, \dots, x_n \rangle$ be an arbitrary skew *PBW* extension. Then,

(i)

$$Gh_A(t) = \frac{1}{(1-t)^n}. \quad (3.1)$$

(ii)

$$Gp_A(t) = \frac{1}{(n-1)!} [t^{n-1} - s_1 t^{n-2} + \dots + (-1)^r s_r t^{n-r-1} + \dots + (n-1)!], \quad (3.2)$$

where $s_1, \dots, s_k, \dots, s_{n-1}$ are the elementary symmetric polynomials in the variables $1 - n, 2 - n, \dots, (n-1) - n$.

Proof. (i) We have

$$Gh_A(t) = \sum_{k=0}^{\infty} (\dim_R A_k) t^k = \sum_{k=0}^{\infty} \binom{n+k-1}{k} t^k = \frac{1}{(1-t)^n}.$$

(ii) Note that

$$\begin{aligned}
\dim_R A_k &= \binom{n+k-1}{k} = \frac{(n+k-1)!}{k!(n-1)!} \\
&= \frac{(k+n-1)(k+n-2)(k+n-3)\cdots(k+n-(n-1))(k+n-n)!}{k!(n-1)!} \\
&= \frac{(k+n-1)(k+n-2)(k+n-3)\cdots(k+n-(n-1))}{(n-1)!} \\
&= \frac{1}{(n-1)!} [k^{n-1} - s_1 k^{n-2} \cdots + (-1)^r s_r k^{n-r-1} + \cdots + (-1)^{n-1} s_{n-1} k^{n-n}] \\
&= \frac{1}{(n-1)!} [k^{n-1} - s_1 k^{n-2} \cdots + (-1)^r s_r k^{n-r-1} + \cdots + (n-1)!],
\end{aligned}$$

where $s_1, \dots, s_k, \dots, s_{n-1}$ are the elementary symmetric polynomials in the variables $1-n, 2-n, \dots, (n-1) - n$. Thus, we found a polynomial $Gp_A(t) \in \mathbb{Q}[t]$ of degree $n-1$ such that

$$\dim_R A_k = Gp_A(k) \text{ for all } k \geq 0. \quad (3.3)$$

□

From Theorem 3.3, and considering the numeral (ii) in Remark 3.2, we can compute the generalized Hilbert series and the generalized Hilbert polynomial for all examples of skew *PBW* extensions described in [13]. In addition, for the skew quantum polynomials, we can interpret some of them as quasi-commutative bijective skew *PBW* extensions of the r -multiparameter quantum torus. Thus, we have the following tables:

Ring	$Gh_A(t)$	$Gp_A(t)$
Habitual polynomial ring $R[x_1, \dots, x_n]$	$\frac{1}{(1-t)^n}$	$\frac{1}{(n-1)!} [t^{n-1} + \dots + 1]$
Ore extension of bijective type $R[x_1; \sigma_1, \delta_1] \cdots [x_n; \sigma_n, \delta_n]$	$\frac{1}{(1-t)^n}$	$\frac{1}{(n-1)!} [t^{n-1} + \dots + 1]$
Weyl algebra $A_n(K)$	$\frac{1}{(1-t)^n}$	$\frac{1}{(n-1)!} [t^{n-1} + \dots + 1]$
Extended Weyl algebra $B_n(K)$	$\frac{1}{(1-t)^n}$	$\frac{1}{(n-1)!} [t^{n-1} + \dots + 1]$
Enveloping algebra of a Lie algebra \mathcal{G} of dimension n , $\mathcal{U}(\mathcal{G})$	$\frac{1}{(1-t)^n}$	$\frac{1}{(n-1)!} [t^{n-1} + \dots + 1]$
Tensor product $R \otimes_K \mathcal{U}(\mathcal{G})$	$\frac{1}{(1-t)^n}$	$\frac{1}{(n-1)!} [t^{n-1} + \dots + 1]$
Crossed product $R * \mathcal{U}(\mathcal{G})$	$\frac{1}{(1-t)^n}$	$\frac{1}{(n-1)!} [t^{n-1} + \dots + 1]$
Algebra of q -differential operators $D_{q,h}[x, y]$	$\frac{1}{1-t}$	1
Algebra of shift operators S_h	$\frac{1}{1-t}$	1
Mixed algebra D_h	$\frac{1}{(1-t)^2}$	$t + 1$
Discrete linear systems $K[t_1, \dots, t_n][x_1, \sigma_1] \cdots [x_n; \sigma_n]$	$\frac{1}{(1-t)^n}$	$\frac{1}{(n-1)!} [t^{n-1} + \dots + 1]$
Linear partial shift operators $K[t_1, \dots, t_n][E_1, \dots, E_n]$	$\frac{1}{(1-t)^n}$	$\frac{1}{(n-1)!} [t^{n-1} + \dots + 1]$
Linear partial shift operators $K(t_1, \dots, t_n)[E_1, \dots, E_n]$	$\frac{1}{(1-t)^n}$	$\frac{1}{(n-1)!} [t^{n-1} + \dots + 1]$
L. P. Differential operators $K[t_1, \dots, t_n][\partial_1, \dots, \partial_n]$	$\frac{1}{(1-t)^n}$	$\frac{1}{(n-1)!} [t^{n-1} + \dots + 1]$
L. P. Differential operators $K(t_1, \dots, t_n)[\partial_1, \dots, \partial_n]$	$\frac{1}{(1-t)^n}$	$\frac{1}{(n-1)!} [t^{n-1} + \dots + 1]$
L. P. Difference operators $K[t_1, \dots, t_n][\Delta_1, \dots, \Delta_n]$	$\frac{1}{(1-t)^n}$	$\frac{1}{(n-1)!} [t^{n-1} + \dots + 1]$
L. P. Difference operators $K(t_1, \dots, t_n)[\Delta_1, \dots, \Delta_n]$	$\frac{1}{(1-t)^n}$	$\frac{1}{(n-1)!} [t^{n-1} + \dots + 1]$
L. P. q -dilation operators $K[t_1, \dots, t_n][H_1^{(q)}, \dots, H_m^{(q)}]$	$\frac{1}{(1-t)^m}$	$\frac{1}{(m-1)!} [t^{m-1} + \dots + 1]$
L. P. q -dilation operators $K(t_1, \dots, t_n)[H_1^{(q)}, \dots, H_m^{(q)}]$	$\frac{1}{(1-t)^m}$	$\frac{1}{(m-1)!} [t^{m-1} + \dots + 1]$
L. P. q -differential operators $K[t_1, \dots, t_n][D_1^{(q)}, \dots, D_m^{(q)}]$	$\frac{1}{(1-t)^m}$	$\frac{1}{(m-1)!} [t^{m-1} + \dots + 1]$
L. P. q -differential operators $K(t_1, \dots, t_n)[D_1^{(q)}, \dots, D_m^{(q)}]$	$\frac{1}{(1-t)^m}$	$\frac{1}{(m-1)!} [t^{m-1} + \dots + 1]$
Diffusion algebras	$\frac{1}{(1-t)^n}$	$\frac{1}{(n-1)!} [t^{n-1} + \dots + 1]$
Additive analogue of the Weyl algebra $A_n(q_1, \dots, q_n)$	$\frac{1}{(1-t)^n}$	$\frac{1}{(n-1)!} [t^{n-1} + \dots + 1]$
Multiplicative analogue of the Weyl algebra $\mathcal{O}_n(\lambda_{j_i})$	$\frac{1}{(1-t)^{n-1}}$	$\frac{1}{(n-2)!} [t^{n-2} + \dots + 1]$
Quantum algebra $\mathcal{U}'(\mathfrak{so}(3, K))$	$\frac{1}{(1-t)^3}$	$\frac{1}{2} [t^2 + 3t + 1]$
3-dimensional skew polynomial algebras	$\frac{1}{(1-t)^3}$	$\frac{1}{2} [t^2 + 3t + 1]$
Dispin algebra $\mathcal{U}(\mathfrak{osp}(1, 2))$	$\frac{1}{(1-t)^3}$	$\frac{1}{2} [t^2 + 3t + 1]$
Woronowicz algebra $W_\nu(\mathfrak{sl}(2, K))$	$\frac{1}{(1-t)^3}$	$\frac{1}{2} [t^2 + 3t + 1]$
Complex algebra $V_q(\mathfrak{sl}(3, \mathbb{C}))$	$\frac{1}{(1-t)^6}$	$\frac{1}{120} [t^5 + 15t^4 + 85t^3 + 217t^2 + 274t + 120]$
Algebra \mathcal{U}	$\frac{1}{(1-t)^{2n}}$	$\frac{1}{(2n-1)!} [t^{2n-1} + \dots + 1]$
Manin algebra $\mathcal{O}_q(M_2(K))$	$\frac{1}{(1-t)^3}$	$\frac{1}{2} [t^2 + 3t + 1]$
Coordinate algebra of the quantum group $SL_q(2)$	$\frac{1}{(1-t)^3}$	$\frac{1}{2} [t^2 + 3t + 1]$
q -Heisenberg algebra $\mathbf{H}_n(q)$	$\frac{1}{(1-t)^{2n}}$	$\frac{1}{(2n-1)!} [t^{2n-1} + \dots + 1]$
Quantum enveloping algebra of $\mathfrak{sl}(2, K)$, $\mathcal{U}_q(\mathfrak{sl}(2, K))$	$\frac{1}{(1-t)^2}$	$t + 1$
Hayashi algebra $W_q(J)$	$\frac{1}{(1-t)^{2n}}$	$\frac{1}{(2n-1)!} [t^{2n-1} + \dots + 1]$
Differential operators on a quantum space S_q , $D_q(S_q)$	$\frac{1}{(1-t)^n}$	$\frac{1}{(n-1)!} [t^{n-1} + \dots + 1]$
Witten's Deformation of $\mathcal{U}(\mathfrak{sl}(2, K))$	$\frac{1}{1-t}$	1
Quantum Weyl algebra of Maltsiniotis A_n^q, λ	$\frac{1}{(1-t)^2}$	$t + 1$
Quantum Weyl algebra $A_n(q, p_{i,j})$	$\frac{1}{(1-t)^2}$	$t + 1$
Multiparameter Weyl algebra $A_n^{Q,1}(K)$	$\frac{1}{(1-t)^2}$	$t + 1$
Quantum symplectic space $\mathcal{O}_q(\mathfrak{sp}(K^{2n}))$	$\frac{1}{(1-t)^2}$	$t + 1$
Quadratic algebras in 3 variables	$\frac{1}{1-t}$	1

Table 1: Hilbert series and Hilbert polynomial for some examples of bijective skew PBW extensions.

Ring	$Gh_A(t)$	$Gp_A(t)$
n -Multiparametric skew quantum space $R_{\mathbf{q},\sigma}[x_1, \dots, x_n]$	$\frac{1}{(1-t)^n}$	$\frac{1}{(n-1)!} [t^{n-1} + \dots + 1]$
n -Multiparametric quantum space $R_{\mathbf{q}}[x_1, \dots, x_n]$	$\frac{1}{(1-t)^n}$	$\frac{1}{(n-1)!} [t^{n-1} + \dots + 1]$
n -Multiparametric skew quantum space $K_{\mathbf{q},\sigma}[x_1, \dots, x_n]$	$\frac{1}{(1-t)^n}$	$\frac{1}{(n-1)!} [t^{n-1} + \dots + 1]$
n -Multiparametric quantum space $K_{\mathbf{q}}[x_1, \dots, x_n]$	$\frac{1}{(1-t)^n}$	$\frac{1}{(n-1)!} [t^{n-1} + \dots + 1]$
Ring of skew quantum polynomials $R_{\mathbf{q},\sigma}[x_1^{\pm 1}, \dots, x_r^{\pm 1}, x_{r+1}, \dots, x_n]$	$\frac{1}{(1-t)^{n-r}}$	$\frac{1}{(n-r-1)!} [t^{n-r-1} + \dots + 1]$
Ring of quantum polynomials $R_{\mathbf{q}}[x_1^{\pm 1}, \dots, x_r^{\pm 1}, x_{r+1}, \dots, x_n]$	$\frac{1}{(1-t)^{n-r}}$	$\frac{1}{(n-r-1)!} [t^{n-r-1} + \dots + 1]$
Algebra of skew quantum polynomials $K_{\mathbf{q},\sigma}[x_1^{\pm 1}, \dots, x_r^{\pm 1}, x_{r+1}, \dots, x_n]$	$\frac{1}{(1-t)^{n-r}}$	$\frac{1}{(n-r-1)!} [t^{n-r-1} + \dots + 1]$
Algebra of quantum polynomials $\mathcal{O}_{\mathbf{q}} = K_{\mathbf{q}}[x_1^{\pm 1}, \dots, x_r^{\pm 1}, x_{r+1}, \dots, x_n]$	$\frac{1}{(1-t)^{n-r}}$	$\frac{1}{(n-r-1)!} [t^{n-r-1} + \dots + 1]$

Table 2: Hilbert series and Hilbert polynomials of some skew quantum polynomials.

4 Generalized Gelfand-Kirillov dimension

With respect to the Gelfand-Kirillov dimension, the classical definition over fields is not good since, in general, for a ring R a finitely generated R -module is not free. Whence, we have to replace the classical dimension of free modules with other invariant. Next we will show that for our purposes the Goldie dimension works properly, assuming that R is a left noetherian domain. A similar problem was considered in [4] for algebras over commutative noetherian domains replacing the vector space dimension with the reduced rank.

The following two remarks induce our definition.

- (i) If R is a left noetherian domain, then R is a left Ore domain and hence $\text{udim}({}_R R) = 1$. From this we get the following conclusion: let V be a free R -module of finite dimension, i.e., $\dim_R V = k$, then $\text{udim}(V) = k$: in fact, $V \cong R^k$, and from this we obtain $\text{udim}(V) = \text{udim}({}_R R \oplus \dots \oplus {}_R R) = \text{udim}({}_R R) + \dots + \text{udim}({}_R R) = k$ (see [14]).
- (ii) Let R be a left noetherian domain and $A = \sigma(R)\langle x_1, \dots, x_n \rangle$ be a skew PBW extension of R , then (3.1) takes the following form:

$$Gh_A(t) = \sum_{k=0}^{\infty} (\text{udim} A_k) t^k = \sum_{k=0}^{\infty} (\dim_R A_k) t^k = \sum_{k=0}^{\infty} \binom{n+k-1}{k} t^k = \frac{1}{(1-t)^n}.$$

Definition 4.1. Let B be a FSG ring such that B_0 is a left noetherian domain. The generalized Gelfand-Kirillov dimension of B is defined by

$$\text{GGKdim}(B) := \sup_V \overline{\lim}_{k \rightarrow \infty} \log_k \text{udim} V^k,$$

where V ranges over all frames of B and $V^k := {}_{B_0} \langle v_1 \cdots v_k | v_i \in V \rangle$ (a frame of B is a finite dimensional B_0 -free submodule of B such that $1 \in V$).

Remark 4.2. (i) Note that B has at least one frame: B_0 is a frame of dimension 1. We say that V is a *generating frame* of B if the subring of B generating by V and B_0 is B . For example, if R is a left noetherian domain and $A = \sigma(R)\langle x_1, \dots, x_n \rangle$ is a skew PBW extension of R , then $V := {}_R \langle 1, x_1, \dots, x_n \rangle$ is a generating frame of A .

(ii) In a similar way as was observed in Remark 3.2, the notion of generalized Gelfand-Kirillov dimension of a finitely semi-graded ring B depends on the semi-graduation, in particular, depends on B_0 . Note that this type of consideration was made in [4] for an alternative notion of the Gelfand-Kirillov dimension using the reduced rank.

(iii) If B is a finitely graded K -algebra, then the classical Gelfand-Kirillov dimension of B coincides with the just defined above notion, i.e., $\text{GGKdim}(B) = \text{GKdim}(B)$.

Proposition 4.3. *Let B be a FSG ring such that B_0 is a left noetherian domain. Let V be a generating frame of B , then*

$$\text{GGKdim}(B) = \overline{\lim}_{k \rightarrow \infty} \log_k(\text{udim}V^k). \quad (4.1)$$

Moreover, this equality does not depend on the generating frame V .

Proof. It is clear that $\overline{\lim}_{k \rightarrow \infty} \log_k(\text{udim}V^k) \leq \text{GGKdim}(B)$. Let W be any frame of B ; since $\dim_{B_0} W < \infty$, then there exists m such that $W \subseteq V^m$, and hence for every k we have $W^k \subseteq V^{km}$, but observe that V^{km} is a finitely generated left B_0 -module, and since B_0 is left noetherian, then V^{km} is a left noetherian B_0 -module, so $\text{udim}V^{km} < \infty$. From this, $\text{udim}W^k \leq \text{udim}V^{km}$. Therefore, $\log_k(\text{udim}W^k) \leq \log_k(\text{udim}V^{km}) = (1 + \log_k m) \log_{km}(\text{udim}V^{km})$. Since $\overline{\lim}_{k \rightarrow \infty} (1 + \log_k m) = 1$, we get that $\overline{\lim}_{k \rightarrow \infty} \log_k(\text{udim}W^k) \leq \overline{\lim}_{k \rightarrow \infty} \log_{km}(\text{udim}V^{km})$. But observe that $\overline{\lim}_{k \rightarrow \infty} \log_{km}(\text{udim}V^{km}) \leq \overline{\lim}_{k \rightarrow \infty} \log_k(\text{udim}V^k)$, whence

$$\text{GGKdim}(B) = \sup_W \overline{\lim}_{k \rightarrow \infty} \log_k(\text{udim}W^k) \leq \overline{\lim}_{k \rightarrow \infty} \log_k(\text{udim}V^k).$$

The proof of the second statement is completely similar. \square

Next we present the main result of the present subsection.

Theorem 4.4. *Let R be a left noetherian domain and $A = \sigma(R)\langle x_1, \dots, x_n \rangle$ be a skew PBW extension of R . Then,*

$$\text{GGKdim}(A) = \overline{\lim}_{k \rightarrow \infty} \log_k \left(\sum_{i=0}^k \dim_R A_i \right) = 1 + \deg(Gp_A(t)) = n.$$

Proof. According to (4.1), $\text{GGKdim}(A) = \overline{\lim}_{k \rightarrow \infty} \log_k(\text{udim}V^k)$, with

$$V :=_R \langle 1, x_1, \dots, x_n \rangle = A_0 \oplus A_1;$$

note that $V^k \subseteq A_0 \oplus A_1 \oplus \dots \oplus A_k$, from this and using (3.3) we get

$$\begin{aligned} \text{GGKdim}(A) &\leq \overline{\lim}_{k \rightarrow \infty} \log_k(\text{udim}(\sum_{i=0}^k A_i)) = \overline{\lim}_{k \rightarrow \infty} \log_k(\sum_{i=0}^k \text{udim}A_i) = \\ &= \overline{\lim}_{k \rightarrow \infty} \log_k(\sum_{i=0}^k \dim_R A_i) = \overline{\lim}_{k \rightarrow \infty} \log_k(\sum_{i=0}^k Gp_A(i)) = \\ &= \overline{\lim}_{k \rightarrow \infty} \log_k(Gp_A(0) + Gp_A(1) + \dots + Gp_A(k)), \end{aligned}$$

but according to (3.2), every coefficient in $Gp_A(t)$ is positive, so $Gp_A(i)$ is positive for every $0 \leq i \leq k$, moreover, $Gp_A(i) \leq Gp_A(k)$, so $Gp_A(0) + Gp_A(1) + \dots + Gp_A(k) \leq (k+1)Gp_A(k)$ and hence

$$\text{GGKdim}(A) \leq \overline{\lim}_{k \rightarrow \infty} \log_k((k+1)Gp_A(k)) = \overline{\lim}_{k \rightarrow \infty} \log_k(k+1) + \overline{\lim}_{k \rightarrow \infty} \log_k(Gp_A(k)) = 1 + \overline{\lim}_{k \rightarrow \infty} \log_k(Gp_A(k)).$$

Observe that every summand of $Gp_A(k)$ in the bracket of (3.2) is $\leq k^{n-1}$ for k enough large, so $Gp_A(k) \leq \frac{n}{(n-1)!} k^{n-1}$ for $k \gg 0$ and this implies that

$$\text{GGKdim}(A) \leq 1 + \overline{\lim}_{k \rightarrow \infty} \log_k \frac{n}{(n-1)!} + \overline{\lim}_{k \rightarrow \infty} \log_k k^{n-1} = 1 + 0 + n - 1 = 1 + \deg(Gp_A(t)) = n.$$

Now we have to prove that $\text{GGKdim}(A) \geq n$. Note that $W := V^{k^{n-1}}$ is a frame of A and $\text{udim}W^k = \text{udim}V^{k^n} \geq k^n$, therefore, $\log_k(\text{udim}W^k) \geq \log_k k^n = n$, and hence

$$\text{GGKdim}(A) \geq \overline{\lim}_{k \rightarrow \infty} \log_k(\text{udim}W^k) = \overline{\lim}_{k \rightarrow \infty} \log_k(\text{udim}V^{k^n}) \geq \overline{\lim}_{k \rightarrow \infty} n = n. \quad \square$$

5 Non-commutative schemes associated to SG rings

The purpose of this section is to extend the notion of non-commutative projective scheme to the case of semi-graded rings. We will assume that the ring B satisfies the following conditions:

- (C1) B is left noetherian SG .
- (C2) B_0 is left noetherian.
- (C3) For every n , B_n is a finitely generated left B_0 -module.
- (C4) $B_0 \subset Z(B)$.

Remark 5.1. (i) From (C4) we have that B_0 is a commutative noetherian ring.

(ii) All important examples of skew PBW extensions satisfy (C1) and (C2). Indeed, let $A = \sigma(R)\langle x_1, \dots, x_n \rangle$ be a bijective skew PBW extension of R , assuming that R is left noetherian, then A is also left noetherian (Theorem 1.5); in addition, by Proposition 2.7, A also satisfies (C3).

(iii) With respect to condition (C4), it is satisfied for finitely graded K -algebras since in such case $B_0 = K$. On the other hand, let $A = \sigma(R)\langle x_1, \dots, x_n \rangle$ be a skew PBW extension of R , then in general $R = A_0 \not\subset Z(A)$, unless A be a K -algebra, with $A_0 = K$ a commutative ring.

(iv) It is important to remark that some results below can be proved without assuming all conditions (C1)-(C4). For example, in Definition 5.6 we only need (C1).

Proposition 5.2. *Let $\text{sgr} - B$ be the collection of all finitely generated semi-graded B -modules, then $\text{sgr} - B$ is an abelian category where the morphisms are the homogeneous B -homomorphisms.*

Proof. It is clear that $\text{sgr} - B$ is a category. $\text{sgr} - B$ has kernels and co-kernels: Let M, M' be objects of $\text{sgr} - B$ and let $f : M \rightarrow M'$ be an homogeneous B -homomorphism; let $L := \ker(f)$, since B is left noetherian and M is finitely generated, then L is a finitely generated semi-graded B -module; let $M'/\text{Im}(f)$ be the co-kernel of f , note that $\text{Im}(f)$ is semi-graded, so $M'/\text{Im}(f)$ is a semi-graded finitely generated B -module.

$\text{sgr} - B$ is normal and co-normal: Let $f : M \rightarrow M'$ be a monomorphism in $\text{sgr} - B$, then f is the kernel of the canonical homomorphism $j : M' \rightarrow M'/\text{Im}(f)$. Now, let $f : M \rightarrow M'$ be an epimorphism in $\text{sgr} - B$, then f is the co-kernel of the inclusion $\iota : \ker(f) \rightarrow M'$.

$\text{sgr} - B$ is additive: the trivial module 0 is an object of $\text{sgr} - B$; if $\{M_i\}$ is a finite family of objects of $\text{sgr} - B$, then its co-product $\bigoplus M_i$ in the category of left B -modules is an object of $\text{sgr} - B$, with semi-graduation given by

$$(\bigoplus M_i)_p := \bigoplus (M_i)_p, p \in \mathbb{Z}.$$

Thus, $\text{sgr} - B$ has finite co-products. Finally, for any objects M, M' of $\text{sgr} - B$, $\text{Mor}(M, M')$ is an abelian group and the composition of morphisms is bilinear with respect the operations in these groups. \square

Definition 5.3. *Let M be an object of $\text{sgr} - B$.*

(i) *For $s \geq 0$, $B_{\geq s}$ is the least two-sided ideal of B that satisfies the following conditions:*

- (a) $B_{\geq s}$ contains $\bigoplus_{p \geq s} B_p$.
- (b) $B_{\geq s}$ is semi-graded as left ideal of B .
- (c) $B_{\geq s}$ is a direct summand of B .

(ii) *An element $x \in M$ is torsion if there exist $s, n \geq 0$ such that $B_{\geq s}^n x = 0$; the set of torsion elements of M is denoted by $T(M)$; M is torsion if $T(M) = M$ and torsion-free if $T(M) = 0$.*

(iii) *For $s, n \geq 0$, $M_{s,n}$ will denote the least semi-graded submodule of M containing $B_{\geq s}^n M$.*

Remark 5.4. (i) Observe that if B is \mathbb{N} -graded, then $B_{\geq s} = \bigoplus_{p \geq s} B_p$.

(ii) Note that $T(M)$ is a submodule of M : In fact, let $x, y \in T(M)$, then there exist $r, s, n, m \geq 0$ such that $B_{> s}^n x = 0$ and $B_{> r}^m y = 0$; observe that $B_{\geq r+s} \subseteq B_{\geq s}, B_{\geq r}$, so $B_{\geq r+s}^{n+m} x \subseteq B_{> s}^n x = 0$ and $B_{\geq r+s}^{n+m} y \subseteq B_{> r}^m y = 0$, whence $B_{\geq r+s}^{n+m}(x+y) = 0$, i.e., $x+y \in T(M)$; if $b \in B$, then $B_{\geq s}^n b \subseteq B_{\geq s}^n$, so $B_{\geq s}^n bx \subseteq B_{\geq s}^n x = 0$, i.e., $bx \in T(M)$.

(iii) Since M is noetherian, $M_{s,n}$ is finitely generated, i.e., $M_{s,n}$ is an object of $\text{sgr} - B$. Moreover, $M/M_{s,n}$ is torsion because $B_{\geq s}^n M \subseteq M_{s,n}$. In addition, note that $M_{s,n}$ is a direct summand of M .

(iv) If we assume that B is a domain, and hence, a left Ore domain, an alternative notion of torsion can be defined as in the classical case of commutative domains: An element $x \in M$ is torsion if there exists $b \neq 0$ in B such that $bx = 0$; the set $t(M)$ of torsion elements of M is in this case also a submodule of M . In addition, note that $T(M) \subseteq t(M)$: Since $B_{\geq s} \neq 0$, let $b \neq 0$ in $B_{\geq s}$, then $b^n x = 0$ and $b^n \neq 0$, i.e., $x \in t(M)$.

(v) It is clear that the collection \mathcal{T} of modules M in $\text{sgr} - B$ such that $t(M) = M$ conforms a full subcategory of $\text{sgr} - B$. Moreover, let $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ be a short exact sequence in $\text{sgr} - B$; it is obvious that $t(M) = M$ if and only if $t(M') = M'$ and $t(M'') = M''$, i.e., the collection \mathcal{T} is a Serre subcategory of $\text{sgr} - B$. The next lemma shows that this property is satisfied also by the torsion modules introduced in Definition 5.3.

Theorem 5.5. *The collection $\text{stor} - B$ of torsion modules forms a Serre subcategory of $\text{sgr} - B$, and the quotient category*

$$\text{qsgr} - B := \text{sgr} - B / \text{stor} - B$$

is abelian.

Proof. It is obvious that $\text{stor} - B$ is a full subcategory of $\text{sgr} - B$. Let $0 \rightarrow M' \xrightarrow{\iota} M \xrightarrow{j} M'' \rightarrow 0$ be a short exact sequence in $\text{sgr} - B$.

Suppose that M is in $\text{stor} - B$ and let $x' \in M'$, then $\iota(x') \in M$ and there exist $s, n \geq 0$ such that $\iota(B_{> s}^n x') = B_{> s}^n \iota(x') = 0$, but since ι is injective, then $B_{> s}^n x' = 0$. This means that $x' \in T(M')$, so $T(M') = M'$, i.e., M' is in $\text{stor} - B$. Now let $x'' \in M''$, then there exists $x \in M$ such that $j(x) = x''$; there exist $r, m \geq 0$ such that $B_{> r}^m x = 0$, whence $B_{> r}^m x'' = 0$, this implies that $x'' \in T(M'')$. Thus, $T(M'') = M''$, i.e., M'' is in $\text{stor} - B$.

Conversely, suppose that M' and M'' are in $\text{stor} - B$; let $x \in M$, then there exist $s, n \geq 0$ such that $B_{> s}^n j(x) = 0$, i.e., $j(B_{> s}^n x) = 0$. Therefore, $B_{> s}^n x \subseteq \ker(j) = \text{Im}(\iota)$, but since M' is torsion, then $\text{Im}(\iota)$ is also a torsion module. Because B is left noetherian, there exist $a_1, \dots, a_l \in B_{\geq s}^n$ such that $B_{\geq s}^n = Ba_1 + \dots + Ba_l$; there exist $r_i, m_i \geq 0, 1 \leq i \leq l$, such that $B_{> r_i}^{m_i} a_i x = 0$. Without loss of generality we can assume that $r_1 \geq r_i$ for every i , so $B_{\geq r_1} \subseteq B_{\geq r_i}$ and hence $B_{\geq r_1}^{m_1} \subseteq B_{\geq r_1}^{m_i}, B_{\geq r_1}^{m_2} \subseteq B_{\geq r_1}^{m_2}, \dots, B_{\geq r_1}^{m_l} \subseteq B_{\geq r_1}^{m_l}$; from this we get that $B_{\geq r_1}^{m_1} a_1 x = 0, B_{\geq r_1}^{m_2} a_2 x = 0, \dots, B_{\geq r_1}^{m_l} a_l x = 0$, let $m := \max\{m_1, \dots, m_l\}$, then $B_{\geq r}^m a_i x = 0$ for every $1 \leq i \leq l$, with $r := r_1$. Therefore, $B_{\geq r}^m B_{\geq s}^n x = B_{\geq r}^m (Ba_1 + \dots + Ba_l)x = B_{\geq r}^m a_1 x + \dots + B_{\geq r}^m a_l x = 0$, i.e., $B_{\geq r}^m B_{\geq s}^n x = 0$. Observe that $B_{\geq r+s}^m \subseteq B_{\geq r}^m$ and $B_{\geq r+s}^n \subseteq B_{\geq s}^n$, so $B_{\geq r+s}^{m+n} \subseteq B_{\geq r}^m B_{\geq s}^n$ and hence $B_{\geq r}^{m+n} x = 0$, i.e., $x \in T(M)$. We have proved that $T(M) = M$, i.e., M is in $\text{stor} - B$.

The second statement of the theorem is a well known property of abelian categories. We want to recall that the objects of $\text{qsgr} - B$ are the objects of $\text{sgr} - B$; moreover, given M, N objects of $\text{qsgr} - B$ the set of morphisms from M to N in the category $\text{qsgr} - B$ is defined by

$$\text{Hom}_{\text{qsgr} - B}(M, N) := \varinjlim \text{Hom}_{\text{sgr} - B}(M', N/N')$$

where the direct limit is taken over all $M' \subseteq M, N' \subseteq N$ in $\text{sgr} - B$ with $M/M' \in \text{stor} - B$ and $N' \in \text{stor} - B$ (see [10], [5], or also [16] Proposition 2.13.4). More exactly, the limit is taken over the set \mathcal{P} of all pairs (M', N') in $\text{sgr} - B$ such that $M' \subseteq M, N' \subseteq N, M/M' \in \text{stor} - B$ and $N' \in \text{stor} - B$. The set \mathcal{P} is partially ordered with order defined by

$(M', N') \leq (M'', N'')$ if and only if $M'' \subseteq M'$ and $N' \subseteq N''$.

\mathcal{P} is directed: Indeed, given $(M', N'), (M'', N'') \in \mathcal{P}$ we apply Proposition 2.6 and the fact that B is left noetherian to conclude that $(M' \cap M'', N' + N'') \in \mathcal{P}$, and this couple satisfies $(M', N') \leq (M' \cap M'', N' + N'')$, $(M'', N'') \leq (M' \cap M'', N' + N'')$.

□

We have all ingredients in order to define non-commutative schemes associated to semi-graded rings.

Definition 5.6. *We define*

$$\text{sproj}(B) := (\text{qsg}r - B, \pi(B))$$

and we call it the non-commutative semi-projective scheme associated to B .

6 Serre-Artin-Zhang-Verevkin theorem for semi-graded rings

We conclude the paper investigating the non-commutative version of Serre-Artin-Zhang-Verevkin theorem for semi-graded rings. For this goal some preliminaries are needed.

Definition 6.1. *Let M be a semi-graded B -module, $M = \bigoplus_{n \in \mathbb{Z}} M_n$. Let $i \in \mathbb{Z}$, the semi-graded module $M(i)$ defined by $M(i)_n := M_{i+n}$ is called a shift of M , i.e.,*

$$M(i) = \bigoplus_{n \in \mathbb{Z}} M(i)_n = \bigoplus_{n \in \mathbb{Z}} M_{i+n}.$$

Remark 6.2. Note that for every $i \in \mathbb{Z}$, $M \cong M(i)$ as B -modules. The isomorphism is given by

$$\begin{aligned} M &= \bigoplus_{n \in \mathbb{Z}} M_n \xrightarrow{\phi_i} \bigoplus_{n \in \mathbb{Z}} M_{i+n} = M(i) \\ m_{n_1} + \cdots + m_{n_t} \in M_{n_1} + \cdots + M_{n_t} &\mapsto m_{n_1} + \cdots + m_{n_t} \in M(i)_{n_1-i} + \cdots + M(i)_{n_t-i}. \end{aligned}$$

ϕ_i is not homogeneous for $i \neq 0$.

The next proposition shows that the shift of degrees is an autoequivalence.

Proposition 6.3. *Let $s : \text{sgr} - B \rightarrow \text{sgr} - B$ defined by*

$$\begin{aligned} M &\mapsto M(1), \\ M \xrightarrow{f} N &\mapsto M(1) \xrightarrow{f(1)} N(1), \\ f(1)(m) &:= f(m), \quad m \in M(1). \end{aligned}$$

Then,

- (i) s is an autoequivalence.
- (ii) For every $d \in \mathbb{Z}$, $s^d(M) = M(d)$.
- (iii) s induces an autoequivalence of $\text{qsg}r - B$ also denoted by s .

Proof. (i) and (ii) are evident. For (iii) we only have to observe that if M is an object of $\text{stor} - B$, then $s(M)$ is also an object of $\text{stor} - B$. □

Definition 6.4. *Let M, N be objects of $\text{sgr} - B$. Then*

- (i) $\underline{\text{Hom}}_B(M, N) := \bigoplus_{d \in \mathbb{Z}} \text{Hom}_{\text{sgr} - B}(M, N(d))$.
- (ii) $\underline{\text{Ext}}_B^i(M, N) := \bigoplus_{d \in \mathbb{Z}} \text{Ext}_{\text{sgr} - B}^i(M, N(d))$.

Remark 6.5. Note that $\underline{Hom}_B(M, N) \hookrightarrow Hom_B(M, N)$. In fact, we have the group homomorphism $\iota : \underline{Hom}_B(M, N) \rightarrow Hom_B(M, N)$ given by $(\dots, 0, f_{d_1}, \dots, f_{d_t}, 0, \dots) \mapsto f_{d_1} + \dots + f_{d_t}$; observe that $f_{d_1} + \dots + f_{d_t} = 0$ if and only if $f_{d_1} = \dots = f_{d_t} = 0$. Indeed, let $m \in M$ be homogeneous of degree p , then $0 = (f_{d_1} + \dots + f_{d_t})(m) = f_{d_1}(m) + \dots + f_{d_t}(m) \in N_{d_1+p} \oplus \dots \oplus N_{d_t+p}$, whence, for every j , $f_{d_j}(m) = 0$. This means that $f_{d_j} = 0$ for $1 \leq j \leq t$, and hence, ι is injective.

Proposition 6.6. *Let M and N be semi-graded B -modules such that every of its homogeneous components are B_0 -modules. Then,*

- (i) $Hom_{\text{sgr}-B}(M, N)$ is a B_0 -module.
- (ii) $\underline{Hom}_B(M, N)$ is a B_0 -module.
- (iii) $Ext_{\text{sgr}-B}^i(M, N)$ is a B_0 -module for every $i \geq 1$.
- (iv) $\underline{Ext}_B^i(M, N)$ is a B_0 -module for every $i \geq 1$.

Proof. (i) If $f \in Hom_{\text{sgr}-B}(M, N)$ and $b_0 \in B_0$, then product $b_0 \cdot f$ defined by $(b_0 \cdot f)(m) := b_0 \cdot f(m)$, $m \in M$, is an element of $Hom_{\text{sgr}-B}(M, N)$: In fact, $b_0 \cdot f$ is obviously additive; let $b \in B$, then $(b_0 \cdot f)(b \cdot m) = b_0 \cdot f(b \cdot m) = b_0[b \cdot f(m)] = (b_0b) \cdot f(m) = (bb_0) \cdot f(m) = b \cdot (b_0 \cdot f(m)) = b \cdot (b_0 \cdot f)(m)$; $b_0 \cdot f$ is homogeneous: Let $m \in M_p$, then $(b_0 \cdot f)(m) = b_0 \cdot f(m) \in b_0 \cdot N_p \subseteq N_p$, for every $p \in \mathbb{Z}$. It is easy to check that $Hom_{\text{sgr}-B}(M, N)$ is a B_0 -module with the defined product.

(ii) This follows from (i).

(iii) Taking a projective resolution of M in the abelian category $\text{sgr}-B$ and applying the functor $Hom_{\text{sgr}-B}(-, N)$, it is easy to verify using (i) that in the complex defining $Ext_{\text{sgr}-B}^i(M, N)$ the kernels and the images are B_0 -modules, i.e., every abelian group $Ext_{\text{sgr}-B}^i(M, N)$ is a B_0 -module.

(iv) This follows from (iii). □

Definition 6.7. *Let $i \geq 0$; we say that B satisfies the $s\text{-}\chi_i$ condition if for every finitely generated semi-graded B -module N and for any $j \leq i$, $\underline{Ext}_B^j(B/B_{\geq 1}, N)$ is finitely generated as B_0 -module. The ring B satisfies the $s\text{-}\chi$ condition if it satisfies the $s\text{-}\chi_i$ condition for all $i \geq 0$.*

Remark 6.8. (i) By Proposition 6.6, $\underline{Ext}_B^j(B/B_{\geq 1}, N)$ is a B_0 -module.

(ii) In the theory of graded rings and modules the conditions defined above are usually denoted simply by χ_i and χ . In this situation, $B/B_{\geq 1} \cong B_0$.

(iii) Observe that in the case of finitely graded K -algebras, $B_0 = K$, $B/B_{\geq 1} \cong K$ and the condition $s\text{-}\chi_i$ means that $\dim_K \underline{Ext}_B^j(K, N) < \infty$.

Definition 6.9. *Let s be the autoequivalence of $\text{qsgr}-B$ defined by the shifts of degrees. We define*

$$\Gamma(\pi(B))_{\geq 0} := \bigoplus_{d=0}^{\infty} Hom_{\text{qsgr}-B}(\pi(B), s^d(\pi(B))).$$

Following the ideas in the proof of Theorem 4.5 in [3] and Proposition 4.11 in [15] we get the following key lemma.

Lemma 6.10. *Let B be a ring that satisfies (C1)-(C4).*

- (i) $\Gamma(\pi(B))_{\geq 0}$ is a \mathbb{N} -graded ring.
- (ii) Let $\underline{B} := \bigoplus_{d=0}^{\infty} Hom_{\text{sgr}-B}(B, s^d(B))$. Then, \underline{B} is a \mathbb{N} -graded ring and there exists a ring homomorphism $\underline{B} \rightarrow \Gamma(\pi(B))_{\geq 0}$.
- (iii) For any object M of $\text{sgr}-B$

$$\Gamma(M)_{\geq 0} := \bigoplus_{d=0}^{\infty} Hom_{\text{sgr}-B}(B, s^d(M))$$

is a graded \underline{B} -module, and

$$\Gamma(\pi(M))_{\geq 0} := \bigoplus_{d=0}^{\infty} \text{Hom}_{\text{qsgr}-B}(\pi(B), s^d(\pi(M)))$$

is a graded $\Gamma(\pi(B))_{\geq 0}$ -module.

(iv) \underline{B} has the following properties:

- (a) $(\underline{B})_0 \cong B_0$ and \underline{B} satisfies (C2).
- (b) \underline{B} satisfies (C3). More generally, let N be a finitely generated graded \underline{B} -module, then every homogeneous component of N is finitely generated over $(\underline{B})_0$.
- (c) \underline{B} satisfies (C1).
- (d) If B is a domain, then \underline{B} is also a domain.

(v) If B is a domain, then

- (a) $\Gamma(\pi(B))_{\geq 0}$ satisfies (C2).
- (b) $\Gamma(\pi(B))_{\geq 0}$ satisfies (C3). More generally, let N be a finitely generated graded $\Gamma(\pi(B))_{\geq 0}$ -module, then every homogeneous component of N is finitely generated over $(\Gamma(\pi(B))_{\geq 0})_0$.
- (c) $\Gamma(\pi(B))_{\geq 0}$ satisfies (C1).
- (d) If \underline{B} satisfies \mathcal{X}_1 , then $\Gamma(\pi(B))_{\geq 0}$ satisfies \mathcal{X}_1 .
- (e) $\Gamma(\pi(B))_{\geq 0}$ is a domain.

Proof. (i) Since $\text{qsgr} - B$ is an abelian category, $\text{Hom}_{\text{qsgr}-B}(\pi(B), s^d(\pi(B)))$ is an abelian group; the product in $\Gamma(\pi(B))_{\geq 0}$ is defined by distributive law and the following rule:

$$\begin{aligned} \text{If } f \in \text{Hom}_{\text{qsgr}-B}(\pi(B), s^n(\pi(B))) \text{ and } g \in \text{Hom}_{\text{qsgr}-B}(\pi(B), s^m(\pi(B))), \text{ then} \\ f \star g := s^n(g) \circ f \in \text{Hom}_{\text{qsgr}-B}(\pi(B), s^{m+n}(\pi(B))). \end{aligned}$$

This product is associative: In fact, if $h \in \text{Hom}_{\text{qsgr}-B}(\pi(B), s^p(\pi(B)))$, then

$$(f \star g) \star h = [s^n(g) \circ f] \star h = s^{m+n}(h) \circ s^n(g) \circ f = f \star (g \star h).$$

It is clear that the product is \mathbb{N} -graded and the unity of $\Gamma(\pi(B))_{\geq 0}$ is i_B taken in $d = 0$ (observe that we have simplified the notation avoiding the bar notation for the morphisms in the category $\text{qsgr} - B$).

(ii) The proof of that \underline{B} is a \mathbb{N} -graded ring is as in (i). For the second assertion we can apply the quotient functor π to define the function

$$\begin{aligned} \underline{B} &\xrightarrow{\rho} \Gamma(\pi(B))_{\geq 0} & (6.1) \\ (f_0, \dots, f_d, 0, \dots) &\mapsto (\pi(f_0), \dots, \pi(f_d), 0, \dots) \end{aligned}$$

which is a ring homomorphism since π is additive (π is exact) and $s\pi = \pi s$.

(iii) The proof of both assertions are as in (i), we only illustrate the product in the first case:

$$\begin{aligned} \text{If } f \in \text{Hom}_{\text{sgr}-B}(B, s^n(B)) \text{ and } g \in \text{Hom}_{\text{sgr}-B}(B, s^m(M)), \text{ then} \\ f \star g := s^n(g) \circ f \in \text{Hom}_{\text{sgr}-B}(B, s^{m+n}(M)). \end{aligned}$$

(iv) (a) Note that $(\underline{B})_0 = \text{Hom}_{\text{sgr}-B}(B, B)$, and consider the function

$$B_0 \xrightarrow{\alpha} \text{Hom}_{\text{sgr}-B}(B, B), \alpha(x) = \alpha_x, \alpha_x(b) := bx, x \in B_0, b \in B;$$

since $B_0 \subset Z(B)$ this function is a ring homomorphism, moreover, bijective. Thus, $(\underline{B})_0$ is a commutative noetherian ring, so $(\underline{B})_0$ satisfies (C2). In addition, observe that the structure of B_0 -module of $\text{Hom}_{\text{sgr}-B}(B, B)$ induced by α coincides with the structure defined in Proposition 6.6.

(b) Note that the function $\text{Hom}_{\text{sgr}-B}(B, B(d)) \xrightarrow{\lambda} B_d$ defined by $f \mapsto f(1)$ is an injective B_0 -homomorphism. Since B_0 is noetherian and B satisfies C3, then $\text{Hom}_{\text{sgr}-B}(B, B(d))$ is finitely generated over $B_0 \cong (\underline{B})_0$.

For the second part, let N be generated by x_1, \dots, x_r , with $x_i \in N_{d_i}$, $1 \leq i \leq r$. Let $x \in N_d$, then there exist $f_1, \dots, f_r \in \underline{B}$ such that $x = f_1 \cdot x_1 + \dots + f_r \cdot x_r$, from this we can assume that $f_i \in (\underline{B})_{d-d_i}$; by the just proved property (C3) for \underline{B} we obtain that every $(\underline{B})_{d-d_i}$ is finitely generated as $(\underline{B})_0$ -module, this implies that N_d is finitely generated over $(\underline{B})_0$.

(c) By (ii), \underline{B} is not only SG but \mathbb{N} -graded.

\underline{B} is left noetherian: We will adapt a proof given in [3]. Let I be a graded left ideal of \underline{B} ; let $f \in \underline{B}$ be homogeneous of degree d_f , then f induces a morphism $s^{-d_f}(B) \xrightarrow{f_-} B$; thus, given a finite set F of homogeneous elements of I , let $P_F := \bigoplus_{f \in F} s^{-d_f}(B)$, $f_F := \sum_{f \in F} f_- : P_F \rightarrow B$ and let $N_F := \text{Im}(f_F)$. Since B is left noetherian we can choose a finite set F_0 such that N_{F_0} is maximal among such images. Let $N := N_{F_0}$ and $P := P_{F_0}$; we define $N'' := \Gamma(N)_{\geq 0} := \bigoplus_{d=0}^{\infty} \text{Hom}_{\text{sgr}-B}(B, s^d(N))$. According to (iii), N'' is a \mathbb{N} -graded \underline{B} -module. Given any element $f \in I$ homogeneous of degree d_f we have the morphism f_- , but since N is maximal the image of this morphism is included in N , and this implies that $f \in N''$, so $I \subseteq N''$. On the other hand, given $f \in I$ homogeneous of degree d_f the \mathbb{N} -graded \underline{B} -homomorphism $s^{-d_f}(\underline{B}) \xrightarrow{f_-} \underline{B}$ defined by $f_-(h) := hf$ has his image in I . Therefore, $N' \subseteq I$, where N' is the image of the induced morphism $P'' \rightarrow \underline{B}$, with $P'' := \bigoplus_{f \in F_0} s^{-d_f}(\underline{B})$. Thus, we have $N' \subseteq N''$, where both are \mathbb{N} -graded \underline{B} -modules, whence we have the \mathbb{N} -graded \underline{B} -module N''/N' . If we prove that N''/N' is noetherian, then since $I/N' \subseteq N''/N'$ we get that I/N' is also noetherian, whence, I/N' is finitely generated; but N' is a finitely generated left ideal of \underline{B} , so I is finitely generated.

N''/N' is noetherian: Note first that N''/N' is a module over $(\underline{B})_0$; if we prove that N''/N' is noetherian over $(\underline{B})_0$, then it is also noetherian over \underline{B} . According to (a), we only need to show that N''/N' is finitely generated over $(\underline{B})_0$. But this follows from (b) since N''/N' is right bounded (i.e., there exists $n \gg 0$ such that the homogeneous component of N''/N' of degree $k \geq n$ is zero, see [3]).

(d) If B is a domain, then \underline{B} is also a domain: Suppose there exist $f, g \neq 0$ in \underline{B} such that $f \star g = 0$, let $f_n \neq 0$ and $g_m \neq 0$ the nonzero homogeneous components of f and g of lowest degree, thus $f_n \in \text{Hom}_{\text{sgr}-B}(B, s^n(B))$, $g_m \in \text{Hom}_{\text{sgr}-B}(B, s^m(B))$ and $0 = f_n \star g_m = s^n(g_m) \circ f_n \in \text{Hom}_{\text{sgr}-B}(B, s^{m+n}(B))$; since $f_n \neq 0$ we have $f_n(1) \neq 0$, also $g_m(1) \neq 0$ and hence $s^n(g_m)(1) \neq 0$, so $0 = s^n(g_m)(f_n(1)) = f_n(1)s^n(g_m)(1)$, but this is impossible since B is a domain.

(v) We set $\Gamma := \Gamma(\pi(B))_{\geq 0}$. Then,

(a) Γ satisfies (C2): We divide the proof of this statement in two steps.

Step 1. Adapting the proof of Proposition 5.3.7 in [16] we will show that

$$\Gamma_0 = \text{Hom}_{\text{qsgr}-B}(\pi(B), \pi(B)) = \varinjlim \text{Hom}_{\text{sgr}-B}(B_{s,n}, B),$$

where the direct limit is taken over the homomorphisms of abelian groups

$$\text{Hom}_{\text{sgr}-B}(B_{s,n}, B) \rightarrow \text{Hom}_{\text{sgr}-B}(B_{r,m}, B)$$

induced by the inclusion homomorphism $B_{r,m} \rightarrow B_{s,n}$, with $r \geq s$ and $m \geq n$. Observe that the collection of couples (s, n) is a partially ordered directed set.

Note first that $\text{Hom}_{\text{qsgr}-B}(\pi(B), \pi(B)) = \varinjlim \text{Hom}_{\text{sgr}-B}(M', B)$, where the direct limit is taken over all $M' \subseteq B$ with $B/M' \in \text{stor} - B$. In fact, we know that

$$\text{Hom}_{\text{qsgr}-B}(\pi(B), \pi(B)) = \varinjlim \text{Hom}_{\text{sgr}-B}(M', B/N'),$$

where the direct limit is taken over all $(M', N') \in \mathcal{P}$, but since B is a domain, $N' = 0$.

Now let $\bar{f} \in \varinjlim Hom_{\text{sgr}-B}(M', B)$, so $f \in Hom_{\text{sgr}-B}(M', B)$ for some $M' \subseteq B$ such that $T(B/M') = B/M'$; since B/M' is finitely generated, we can reasoning as in the proof of Theorem 5.5 and find $s, n \geq 0$ such that $B_s^n B \subseteq M'$, i.e., $B_{s,n} \subseteq M'$. From this we get that $\bar{f} = \overline{f'}$, where $\overline{f'} \in Hom_{\text{sgr}-B}(B_{s,n}, B)$, with $f' := f\iota$ and $\iota : B_{s,n} \hookrightarrow M'$ the inclusion. Since $B/B_{s,n}$ is torsion we obtain that $\varinjlim Hom_{\text{sgr}-B}(M', B) = \varinjlim Hom_{\text{sgr}-B}(B_{s,n}, B)$.

Step 2. Considering $s, n = 0$ in the limit above we obtain a ring homomorphism

$$(\underline{B})_0 = Hom_{\text{sgr}-B}(B, B) \xrightarrow{\gamma} Hom_{\text{qsgr}-B}(\pi(B), \pi(B)) = \Gamma_0;$$

since $(\underline{B})_0$ is noetherian we can prove that γ is surjective. Let $\bar{f} \in \Gamma_0$ with $f \in Hom_{\text{sgr}-B}(B_{s,n}, B)$, consider the commutative triangles

$$\begin{array}{ccc} & B & \\ f \nearrow & & \nwarrow f' \\ B_{s,n} & \xrightarrow{\iota} & B_{0,0} = B \end{array}, \quad \begin{array}{ccc} & B & \\ f \nearrow & & \nwarrow f \\ B_{s,n} & \xrightarrow{i} & B_{s,n} \end{array}$$

where f' is defined by $f'(x+l) := f(x)$, with $x \in B_{s,n}$, $l \in L$ and $B = B_{s,n} \oplus L$. Thus, $i^*(f) = fi = f$ and $\iota^*(f') = f'\iota = f$, so $\bar{f} = \overline{f'} = \gamma(f')$.

From this we conclude that Γ_0 is a commutative noetherian ring, and hence, Γ satisfies (C2).

(b) Γ satisfies (C3): Since Γ is graded, Γ_d is a Γ_0 -module for every d , but by (a) we have a ring homomorphism $B_0 \cong (\underline{B})_0 \xrightarrow{\gamma} \Gamma_0$, so the idea is to prove that Γ_d is finitely generated over B_0 . For this we will show that there exists a surjective B_0 -homomorphism $(\underline{B})_d \xrightarrow{\beta} \Gamma_d$. Note that $\Gamma_d = Hom_{\text{qsgr}-B}(\pi(B), \pi(B(d))) = \varinjlim Hom_{\text{sgr}-B}(B_{s,n}, B(d))$ (the proof of this is as the step 1 in (a)); let $f \in (\underline{B})_d = Hom_{\text{sgr}-B}(B, B(d))$, we define $\beta(f) := \overline{f\iota}$, where $\iota : B_{s,n} \rightarrow B = B_{0,0}$; we can repeat the proof of the step 2 in (a) and conclude that β is a surjective B_0 -homomorphism.

Additionally, let N be a finitely generated graded Γ -module, says N generated by a finite set of homogeneous elements x_1, \dots, x_r , with $x_i \in N_{d_i}$, $1 \leq i \leq r$. Let $x \in N_d$, then there exist $f_1, \dots, f_r \in \Gamma$ such that $x = f_1 \cdot x_1 + \dots + f_r \cdot x_r$, from this we can assume that $f_i \in \Gamma_{d-d_i}$, but as was observed before, every Γ_{d-d_i} is finitely generated as Γ_0 -module, so N_d is finitely generated over Γ_0 for every d .

(c) Γ satisfies (C1): By (iii), Γ is not only SG but \mathbb{N} -graded.

Γ is left noetherian: We will adapt the proof of (iv)-(c). Let I be a graded left ideal of Γ ; let $f \in \Gamma$ be homogeneous of degree d_f , then f induces a morphism $s^{-d_f}(\pi(B)) \xrightarrow{f_-} \pi(B)$; thus, given a finite set F of homogeneous elements of I , let $P_F := \bigoplus_{f \in F} s^{-d_f}(\pi(B))$, $f_F := \sum_{f \in F} f_- : P_F \rightarrow \pi(B)$ and let $N_F := Im(f_F)$. Since $\pi(B)$ is a noetherian object of $\text{qsgr}-B$ we can choose a finite set F_0 such that N_{F_0} is maximal among such images. Let $\pi(N) := N_{F_0}$ and $\pi(P) := P_{F_0}$; we define $N'' := \Gamma(\pi(N))_{\geq 0} := \bigoplus_{d=0}^{\infty} Hom_{\text{qsgr}-B}(\pi(B), s^d(\pi(N)))$. According to (iii), N'' is a \mathbb{N} -graded Γ -module. Given any element $f \in I$ homogeneous of degree d_f we have the morphism f_- , but since N is maximal the image of this morphism is included in N , and this implies that $f \in N''$, so $I \subseteq N''$. On the other hand, given $f \in I$ homogeneous of degree d_f the \mathbb{N} -graded Γ -homomorphism $s^{-d_f}(\Gamma) \xrightarrow{f_-} \Gamma$ defined by $f_-(h) := hf$ has his image in I . Therefore, $N' \subseteq I$, where N' is the image of the induced morphism $P'' \rightarrow \Gamma$, with $P'' := \bigoplus_{f \in F_0} s^{-d_f}(\Gamma)$. Thus, we have $N' \subseteq N''$, where both are \mathbb{N} -graded Γ -modules, whence we have the \mathbb{N} -graded Γ -module N''/N' . If we prove that N''/N' is noetherian, then since $I/N' \subseteq N''/N'$ we get that I/N' is also noetherian, whence, I/N' is finitely generated; but N' is a finitely generated left ideal of Γ , so I is finitely generated.

N''/N' is noetherian: Note first that N''/N' is a module over Γ_0 ; if we prove that N''/N' is noetherian over Γ_0 , then it is also noetherian over Γ . According to (a), we only need to show that N''/N' is finitely generated over Γ_0 . But this follows from (b) since N''/N' is right bounded.

(d) Γ satisfies \mathcal{X}_1 : Let N be a finitely generated graded Γ -module, we have $\underline{Ext}_\Gamma^j(\Gamma/\Gamma_{\geq 1}, N) = \underline{Ext}_\Gamma^j(\Gamma_0, N)$, so we must prove that $\underline{Ext}_\Gamma^j(\Gamma_0, N)$ is finitely generated as Γ_0 -module for $j = 0, 1$. By the surjective homomorphism $(\underline{B})_0 \xrightarrow{\gamma} \Gamma_0$ in the step 2 in (a), it is enough to show that $\underline{Ext}_\Gamma^j(\Gamma_0, N)$ is finitely generated over $(\underline{B})_0$. Observe that γ is also a graded homomorphism of left (\underline{B}) -modules; moreover, N is a finitely generated graded left (\underline{B}) -module since the homomorphism ρ in (ii) is surjective; the proof of this last statement is as in the step 2 of (a), using of course that B is a domain, we include it for completeness: It is enough to consider $\bar{f} \in \Gamma_d = \varinjlim Hom_{\text{sgr}-B}(B_{s,n}, B(d))$, with $f \in Hom_{\text{sgr}-B}(B_{s,n}, B(d))$ for some $s, n \geq 0$; we define $f' : B_{0,0} \rightarrow B(d)$, $f'(x) := f(y)$, where $B = B_{s,n} \oplus L$ and $x = y + l$ with $y \in B_{s,n}$ and $l \in L$; therefore, $\rho(f') = \pi(f') = \bar{f}$ since we have $f'\iota = f$.

Now we can apply the functor $\underline{Ext}_B^j(\cdot, N)$ and get the injective homomorphism of left $(\underline{B})_0$ -modules $\underline{Ext}_B^j(\Gamma_0, N) \rightarrow \underline{Ext}_B^j((\underline{B})_0, N)$, but since \underline{B} satisfies \mathcal{X}_1 , $\underline{Ext}_B^j((\underline{B})_0, N)$ is finitely generated over $(\underline{B})_0$, so $\underline{Ext}_B^j(\Gamma_0, N)$ is finitely generated since $(\underline{B})_0$ is left noetherian. From the injective $(\underline{B})_0$ -homomorphism $\underline{Ext}_B^j(\Gamma_0, N) \rightarrow \underline{Ext}_\Gamma^j(\Gamma_0, N)$ we conclude that $\underline{Ext}_\Gamma^j(\Gamma_0, N)$ is also finitely generated over $(\underline{B})_0$.

(e) Γ is a domain: Suppose there exist $f, g \neq 0$ in Γ such that $f \star g = 0$, let $f_n \neq 0$ and $g_m \neq 0$ the nonzero homogeneous components of f and g of lowest degree, thus

$$f_n \in Hom_{\text{qsgr}-B}(\pi(B), s^n(\pi(B))), g_m \in Hom_{\text{qsgr}-B}(\pi(B), s^m(\pi(B)))$$

and $0 = f_n \star g_m = s^n(g_m) \circ f_n \in Hom_{\text{qsgr}-B}(\pi(B), s^{m+n}(\pi(B)))$; note that the representative elements of f_n and g_m in $Hom_{\text{sgr}-B}(B_{0,0}, B(n)) \cong Hom_{\text{sgr}-B}(B, B) = (\underline{B})_0$ and $Hom_{\text{sgr}-B}(B_{0,0}, B(m)) \cong Hom_{\text{sgr}-B}(B, B) = (\underline{B})_0$, respectively, are non zero, but this is impossible since $(\underline{B})_0$ is a domain and the representative element of $f \star g$ in $Hom_{\text{sgr}-B}(B_{0,0}, B(n+m)) \cong Hom_{\text{sgr}-B}(B, B) = (\underline{B})_0$ is zero. \square

Proposition 6.11. *Let S be a commutative noetherian ring and $\rho : C \rightarrow D$ be a homomorphism of \mathbb{N} -graded left noetherian S -algebras. If the kernel and cokernel of ρ are right bounded, then $D \otimes_C -$ defines an equivalence of categories $\text{qgr} - C \simeq \text{qgr} - D$, where \otimes denotes the graded tensor product.*

Proof. The proof of Proposition 2.5 in [3] applies since it is independent of the notion of torsion. \square

We are prepared for proving the main theorem of the present section.

Theorem 6.12. *If B is a domain that satisfies (C1)-(C4) and \underline{B} satisfies the condition \mathcal{X}_1 then there exists an equivalence of categories*

$$\text{qgr} - \underline{B} \simeq \text{qgr} - \Gamma(\pi(B))_{\geq 0}.$$

Proof. Note that the ring homomorphism in (6.1) satisfies the conditions of Proposition 6.11, with $S = B_0$, $C = \underline{B}$ and $D = \Gamma(\pi(B))_{\geq 0}$. In fact, from Lemma 6.10 we know that \underline{B} and $\Gamma(\pi(B))_{\geq 0}$ are \mathbb{N} -graded left noetherian rings and B_0 -modules; moreover, they are B_0 -algebras: We check this for \underline{B} , the proof for $\Gamma(\pi(B))_{\geq 0}$ is similar. If $f \in Hom_{\text{sgr}-B}(B, B(n))$, $g \in Hom_{\text{sgr}-B}(B, B(m))$, $x \in B_0$ and $b \in B$, then

$$\begin{aligned} [x \cdot (f \star g)](b) &= x \cdot (s^n(g) \circ f)(b) = xg(n)(f(b)); \\ [f \star (x \cdot g)](b) &= [s^n(x \cdot g) \circ f](b) = (x \cdot g)(n)(f(b)) = xg(n)(f(b)). \end{aligned}$$

Finally, we can apply the proof of part S10 in Theorem 4.5 in [3] to conclude that the kernel and cokernel of ρ are right bounded. \square

Remark 6.13. Considering the above developed theory for graded rings and right modules instead of semi-graded rings and left modules it is possible to prove that $\underline{B} \cong B$. Thus, in such case we get from the previous theorem the Artin-Zhang-Verevkin equivalence $\text{qgr} - B \simeq \text{qgr} - \Gamma(\pi(B))_{\geq 0}$.

Example 6.14. The examples of skew *PBW* extensions below are semi-graded (non \mathbb{N} -graded) domains and satisfy the conditions (C1)-(C4); in each case we will prove that \underline{B} satisfies the condition \mathcal{X}_1 ; therefore, for these algebras Theorem 6.12 is true. In every example $B_0 = K$ is a field, we indicate the relations defining B (see [13]) and the associated graded ring $Gr(B)$ (Proposition 2.6):

(i) Enveloping algebra of a Lie K -algebra \mathcal{G} of dimension n , $\mathcal{U}(\mathcal{G})$:

$$\begin{aligned} x_i k - k x_i &= 0, \quad k \in K; \\ x_i x_j - x_j x_i &= [x_i, x_j] \in \mathcal{G} = Kx_1 + \cdots + Kx_n, \quad 1 \leq i, j \leq n; \\ Gr(B) &= K[x_1, \dots, x_n]. \end{aligned}$$

(ii) Quantum algebra $\mathcal{U}'(so(3, K))$, with $q \in K - \{0\}$:

$$x_2 x_1 - q x_1 x_2 = -q^{1/2} x_3, \quad x_3 x_1 - q^{-1} x_1 x_3 = q^{-1/2} x_2, \quad x_3 x_2 - q x_2 x_3 = -q^{1/2} x_1;$$

in this case $Gr(B) = K_{\mathbf{q}}[x_1, x_2, x_3]$ is the 3-multiparametric quantum space, i.e., a quantum polynomial ring in 3 variables, with

$$\mathbf{q} = \begin{bmatrix} 1 & q & q^{-1} \\ q^{-1} & 1 & q \\ q & q^{-1} & 1 \end{bmatrix}.$$

(iii) Dispin algebra $\mathcal{U}(osp(1, 2))$:

$$\begin{aligned} x_2 x_3 - x_3 x_2 &= x_3, \quad x_3 x_1 + x_1 x_3 = x_2, \quad x_1 x_2 - x_2 x_1 = x_1; \\ Gr(B) &= K_{\mathbf{q}}[x_1, x_2, x_3], \quad \text{with } \mathbf{q} = \begin{bmatrix} 1 & 1 & -1 \\ 1 & 1 & 1 \\ -1 & 1 & 1 \end{bmatrix}. \end{aligned}$$

(iv) Woronowicz algebra $\mathcal{W}_{\nu}(\mathfrak{sl}(2, K))$, where $\nu \in K - \{0\}$ is not a root of unity:

$$\begin{aligned} x_1 x_3 - \nu^4 x_3 x_1 &= (1 + \nu^2) x_1, \quad x_1 x_2 - \nu^2 x_2 x_1 = \nu x_3, \quad x_3 x_2 - \nu^4 x_2 x_3 = (1 + \nu^2) x_2; \\ Gr(B) &= K_{\mathbf{q}}[x_1, x_2, x_3], \quad \text{with } \mathbf{q} = \begin{bmatrix} 1 & \nu^{-2} & \nu^{-4} \\ \nu^2 & 1 & \nu^4 \\ \nu^4 & \nu^{-4} & 1 \end{bmatrix}. \end{aligned}$$

(v) Eight types of 3-dimensional skew polynomial algebras, with $\alpha, \beta, \gamma \in K - \{0\}$:

$$\begin{aligned} &x_2 x_3 - x_3 x_2 = x_3, \quad x_3 x_1 - \beta x_1 x_3 = x_2, \quad x_1 x_2 - x_2 x_1 = x_1; \\ &x_2 x_3 - x_3 x_2 = 0, \quad x_3 x_1 - \beta x_1 x_3 = x_2, \quad x_1 x_2 - x_2 x_1 = 0; \\ &x_2 x_3 - x_3 x_2 = x_3, \quad x_3 x_1 - \beta x_1 x_3 = 0, \quad x_1 x_2 - x_2 x_1 = x_1; \\ &x_2 x_3 - x_3 x_2 = x_3, \quad x_3 x_1 - \beta x_1 x_3 = 0, \quad x_1 x_2 - x_2 x_1 = 0; \\ &x_2 x_3 - x_3 x_2 = x_1, \quad x_3 x_1 - \alpha x_1 x_3 = x_2, \quad x_1 x_2 - x_2 x_1 = x_3; \\ &x_2 x_3 - x_3 x_2 = 0, \quad x_3 x_1 - \alpha x_1 x_3 = 0, \quad x_1 x_2 - x_2 x_1 = x_3; \\ &x_2 x_3 - x_3 x_2 = -x_2, \quad x_3 x_1 - \alpha x_1 x_3 = x_1 + x_2, \quad x_1 x_2 - x_2 x_1 = 0; \\ &x_2 x_3 - x_3 x_2 = x_3, \quad x_3 x_1 - \alpha x_1 x_3 = x, \quad x_1 x_2 - x_2 x_1 = 0; \\ Gr(B) &= K_{\mathbf{q}}[x_1, x_2, x_3], \quad \text{where } \mathbf{q} \text{ is an appropriate matrix in every case.} \end{aligned}$$

Observe that in every example, $Gr(B)$ is a noetherian Artin-Schelter regular algebra, and hence, $Gr(B)$ satisfies the \mathcal{X}_1 condition (see [15]). From this we will conclude that \underline{B} also satisfies such condition.

In fact, note first that in general there is an injective \mathbb{N} -graded homomorphism of B_0 -algebras $\eta : \underline{B} \rightarrow Gr(B)$ defined by

$$\begin{aligned} \bigoplus_{d=0}^{\infty} Hom_{sgr-B}(B, B(d)) &\xrightarrow{\eta} \bigoplus_{d=0}^{\infty} Gr(B)_d = \bigoplus_{d=0}^{\infty} \frac{B_0 \oplus \cdots \oplus B_d}{B_0 \oplus \cdots \oplus B_{d-1}} \\ f_0 + \cdots + f_d &\mapsto \overline{f_0(1)} + \cdots + \overline{f_d(1)}, \end{aligned}$$

with $f_i \in \text{Hom}_{\text{sgr}-B}(B, B(i))$, $0 \leq i \leq d$. We only check that η is multiplicative, the other conditions can be proved also easily: $\eta(f_n \star g_m) = \eta(s^n(g_m) \circ f_n) = \overline{(s^n(g_m) \circ f_n)(1)} = \overline{s^n(g_m)(f_n(1))} = \overline{g_m(f_n(1))} = \overline{f_n(1)g_m(1)} = \overline{f_n(1)} \overline{g_m(1)} = \eta(f_n)\eta(g_m)$.

Thus, in the examples above $K = B_0$, $(\underline{B})_0 \cong B_0 \cong \text{Gr}(B)_0$ and the kernel and cokernel of η are right bounded, so we can apply the part (5) of Lemma 8.2 in [3] and conclude that \underline{B} satisfies \mathcal{X}_1 .

We finish remarking that for the listed examples we can apply Proposition 6.11 and Theorem 6.12 and obtain that

$$\begin{aligned} \text{qgr} - K[x_1, x_2, x_3] &\simeq \text{qgr} - \Gamma(\pi(B))_{\geq 0}, \text{ with } B = \mathcal{U}(\mathcal{G}); \\ \text{qgr} - K_{\mathbf{q}}[x_1, x_2, x_3] &\simeq \text{qgr} - \Gamma(\pi(B))_{\geq 0}, \end{aligned}$$

with $B = \mathcal{U}'(\text{so}(3, K)), \mathcal{U}(\text{osp}(1, 2)), \mathcal{W}_{\nu}(\mathfrak{sl}(2, K))$ or any of eight types of 3-dimensional skew polynomial algebras above, and \mathbf{q} an appropriate matrix in every case.

Acknowledgements

The authors are grateful to Efim Zelmanov, Ivan Shestakov, Iryna Kashuba, Daniel Rogalski, Andrea Solotar, Mariano Suárez, Pablo Zadunaisky, Blas Torrecillas y Juan Cuadra for valuable comments and suggestions.

References

- [1] **Acosta, J.P., Chaparro, C., Lezama, O., Ojeda, I., and Venegas, C.**, *Ore and Goldie theorems for skew PBW extensions*, Asian-European Journal of Mathematics, 6 (4), 2013, 1350061-1; 1350061-20.
- [2] **Artin, M., Tate, J. and Van den Bergh, M.**, *Some algebras associated to automorphisms of elliptic curves*, The Grothendieck Festschrift, Vol. I, Birkhäuser Boston, 1990, 33-85.
- [3] **Artin, M. and Zhang J. J.**, *Noncommutative projective schemes*, Adv. Math. 109 (2), 1994, 228-287.
- [4] **Beaulieu, R. A., Jensen, A. and Jøndrup, S.**, *Towards a more general notion of Gelfand-Kirillov dimension*, Israel Journal of Mathematics, 93, 1996, 73-92.
- [5] **Gabriel, P.**, *Des catégories abéliennes*, Bull. Soc. Math. France 90, 1962, 323-448.
- [6] **Gaddis, J. D.**, *PBW deformations of Artin-Schelter regular algebras and their homogenizations*, Ph.D. thesis, The University of Wisconsin-Milwaukee, 2013.
- [7] **Gallego, C. and Lezama, O.**, *d-Hermite rings and skew PBW extensions*, to appear in São Paulo J. Math. Sci., 2016, DOI 10.1007/s40863-015-0010-8.
- [8] **Ginzburg, V.**, *Calabi–Yau algebras*, 2006, arXiv:math.AG/0612139v3.
- [9] **Goetz, P.**, *The Noncommutative Algebraic Geometry of Quantum Projective Spaces*, Ph.D. Thesis, University of Oregon, 2003.
- [10] **Grothendieck, A.**, *Sur quelques points d’algèbre homologique*, Tôhoku Math. J. (2) 9 (1957), 119-221; English translation by Marcia L. Barr and Michael Barr, September 10, 2010.
- [11] **Kanazawa, A.**, *Non-commutative projective Calabi-Yau schemes*, Journal of Pure and Applied Algebra, 19 (7), 2015, 2771-2780.

- [12] **Lezama, O. and Gallego, C.**, *Gröbner bases for ideals of sigma-PBW extensions*, *Communications in Algebra*, 39 (1), 2011, 50-75.
- [13] **Lezama, O. & Reyes, M.**, *Some homological properties of skew PBW extensions*, *Comm. in Algebra*, 42, (2014), 1200-1230.
- [14] **McConnell, J. and Robson, J.**, *Non-commutative Noetherian Rings*, *Graduate Studies in Mathematics*, AMS, 2001.
- [15] **Rogalski, D.**, *An introduction to non-commutative projective algebraic geometry*, arXiv:1403.3065 [math.RA].
- [16] **Smith, S.P.**, *Non-commutative algebraic geometry*, Department of Mathematics, University of Washington, 2000.
- [17] **Verevkin, A. B.**, *Serré injective sheaves*, *Mat. Zametki*, 1992, 52 (4), 35-41.
- [18] **Verevkin, A. B.**, *On a noncommutative analogue of the category of coherent sheaves on a projective scheme*, *Amer. Math. Soc. Transl.*, 1992, 151 (2), 41-53.