

On similarity of jet quenching and charmonia suppression

Martin Spousta^a

^aCharles University, Faculty of Mathematics and Physics, Prague, Czech Republic

Abstract

We quantify the magnitude and the color charge dependence of the medium induced parton energy loss in lead-lead collisions at the LHC using the data on inclusive jet suppression. The extracted color charge dependence shows that the difference between the in-medium loss of quarks and gluons is consistent with the difference between the radiation of quarks and gluons in the vacuum. Then, we examine the energy loss of prompt charmonia and we point to a remarkable similarity between the quenching of light-quark-initiated jets and the prompt charmonia suppression. Finally, we discuss possible sources of this similarity.

1. Introduction

Collisions between lead nuclei at the LHC produce colored medium where relevant degrees of freedom are deconfined quarks and gluons [1]. Hard scattering interactions in perturbative quantum chromodynamics (pQCD) lead to production of two highly virtual back-to-back partons (in the leading order of pQCD) which subsequently evolve as parton showers, hadronize, and are experimentally observed as back-to-back dijet. If partons traverse on their path a colored medium they lose energy which can be seen as modification of jet yields and jet properties. This phenomenon is commonly called “jet quenching” [2–4]. The first evidence of jet quenching at the LHC was provided by the measurement of jet pairs [5, 6]. Often, the magnitude of jet quenching is quantified by the nuclear modification factor, R_{AA} , which is the ratio of per-event-normalized yields in lead-lead (Pb+Pb) collisions to a cross-section in proton-proton (pp) collisions scaled by the nuclear overlap function. A precise measurement of jet R_{AA} was recently provided in Ref. [7]. Understanding the jet quenching means understanding the interaction of energetic color charges with the deconfined medium and the properties of this medium.

Not only jets but also charmonia can be used as tools to reveal the properties of the medium and its interactions. The production of charmonia in elementary collisions is often described in a nonrelativistic QCD effective-field-theory (EFT) framework [8]. In that theory, first, the $c\bar{c}$ pair

is produced either in a color singlet or color octet state. This “pre-resonant” $c\bar{c}$ pair then binds into a physical charmonium by non-perturbative evolution described in terms of long-distance matrix elements. The pre-resonant pair in the color octet state changes its color and spin by radiating off gluon(s) when evolving to the physical quarkonium state while the pair in the color singlet state retains these quantum numbers unchanged.

In nucleus-nucleus collisions, the production of charmonia was observed to be strongly suppressed with respect to proton-proton yield at LHC, RHIC, and SPS [9]. Generally, charmonia suppression can be described using the EFT framework at finite temperature in the limit of weakly coupled medium or in the lattice-QCD in the strongly coupled regime [10]. In the EFT framework at finite temperature the thermal part of $c\bar{c}$ potential has both a real and imaginary part [11]. The real part of the $c\bar{c}$ potential is connected with color screening of the potential in the deconfined medium. The imaginary part of the potential is connected with thermal decay and it is argued that it may be a dominant mechanism for the charmonia suppression within the EFT framework [12]. Since heavy quarks are not fully thermalized in the medium, further model assumptions need to be made in order to predict charmonium production rates. Examples of these models are transport models, statistical hadronization models, collisional dissociation models, or comover models which are reviewed in Ref. [9]. Different mechanisms leading to charmonia suppression that

are implemented in these models play different role at different kinematic regions and center-of-mass energies which makes the interpretation of similarities or differences in the charmonia suppression between the LHC, RHIC, and SPS complicated.

In proton-nucleus (or deuteron-nucleus) collisions, the production of charmonia was observed to be also modified with respect to proton-proton yield although the modifications are generally smaller than those seen in nucleus-nucleus collisions [9]. These modifications are due to an interplay of various effects such as effects of nuclear modifications to parton distribution functions (shadowing, anti-shadowing, EMC effect), multiple parton scattering, or nuclear absorption of bound state. These “cold nuclear matter effects” (CNM effects) are present in nucleus-nucleus collisions as well with a strength that depends on the choice of kinematic region. Therefore, they need to be considered when interpreting the nucleus-nucleus data.

This paper elaborates on the quantification of the jet quenching and it explores a similarity in the suppression of jets and prompt charmonia (that is those not coming from weak decay of B -hadrons) in $\sqrt{s_{NN}} = 2.76$ TeV Pb+Pb collisions at the LHC. The work is organized as follows. First, the effective suppression parameters of high- p_T quarks and gluons are extracted from jet suppression measurements. Then, these parameters are used to model the suppression of prompt charmonia. The result is compared with several recent measurements. Finally, the connection between the energy loss of jets and the charmonia suppression is discussed.

2. Single jet suppression

It was shown in a recent paper on the interpretation of single jet suppression measurements at the LHC [13] that many of the features seen in the data are driven by a primordial parton spectrum and the different quark and gluon energy loss. Further, it is concluded in that study that the suppression data on jets can be described by an effective quenching model (hereafter called EQ model) in which partons lose energy depending on their initial transverse momentum, $p_{T,ini}$, and an effective color factor, c_F , such that the total energy lost by a parton is

$$\Delta p_T = c_F \cdot s \cdot \left(\frac{p_{T,ini}}{p_{T,0}} \right)^\alpha. \quad (1)$$

Here s and α are free parameters of the model, $p_{T,ini}$ is the transverse momentum of a parton ini-

tiating a jet and $p_{T,0}$ is an arbitrary scale, here and in Ref. [13] set to 40 GeV. The parameter c_F was fixed to be 1 and 9/4 for light-quark-initiated jets and gluon-initiated jets, respectively. Input to the model was the parameterization of the unquenched jet- p_T spectra which was obtained from the PYTHIA8 generator [14]. The EQ model successfully described the representative data on the single jet suppression at high- p_T : the inclusive jet R_{AA} in various rapidity intervals [7], trends in the jet fragmentation functions [15, 16] at $p_T \gtrsim 10$ GeV and the inclusive charged particle R_{AA} at $p_T \gtrsim 20$ GeV [17].

The EQ model represents a model with minimal assumptions on the physics of the jet quenching process. The dynamics of the quenching process as well as the properties of the medium are encapsulated in a few free parameters. The fact that not only the inclusive jet suppression but also the jet structure is largely captured by the model speaks in favor of a physics picture in which a significant part of a parton shower remains unresolved by the medium. That may happen if the parton shower loses its energy coherently and subsequently fragments as in the vacuum. Indeed, it was recognized in several theoretical papers that such color coherence effects play an important role in the jet quenching process [18–21]. If the parton shower, or its large part, radiates as one object, one can ask if it is possible to find some similarities between the suppression of jets and a suppression of other objects with an internal structure. One such candidate are the charmonia. Before turning the attention to that question, the original model calls for an extension: the color factor should be extracted from the data and a possible dependence on the input spectra should be quantified.

The choice of $c_F = C_A/C_F = 9/4$ used in the EQ model for gluons represents an assumption on the difference in the probability to radiate a gluon from a gluon and quark source in the vacuum in the $Q \rightarrow \infty$ limit [22] or the soft limit [23]. The value of c_F can be different due to both the neglected non-leading corrections [24] and the presence of the deconfined medium. Thus, a global fit has been performed to extract s , α and c_F simultaneously. This has been done by minimizing the difference between the EQ model and measured jet R_{AA} [7] in different rapidity intervals and centrality bins. This procedure follows the logic of extracting this factor in the vacuum [25, 26].

To test for a sensitivity of the extracted sup-

$s = x \cdot N_{\text{part}} + y$	$x = (12.3 \pm 1.4) \cdot 10^{-3} \text{ GeV},$ $y = 1.5 \pm 0.2 \text{ GeV}$
α	0.52 ± 0.02
c_F	1.78 ± 0.12

Table 1: Parameters of the effective quenching model extracted from data. The $\chi^2/n_{\text{dof}} = 0.63$ for $n_{\text{dof}} = 287$. For details see the text.

pression parameters on the input jet spectra, the POWHEG MC generator [27, 28] interfaced to PYTHIA8 was used as an alternative to PYTHIA8 to provide a full next-to-leading order (NLO) jet p_T spectra. This NLO simulation was repeated three times using different input PDF sets (CT10 [29], MSTW2008NLO [30], and NNPDF2.0 [31]) which were selected to correspond to those used in the precise measurement of the jet cross-section in pp collisions at $\sqrt{s} = 7 \text{ TeV}$ [32]. These three sets of jet- p_T spectra were then parameterized and used along with the nominal PYTHIA8 as an input to the model which allowed to quantify uncertainties on extracted parameters.

The values of parameters from the global fit are summarized in Tab. 1. The global fit has revealed three interesting properties related to the energy loss: 1) the magnitude of the energy loss, s , depends linearly on the N_{part} (thus s is parameterized as a function of N_{part} in Tab. 1); 2) the power α is constant as a function of N_{part} (already observed in Ref. [13]); 3) c_F is consistent with the value calculated and measured in the vacuum which is ≈ 1.8 at the jet hardness $Q = 100 \text{ GeV}$ [24, 25]. This represents a precise effective quantification of the jet quenching in $\sqrt{s_{\text{NN}}} = 2.76 \text{ TeV}$ Pb+Pb data. Extracting c_F from the data using a model with minimal assumptions on the physics of the jet quenching should help to constrain the theory of this phenomenon.

3. Quarkonia suppression

There is no unique interpretation of the charmonia suppression measurements as of now [9]. New measurements at the LHC [33–38] should provide more insight to the mechanism of the charmonia suppression. The new precise measurements of the prompt J/ψ in the muon channel [33, 34] showed that the nuclear modification factor, $R_{\text{AA}}^{J/\psi}$, reaches a value of ~ 0.2 in the most central collisions ($N_{\text{part}} \gtrsim 350$), continuously grows up to a value

of $\sim 0.6 - 0.7$ reached in the most peripheral collisions ($N_{\text{part}} \lesssim 50$). The $R_{\text{AA}}^{J/\psi}$ exhibits only a weak (if any) dependence on the J/ψ momentum in the region of $p_T = 6.5 - 30 \text{ GeV}$ and $|y| < 2.4$. The dependence of $R_{\text{AA}}^{J/\psi}$ on the rapidity is also weak. More recently, a prompt production of $\psi(2S)$ was also measured in terms of a double ratio of measured yields, $(N_{\psi(2S)}/N_{J/\psi})|_{\text{Pb+Pb}}/(N_{\psi(2S)}/N_{J/\psi})|_{pp} = R_{\text{AA}}^{\psi(2S)}/R_{\text{AA}}^{J/\psi}$ [37]. It was shown that $\psi(2S)$ yields are suppressed by a factor of ~ 2 with respect to J/ψ in the range $|y| < 1.6$ and $6.5 < p_T < 30 \text{ GeV}$.

To test the idea of similarity in the physics of jet quenching and prompt charmonia suppression the EQ model described in the previous section has been employed. The input to the model are p_T spectra of J/ψ and $\psi(2S)$ and effective parameters obtained from the analysis of jet R_{AA} . The p_T spectra were obtained from PYTHIA8 (the same initial setup as in Ref. [13] was used) which was reweighted to reproduce the data measured in pp collisions at the 2.76 TeV [39]. It was found that PYTHIA reproduces well the shape of the J/ψ p_T spectra even without reweighting, while a reweighting was needed to reproduce the p_T spectra of $\psi(2S)$ (weights range from 0.2 to 6). The realistic p_T spectra were then used as an input to the EQ model which was run with two different settings of the color factor: first, corresponding to the color factor for the energy loss of light-quark initiated jets (defined to be one), and second, corresponding to the color factor extracted for the energy loss of gluon-initiated jets. The comparison of the data with the model is shown in Fig. 1. An excellent agreement of the model with the data for the case of the light-quark energy loss is seen. The N_{part} dependence of $R_{\text{AA}}^{J/\psi}$, its p_T and rapidity dependence are reproduced.

The ability of the model to reproduce the suppression of $\psi(2S)$ is shown in the Fig. 2 where the model is compared with the data on the ratio of nuclear modification factors, $R_{\text{AA}}^{\psi(2S)}/R_{\text{AA}}^{J/\psi}$, from Ref. [37]. Remarkably, a good agreement between the model and the data is seen. The model reproduces the measurement well except for the most peripheral collisions. Here, however, no significant $\psi(2S)$ signal was measured by CMS and consequently only a limit at 95% confidence level on the value of the ratio was provided.

The striking similarity between the measured J/ψ and $\psi(2S)$ suppression and the energy loss of jets suggests that the radiative energy loss may




Figure 1: The $R_{AA}^{J/\psi}$ predicted by the model compared to the data measured differentially in p_T of J/ψ (*left*), rapidity of J/ψ (*middle*), and the number of participants (*right*) [33, 34]. The model is evaluated in two configurations of the effective color factor – the color factor corresponding to light quarks and gluons.




Figure 2: The ratio of $R_{AA}^{\psi(2S)}$ to $R_{AA}^{J/\psi}$ predicted by the model compared to the data [37]. The model is evaluated in two configurations of the effective color factor – the color factor corresponding to light quarks and gluons.




Figure 3: The $R_{AA}^{J/\psi}$ predicted by the model compared to the data measured by ALICE at low p_T in the forward rapidity region [40].

be a dominant contribution to the energy loss of charmonia in the studied kinematic region as we will discuss in the next section. Now, we will discuss impact of feed-down contributions from excited states on presented results and the role of CNM effects. The measured charmonia production generally includes direct production from the hard interaction, as well as charmonium feed-down from excited states. While the feed-down contribution from excited resonances to $\psi(2S)$ state is minimal, the feed-down contribution to J/ψ is significant [9]. Thus, it is important to evaluate the impact of feed-down on presented results. The size of feed-down contribution from $\psi(2S)$ and χ_c decays to J/ψ were estimated to be 8% and 25%, respectively [41]. The

p_T dependence of the dominant feed-down contribution of χ_c to J/ψ was then evaluated in pp collisions at the LHC [42, 43]. In particular, for the central rapidity region of $|y| < 0.75$, it was found that the contribution from χ_c is on average 25% varying by only 2% in the p_T interval of 10 – 30 GeV. Also measured in pp collisions was the p_T spectrum of χ_c [43]. To access the impact of the feed-down on presented results, the measured χ_c and $\psi(2S)$ spectra and fractions of prompt J/ψ produced in χ_c and $\psi(2S)$ decays, respectively, were used to reweight the initial spectrum of J/ψ such that its shape reflects a contribution from excited states. The R_{AA} of J/ψ was then recalculated using this new spectrum. The new reweighted spectrum characterizes a physics picture in which it is always the excited

state which loses the energy in the medium while the J/ψ resulting from the decay of that excited state does not lose the energy. More realistic scenario is that only a fraction of χ_c or $\psi(2S)$ loses the energy while the rest first decays and then it loses the energy as J/ψ . Since we do not know such a fraction, the difference of reweighted R_{AA} and original R_{AA} was used as an estimate of the uncertainty of presented results. Maximal difference between the reweighted and original R_{AA} was found to be less than 10%. This estimate is similar to the estimate of the impact of feed-down on the nuclear modification factor measured in d+Au collisions at RHIC which was found to be 5% [44]. The uncertainty in the determination of R_{AA} due to the feed-down effects was combined with the uncertainty in the determination of R_{AA} due to uncertainties in parameters of the EQ model and is plotted on Figures 1 and 2 as a shaded band.

Besides feed-down effects, the interpretation of presented results might be obscured by initial-state effects or effects from final-state nuclear interactions that are unrelated to the presence of the hot deconfined medium, i.e. by CNM effects. Sizable suppression of J/ψ was seen in proton-lead (p+Pb) collisions in the forward rapidity region or at low p_T ($p_T \lesssim 6$ GeV) at the LHC [45–47] which clearly suggests that these effects come into the play. On the contrary, at high p_T ($p_T \gtrsim 6$ GeV) in the midrapidity region at the LHC, the nuclear modification factor measured in p+Pb collisions is consistent with unity within the precision of measurements [39, 45]. This observation is consistent with the estimate done using EKS98 nPDFs [48] which quantifies the impact of CNM effects, namely the shadowing, on the R_{AA} to be less than 10% in the kinematic region probed in this paper [9]. While this suggests that CNM effects should not affect the interpretation of the similarity of the jet quenching and charmonia suppression, there might still be a group of CNM effects that is common for both, charmonia and jets. Suggestive of that is a sizable suppression seen in the most peripheral bin for both, the charmonia and jets.

Figures 1 and 2 represent a selection of all available LHC data on prompt J/ψ and $\psi(2S)$ for which corresponding pp reference spectra exist which allows an exact data-to-model comparison. Besides those high- p_T R_{AA} measurements, ALICE Collaboration provided also measurement of $R_{AA}^{J/\psi}$ [36] for which the corresponding pp spectra were measured

[49]. This measurement presents R_{AA} of inclusive J/ψ (that is with no separation of non-prompt J/ψ contribution which is however smaller than 15% [50]) in the forward rapidity region and at low p_T . As discussed before, in this kinematic region CNM effects play a significant role. It was also shown in past that recombination and statistical hadronization processes likely play a significant role in this kinematic region [51–54]. Thus, we cannot expect that the model presented here will reproduce the data. Nevertheless, it is instructive to quantify a departure of the model from the data. This is shown in Fig. 3. One can see that the model can reproduce the $R_{AA}^{J/\psi}$ at $p_T \gtrsim 6$ GeV where ALICE data in the forward region agrees with CMS data measured in the midrapidity region. On the contrary, a significant departure of the model from the data is seen at low p_T .

4. Discussion

The ability of the EQ model to reproduce the suppression data on J/ψ and $\psi(2S)$ suggests that the measured shape of the suppression at high- p_T is driven by the shape of the initial spectra of charmonia and that the radiative energy loss may play a dominant role at high- p_T . We will leave the full analysis of this observation and the observation of similarity between the light-quark suppression and the charmonia suppression for separate works. Here, we will only summarize straightforward leading order calculations searching for a similarity between the radiation of quarks and charmonium. In particular, we discuss two basic scenarios: 1) charmonium is produced early in the collision in the color octet state and radiates coherently; 2) charmonium is produced late, from quarks or gluons that are already quenched.

For the first scenario, ratio is calculated of the probability of radiating a gluon from a charmonium in the color octet state, P_ψ , to the probability of radiating a gluon from a single light quark, P_q . This ratio is calculated in the collinear limit, leading to no change in the spin of charmonium due to the radiation, and under the assumption that the $c\bar{c}$ pair is not dissociated and radiates coherently in the range of virtualities from Q_{max}^ψ to Q_{min}^ψ . The result is

$$\frac{P_\psi(Q_{min}^\psi, Q_{max}^\psi)}{P_q(Q_{min}, Q_{max})} = \frac{c_\psi}{C_F} \left(1 + \ln \frac{k_{max}}{k_{min}} \cdot \ln^{-1} \frac{Q_{max}}{Q_{min}} \right), \quad (2)$$

where $c_\psi = C_A$ is the color factor for the radiation from charmonium. Constant k relates virtualities in the two processes, $Q^\psi = kQ$ ¹. The ratio (2) is equal to one if the virtuality range differs between the two processes (e.g. $k_{max}/k_{min} \approx 1/9$ for $Q_{min} = 0.2$ GeV and $Q_{max} = 10$ GeV). If this is the case, Eq. (2) predicts that the R_{AA} of J/ψ should start to deviate from the R_{AA} of light-quark initiated jets at higher transverse momenta due to the slow logarithmic dependence on the virtuality. In particular, at $p_T = 50$ GeV, the R_{AA} of light-quark initiated jets should be 1.3 times larger compared to the measured R_{AA} of J/ψ . This prediction can be tested in the LHC run 2 data. Not quantified in the formula is the dead cone effect which may also decrease the resulting radiation of the charmonium [56, 57]. Another factor not accounted for in this scenario is a presence of the charmonium in the color singlet state which might however represent a small contribution compared to the color octet state [58]. Irrespective of these important details, Eq. (2) implies that a simple expectation that the radiative energy loss of charmonia should be similar to the radiative energy loss of gluon-initiated jets is not valid since not only the color factors enter here but also kinematic range over which the two systems radiate need to be considered.

In the second scenario, the charmonium is produced late, from quarks or gluons that are already quenched. While the correlation between the p_T of the initial parton and the p_T of charmonium is rather strong, less steep spectra of charmonia would lead to generally larger R_{AA} of charmonia than the R_{AA} of partons. This was explicitly checked by modeling using PYTHIA. In this scenario, the similarity between the suppression of charmonia and light quarks would be accidental, steaming from an intriguing combination of quark and gluon energy loss which disfavors this scenario.

5. Summary

This paper has presented a quantification of the jet quenching in $\sqrt{s_{NN}} = 2.76$ TeV lead-lead collisions which is based on an extension of the effective quenching model [13]. In particular, extracted from the data was the effective color factor, c_F , characterizing a difference in the probability to radiate

¹Equation (2) holds also for the case of massive splitting functions calculated in the quasi-collinear limit [55].

a gluon from a gluon and quark source. The extracted value, $c_F = 1.78 \pm 0.12$, is consistent with the value obtained for vacuum [24, 25]. Using the quantification of the energy loss of jets, a similarity between the suppression of jets and prompt charmonia was explored. A striking similarity between the measured J/ψ and $\psi(2S)$ suppression and the energy loss of light-quark initiated jets was seen. While this observation requires a thorough theoretical analysis, it suggests that the radiative energy loss may be a dominant contribution to the charmonia suppression at $p_T \gtrsim 6$ GeV at the LHC. Quantification of the magnitude of the jet quenching together with observations made in this paper should improve the understanding of physics mechanism behind both, the jet quenching and charmonia suppression. They may also contribute to better understanding of general aspects of charmonia formation.

Acknowledgment

I'm grateful to Alexander Kupčo, Konrad Tywoniuk, Brian Cole, Jiří Dolejší, Daniel Scheirich, Yen-Jie Lee and Camelia Mironov for useful discussions. This work was supported by Charles University Research Development Scheme (PRVOUK) P45, UNCE 204020/2012 and MSMT grant INGO II LG15052.

References

- [1] F. Karsch, E. Laermann, Thermodynamics and in medium hadron properties from lattice QCD [arXiv:hep-lat/0305025](#).
- [2] U. A. Wiedemann, Jet Quenching in Heavy Ion Collisions (2010) 521–562 [Landolt-Bornstein23,521(2010)]. [arXiv:0908.2306](#), [doi:10.1007/978-3-642-01539-7_17](#).
- [3] A. Majumder, M. Van Leeuwen, The Theory and Phenomenology of Perturbative QCD Based Jet Quenching, *Prog. Part. Nucl. Phys.* A66 (2011) 41–92. [arXiv:1002.2206](#), [doi:10.1016/j.pnpnp.2010.09.001](#).
- [4] Y. Mehtar-Tani, J. G. Milhano, K. Tywoniuk, Jet physics in heavy-ion collisions, *Int. J. Mod. Phys. A*28 (2013) 1340013. [arXiv:1302.2579](#), [doi:10.1142/S0217751X13400137](#).
- [5] ATLAS Collaboration, Observation of a Centrality-Dependent Dijet Asymmetry in Lead-Lead Collisions at $\sqrt{s_{NN}} = 2.77$ TeV with the ATLAS Detector at the LHC, *Phys. Rev. Lett.* 105 (2010) 252303. [arXiv:1011.6182](#), [doi:10.1103/PhysRevLett.105.252303](#).
- [6] CMS Collaboration, Observation and studies of jet quenching in PbPb collisions at nucleon-nucleon center-of-mass energy = 2.76 TeV, *Phys. Rev. C*84 (2011) 024906. [arXiv:1102.1957](#), [doi:10.1103/PhysRevC.84.024906](#).

- [7] ATLAS Collaboration, Measurements of the Nuclear Modification Factor for Jets in Pb+Pb Collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS Detector, *Phys.Rev.Lett.* 114 (7) (2015) 072302. [arXiv:1411.2357](#), [doi:10.1103/PhysRevLett.114.072302](#).
- [8] G. T. Bodwin, E. Braaten, G. P. Lepage, Rigorous QCD analysis of inclusive annihilation and production of heavy quarkonium, *Phys. Rev. D* 51 (1995) 1125–1171, [Erratum: *Phys. Rev. D* 55, 5853 (1997)]. [arXiv:hep-ph/9407339](#), [doi:10.1103/PhysRevD.55.5853](#), [doi:10.1103/PhysRevD.51.1125](#).
- [9] A. Andronic, et al., Heavy-flavour and quarkonium production in the LHC era: from proton-proton to heavy-ion collisions, *Eur. Phys. J. C* 76 (3) (2016) 107. [arXiv:1506.03981](#), [doi:10.1140/epjc/s10052-015-3819-5](#).
- [10] N. Brambilla, et al., Heavy quarkonium physics [arXiv:hep-ph/0412158](#).
- [11] N. Brambilla, J. Ghiglieri, A. Vairo, P. Petreczky, Static quark-antiquark pairs at finite temperature, *Phys. Rev. D* 78 (2008) 014017. [arXiv:0804.0993](#), [doi:10.1103/PhysRevD.78.014017](#).
- [12] N. Brambilla, et al., Heavy quarkonium: progress, puzzles, and opportunities, *Eur. Phys. J. C* 71 (2011) 1534. [arXiv:1010.5827](#), [doi:10.1140/epjc/s10052-010-1534-9](#).
- [13] M. Spousta, B. Cole, Interpreting single jet measurements in Pb + Pb collisions at the LHC, *Eur. Phys. J. C* 76 (2016) 50. [arXiv:1504.05169](#), [doi:10.1140/epjc/s10052-016-3896-0](#).
- [14] T. Sjostrand, S. Ask, J. R. Christiansen, R. Corke, N. Desai, P. Ilten, S. Mrenna, S. Prestel, C. O. Rasmussen, P. Z. Skands, An Introduction to PYTHIA 8.2, *Comput. Phys. Commun.* 191 (2015) 159–177. [arXiv:1410.3012](#), [doi:10.1016/j.cpc.2015.01.024](#).
- [15] CMS Collaboration, Measurement of jet fragmentation in PbPb and pp collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Phys.Rev. C* 90 (2) (2014) 024908. [arXiv:1406.0932](#), [doi:10.1103/PhysRevC.90.024908](#).
- [16] ATLAS Collaboration, Measurement of inclusive jet charged-particle fragmentation functions in Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector, *Phys.Lett. B* 739 (2014) 320–342. [arXiv:1406.2979](#), [doi:10.1016/j.physletb.2014.10.065](#).
- [17] Study of high-pT charged particle suppression in PbPb compared to pp collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Eur. Phys. J. C* 72 (2012) 1945. [arXiv:1202.2554](#), [doi:10.1140/epjc/s10052-012-1945-x](#).
- [18] Y. Mehtar-Tani, C. A. Salgado, K. Tywoniuk, Angular ordering of gluon radiation in QCD media, *Phys. Rev. Lett.* 106 (2011) 122002. [arXiv:1009.2965](#), [doi:10.1103/PhysRevLett.106.122002](#).
- [19] Y. Mehtar-Tani, C. A. Salgado, K. Tywoniuk, Jets in QCD Media: From Color Coherence to Decoherence, *Phys. Lett. B* 707 (2012) 156–159. [arXiv:1102.4317](#), [doi:10.1016/j.physletb.2011.12.042](#).
- [20] J. Casalderrey-Solana, Y. Mehtar-Tani, C. A. Salgado, K. Tywoniuk, New picture of jet quenching dictated by color coherence, *Phys. Lett. B* 725 (2013) 357–360. [arXiv:1210.7765](#), [doi:10.1016/j.physletb.2013.07.046](#).
- [21] J.-P. Blaizot, E. Iancu, Y. Mehtar-Tani, Medium-induced QCD cascade: democratic branching and wave turbulence, *Phys. Rev. Lett.* 111 (2013) 052001. [arXiv:1301.6102](#), [doi:10.1103/PhysRevLett.111.052001](#).
- [22] K. Konishi, A. Ukawa, G. Veneziano, Jet Calculus: A Simple Algorithm for Resolving QCD Jets, *Nucl. Phys. B* 157 (1979) 45–107. [doi:10.1016/0550-3213\(79\)90053-1](#).
- [23] V. A. Khoze, S. Lupia, W. Ochs, Perturbative universality in soft particle production, *Eur. Phys. J. C* 5 (1998) 77–90. [arXiv:hep-ph/9711392](#), [doi:10.1007/s100529800818](#), [doi:10.1007/s100520050249](#).
- [24] A. Capella, I. M. Dremin, J. W. Gary, V. A. Nechitailo, J. Tran Thanh Van, Evolution of average multiplicities of quark and gluon jets, *Phys. Rev. D* 61 (2000) 074009. [arXiv:hep-ph/9910226](#), [doi:10.1103/PhysRevD.61.074009](#).
- [25] D. Acosta, et al., Measurement of charged particle multiplicities in gluon and quark jets in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV, *Phys. Rev. Lett.* 94 (2005) 171802. [doi:10.1103/PhysRevLett.94.171802](#).
- [26] G. Abbiendi, et al., Experimental properties of gluon and quark jets from a point source, *Eur. Phys. J. C* 11 (1999) 217–238. [arXiv:hep-ex/9903027](#), [doi:10.1007/s100520050628](#).
- [27] S. Alioli, P. Nason, C. Oleari, E. Re, A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX, *JHEP* 06 (2010) 043. [arXiv:1002.2581](#), [doi:10.1007/JHEP06\(2010\)043](#).
- [28] S. Alioli, K. Hamilton, P. Nason, C. Oleari, E. Re, Jet pair production in POWHEG, *JHEP* 04 (2011) 081. [arXiv:1012.3380](#), [doi:10.1007/JHEP04\(2011\)081](#).
- [29] H.-L. Lai, M. Guzzi, J. Huston, Z. Li, P. M. Nadolsky, J. Pumplin, C. P. Yuan, New parton distributions for collider physics, *Phys. Rev. D* 82 (2010) 074024. [arXiv:1007.2241](#), [doi:10.1103/PhysRevD.82.074024](#).
- [30] A. D. Martin, W. J. Stirling, R. S. Thorne, G. Watt, Parton distributions for the LHC, *Eur. Phys. J. C* 63 (2009) 189–285. [arXiv:0901.0002](#), [doi:10.1140/epjc/s10052-009-1072-5](#).
- [31] R. D. Ball, L. Del Debbio, S. Forte, A. Guffanti, J. I. Latorre, J. Rojo, M. Ubiali, A first unbiased global NLO determination of parton distributions and their uncertainties, *Nucl. Phys. B* 838 (2010) 136–206. [arXiv:1002.4407](#), [doi:10.1016/j.nuclphysb.2010.05.008](#).
- [32] CMS Collaboration, Measurement of the Inclusive Jet Cross Section in pp Collisions at $\sqrt{s} = 7$ TeV, *Phys.Rev.Lett.* 107 (2011) 132001. [arXiv:1106.0208](#), [doi:10.1103/PhysRevLett.107.132001](#).
- [33] CMS Collaboration, Suppression of non-prompt J/ψ , prompt J/ψ , and $Y(1S)$ in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *JHEP* 05 (2012) 063. [arXiv:1201.5069](#), [doi:10.1007/JHEP05\(2012\)063](#).
- [34] CMS Collaboration, J/ψ results from CMS in PbPb collisions, with 150mub-1 data, CMS-PAS-HIN-12-014.
- [35] ATLAS Collaboration, Measurement of the centrality dependence of J/ψ yields and observation of Z production in lead-lead collisions with the ATLAS detector at the LHC, *Phys. Lett. B* 697 (2011) 294–312. [arXiv:1012.5419](#), [doi:10.1016/j.physletb.2011.02.006](#).
- [36] ALICE Collaboration, Centrality, rapidity and transverse momentum dependence of J/ψ suppression in Pb-Pb collisions at $\sqrt{s_{NN}}=2.76$ TeV, *Phys. Lett. B* 734 (2014) 314–327. [arXiv:1311.0214](#), [doi:10.1016/j.physletb.2014.05.064](#).
- [37] CMS Collaboration, Measurement of Prompt $\psi(2S) \rightarrow J/\psi$ Yield Ratios in Pb-Pb and $p-p$ Collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Phys. Rev. Lett.* 113 (26) (2014) 262301. [arXiv:1410.1804](#), [doi:10.1103/PhysRevLett.](#)

- 113.262301.
- [38] ALICE Collaboration, Inclusive, prompt and non-prompt J/ψ production at mid-rapidity in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, JHEP 07 (2015) 051. [arXiv:1504.07151](#), [doi:10.1007/JHEP07\(2015\)051](#).
- [39] ATLAS Collaboration, Study of J/ψ and $\psi(2S)$ production in $\sqrt{s_{NN}} = 5.02$ TeV $p + Pb$ and $\sqrt{s} = 2.76$ TeV pp collisions with the ATLAS detector, ATLAS-CONF-2015-023.
- [40] ALICE Collaboration, J/ψ suppression at forward rapidity in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Rev. Lett. 109 (2012) 072301. [arXiv:1202.1383](#), [doi:10.1103/PhysRevLett.109.072301](#).
- [41] P. Faccioli, C. Lourenco, J. Seixas, H. K. Woehri, Study of ψ' and χ_c decays as feed-down sources of J/ψ hadro-production, JHEP 10 (2008) 004. [arXiv:0809.2153](#), [doi:10.1088/1126-6708/2008/10/004](#).
- [42] R. Aaij, et al., Measurement of the ratio of prompt χ_c to J/ψ production in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B718 (2012) 431–440. [arXiv:1204.1462](#), [doi:10.1016/j.physletb.2012.10.068](#).
- [43] ATLAS Collaboration, Measurement of χ_{c1} and χ_{c2} production with $\sqrt{s} = 7$ TeV pp collisions at ATLAS, JHEP 07 (2014) 154. [arXiv:1404.7035](#), [doi:10.1007/JHEP07\(2014\)154](#).
- [44] A. Adare, et al., Nuclear Modification of ψ' , χ_c , and J/ψ Production in d+Au Collisions at $\sqrt{s_{NN}}=200$ GeV, Phys. Rev. Lett. 111 (20) (2013) 202301. [arXiv:1305.5516](#), [doi:10.1103/PhysRevLett.111.202301](#).
- [45] ALICE Collaboration, Rapidity and transverse-momentum dependence of the inclusive J/ψ nuclear modification factor in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, JHEP 06 (2015) 055. [arXiv:1503.07179](#), [doi:10.1007/JHEP06\(2015\)055](#).
- [46] ALICE Collaboration, J/ψ production and nuclear effects in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, JHEP 02 (2014) 073. [arXiv:1308.6726](#), [doi:10.1007/JHEP02\(2014\)073](#).
- [47] R. Aaij, et al., Study of J/ψ production and cold nuclear matter effects in pPb collisions at $\sqrt{s_{NN}} = 5$ TeV, JHEP 02 (2014) 072. [arXiv:1308.6729](#), [doi:10.1007/JHEP02\(2014\)072](#).
- [48] K. J. Eskola, V. J. Kolhinen, C. A. Salgado, The Scale dependent nuclear effects in parton distributions for practical applications, Eur. Phys. J. C9 (1999) 61–68. [arXiv:hep-ph/9807297](#), [doi:10.1007/s100520050513](#), [doi:10.1007/s100529900005](#).
- [49] ALICE Collaboration, Inclusive J/ψ production in pp collisions at $\sqrt{s} = 2.76$ TeV, Phys. Lett. B718 (2012) 295–306, [Erratum: Phys. Lett. B748,472(2015)]. [arXiv:1203.3641](#), [doi:10.1016/j.physletb.2012.10.078](#), [doi:10.1016/j.physletb.2015.06.058](#).
- [50] J. Book, J/ψ production in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Nucl. Phys. A931 (2014) 591–595. [doi:10.1016/j.nuclphysa.2014.09.082](#).
- [51] X. Zhao, R. Rapp, Medium Modifications and Production of Charmonia at LHC, Nucl. Phys. A859 (2011) 114–125. [arXiv:1102.2194](#), [doi:10.1016/j.nuclphysa.2011.05.001](#).
- [52] Y.-p. Liu, Z. Qu, N. Xu, P.-f. Zhuang, J/ψ Transverse Momentum Distribution in High Energy Nuclear Collisions at RHIC, Phys. Lett. B678 (2009) 72–76. [arXiv:0901.2757](#), [doi:10.1016/j.physletb.2009.06.006](#).
- [53] E. G. Ferreira, Charmonium dissociation and recombination at LHC: Revisiting comovers, Phys. Lett. B731 (2014) 57–63. [arXiv:1210.3209](#), [doi:10.1016/j.physletb.2014.02.011](#).
- [54] A. Andronic, P. Braun-Munzinger, K. Redlich, J. Stachel, The thermal model on the verge of the ultimate test: particle production in Pb-Pb collisions at the LHC, J. Phys. G38 (2011) 124081. [arXiv:1106.6321](#), [doi:10.1088/0954-3899/38/12/124081](#).
- [55] S. Catani, S. Dittmaier, Z. Trocsanyi, One loop singular behavior of QCD and SUSY QCD amplitudes with massive partons, Phys. Lett. B500 (2001) 149–160. [arXiv:hep-ph/0011222](#), [doi:10.1016/S0370-2693\(01\)00065-X](#).
- [56] Y. L. Dokshitzer, D. E. Kharzeev, Heavy quark colorimetry of QCD matter, Phys. Lett. B519 (2001) 199–206. [arXiv:hep-ph/0106202](#), [doi:10.1016/S0370-2693\(01\)01130-3](#).
- [57] N. Armesto, C. A. Salgado, U. A. Wiedemann, Medium induced gluon radiation off massive quarks fills the dead cone, Phys. Rev. D69 (2004) 114003. [arXiv:hep-ph/0312106](#), [doi:10.1103/PhysRevD.69.114003](#).
- [58] S. P. Baranov, A. V. Lipatov, N. P. Zotov, Prompt charmonia production and polarization at LHC in the NRQCD with k_T -factorization. Part I: $\psi(2S)$ meson, Eur. Phys. J. C75 (9) (2015) 455. [arXiv:1508.05480](#), [doi:10.1140/epjc/s10052-015-3689-x](#).