# Egyptian Fractions with odd denominators

Christian Elsholtz\*

#### Abstract

The number of solutions of the diophantine equation  $\sum_{i=1}^{k} \frac{1}{x_i} = 1$ , in particular when the  $x_i$  are distinct odd positive integers is investigated. The number of solutions S(k) in this case is, for odd k:

$$\exp\left(\exp\left(c_1 \frac{k}{\log k}\right)\right) \le S(k) \le \exp\left(\exp\left(c_2 k\right)\right)$$

with some positive constants  $c_1$  and  $c_2$ . This improves upon an earlier lower bound of  $S(k) \ge \exp\left((1+o(1))\frac{\log 2}{2}k^2\right)$ .

### 1 Introduction

In this paper we study the number of solutions of the diophantine equation

(1.1) 
$$\sum_{i=1}^{k} \frac{1}{x_i} = 1$$

in particular, where the  $x_i$  have some restrictions, such as all  $x_i$  are distinct odd positive integers. Let us first review what is known for distinct positive

<sup>\*</sup>Institut für Analysis und Zahlentheorie, Graz University of Technology, Kopernikusgasse 24, A-8010 Graz, Austria. E-mail: elsholtz@math.tugraz.at

<sup>2010</sup> Mathematics Subject Classification: 11D68, 11D72.

*Key words and phrases*: Egyptian fractions; number of solutions of Diophantine equations.

integers, without further restriction: Let

$$\mathcal{X}_k = \{ (x_1, x_2, \dots, x_k) : \sum_{i=1}^k \frac{1}{x_i} = 1, \quad 0 < x_1 < x_2 < \dots < x_k \}.$$

It is known that

(1.2) 
$$\exp\left(\exp\left(\left((\log 2)(\log 3) + o(1)\right)\frac{k}{\log k}\right)\right) \le |\mathcal{X}_k| \le c_0^{(\frac{5}{3}+\varepsilon)2^{k-3}}$$

where  $c_0 = 1.264...$  is  $\lim_{n \to \infty} u_n^{1/2^n}$ ,  $u_n = 1, u_{n+1} = u_n(u_n + 1)$ .

The lower bound is due to Konyagin [12], the upper bound due to Browning and Elsholtz [3]. Earlier results on the upper and lower bounds were due to Sándor [13] and Erdős, Graham and Straus (see [9], page 32).

The set of solutions has also been investigated with various restrictions on the variables  $x_i$ . A quite general and systematic investigation of expansions of  $\frac{a}{b}$  as a sum of unit fractions with restricted denominators is due to Graham [10]. Elsholtz, Heuberger, Prodinger [7] gave an asymptotic formula for the number of solutions of (1.1), with two main terms, when the  $x_i$  are (not necessarily distinct) powers of a fixed integer t.

Another prominent case is when all denominators  $x_i$  are odd. Sierpiński [16] proved that a nontrivial solution exists. It is known that for k = 9there are exactly 5 solutions, and for k = 11, there are exactly 379,118 solutions (see [15, 2]). Chen, Elsholtz and Jiang [4] showed that for odd denominators  $x_i$  the number of solutions of (1.1) is increasing with a lower bound of  $\sqrt{2}^{k^2(1+o(1))}$ . Other types of restrictions on the denominator have been studied, e.g. by Croot [5] and Martin [11]. The number of solutions of the equation  $\frac{m}{n} = \sum_{i=1}^{k} \frac{1}{x_i}$  have also been estimated by Elsholtz and Tao [8].

In this paper we take inspiration from the proof of Chen et al. [4] for odd denominators, and the proof of Konyagin [12] for lower bounds in the case of unrestricted  $x_i$ . As Konyagin's proof makes crucial use of ingenious identities, involving a lot of even numbers, it seems unclear whether one can generalize it to odd integers. Here is our main result:

**Theorem 1.1.** Let  $s \ge 1$  and let  $\{p_1, \ldots, p_s\}$  denote a set of primes, and let  $P = p_1 \cdots p_s$  be squarefree. Let k be sufficiently large. Moreover, if P is

even, let k be odd. Let

$$\mathcal{X}_{k,P} = \{(x_1, x_2, \dots, x_k) : \sum_{i=1}^k \frac{1}{x_i} = 1, \text{ with distinct positive } x_i \equiv \pm 1 \mod P\}$$

There is some positive constant c(P) such that the following holds:

$$|\mathcal{X}_{k,P}| \ge \exp\left(\exp\left(c(P)\frac{k}{\log k}\right)\right).$$

The case P = 2 is the case of odd denominators:

Corollary 1.2. Let k be odd, and

$$\mathcal{X}_{k,\text{odd}} = \{(x_1, x_2, \dots, x_k) : \sum_{i=1}^k \frac{1}{x_i} = 1, \text{ with odd distinct positive } x_i\}.$$

There is some positive constant c such that the following holds:

$$|\mathcal{X}_{k,\text{odd}}| \ge \exp\left(\exp\left(c\frac{k}{\log k}\right)\right).$$

For comparison, an upper bound of type  $\exp(\exp(c_2 k))$  follows from the unrestricted case, see (1.2).

## 2 Proof

**Lemma 2.1.** Let P > 1 be a squarefree integer. Let  $\omega(n)$  denote the number of distinct prime factors of n, and d(m) the number of divisors of n. The following holds:  $\omega(P^m - 1) \ge d(m) - 6$ .

*Proof.* Due to a result of Bang, Zsigmondy, Birkhoff and Vandiver (see e.g. Schinzel [14]), it is known that for n > 6 the values of  $P^n - 1$  have at least one *primitive* prime factor. (A prime factor of the sequence  $P^n - 1$  is primitive if it divides  $P^n - 1$ , but does not divide any  $P^m - 1$  with m < n.).

Let  $m = m_1 m_2$ . For each divisor  $m_1$  one has the factorization

$$P^{m} - 1 = (P^{m_1} - 1)(P^{m_1m_2 - m_1} + P^{m_1m_2 - 2m_1} + \dots + P^{m_1} + 1),$$

hence the number of prime factors of  $P^m - 1$  is at least the sum of the number of primitive prime factors of  $P^{m_1} - 1$ , for all possible divisors  $m_1$  of m.

**Lemma 2.2.** For  $X \ge 3$ , there exists a natural number m < X such that  $d(m) > \exp\left((\ln 2 + o(1))\frac{\ln X}{\ln \ln X}\right)$  as  $X \to \infty$ .

This follows from a theorem of Wigert [17], but can also be seen directly. Let  $P_r = \prod_{i=1}^r q_i$  be the product over the first primes, and choose  $m = P_r$ , if  $P_r \leq X < P_{r+1}$ . Then  $d(m) = 2^r = \exp\left((\ln 2 + o(1))\frac{\ln m}{\ln \ln m}\right) = \exp\left((\ln 2 + o(1))\frac{\ln X}{\ln \ln X}\right)$ . Taking the first r odd primes, one can also find an odd number m of this type.

**Lemma 2.3.** For every  $a, b, n_0 \in \mathbb{N}$  the following holds: every positive integer can be written as a finite sum of distinct fractions of the form  $\frac{1}{an+b}$ ,  $n \ge n_0$ .

This result with  $n_0 = 0$  was originally proved by van Albada and van Lint [1]. The result for general  $n_0$  easily follows by using the progression  $a'n + b' = an + (an_0 + b), n \ge 0$ .

As an easy consequence we have:

Lemma 2.4. There exist distinct positive integers

$$l_1, \ldots, l_{r_1}, m_1, \ldots, m_{r_2}, n_1, \ldots, n_{r_3},$$

all larger than 1, in the residue class  $1 \mod 3P(P^2-1)$  such that the following holds:

$$\sum_{i=1}^{r_1} \frac{1}{l_i} = P - 2, \qquad \sum_{i=1}^{r_2} \frac{1}{m_i} = 1, \qquad \sum_{i=1}^{r_3} \frac{1}{n_i} = P,$$

If P = 2, then  $r_1 = 0$ , otherwise  $r_1, r_2, r_3 > 0$ . Moreover, it is clear that  $r_2 \equiv 1 \mod P$ .

*Proof of Theorem.* The idea employed in [4] and [12] is to write 1 as a sum of fractions where one denominator has a large number of divisors, and to split this fraction recursively into several fractions, where (at least) one of these has again a large number of divisors.

Here we show that it is possible to have, for any given  $t \in \mathbb{N}$ , the fraction  $\frac{1}{P^t-1}$  as one of these fractions. Let us start with the trivial decomposition

$$1 = \frac{1}{P-1} + \frac{P-2}{P-1}.$$

In order to avoid that the denominator P-1 occurs more than once we use Lemma 2.4 to write the integer P-2 as a sum of distinct unit fractions, with  $l_i > 1$ :  $P-2 = \sum_{i=1}^{r_1} \frac{1}{l_i}$ .

Next we observe that any fraction  $\frac{1}{P^n-1}$  can be decomposed to obtain a sum of unit fractions containing a)  $\frac{1}{P^{2n}-1}$  or b)  $\frac{1}{P^{n+1}-1}$ .

(a) 
$$\frac{1}{P^n - 1} = \frac{1}{P^n + 1} + \frac{1}{P^{2n} - 1} + \sum_{i=1}^{r_2} \frac{1}{(P^{2n} - 1)m_i}$$

By Lemma 2.4

$$1 = \sum_{i=1}^{r_2} \frac{1}{m_i}, \quad m_i \equiv 1 \text{ mod } 3P(P^2 - 1), m_i > 1 \text{ and distinct.}$$

Note that all occurring denominators are distinct, with the possible exception that  $P^n + 1 = P^{2n} - 1$  holds if P = 2, n = 1. In this case, one rewrites  $\frac{1}{P+1} = \frac{1}{3} = \sum_{i=1}^{r_2} \frac{1}{3m_i}$ . These denominators have not been used before, as the  $l_i$  or  $m_i$  are congruent to 1 mod 3, whereas the new denominators  $3m_i$  are not.

(b) 
$$\frac{1}{P^n-1} = \frac{1}{P^{n+1}-1} + \frac{P-1}{(P^n-1)(P^{n+1}-1)} + \frac{P-1}{(P^{n+1}-1)}$$

Note that these three fractions are unit fractions, as the denominators are divisible by P - 1. These three fractions are distinct, unless n = 1. In this case the fraction  $\frac{1}{P^2-1}$  occurs twice and one of these is rewritten as  $\frac{1}{P^2-1} = \sum_{i=1}^{r_2} \frac{1}{(P^2-1)m_i}$ . These denominators have not been used before, as the previous denominators  $l_i$  and  $m_i$  were by construction congruent to 1 mod  $P^2 - 1$ . Also,  $P^n + 1$ ,  $P^{2n} - 1$ ,  $(P^{2n} - 1)m_i$  are new.

For constructing a solution with  $\frac{1}{P^t-1}$  we write t in binary. The first binary digit is of course 1. For the positions  $i \ge 2$  we perform two different types of steps, corresponding to (a) and (b) above:

1) If the *i*-th leading position is a 0, then we take the "doubling" a).

2) If the *i*-th leading position is a 1, then we first take the doubling a), followed by an "addition" b),

For example, if  $t = 53 = 110101_2$  and starting from left to right:

i = 1		2	3		4	5		6
1		1	0		1	0		1
	a	b	a	a	b	a	a	b
n = 1	2	3	6	12	13	26	52	53

Generally, any integer t can be obtained in at most  $2\frac{\log t}{\log 2}$  such steps a) or b). In other words, starting from n = 1 we can obtain a decomposition

$$1 = \frac{1}{P^t - 1} + \sum_{i=1}^{k'-1} \frac{1}{x_i}$$

with  $k' = O(r_1 + r_2 \log t + r_3) = O_P(\log t)$  unit fractions. Observe that all denominators have been rearranged to be distinct.

We next come to the most crucial step, which determines the number of solutions:

**Lemma 2.5.** Let  $\sum_{i=1}^{r_3} \frac{1}{n_i} = P$  (by Lemma 2.4).

a) For any divisor  $d|(P^t - 1)$  the following is an identity.

$$\frac{1}{P^t - 1} = \frac{1}{P^t - 1 + Pd} + \sum_{i=1}^{r_3} \frac{1}{\frac{P^t - 1}{d}(P^t - 1 + Pd)n_i}$$

- b) The number of divisors  $d|P^t 1$  with  $d \equiv 1 \mod P$  is at least  $2^{\frac{\omega(P^t 1)}{P}}$
- c) If  $d \equiv 1 \mod P$ , then all denominators are  $\pm 1 \mod P$ .

Part a) and c) are easy to verify. For part b) observe: For any P prime factors  $p_k$ , being coprime to P, there is at least one subset of these primes, whose product is 1 mod P. Indeed, the sequence  $a_1 = p_1, a_2 = p_1 p_2, ..., a_P =$  $\prod_{k=1}^{P} p_k$  must have two members  $a_i, a_j$ , say, which are equivalent modulo P. Then  $\frac{a_i}{a_i} = \prod_{k=i+1}^{j} p_k \equiv 1 \mod P$ . Therefore, the number of divisors  $d \equiv 1 \mod P$  is at least  $2^{\frac{\omega(P^t-1)}{P}}$ . (Clearly, this argument can be refined (see e.g. [6]), but this would not improve our final result.) All solutions produced in this way are distinct, as each solution has a unique denominator  $P^t - 1 + Pd$ . Moreover, as all these denominators are greater than  $P^t$ , and as in our application t will be chosen large, these new denominators are greater than those that have been used before.

We choose t as a product of the first primes. By Lemma 2.2 the number of divisors, and hence the number of solutions satisfies:

$$|\mathcal{X}_{k,P}| \ge 2^{\omega(P^t - 1)/P} \ge 2^{(d(t) - 6)/P} \ge 2^{\exp\left(\frac{\log 2 + o(1)}{P} \frac{\log t}{\log \log t}\right)} \ge \exp\left(\exp(c(P)k/\log k)\right)$$

Recall that the number of fractions is  $k = O_P(\log t)$ .

Finally let us comment on the condition that k is odd, (see statement of the Theorem), when P is even. By multiplying equation (1.1) by its common denominator, and reducing modulo P it is clear that this condition is necessary. The condition is also sufficient as in view of step a) we can replace one fraction by  $r_2 + 2$  fractions. Again, by the same argument  $r_2 \equiv 1 \mod P$ , so that effectively we replace one fraction by 3 fractions (modulo P). Iterating this, we can reach any residue class modulo P, when P is odd, and the odd residue classes, when P is even. The number of extra fractions required is  $O(P r_2) = O_P(1)$ . This does not influence the overall result. In any case, the theorem is valid for sufficiently large  $k \geq k_P$ , with this necessary and sufficient congruence obstruction.

**Remark 2.6.** We have not worked out the constant c(P). One may observe that c(P) might be as small as  $\frac{1}{r_2}$ . To estimate  $r_2$  one observes that  $\sum_{i=1}^{r_2} \frac{1}{i \, 3P(P^2-1)-1} \geq \sum_{i=1}^{r_2} \frac{1}{m_i} \approx \frac{\log r_2}{3P(P^2-1)} > P$  must hold. Hence  $r_2$  appears to be at least of exponential growth in P. Taking denominators  $x_i$  only coprime to P, but not necessarily restricted to  $x_i \equiv \pm 1 \mod 3P(P^2-1)$  would improve this constant c(P).

I would like to thank Sergei Konyagin for insightful comments on the problem during a conference at CIRM (Luminy). The paper has been completed during a very pleasant stay at Forschungsinstitut Mathematik (FIM) at ETH Zürich.

#### References

- P.J. van Albada, J.H. van Lint, Reciprocal bases for the integers. Amer. Math. Monthly 70 (1963), 170–174.
- [2] R. Arce-Nazario, R. Castro, F. Figueroa, R. On the number of solutions of  $\sum_{i=1}^{11} \frac{1}{x_i} = 1$  in distinct odd natural numbers, J. Number Theory 133 no. 6, (2013), 2036–2046.
- [3] T. D. Browning and C. Elsholtz. The number of representations of rationals as a sum of unit fractions. Illinois J. Math. 55 (2011), no. 2, 685–696.
- [4] Y.-G. Chen, C. Elsholtz, and L.-L. Jiang. Egyptian fractions with restrictions. Acta Arith. 154 (2012), no. 2, 109–123.
- [5] E.S. Croot, On a coloring conjecture about unit fractions. Ann. of Math.
  (2) 157 (2003), no. 2, 545–556.
- [6] M. Drmota, M. Skałba, Equidistribution of divisors and representations by binary quadratic forms, Int. J. Number Theory 9 (2013), 2011–2018.
- [7] C. Elsholtz, C. Heuberger, and H. Prodinger. The number of Huffman codes, compact trees, and sums of unit fractions. IEEE Trans. Inform. Theory 59 (2013), no. 2, 1065–1075.
- [8] C. Elsholtz and T. Tao. Counting the number of solutions to the Erdős-Straus equation on unit fractions. J. Aust. Math. Soc. 94 (2013), no. 1, 50–105.
- [9] P. Erdős and R.L. Graham, Old and new problems and results in combinatorial number theory. Monographie No.28 de L'Enseignement Mathématique. Genève, 128 p. (1980).
- [10] R. L. Graham, On finite sums of unit fractions, Proc. London Math. Soc. 14 (1964), 193–207.

- [11] G. Martin, Dense Egyptian fractions, Trans. Amer. Math. Soc. 351 (1999), 3641–3657.
- [12] S. V. Konyagin, Double Exponential Lower Bound for the Number of Representations of Unity by Egyptian Fractions, Math. Notes 95 (2014), no. 1-2, 277–281.
- [13] C. Sándor, On the number of solutions of the Diophantine equation  $\sum_{i=1}^{n} \frac{1}{x_i} = 1$ . Period. Math. Hungar. 47 (2003), no. 1-2, 215–219.
- [14] A. Schinzel, On primitive prime factors of  $a^n b^n$ , Proc. Camb. Phil. Soc. 58 (1962), 555–562.
- [15] P. Shiu, Egyptian fraction representations of 1 with odd denominators, Math. Gaz., 93 (2009), pp. 271–276.
- [16] W. Sierpiński, Sur les décompositions de nombres rationnels en fractions primaires, Mathesis 65 (1956), 16–32.
- [17] S. Wigert, Sur l'ordre grandeur du nombre de diviseurs d'un entier. Ark. Mat. 3, no. 18 (1907), 1–9.