TSIRELSON'S PROBLEM AND AN EMBEDDING THEOREM FOR GROUPS ARISING FROM NON-LOCAL GAMES

WILLIAM SLOFSTRA

ABSTRACT. Tsirelson's problem asks whether the commuting operator model for two-party quantum correlations is equivalent to the tensor-product model. We give a negative answer to this question by showing that there are nonlocal games which have perfect commuting-operator strategies, but do not have perfect tensor-product strategies. The weak Tsirelson problem, which is known to be equivalent to Connes embedding problem, remains open.

The examples we construct are instances of (binary) linear system games. For such games, previous results state that the existence of perfect strategies is controlled by the *solution group* of the linear system. Our main result is that every finitely presented group embeds in some solution group. As an additional consequence, we show that the problem of determining whether a linear system game has a perfect commuting-operator strategy is undecidable.

CONTENTS

1. Introduction	2
2. Quantum correlation sets and linear system games	6
2.1. Quantum correlation sets	6
2.2. Linear system games, hypergraphs, and solution groups	9
3. The embedding theorem and consequences	11
4. Presentations by involutions	14
5. The wagon wheel embedding	18
6. Pictures for groups generated by involutions	20
6.1. Pictures as planar graphs	21
6.2. Groups and labellings of pictures	25
7. Pictures over solution groups and hypergraphs	28
8. A category of hypergraphs	31
9. Cycles and outer faces	38
10. Pictures over suns	41
11. Stellar cycles and constellations	48
11.1. Structure of constellations	49
11.2. Stellar cycles	50
11.3. Covers versus copies	52
11.4. Cycles covered by other cycles	54
11.5. Proof of Theorem 11.4	55

Partially supported by NSERC grant number 2018-03968.

12. Proof of the embedding theorem References

1. INTRODUCTION

In a two-player non-local game, the players, commonly called Alice and Bob, are physically separated and unable to communicate. They each receive a question chosen at random from a finite question set, and reply with a response from a finite answer set. If the joint answers meet a predetermined winning condition dependent on the joint questions, then Alice and Bob win; otherwise they lose. The rules of the game, including the winning condition and distribution on questions, are completely known to Alice and Bob, and they can arrange in advance a strategy which will maximize their success probability. However, since they cannot communicate during the game, they may not be able to play perfectly, i.e. win with probability one.

Classically, Alice and Bob's strategy for a non-local game is described by a local hidden variable model. Bell's famous theorem states that Alice and Bob can achieve better results than is possible with a local hidden variable model if they share an entangled quantum state [Bel64]. Since Bell's discovery, non-local games have been heavily studied¹ in physics, mathematics, and computer science; see [CHSH69, FC72, Cir80, AGR82, CHTW04, BPA⁺08, NPA08, JPPG⁺¹⁰, KV11, KKM⁺¹¹, HBD⁺¹⁵ for a small sample of results. Despite this, a number of foundational questions remain open, chief among which is Tsirelson's problem: A quantum strategy for a non-local game can be described as a set of measurement operators on Hilbert spaces H_A and H_B , along with a quantum state in the joint space $H = H_A \otimes H_B$. We refer to this as the *tensor-product model*. While the tensor-product model is often the default, there is another choice: a quantum strategy can be described as a set of measurements and quantum state on a shared Hilbert space H, with the property that Alice's measurement operators commute with Bob's measurement operators. This *commuting-operator model* is used, for instance, in algebraic quantum field theory [HK64]. The observable consequences of a strategy in either model are captured by the correlation matrix of the strategy. If H is finite-dimensional, every correlation matrix arising from a commutingoperator strategy can be realized using a tensor-product strategy. Tsirelson's problem asks whether this is true for a general Hilbert space.

This problem has an interesting history. Tsirelson originally stated the problem in a survey on Bell inequalities [Tsi93], and claimed without proof

¹Usually under the name "Bell tests" or "Bell inequalities". The term "non-local games" is more recent.

that the two models gave rise to the same set of correlation matrices. He later retracted this claim, and posted the question to a list of open problems in quantum information theory [Tsi06]. Tsirelson and subsequent authors [NPA08, SW08, JNP+11, Fri12, NCPGV12, PT15, DP16, PSS+16] studying Tsirelson's problem have considered several different sets of tensor-product strategies. Paulsen and Todorov [PT15] (see also Dykema and Paulsen [DP16]) observe that these variations lead to a hierarchy of sets of correlation matrices

(1.1)
$$C_q \subseteq C_{qs} \subseteq C_{qa} \subseteq C_{qc},$$

where C_q is the set of correlations arising from tensor-product strategies on finite-dimensional Hilbert spaces, C_{qs} is the set of correlations arising from tensor-product strategies (on possibly infinite-dimensional Hilbert spaces) with a vector state, $C_{qa} = \overline{C_{qs}}$ is the set of correlations which are limits of correlations in C_{qs} , and C_{qc} is the set of correlations arising from commuting-operator strategies. If we restrict to non-local games with question sets of size n and answer sets of size m, then all these sets are convex subsets of $\mathbb{R}^{m^2n^2}$, and prior to this paper, none of the inclusions (including $C_q \subseteq C_{qc}$) were known to be strict. Thus for each $t \in \{q, qs, qa\}$ there is a Tsirelson problem asking whether $C_{qc} = C_t$. Ozawa, building on the work of Junge, Navascués, Palazuelos, Pérez-García, Sholz, and Werner [JNP+11] and work of Fritz [Fri12], has shown that $C_{qc} = C_{qa}$ if and only if Connes' embedding conjecture is true [Oza13]. At the other end of the hierarchy, if C_q was equal to C_{qc} then every correlation matrix, whether commuting operator or tensor-product, would arise from a finite-dimensional Hilbert space. The "middle" version, which asks whether $C_{qc} = C_{qs}$, seems closest to Tsirelson's original claim.

The first main result of this paper is that there is a non-local game which can be played perfectly with a commuting-operator strategy, but which cannot be played perfectly using a tensor-product strategy with a vector state. Thus we resolve the middle version of Tsirelson's problem by showing that $C_{qc} \neq C_{qs}$. This also shows that $C_{qc} \neq C_q$. As a result, this game is interesting from the perspective of quantum information and computation, where a non-local game is often regarded as a computational scenario in which better results can be achieved with entanglement as a resource. From this point of view, it is natural to ask how much entanglement is needed to play a game optimally, and in particular, whether every game can be played optimally on a finitedimensional Hilbert space. The game we construct shows that this is not possible, at least if we allow commuting-operator strategies. Previously-known examples of this type have involved either quantum questions [LTW13, RV15] or infinite answer sets [MV14].

One reason we would have desired that every game have an optimal strategy on a finite-dimensional Hilbert space is that it would make it possible to

determine the optimal winning probability of a non-local game over entangled strategies. At present the only known methods for this task, aside from bruteforce search over strategies, are variants of the Navascués-Pironio-Acín (NPA) hierarchy [NPA08, DLTW08]. Given a non-local game, the NPA hierarchy provides a sequence of upper bounds which converge to the optimal winning probability in the commuting-operator model. However, the hierarchy does not provide, outside of special cases, a stopping criterion, i.e. a way to tell if the value will fall below a given threshold. Our second main result is that it is undecidable to determine if a non-local game can be played perfectly with a commuting-operator strategy. In particular, this implies that there is no stopping criterion for the NPA hierarchy which applies to all games.

The games we consider are binary linear system games, so named because they arise from linear systems over \mathbb{Z}_2 . Such games have been studied previously in [CM14, Ark12, Ji13]. Cleve and Mittal implicitly associate a certain group to every linear system over \mathbb{Z}_2 , such that perfect tensor-product strategies for the game correspond to certain finite-dimensional representations of the group [CM14]. We call this group, which is analogous to the solution space of a linear system, the *solution group*. In [CLS16] it is shown that perfect commuting-operator strategies for a binary linear system game correspond to certain possibly-infinite-dimensional representations of the solution group. Solution groups form an interesting class of groups. They are finitely presented, but their presentations must satisfy a property which in [CLS16] is called *local compatibility*: if

$x_1 \cdots x_n = 1$

is a relation, where x_i , $1 \leq i \leq n$, are not necessarily distinct generators of the group, then the presentation must also contain the relations $x_i x_j = x_j x_i$ for all $1 \leq i, j \leq n$. This condition is natural from the perspective of quantum mechanics, where two observables commute if and only if the observables correspond to quantities which can be measured (or known) simultaneously. Group relations of this exact type can be found in contextuality theorems of Mermin and Peres [Mer90, Per90, Mer93]. Local compatibility is a priori a strong constraint on group presentations. Our primary result, on which our other two results are based, is that any finitely presented group can be embedded in a solution group. This embedding theorem allows us to extend results from combinatorial group theory to solution groups. In particular, the embedding theorem implies the existence of a non-residually-finite solution group, and this leads directly to the separation $C_{qc} \neq C_{qs}$. Using non-residually-finite groups to recognize infinite-dimensional state spaces was previously proposed in [Fri13].

There have been a number of related results since this paper was first posted. The proof of the embedding theorem takes up most of this paper, and it's

natural to ask whether there might be a shorter proof. In a follow-up paper [Slo17], we give a shorter and more direct proof of an embedding theorem for a restricted class of finitely-presented groups which is still sufficient for our primary applications. This shorter proof allows us to keep control over the dimensions of approximate representations in the embedding, leading to the finer separation $C_{qa} \neq C_{qs}$. It is also natural to ask how large the question and answer sets need to be to separate C_{qs} from C_{qa} or C_{qc} . The group embedding theorems in this paper and [Slo17] give answer sets of size 8, but question sets with hundreds of questions. Dykema, Paulsen, and Prakash have made dramatic progress on this question, showing that $C_{qa} \neq C_{qs}$ for correlations with only five questions and two answers [DPP17]. Other related results can be found in [GHJ17, Har18]. Also, Coladangelo and Stark have shown that $C_{qs} \neq C_q$ [CS18], leaving the question of whether $C_{qa} = C_{qc}$ as the last unknown separation in Equation (1.1). Finally, Tobias Fritz has used the existence of a solution group with an undecidable word problem to show that quantum logic is undecidable [Fri16].

The rest of this paper is structured as follows. In the next section, we first recall the basic definitions of quantum correlation sets and non-local games, and then define binary linear system games and the solution group of a linear system. In Section 3, we state the embedding theorem for solution groups, and prove our two main results as corollaries. The rest of the paper is concerned with the proof of the embedding theorem. The main technical tool used is pictures of groups. We give an overview of pictures tailored to solution groups in Sections 6 and 7; expert readers will want to skip or briefly review these sections on first reading. In dealing with pictures, it is more convenient to use hypergraphs instead of linear systems, and we introduce hypergraphs into our definitions immediately in Section 2.2. In Section 8 we develop a notion of morphisms between hypergraphs; the concept is similar to graph minors, but differs from standard notions of hypergraph minors as in [RS10]. The more technical aspects of the proof are contained in Sections 9-11; we suggest that the reader skim these sections on first reading and proceed to Section 12. For a second reading, Corollary 10.4 might provide a good initial target.

1.1. Acknowledgements. I thank Richard Cleve, Jason Crann, Zhengfeng Ji, Li Liu, Andrew Marks, Vern Paulsen, and Vincent Russo for many helpful discussions. I thank Richard Cleve, Li Liu, and the anonymous referees for detailed comments on the manuscript, and Tobias Fritz and Doug Park for helpful suggestions. Any remaining errors are my own.

2. QUANTUM CORRELATION SETS AND LINEAR SYSTEM GAMES

2.1. Quantum correlation sets. In quantum mechanics, a physical system corresponds to a Hilbert space H, and the state of the system is recorded in a unit vector $v \in H$. A measurement in a system with finite outcome set $[m] := \{1, \ldots, m\}$ can be represented by a family $\{P_i\}_{i=1}^m$ of self-adjoint projections P_i such that $\sum_{i=1}^m P_i = \mathbb{1}$. Such a family is called a *projection-valued measure (PVM)*. The probability of measuring outcome *i* with PVM $\{P_i\}_{i=1}^m$ and state v is $v \cdot P_i v$, where $v \cdot w$ is the inner product in H.

As described in the introduction, there are two natural ways to represent a bipartite physical system (i.e. a system composed of two physically-separated subsystems) in quantum mechanics. In the tensor-product formalism, the subsystems correspond to Hilbert spaces H_A and H_B , and the Hilbert space of the joint system is $H_A \otimes H_B$. The state of the joint system is a unit vector $v \in H_A \otimes H_B$. If $\{P_a\}_{a=1}^{m_A}$, is a measurement on H_A and $\{Q_b\}_{b=1}^{m_B}$ is a measurement on H_B , then the joint measurement is represented by $\{P_a \otimes Q_b\}_{(a,b)\in[m_A]\times[m_B]}$, and the probability of measuring $(a,b) \in [m_A] \times [m_B]$ is $v \cdot P_a \otimes Q_b v$. Let m_A, m_B, n_A, n_B be positive integers. A function $p \in \mathbb{R}^{[m_A] \times [m_B] \times [n_A] \times [n_B]}$ is a tensor-product correlation if there are Hilbert spaces H_A and H_B , a unit vector $v \in H_A \otimes H_B$, a PVM $\{P_a^x\}_{a=1}^{m_A}$ on H_A for every $x \in [n_A]$, and a PVM $\{Q_b^y\}_{b=1}^{m_B}$ on H_B for every $y \in [n_B]$, such that

$$p(a,b|x,y) = v \cdot P_a^x \otimes Q_b^y v$$

for every $(a, b, x, y) \in [m_A] \times [m_B] \times [n_A] \times [n_B]$. In other words, p is a tensorproduct correlation if there is a bipartite system and state vector, along with a set of n_A (resp. n_B) measurements on the first (resp. second) subsystem, each with m_A (resp. m_B) outcomes, such that p(a, b|x, y) is the probability of measuring outcome (a, b) on measurements (x, y). A tensor-product correlation p is *finite-dimensional* if the Hilbert spaces H_A and H_B can be chosen to be finite-dimensional. The set of tensor-product correlations for a given tuple (m_A, m_B, n_A, n_B) is denoted by $C_{qs}(m_A, m_B, n_A, n_B)$, and the set of finite-dimensional correlations is denoted by $C_{qa}(m_A, m_B, n_A, n_B)$. The closure of $C_{qs}(m_A, m_B, n_A, n_B)$ is denoted by $C_{qa}(m_A, m_B, n_A, n_B)$.² By a result of Scholz-Werner, $C_{qa}(m_A, m_B, n_A, n_B)$ is also the closure of $C_q(m_A, m_B, n_A, n_B)$ [SW08].

The other natural way to represent a bipartite system is via the commutingoperator formalism. In this framework, each subsystem is represented by subalgebras \mathcal{A} and \mathcal{B} of bounded operators on the Hilbert space H of the joint system, such that ab = ba for all elements $a \in \mathcal{A}$ and $b \in \mathcal{B}$. If $\{P_a\}_{a=1}^{m_A}$ and $\{Q_b\}_{b=1}^{m_B}$ are PVMs in \mathcal{A} and \mathcal{B} respectively, then $\{P_aQ_b\}_{(a,b)\in[m_A]\times[m_B]}$

²The letters qa and qs can be read as "quantum-approximable" and "quantum-spatial". The latter term refers to the spatial tensor product of C^* -algebras (see below).

is also a PVM, and the probability of measuring outcome (a, b) on state $v \in H$ is $v \cdot P_a Q_b v$. Let m_A, m_B, n_A, n_B be positive integers. A function $p \in \mathbb{R}^{[m_A] \times [m_B] \times [n_A] \times [n_B]}$ is a commuting-operator correlation if there is a Hilbert space H, a unit vector $v \in H$, a PVM $\{P_a^x\}_{a=1}^{m_A}$ on H_A for every $x \in [n_A]$, and a PVM $\{Q_b^y\}_{b=1}^{m_B}$ on H_B for every $y \in [n_B]$, such that $P_a^x Q_b^y = Q_b^y P_a^x$ and

$$p(a,b|x,y) = v \cdot P_a^x Q_b^y v$$

for all $(a, b, x, y) \in [m_A] \times [m_B] \times [n_A] \times [n_B]$. The set of commuting operator correlations for a given tuple (m_A, m_B, n_A, n_B) is denoted by $C_{qc}(m_A, m_B, n_A, n_B)$. It is not hard to see that this set is closed.

We could also use positive-operator valued measures (POVMs) instead of PVMs to model measurements in the above definitions. However, any correlation achievable with POVMs can also be achieved with PVMs. For tensorproduct correlations this follows immediately from the Naimark dilation theorem, while for commuting-operator correlations this fact can be found in [Fri12, Section 3] or [PT15]. We stick with PVMs because of the clearer connection with operator algebras below.

A two-player non-local game with question sets $[n_A]$, $[n_B]$ and answer sets $[m_A], [m_B]$ is specified by a probability distribution π on $[n_A] \times [n_B]$ and a function $V : [m_A] \times [m_B] \times [n_A] \times [n_B] \rightarrow \{0, 1\}$. In the operational interpretation of the game, a referee chooses a pair of questions $(x, y) \in [n_A] \times [n_B]$ at random according to π , giving x to the first player (Alice), and y to the second player (Bob). The players respond with answers $a \in [m_A]$ and $b \in [m_B]$ respectively. The players win the game if V(a, b|x, y) = 1, and lose if V(a, b|x, y) = 0. The players' strategy in such a game can be described by the function $p \in \mathbb{R}^{[m_A] \times [m_B] \times [n_A]}$, where p(a, b|x, y) is the probability that the players output (a, b) on inputs (x, y). The winning probability on a strategy p is given by

$$\sum_{x=1}^{n_A} \sum_{y=1}^{n_B} \pi(x, y) \sum_{a=1}^{m_A} \sum_{b=1}^{m_B} V(a, b|x, y) p(a, b|x, y).$$

The game has a perfect tensor-product strategy (resp. perfect commutingoperator strategy) if there is a strategy in $C_{qs}(m_A, m_B, n_A, n_B)$ (resp. $C_{qc}(m_A, m_B, n_A, n_B)$) for which the winning probability is 1. We will show in Section 3 that there is a game with a perfect commuting-operator strategy, but no perfect tensor-product strategy, demonstrating that $C_{qs}(m_A, m_B, n_A, n_B) \neq C_{qc}(m_A, m_B, n_A, n_B)$ for some (m_A, m_B, n_A, n_B) .

For simplicity, many authors work only with the correlation sets $C_t(m, n) := C_t(m, m, n, n), t \in \{q, qs, qa, qc\}$. We work with games which typically have $m_A \neq m_B$ and $n_A \neq n_B$. However, by padding the game with irrelevant questions and answers, we can assume that $m_A = m_B$ and $n_A = n_B$. Indeed,

suppose that π, V is the data of game with question sets $[n_A], [n_B]$ and answer sets $[m_A], [m_B]$. Let $m = \max(m_A, m_B)$ and $n = \max(n_A, n_B)$, extend π to a probability distribution $\tilde{\pi}$ on $[n] \times [n]$ by setting $\tilde{\pi}(x, y) = 0$ if $x > n_A$ or $y > n_B$, and similarly extend V to a function $\tilde{V} : [m]^2 \times [n]^2 \to \{0, 1\}$ by setting $\tilde{V}(a, b|x, y) = 0$ if $(a, b, x, y) \notin [m_A] \times [m_B] \times [n_A] \times [n_B]$. Then it is easy to see that the game determined by $\tilde{\pi}, \tilde{V}$ has a perfect strategy in $C_t(m, n)$ if and only if the game determined by π, V has a perfect strategy in $C_t(m_A, m_B, n_A, n_B)$. Hence our argument will imply that $C_{qs}(m, n) \neq C_{qc}(m, n)$ for some m, n.

We also note that it is not necessary for the questions and answers in a non-local game to be integers: the question and answer sets $[n_A]$, $[n_B]$, $[m_A]$, $[m_B]$ can be replaced with any finite sets of the same size.

To finish the section, we note that the different correlation sets can also be defined using different tensor products of C^* -algebras. A PVM $\{P_a\}_{a=1}^m$ on a Hilbert space H can be thought of as the spectral projections of the unitary matrix $U = \sum_{a=1}^m e^{2\pi(a-1)/m}P_a$, and in this way PVMs with m outcomes correspond to representations of the C^* -group algebra $C^*\mathbb{Z}_m$. Similarly, collections of n PVMs, each with m outcomes, correspond to representations of $C^*\mathbb{Z}_m^{*n}$, where \mathbb{Z}_m^{*n} is the free product of n copies of \mathbb{Z}_m . Let p_a^x , $a \in [m_A]$, $x \in [n_A]$ be the ath spectral projection in the xth factor of \mathbb{Z}_m in $\mathcal{A} = C^*\mathbb{Z}_{mA}^{*n_A}$, and similarly let q_b^y , $b \in [m_B]$, $y \in [n_B]$ be the bth spectral projection in the yth factor of \mathbb{Z}_m in $\mathcal{B} = C^*\mathbb{Z}_{mB}^{*n_B}$. Then $C_{qc}(m_A, m_B, n_A, n_B)$ consists of the functions $p \in \mathbb{R}^{[m_A] \times [m_B] \times [n_A] \times [n_B]}$ of the form $p(a, b|x, y) = f(p_a^x q_b^y)$ for some state f on the maximal tensor product $\mathcal{A} \otimes_{max} \mathcal{B}$, and $C_{qa}(m_A, m_B, n_A, n_B)$ consists of the functions $p \in \mathbb{R}^{[m_A] \times [m_B] \times [n_A] \times [n_B]}$ of the form $p(a, b|x, y) = f(p_a^x q_b^y)$ for some state f on the minimal (also known as the spatial) tensor product $\mathcal{A} \otimes_{min} \mathcal{B}$ [SW08, Fri12, JNP+11].

A well-known result of Kirchberg states that Connes embedding problem is equivalent to what is known as Kirchberg's conjecture, that $C^* \mathcal{F}_n \otimes_{max} C^* \mathcal{F}_n = C^* \mathcal{F}_n \otimes_{min} C^* \mathcal{F}_n$ for all (or some) $n \geq 2$, where \mathcal{F}_n is the free group on ngenerators [Kir93]. In Kirchberg's conjecture, the group \mathcal{F}_n can be replaced with the group \mathbb{Z}_m^{*n} for $m, n \geq 2$, $(m, n) \neq (2, 2)$ (see for instance the surveys in [Fri12, Oza13]). Hence an affirmative answer to the the Connes embedding problem would imply that $C_{qc}(m, n) = C_{qa}(m, n)$ for all (m, n). Ozawa has shown that the converse is also true, so $C_{qc}(m, n) = C_{qa}(m, n)$ for all (m, n) if and only if Connes embedding problem is true.

The set $C_{qs}(m_A, m_B, n_A, n_B)$ consists of all functions $p \in \mathbb{R}^{[m_A] \times [m_B] \times [n_A] \times [n_B]}$ of the form $p(a, b|x, y) = v \cdot \phi_A(p_a^x) \otimes \phi_B(q_b^y)v$ for any pair of representations ϕ_A and ϕ_B of \mathcal{A} and \mathcal{B} on Hilbert spaces H_A and H_B respectively, and unit vector $v \in H_A \otimes H_B$. A state on a tensor-product $\mathcal{A} \otimes \mathcal{B}$ of the form $T \mapsto$ $v \cdot (\phi_A \otimes \phi_B)(T)v$ is called a spatial state; for the minimal tensor product, these states are dense in the set of all states. Hence separating $C_{qc}(m_A, m_B, n_A, n_B)$ from $C_{qs}(m_A, m_B, n_A, n_B)$ shows in particular that there are states on $\mathcal{A} \otimes_{max} \mathcal{B}$ which do not come from spatial states.

2.2. Linear system games, hypergraphs, and solution groups. Binary linear system games are based on linear systems Ax = b over \mathbb{Z}_2 . It is convenient to think of linear systems in terms of hypergraphs. By a hypergraph, we mean a triple $\mathcal{H} = (V, E, A)$, where $V = V(\mathcal{H})$ and $E = E(\mathcal{H})$ are finite sets of vertices and edges respectively, and $A \in \mathbb{Z}_{\geq 0}^{V \times E}$ is the incidence matrix between V and E, so $A_{ve} \geq 0$ is the degree of incidence between edge e and vertex v. We say that v and e are incident if $A_{ve} > 0$. If $v \in V$, then the degree of v is $|v| = \sum_e A_{ve}$. Similarly if $e \in E$ then $|e| = \sum_v A_{ve}$. We say that \mathcal{H} is simple if $A_{ve} \leq 1$ for all $v \in V$ and $e \in E$, k-regular if |v| = k for all $v \in V$, and a graph if |e| = 2 for all $e \in E$.

Note that this definition of hypergraphs allows both isolated vertices and isolated edges, i.e. vertices (resp. edges) which are incident to no edges (resp. vertices). A \mathbb{Z}_2 -vertex labelling of \mathcal{H} is a function $b: V \to \mathbb{Z}_2$. With these conventions, there is a correspondence between linear systems Ax = b and simple hypergraphs \mathcal{H} with a vertex labelling b. From this point of view, the edges of a hypergraph correspond to the variables of a linear system, and the vertices correspond to constraints. Similarly, pairs (\mathcal{H}, b) where \mathcal{H} is not necessarily simple correspond to linear systems Ax = b over \mathbb{Z}_2 with a choice of non-negative integer representatives for the coefficients A_{ve} .

To any $m \times n$ linear system Ax = b, we can associate a linear system nonlocal game \mathcal{G} , and a group Γ [CM14, CLS16]. In the game \mathcal{G} , Alice receives the index $1 \leq x \leq m$ of a row of A, and returns a function $a \in \mathbb{Z}_2^{V_x}$, where $V_x := \{1 \leq j \leq n : A_{ij} \neq 0\}$. Bob receives the index $1 \leq y \leq n$ of a column of A, and returns $b \in \mathbb{Z}_2$. The players win if $\sum_{j \in V_x} a(j) = b_x$, and either $y \notin V_x$, or $y \in V_x$ and a(y) = b. In other words, Alice outputs values for every variable in the *x*th equation, Bob outputs a value for the *y*th variable, and the players win if Alice's answer satisfies the *x*th equation and their answers are consistent with each other. The group Γ is the focus of attention of this paper, and is defined as follows.

Definition 2.1. Let $\mathcal{H} = (V, E, I)$ be a (not necessarily simple) hypergraph and let b be a function $V \to \mathbb{Z}_2 : v \mapsto b_v$. The solution group $\Gamma = \Gamma(\mathcal{H}; b)$ associated to \mathcal{H} and b is the group generated by $\{x_e, e \in E\} \cup \{J\}$, subject to relations:

- (1) $x_e^2 = 1$ for all $e \in E$ and $J^2 = 1$ (i.e. Γ is generated by involutions)
- (2) $[x_e, J] = 1$ for all $e \in E$ (i.e. J is central),
- (3) $[x_e, x_{e'}] = 1$ if there is some vertex v incident to both e and e', and

(4)

$$\prod_{e} x_{e}^{A_{ve}} = J^{b_{v}} \text{ for all } v \in V.$$

The null solution group is the group $\Gamma(\mathcal{H}) := \Gamma(\mathcal{H}, 0)$.

We call the last two types of relations commuting relations and linear relations respectively. The definition of the linear relations assumes that E is ordered, but the choice of order is irrelevant because of the commuting relations. Note that if v and e are incident and A_{ve} is even, then the linear relations

$$\prod_{e'} x_{e'}^{A_{ve'}} = J^{b_v} \text{ and } \prod_{e' \neq e} x_{e'}^{A_{ve'}} = J^{b_v}$$

are equivalent. However, the fact that $A_{ve} > 0$ might still lead to commuting relations that wouldn't hold otherwise.




FIGURE 1. The hypergraph from Example 2.2. Edges with degree two are drawn as lines, while edges with degree not equal to two are drawn as shaded regions.

Example 2.2. Consider the hypergraph \mathcal{H} with matrix

$$A = \begin{pmatrix} 1 & 0 & 0 & 0 & 2 \\ 1 & 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 1 & 0 \end{pmatrix},$$

represented visually in Figure 1, and let b = (0, 0, 1). Then $\Gamma(\mathcal{H}, b)$ is the group generated by x_1, \ldots, x_5, J , satisfying the relations

$$J^2 = 1, x_i^2 = [x_i, J] = 1$$
 for all $1 \le i \le 5$,

commuting relations

$$[x_i, x_j] = 1 \text{ for } (i, j) \in \{(1, 5), (1, 2), (1, 4), (2, 4), (1, 3), (3, 4)\},\$$

10

and linear relations

$$x_1 x_5^2 = x_1 x_2 x_4 = 1, x_1 x_3 x_4 = J.$$

In this example, the first linear relation implies $x_1 = 1$, so $\Gamma(\mathcal{H}, b)$ is isomorphic to $(\mathbb{Z}_2 * \mathbb{Z}_2) \times \mathbb{Z}_2$.

The one-dimensional representations π of $\Gamma(\mathcal{H}; b)$ in which $\pi(J) \neq 1$ correspond to the solutions of the linear system Ax = b. Higher-dimensional representations of $\Gamma(\mathcal{H}; b)$ with this property can be thought of as quantum solutions of Ax = b. This is justified by the following theorem, which relates solution groups to non-local games.

Theorem 2.3 ([CM14, CLS16]). Let Ax = b be a linear system over \mathbb{Z}_2 , where A is a non-negative integral matrix, let \mathcal{G} be the associated linear system nonlocal game, and let Γ be the corresponding solution group. Then:

- \mathcal{G} has a perfect quantum commuting-operator strategy if and only if $J \neq 1$ in Γ .
- \mathcal{G} has a perfect quantum tensor-product strategy if and only if \mathcal{G} has a perfect finite-dimensional quantum strategy, and this happens if and only if Γ has a finite-dimensional representation π with $\pi(J) \neq \mathbb{1}$.

The first part of this theorem is due to [CLS16], while the second part is due to [CM14].

3. The embedding theorem and consequences

In light of Theorem 2.3, we would like to understand the structure (or lack thereof) of solution groups for linear system games. Recall that a presentation $\langle S:R\rangle$ of a group G is a set S and subset R of the free group $\mathcal{F}(S)$ generated by S, such that $G = \mathcal{F}(S)/(R)$, where (R) is the normal subgroup generated by R. A group is finitely presented if it has a presentation $\langle S:R\rangle$ where both S and R are finite. Our primary result is that understanding solution groups is as hard as understanding finitely presented groups.

Theorem 3.1. Let G be a finitely presented group, let $J' \in G$ be a central element with $(J')^2 = 1$, and let w_1, \ldots, w_n , $n \ge 0$ be a sequence of elements in G such that $w_i^2 = 1$ for all $1 \le i \le n$. Then there is a hypergraph \mathcal{H} , a vertex labelling function $b: V(\mathcal{H}) \to \mathbb{Z}_2$, a sequence of edges e_1, \ldots, e_n in \mathcal{H} , and a homomorphism $\phi: G \to \Gamma(\mathcal{H}, b)$ such that ϕ is an embedding, $\phi(J') = J$, and $\phi(w_i) = x_{e_i}$ for all $1 \le i \le n$.

When J' = 1 and n = 0, Theorem 3.1 states that every finitely presented group G embeds in some solution group Γ . The full version of Theorem 3.1 says that if we also specify a list of involutions in G, then we can construct the

embedding so that these involutions map to generators of Γ , and in particular we can map a specified central involution to the element $J \in \Gamma$. This last condition is essential for our applications.

The hypergraph \mathcal{H} , vertex labelling b and homomorphism ϕ in Theorem 3.1 can be explicitly constructed from a presentation of G and a choice of representatives for w_1, \ldots, w_n and J'. The construction is recursive, in the sense that there is a Turing machine which, given a presentation for G and words representing w_1, \ldots, w_n , and J' as input, outputs \mathcal{H} , b, and ϕ as in the theorem. In particular, the construction does not depend on whether the elements w_1, \ldots, w_n, J are trivial or non-trivial in G. This construction is described in Sections 4 and 5. Theorem 3.1 is proved at the beginning of Section 5 via reduction to another embedding theorem. The proof of this latter embedding theorem (and hence the proof of Theorem 3.1) is completed in Section 12.

In the remainder of this section, we prove two consequences of Theorem 3.1. The first is an answer to Tsirelson's problem.

Corollary 3.2. There is a linear system non-local game which has a perfect quantum commuting-operator strategy, but does not have a perfect quantum tensor-product strategy.

For the proof, we recall the notion of an HNN extension. Let $\alpha : K_0 \to K_1$ be an isomorphism between two subgroups K_0 and K_1 of a group $H = \langle S : R \rangle$. The HNN extension of H by α is the group with presentation

$$G = \langle S \cup \{t\} : R \cup \{tkt^{-1} = \alpha(k) : k \in K_0\} \rangle,$$

where t is an indeterminate not in S. It is well-known that the natural map $H \hookrightarrow G$ is an injection, so H is a subgroup of G (see, e.g., [LS77, Section IV.2]).

Proof of Corollary 3.2. Suppose that G is a finitely presented group with a central element J' of order two, such that $\pi(J') = 1$ for every finite-dimensional representation π of G. By Theorem 3.1, there is an embedding of G in a solution group Γ which identifies J' with J. In particular, this implies that $J \neq 1$ in Γ , so the associated linear system non-local game G must have a perfect quantum commuting-operator strategy by the first part of Theorem 2.3. If π is a finite-dimensional representation of Γ , then $\pi(J) = \pi|_G(J') = 1$. By the second part of Theorem 2.3, the associated linear system non-local game does not have a perfect quantum tensor-product strategy.

To finish the proof, we construct a group G with the above property. Consider Higman's group

$$H_0 = \langle a, b, c, d : aba^{-1} = b^2, bcb^{-1} = c^2, cdc^{-1} = d^2, dad^{-1} = a^2 \rangle.$$

It is well-known that H_0 has no non-trivial linear representations, and that the generators a, b, c, d of H_0 have infinite order [Hig51, Ber94]. Let $H = H_0 \times \mathbb{Z}_2$, and let $J \in H$ denote the generator of the \mathbb{Z}_2 -factor. Let G be the HNN extension of H by the automorphism of $\langle a, J \rangle \cong \mathbb{Z} \times \mathbb{Z}_2$ sending $J \mapsto J$ and $a \mapsto aJ$. By the properties of the HNN extension, H is a subgroup of G, and in particular J is non-trivial in G. Furthermore, we can construct a presentation for G from a presentation of H by adding a generator x and relations [x, J] = 1and [x, a] = J. The former relation implies that J is central in G. Finally, if π is a finite-dimensional representation of G, then $\pi|_{H_0}$ is trivial, and in particular, $\pi(a) = 1$. But this implies that

$$\pi(J) = \pi([x, a]) = \mathbb{1},$$

as required.

We note that any non-residually-finite group can be used in place of Higman's group in the above proof.

It would be interesting to know the smallest linear system for which the corresponding game can be played perfectly only with commuting-operator strategies. No effort is made to reduce the size of the linear system in the proof of Theorem 3.1, and the main construction from Section 5 depends on the number of variables and the total length of the relations in the presentation of G. If we naively follow the proof through for the group in Corollary 3.2, we get a linear system with roughly 600 variables and 450 linear relations. Making some obvious improvements in Section 4 can get this down to 400 variables and 300 relations.

The second consequence concerns the difficulty of determining whether a non-local game has a perfect commuting-operator strategy.

Corollary 3.3. It is undecidable to determine if a binary linear system game has a perfect commuting-operator strategy.

Proof. By Theorem 2.3, determining if a binary linear system game has a perfect strategy is equivalent to determining if $J \neq 1$ in a solution group. Because Theorem 3.1 is constructive, this is in turn equivalent to the following decision problem: given a finite group presentation $G = \langle S : R \rangle$ and a word J in the generators S such that $J \in Z(G)$ and $J^2 = 1$, decide if J = 1 in G.

We claim that the word problem for groups can be reduced to this latter problem. Indeed, given a finitely presented group $K = \langle S : R \rangle$ and a word $w \in \mathcal{F}(S)$ in the generators of K, it is possible to recursively construct a finitely presented group L_w with the property that K is a subgroup of L_w if $w \neq 1$, and L_w is trivial if w = 1 (see [LS77, pg. 190], where this construction is attributed to Rabin). The presentation of L_w can be constructed by adding finitely many generators and relations to the presentation of K. Let H_w be the result of

applying this construction to the group $K \times \mathbb{Z}$, and let z be the generator of H_w corresponding to the generator of the \mathbb{Z} -factor in $K \times \mathbb{Z}$. If w = 1, then H_w is trival, and z = 1. If $w \neq 1$, then $K \times \mathbb{Z}$ is a subgroup of H_w , and the order of z is infinite. Finally, construct a group G_w by adding two generators x and J to the presentation of H_w , along with relations $J^2 = [x, J] = 1$, [s, J] = 1 for all generators s of H_w , and [x, z] = J. As in Corollary 3.2, if $w \neq 1$ then G_w is the HNN extension of the group $H_w \times \mathbb{Z}_2$ by the automorphism of the subgroup $\langle z, J \rangle \cong \mathbb{Z} \times \mathbb{Z}_2$ sending $J \mapsto J$ and $z \mapsto zJ$. Thus, if $w \neq 1$ then $J \neq 1$ in G_w . If w = 1, then z = 1 in H_w , and consequently J = [x, z] = 1 in G_w . This completes the reduction.

Although not used in either of the above corollaries, Theorem 3.1 also allows us to embed a finitely presented group G in a solution group Γ so that a given set of involutions of G become generators of Γ . This can be used to prove that other tasks involving solution groups are undecidable. For instance, it is undecidable to determine if a generator x_e of a null solution group is nontrivial.

4. Presentations by involutions

We are interested primarily in groups which (a) have a distinguished central element of order \leq two, and (b) are generated by involutions. For clarity in subsequent sections, we encode these conditions in two formal definitions.

Definition 4.1. A group over \mathbb{Z}_2 is a group G with a distinguished central element $J = J_G$ such that $J^2 = 1$.

A morphism $G_1 \to G_2$ over \mathbb{Z}_2 is a group homomorphism sending $J_{G_1} \mapsto J_{G_2}$. Similarly, an embedding over \mathbb{Z}_2 is an injective morphism over \mathbb{Z}_2 .

Note that J is allowed to be the identity in this definition. This is so that we can construct groups over \mathbb{Z}_2 by starting with some finite presentation, picking an element $J' \in \mathcal{F}(S)$, and adding relations $(J')^2 = 1$ and [J', s] = 1for all $s \in S$. By allowing J = 1, we can do this even if J' becomes trivial.

Elements of $\mathcal{F}(S)$ are represented by words over $\{s, s^{-1} : s \in S\}$. Every element $r \in \mathcal{F}(S)$ can be represented uniquely as $s_1^{a_1} \cdots s_n^{a_n}$, where $a_i \in \{\pm 1\}$, and $a_i = a_{i+1}$ whenever $s_i = s_{i+1}$. A word meeting these conditions is said to be *reduced*. The number *n* is the *length* of *r*. The element *r* is said to be *cyclically reduced* if, in addition, $s_n = s_1$ implies that $a_n = a_1$.

Definition 4.2. Given a set S, let $\mathcal{F}_2(S) = \langle S : s^2 = 1, s \in S \rangle$. A presentation by involutions over \mathbb{Z}_2 for a group G is a set of generators S and a set of relations $R \subset \mathcal{F}_2(S) \times \mathbb{Z}_2$ such that $G = \mathcal{F}_2(S) \times \mathbb{Z}_2/(R)$, where (R) is the normal subgroup generated by R. We denote presentations of this form by $\operatorname{Inv}\langle S : R \rangle$, and write $G = \operatorname{Inv}\langle S : R \rangle$ when the meaning is clear.

We use J (written in multiplicative notation) to denote the generator of the \mathbb{Z}_2 -factor in $\mathcal{F}_2(S) \times \mathbb{Z}_2$. If $G = \text{Inv}\langle S : R \rangle$, we can regard G as a group over \mathbb{Z}_2 by letting $J = J_G$ be the image of $J \in \mathcal{F}_2(S) \times \mathbb{Z}_2$ in G.

Elements of $\mathcal{F}_2(S)$ are represented by words over S. Every element $r \in \mathcal{F}_2(S) \times \mathbb{Z}_2$ can be represented uniquely as $J^a s_1 \cdots s_n$, where $a \in \mathbb{Z}_2$ and s_1, \ldots, s_n is a sequence in S with $s_i \neq s_{i+1}$. Again, a word of this form is said to be reduced, and n is called the length of r. If, in addition, $s_n \neq s_1$ then we say that r is cyclically reduced. We say that a set of relations R is cyclically reduced if every element of R is cyclically reduced.

If R is a set of relations, the symmetrization of R is the set of relations R^{sym} containing all relations of the form

 $J^{a}s_{i}s_{i+1}\cdots s_{n}s_{1}\cdots s_{i-1}$ and $J^{a}s_{i}s_{i-1}\cdots s_{1}s_{n}\cdots s_{i+1}, 1 \leq i \leq n$,

for every relation $J^a s_1 \cdots s_n$ in R.

Every group presented by involutions over \mathbb{Z}_2 has a presentation $\operatorname{Inv}\langle S:R \rangle$ where R is cyclically reduced. The presentations $\operatorname{Inv}\langle S:R \rangle$ and $\operatorname{Inv}\langle S:R \rangle$ $R^{sym}\rangle$ are equivalent (i.e. they define isomorphic groups), and if R is cyclically reduced then R^{sym} is cyclically reduced.

By definition, solution groups are examples of groups presented by involutions over \mathbb{Z}_2 . Theorem 3.1 states that every finitely presented group over \mathbb{Z}_2 embeds (over \mathbb{Z}_2) in a solution group. The first step in proving Theorem 3.1 is showing that every finitely presented group embeds in a group presented by involutions.

Proposition 4.3. Suppose (G, J) is a group over \mathbb{Z}_2 with finite presentation $\langle S: R \rangle$, and $J' \in \mathcal{F}(S)$ is a representative of J_G . Let T be the set of indeterminates $\{z_{s1}, z_{s2}: s \in S\}$, and choose integers $k_s \geq 1$ for all $s \in S$. Finally, let $\phi: \mathcal{F}(S) \to \mathcal{F}_2(T) \times \mathbb{Z}_2$ be the morphism sending $s \mapsto (z_{s1}z_{s2})^{k_s}$. Then the induced morphism

 $\phi: G \to K := \operatorname{Inv}\langle T: R' \rangle$, where $R' := \{\phi(r): r \in R\} \cup \{J_K \phi(J')\}$

is an embedding over \mathbb{Z}_2 .

Furthermore, if $R \cup \{J'\}$ is cyclically reduced then R' is cyclically reduced.

Proof. Let s_1, \ldots, s_n be a list of the elements of S, and let m_i be the order of s_i in G. For convenience, we write z_{ij} in place of $z_{s_i,j}$. For each $0 \le r \le n$, let $K_r := \langle S \cup \{z_{11}, z_{12}, \ldots, z_{r1}, z_{r2}, J\} : R \cup \{z_{ij}^2 = [z_{ij}, J] = 1 : 1 \le i \le r, j = 1, 2\} \cup \{s_j = (z_{i1}z_{i2})^{k_{s_j}}\} \cup \{J = J'\}\rangle.$

The presentation K_n is equivalent to the presentation $\text{Inv}\langle T : R' \rangle$ of the group K, so we just need to show that the natural map $G \to K_n$ is an inclusion. But this follows from the fact that each K_i , $1 \leq i \leq n$, is an amalgamated product

of K_{i-1} with either a dihedral group, or the product of a dihedral group and \mathbb{Z}_2 . Indeed, suppose that the natural map of $G \to K_{i-1}$ is an inclusion, and let

$$D_i := \langle z_{i1}, z_{i2} : z_{i1}^2 = z_{i2}^2 = (z_{i1}z_{i2})^{k_{s_i}m_i} = 1 \rangle,$$

the dihedral group of order $2k_{s_i}m_i$ (if m_i is infinite, then the last relation is omitted, so that D_i is the infinite dihedral group). If $J_G \notin \langle s_i \rangle$, then $\langle s_i, J_G \rangle \cong \mathbb{Z}_{m_i} \times \mathbb{Z}_2$, and K_i is the amalgamated product of K_{i-1} with $D_i \times \mathbb{Z}_2$ over $\langle s_i, J_G \rangle$, where we identify $s_i \in G \subseteq K_{i-1}$ with $(z_{i1}z_{i2})^{k_{s_i}}$, and J_G with the generator of \mathbb{Z}_2 in $D_i \times \mathbb{Z}_2$. If, on the other hand, $J_G \in \langle s_i \rangle$, then $J_G = s_i^a$, where a = 0 or $m_i/2$. In both cases, $(z_{i1}z_{i2})^{k_{s_i}a}$ is central in D_i , so K_i is the amalgamated product of K_{i-1} with D_i over $\langle s_i \rangle$, where we again identify s_i with $(z_{i1}z_{i2})^{k_{s_i}}$ (it's necessary to check that z_{i1} and z_{i2} commute with s_i^a because of the relations $[z_{ij}, J]$ in K_i). It follows that the natural map $G \to K_i$ is an inclusion. Since $G \cong K_0$, we ultimately conclude that the natural map $G \to K_n$ is an inclusion as desired.

Finally, it is easy to see that if $r \in R$ is cyclically reduced, then $\phi(r)$ is cyclically reduced. If J' is cyclically reduced, then $J\phi(J')$ is also cyclically reduced.

Definition 4.4. A relation $r = J^a s_1 \cdots s_n \in \mathcal{F}_2(S) \times \mathbb{Z}_2$ is odd (resp. even) if a is odd (resp. even). Equivalently, a relation is odd (resp. even) if it is of the form r' = J (resp. r' = 1) for some $r' \in \mathcal{F}_2(S)$.

The even part of the relation $r = J^a s_1 \cdots s_n$ is $r^+ = s_1 \cdots s_n$. If $\text{Inv}\langle S : R \rangle$ is a presentation by involutions over \mathbb{Z}_2 , then the corresponding even presentation over \mathbb{Z}_2 is $\text{Inv}\langle S : R^+ \rangle$, where $R^+ = \{r^+ : r \in R\}$.

Similarly, if G is any group over \mathbb{Z}_2 , then the even quotient is $G^+ := G/(J_G) \times \mathbb{Z}_2$. The group G^+ is regarded as a group over \mathbb{Z}_2 with J_{G^+} equal to the generator of the \mathbb{Z}_2 factor.

It is easy to see that if $G = \text{Inv}\langle S : R \rangle$, then $G^+ = \text{Inv}\langle S : R^+ \rangle$. For instance, the null solution group $\Gamma(\mathcal{H}) = \Gamma(\mathcal{H}, 0)$ of a hypergraph \mathcal{H} is the even quotient $\Gamma(\mathcal{H}, b)^+$ of the solution group $\Gamma(\mathcal{H}, b)$ for any b.

Definition 4.5. Let $J^a s_1 \cdots s_n$ be a reduced word for an element $r \in \mathcal{F}_2(S) \times \mathbb{Z}_2$. The multiplicity of $s \in S$ in r is

$$mult(s; r) := |\{1 \le i \le n : s_i = s\}|.$$

We say that $s \neq t \in S$ are adjacent in *r* if either $\{s, t\} = \{s_i, s_{i+1}\}$ for some i = 1, ..., n-1 or $\{s, t\} = \{s_1, s_n\}$.

The reason that we introduce numbers k_s in Proposition 4.3 is that, for the proof of Theorem 3.1, we would like to work with relations r where mult(s; r) is even for all s. In fact, we will be able to handle slightly more general presentations, which we now define.

Definition 4.6. We say that a presentation Inv(S : R) by involutions over \mathbb{Z}_2 is collegial if

- (a) the presentation is finite and cyclically reduced,
- (b) $R \cap \{1, J\} = R \cap S = \emptyset$, and
- (c) if $\operatorname{mult}(s; r_0)$ is odd for some $r_0 \in R$, and t is adjacent to s in some $r_1 \in R$, then $\operatorname{mult}(t; r')$ is even for all $r' \in R$.

Remark 4.7. Note that if $\text{Inv}\langle S : R \rangle$ is collegial, then every relation $r \in R$ must have length at least four, i.e. $r = J^a s_1 \cdots s_n$ where $n \ge 4$. This is because relations of length zero and one are explicitly excluded by condition (b), relations of the form s^2 , sts, s^2t , or ts^2 are not cyclically reduced, and relations st and str, where s, t, r are distinct, do not satisfy condition (c).

Corollary 4.8. Let G be a finitely presented group over \mathbb{Z}_2 , with a sequence of elements $w_1, \ldots, w_n \in G$ such that $w_i^2 = 1$ for all $1 \leq i \leq n$. Then there is collegial presentation $\text{Inv}\langle S: R \rangle$, and an embedding $\phi: G \to K := \text{Inv}\langle S: R \rangle$ over \mathbb{Z}_2 such that $\phi(w_i) \in S \subset K$ for all $1 \leq i \leq n$.

Proof. We can find a cyclically reduced presentation $\langle S_0 : R_0 \rangle$ for G in which J_G is a generator, $1 \notin R_0$, and each w_i has a representative $w'_i \in \mathcal{F}(S_0) \setminus \{1\}$ (it is always possible to find such a presentation, since if necessary we can add an extra generator z, along with the relation z = 1, and use this as a representative of the identity). In particular, this gives us a presentation where J_G is represented by a cyclically reduced non-identity element of $\mathcal{F}(S_0)$, namely itself.

Applying Proposition 4.3 to this presentation with $k_s = 2$ (or any other even number) for all $s \in S_0$ gives us an embedding ϕ of G in a finite presentation $\operatorname{Inv}\langle T: R' \rangle$, where R' is cyclically reduced. Since all k_s 's are even, $\operatorname{mult}(t; \phi(r))$ is even for every $r \in \mathcal{F}(S_0)$ and $t \in T$, and every relation in R' has length ≥ 4 . We conclude that $\operatorname{Inv}\langle T: R' \rangle$ is collegial.

Now let

$$S = T \cup \{\overline{w}_1, \dots, \overline{w}_n\},\$$

where $\overline{w}_1, \ldots, \overline{w}_n$ are new indeterminates, and set

$$R = R' \cup \{\overline{w}_i \phi(w'_i) : 1 \le i \le n\},\$$

where $\phi : \mathcal{F}(S_0) \to \mathcal{F}_2(T) \times \mathbb{Z}_2$ as in Proposition 4.3. Since \overline{w}_i does not appear in $\phi(w'_i)$, the relation $r = \overline{w}_i \phi(w'_i)$ is cyclically reduced. Furthermore, none of the \overline{w}_i 's are adjacent, and $\operatorname{mult}(s; r)$ is even for all $s \in T$ and $r \in R$, so $\operatorname{Inv}\langle S : R \rangle$ is collegial. But $\operatorname{Inv}\langle S : R \rangle$ is plainly equivalent to $\operatorname{Inv}\langle T : R' \rangle$, so the corollary follows.

Note that the proof of Corollary 4.8 is constructive, in the sense that there is a Turing machine which outputs S, R, and ϕ (as specified by a set of representatives in $\mathcal{F}_2(S)$ for the images through ϕ of the generators of G) given a finite presentation for G and a set of representatives for w_1, \ldots, w_n .

5. The wagon wheel embedding

Using Corollary 4.8, the proof of Theorem 3.1 reduces to the following:

Theorem 5.1. Let G be a group with a collegial presentation $\mathcal{I} = \text{Inv}\langle S : R \rangle$. Then there is a hypergraph $\mathcal{W} := \mathcal{W}(\mathcal{I})$ and vertex labelling $b := b(\mathcal{I})$ such that $S \subset E(\mathcal{W})$, and the resulting map

$$\mathcal{F}(S) \times \mathbb{Z}_2 \to \Gamma(\mathcal{W}, b) : s \mapsto x_s$$

descends to an embedding $G \hookrightarrow \Gamma(\mathcal{W}, b)$ over \mathbb{Z}_2 .

Proof of Theorem 3.1 using Theorem 5.1. Let G be a group over \mathbb{Z}_2 with elements w_1, \ldots, w_n such that $w_i^2 = 1$ for $i = 1, \ldots, n$. By Corollary 4.8, there is a collegial presentation $\mathcal{I} := \text{Inv}\langle S : R \rangle$ and an embedding $\phi_1 : G \to K := \text{Inv}\langle S : R \rangle$ over \mathbb{Z}_2 with $\phi_1(w_i) \in S$ for all $i = 1, \ldots, n$.

By Theorem 5.1, there is an embedding $\phi_2 : K \to \Gamma(\mathcal{W}(\mathcal{I}), b(\mathcal{I}))$ over \mathbb{Z}_2 with $\phi_2(s) = x_s$ for all $s \in S$. The composition $\phi_2 \circ \phi_1$ satisfies the conditions of Theorem 3.1.

Although we are still very far from being able to prove Theorem 5.1, in this section we shall describe the hypergraph $\mathcal{W}(\mathcal{I})$, which we call the *wagon wheel* hypergraph of \mathcal{I} . The proof of Theorem 5.1 will be given in Section 12.

The wagon wheel hypergraph can be defined for any (not necessarily collegial) presentation $\text{Inv}\langle S, R \rangle$. Let $R = \{r_1, \ldots, r_m\}$, let n_i be the length of r_i , and write $r_i = J^{p_i} s_{i1} \cdots s_{in_i}$, where $s_{ij} \in S$. The wagon wheel hypergraph is a simple hypergraph \mathcal{W} with vertex set

$$V := \{ (i, j, k) : 1 \le i \le m, j \in \mathbb{Z}_{n_i}, 1 \le k \le 3 \},\$$

and edge set

$$E := S \sqcup \{a_{ij}, b_{ij}, c_{ij}, d_{ij} : 1 \le i \le m, j \in \mathbb{Z}_{n_i}\}.$$

As a result, if $M := \sum_{i=1}^{k} n_i$, then \mathcal{W} has 3M vertices and 4M + |S| edges. \mathcal{W} has the following incidence relations for every $1 \le i \le m$ and $1 \le j \le n_i$:

- $s \in S$ is incident with (i, j, 1) if and only if $s_{ij} = s$,
- a_{ij} is incident with (i, j 1, 2) and (i, j, 1),
- b_{ij} is incident with (i, j, 1) and (i, j, 2),
- c_{ij} is incident with (i, j, 2) and (i, j, 3), and




FIGURE 2. The portion of the wagon wheel hypergraph containing vertices V_i and all incident edges. To save space, (i, j, k)is written as j, k, and s_{ij}, a_{ij}, \ldots are written as s_j, a_j, \ldots

• d_{ij} is incident with (i, j - 1, 3) and (i, j, 3).

Note that the only edges incident with vertices

$$V_i := \{ (i, j, k) : j \in \mathbb{Z}_{n_i}, 1 \le k \le 3 \}$$

are the edges in

$$E_{i} := \{a_{ij}, b_{ij}, c_{ij}, d_{ij} : j \in \mathbb{Z}_{n_{i}}\}$$

and the edges s_{i1}, \ldots, s_{in_i} . Furthermore, all the edges in E_i are incident with exactly two vertices, both belonging to V_i . The portion of the hypergraph \mathcal{W} incident with V_i is shown in Figure 2. An example of a wagon wheel hypergraph for a small (non-collegial) presentation is shown in Figure 3.

We also need to define the vertex labelling in Theorem 5.1.

Definition 5.2. An \mathcal{I} -labelling of \mathcal{W} is a vertex labelling $b : V \to \mathbb{Z}_2$ such that $|b^{-1}(1) \cap V_i| = p_i \mod 2$ for all $1 \le i \le m$.

For Theorem 5.1, we can choose any \mathcal{I} -labelling. This is because all \mathcal{I} -labellings are equivalent in the following sense:

Lemma 5.3. Let b and b' be two \mathcal{I} -labellings of \mathcal{W} . Then there is an isomorphism $\Gamma(\mathcal{W}, b) \to \Gamma(\mathcal{W}, b')$ which sends $x_s \mapsto x_s$ for all $s \in S$.




FIGURE 3. An example of the wagon wheel hypergraph $\mathcal{W}(\mathcal{I})$ when $\mathcal{I} = \text{Inv}\langle x, y, z, u, v : xyxz = xuvu = 1 \rangle$.

Proof. Suppose \mathcal{H} is a hypergraph with incidence matrix $A(\mathcal{H})$ and vertex labelling $b^{(0)}$. Given $e \in E(\mathcal{H})$, let $b^{(1)}$ be the vertex labelling with $b_v^{(1)} = b_v^{(0)} + A(\mathcal{H})_{ve}$ (i.e. we toggle the sign of all vertices incident with e according to multiplicity). Then there is an isomorphism

$$\Gamma(\mathcal{H}, b^{(0)}) \to \Gamma(\mathcal{H}, b^{(1)}) : x_f \mapsto \begin{cases} x_f & f \neq e \\ Jx_e & f = e \end{cases}$$

For \mathcal{W} , since $|b^{-1}(1) \cap V_i|$ and $|(b')^{-1}(1) \cap V_i|$ have the same parity, it is easy to see that $b|_{V_i}$ can be transformed to $b'|_{V_i}$ by toggling signs of vertices incident to edges $e \in E_i$ as necessary. The lemma follows.

Finally, note that it is not hard to write down a Turing machine to construct the wagon wheel hypergraph $\mathcal{W}(\mathcal{I})$ and choose an \mathcal{I} -labelling *b* for a given input presentation \mathcal{I} . Since Corollary 4.8 is also constructive, the proof of Theorem 3.1 is also constructive as claimed.

6. Pictures for groups generated by involutions

In this section we give an overview of the main technical tool used in the proof of Theorem 5.1: pictures of groups. These pictures, which are dual to the somewhat better known van Kampen diagrams, are a standard tool in combinatorial group theory. The purpose of pictures is to encode derivations of group identities from a set of starting relations; see [Sho07] for additional background. Here we introduce a variant adapted to groups generated by involutions.

6.1. Pictures as planar graphs. By a curve, we shall mean the image of an piecewise (regular) real-analytic function³ γ from a closed interval [a, b] (where a < b) to either the plane or the sphere. A curve γ is simple if $\gamma(s) \neq \gamma(t)$ for all $a \leq s < t \leq b$, except possibly when s = a and t = b. The points $\gamma(a)$ and $\gamma(b)$ are called the endpoints of the curve. A curve has either one or two endpoints; if $\gamma(a) = \gamma(b)$ then the curve is said to be *closed*. A connected region (in the plane or on the sphere) is *simple* if its boundary is a simple closed curve.

Definition 6.1. A picture is a collection (V, E, D), where

- (a) \mathcal{D} is a closed simple region,
- (b) V is a finite collection of points, called vertices, in \mathcal{D} ,
- (c) E is a finite collection of simple curves, called edges, in \mathcal{D} , and
- (d) for all edges $e \in E$ and points p of e,
 - (i) if e is not closed and p is an endpoint of e, then either $p \in V$, or p belongs to the boundary of \mathcal{D} and is not the endpoint of any other edge;
 - (ii) if e is closed and p is an endpoint of e, then p does not belong to the boundary of D;
 - (iii) if p is not an endpoint of e, then $p \notin V$, and p does not belong to any other edge or the boundary of \mathcal{D} .

If an edge e contains a vertex v, then we say that e and v are incident. If e contains a point of the boundary of \mathcal{D} , then we say that e is incident with the boundary. A picture is closed if no edges are incident with the boundary of \mathcal{D} . The size of a G-picture \mathcal{P} is the number of vertices in \mathcal{P} .

According to this definition, a picture is a type of planar embedding of a graph, albeit a graph where we can have multiple edges between vertices, loops at a vertex, and even closed loops which are not incident to any vertex. From this point of view, the boundary of \mathcal{D} can be regarded as a special type of vertex; if we think of the picture as drawn on a sphere, then this vertex would naturally be drawn at infinity. An illustration of these two equivalent points of view is shown in Figure 4. However, it is more convenient not to include the boundary of \mathcal{D} in the vertex set of a picture, and we stick with the convention

³Using real-analytic curves has some advantages in streamlining the following definitions; for instance, we can use the fact that two real-analytic curves intersect in at most finitely many points. However, the arguments in the subsequent sections also work with other formal definitions of pictures, such as the definition in [Sho07] using smooth curves and "fat" vertices.




FIGURE 4. A picture embedded in a disk (left) and on the plane with the exterior of the disk shrunk down to a special vertex at infinity (right).




FIGURE 5. A closed picture embedded in the sphere seen (up to isotopy) with two different choices for the location of the point at infinity. Faces are distinguished by different colours.

of treating the boundary separately. In particular, the picture in Figure 4 has size 7.

There is one important exception where we want to forget the boundary of \mathcal{D} , and that is when the picture is closed. If this happens, we often want to think of the picture as embedded in the sphere, without the point at infinity being marked. An example of a closed picture on the sphere, as seen from two different positions, is given in Figure 5. To handle this case, we allow \mathcal{D} to be the whole sphere, in which case the boundary is empty. Also note that we consider two pictures equal if they differ up to isotopy, either in the plane or on the sphere as appropriate. Such isotopies are allowed to move the boundary of the simple region, as well as the location of endpoints of edges on the boundary, as long as endpoints are not identified.

If \mathcal{P} is a picture in \mathcal{D}_0 and \mathcal{D} is a closed simple subregion of \mathcal{D}_0 , then the portion of \mathcal{P} contained in \mathcal{D} can be interpreted as a picture inside \mathcal{D} . We do, however, have to make sure that boundary edges of the picture inside \mathcal{D} do not have a common endpoint. This leads to two natural notions of the restriction of \mathcal{P} to \mathcal{D} .

Definition 6.2. Let \mathcal{P} be a picture in \mathcal{D}_0 , and let \mathcal{D} be a closed simple region in \mathcal{D}_0 with interior \mathcal{D}° . Given $\epsilon > 0$, let \mathcal{D}^{ϵ} denote the ϵ -relaxation of \mathcal{D} (the set of points which are at most distance ϵ from \mathcal{D}), and let $\mathcal{D}^{-\epsilon}$ denote the ϵ -contraction (the set of points inside \mathcal{D} which are at least distance ϵ from the boundary of \mathcal{D}). The boundary of $\mathcal{D}^{\pm \epsilon}$ will be a simple closed curve for small $\epsilon > 0.^4$

Recall that a curve intersects the boundary of \mathcal{D} transversally if, in every small disk around the intersection point, there are points of the curve which lie both on the interior and the exterior of \mathcal{D} .⁵ We say that \mathcal{D} is transverse to \mathcal{P} if every edge which intersects the boundary of \mathcal{D} does so transversally, and the boundary of \mathcal{D} does not contain any vertices of \mathcal{P} .

The restriction of \mathcal{P} to a transverse region \mathcal{D} is the picture $\operatorname{res}(\mathcal{P}, \mathcal{D})$ with vertex set $V(\mathcal{P}) \cap \mathcal{D}$, and whose edges are the closures of the connected components of $e \cap \mathcal{D}^o$, for $e \in E(\mathcal{P})$. In other words, edges are cut off at the boundary, and edges which intersect the boundary at multiple points may be cut into multiple edges.⁶

For a general region \mathcal{D} , the contraction $\mathcal{D}^{-\epsilon}$ will be transverse to \mathcal{P} for small enough $\epsilon > 0$. The restrictions $\operatorname{res}(\mathcal{P}, \mathcal{D}^{-\epsilon})$ are thus well-defined, and can be identified via isotopy with pictures in \mathcal{D} . These pictures belong to a single isotopy class $\operatorname{res}(\mathcal{P}, \mathcal{D})$, which we identify as the restriction of \mathcal{P} to \mathcal{D} .⁷

⁵Transversality is typically defined in terms of tangent lines to the curves, but this more permissive definition is fine for our purposes.

⁴In particular, the boundary curves will be piecewise real-analytic. When the boundary curve γ of \mathcal{D} is real-analytic, this follows from the tubular neighbourhood theorem: for small enough ϵ , the boundary curves of $D^{\pm \epsilon}$ will be normal translates of γ . If γ is piecewise real-analytic, then the situation becomes more complicated at the corners of γ . When two real-analytic segments of γ meet at an acute angle, the boundary of $D^{\pm \epsilon}$ will consist of the normal translates of the two segments, cut off at their intersection point. For two segments meeting at an obtuse angle, the normal translates of each segment will be joined by a portion of the circle of radius ϵ centered at the corner.

⁶Since boundaries and edges are both real-anlytic, there will be only finitely many edges in this picture, as required by our definition.

⁷This definition also uses the fact that edges and boundaries are real-analytic. For smooth curves, more care must be taken: the contraction may fail to be transverse to \mathcal{P} for arbitrarily small ϵ , and even if we only look at curves transverse to \mathcal{P} in \mathcal{D}^o , there can be curves which are arbitrarily close to the boundary with arbitrarily many intersections with \mathcal{P} (so the isotopy class may not be unique).




FIGURE 6. Pictures can be restricted to a region homotopic to a disk and bounded by a simple closed curve which is transverse to the picture.




FIGURE 7. The germ of the region enclosed by a simple cycle is computed by first taking an ϵ -relaxation of the region. In this case the cycle is facial, so the restriction would be empty.

Similarly, the germ of \mathcal{D} in \mathcal{P} is the isotopy class germ $(\mathcal{P}, \mathcal{D})$ of res $(\mathcal{P}, \mathcal{D}^{\epsilon})$ for small $\epsilon > 0$.

Unless otherwise noted, we assume that subregions are closed. If \mathcal{D} is transverse to \mathcal{P} , then res $(\mathcal{P}, \mathcal{D})$ and germ $(\mathcal{P}, \mathcal{D})$ agree. An example of this type of restriction is shown in Figure 6. In general, vertices in the boundary of \mathcal{D} will not appear in res $(\mathcal{P}, \mathcal{D})$, but are preserved, along with all their outgoing edges, in germ $(\mathcal{P}, \mathcal{D})$. One way to get a simple region is to take a region \mathcal{D} enclosed by a simple cycle in \mathcal{P} . An example of this type, in which the germ is different than the restriction, is given in Figure 7.

Definition 6.3. Let \mathcal{P} be a picture in \mathcal{D}_0 . A simple cycle in \mathcal{P} is a collection of edges whose union is a simple closed curve.

A closed loop is an edge of \mathcal{P} which is not incident to any vertex or to the boundary, and thus forms a simple cycle by itself.

A face of \mathcal{P} is an open connected region \mathcal{D} of \mathcal{D}_0 which does not contain any points of \mathcal{P} , and such that the boundary of \mathcal{D} is a union of points of \mathcal{P} and

points in the boundary of \mathcal{D}_0 . An outer face is a face whose boundary contains points of the boundary of \mathcal{D}_0 .

A simple cycle is facial if it is the boundary of a face.

Every simple cycle in the disk bounds a unique simple region (the interior of the cycle), while a simple cycle on the sphere bounds two simple regions. A face does not have to be simple, but a facial cycle always bounds a simple face by definition.

6.2. Groups and labellings of pictures. Recall that r^+ refers to the even part of a relation $r \in \mathcal{F}_2(S) \times \mathbb{Z}_2$ (see Definition 4.4).

Definition 6.4. Let $G = \text{Inv}\langle S : R \rangle$. A *G*-picture is a picture \mathcal{P} with every vertex *v* labelled by a relation $r(v) \in R$ and every edge *e* labelled by a generator $s(e) \in S$, such that if e_1, \ldots, e_n is the sequence of edges incident to *v*, read in counter-clockwise order with multiplicity from some starting point, then $s(e_1)s(e_2)\cdots s(e_n) \in \{r(v)^+\}^{sym}$.

The boundary of \mathcal{P} is the cyclic word $\operatorname{bd}(\mathcal{P}) = s(e_1) \cdots s(e_n)$ over S, where e_1, \ldots, e_n is the list of edges incident with the boundary, read in counterclockwise order around the boundary of the disc, with multiplicity. If \mathcal{P} is closed then we say that $\operatorname{bd}(\mathcal{P}) = 1$, the empty word.

Two pictures \mathcal{P}_1 and \mathcal{P}_2 are equivalent if $\mathrm{bd}(\mathcal{P}_1) = \mathrm{bd}(\mathcal{P}_2)$.

The sign of a picture \mathcal{P} is sign $(\mathcal{P}) = |\{v \in V(\mathcal{P}) : r(v) \text{ is odd}\}| \mod 2$.

If \mathcal{D} is a simple region and \mathcal{P} is a *G*-picture, then res $(\mathcal{P}, \mathcal{D})$ and germ $(\mathcal{P}, \mathcal{D})$ both inherit the structure of a *G*-picture from \mathcal{P} via restricting the labelling functions.

Typically pictures use directed edges to represent inverses of generators, but this is not necessary for groups generated by involutions. As previously mentioned, the point of pictures is that they capture relations in the group, in the following sense:

Proposition 6.5 (van Kampen lemma). Let $G = \text{Inv}\langle S : R \rangle$, let r be a word over S, and let $a \in \mathbb{Z}_2$. Then $r = J^a$ in G if and only if there is a G-picture \mathcal{P} with $\text{bd}(\mathcal{P}) = r$ and $\text{sign}(\mathcal{P}) = a$.

The original version of the van Kampen lemma goes back to [VK33]. We give a proof of this version for the convenience of the reader.

Proof of Proposition 6.5. Given any word $w = s_1 \cdots s_n$ over S, we abuse notation slightly and let w^{-1} refer to the word $s_n \cdots s_1$. Because J is central, it is not hard to see that $r = J^a$ in G if and only if there is a sequence r_0, \ldots, r_n of (not necessarily reduced) words over S such that

• for every i = 1, ..., n, the word r_i can be constructed from r_{i-1} by either replacing a subword w_0 with w_1 , where $w_0 w_1^{-1}$ is the even part

 w^+ of some relation $w \in R^{sym}$, or inserting or deleting s^2 for some $s \in S$;

- *a* is (the parity of) the number of replacements $w_0 \to w_1$ in the point above, in which the corresponding relation $w \in \mathbb{R}^{sym}$ is odd; and
- $r_0 = r$ and $r_n = 1$.

From any such sequence, it is not hard to construct a *G*-picture \mathcal{P} with $bd(\mathcal{P}) = r$ and $sign(\mathcal{P}) = a$. Indeed, for each $i = 1, \ldots, n$ we can build a *G*-picture \mathcal{P}_i in a rectangle with $bd(\mathcal{P}_i) = r_{i-1}r_i^{-1}$, where the boundary edges labelled by r_{i-1} are connected to the top of the rectangle, and the boundary edges labelled by r_i^{-1} are connected to the bottom of the rectangle, using the following recipe:

- If $r_{i-1} = xw_0y$ and $r_i = xw_1y$, where x, y are words over S and $w_0w_1^{-1} \in \{w^+\}^{sym}$ for some $w \in R$, then \mathcal{P}_i is a G-picture with a single vertex labelled by w. This vertex is connected to the top half of the rectangle by edges labelled (from right to left) by w_0 and to the bottom half by edges labelled by w_1 . To the right and left of the vertex we add edges connecting the top to the bottom of the rectangle, labelled by x and y respectively.
- If $r_{i-1} = xy$ and $r_i = xs^2y$ for some $s \in S$, then \mathcal{P}_i is a picture with edges labelled (from right to left) by x and y connecting the top of the rectangle to the bottom, and an edge labelled by s with both endpoints incident to the bottom of the rectangle.
- If $r_{i-1} = xs^2y$ and $r_i = xy$ for some $s \in S$, then \mathcal{P}_i is a picture with edges labelled (from right to left) by x and y connecting the top of the rectangle to the bottom, and an edge labelled by s with both endpoints incident to the top of the rectangle.

Note that \mathcal{P}_n has no edges incident to the bottom of the rectangle. Putting these pictures together from top to bottom gives a picture \mathcal{P} with $\mathrm{bd}(\mathcal{P}) = r$ and $\mathrm{sign}(\mathcal{P}) = a$ as desired.

An example of this process with

$$G = \text{Inv}\langle a, b, c, d, f, g : adg = 1, bcf = 1, dfg = J \rangle$$

is given in Figure 8. The figure shows a picture \mathcal{P} with $bd(\mathcal{P}) = abc$ and $sign(\mathcal{P}) = 1$, corresponding to the identity abc = J. The picture is constructed by concatenating pictures $\mathcal{P}_1, \ldots, \mathcal{P}_7$ (demarcated by the dashed lines) corresponding to the derivation $abc = abca^2 = afa^2 = ad^2fa^2 = ad^2f = Jadg = Jg^2 = J$. As can be seen from the example, the algorithm outlined above can

26




FIGURE 8. A *G*-picture \mathcal{P} with $bd(\mathcal{P}) = abc$ and $sign(\mathcal{P}) = 1$ for $G = Inv\langle a, b, c, d, f, g : adg, bcf, Jdfg \rangle$

create pictures with closed loops, but these are allowed in our definition of G-pictures.

Conversely, suppose we are given a G-picture with $\operatorname{bd}(\mathcal{P}) = r$ and $\operatorname{sign}(\mathcal{P}) = a$. After isotopy, we can assume that \mathcal{P} is a picture in a rectangle in the plane, with all boundary edges incident to the top of the rectangle. We can also assume that every edge intersects every horizontal line through the rectangle in a finite number of points, and intersects only finitely many horizontal lines non-transversely. Say that a point on the interior of the diagram is a *critical point* if it is a vertex of \mathcal{P} , or a point on an edge which intersects a horizontal line non-transversely. By moving critical points up or down, we can assume that every horizontal line through the rectangle hits at most one critical point. We can then cut \mathcal{P} along horizontal lines into a sequence of pictures $\mathcal{P}_1, \ldots, \mathcal{P}_n$ each of which contains a single critical point. As in the above argument, this sequence can be turned into a derivation that $r = J^a$ in G.

Definition 6.6. Let \mathcal{D} be a simple region of a *G*-picture \mathcal{P}_0 , and let $\mathcal{P}' = \operatorname{res}(\mathcal{P}, \mathcal{D})$ or $\operatorname{germ}(\mathcal{P}, \mathcal{D})$. If \mathcal{P}' is equivalent to a picture \mathcal{P}'' , then we can cut out \mathcal{P}' and glue in \mathcal{P}'' in its place to get a new picture \mathcal{P}_1 . We refer to the process $\mathcal{P}_0 \Rightarrow \mathcal{P}_1$ as surgery.

If \mathcal{P}' has size zero,⁸ then we call $\mathcal{P}_0 \Rightarrow \mathcal{P}_1$ a null surgery. We say that \mathcal{P}_0 and \mathcal{P}_1 are equivalent via null surgeries if there is a sequence of null surgeries transforming \mathcal{P}_0 to \mathcal{P}_1 .

⁸Size zero pictures do not have vertices, but they can still have edges.




FIGURE 9. Surgery for an $S_4 \times \mathbb{Z}_2$ -picture with boundary relation $s_1s_2s_3s_3s_1 = s_2s_1s_2$.

Example 6.7. Consider the Coxeter group $S_4 \times \mathbb{Z}_2$ over \mathbb{Z}_2 . This group has presentation

 $G = \text{Inv}\langle s_1, s_2, s_3 : s_1 s_3 = s_3 s_1, s_1 s_2 s_1 = s_2 s_1 s_2, s_2 s_3 s_2 = s_3 s_2 s_3 \rangle,$

and in particular is presented by involutions. An example of a surgery for this group is shown in Figure 9.

7. Pictures over solution groups and hypergraphs

Let \mathcal{H} be a hypergraph with vertex labelling function $b : V(\mathcal{H}) \to \mathbb{Z}_2$. The solution group $\Gamma(\mathcal{H}, b)$ is finitely presented by involutions over \mathbb{Z}_2 ; to talk about Γ -pictures, we just need to pick a presentation of $\Gamma(\mathcal{H}, b)$. One candidate is the presentation from Definition 2.1. This presentation contains two types of relations: *linear relations* of the form $\prod x_e = J^a$, and *commuting relations* of the form $x_e x_{e'} = x_{e'} x_e$. However, it will be more convenient to use a presentation without commuting relations:

Definition 7.1. As a group presented by involutions over \mathbb{Z}_2 , we let $\Gamma(\mathcal{H}, b) =$ Inv $\langle S, R \rangle$, where $S = \{x_e : e \in E(\mathcal{H})\}$ and

$$R = \{J^{b_v} x_{e_1} \cdots x_{e_n} : all \ v \in V and all orderings \ e_1, \dots, e_n\}$$




FIGURE 10. In a solution group with linear relation $x_1x_2x_3x_4 = J^a$, the commuting relation $x_1x_2 = x_2x_1$ can be represented pictorially in two different ways, depending on whether we use the presentation from Definition 2.1 (on the left), or the presentation from Definition 7.1 (on the right).

of the edges incident to v listed with multiplicity}

Example 7.2. Consider the solution group $\Gamma(\mathcal{H}, b)$ for the simple hypergraph \mathcal{H} with a single vertex incident to four edges. In the presentation in Definition 2.1, $\Gamma(A, b)$ is generated by x_1, x_2, x_3, x_4, J subject to relations $J^2 = 1, x_i^2 = [x_i, J] = 1$ for $1 \leq i \leq 4$, commuting relations $[x_i, x_j] = 1$ for $1 \leq i < j \leq 4$, and the single linear relation $x_1x_2x_3x_4 = J^b$. In the presentation in Definition 7.1, the commuting relations are omitted, and the single linear relation is replaced by the 12 relations $x_{\sigma(1)}x_{\sigma(2)}x_{\sigma(3)}x_{\sigma(4)} = J^b$, where σ is a permutation of $\{1, 2, 3, 4\}$ (the other relations remain the same).

In general, all the linear relations in Definition 7.1 can be recovered from a single linear relation and the commuting relations, while a commuting relation can be recovered from two linear relations, as shown in Figure 10 (for the group in Example 7.2). Consequently, the presentations of $\Gamma(\mathcal{H}, b)$ in 2.1 and 7.1 are equivalent. We will always use the presentation in Definition 7.1 when working with Γ -pictures.

Definition 7.3. Let \mathcal{H} be a hypergraph. An \mathcal{H} -picture is a triple (\mathcal{P}, h_V, h_E) , where \mathcal{P} is a picture, and h_V and h_E are labelling functions $V(\mathcal{P}) \to V(\mathcal{H})$ and $E(\mathcal{P}) \to E(\mathcal{H})$ respectively, such that for all $v \in V(\mathcal{P})$ and $e' \in E(\mathcal{H})$, if we list the edges e_1, \ldots, e_n of \mathcal{P} incident to v with multiplicity then $A_{h_V(v)e'} =$ $|\{1 \leq i \leq n : h_E(e_i) = e'\}|$. For convenience, we usually write \mathcal{P} for the triple (\mathcal{P}, h_V, h_E) , and h for the labelling functions h_V and h_E .

The boundary of an \mathcal{H} -picture \mathcal{P} is the cyclic word $\operatorname{bd}(\mathcal{P}) = h(e_1) \cdots h(e_n)$ over $E(\mathcal{H})$, where as before e_1, \ldots, e_n is the sequence of edges incident with the boundary, read counter-clockwise with multiplicity. The character of \mathcal{P} is the vector $\operatorname{ch}(\mathcal{P}) \in \mathbb{Z}_2^{V(\mathcal{H})}$ with $\operatorname{ch}(\mathcal{P})_v = |h_V^{-1}(v)| \mod 2$.

It is easy to see that there is a one-to-one correspondence between $\Gamma(\mathcal{H}, b)$ pictures and \mathcal{H} -pictures. If \mathcal{P} is an \mathcal{H} -picture, then the sign of the corresponding Γ -picture is the standard dot product $\operatorname{ch}(\mathcal{P}) \cdot b$. The leads to the following restatement of the van Kampen lemma for \mathcal{H} -pictures.


Proposition 7.4 (van Kampen lemma). Let $\Gamma(\mathcal{H}, b)$ be a solution group. Then $x_{e_1} \cdots x_{e_n} = J^a$ in $\Gamma(\mathcal{H}, b)$ if and only if there is an \mathcal{H} -picture \mathcal{P} with $bd(\mathcal{P}) = e_1 \cdots e_n$ and $ch(\mathcal{P}) \cdot b = a$.

Example 7.5. Consider the solution group for the linear system

(1)
$$x + y + z = 1$$

(2) $x + y + z = 0$

The underlying hypergraph \mathcal{H} of this system is shown below.



This drawing of the hypergraph is also a closed \mathcal{H} -picture \mathcal{P} with character $\operatorname{ch}(\mathcal{P}) = (1,1)$. Since b = (1,0) and $\operatorname{ch}(\mathcal{P}) \cdot b = 1$, van Kampen's lemma tells us that J = 1.

Remark 7.6. Given a hypergraph \mathcal{H} with incidence matrix A, let $X \subset \mathbb{Z}_2^V$ be the set of vertex labellings b such that $J \neq 1$ in $\Gamma(\mathcal{H}, b)$, and let $Y \subset \mathbb{Z}_2^V$ be the set of characters $ch(\mathcal{P})$ of \mathcal{H} -pictures \mathcal{P} . It is not hard to see that X and Yare subspaces of \mathbb{Z}_2^V . Proposition 7.4 states that Y is the orthogonal subspace to X with respect to the standard bilinear product. By Theorem 2.3, X can be regarded as a quantum-information-theoretic analogue of the columnspace of A. From this point of view, Y is then an analogue of the left nullspace of A.

If edge e is incident to vertex v in an \mathcal{H} -picture \mathcal{P} , then h(e) will be incident to h(v) in \mathcal{H} . Consequently, the labelling function h in Definition 7.3 can be seen as a type of weak hypergraph homomorphism. If \mathcal{P} is closed, and \mathcal{H} and \mathcal{P} are simple loopless graphs, then h will be an actual graph homomorphism if and only if $h(v) \neq h(v')$ for all adjacent vertices v and v' in \mathcal{P} . Furthermore, if this happens then h must be a planar graph cover. This motivates the following definition:

Definition 7.7. Let \mathcal{H} be a hypergraph. A closed \mathcal{H} -picture \mathcal{P} is a cover of \mathcal{H} if every edge of \mathcal{P} is incident with two distinct vertices v and v' such that $h(v) \neq h(v')$.

30

Size, equivalence, restriction to a region, surgery, and null surgery are all defined for \mathcal{H} -pictures via the correspondence with Γ -pictures. For instance, the size of an \mathcal{H} -picture is simply the number of vertices in the picture.

Definition 7.8. Let \mathcal{H} be a hypergraph, and let $b : V(\mathcal{H}) \to \mathbb{Z}_2$ be a vertexlabelling function. Two \mathcal{H} -pictures \mathcal{P}_0 and \mathcal{P}_1 are b-equivalent (resp. characterequivalent) if $bd(\mathcal{P}_0) = bd(\mathcal{P}_1)$ and $ch(\mathcal{P}_0) \cdot b = ch(\mathcal{P}_1) \cdot b$ (resp. $ch(\mathcal{P}_0) = ch(\mathcal{P}_1)$).

An \mathcal{H} -picture \mathcal{P} is b-minimal (resp. character-minimal) if \mathcal{P} has minimum size among all b-equivalent (resp. character-equivalent) pictures.

Two pictures are character-equivalent if and only if they are b-equivalent for all vertex-labelling functions b. Thus a b-minimal picture is also character-minimal.

If $\mathcal{P}_0 \Rightarrow \mathcal{P}_1$ is a surgery in which a region \mathcal{P}' is replaced by an equivalent region \mathcal{P}'' , then $\mathrm{bd}(\mathcal{P}_0) = \mathrm{bd}(\mathcal{P}_1)$, and $\mathrm{ch}(\mathcal{P}_1) = \mathrm{ch}(\mathcal{P}_0) - \mathrm{ch}(\mathcal{P}') + \mathrm{ch}(\mathcal{P}'')$. So if \mathcal{P}' and \mathcal{P}'' are *b*-equivalent (resp. character equivalent), then \mathcal{P}_0 and \mathcal{P}_1 will be *b*-equivalent (resp. character-equivalent). We conclude that if \mathcal{P} is *b*-minimal (resp. character-minimal) then $\mathrm{res}(\mathcal{P}, \mathcal{D})$ and $\mathrm{germ}(\mathcal{P}, \mathcal{D})$ will be *b*-minimal (resp. character-minimal) for all simple regions \mathcal{D} (as otherwise we could replace $\mathrm{res}(\mathcal{P}, \mathcal{D})$ or $\mathrm{germ}(\mathcal{P}, \mathcal{D})$ with smaller pictures).

We can also remove closed loops without changing the character-equivalence class:

Lemma 7.9. Suppose \mathcal{P} is a picture, and let \mathcal{P}' be the same picture but with all closed loops deleted. Then \mathcal{P} and \mathcal{P}' are character-equivalent and have the same size.

8. A CATEGORY OF HYPERGRAPHS

Definition 8.1. Let $\mathcal{H} = (V, E, A)$ be a hypergraph. A subhypergraph of \mathcal{H} is a hypergraph $\mathcal{H}' = (V', E', A')$ with $V' \subset V$, $E' \subset E$, and $A'_{ve} = A_{ve}$ for all $v \in V'$ and $e \in E'$.

In other words, a subhypergraph is simply a subset of $V(\mathcal{H}) \cup E(\mathcal{H})$. Although this definition is substantially less restrictive than other notions of subhypergraphs in the literature, it is natural in the context of hypergraphs with isolated edges.

Definition 8.2. If \mathcal{H}' is a subhypergraph of \mathcal{H} , then the neighbourhood $\mathcal{N}(\mathcal{H}')$ of \mathcal{H}' is the subhypergraph with $V(\mathcal{N}(\mathcal{H}')) = V(\mathcal{H}')$, and

 $E(\mathcal{N}(\mathcal{H}')) = E(\mathcal{H}') \cup \{e \in E(\mathcal{H}) : e \text{ is incident in } \mathcal{H} \text{ to some vertex } v \in V(\mathcal{H}')\}.$ We say that \mathcal{H}' is open if $\mathcal{H}' = \mathcal{N}(\mathcal{H}').$

The proof of the following proposition is elementary, and we omit it.

Proposition 8.3. Let \mathcal{H} be a hypergraph. The collection of open subhypergraphs of \mathcal{H} forms a topology on $V(\mathcal{H}) \cup E(\mathcal{H})$. A subhypergraph \mathcal{H}' is closed in this topology if and only if, for all $v \in V(\mathcal{H})$, if v is incident to $e \in E(\mathcal{H}')$ then $v \in V(\mathcal{H}')$.

We use this topology when referring to closed subhypergraphs and the closure of a subhypergraph.

Definition 8.4. Let \mathcal{H}_1 and \mathcal{H}_2 be hypergraphs. A generalized morphism $\phi : \mathcal{H}_1 \to \mathcal{H}_2$ consists of a pair of functions

$$\phi_V: V(\mathcal{H}_1) \to V(\mathcal{H}_2) \cup \{\varepsilon\} \text{ and } \phi_E: E(\mathcal{H}_1) \to E(\mathcal{H}_2) \cup \{\varepsilon\},\$$

where ε is a special symbol not occurring in $V(\mathcal{H}_i) \cup E(\mathcal{H}_i)$, i = 1, 2, such that for all $v \in V(\mathcal{H}_1)$,

(1) if $\phi_V(v) \neq \varepsilon$, then

$$\sum_{e \in \phi_E^{-1}(e')} A(\mathcal{H}_1)_{ve} = A(\mathcal{H}_2)_{\phi_V(v)e'},$$

for all $e' \in E(\mathcal{H}_2)$, and

(2) if $\phi_V(v) = \varepsilon$, then

$$\sum_{e \in E(\mathcal{H}_1) \setminus \phi_E^{-1}(\varepsilon)} A(\mathcal{H}_1)_{v \epsilon}$$

is even, and $\phi_E(e_1) = \phi_E(e_2)$ for all edges $e_1, e_2 \in E(\mathcal{H}_1) \setminus \phi_E^{-1}(\varepsilon)$ incident to v.

When there is no confusion, we write ϕ for both ϕ_V and ϕ_E . The composition $\phi_2 \circ \phi_1$ of two morphisms $\phi_1 : \mathcal{H}_1 \to \mathcal{H}_2$ and $\phi_2 : \mathcal{H}_2 \to \mathcal{H}_3$ is defined by setting $\phi_2(\varepsilon) = \varepsilon$.

Proposition 8.5.

- (a) If v and e are incident in \mathcal{H}_1 , and $\phi : \mathcal{H}_1 \to \mathcal{H}_2$ is a generalized morphism with $\phi(v) \in V(\mathcal{H}_2)$, $\phi(e) \in E(\mathcal{H}_2)$, then $\phi(v)$ and $\phi(e)$ are incident in \mathcal{H}_2 .
- (b) If $\phi_1 : \mathcal{H}_1 \to \mathcal{H}_2$ and $\phi_2 : \mathcal{H}_2 \to \mathcal{H}_3$ are generalized morphisms, then $\phi_2 \circ \phi_1$ is a generalized morphism.

Proof. For part (a), let $e' = \phi(e) \in E(\mathcal{H}_2)$. If v and e are incident in \mathcal{H}_1 , and $\phi(v) \neq \varepsilon$, then $A(\mathcal{H}_2)_{\phi(v)e'} \geq A(\mathcal{H}_1)_{ve} > 0$. So e' and $\phi(v)$ are incident. For part (b), let $\phi = \phi_2 \circ \phi_1$. If $\phi_1(v) = \varepsilon$, then $\phi(v) = \phi_2(\varepsilon) = \varepsilon$ by definition. So

32

if $\phi(v) \neq \varepsilon$, then $\phi_1(v) \neq \varepsilon$, and

$$\sum_{e \in \phi^{-1}(e')} A(\mathcal{H}_1)_{ve} = \sum_{e'' \in \phi_2^{-1}(e')} \sum_{e \in \phi_1^{-1}(e'')} A(\mathcal{H}_1)_{ve}$$
$$= \sum_{e'' \in \phi_2^{-1}(e')} A(\mathcal{H}_2)_{\phi_1(v)e''} = A(\mathcal{H}_3)_{\phi(v)e'}$$

for all $e' \in E(\mathcal{H}_3)$.

Next, suppose $\phi(v) = \varepsilon$ and $\phi_1(v) \neq \varepsilon$. If $e_1, e_2 \in E(\mathcal{H}_1) \setminus \phi^{-1}(\varepsilon)$ are both incident to v, then $\phi(e_1)$ and $\phi(e_2)$ belong to $E(\mathcal{H}_2) \setminus \phi_2^{-1}(\varepsilon)$ and are incident to $\phi_1(v)$ by part (a). Thus $\phi(e_1) = \phi(e_2)$. Similarly,

$$\sum_{e \in E(\mathcal{H}_1) \setminus \phi^{-1}(\varepsilon)} A(\mathcal{H}_1)_{ve} = \sum_{e' \in E(\mathcal{H}_2) \setminus \phi_2^{-1}(\varepsilon)} \sum_{e \in \phi_1^{-1}(e')} A(\mathcal{H}_1)_{ve}$$
$$= \sum_{e' \in E(\mathcal{H}_2) \setminus \phi_2^{-1}(\varepsilon)} A(\mathcal{H}_2)_{\phi_1(v)e'}$$

is even, since $\phi_2(\phi_1(v)) = \varepsilon$.

Finally, suppose $\phi(v) = \phi_1(v) = \varepsilon$. If $E(\mathcal{H}_1) \setminus \phi^{-1}(\varepsilon)$ does not contain any edges incident with v, then part (2) of Definition 8.4 is trivially satisfied. Suppose on the other hand that $E(\mathcal{H}_1) \setminus \phi^{-1}(\varepsilon)$ contains edges incident with v. All such edges belong to $E(\mathcal{H}_1) \setminus \phi_1^{-1}(\varepsilon)$, and hence are sent by ϕ_1 to some common edge $e' \in E(\mathcal{H}_2)$, where $\phi_2(e') \neq \varepsilon$. Conversely, any edge of $E(\mathcal{H}_1) \setminus \phi_1^{-1}(\varepsilon)$ incident with v is sent by ϕ_1 to e', and hence belongs to $E(\mathcal{H}_1) \setminus \phi^{-1}(\varepsilon)$. We conclude that all edges of $E(\mathcal{H}_1) \setminus \phi^{-1}(\varepsilon)$ incident with vare sent to $\phi_2(e')$, and that

$$\sum_{e \in E(\mathcal{H}_1) \setminus \phi^{-1}(\varepsilon)} A(\mathcal{H}_1)_{ve} = \sum_{e \in E(\mathcal{H}_1) \setminus \phi_1^{-1}(\varepsilon)} A(\mathcal{H}_1)_{ve}$$

is even. Consequently ϕ is a generalized morphism.

Example 8.6. If e is an edge in a hypergraph \mathcal{H} , we can delete e to get a new hypergraph $\mathcal{H} \setminus e$. There is a generalized morphism $\mathcal{H} \to \mathcal{H} \setminus e$ which sends $e \mapsto \varepsilon$. Similarly, we construct other generalized morphisms by identifying edges, deleting isolated vertices, and collapsing vertices incident to an even number of edges (deleting a vertex and identifying all incident edges). An example of a series of these operations is shown in Figure 11.

Another way we can construct generalized morphisms is through subhypergraphs.

Proposition 8.7. Let \mathcal{H} be a hypergraph.




FIGURE 11. A generalized morphism ϕ between two hypergraphs. The morphism sends an edge or vertex x to x', with the following exceptions: $\phi(i) = \phi(2) = \phi(4) = \varepsilon$, $\phi(b) = a'$, $\phi(d) = \phi(f) = \phi(h) = c'$, and $\phi(8) = 6'$.

(a) If \mathcal{H}' is closed subhypergraph, then the function

$$r: V(\mathcal{H}) \cup E(\mathcal{H}) \to V(\mathcal{H}') \cup E(\mathcal{H}') \cup \{\varepsilon\} : x \mapsto \begin{cases} x & x \in V(\mathcal{H}') \cup E(\mathcal{H}') \\ \varepsilon & otherwise \end{cases}$$

is a generalized morphism $\mathcal{H} \to \mathcal{H}'$.

(b) If \mathcal{H}' is an open subhypergraph, then the inclusion map

 $\iota: V(\mathcal{H}') \cup E(\mathcal{H}') \to V(\mathcal{H}) \cup E(\mathcal{H}): x \mapsto x$

is a generalized morphism $\mathcal{H}' \to \mathcal{H}$.

Proof. If \mathcal{H}' is any subhypergraph, then part (1) of Definition 8.4 holds for r and part (2) holds vacuously for ι . If \mathcal{H}' is closed and $r(v) = \epsilon$, then $r(e) = \epsilon$ for all edges e incident to v, so part (2) of Definition 8.4 holds for r. If \mathcal{H}' is open, $v \in V(\mathcal{H}')$, and $e' \in E(\mathcal{H})$ is incident to v, then e' belongs to \mathcal{H}' , and hence part (1) holds for ι .

Definition 8.8. A subhypergraph \mathcal{H}' of \mathcal{H} is a retract if there is a generalized morphism $r : \mathcal{H} \to \mathcal{H}'$ such that $r|_{\mathcal{H}'}$ is the identity.

Part (a) of Proposition 8.7 shows that every closed subhypergraph is a retract.

Morphisms can also be constructed by gluing together morphisms over open subhypergraphs. **Proposition 8.9.** Let $\{\mathcal{H}_i\}_{i\in I}$ be a family of open subhypergraphs of \mathcal{H} such that $\bigcup \mathcal{H}_i = \mathcal{H}$, and let $\{\phi_i\}_{i\in I}$ be a family of generalized morphisms $\phi_i : \mathcal{H}_i \to \mathcal{H}'$ such that $\phi_i|_{\mathcal{H}_i \cap \mathcal{H}_j} = \phi_j|_{\mathcal{H}_i \cap \mathcal{H}_j}$. Then there is a unique generalized morphism $\phi : \mathcal{H} \to \mathcal{H}'$ such that $\phi_{|\mathcal{H}_i} = \phi_i$.

Proof. Clearly ϕ is uniquely defined as a function. Given $v \in V(\mathcal{H})$, find \mathcal{H}_i with $v \in V(\mathcal{H}_i)$, so $\phi(v) = \phi_i(v)$. Since \mathcal{H}_i is open, if $A_{ve}(\mathcal{H}) > 0$ then $e \in \mathcal{H}_i$ and $A(\mathcal{H}_i)_{ve} = A(\mathcal{H})_{ve}$. Consequently, if $\phi(v) \neq \varepsilon$ then

$$\sum_{e \in \phi^{-1}(e')} A(\mathcal{H})_{ve} = \sum_{\substack{e \in \phi^{-1}(e')\\e \in E(\mathcal{H}_i)}} A(\mathcal{H}_i)_{ve} = \sum_{e \in \phi_i^{-1}(e')} A(\mathcal{H}_i)_{ve} = A(\mathcal{H}')_{\phi(v)e'}$$

since ϕ_i is a morphism. Similarly, if $\phi(v) = \varepsilon$, then

$$\sum_{e \in E(\mathcal{H}) \setminus \phi^{-1}(\varepsilon)} A(\mathcal{H})_{ve} = \sum_{e \in E(\mathcal{H}_i) \setminus \phi^{-1}(\varepsilon)} A(\mathcal{H}_i)_{ve} = \sum_{e \in E(\mathcal{H}_i) \setminus \phi_i^{-1}(\varepsilon)} A(\mathcal{H}_i)_{ve}$$

is even, and $\phi(e_1) = \phi_i(e_1) = \phi_i(e_2) = \phi(e_2)$ for all edges $e_1, e_2 \in E(\mathcal{H}) \setminus \phi^{-1}(\varepsilon)$ incident to v.

By Proposition 8.5, part (b), hypergraphs with generalized morphisms form a category. The next proposition shows that the map $\mathcal{H} \mapsto \Gamma(\mathcal{H})$ extends to a functor from the category of hypergraphs to the category of groups over \mathbb{Z}_2 .

Proposition 8.10. Let $\phi : \mathcal{H}_1 \to \mathcal{H}_2$ be a generalized morphism. Then there is a morphism $\phi : \Gamma(\mathcal{H}_1) \to \Gamma(\mathcal{H}_2)$ over \mathbb{Z}_2 defined by

$$\phi(x_e) = \begin{cases} 1 & \phi_E(e) = \varepsilon \\ x_{\phi(e)} & otherwise \end{cases}$$

Proof. The morphism, if it exists, will be uniquely determined by the values $\phi(x_e), e \in E(\mathcal{H}_1)$. To show that the morphism is well-defined, we need to show that

(8.1)
$$\prod_{i=1}^{n} \phi(x_{e_i}) = 1$$

for every vertex v of \mathcal{H}_1 and ordering e_1, \ldots, e_n of the edges incident to v, listed with multiplicity. Suppose that $\phi(v) = \varepsilon$. If $\phi(e_i) = \varepsilon$ for all $1 \le i \le n$, then $\phi(x_{e_i}) = 1$ for all i, so Equation (8.1) holds. If $\phi(e_i) \ne \varepsilon$ for some i, then

$$\prod_{j=1}^{n} \phi(x_{e_j}) = \prod_{\phi(e_j) \neq \varepsilon} x_{\phi(e_j)} = x_{\phi(e_i)}^K = 1,$$

where the last equality holds because $K = \sum_{e \in E(\mathcal{H}_1) \setminus \phi_E^{-1}(\varepsilon)} A_{ve}$ is even. Hence Equation (8.1) holds in this case as well.




FIGURE 12. In the hypergraph of the cube (left), the open neighbourhood of the bottom face is a retract. In the hypergraph on the right, this is no longer the case.

If $\phi(v) \neq \varepsilon$, then for all $1 \leq i \leq n$, either $\phi(e_i) = \varepsilon$ or $\phi(e_i)$ is incident to $\phi(v)$. We conclude that $\phi(x_{e_1}), \ldots, \phi(x_{e_n})$ commute in $\Gamma(\mathcal{H}_2)$. Consequently

$$\prod_{i=1}^{n} \phi(x_{e_i}) = \prod_{e' \in E(\mathcal{H}_2)} \prod_{e \in \phi_E^{-1}(e')} \phi(x_e)^{A_{ve}} = \prod_{e' \in E(\mathcal{H}_2)} x_{e'}^{A_{\phi(v)e'}} = 1$$

We conclude that the morphism ϕ is well-defined.

As a consequence, open retracts are special:

Corollary 8.11. If \mathcal{H}' is an open subhypergraph of \mathcal{H} and a retract of \mathcal{H} then $\Gamma(\mathcal{H}')$ is a (semidirect factor) subgroup of $\Gamma(\mathcal{H})$.

Proof. Let r be the retraction morphism $\mathcal{H} \to \mathcal{H}'$, and let ι be the inclusion $\mathcal{H}' \to \mathcal{H}$. Then $r \circ \iota$ is the identity on \mathcal{H}' , so the composition $r \circ \iota : \Gamma(\mathcal{H}') \to \Gamma(\mathcal{H}) \to \Gamma(\mathcal{H})$ is the identity morphism, and consequently $\iota : \Gamma(\mathcal{H}') \to \Gamma(\mathcal{H})$ must be injective. It also follows immediately that $\Gamma(\mathcal{H}) = N \rtimes \Gamma(\mathcal{H}')$, where N is the kernel of $r : \Gamma(\mathcal{H}) \to \Gamma(\mathcal{H}')$.

Example 8.12. Consider the graph \mathcal{G} of a cube, shown on the left in Figure 12, with vertices numbered for reference. The open neighbourhood of $\{1, 2, 3, 4\}$ is the subhypergraph \mathcal{H} with vertex set $\{1, 2, 3, 4\}$ and edge set $\{12, 23, 34, 14, 15, 26, 37, 48\}$. The morphism $\mathcal{G} \to \mathcal{H}$ which is the identity on \mathcal{H} , and sends

$$5 \mapsto 1, \quad 6 \mapsto 2, \quad 7 \mapsto 3, \quad 8 \mapsto 4, \text{ and}$$

$$56 \mapsto 12, \quad 67 \mapsto 23, \quad 78 \mapsto 34, \quad 58 \mapsto 14,$$

is a retract of \mathcal{G} onto \mathcal{H} .

On the other hand, if we subdivide the edge 12 with a vertex as shown on the right of Figure 12, then the open neighbourhood \mathcal{H} of $\{1, 2, 3, 4, 9\}$ is no TSIRELSON'S PROBLEM AND AN EMBEDDING THEOREM




FIGURE 13. The generalized morphism from Figure 11 applied to a picture. A choice must be made in regards to the edges formerly incident to vertex 4. We have chosen to connect 1' - 3' and 5' - 7' instead of 3' - 7' and 1' - 5'.

longer a retract of \mathcal{G} , since any retract must send $6 \mapsto 2$ and $7 \mapsto 3$, but 23 is not an edge of \mathcal{H} . However, the open neighbourhood of $\{5, 6, 7, 8\}$ is still a retract, since we can define the morphism as above but with edges 29 and 39 sent to 67, and vertex 9 sent to ε .

Finally, we can also apply generalized morphisms to pictures.

Proposition 8.13. Let $\phi : \mathcal{H}_1 \to \mathcal{H}_2$ be a generalized morphism, and let \mathcal{P} be an \mathcal{H}_1 -picture with $bd(\mathcal{P}) = e_1 \cdots e_n$. Construct a new picture \mathcal{P}' as follows:

- (1) If e is an edge of \mathcal{P} such that $\phi(h(e)) = \varepsilon$, then delete e from \mathcal{P} .
- (2) For all remaining edges e of \mathcal{P} , change the label from $h(e) \in E(\mathcal{H}_1)$ to $\phi(h(e)) \in E(\mathcal{H}_2)$.
- (3) If v is a vertex of \mathcal{P} such that $\phi(h(v)) = \varepsilon$, then after applying steps (1) and (2), there is an even number of edges incident with v, and all have the same label in $E(\mathcal{H}_2)$. Delete v, and connect up the remaining incident edges so that no pair of edges cross.
- (4) For all remaining vertices v of \mathcal{P} , change the label from $h(v) \in E(\mathcal{H}_1)$ to $\phi(h(v)) \in E(\mathcal{H}_2)$.

Then \mathcal{P}' is an \mathcal{H}_2 -picture of size less than or equal to the size of \mathcal{P} , with $\mathrm{bd}(\mathcal{P}') = \phi(e_1) \cdots \phi(e_n)$, where ε is regarded as the empty word.

Proof. To show that this picture is an \mathcal{H}_2 -picture, we need to show that there are $A(\mathcal{H}_2)_{v'e'}$ edges labelled by e' incident to any vertex labelled by v' in \mathcal{P}' .

But this follows immediately from the construction and part (1) of Definition 8.4. Since the construction does not add any vertices, the size of \mathcal{P}' must be at most the size of \mathcal{P} .

An example is given in Figure 13. Since step (3) in Proposition 8.13 requires choosing a matching on edges, the graph \mathcal{P}' we end up with is not unique. One exception is when ϕ is one of the morphisms r or ι from Proposition 8.7; in this case, there is always a unique choice, and hence \mathcal{P}' is uniquely determined. However, in general the different choices can differ by a sequence of null-surgeries.

Definition 8.14. If $\phi : \mathcal{H}_1 \to \mathcal{H}_2$ is a generalized morphism and \mathcal{P} is an \mathcal{H}_1 -picture, we use $\phi(\mathcal{P})$ to denote either the null-surgery equivalence class of pictures constructed in Proposition 8.13, or some arbitrarily chosen representative of this class.

If $r : \mathcal{H} \to \mathcal{H}'$ is the retraction morphism onto a closed subhypergraph, and \mathcal{P} is a \mathcal{H} -picture, then we also denote $r(\mathcal{P})$ by $\mathcal{P}[\mathcal{H}']$.

If ι is the inclusion of an open subhypergraph \mathcal{H}' in \mathcal{H} , and \mathcal{P} is an \mathcal{H}' picture, then \mathcal{P} and $\iota(\mathcal{P})$ are essentially identical. In this case, Proposition 8.13 states the obvious fact that every picture over \mathcal{H}' can be regarded as a picture over \mathcal{H} . Note that if such a picture is character (or b)-minimal as an \mathcal{H} -picture, then it is also minimal as an \mathcal{H}' -picture; however, the converse is not true.

9. Cycles and outer faces

In this section, we lay the foundation for the proof of Theorem 5.1 by looking at the interaction between pictures and cycles in hypergraphs.

Definition 9.1. A cycle in a hypergraph \mathcal{H} is a closed subhypergraph \mathcal{C} which is a simple connected 2-regular graph. A cycle \mathcal{C} is cubic if every vertex $v \in V(\mathcal{C})$ has degree three in \mathcal{H} .

A C-cycle in an H-picture is a simple cycle C such that every edge of C is labelled by an edge of C.

Lemma 9.2. Let C be a cycle in a hypergraph \mathcal{H} , and suppose \mathcal{P} is an \mathcal{H} -picture such that the edges of C do not appear in $bd(\mathcal{P})$. Then every connected component of $\mathcal{P}[C]$ is a C-cycle.

Proof. $\mathcal{P}[\mathcal{C}]$ is a closed \mathcal{C} -picture in which every vertex has degree two. As a result, every connected component will be a simple closed curve.

In several upcoming proofs, we will use the following measure of the complexity of \mathcal{P} .

38

Definition 9.3. If C is a cycle in H, and P is an H-picture, then we let #Cycle(P,C) denote the number of C-cycles in P. If Φ is a collection of cycles, we let

$$\#\operatorname{Cycle}(\mathcal{P}, \Phi) = \sum_{\mathcal{C} \in \Phi} \#\operatorname{Cycle}(\mathcal{P}, \mathcal{C}).$$

If we include incident vertices, then a C-cycle C is a closed C-picture, and thus the definition of cover from Definition 7.7 applies. If C is both facial and a cover of C, then we say that C is a *facial cover*. Note that closed loops are not covers.

Proposition 9.4. If C is a cubic cycle in H, and P is a character-minimal H-picture with no closed loops, then every facial C-cycle in P is a facial cover.

Proof. Suppose that C is a facial C-cycle. Since C is not a loop at a vertex, and \mathcal{P} has no closed loops, every edge of C must be incident to two distinct vertices of \mathcal{P} . Suppose that C has an edge e connecting two vertices u_0 and u_1 with $h(u_0) = v = h(u_1)$. By hypothesis, v has degree 3 in \mathcal{H} , and thus is incident with e' = h(e), another edge f' of C, and an edge g' which is not in C. Hence each vertex u_i is incident with an edge f_i belonging to C with $h(f_i) = f'$, and an edge g_i not in C with $h(g_i) = g'$. Note that the edges f_0 and f_1 could be equal, as could g_0 and g_1 . Since C is facial, g_0 and g_1 must lie on the same side of C. Thus there is a surgery which removes u_0 and u_1 , connecting g_0 with g_1 and f_0 with f_1 , as shown below:



But this means that \mathcal{P} is not character-minimal.

A connected closed cover C of a cycle \mathcal{C} is determined up to isotopy by an orientation of \mathcal{C} , and the *ply*, that is, the size of $h^{-1}(v)$ for any $v \in V(\mathcal{C})$, where $h: C \to \mathcal{C}$ is the labelling function.

Definition 9.5. A C-cycle C is a copy of C if the labelling function $h : C \to C$ is a graph isomorphism, or equivalently if $|h^{-1}(v)| = 1$ for all $v \in V(C)$.

We are also interested in how different cycles interact.

Lemma 9.6. Let C be a cycle in \mathcal{H} , and let \mathcal{D} be a simple region in an \mathcal{H} picture \mathcal{P} . If C is a facial C-cycle in \mathcal{P} , then either C is contained in \mathcal{D} , or the
edges of germ (C, \mathcal{D}) form the boundaries of simple outer faces of germ $(\mathcal{P}, \mathcal{D})$.

Proof. Suppose C is not contained in \mathcal{D} . If germ (C, \mathcal{D}) is empty, then the lemma is vacuously true. Otherwise, choose $\epsilon > 0$ such that germ $(\mathcal{P}, \mathcal{D}) =$

res $(\mathcal{P}, \mathcal{D}^{\epsilon})$, and in particular such that C intersects the boundary of \mathcal{D}^{ϵ} transversally. Then $C \cap \mathcal{D}^{\epsilon}$ is divided into a number of segments which start and end on the boundary of \mathcal{D}^{ϵ} . Thus every face of germ (C, \mathcal{D}) will be a simple region whose boundary contains some (possibly disconnected) portion of the boundary of \mathcal{D}^{ϵ} . Each edge of germ (C, \mathcal{D}) will be incident to exactly two such faces, and one of these two faces will be contained in $\mathcal{F} \cap \mathcal{D}^{\epsilon}$, where \mathcal{F} is a face in \mathcal{P} bounded by C. Since \mathcal{F} is a face, any face of germ (C, \mathcal{D}) contained in $\mathcal{F} \cap \mathcal{D}^{\epsilon}$ will be a face of germ $(\mathcal{P}, \mathcal{D})$.

Lemma 9.7. Let C be a simple cycle in a picture \mathcal{P} , let \mathcal{D} be a simple region bounded by C, and let \mathcal{F} be a simple outer face of germ $(\mathcal{P}, \mathcal{D})$. Then the edges of germ $(\mathcal{P}, \mathcal{D})$ in the boundary of \mathcal{F} form a single simple path P with at least two edges.

Furthermore, if we write the edges of P in order as e_1, \ldots, e_n , then e_1 and e_n are incident with the boundary of \mathcal{D} , and e_2, \ldots, e_{n-1} belong to C.

Proof. Every edge of $\operatorname{bd}(\operatorname{germ}(\mathcal{P}, \mathcal{D}))$ is incident to a vertex in C. If \mathcal{F} is simple, then $\operatorname{bd}(\operatorname{germ}(\mathcal{P}, \mathcal{D}))$ must contain at least two edges. If these edges are incident with two or more vertices of C, then it follows from the definition of $\operatorname{germ}(\mathcal{P}, \mathcal{D})$ that every outer face is a simple region with boundary of the required form. If all the edges of $\operatorname{bd}(\operatorname{germ}(\mathcal{P}, \mathcal{D}))$ are incident with a single vertex in C, then all but one of the outer faces is simple, and the simple outer faces are bounded by a path formed by two edges of $\operatorname{bd}(\operatorname{germ}(\mathcal{P}, \mathcal{D}))$.

Definition 9.8. Let \mathcal{P} be a picture in \mathcal{D}_0 , where \mathcal{D}_0 is not the whole sphere. An outer quadrilateral of \mathcal{P} is the closure of a simple outer face \mathcal{F} of \mathcal{P} , whose boundary contains exactly three edges of \mathcal{P} , together forming a path of length three.

An outer quadrilateral must have a fourth side consisting of points in the boundary of \mathcal{D}_0 , hence the name. A picture with highlighted outer quadrilaterals is shown in Figure 14.

Proposition 9.9. Let C be a cycle in \mathcal{H} , and let C' be a cubic cycle such that $|E(C) \cap E(C')| \leq 1$. If \mathcal{D} is a region bounded by a C'-cycle C', and C is a facial C-cycle, then either C is contained in \mathcal{D} , or the edges of germ (C, \mathcal{D}) form outer quadrilaterals of germ $(\mathcal{P}, \mathcal{D})$.

Proof. By Lemma 9.6, we can assume that \mathcal{F} is a simple outer face of germ $(\mathcal{P}, \mathcal{D})$ formed by the edges of germ (C, \mathcal{D}) . By Lemma 9.7, the edges of germ (C, \mathcal{D}) in the boundary of \mathcal{F} form a single path $e_0, e_1, \ldots, e_{n+1}$, where e_0 and e_{n+1} are

40





FIGURE 14. A picture in the disk, with four outer faces, two of which (shaded) form outer quadrilaterals.

incident with the boundary and e_1, \ldots, e_n belong to C', as shown below:



Since C' is cubic, it follows as well that $n \ge 1$. Since $|E(\mathcal{C}) \cap E(\mathcal{C}')| \le 1$ and the edges e_1, \ldots, e_n belong to both C and C', we must have $h(e_1) = \ldots =$ $h(e_n) = e'$, the unique element of $E(\mathcal{C}) \cap E(\mathcal{C}')$. Since \mathcal{C} (and \mathcal{C}') are simple graphs, the edge e' is incident with exactly two vertices v and v', and both $A_{ve'} = A_{v'e'} = 1$. Thus n must be one, and we conclude that the closure of \mathcal{F} is an outer quadrilateral. \Box

10. Pictures over suns

In this section, we look at pictures over a specific family of hypergraphs:

Definition 10.1. The sun of size n, where $n \ge 3$, is the hypergraph with vertex set $\{1, \ldots, n\}$, edge set $\{e_i, f_i : 1 \le i \le n\}$, and incidence relation

$$A_{if_j} = \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases} \quad and \quad A_{ie_j} = \begin{cases} 1 & i \cong j \text{ or } i \cong j+1 \mod n \\ 0 & otherwise \end{cases}$$

Note that a sun has a unique cycle. The sun of size n = 6 is shown in Figure 15.

Proposition 10.2. Let \mathcal{H} be a sun, and let \mathcal{P} be a character-minimal \mathcal{H} picture such that $bd(\mathcal{P})$ does not contain any edges from the cycle \mathcal{C} of \mathcal{H} . Then \mathcal{P} is character-equivalent to a character-minimal picture \mathcal{P}' with no closed loops, in which every \mathcal{C} -cycle is a facial cover. Furthermore, \mathcal{P}' can be chosen so that every outer quadrilateral of \mathcal{P} is an outer quadrilateral of \mathcal{P}' .




FIGURE 15. The sun of size 6.

The last statement in this proposition implies in particular that if an edge e is part of an outer quadrilateral in \mathcal{P} , then e is also an edge of \mathcal{P}' , with the same label. The proof of Proposition 10.2 uses the following lemma.


Lemma 10.3. If \mathcal{P} is a closed picture over a sun \mathcal{H} , then $ch(\mathcal{P}) = 0$.

Proof. Let n be the size of \mathcal{H} , and let \mathcal{U} be the closure of the complement of the cycle \mathcal{C} of \mathcal{H} , or in other words, the subhypergraph with vertex set $\{1, \ldots, n\}$ and edge set $\{f_1, \ldots, f_n\}$. Then every vertex in $\mathcal{P}[\mathcal{U}]$ has degree one, so if \mathcal{P} is closed then $\mathcal{P}[\mathcal{U}]$ is a matching. Furthermore, if e is an edge of $\mathcal{P}[\mathcal{U}]$ with $h(e) = f_i$, and endpoints at vertices v and v', then h(v) = h(v') = i. We conclude that \mathcal{P} has an even number of vertices labelled by i, so $ch(\mathcal{P})_i = 0$ for every i.


Proof of Proposition 10.2. Without loss of generality, we can assume that every picture lies in a bounded region, rather than a sphere. As a consequence, every simple closed curve will bound a unique simple region. Suppose that Cis a C-cycle in some \mathcal{H} -picture \mathcal{P} , and let \mathcal{D} be the simple region bounded by C. For the purpose of this proof, we let $NE(C, \mathcal{P})$ be the number of edges ein res $(\mathcal{P}, \mathcal{D})$ with $h(e) \in \mathcal{U}$, where \mathcal{U} is the closure of the complement of C, as in the proof of Lemma 10.3. We then set

$$NE(\mathcal{P}) := \sum_{C \text{ a } \mathcal{C}\text{-cycle}} NE(C, \mathcal{P}).$$

Now suppose we start with some character-minimal picture $\mathcal{P} = \mathcal{P}_0$ with no edges from \mathcal{C} in $\mathrm{bd}(\mathcal{P}_0)$. Our strategy will be to reduce $NE(\mathcal{P}_0)$ via a sequence of surgeries. Let C be a \mathcal{C} -cycle in \mathcal{P}_0 , and let \mathcal{D} be the simple region bounded by C. Suppose e is an edge contained in \mathcal{D} and incident with C, such that $h(e) = f_i$ for some i. Clearly e cannot be incident with the boundary of \mathcal{P}_0 , and since e is not a closed loop, we conclude that e is incident with two distinct vertices v_0 and v_1 with $h(v_0) = h(v_1) = i$. Let e, a_0 , a_1 be the sequence of edges incident with v_0 , as they appear in counter-clockwise order, and let e, b_0 , b_1 be the edges incident with v_1 as they appear in clockwise order. Then a_0 , e, and b_0 all lie in the boundary of a common face, as do a_1 , e, and b_1 (all these edges may lie in a common non-simple face, so we have to be careful with wording here). If $h(a_0) = h(b_0)$ and $h(a_1) = h(b_1)$, then we could delete e, v_0 , and v_1 , and connect a_i with b_i (shown in the diagram below, where we assume without loss of generality that $h(a_0) = e_i$)




to get a character-equivalent picture of smaller size, contradicting minimality. We conclude that we must have $\{h(e), h(a_j), h(b_j)\} = \{e_{i-1}, e_i, f_i\}$ for j = 0, 1. Thus we can perform the surgery where we make v_j incident to e, a_j , and b_j , for j = 0, 1, (shown below, again with the assumption that $h(a_0) = e_i$)



to get a new picture \mathcal{P}_1 . Since the surgery did not change the number or labels of vertices, \mathcal{P}_1 will be character-minimal and character-equivalent to \mathcal{P}_0 . Since $\mathrm{bd}(\mathcal{P}_0)$ does not contain any edges from \mathcal{C} , the boundary of an outer quadrilateral of \mathcal{P}_0 must contain edges e_1, e_2, e_3 , where $h(e_2) \in \mathcal{C}$, and e_1 and e_3 are the two unique edges of $h^{-1}(\mathcal{U})$ which are incident with the endpoints of e_2 . Since e is not incident to the boundary, the edges a_0, a_1, b_0 , and b_1 and the vertices v_0 and v_1 do not belong to an outer quadrilateral of \mathcal{P}_0 , and hence every outer quadrilateral of \mathcal{P}_0 will be an outer quadrilateral of \mathcal{P}_1 .


Recall that e lies in \mathcal{D} . Without loss of generality, we can assume that v_0 belongs to C. By Lemma 9.2, v_1 must also belong to a \mathcal{C} -cycle. This creates two possible outcomes for $NE(\mathcal{P}_1)$. First, suppose that both v_0 and v_1 lie on C. Then the surgery will "pinch" the region bounded by C, creating two

C-cycles C_0 and C_1 connected by e, as shown below.



Any edge $e' \neq e$ in \mathcal{D} will end up in the region bounded by C_0 or the region bounded by C_1 , so $NE(C, \mathcal{P}_0) = NE(C_1, \mathcal{P}_1) + NE(C_2, \mathcal{P}_1) + 1$.

The other possibility is that v_1 lies on a different C-cycle C', where (since C-cycles cannot cross) C' lies in the interior of the region bounded by C. In this case, the surgery will connect C and C' to form a new cycle C'', as shown below.



The edge e, along with all the edges inside the cycle C', will end up on the outside of C''. The only edges remaining in the region bounded by C'' are edges that belonged to \mathcal{D} , so $NE(\mathcal{C}'', \mathcal{P}_1) = NE(C, \mathcal{P}_0) - NE(C', \mathcal{P}_0) - 1$. In both cases, no other cycles are changed by the surgery, so we conclude that $NE(\mathcal{P}_0) > NE(\mathcal{P}_1)$.

Iterating this procedure, we get a sequence $\mathcal{P}_0, \mathcal{P}_1, \mathcal{P}_2, \ldots$ of character-minimal pictures, all character-equivalent, such that the outer quadrilaterals of \mathcal{P}_i are outer quadrilaterals of \mathcal{P}_{i+1} , and $NE(\mathcal{P}_i) > NE(\mathcal{P}_{i+1})$. Since $NE(\mathcal{P}_i)$ cannot decrease indefinitely, this process must terminate at a picture \mathcal{P}_n with the property that if $e \in h^{-1}(\mathcal{U})$ is incident with a \mathcal{C} -cycle C, then e is not contained in the region bounded by C. Equivalently, we can say that $\operatorname{res}(\mathcal{P}_n, \mathcal{D})$ is closed for every simple region \mathcal{D} bounded by a \mathcal{C} -cycle.

Let \mathcal{P}' be the picture \mathcal{P}_n with all closed loops deleted. By Lemma 7.9, \mathcal{P}' is character-minimal and character-equivalent to \mathcal{P}_n (and hence \mathcal{P}_0). In addition, it is easy to see that all outer quadrilaterals of \mathcal{P}_n will be outer quadrilaterals of \mathcal{P}' , and that $\operatorname{res}(\mathcal{P}', \mathcal{D})$ will be closed for every simple region \mathcal{D} bounded by a \mathcal{C} -cycle in \mathcal{P}' . Suppose \mathcal{D} is a simple region bounded by a \mathcal{C} cycle such that $\operatorname{res}(\mathcal{P}', \mathcal{D})$ is non-empty. Since \mathcal{P}' does not contain any closed loops, $\operatorname{res}(\mathcal{P}', \mathcal{D})$ must contain a vertex. But since $\operatorname{res}(\mathcal{P}', \mathcal{D})$ is closed, we must

44


have $ch(res(\mathcal{P}', \mathcal{D})) = 0$ by Lemma 10.3, and hence we can delete $res(\mathcal{P}', \mathcal{D})$ from \mathcal{P}' to get a character-equivalent picture of smaller size, contradicting the minimality of \mathcal{P}' . We conclude that $\operatorname{res}(\mathcal{P}', \mathcal{D})$ must be empty for every simple region bounded by a \mathcal{C} -cycle, and hence every \mathcal{C} -cycle in \mathcal{P}' is facial. By Proposition 9.4, every \mathcal{C} -cycle in \mathcal{P}' is a facial cover, as required.

The following collorary is not needed for the proof of Theorem 5.1, but it does serve as a good example of how we will apply Proposition 10.2 in the following section.

Corollary 10.4. Let \mathcal{H} be the sun of size n, let $b: V(\mathcal{H}) \to \mathbb{Z}_2$ be a vertex labelling function, and set $b_0 := \sum_{i=1}^n b_i$. Then the subgroup of $\Gamma = \Gamma(\mathcal{H}, b)$ generated by $S = \{x_{f_1}, \ldots, x_{f_n}\}$ is isomorphic to $K = \text{Inv}\langle f_1, \ldots, f_n: f_1 \cdots f_n =$ $J^{b_0}\rangle$.

Proof. Clearly there is an \mathcal{H} -picture \mathcal{P} with $\mathrm{bd}(\mathcal{P}) = f_1 \cdots f_n$ and $\mathrm{ch}(\mathcal{P}) =$


(1,...,1), so $x_{f_1} \cdots x_{f_n} = J^{b_0}$ in Γ by Proposition 7.4. Conversely, suppose that $x_{f_{i_1}} \cdots x_{f_{i_k}} = J^c$ holds in Γ . By Proposition 7.4 again, there is an \mathcal{H} -picture \mathcal{P} with $\mathrm{bd}(\mathcal{P}) = f_{i_1} \cdots f_{i_k}$ and $\mathrm{ch}(\mathcal{P}) \cdot b = c$. Let \mathfrak{P} be the set of character-minimal pictures which are character-equivalent to \mathcal{P} , and in which every \mathcal{C} -cycle is a facial cover. By Proposition 10.2, \mathfrak{P} is non-empty, and since all elements of \mathfrak{P} have the same number of vertices, $\#\text{Cycle}(\mathcal{P}',\mathcal{C})$ is bounded for $\mathcal{P}' \in \mathfrak{P}$. Let \mathcal{P}' be an element of \mathfrak{P} which maximizes $\#\text{Cycle}(\mathcal{P}', \mathcal{C})$. If C is a C-cycle in \mathcal{P}' which is not a copy of \mathcal{C} , then it is possible to cut C into two C-cycles as shown below, where the interior of C is a face of \mathcal{P} .



Since this surgery does not change the character or modify any other \mathcal{C} -cycle in \mathcal{P} , we get an element of \mathfrak{P} with more \mathcal{C} -cycles than \mathcal{P}' , a contradiction. We conclude that every \mathcal{C} -cycle in \mathcal{P}' is a facial copy.

Since \mathcal{C} is cubic, each \mathcal{C} -cycle in \mathcal{P}' must bound a unique face, even if \mathcal{P} is a picture in a sphere. Let \mathcal{P}'' be the picture constructed by contracting each

one of these faces to a vertex, as shown below (for n = 6).



Since every edge in \mathcal{P}' labelled by an e_i must belong to a \mathcal{C} -cycle by Lemma 9.2, every edge remaining in \mathcal{P}'' is labelled by an f_i for some $1 \leq i \leq n$. If we label each vertex by the single relation $J^{b_0}f_1\cdots f_n$, then \mathcal{P}'' is a K-picture with $\mathrm{bd}(\mathcal{P}'') = f_{i_1}\cdots f_{i_k}$ and $\mathrm{sign}(\mathcal{P}'') = c$. By Proposition 6.5, the relation $f_{i_1}\cdots f_{i_k} = J^c$ holds in K.


The proof techniques of Proposition 10.2 and Corollary 10.4 are illustrated by the following example.

Example 10.5. Let \mathcal{H} be the sun of size 4, and let $K = \text{Inv}\langle f_1, \ldots, f_4 : f_1 f_2 f_3 f_4 = 1 \rangle$. It is not hard to see that the relation $w := f_1 f_2 f_3 f_4 (f_1 f_2 f_3)^2 = 1$ holds in K. Given a K-picture with boundary word w, we can replace every vertex with a C-cycle to get an equivalent \mathcal{H} -picture, for instance as shown in Figure 16c. However, there are \mathcal{H} -pictures with boundary word w, as shown in Figure 16a, which do not come from a K-picture in this way. Nonetheless, if \mathcal{P} is a minimal \mathcal{H} -picture whose boundary contains only f_i 's, then the proofs of Proposition 10.2 and Corollary 10.4 provide a method to transform \mathcal{P} to an equivalent picture which does come from a K-picture. The transformation process is shown in Figure 16.

To finish the section, we prove one more technical lemma:

Lemma 10.6. Let \mathcal{P} be a character-minimal picture with no closed loops over the sun \mathcal{H} of size n. Then there is no cycle in \mathcal{P} with edge labels contained in $\{e_i, f_i, f_{i+1}\}$ for some fixed $1 \leq i \leq n$ (where $f_{n+1} := f_1$).

Proof. Suppose C is a cycle of this form, and let \mathcal{U}_i be the closure of the subhypergraph $\{f_i, f_{i+1}\}$. Since \mathcal{P} has no closed loops, C must have at least one vertex. Since \mathcal{H} is simple, C must have at least two vertices, and there cannot be two consecutive edges of C with the same label. Thus $C[\mathcal{U}_i]$ is a (non-empty) matching between the vertices of C, where paired vertices must have the same label (either i or i+1, depending on whether the edge connecting them is labelled by an f_i or f_{i+1}). Since every vertex of C occurs in $C[\mathcal{U}_i]$, C must consist of a sequence of vertices $v_1, \ldots, v_{2n}, v_{2n+1} = v_1$, such that v_{2i-1}



(A) A single C-cycle with an edge labelled by f_i in its interior.

(B) We remove the edge in the interior of the C-cycle using the surgery highlighted in blue. We now have a picture with two facial covers of C.

(C) By cutting and reattaching the edges highlighted in red, we can turn the two facial covers into three facial copies of C.

FIGURE 16. Using surgery, we turn a picture over the sun of size 4 into an equivalent picture where all C-cycles are facial copies.

and v_{2i} are connected by an edge b_i with $h(b_i) \in \{f_i, f_{i+1}\}$, and v_{2i} and v_{2i+1} are connected by an edge labelled by e_i , i = 1, ..., n. Let a_i be the edge incident to v_i which is not in C. Since every vertex v_i is incident to an edge of C labelled by e_i , we conclude that

$$h(a_i) = \begin{cases} e_{i-1} & h(v_i) = i \\ e_{i+1} & h(v_i) = i+1 \end{cases},$$

where $e_0 := e_n$ and $e_{n+1} := e_1$. But since $h(v_{2i-1}) = h(v_{2i})$, this means that $h(a_{2i-1}) = h(a_{2i})$ for all i = 1, ..., n. Thus if we delete all the vertices and edges of C, we can connect a_{2i-1} and a_{2i} along the path previously taken by b_i to get a new \mathcal{H} -picture \mathcal{P}' which is character-equivalent to \mathcal{P} . But the size of \mathcal{P}' would then be strictly smaller than the size of \mathcal{P} , contradicting minimality.

11. Stellar cycles and constellations

In this section, we extend the results for suns from the previous section to more complicated hypergraphs through the notion of a *constellation*. The result is the "constellation theorem", which will be the key to the proof of Theorem 5.1.

Definition 11.1. Let \mathcal{H} be a hypergraph with vertex labelling function $b : V(\mathcal{H}) \to \mathbb{Z}_2$. A cycle \mathcal{C} in \mathcal{H} is b-stellar if

- (a) the neighbourhood $\mathcal{N}(\mathcal{C})$ is isomorphic to a sun,
- (b) $\mathcal{N}(\mathcal{C})$ is a retract of \mathcal{H} , and
- (c) $b_v = 0$ for all $v \in V(\mathcal{C})$.

Definition 11.2. Let \mathcal{H} be a hypergraph with vertex labelling $b : V(\mathcal{H}) \to \mathbb{Z}_2$. A b-constellation is a collection Φ of subhypergraphs of \mathcal{H} satisfying the following properties:

- (a) If $C \in \Phi$, then C is a cycle, the neighbourhood $\mathcal{N}(C)$ is isomorphic to a sun, and C is either:
 - (i) b-stellar, or
 - (ii) a sequence of edges $e_1e_2\cdots e_n$ (in order), $n \ge 3$, such that e_k belongs to a b-stellar cycle $\mathcal{C}' \in \Phi$ for all $3 \le k \le n$.
- (b) For every element $C \in \Phi$, either:
 - (i) there is an edge e in C which does not belong to any cycle in $\Phi \setminus \{C\}$, or
 - (ii) there is another cycle $\mathcal{C}' \in \Phi$ such that $E(\mathcal{C}) \cap E(\mathcal{C}') \neq \emptyset$, and \mathcal{C}' contains an edge e which does not belong to any cycle in $\Phi \setminus \{\mathcal{C}'\}$.
- (c) If $C_0, C_1 \in \Phi$, where $C_0 \neq C_1$, then $|E(C_0) \cap E(C_1)| \leq 1$, and if neither C_0 or C_1 is b-stellar, then $E(C_0) \cap E(C_1) = \emptyset$.

If Φ is a b-constellation, then a cycle C in a picture \mathcal{P} is a Φ -cycle if C is a \mathcal{C} -cycle for some $\mathcal{C} \in \Phi$.

Roughly speaking, property (a) in Definition 11.2 says that every cycle C in Φ is either stellar or (mostly) covered by other stellar cycles, while properties (b) and (c) state that the cycles in Φ do not overlap too much.

Example 11.3. Consider the hypergraph \mathcal{H} shown in Figure 17, and let b be the vertex labelling function with $b_9 = 1$ and $b_v = 0$ for $v \neq 9$. Let C_1 , C_2 , and C_3 be the cycles with vertex sets $\{1, 2, 5, 6\}$, $\{1, 4, 5, 8\}$, and $\{3, 4, 7, 8\}$ respectively, and let C_4 be the cycle with vertex set $\{1, 2, 3, 4, 9\}$. As in Example




FIGURE 17. A hypergraph based on Figure 12, but where the open neighbourhood of each face is a sun.

8.12, the neighbourhoods $\mathcal{N}(C_i)$, i = 1, 2, 3, are retracts of \mathcal{H} , and hence C_1, C_2 , and C_3 are b-stellar. The cycle C_4 is not a retract, nor is $b_v = 0$ for all $v \in V(C_4)$, so C_4 is not b-stellar. But $\mathcal{N}(C_4)$ is a sun, and the edges 12, 14, and 34 belong to C_1 , C_2 , and C_3 respectively. Thus $\Phi = \{C_1, \ldots, C_4\}$ is a b-constellation.

The cycle with vertex set $\{5, 6, 7, 8\}$ is also b-stellar, and could be added to this constellation, but the cycle with vertex set $\{2, 3, 6, 7, 9\}$ cannot be added since it is not b-stellar and shares edges with C_4 .

We can now state the main theorem of this section:

Theorem 11.4 (Constellation theorem). Let \mathcal{H} be a hypergraph with vertex labelling b, and let Φ be a b-constellation. Let \mathcal{P} be an \mathcal{H} -picture such that:

- (p.1) bd (\mathcal{P}) does not contain any edges from any cycle $\mathcal{C} \in \Phi$, and
- (p.2) either b = 0 or \mathcal{P} is closed.

Then \mathcal{P} is b-equivalent to a picture \mathcal{P}' such that all Φ -cycles in \mathcal{P}' are facial copies.

The rest of this section is concerned with the proof of Theorem 11.4. To aid the reader, the proof is split into a number of lemmas, which are in turn grouped into subsections. In all the lemmas, \mathcal{H} will be a hypergraph and b will be a vertex labelling. We refer to the hypotheses (p.1) and (p.2) of Theorem 11.4 as necessary.

11.1. Structure of constellations. We start by proving two lemmas about constellations.

Lemma 11.5. If Φ is a b-constellation in \mathcal{H} , and e is an edge of \mathcal{H} , then there are at most two cycles in Φ containing e.

Proof. Suppose \mathcal{C} is an element of Φ containing e. Since $\mathcal{N}(\mathcal{C})$ is a sun, \mathcal{C} is cubic. Thus if v is an endpoint of e, then there is a unique edge f incident to v and not contained in \mathcal{C} . If $\mathcal{C}' \in \Phi \setminus \{\mathcal{C}\}$ contains e, then \mathcal{C}' must also contain f, since $|E(\mathcal{C}) \cap E(\mathcal{C}')| \leq 1$. Hence if \mathcal{C}' and $\mathcal{C}'' \in \Phi \setminus \{\mathcal{C}\}$ both contain e, then $\{e, f\} \subset E(\mathcal{C}') \cap E(\mathcal{C}'')$, and consequently $\mathcal{C}' = \mathcal{C}''$.

Lemma 11.6. Let Φ be a b-constellation in \mathcal{H} , and let $\Phi' \subseteq \Phi$. If all elements of Φ' are b-stellar or Φ' contains all b-stellar elements of Φ , then Φ' is a b-constellation.

Proof. Every subset of Φ satisfies part (c) of Definition 11.2. If all elements of Φ' are *b*-stellar, or Φ' contains all *b*-stellar cycles in Φ , then Φ' also satisfies part (a). If *e* is an edge of $\mathcal{C} \in \Phi'$ such that *e* does not belong to any element of $\Phi \setminus \{\mathcal{C}\}$, then clearly *e* does not belong to any element of $\Phi' \setminus \{\mathcal{C}\}$. Suppose every edge of $\mathcal{C} \in \Phi'$ belongs to some element of $\Phi \setminus \{\mathcal{C}\}$. Then by definition there is a cycle $\mathcal{C}' \in \Phi$ such that $E(\mathcal{C}) \cap E(\mathcal{C}') \neq \emptyset$ and \mathcal{C}' contains an edge *e* which does not belong to any cycle in $\Phi \setminus \{\mathcal{C}\}$. If \mathcal{C}' belongs to Φ' , then we are done. If \mathcal{C}' does not belong to Φ' , then let *e'* be the unique edge of $E(\mathcal{C}) \cap E(\mathcal{C}')$. By Lemma 11.5, the only cycles of Φ which contain *e'* are \mathcal{C} and \mathcal{C}' , and hence no element of $\Phi' \setminus \{\mathcal{C}\}$ contains *e'*.

11.2. Stellar cycles. Now we turn to the core of the argument: showing that C-cycles can be turned into facial covers if C is stellar. The proof relies on hypothesis (p.2) in the following way:

Lemma 11.7. Suppose C is a b-stellar cycle in \mathcal{H} , and \mathcal{P} is a b-minimal \mathcal{H} picture satisfying hypothesis (p.2). Then every C-cycle in \mathcal{P} bounds a simple
region \mathcal{D} such that $ch(germ(\mathcal{P}, \mathcal{D})) \cdot b = 0$.

Proof. If b = 0, then the lemma is vacuously true. If \mathcal{P} is closed and $ch(\mathcal{P}) \cdot b = 0$, then \mathcal{P} is *b*-equivalent to the empty picture. But since \mathcal{P} is *b*-minimal, in this case \mathcal{P} must have size zero, and again the lemma is vacuously true.

Suppose that \mathcal{P} is closed and $ch(\mathcal{P}) \cdot b = 1$. Since \mathcal{P} is closed, we can think of \mathcal{P} as a picture in the sphere, in which case every \mathcal{C} -cycle bounds two simple regions \mathcal{D}_1 and \mathcal{D}_2 . Every vertex of \mathcal{P} appears in one of germ $(\mathcal{P}, \mathcal{D}_1)$ or germ $(\mathcal{P}, \mathcal{D}_2)$, with only the vertices of C appearing in both. Hence

$$\operatorname{ch}(\mathcal{P}) = \operatorname{ch}(\operatorname{germ}(\mathcal{P}, \mathcal{D}_1)) + \operatorname{ch}(\operatorname{germ}(\mathcal{P}, \mathcal{D}_2)) - \operatorname{ch}(C).$$

Since \mathcal{C} is b-stellar, $b_v = 0$ for all $v \in V(\mathcal{C})$, so $ch(\mathcal{C}) \cdot b = 0$. Consequently,

$$\operatorname{ch}(\operatorname{germ}(\mathcal{P}, \mathcal{D}_1)) \cdot b + \operatorname{ch}(\operatorname{germ}(\mathcal{P}, \mathcal{D}_2)) \cdot b = \operatorname{ch}(\mathcal{P}) \cdot b = 1$$

and we conclude that one of $ch(germ(\mathcal{P}, \mathcal{D}_i)) \cdot b, i = 1, 2$ must be 0.

50

Lemma 11.8. Let Φ be a b-constellation in which every cycle $C \in \Phi$ is bstellar. Let \mathcal{P} be an \mathcal{H} picture satisfying (p.1) and (p.2). Then \mathcal{P} is bequivalent to a b-minimal picture \mathcal{P}' with no closed loops, such that every Φ -cycle in \mathcal{P}' is facial.

Proof. The proof is by induction on the size of Φ . If Φ is empty, then the lemma is true by Definition 7.8 and Lemma 7.9. Suppose the lemma is true for all *b*-constellations of size *m*, where $m \geq 0$, and let Φ be a *b*-constellation of size m + 1 in which every cycle is *b*-stellar. If we pick an element $\mathcal{C} \in \Phi$, then $\Phi' := \Phi \setminus \{\mathcal{C}\}$ will be a *b*-constellation of size *m* by Lemma 11.6, so that every \mathcal{H} -picture satisfying (p.1) and (p.2) (with respect to Φ) will be *b*-equivalent to some *b*-minimal picture \mathcal{P} , also satisfying (p.1) and (p.2), such that every Φ' -cycle in \mathcal{P} is facial. Thus, to show that the lemma holds for Φ , we can assume that we start with a picture \mathcal{P} of this form.

Let $NF(\mathcal{P})$ denote the number of non-facial \mathcal{C} -cycles in \mathcal{P} , where $\mathcal{C} \in \Phi$ is the cycle chosen above. Similarly to the proof of Proposition 10.2, our strategy will be to perform a sequence of surgeries, starting from $\mathcal{P}_0 := \mathcal{P}$, each of which decreases $NF(\mathcal{P})$. Suppose that \mathcal{P} has a non-facial \mathcal{C} -cycle C. By Lemma 11.7, there is a simple region \mathcal{D} bounded by C such that $\operatorname{ch}(\operatorname{germ}(\mathcal{P},\mathcal{D})) \cdot b = 0$. Let $f: \mathcal{H} \to \mathcal{N}(\mathcal{C})$ be a retract onto $\mathcal{N}(\mathcal{C})$, so that $\widehat{\mathcal{P}} := f(\operatorname{germ}(\mathcal{P}, \mathcal{D}))$ is an $\mathcal{N}(\mathcal{C})$ -picture. Recall that, since $\mathcal{N}(\mathcal{C})$ is open, an $\mathcal{N}(\mathcal{C})$ -picture like $\widehat{\mathcal{P}}$ can be regarded as an \mathcal{H} -picture. By the definition of germ, the labels of edges and vertices in the boundary of the outer faces of germ $(\mathcal{P}, \mathcal{D})$ must belong to $\mathcal{N}(\mathcal{C})$. Since f is a retract, we conclude that the closures of the outer faces of $\operatorname{germ}(\mathcal{P}, \mathcal{D})$ will be identical to the closures of the outer faces of $\widehat{\mathcal{P}}$, and in particular, $\mathrm{bd}(\widehat{\mathcal{P}}) = \mathrm{bd}(\mathrm{germ}(\mathcal{P}, \mathcal{D}))$. The construction in Proposition 8.13 relabels or deletes vertices not in $\mathcal{N}(\mathcal{C})$, so $\widehat{\mathcal{P}}$ might not be character-equivalent to germ $(\mathcal{P}, \mathcal{D})$. But since \mathcal{C} is b-stellar, $b_v = 0$ for all $v \in V(\mathcal{C})$, and consequently, $ch(\widehat{\mathcal{P}}) \cdot b = 0$. We conclude that $\widehat{\mathcal{P}}$ is b-equivalent to germ $(\mathcal{P}, \mathcal{D})$. Since the size of $\widehat{\mathcal{P}}$ is less than or equal to the size of germ $(\mathcal{P}, \mathcal{D})$, and germ $(\mathcal{P}, \mathcal{D})$ is b-minimal, we also conclude that $\widehat{\mathcal{P}}$ is b-minimal as an \mathcal{H} -picture. It follows that $\widehat{\mathcal{P}}$ is b-minimal (and hence character-minimal) as an $\mathcal{N}(\mathcal{C})$ -picture. From, again, the definition of germ, we know that $\operatorname{bd}(\operatorname{germ}(\mathcal{P},\mathcal{D})) = \operatorname{bd}(\widehat{\mathcal{P}})$ does not contain any edges from \mathcal{C} . Finally, $\mathcal{N}(\mathcal{C})$ is a sun, so we can apply Proposition 10.2 to $\widehat{\mathcal{P}}$ to get an $\mathcal{N}(\mathcal{C})$ -picture $\widehat{\mathcal{P}}'$ such that $\widehat{\mathcal{P}}'$ is character-equivalent to and of the same size as $\widehat{\mathcal{P}}$, every \mathcal{C} -cycle in $\widehat{\mathcal{P}}'$ is facial, every outer quadrilateral of $\widehat{\mathcal{P}}$ is an outer quadrilateral of $\widehat{\mathcal{P}}'$, and $\widehat{\mathcal{P}}'$ has no closed loops. In particular, $\widehat{\mathcal{P}}'$ is *b*-minimal and b-equivalent to germ(\mathcal{P}, \mathcal{D}), and every outer quadrilateral of germ(\mathcal{P}, \mathcal{D}) is an outer quadrilateral of $\widehat{\mathcal{P}}'$.

Let \mathcal{P}_1 be the result of replacing germ $(\mathcal{P}, \mathcal{D})$ with $\widehat{\mathcal{P}}'$. Clearly \mathcal{P}_1 is bminimal and b-equivalent to \mathcal{P} . Suppose C' is a \mathcal{C}' -cycle in \mathcal{P}_1 , where $\mathcal{C}' \in \Phi'$. Now $\operatorname{bd}(\mathcal{P}) = \operatorname{bd}(\mathcal{P}_1)$ does not contain any edges from the cycles of Φ , so by Lemma 9.2, every edge of \mathcal{P} which is labelled by an edge of \mathcal{C}' belongs to a unique \mathcal{C}' -cycle, and the same holds for \mathcal{P}_1 . If \mathcal{C}' contains a boundary edge e of $\widehat{\mathcal{P}}'$, then there is a \mathcal{C}' -cycle C'' in \mathcal{P} which also contains e. By hypothesis, C''is facial, and hence by Proposition 9.9, the edges of $\operatorname{germ}(C'', \mathcal{D})$ form outer quadrilaterals in germ(\mathcal{P}, \mathcal{D}). But since the outer quadrilaterals of germ(\mathcal{P}, \mathcal{D}) are also outer quadrilaterals of $\widehat{\mathcal{P}}'$, the edges of C'' are unchanged in \mathcal{P}_1 . Since e belongs to a unique \mathcal{C}' -cycle in \mathcal{P}_1 , we must have C' = C'', and hence C'is facial in \mathcal{P}_1 . On the other hand, if C' does not contain a boundary edge of $\widehat{\mathcal{P}}'$, then C' either does not intersect \mathcal{D} , or lies entirely inside of \mathcal{D} . In the former case, C' will also be a facial \mathcal{C}' -cycle in \mathcal{P} , and will remain facial in \mathcal{P}_1 . In the latter case, C' would have to consist only of edges labelled by $E' = E(\mathcal{C}') \cap E(\mathcal{N}(\mathcal{C}))$. Since $\mathcal{N}(\mathcal{C})$ is a sun and $|E(\mathcal{C}') \cap E(\mathcal{C})| \leq 1$, the intersection E' is either empty, or is equal to (in the notation of Definition 10.1) $\{e_i, f_i, f_{i+1}\}$ for some *i*. But since $\widehat{\mathcal{P}}'$ is character-minimal and has no closed loops, Lemma 10.6 implies that $\widehat{\mathcal{P}}'$ does not contain any cycles of this form. We conclude that every \mathcal{C}' -cycle in \mathcal{P}_1 remains facial. Finally, every \mathcal{C} -cycle in \mathcal{P}_1 belongs either to \mathcal{P}' or is inherited unchanged from \mathcal{P} , with the consequence that $NF(\mathcal{P}_1) < NF(\mathcal{P})$.

Iterating this procedure, we get a sequence $\mathcal{P}_0 = \mathcal{P}, \mathcal{P}_1, \ldots, \mathcal{P}_k$ of *b*-minimal pictures, all *b*-equivalent, such that all Φ' -cycles in \mathcal{P}_i are facial for $1 \leq i \leq k$, and such that all Φ -cycles in \mathcal{P}_k are facial. Deleting all closed loops from \mathcal{P}_k finishes the proof.

11.3. Covers versus copies. For the next lemma, we show that if all Φ -cycles are facial covers, then we can turn Φ -cycles into facial copies.

Lemma 11.9. Let Φ be a b-constellation. Suppose that \mathcal{P} is an \mathcal{H} -picture satisfying (p.1), and such that all Φ -cycles in \mathcal{P} are facial covers. Then \mathcal{P} is character-equivalent to a picture \mathcal{P}' in which all Φ -cycles are facial copies.

Proof. The proof is similar to the proof of Corollary 10.4. Let \mathfrak{P} be the set of pictures which are character-equivalent to \mathcal{P} , have the same size as \mathcal{P} , and in which all Φ -cycles are facial covers. Since all elements of \mathfrak{P} have the same number of vertices, $\#\text{Cycle}(\mathcal{P}', \Phi)$ is bounded across $\mathcal{P}' \in \mathfrak{P}$. Let \mathcal{P}' be an element of \mathfrak{P} which maximizes $\#\text{Cycle}(\mathcal{P}', \Phi)$.

Suppose that C is a C-cycle in \mathcal{P}' which is not a copy of C, where $C \in \Phi$ has an edge e which does not belong to any other cycle in Φ . Since C is a cover, there are two distinct edges e_1 and e_2 in C with $h(e_1) = h(e_2) = e$. As in the proof of Corollary 10.4, we can cut C at e_1 and e_2 to get a new picture




FIGURE 18. Surgery in the proof of Lemma 11.9. Edges labelled by e and f are cut and reconnected to turn three Φ -cycles into four Φ -cycles. The edges of C are dashed, while the edges of C_1 and C_2 are dotted. Interiors of cycles are faces in \mathcal{P}' and \mathcal{P}'' .

 \mathcal{P}'' , character-equivalent to \mathcal{P}' , in which C has been replaced by two facial covers. By Lemma 9.2 and the hypothesis on e, the edges e_1 and e_2 are not contained in any other Φ -cycle. Thus all other Φ -cycles in \mathcal{P}' are unchanged in \mathcal{P}'' , and \mathcal{P}'' will be an element of \mathfrak{P} with #Cycle $(\mathcal{P}', \Phi) < \#$ Cycle (\mathcal{P}'', Φ) , a contradiction. We conclude that if $\mathcal{C} \in \Phi$ has an edge which does not belong to any other cycle in Φ , then all \mathcal{C} -cycles in \mathcal{P}' are facial copies.

Now suppose that C is a C-cycle in \mathcal{P}' which is not a copy of C, where every edge of $C \in \Phi$ belongs to some other cycle of Φ . By Definition 11.2, part (b), there is another cycle $\mathcal{C}' \in \Phi$ such that $E(\mathcal{C}) \cap E(\mathcal{C}') = \{e\}$, and \mathcal{C}' contains an edge e' which does not belong to any cycle in $\Phi \setminus \{C'\}$. As above, there are distinct edges e_1 and e_2 in C such that $h(e_1) = h(e_2) = e$. By Lemma 9.2, each edge e_i belongs to a unique \mathcal{C}' -cycle C_i , i = 1, 2. Since C_1 and C_2 are copies of \mathcal{C}' , we must have $C_1 \neq C_2$. Let f_i be the unique edge in C_i with $h(f_i) = e'$. Thus we can construct a new picture \mathcal{P}'' by cutting and reconnecting e_i and f_i , i = 1, 2, as shown in Figure 18.

By Lemmas 9.2 and 11.5, C and C_i are the only Φ -cycles containing e_i , while C_i is the only Φ -cycle containing f_i . Thus all Φ -cycles in \mathcal{P}'' , aside from those shown in Figure 18, are Φ -cycles in \mathcal{P}' , and hence will be facial covers. Because C is a cover and C_1 and C_2 are copies, the new cycles created in Figure 18 will be facial covers. We conclude again that \mathcal{P}'' will be an element of \mathfrak{P} with #Cycle(\mathcal{P}', Φ) < #Cycle(\mathcal{P}'', Φ), a contradiction. Thus every Φ -cycle in \mathcal{P}' is a facial copy.

11.4. Cycles covered by other cycles. Finally, we prove two lemmas that will allow us to handle non-stellar cycles.

Lemma 11.10. Let $\Phi' \subseteq \Phi$ be a pair of b-constellations. Suppose that \mathcal{P} is an \mathcal{H} -picture satisfying (p.1) with respect to Φ , such that all Φ' -cycles in \mathcal{P} are facial covers. Let \mathcal{C}_0 be a connected closed subhypergraph of $\mathcal{C} \in \Phi \setminus \Phi'$,⁹ such that every edge of \mathcal{C}_0 is contained in an element of Φ' . If C is a \mathcal{C} -cycle in \mathcal{P} , then:

- (a) $C[\mathcal{C}_0]$ is a cover of \mathcal{C}_0 , and
- (b) if C_0 is a connected component of $C[C_0]$, then all edges not contained in C and incident with a vertex of C_0 lie on the same side of C.

Proof. Let e be an edge of C with $h(e) \in E(\mathcal{C}_0)$. By hypothesis and Lemma 9.2, e belongs to a \mathcal{C}' -cycle C', where $\mathcal{C}' \in \Phi'$. Since C' is a facial cover, e has two distinct endpoints a and b with $h(a) \neq h(b)$. We conclude that $C[\mathcal{C}_0]$ is a cover of \mathcal{C}_0 .

Now let x and y be the edges of \mathcal{P} incident to a and b respectively, but not contained in C. Since $|E(\mathcal{C}) \cap E(\mathcal{C}')| \leq 1$ and \mathcal{C} is simple, we conclude that x and y belong to C'. Let \mathcal{D} be a simple region bounded by C. By Proposition 9.9, since C' is facial the edges x, y, and e either belong to \mathcal{D} , or form an outer quadrilateral in germ $(\mathcal{P}, \mathcal{D})$. It follows that x and y lie on the same side of C. Since C_0 is connected, all edges not contained in C and incident to C_0 lie on the same side of C.

Lemma 11.11. Let $\Phi' \subseteq \Phi$ be a pair of b-constellations. Suppose that \mathcal{P} is a b-minimal picture with no closed loops satisfying (p.1) and (p.2), such that every Φ' -cycle is a facial cover. If every edge of $\mathcal{C} \in \Phi$ is contained in some element of Φ' , then every \mathcal{C} -cycle in \mathcal{P} is a facial cover.

Proof. Let C be a C-cycle. Applying Lemma 11.10 with $C_0 = C$, we get immediately that C is a cover, and that the edges incident to C all lie on the same side of C.

Suppose b = 0, and let \mathcal{D} be a simple region bounded by C. If all the edges incident to C lie in \mathcal{D} , then germ $(\mathcal{P}, \mathcal{D})$ is closed, and ch $(\text{germ}(\mathcal{P}, \mathcal{D})) \cdot b = 0$. Since \mathcal{P} is *b*-minimal, germ $(\mathcal{P}, \mathcal{D})$ must be *b*-minimal, and this implies that germ $(\mathcal{P}, \mathcal{D})$ must have size zero, in contradiction of the fact that germ $(\mathcal{P}, \mathcal{D})$ contains C. Thus all the edges incident to C lie outside the interior of C, and we conclude that res $(\mathcal{P}, \mathcal{D})$ is closed. But once again, this implies that res $(\mathcal{P}, \mathcal{D})$ must have size zero, and since \mathcal{P} has no closed loops, this implies that res $(\mathcal{P}, \mathcal{D})$ is empty. We conclude that C is facial.

54

⁹In other words, C_0 is either equal to C, or a path in C.

Now suppose that \mathcal{P} is closed. Similarly to the proof of Lemma 11.7, the fact that \mathcal{P} is *b*-minimal with size greater than zero implies that $ch(\mathcal{P}) \cdot b = 1$. Now *C* bounds two simple regions \mathcal{D}_1 and \mathcal{D}_2 in the sphere, where we assume that all edges incident to *C* are contained in \mathcal{D}_1 , so $res(\mathcal{P}, \mathcal{D}_2)$ is closed. If $ch(res(\mathcal{P}, \mathcal{D}_2)) \cdot b = 1$, then $res(\mathcal{P}, \mathcal{D}_2)$ is *b*-equivalent to \mathcal{P} , contradicting the *b*-minimality of \mathcal{P} . Thus $ch(res(\mathcal{P}, \mathcal{D}_2)) \cdot b = 0$, and as above, $res(\mathcal{P}, \mathcal{D}_2)$ must be empty. We conclude again that *C* is facial. \Box


11.5. **Proof of Theorem 11.4.** Given a *b*-constellation Φ , let Φ_0 be the set of *b*-stellar cycles in Φ , and let Φ_1 be the set of cycles $C \in \Phi$ such that every edge of C belongs to a cycle in Φ_0 . Then $\Phi_0 \subseteq \Phi_1$, and by Lemma 11.6, Φ_0 and Φ_1 are *b*-constellations.

Given a picture \mathcal{P}_0 satisfying (p.1) and (p.2), Lemma 11.8 states that we can find a *b*-equivalent picture \mathcal{P}_1 which is *b*-minimal and has no closed loops, such that every Φ_0 -cycle in \mathcal{P}_1 is facial. By Proposition 9.4, every Φ_0 -cycle in \mathcal{P}_1 is a facial cover. By Lemma 11.11, every Φ_1 -cycle in \mathcal{P}_1 is also a facial cover.

This leaves the cycles in $\Phi \setminus \Phi_1$. By definition, any element of $\Phi \setminus \Phi_1$ is non-stellar. If a non-stellar cycle $\mathcal{C} \in \Phi$ shares an edge with another cycle $\mathcal{C}' \in \Phi$, then \mathcal{C}' must be *b*-stellar. Hence $\Phi \setminus \Phi_1$ consists of the non-stellar cycles $\mathcal{C} \in \Phi$ which have an edge *e* not contained in any element of $\Phi \setminus \{\mathcal{C}\}$. For the purpose of this proof, we say that such an edge $e \in \mathcal{C}$ is *independent*. By part (a) of Definition 11.2, every element $\mathcal{C} \in \Phi \setminus \Phi_1$ has either one or two independent edges. In the latter case, the two edges will be incident with a common vertex of \mathcal{C} .

Suppose that C is a C-cycle in \mathcal{P}_1 , where $\mathcal{C} \in \Phi \setminus \Phi'$. Let \mathcal{C}_0 be the path containing the non-independent edges of \mathcal{C} , regarded as a closed subhypergraph. By Lemma 11.10, $C[\mathcal{C}_0]$ is a cover of \mathcal{C}_0 , and since \mathcal{C}_0 is a path rather than a cycle, all connected components of $C[\mathcal{C}_0]$ are copies of \mathcal{C}_0 . If $C[\mathcal{C}_0]$ is non-empty, we can write C as a sequence $C_1C_2\cdots C_{2k-1}C_{2k}$ of paths C_i , where $k \geq 1$, the path C_{2i-1} is a connected component of $C[\mathcal{C}_0]$ for all $i = 1, \ldots, k$, and the edges of C_{2i} are labelled by independent edges of \mathcal{C} for all $i = 1, \ldots, k$.

Let v_1 (resp. v_2) be the first (resp. last) vertex of C_1 , and let f_1 (resp. f_2) be the edge of C_{2k} (resp. C_2) which is incident to v_1 (resp. v_2). The labels $a_i = h(v_i)$ are the endpoints of the path C_0 in C. If C has one independent edge e, then e will join a_1 and a_2 , and $h(f_1) = h(f_2) = e$. By Lemma 11.10, every edge incident to C_1 lies on the same side of C, so we can cut and reconnect f_1 and f_2 as shown in Figure 19a. If C has two independent edges, then we can do something similar. In this case, there will be an independent edge $e_i = h(f_i)$ incident to a_i , i = 1, 2, and both edges will be incident with a third vertex c. Let g be the edge incident to c not in C. Since C is cubic, any cycle containing g must also contain either e_1 or e_2 . Since these edges are independent, g is



(B) two independent edges.

FIGURE 19. Surgery to cut apart the cycle C in the proof of Theorem 11.4.

not contained in any element of Φ . We can then cut f_1 and f_2 and reconnect them by adding two vertices labelled by c, and a edge labelled by g, as shown in Figure 19b.

These surgeries are not as well behaved as those considered previously: When \mathcal{C} has two independent edges, we end up increasing the size, so the result will no longer be *b*-minimal. When \mathcal{C} has a single independent edge, it is possible that $f_1 = f_2$, in which case we create a closed loop. However, in both cases we split C into two cycles $C^{(1)}$ and $C^{(2)}$, where $C^{(1)}$ is a facial copy of \mathcal{C} , and $C^{(2)}[\mathcal{C}_0]$ has fewer connected components than $C[\mathcal{C}_0]$. Furthermore, we do not change any other \mathcal{C} -cycle. And since we only change or add edges whose labels are not contained in any element of $\Phi \setminus \{\mathcal{C}\}$, we conclude that C is the only Φ -cycle changed by this surgery. As a result we may repeat this type of surgery to get a picture \mathcal{P}_2 (not necessarily *b*-minimal, and possibly containing closed loops) which is character-equivalent to \mathcal{P}_1 , and in which every Φ -cycle is either a facial cover, or labelled only by independent edges.

Let \mathfrak{P} be the collection of pictures which are character-equivalent to \mathcal{P}_2 , and in which every Φ -cycle is either a facial cover or labelled only by independent edges. Let \mathcal{P}_3 be an element of \mathfrak{P} of minimum size, and let \mathcal{P}_4 be the picture

 \mathcal{P}_3 with all closed loops deleted. Clearly \mathcal{P}_4 is also an element of \mathfrak{P} of minimum size. Suppose \mathcal{P}_4 has a \mathcal{C} -cycle C which is not a facial cover for some $\mathcal{C} \in \Phi$. By definition, C is labelled by independent edges of C. Since \mathcal{P}_4 has no closed loops and \mathcal{C} is simple, C has at least two edges. Consequently \mathcal{C} must have two independent edges, say e_1 and e_2 . As above, let c be the vertex incident to both e_1 and e_2 , and let g be the edge incident to c and not in C. Since every edge of C is labelled by e_1 or e_2 , every vertex of C must be labelled by c. We can now argue similarly to Lemma 10.6: C must consist of a sequence of edges f_1, \ldots, f_{2k} , where $h(f_{2i-1}) = e_1$ and $h(f_{2i}) = e_2$ for all $1 \le i \le k$. Let v_1, \ldots, v_{2k} be the vertices of C in order, so f_i has endpoints v_i and v_{i+1} , where $v_{2k+1} := v_1$, and let g_i be the edge of \mathcal{P}_4 incident to v_i with $h(g_i) = g$. Let \mathcal{P}_5 be the picture \mathcal{P}_4 with all the edges and vertices of C deleted, and g_{2i-1} and g_{2i} joined into a single edge along the path taken by f_{2i-1} . Since the edges e_1 , e_2 , and g do not belong to any element of $\Phi \setminus \{\mathcal{C}\}$, this process does not create or change any other Φ -cycle. Thus $\mathcal{P}_5 \in \mathfrak{P}$, in contradiction of the minimality of \mathcal{P}_4 . We conclude that every Φ -cycle in \mathcal{P}_4 is a facial cover, and the theorem follows from Lemma 11.9.

12. Proof of the embedding theorem

In this section we finish the proof of Theorem 5.1 (and thus complete the proof of Theorem 3.1). We continue with the notation from Section 5, so $\mathcal{I} := \text{Inv}\langle S : R \rangle$ is a presentation by involutions over \mathbb{Z}_2 , the set of relations is written as $R = \{r_1, \ldots, r_m\}$ where $r_i = J^{p_i} s_{i1} \cdots s_{in_i}$, G is the group with presentation \mathcal{I} , and $\mathcal{W} := \mathcal{W}(\mathcal{I})$ is the corresponding wagon wheel hypergraph. Although we do not yet assume that \mathcal{I} is collegial, for convenience we assume that the length n_i of the relation r_i is at least 4 for all $1 \leq i \leq m$ (by Remark 4.7, this assumption holds if \mathcal{I} is collegial). We also use the same notation for the vertices and edges of \mathcal{W} , so

$$V = \bigcup_{i=1}^{m} V_i, \text{ where } V_i := \{(i, j, k) : j \in \mathbb{Z}_{n_i}, 1 \le k \le 3\}, \text{ and}$$
$$E = S \sqcup \bigcup_{i=1}^{m} E_i, \text{ where } E_i := \{a_{ij}, b_{ij}, c_{ij}, d_{ij} : j \in \mathbb{Z}_{n_i}\}.$$

In addition, we make the following definitions:

- Let \mathcal{W}_i be the closed subhypergraph of \mathcal{W} containing vertices V_i and edges E_i . (The open neighbourhood $\mathcal{N}(\mathcal{W}_i)$ is shown in Figure 2.)
- Let \mathcal{A}_i , $1 \leq i \leq m$ be the cycle containing edges a_{i1} , b_{i1} , a_{i2} , b_{i2} , ..., a_{in_i} , b_{in_i} .
- Let \mathcal{B}_i , $1 \leq i \leq m$, be the cycle containing edges d_{i1}, \ldots, d_{in_i} .




FIGURE 20. To retract $\mathcal{N}(\mathcal{W}_i)$ onto $\mathcal{N}(\mathcal{B}_i)$, we remove intermediary vertices to get a simplified wagon wheel shape, and then fold this wagon wheel onto the central cycle. In the example shown above, $n_i = 4$.

- Let C_{ij} , $1 \leq i \leq m$, $j \in \mathbb{Z}_{n_i}$, be the cycle containing edges a_{ij} , b_{ij} , c_{ij} , d_{ij} , and $c_{i,j-1}$.
- Let $\Phi = \{\mathcal{C}_{ij} : 1 \leq i \leq m, j \in \mathbb{Z}_{n_i}\} \cup \{\mathcal{B}_i : 1 \leq i \leq m\}.$

Before we can prove Theorem 5.1, we need some preliminary lemmas.

Lemma 12.1. If b is an \mathcal{I} -labelling of \mathcal{W} , then there is a well-defined morphism $G \to \Gamma(\mathcal{W}, b)$ over \mathbb{Z}_2 sending $s \mapsto x_s$ for all $s \in S$.

Proof. There is a well-defined morphism $\mathcal{F}(S) \times \mathbb{Z}_2 \to \Gamma(\mathcal{W}, b)$ over \mathbb{Z}_2 sending $s \mapsto x_s$. As can be seen from Figure 2, there is a $\mathcal{N}(\mathcal{W}_i)$ -picture \mathcal{P} with $\mathrm{bd}(\mathcal{P}) = s_{i1} \cdots s_{in_i}$ and $\mathrm{ch}(\mathcal{P}) \cdot b = \sum_{v \in V_i} b_v = p_i$. By Proposition 7.4, the relation r_i holds in $\Gamma(\mathcal{W}, b)$ for all $1 \leq i \leq n$.

Lemma 12.2. $\mathcal{N}(\mathcal{B}_i)$ is a retract of \mathcal{W} for all $1 \leq i \leq n$.

Proof. Define $r : \mathcal{W} \to \mathcal{N}(\mathcal{B}_i)$ by

$$r((i', j, k)) = \begin{cases} \varepsilon & i' \neq i \\ \varepsilon & i = i' \text{ and } k = 1 \\ (i, j, 3) & i = i' \text{ and } k = 2, 3 \end{cases}$$

$$r(e) = \begin{cases} \varepsilon & e \in S \\ \varepsilon & e \in E_{i'} \text{ with } i \neq i' \\ c_{ij} & e = c_{ij} \\ d_{ij} & e \in \{a_{ij}, b_{ij}, d_{ij}\} \end{cases}$$

It is clear that r is the identity on $\mathcal{N}(\mathcal{B}_i)$. The only vertices mapped to ε which are incident with edges in $E(\mathcal{W}) \setminus r^{-1}(\varepsilon)$ are the vertices (i, j, 1) for $j \in \mathbb{Z}_{n_i}$.

For these vertices, the incident edges a_{ij} and b_{ij} are identified as required by condition (2) of Definition 8.4.

Since r is the identity on $\mathcal{N}(\mathcal{B}_i)$, condition (1) of Definition 8.4 holds for the vertices (i, j, 3), $1 \leq j \leq n_i$. The vertices (i, j, 2) are incident with three edges of $E(\mathcal{W}) \setminus r^{-1}(\varepsilon)$, namely $a_{i,j+1}, c_{ij}$, and $b_{i,j}$, and these edges are mapped to the three edges $d_{i,j+1}, c_{ij}$, and $d_{i,j}$ incident to (i, j, 3). We conclude that condition (1) of Definition 8.4 also holds for the vertices (i, j, 2), and hence r is a generalized morphism.

The map r can be visualized as deleting everything outside of \mathcal{W}_i to get a simplified wagon wheel shape, and then folding this wagon wheel onto $\mathcal{N}(\mathcal{B}_i)$. This is depicted in Figure 20.

Lemma 12.3. Let $s = s_{ij}$ for some $1 \le i \le m$ and $j \in \mathbb{Z}_{n_i}$. If R is cyclically reduced, and $\operatorname{mult}(s; r_{i'})$ is even for all $1 \le i' \le m$, then $\mathcal{N}(\mathcal{C}_{ij})$ is a retract of \mathcal{W} .

Proof. We start by showing that $\mathcal{N}(\mathcal{C}_{ij})$ is a retract of $\mathcal{N}(\mathcal{W}_i)$. Since \mathcal{W}_i depends only on the cyclic order of $s_{i1} \cdots s_{in_i}$, we can assume without loss of generality that j = 1. Suppose that $\operatorname{mult}(s; r_i) = 2k$, and let $1 = j_1 < j_2 < \cdots < j_{2k} \leq n_i$ be a list of the indices $1 \leq l \leq n_i$ such that $s_{il} = s$. Since R is cyclically reduced, $j_{i+1} > j_i + 1$ for all $i = 1, \ldots, 2k - 1$, and $j_{2k} < n_i$. For convenience, let

$$\mathcal{J}_{r} = \{1, j_{2}, \dots, j_{2k}\}, \quad \mathcal{J}_{l} = \{0, j_{2} - 1, \dots, j_{2k} - 1\},$$
$$\mathcal{J}_{r}^{odd} = \{1, j_{3}, j_{5} \dots, j_{2k-1}\}, \quad \mathcal{J}_{l}^{odd} = \{0, j_{3} - 1, j_{5} - 1, \dots, j_{2k-1} - 1\},$$
$$\mathcal{J}_{r}^{even} = \{j_{2}, j_{4}, \dots, j_{2k}\}, \quad \text{and} \quad \mathcal{J}_{l}^{even} = \{j_{2} - 1, j_{4} - 1, \dots, j_{2k} - 1\}.$$

These sets represent the indices of vertices and edges on the right and left of the cycles C_{ij_p} . To talk about edges which do not belong to these cycles, we also define

 $\overline{\mathcal{J}}_r = \{j_1 + 1, j_1 + 2, \dots, j_2 - 1, j_3 + 1, \dots, j_4 - 1, \dots, j_{2k-1} + 1, \dots, j_{2k} - 1\} \text{ and}$ $\overline{\mathcal{J}}_l = \{j_2 + 1, j_2 + 2, \dots, j_3 - 1, j_4 + 1, \dots, j_5 - 1, \dots, j_{2k} + 1, \dots, n_i\}.$ Define $q_i : \mathcal{N}(\mathcal{W}_i) \to \mathcal{N}(\mathcal{C}_{i1})$ by

$$q_i((i, j, k)) = \begin{cases} \varepsilon & k = 1 \text{ and } j \notin \mathcal{J}_r \\ (i, 1, 1) & k = 1 \text{ and } j \in \mathcal{J}_r \\ \varepsilon & k = 2, 3 \text{ and } j \notin \mathcal{J}_r \cup \mathcal{J}_l \\ (i, 0, k) & k = 2, 3 \text{ and } j \in \mathcal{J}_l^{odd} \cup \mathcal{J}_r^{even} \\ (i, 1, k) & k = 2, 3 \text{ and } j \in \mathcal{J}_r^{odd} \cup \mathcal{J}_l^{even}. \end{cases}$$

,




FIGURE 21. To retract $\mathcal{N}(\mathcal{W}_i)$ onto $\mathcal{N}(\mathcal{C}_{i1})$, we remove intermediary vertices and edges, and then fold up the remaining cycles like a napkin. In the example above, $n_i = 4$ and $\text{mult}(s; r_i) = 2$. In general, if $\text{mult}(s; r_i) = 2k$ then we make k folds.

$$q_{i}(a_{ij}) = \begin{cases} a_{i1} \quad j \in \mathcal{J}_{r}^{odd} \\ b_{i1} \quad j \in \mathcal{J}_{r}^{even} \\ b_{i0} \quad j \in \overline{\mathcal{J}}_{l} \\ a_{i2} \quad j \in \overline{\mathcal{J}}_{r} \end{cases}, \qquad q_{i}(b_{ij}) = \begin{cases} a_{i1} \quad j \in \mathcal{J}_{r}^{even} \\ b_{i1} \quad j \in \mathcal{J}_{r}^{odd} \\ b_{i0} \quad j \in \overline{\mathcal{J}}_{l} \\ a_{i2} \quad j \in \overline{\mathcal{J}}_{r} \end{cases}$$
$$q_{i}(c_{ij}) = \begin{cases} c_{i0} \quad j \in \mathcal{J}_{l}^{odd} \cup \mathcal{J}_{r}^{even} \\ c_{i1} \quad j \in \mathcal{J}_{r}^{odd} \cup \mathcal{J}_{l}^{even} \\ \varepsilon \quad j \notin \mathcal{J}_{r} \cup \mathcal{J}_{l} \end{cases}, \qquad q_{i}(d_{ij}) = \begin{cases} d_{i1} \quad j \in \mathcal{J}_{r} \\ d_{i0} \quad j \in \overline{\mathcal{J}}_{l} \\ d_{i2} \quad j \in \overline{\mathcal{J}}_{r} \end{cases}, \\ d_{i2} \quad j \in \overline{\mathcal{J}}_{r} \end{cases}$$
$$q_{i}(s') = \begin{cases} s \quad s' = s \\ \varepsilon \quad s' \in S \setminus \{s\} \end{cases}.$$

The map q_i can be visualized as deleting vertices and edges c_{ij} (the spokes of the wagon wheel) not in the cycles C_{ij_p} , and then folding up the cycles C_{ij_p} onto C_{i1} , alternating the directions of the folds after each C_{ij_p} like a napkin. The smallest example, when k = 1, is depicted in Figure 21. As in the proof of Lemma 12.2, it follows that q_i is a retract.

Now we look at $\mathcal{W}_{i'}$ for $i' \neq i$. If $\operatorname{mult}(s; r_{i'}) = 0$, then define

$$q_{i'}: \mathcal{N}(\mathcal{W}_{i'}) \to \mathcal{N}(\mathcal{C}_{ij}): x \mapsto \varepsilon.$$

If mult(s; $r_{i'}$) > 0, then find j' such that $s_{i'j'} = s$, and let $f : \mathcal{N}(\mathcal{W}_{i'}) \to \mathcal{N}(\mathcal{C}_{i'j'})$ be the retract defined above onto $\mathcal{N}(\mathcal{C}_{i'j'})$. Now $\mathcal{N}(\mathcal{C}_{i'j'})$ and $\mathcal{N}(\mathcal{C}_{ij})$ are both suns, so there is an isomorphism $g : \mathcal{N}(\mathcal{C}_{i'j'}) \to \mathcal{N}(\mathcal{C}_{ij})$ with g(s) = s, and we let $q_{i'} = g \circ f$.

The morphisms $q_{i'}$, $1 \leq i' \leq m$ all send $s' \in S$ to either ε if $s' \neq s$, or to s if s' = s. If $i' \neq i''$, then the open subhypergraphs $\mathcal{N}(\mathcal{W}_{i'})$ and $\mathcal{N}(\mathcal{W}_{i''})$ have no vertices in common. All common edges of $\mathcal{N}(\mathcal{W}_{i'})$ and $\mathcal{N}(\mathcal{W}_{i''})$ belong to S, so $q_{i'}$ and $q_{i''}$ agree on the intersection. Every vertex of \mathcal{W} belongs to some \mathcal{W}_i . There may be elements s' of S which do not appear in any relation $r_{i'}$, and hence do not belong to any $\mathcal{N}(\mathcal{W}_i)$; for these edges, we can add additional morphisms which send $s' \mapsto \varepsilon$. By Proposition 8.9, there is a morphism $q: \mathcal{W} \to \mathcal{N}(\mathcal{C}_{ij})$ which agrees with $q_{i'}$ on $\mathcal{N}(\mathcal{W}_{i'})$, and in particular is a retract.

Lemma 12.4. Suppose \mathcal{I} is collegial. Then there is an \mathcal{I} -labelling b of \mathcal{W} such that Φ is a b-constellation.

Proof. Let b be any \mathcal{I} -labelling such that

- $|b^{-1}(1) \cap V(\mathcal{W}_i)| \le 1$ for all $1 \le i \le m$,
- b((i, j, 2)) = b((i, j, 3)) = 0 for all $1 \le i \le m$ and $j \in \mathbb{Z}_{n_i}$, and
- if b((i, j, 1)) = 1, then either mult $(s_{ij}, r_{i'})$ is odd for some $1 \le i' \le m$, or mult $(s_{ij'}, r_{i'})$ is even for all $j' \in \mathbb{Z}_{n_i}$ and $1 \le i' \le m$.

We will show that Φ is a *b*-constellation. First, we observe that \mathcal{B}_i is *b*-stellar for all $1 \leq i \leq n$. Indeed, $\mathcal{N}(\mathcal{B}_i)$ is a sun, and by Lemma 12.2, $\mathcal{N}(\mathcal{B}_i)$ is a retract of \mathcal{W} . Finally, $b|_{\mathcal{B}_i} = 0$, so \mathcal{B}_i is *b*-stellar. Similarly, Lemma 12.3 implies that \mathcal{C}_{ij} will be *b*-stellar for $1 \leq i \leq m$ and $j \in \mathbb{Z}_{n_i}$ as long as b((i, j, 1)) = 0and $\text{mult}(s_{ij}; r_{i'})$ is even for all $1 \leq i' \leq m$.

Suppose C_{ij} is not b-stellar. If $\operatorname{mult}(s_{ij}; r_{i'})$ is odd for some $1 \leq i' \leq m$, then by Definition 4.6, $\operatorname{mult}(s_{i,j+1}; r_{i''})$ and $\operatorname{mult}(s_{i,j-1}; r_{i''})$ are even for all $1 \leq i'' \leq m$. By the definition of b, we must have b((i, j + 1, 1)) = b((i, j - 1, 1)) = 0. Thus $C_{i,j+1}$ and $C_{i,j-1}$ are b-stellar, so C_{ij} (which consists of edges $a_{ij}, b_{ij}, c_{ij}, d_{ij}, c_{i,j-1}$) shares edges c_{ij}, d_{ij} , and $c_{i,j-1}$ with b-stellar cycles. If $\operatorname{mult}(s_{ij}; r_{i'})$ is even for all $1 \leq i' \leq m$, then we must have b((i, j, 1)) = 1. By the definition of b, this means that $\operatorname{mult}(s_{ij'}, r_{i'})$ is even for all $1 \leq i' \leq m$ and $j' \in \mathbb{Z}_{n_i}$, and b((i, j + 1, 1)) = b((i, j - 1, 1)) = 0. It follows that $C_{i,j+1}$ and $C_{i,j-1}$ are b-stellar, and once again C_{ij} will share edges c_{ij}, d_{ij} , and $c_{i,j-1}$ with b-stellar cycles. Thus Φ satisfies condition (a) of Definition 11.2.

The cycle C_{ij} is the only cycle in Φ containing edges a_{ij} and b_{ij} . Every edge d_{ij} of \mathcal{B}_i is also contained in C_{ij} , so Φ satisfies condition (b) of Definition 11.2.

Finally, it is easy to see that $|E(\mathcal{C}_{ij}) \cap E(\mathcal{D}_{i'})| \leq 1$ for all $1 \leq i, i' \leq m$ and $1 \leq i' \leq m$, and (since $n_i \geq 4$ for all $1 \leq i \leq m$) that $|E(\mathcal{C}_{ij}) \cap E(\mathcal{C}_{i'j'})| \leq 1$ for all distinct (i, j) and (i', j'). We showed above that if \mathcal{C}_{ij} is not b-stellar, then $\mathcal{C}_{i,j+1}$ and $\mathcal{C}_{i,j-1}$ are b-stellar, so there is no pair of non-b-stellar cycles in Φ with a common edge. Thus Φ satisfies condition (c) of Definition 11.2. \Box

Lemma 12.5. Let b be an \mathcal{I} -labelling, and let \mathcal{P} be a \mathcal{W} -picture in which all Φ -cycles are facial copies, and such that all edges in $\mathrm{bd}(\mathcal{P})$ belong to S. Then there is a G-picture \mathcal{P}' with $\mathrm{bd}(\mathcal{P}') = \mathrm{bd}(\mathcal{P})$ and $\mathrm{sign}(\mathcal{P}') = \mathrm{ch}(\mathcal{P}) \cdot b$.

Proof. Let $\mathcal{W} \setminus S$ denote the closed subhypergraph containing all vertices of \mathcal{W} and all edges except those in S. Equivalently, $\mathcal{W} \setminus S$ is the subhypergraph with connected components \mathcal{W}_i , $1 \leq i \leq m$. Suppose that \mathcal{P}_0 is a connected component of $\mathcal{P}[\mathcal{W} \setminus S]$. Since $\mathrm{bd}(\mathcal{P})$ does not contain any edges from outside S, \mathcal{P}_0 must be closed. Since \mathcal{P}_0 is connected, there must be some $1 \leq i \leq m$ such that \mathcal{P}_0 is a \mathcal{W}_i -picture. By Lemma 9.2, \mathcal{P}_0 contains a Φ -cycle. By hypothesis, every Φ -cycle in \mathcal{P}_0 is a facial copy, so every \mathcal{C}_{ij} -cycle will contain an edge labelled by d_{ij} . Thus \mathcal{P}_0 contains an edge labelled by d_{ij} for some $j \in \mathbb{Z}_{n_i}$, and by Lemma 9.2 again, \mathcal{P}_0 contains a \mathcal{B}_i -cycle B. Since $\mathcal{B}_i \in \Phi$, B is also a facial copy, and hence consists of edges \hat{d}_j , $j \in \mathbb{Z}_{n_i}$, such that $h(\hat{d}_j) = d_{ij}$. Let \hat{c}_j , $j \in \mathbb{Z}_{n_i}$, be the third edge incident to the common endpoint of \hat{d}_j and \hat{d}_{j+1} . Every edge \hat{d}_j is contained in a unique \mathcal{C}_{ij} -cycle C_j , which will consist of edges \hat{c}_j , \hat{d}_j , \hat{c}_{j-1} , and two additional edges \hat{a}_j and \hat{b}_j with $h(\hat{a}_j) = a_{ij}$ and $h(\hat{b}_j) = b_{ij}$. Let

$$\mathcal{P}_0' = B \cup \bigcup_{j \in \mathbb{Z}_{n_i}} C_j.$$

The common endpoint of \hat{a}_j and \hat{b}_j has degree two in $\mathcal{P}[\mathcal{W} \setminus S]$, while the endpoints of \hat{c}_j have degree three and are incident with \hat{c}_j , \hat{d}_j , \hat{d}_{j+1} and \hat{c}_j , \hat{a}_{j+1} , \hat{b}_j respectively. Hence \mathcal{P}'_0 contains every edge of $\mathcal{P}[\mathcal{W} \setminus S]$ incident to a vertex of \mathcal{P}'_0 . Thus \mathcal{P}'_0 is a maximal connected subgraph of $\mathcal{P}[\mathcal{W} \setminus S]$, so $\mathcal{P}'_0 = \mathcal{P}_0$. In particular, we conclude that $|h^{-1}(v) \cap V(\mathcal{P}_0)| = 1$ for all $v \in V(\mathcal{W}_i)$. Since bis an \mathcal{I} -labelling, it follows that $ch(\mathcal{P}_0) \cdot b = a_i$.

Now let $\mathcal{D}(B)$ be the closure of the face bounded by B, let $\mathcal{D}(C_j)$ be the closure of the face bounded by $C_j, j \in \mathbb{Z}_{n_i}$, and let

$$\mathcal{D}(\mathcal{P}_0) := \mathcal{D}(B) \cup \bigcup_{j \in \mathbb{Z}_{n_i}} \mathcal{D}(C_j).$$


Clearly \mathcal{P}_0 is contained in $\mathcal{D}(\mathcal{P}_0)$, and conversely every edge or vertex of \mathcal{P} in $\mathcal{D}(\mathcal{P}_0)$ belongs to \mathcal{P}_0 . Since \hat{a}_j and \hat{b}_j belong to the boundary of $\mathcal{D}(C_j)$ and are not contained in B or any $C_{j'}$, $j' \neq j$, we conclude that \hat{a}_j and \hat{b}_j belong to the boundary of $\mathcal{D}(\mathcal{P}_0)$. Every other edge of C_j belongs either to B, to C_{j+1} , or to C_{j-1} , so the boundary of $\mathcal{D}(\mathcal{P}_0)$ does not contain any other edges of C_j . Similarly, the boundary does not contain any edges of B, or any of the vertices $(i, j, 3), j \in \mathbb{Z}_{n_i}$. We conclude that $\mathcal{D}(\mathcal{P}_0)$ is bounded by the \mathcal{A}_i -cycle $A = \hat{a}_1 \hat{b}_1 \cdots \hat{a}_{n_i} \hat{b}_{n_i}$. In particular, $\mathcal{D}(\mathcal{P}_0)$ is a simple region, and $\operatorname{bd}(\operatorname{germ}(\mathcal{P}, \mathcal{D}(\mathcal{P}_0))) = s_{i1} \cdots s_{in_i}$ or $s_{in_i} \cdots s_{i1}$ depending on the orientation of A.

Since $\mathcal{D}(\mathcal{P}_0)$ contains only edges and vertices of \mathcal{P}_0 , if \mathcal{P}_1 is another connected component of $\mathcal{P}[\mathcal{W} \setminus S]$, then $\mathcal{D}(\mathcal{P}_0)$ and $\mathcal{D}(\mathcal{P}_1)$ are completely disjoint. Thus, as in the proof of Corollary 10.4, we can collapse each region $\mathcal{D}(\mathcal{P}_0)$ to a single vertex labelled by $r_i = J^{a_i} s_{i1} \cdots s_{in_i}$ to form a *G*-picture \mathcal{P}' . Since the edges of *S* will be unchanged, $\mathrm{bd}(\mathcal{P}) = \mathrm{bd}(\mathcal{P}')$. Since every vertex of \mathcal{P} belongs to a unique connected component of $\mathcal{P}[\mathcal{W} \setminus S]$, we conclude that $\mathrm{sign}(\mathcal{P}') = \mathrm{ch}(\mathcal{P}) \cdot b$.

Proof of Theorem 5.1. Let G be the group with presentation $\operatorname{Inv}\langle S: R\rangle$, and suppose that $\mathcal{I} = \operatorname{Inv}\langle S: R\rangle$ is collegial. Let $\mathcal{I}^+ = \operatorname{Inv}\langle S: R^+\rangle$, and let G^+ be the even quotient of G, as in Definition 4.4. By Lemma 12.4, we can choose an \mathcal{I} -labelling b of \mathcal{W} such that Φ is a b-constellation. Every b-stellar cycle is also 0-stellar, so Φ is also a 0-constellation by Definition 11.2. By Lemma 12.1, there are morphisms $\phi: G \to \Gamma(\mathcal{W}, b)$ and $\phi^+: G^+ \to \Gamma(\mathcal{W}, 0)$, both sending $s \mapsto x_s$ (note that 0 is an \mathcal{I}^+ -labelling).

To show that ϕ is injective, we start with ϕ^+ . If $\phi^+(w) = 1$ for some $w \in \mathcal{F}_2(S)$, then by Proposition 7.4 there is a \mathcal{W} -picture \mathcal{P} with $\mathrm{bd}(\mathcal{P}) = w$. By Theorem 11.4 (b = 0 case), we can choose \mathcal{P} so that all Φ -cycles in \mathcal{P} are facial copies. By Lemma 12.5, there is a G^+ -picture \mathcal{P}' such that $\mathrm{bd}(\mathcal{P}') = \mathrm{bd}(\mathcal{P})$, so w = 1 in G^+ . Since $\phi^+(J^aw) = 1$ for $a \in \mathbb{Z}_2$ and $w \in \mathcal{F}_2(S)$ if and only if a = 0 and $\phi^+(w) = 1$, it follows that ϕ^+ is injective.

Now there is a commutative diagram



where q_1 and q_2 are the quotient maps $G \to G^+ = (G/(J_G)) \times \mathbb{Z}_2$ and $\Gamma(\mathcal{W}, b) \to \Gamma(\mathcal{W}, 0) = (\Gamma(\mathcal{W}, b)/(J_{\Gamma})) \times \mathbb{Z}_2$ by J_G and $J_{\Gamma} := J_{\Gamma(\mathcal{W}, b)}$ respectively. Since J_G is central of order \leq two, we conclude that ker $q_1 = \{1, J_G\}$. Since ϕ^+ is injective, if $\phi(w) = 1$ for $w \in G$, then $q_1(w) = 1$, and consequently $w \in \{1, J_G\}$. Thus it remains only to show that $\phi(J_G) = 1$ if and only if $J_G = 1$ in G. By definition, $\phi(J_G) = J_{\Gamma}$, and if $J_{\Gamma} = 1$, then by Proposition 7.4 there is a closed \mathcal{W} -picture \mathcal{P} with $ch(\mathcal{P}) \cdot b = 1$. Since \mathcal{P} is closed, Theorem 11.4 again implies that we can choose \mathcal{P} so that all Φ -cycles in \mathcal{P} are facial copies. By Lemma 12.5, there is a closed G-picture \mathcal{P}' such that $sign(\mathcal{P}') = 1$, and we conclude that $J_G = 1$ in G. Thus ϕ is injective. \Box

References

- [AGR82] Alain Aspect, Philippe Grangier, and Gérard Roger, Experimental Realization of Einstein-Podolsky-Rosen-Bohm gedankenexperiment : A New Violation of Bell's Inequalities, Physical Review Letters 49 (1982), no. 2, 91–94.
- [Ark12] Alex Arkhipov, *Extending and Characterizing Quantum Magic Games*, arXiv:1209.3819 [quant-ph].
- [Bel64] J. S. Bell, On the Einstein Podolsky Rosen paradox, Physics 1 (1964), no. 3, 195–200.
- [Ber94] A. J. Berrick, Groups with no nontrivial linear representations, Bull. Austral. Math. Soc. 50 (1994), 1–11.
- [BPA⁺08] Nicolas Brunner, Stefano Pironio, Antonio Acín, Nicolas Gisin, André Allan Méthot, and Valerio Scarani, *Testing the Dimension of Hilbert Spaces*, Physical Review Letters **100** (2008), no. 21, 210503–4.
- [CHSH69] John F. Clauser, Michael A. Horne, Abner Shimony, and Richard A. Holt, Proposed Experiment to Test Local Hidden-Variable Theories, Physical Review Letters 23 (1969), no. 15, 880–884.
- [CHTW04] R. Cleve, P. Høyer, B. Toner, and J. Watrous, Consequences and limits of nonlocal strategies, 19th IEEE Annual Conference on Computational Complexity, 2004. Proceedings, 2004, pp. 236–249.
- [Cir80] B. S. Cirel'son, Quantum generalizations of Bell's inequality, Letters in Mathematical Physics 4 (1980), no. 2, 93–100.
- [CLS16] Richard Cleve, Li Liu, and William Slofstra, *Perfect commuting-operator* strategies for linear system games, arXiv:1606.02278.
- [CM14] Richard Cleve and Rajat Mittal, Characterization of Binary Constraint System Games, Automata, Languages, and Programming, Lecture Notes in Computer Science, no. 8572, Springer Berlin Heidelberg, 2014, arXiv:1209.2729, pp. 320– 331.
- [CS18] Andrea Coladangelo and Jalex Stark, Unconditional separation of finite and infinite-dimensional quantum correlations, arXiv:1804.05116.
- [DLTW08] Andrew C. Doherty, Yeong-Cherng Liang, Ben Toner, and Stephanie Wehner, The Quantum Moment Problem and Bounds on Entangled Multi-prover Games, 2012 IEEE 27th Conference on Computational Complexity (Los Alamitos, CA, USA), vol. 0, IEEE Computer Society, 2008, pp. 199–210.
- [DP16] Kenneth J. Dykema and Vern Paulsen, Synchronous correlation matrices and Connes' embedding conjecture, Journal of Mathematical Physics 57 (2016), no. 1, 015214.
- [DPP17] Ken Dykema, Vern I. Paulsen, and Jitendra Prakash, Non-closure of the set of quantum correlations via graphs, arXiv:1709.05032.
- [FC72] Stuart J. Freedman and John F. Clauser, Experimental Test of Local Hidden-Variable Theories, Physical Review Letters 28 (1972), no. 14, 938–941.
- [Fri12] Tobias Fritz, Tsirelson's problem and Kirchberg's conjecture, Reviews in Mathematical Physics 24 (2012), no. 05, 1250012.
- [Fri13] _____, On infinite-dimensional state spaces, Journal of Mathematical Physics 54 (2013), no. 5, 052107.
- [Fri16] _____, Quantum logic is undecidable, arXiv:1607.05870.
- [GHJ17] Li Gao, Samuel J. Harris, and Marius Junge, *Quantum Teleportation and Super-dense Coding in Operator Algebras*, arXiv:1709.02785.

- [Har18] Samuel J. Harris, *Bipartite matrix-valued tensor product correlations that are not finitely representable*, arXiv:1806.08745.
- [HBD⁺15] B. Hensen, H. Bernien, A. E. Dréau, A. Reiserer, N. Kalb, M. S. Blok, J. Ruitenberg, R. F. L. Vermeulen, R. N. Schouten, C. Abellán, W. Amaya, V. Pruneri, M. W. Mitchell, M. Markham, D. J. Twitchen, D. Elkouss, S. Wehner, T. H. Taminiau, and R. Hanson, *Loophole-free Bell inequality violation using electron spins separated by 1.3 kilometres*, Nature **526** (2015), no. 7575, 682–686.
- [Hig51] Graham Higman, A Finitely Generated Infinite Simple Group, Journal of the London Mathematical Society s1-26 (1951), no. 1, 61–64.
- [HK64] Rudolf Haag and Daniel Kastler, An Algebraic Approach to Quantum Field Theory, Journal of Mathematical Physics 5 (1964), no. 7, 848–861.
- [Ji13] Zhengfeng Ji, Binary Constraint System Games and Locally Commutative Reductions, arXiv:1310.3794 [quant-ph].
- [JNP⁺11] M. Junge, M. Navascués, C. Palazuelos, D. Pérez-García, V. B. Scholz, and R. F. Werner, *Connes' embedding problem and Tsirelson's problem*, Journal of Mathematical Physics 52 (2011), no. 1, 012102.
- [JPPG⁺10] M. Junge, C. Palazuelos, D. Pérez-García, I. Villanueva, and M. M. Wolf, Unbounded Violations of Bipartite Bell Inequalities via Operator Space Theory, Communications in Mathematical Physics **300** (2010), no. 3, 715–739.
- [Kir93] Eberhard Kirchberg, On non-semisplit extensions, tensor products and exactness of group C^{*}-algebras, Inventiones mathematicae **112** (1993), no. 1, 449– 489.
- [KKM⁺11] J. Kempe, H. Kobayashi, K. Matsumoto, B. Toner, and T. Vidick, *Entangled Games Are Hard to Approximate*, SIAM Journal on Computing 40 (2011), no. 3, 848–877.
- [KV11] Julia Kempe and Thomas Vidick, Parallel Repetition of Entangled Games, Proceedings of the Forty-third Annual ACM Symposium on Theory of Computing (New York, NY, USA), STOC '11, ACM, 2011, pp. 353–362.
- [LS77] R.C. Lyndon and P.E. Schupp, *Combinatorial group theory*, Classics in Mathematics, Springer, 1977.
- [LTW13] Debbie Leung, Ben Toner, and John Watrous, Coherent state exchange in multi-prover quantum interactive proof systems, Chicago Journal of Theoretical Computer Science 19 (2013), no. 1, 1–18.
- [Mer90] N. David Mermin, Simple unified form for the major no-hidden-variables theorems, Physical Review Letters 65 (1990), no. 27, 3373–3376.
- [Mer93] _____, Hidden variables and the two theorems of John Bell, Reviews of Modern Physics **65** (1993), no. 3, 803–815.
- [MV14] Laura Mančinska and Thomas Vidick, Unbounded Entanglement Can Be Needed to Achieve the Optimal Success Probability, Automata, Languages, and Programming, Lecture Notes in Computer Science, no. 8572, Springer Berlin Heidelberg, 2014, pp. 835–846.
- [NCPGV12] M. Navascués, T. Cooney, D. Pérez-García, and N. Villanueva, A Physical Approach to Tsirelson's Problem, Foundations of Physics 42 (2012), no. 8, 985–995.
- [NPA08] Miguel Navascués, Stefano Pironio, and Antonio Acín, A convergent hierarchy of semidefinite programs characterizing the set of quantum correlations, New Journal of Physics 10 (2008), no. 7, 073013.

- [Oza13] Narutaka Ozawa, About the Connes embedding conjecture, Japanese Journal of Mathematics 8 (2013), no. 1, 147–183.
- [Per90] Asher Peres, Incompatible results of quantum measurements, Physics Letters A 151 (1990), no. 3, 107–108.
- [PSS⁺16] Vern I. Paulsen, Simone Severini, Daniel Stahlke, Ivan G. Todorov, and Andreas Winter, *Estimating quantum chromatic numbers*, Journal of Functional Analysis **270** (2016), no. 6, 2188–2222.
- [PT15] Vern I. Paulsen and Ivan G. Todorov, *Quantum Chromatic Numbers Via Operator Systems*, The Quarterly Journal of Mathematics **66** (2015), no. 2, 677–692.
- [RS10] Neil Robertson and Paul Seymour, Graph minors XXIII. Nash-Williams' immersion conjecture, Journal of Combinatorial Theory, Series B 100 (2010), no. 2, 181–205.
- [RV15] Oded Regev and Thomas Vidick, *Quantum XOR Games*, ACM Trans. Comput. Theory **7** (2015), no. 4, 15:1–15:43.
- [Sho07] Hamish Short, *Diagrams and groups*, The Geometry of the Word Problem for Finitely Generated Groups, Birkhäuser Basel, Basel, 2007.
- [Slo17] William Slofstra, The set of quantum correlations is not closed, arXiv:1703.08618.
- [SW08] V. B. Scholz and R. F. Werner, *Tsirelson's Problem*, arXiv:0812.4305.
- [Tsi93] B.S. Tsirelson, Some results and problems on quantum Bell-type inequalities, Hadronic Journal Supplement 8 (1993), 329–345.
- [Tsi06] B.S. Tsirelson, Bell inequalities and operator algebras, problem statement for website of open problems at TU Braunschweig (2006), available at http://web.archive.org/web/20090414083019/http://www.imaph.tu-bs.de/qi/problems/33.ht
- [VK33] E. R. Van Kampen, On Some Lemmas in the Theory of Groups, American Journal of Mathematics 55 (1933), no. 1, 268–273.

INSTITUTE FOR QUANTUM COMPUTING AND DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF WATERLOO, CANADA

E-mail address: weslofst@uwaterloo.ca